Title: Partial genome characterization of a novel potentially zoonotic parapoxvirus in a horse,

Finland

Authors: Jenni Virtanen, Maria Hautaniemi, Lara Dutra, Ilya Plyusnin, Katja Hautala, Teemu

Smura, Olli Vapalahti, Tarja Sironen, Ravi Kant, Paula M. Kinnunen

Author affiliations: University of Helsinki, Helsinki, Finland (J. Virtanen, L. Dutra, I. Plyusnin, K.

Hautala, T. Smura, O. Vapalahti, T. Sironen, R. Kant, P.M. Kinnunen), Finnish Food Authority,

Helsinki (M. Hautaniemi), Medical University of Gdansk, 81–519 Gdynia, Poland (R. Kant)

Abstract

We report a sequencing protocol and 121 kb poxvirus sequence from a clinical sample of a horse

with dermatitis. Based on phylogenetic analyses, the virus is a novel parapoxvirus. We show

association with recent epidemic, and previous data suggest zoonotic potential. Further

characterization and development of diagnostic protocols are necessary.

Text

Parapoxviruses (PPVs) usually cause contagious skin infections in ruminants and occasionally

infect other species such as humans (*1*). The genus *Parapoxvirus* currently encompasses the

following recognized species: Orf virus (ORFV), Bovine Papular Stomatitis Virus (BPSV),

Pseudocowpoxvirus (PCPV), Red deerpox virus (RDPV), and Grey sealpox virus (GSEPV) (*2*). Of

these, ORFV, BPSV, PCPV, and RDPV are famously zoonotic. PPV genomes are usually 130-140

kb (*2*). Recently, poxviruses have emerged in humans and horses (*3, 4*).

A severe infection caused by a parapox-like virus (F14.1158H) was first verified from a horse

euthanized in Finland in 2013 (*5*). According to the short sequences (1.1 kb in total) obtained from

envelope phospholipase (ORF011) and RNA polymerase subunit RPO147 (ORF056) genes,

F14.1158H is most closely related to PPVs and is similar to the 585 bp sequences detected in

lesions from humans after contact with horses and donkeys in the United States (*5, 6*). However, the

actual classification remained unclear due to limited sequence data and lack of amplification in

numerous PPV-PCR assays (*5*).

No other clinical cases were confirmed until 2022 when an epidemic of dermatitis emerged in

horses across Finland. Parapoxvirus infection was subsequently identified in several cases using

pan-PPV-PCR (*7*) and Sanger sequencing (Appendix). Partial ORF011 sequences were 97%

identical to the 2013 case with identity of 79-87% to other parapoxviruses (Table A1). This

highlighted the need to properly characterise F14.1158H.

To better characterise the virus, DNA extracted directly from a skin lesion of the 2013 equine case

(*5*) was subjected to next-generation sequencing (NGS) with two different protocols (Appendix).

First protocol relying on a pool of poxvirus primers was insufficient to acquire enough sequence

data. With a PCR-free approach, utilising enrichment of the viral DNA, we acquired as much as 121

kb of nucleotide sequence, almost full genome, with coverage values of approximately 100 in 5

contigs (BioProject: PRJNA922554 and GenBank: OQ248663-OQ248667). Overall GC-content

was 68.4%, which is similar to PPVs *(2)*. Due to the lack of a reference genome, the data could not

be fully assembled and oriented. The lack of high-quality DNA and unsuccessful virus isolation

attempts (Appendix) further complicated the sequencing process. In the future, it is important to fill

the gaps to ensure finalization of a complete genome that can be used as a reference. In another

study, a combination of short- and long-read sequencing was used to recover the full genome of the

GSEPV *(8)*. However, with our clinical sample, the small amount of DNA available for sequencing

led to an alternative approach.

Phylogenetic analysis was conducted for the following poxvirus core genes *(9)* (ORF numbers

designated according to PPV ORFs *(10)*): DNA polymerase (ORF025), early transcription factor

VETFL (ORF083), DNA-directed RNA polymerase subunit RPO132 (ORF101), and DNA

topoisomerase type 1 (ORF062) (Figure). Consistent with earlier observations based on partial

sequences of ORF11 and ORF056, F14.1158H grouped clearly closer to PPVs than other

poxviruses (Figure A), although distinctly separate from the five currently recognized species

(Figure B-D). Amino acid sequences were more similar to PPV species than to other

Chordopoxviruses. For example, amino acid identity of DNA polymerase was 46-60% between

F14.1158H and viruses from other genera, 76-80% between F14.1158H and PPVs, and 84-95%

between the previously recognized PPV species (Table A3). Within the PPVs, F14.1158H generally

showed the second lowest pairwise nucleotide identity of the group (after the most divergent

GSEPV) (Table) with identities to other PPVs being 74-83% (ORF025), 73-83% (ORF083), 78-

87% (ORF101), and 84-91% (ORF062). GSEPV was consistently furthest from F14.1158H

whereas F14.1158Hs identities to other PPVs were similar. A relatively high difference explains

why F14.1158H was not detected by several PCRs designed for detecting PPV, which should be

considered when designing diagnostic protocols. These phylogenetic results and sequence identities

together with the high GC-content and disease characteristics indicate that F14.1158H is a novel PPV, Equine parapoxvirus (EqPPV). The final taxonomic position and the possible differences of human and equine-derived variants *(6)* will require more data.

Taken together, most known PPVs are zoonotic, and any novel virus detected in animals should be treated with concern *(6)*. Thus, considering the tendency of PPVs to cause disease, such as farmyard pox, in humans, EqPPV has a zoonotic potential. Hence, it is important to sample humans and other animals in contact with infected horses. It is also important to establish diagnostic protocols due to low specificity and sensitivity of pan-PPV-PCR for EqPPV (Appendix). Evidently, this virus is of veterinary importance posing a threat for horses and causing financial losses to owners. The information provided here will inform development of proper diagnostic tools enabling also establishment prevention measures.

Acknowledgements:

We thank DVM, PhD, Dipl. ECVP Niina Airas for her expert support regarding equine medicine. We also thank Mira Utriainen for technical assistance.

This study was financially supported by the Niemi Foundation, Finnish Foundation of Veterinary Research, and the Erkki Rajakoski Fund of Hippos Finland.

About the author: Ms. Virtanen is a post-doctoral researcher at the University of Helsinki in the field of clinical microbiology. Her research interests include zoonotic viruses and pathogens in human-animal interface.

Address for Correspondence: Jenni Virtanen, Department of Veterinary Biosciences, Faculty of

Veterinary Medicine, P.O. Box 66, and Department of Virology, Faculty of Medicine, P.O. Box 21,

00014 University of Helsinki, Helsinki, Finland; email: jenni.me.virtanen@helsinki.fi

Conflict of interest declaration: Besides holding the title of Adjunct Professor (Docent) of

University of Helsinki, PMK is an employee of MSD Animal Health. This study was initiated

before her joining the company. Other authors have nothing to declare.

References

1.   Essbauer S, Pfeffer M, Meyer H. Zoonotic poxviruses. Vet Microbiol. 2010 Jan 27;140(3-4):229-36.
2.   C. J. McInnes, I. K. Damon, G. L. Smith, G. McFadden, S. N. Isaacs, R. L. Roper, et al. ICTV Virus
Taxonomy Profile: Poxviridae 2023. Journal of General Virology. 2023;104:00184.
3.   Capobianchi MR, Di Caro A, Piubelli C, Mori A, Bisoffi Z, Castilletti C. Monkeypox 2022 outbreak in
non-endemic countries: Open questions relevant for public health, nonpharmacological intervention and
literature review. Front Cell Infect Microbiol. 2022;12:1005955.
4.   Ehmann R, Brandes K, Antwerpen M, Walter M, K VS, Stegmaier E, et al. Molecular and genomic
characterization of a novel equine molluscum contagiosum-like virus. J Gen Virol. 2021 Mar;102(3).
5.   Airas N, Hautaniemi M, Syrja P, Knuuttila A, Putkuri N, Coulter L, et al. Infection with Possible Novel
Parapoxvirus in Horse, Finland, 2013. Emerg Infect Dis. 2016 Jul;22(7):1242-5.
6.   Osadebe LU, Manthiram K, McCollum AM, Li Y, Emerson GL, Gallardo-Romero NF, et al. Novel
poxvirus infection in 2 patients from the United States. Clin Infect Dis. 2015 Jan 15;60(2):195-202.
7.   Inoshima Y, Morooka A, Sentsui H. Detection and diagnosis of parapoxvirus by the polymerase
chain reaction. J Virol Methods. 2000 Feb;84(2):201-8.
8.   Gunther T, Haas L, Alawi M, Wohlsein P, Marks J, Grundhoff A, et al. Recovery of the first full-length
genome sequence of a parapoxvirus directly from a clinical sample. Sci Rep. 2017 Jun 16;7(1):3734.
9.   Yu Z, Zhang W, Fu H, Zou X, Zhao M, Liang S, et al. Genomic analysis of Poxviridae and exploring
qualified gene sequences for phylogenetics. Comput Struct Biotechnol J. 2021;19:5479-86.
10.   Hautaniemi M, Ueda N, Tuimala J, Mercer AA, Lahdenpera J, McInnes CJ. The genome of
pseudocowpoxvirus: comparison of a reindeer isolate and a reference strain. J Gen Virol. 2010 Jun;91(Pt
6):1560-76.

Table: Nucleotide identity (as percentages) comparison between F14.1158H and other

parapoxviruses and Molluscum contagiosum virus in the four selected core genes.

	Molluscum contagiosum virus (MC515)	Grey sealpox virus	Red deerpox virus	Bovine Papular Stomatitis Virus (AR02)	Orf virus (UPM/HSN-20)	Orf virus (NZ2)	Pseudocowpoxvirus (F00.120R)	Pseudocowpoxvirus (VR634)
DNA polymerase (ORF025)								
Grey sealpox virus	62							
Red deerpox virus	68	79						
Bovine Papular Stomatitis Virus (AR02)	68	79	86					
Orf virus (UPM/HSN-20)	69	79	87	87				
Orf virus (NZ2)	69	79	87	87	99			
Pseudocowpoxvirus (F00.120R)	69	79	87	87	94	94		
Pseudocowpoxvirus (VR634)	70	80	88	88	95	95	98	
F14.1558H	69	74	81	81	82	82	82	83
Early transcription factor (ORF083)								
Grey sealpox virus	65							
Red deerpox virus	73	80						
Bovine Papular Stomatitis Virus (AR02)	74	80	90					
Orf virus (UPM/HSN-20)	73	79	88	89				
Orf virus (NZ2)	73	79	88	89	99			
Pseudocowpoxvirus (F00.120R)	74	80	90	91	95	95		
Pseudocowpoxvirus (VR634)	74	80	90	91	95	95	98	
F14.1558H	74	78	87	87	86	86	87	87
RNA polymerase (ORF101)								
Grey sealpox virus	73							
Red deerpox virus	79	84						
Bovine Papular Stomatitis Virus (AR02)	80	85	93					
Orf virus (UPM/HSN-20)	80	86	93	93				
Orf virus (NZ2)	80	86	93	93	100			
Pseudocowpoxvirus (F00.120R)	80	86	92	94	97	97		
Pseudocowpoxvirus (VR634)	80	86	93	94	98	98	98	
F14.1558H	80	84	89	91	91	91	91	91
Topoisomerase 1 (ORF062)								
Grey sealpox virus	62							
Red deerpox virus	68	79						
Bovine Papular Stomatitis Virus (AR02)	68	79	86					
Orf virus (UPM/HSN-20)	69	79	87	87				
Orf virus (NZ2)	69	79	87	87	99			
Pseudocowpoxvirus (F00.120R)	69	79	87	87	94	94		
Pseudocowpoxvirus (VR634)	70	80	88	88	95	95	98	
F14.1558H	69	74	81	81	82	82	82	83

A

B

C

D

Figure. A) Grouping of F14.1158H among all the genera of the subfamily *Chordopoxvirinae* in a

phylogenetic tree based on amino acid sequences of the DNA polymerase (ORF025) gene. B-D)

Grouping of F141158H among the genus Parapoxvirus in phylogenetic trees based on the

nucleotide sequences of the early transcription factor (ORF083) (B), RNA polymerase (ORF101)

(C), and topoisomerase 1 (ORF062) (D) genes. CRV = Crocodilepox virus, EMCLV = Equine

molluscum contagiosum-like virus, MOCV = Molluscum contagiosum virus, SQPV = Squirrelpox

virus, CPXV = Cowpox virus, MPXV = Monkeypox virus, VACV = Vaccinia virus, YMTV =

Yaba monkey tumor virus, SPPV = Sheeppox virus, MYXV = Myxoma virus, SWPV = Swinepox

virus, DPV = Deerpox virus, FWPV = Fowlpox virus, MsEPV = Melanoplus sanguinipes

entomopoxvirus. MsEPV is used as an outgroup in A and SQPV and MOCV in B-D. Bootstrap

values above 70% are shown next to the nodes.

139