**Title:** Partial genome characterization of a novel potentially zoonotic parapoxvirus in a horse,

Finland

**Authors:** Jenni Virtanen, Maria Hautaniemi, Lara Dutra, Ilya Plyusnin, Teemu Smura, Olli

Vapalahti, Tarja Sironen, Ravi Kant, Paula M. Kinnunen

**Author affiliations:** University of Helsinki, Helsinki, Finland (J. Virtanen, L. Dutra, I. Plyusnin, T.

Smura, O. Vapalahti, T. Sironen, R. Kant, P.M. Kinnunen), Finnish Food Authority, Helsinki (M.

Hautaniemi), Medical University of Gdansk, 81–519 Gdynia, Poland (R. Kant)

**Abstract**

We report a protocol and 121 kb poxvirus sequence from a clinical sample of a horse with

dermatitis diagnosed earlier. Based on phylogenetic analyses, the virus is a novel parapoxvirus.

Because it can cause epidemics and has zoonotic potential, proper characterization and diagnostic

protocols are necessary.

**Text**

Parapoxviruses (PPVs) usually cause contagious pustular skin infections in ruminants and occasionally infect other species such as humans (*1*). The genus *Parapoxvirus* currently encompasses the following five recognized species: Orf virus (ORFV), Bovine Papular Stomatitis Virus (BPSV), Pseudocowpoxvirus (PCPV), Red deerpox virus (RDPV), and Grey sealpox virus (*2*). Of these, ORFV, BPSV, PCPV, and RDPV are known to be zoonotic. PPV genomes are usually 130-140 kb (*2, 3*). Recently, poxviruses have emerged in humans and horses (*4, 5*).

A severe infection caused by a parapox-like virus (strain F14.1158H) was first verified from a horse euthanized in Finland in 2013 (*6*). No other cases were confirmed until 2022 when an epidemic of dermatitis emerged in racehorses across Finland (*7*). Parapoxvirus infection was subsequently identified in several cases (*8*). This brought a need to properly characterise F14.1158H. According to the short sequences (1.1 kb in total) obtained from envelope phospholipase (ORF011) and RNA polymerase subunit RPO147 (ORF056) genes in the previous study, F14.1158H is most closely related to PPVs and is similar to the sequences detected in lesions from humans who had been in contact with horses and donkeys in the United States, raising concerns about zoonotic risk (*6, 9*). However, the actual classification remained unclear due to limited sequence data and lack of amplification in numerous PPV-PCR assays (*6*).

To better characterise the virus, DNA extracted directly from a skin lesion of the first equine case (*6*) was subjected to next-generation sequencing (NGS) with two different protocols (Appendix). First protocol relying on a pool of poxvirus primers was insufficient in acquiring enough sequence data. With a PCR-free approach, utilising enrichment of the viral DNA, we acquired as much as 121 kb of nucleotide sequence, almost full genome, with coverage values of approximately 100 in 5 contigs (BioProject: PRJNA922554 and GenBank: OQ248663-OQ248667). Overall GC-content was 68.4%, which is similar to PPVs (*2*). Due to the lack of a reference genome, the data could not

be fully assembled and oriented. The lack of high-quality DNA and unsuccessful virus isolation

attempts further complicated the sequencing process. In the future, it is important to fill the gaps to

ensure finalization of a complete genome that can be used as a reference. Another approach, a

combination of short- and long-read sequencing, was used to recover the full genome of the Gray

sealpox virus (*10*). However, with our clinical sample, the small amount of DNA available for

sequencing led to an alternative approach.

Phylogenetic analysis was conducted for the following poxvirus core genes (*11*) (ORF numbers

designated according to PPV ORFs (*12*)): DNA polymerase (ORF025), early transcription factor

VETFL (ORF083), DNA-directed RNA polymerase subunit RPO132 (ORF101), and DNA

topoisomerase type 1 (ORF062) (Figure). Consistent with earlier observations based on partial

sequences of ORF11 and ORF056, F14.1158H grouped clearly closer to PPVs than other

poxviruses (Figure A), although distinctly separate from the five currently recognized species

(Figure B-D). Amino acid sequences were more similar to PPV species than to other

Chordopoxviruses. For example, amino acid identity of DNA polymerase was 46-60% between

F14.1158H and viruses from other genera, 76-80% between F14.1158H and PPVs, and 84-95%

between the five previously recognized PPV species (Table S2). Within the PPVs, F14.1158H

generally showed the second lowest pairwise nucleotide identity of the group (after the most

divergent Grey sealpox virus) (Table) with identities to other PPVs being 74-83% (ORF025), 73-

83% (ORF083), 78-87% (ORF101), and 84-91% (ORF062). Sealpox virus was consistently furthest

from F14.1158H whereas F14.1158Hs identities to other PPVs were similar. A relatively high

difference explains why F14.1158H was not detected by several PCRs designed for detecting PPV,

which should be considered when designing diagnostic protocols. These phylogenetic results and

sequence identities together with the high GC-content and disease characteristics indicate that

F14.1158H is a novel PPV. The final taxonomic position and the possible differences of human and

equine-derived variants (*9*) will require more data.

Taken together, most known PPVs are zoonotic and a similar virus to the one reported here has

been found in humans in the United States (*9*). Also, considering the tendency of other PPVs to

cause disease, such as farmyard pox, in humans, this virus has a zoonotic potential. In the future, it

is important to sample humans and other animals in contact with infected horses. Evidently, this

virus is of veterinary importance posing a threat for horses and also causing financial losses to

owners.  Hence, the information provided here will inform development of proper diagnostic tools

enabling also prevention measures to be established.

**Acknowledgements:**

We thank DVM Katja Hautala and DVM, PhD, Dipl. ECVP Niina Airas for their expert support

regarding equine medicine.

This study was financially supported by the Niemi Foundation, Finnish Foundation of Veterinary

Research, and the Erkki Rajakoski Fund of Hippos Finland.

**About the author:** Ms. Virtanen is a post-doctoral researcher at the University of Helsinki in the

field of clinical microbiology. Her research interests include zoonotic viruses and pathogens in

human-animal interface.

Address for Correspondence: Jenni Virtanen, Department of Veterinary Biosciences, Faculty of

Veterinary Medicine, P.O. Box 66, and Department of Virology, Faculty of Medicine, P.O. Box 21,

00014 University of Helsinki, Helsinki, Finland; email: jenni.me.virtanen@helsinki.fi

**Conflict of interest declaration:** Besides holding the title of Adjunct Professor (Docent) of

University of Helsinki, PMK is an employee of MSD Animal Health. This study was initiated

before her joining the company. Other authors have nothing to declare.

## References

1.      Essbauer S, Pfeffer M, Meyer H. Zoonotic poxviruses. Vet Microbiol. 2010 Jan 27;140(3-4):229-36.

2.      International Committee on Taxonomy of Viruses [cited 2023 Jan 5].
https://ictv.global/taxonomy/.

3.      Skinner MA, Buller RM, Damon IK, Lefkowitz EJ, McFadden G, McInnes CJ, et al.
Family: Poxviridae. ICTV Ninth Report. 2009. https://ictv.global/report_9th/dsDNA/poxviridae.

4.      Capobianchi MR, Di Caro A, Piubelli C, Mori A, Bisoffi Z, Castilletti C. Monkeypox 2022
outbreak in non-endemic countries: Open questions relevant for public health, nonpharmacological
intervention and literature review. Front Cell Infect Microbiol. 2022;12:1005955.

5.      Ehmann R, Brandes K, Antwerpen M, Walter M, K VS, Stegmaier E, et al. Molecular and
genomic characterization of a novel equine molluscum contagiosum-like virus. J Gen Virol. 2021
Mar;102(3).

6.      Airas N, Hautaniemi M, Syrja P, Knuuttila A, Putkuri N, Coulter L, et al. Infection with
Possible Novel Parapoxvirus in Horse, Finland, 2013. Emerg Infect Dis. 2016 Jul;22(7):1242-5.

7.      Tukia A. Scabby disease spread like wildfire has rampantly affected trotting stables [In
Finnish, Newspaper article]. Iltalehti. 2022 Mar 16. https://www.is.fi/ravit/art-
2000008686523.html.

8.      The study of the pastern infection epidemic continues — experiences needed now [In
Finnish, Newspaper article]. Hevosurheilu. 2022 Jun 23.
https://www.hevosurheilu.fi/ravit/uutiset/vuohistulehdusepidemian-selvitysty%C3%B6-jatkuu-nyt-
kaivataan-kokemuksia.

9.      Osadebe LU, Manthiram K, McCollum AM, Li Y, Emerson GL, Gallardo-Romero NF, et
al. Novel poxvirus infection in 2 patients from the United States. Clin Infect Dis. 2015 Jan
15;60(2):195-202.

10.    Gunther T, Haas L, Alawi M, Wohlsein P, Marks J, Grundhoff A, et al. Recovery of the first

full-length genome sequence of a parapoxvirus directly from a clinical sample. Sci Rep. 2017 Jun

16;7(1):3734.

11.    Yu Z, Zhang W, Fu H, Zou X, Zhao M, Liang S, et al. Genomic analysis of Poxviridae and

exploring qualified gene sequences for phylogenetics. Comput Struct Biotechnol J. 2021;19:5479-

86.

12.    Hautaniemi M, Ueda N, Tuimala J, Mercer AA, Lahdenpera J, McInnes CJ. The genome of

pseudocowpoxvirus: comparison of a reindeer isolate and a reference strain. J Gen Virol. 2010

Jun;91(Pt 6):1560-76.

138 Table: Nucleotide identity (as percentages) comparison between F14.1158H and other

139 parapoxviruses and Molluscum contagiosum virus in the four selected core genes.

| | Molluscum contagiosum virus (MC515) | Grey sealpox virus | Red deerpox virus | Bovine Papular Stomatitis Virus (AR02) | Orf virus (UPM/HSN-20) | Orf virus (NZ2) | Pseudocowpoxvirus (F00.120R) | Pseudocowpoxvirus (VR634) |
|---|---|---|---|---|---|---|---|---|
| **DNA polymerase (ORF025)** | | | | | | | | |
| Grey sealpox virus | 62 | | | | | | | |
| Red deerpox virus | 68 | 79 | | | | | | |
| Bovine Papular Stomatitis Virus (AR02) | 68 | 79 | 86 | | | | | |
| Orf virus (UPM/HSN-20) | 69 | 79 | 87 | 87 | | | | |
| Orf virus (NZ2) | 69 | 79 | 87 | 87 | 99 | | | |
| Pseudocowpoxvirus (F00.120R) | 69 | 79 | 87 | 87 | 94 | 94 | | |
| Pseudocowpoxvirus (VR634) | 70 | 80 | 88 | 88 | 95 | 95 | 98 | |
| F14.1558H | 69 | 74 | 81 | 81 | 82 | 82 | 82 | 83 |
| **Early transcription factor (ORF083)** | | | | | | | | |
| Grey sealpox virus | 65 | | | | | | | |
| Red deerpox virus | 73 | 80 | | | | | | |
| Bovine Papular Stomatitis Virus (AR02) | 74 | 80 | 90 | | | | | |
| Orf virus (UPM/HSN-20) | 73 | 79 | 88 | 89 | | | | |
| Orf virus (NZ2) | 73 | 79 | 88 | 89 | 99 | | | |
| Pseudocowpoxvirus (F00.120R) | 74 | 80 | 90 | 91 | 95 | 95 | | |
| Pseudocowpoxvirus (VR634) | 74 | 80 | 90 | 91 | 95 | 95 | 98 | |
| F14.1558H | 74 | 78 | 87 | 87 | 86 | 86 | 87 | 87 |
| **RNA polymerase (ORF101)** | | | | | | | | |
| Grey sealpox virus | 73 | | | | | | | |
| Red deerpox virus | 79 | 84 | | | | | | |
| Bovine Papular Stomatitis Virus (AR02) | 80 | 85 | 93 | | | | | |
| Orf virus (UPM/HSN-20) | 80 | 86 | 93 | 93 | | | | |
| Orf virus (NZ2) | 80 | 86 | 93 | 93 | 100 | | | |
| Pseudocowpoxvirus (F00.120R) | 80 | 86 | 92 | 94 | 97 | 97 | | |
| Pseudocowpoxvirus (VR634) | 80 | 86 | 93 | 94 | 98 | 98 | 98 | |
| F14.1558H | 80 | 84 | 89 | 91 | 91 | 91 | 91 | 91 |
| **Topoisomerase 1 (ORF062)** | | | | | | | | |
| Grey sealpox virus | 62 | | | | | | | |
| Red deerpox virus | 68 | 79 | | | | | | |
| Bovine Papular Stomatitis Virus (AR02) | 68 | 79 | 86 | | | | | |
| Orf virus (UPM/HSN-20) | 69 | 79 | 87 | 87 | | | | |
| Orf virus (NZ2) | 69 | 79 | 87 | 87 | 99 | | | |
| Pseudocowpoxvirus (F00.120R) | 69 | 79 | 87 | 87 | 94 | 94 | | |
| Pseudocowpoxvirus (VR634) | 70 | 80 | 88 | 88 | 95 | 95 | 98 | |
| F14.1558H | 69 | 74 | 81 | 81 | 82 | 82 | 82 | 83 |

A

B

C

D

**Figure.** A) Grouping of F14.1158H among all the genera of the subfamily *Chordopoxvirinae* in a

phylogenetic tree based on amino acid sequences of the DNA polymerase (ORF025) gene. B-D)

Grouping of F141158H among the genus Parapoxvirus in phylogenetic trees based on the

nucleotide sequences of the early transcription factor (ORF083) (B), RNA polymerase (ORF101)

(C), and topoisomerase 1 (ORF062) (D) genes. CRV = Crocodilepox virus, EMCLV = Equine

molluscum contagiosum-like virus, MOCV = Molluscum contagiosum virus, SQPV = Squirrelpox

virus, CPXV = Cowpox virus, MPXV = Monkeypox virus, VACV = Vaccinia virus, YMTV =

Yaba monkey tumor virus, SPPV = Sheeppox virus, MYXV = Myxoma virus, SWPV = Swinepox

virus, DPV = Deerpox virus, FWPV = Fowlpox virus, MsEPV = Melanoplus sanguinipes

entomopoxvirus. MsEPV is used as an outgroup in A and SQPV and MOCV in B-D. Bootstrap

values above 70% are shown next to the nodes.

156