TITLE & AUTHORS

HIV-1 Vpr combats the PU.1-driven antiviral response in primary human macrophages.

Maria C. Virgilio[1,2,3], Barka Ramnani[3], Thomas Chen[4], W. Miguel Disbennett[4,5,a], Jay Lubow[4,b],

Joshua D. Welch[2], Kathleen L. Collins[1,3,4,*]

[1]Cellular and Molecular Biology Program, University of Michigan, Ann Arbor, MI, USA

[2]Department of Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor,

MI, USA

[3]Department of Internal Medicine, University of Michigan, Ann Arbor, MI, USA

[4]Department of Microbiology and Immunology, University of Michigan, Ann Arbor, MI, USA

[5]Post-Baccalaureate Research Education Program (PREP), University of Michigan, Ann Arbor,

MI, USA

[a]Present location: University of Pennsylvania, Philadelphia, PA, USA

[b]Present location: Fred Hutchinson Cancer Research Center, Seattle, WA, USA

[*] Correspondence: klcollin@umich.edu

ABSTRACT

HIV-1 Vpr promotes efficient spread of HIV-1 from macrophages to T cells by transcriptionally

downmodulating restriction factors that target HIV-1 Envelope protein (Env). Here we find that

Vpr induces broad transcriptomic changes by targeting PU.1, a transcription factor necessary

for expression of host innate immune response genes, including those that target Env.

Consistent with this, we find silencing PU.1 in infected macrophages lacking Vpr rescues Env.

Vpr downmodulates PU.1 through a proteasomal degradation pathway that depends on physical

interactions with PU.1 and DCAF1, a component of the Cul4A E3 ubiquitin ligase. The capacity

for Vpr to target PU.1 is highly conserved across primate lentiviruses. In addition to impacting

infected cells, we find that Vpr suppresses expression of innate immune response genes in

uninfected bystander cells, and that virion-associated Vpr can degrade PU.1. Together, we

demonstrate Vpr counteracts PU.1 in macrophages to blunt antiviral immune responses and

promote viral spread.

KEYWORDS

Vpr, HIV-1, HIV-2, SIV, PU.1, macrophages, scRNA-seq, interferon, ISG15, TLR

INTRODUCTION

The HIV-1 genome encodes several accessory proteins that counteract innate and adaptive antiviral responses. Although HIV accessory proteins have been widely studied, the role of the Vpr accessory protein has remained enigmatic. Vpr is highly conserved amongst lentiviruses and is necessary for optimal replication in macrophages[1]. In addition, it is unique amongst HIV accessory proteins in that it is packaged in the virus particle at high levels through specific interactions with p6Gag[2]. Studies using human lymphoid tissue, which is rich in both T cells and macrophages, have found that loss of Vpr decreases virus production but only when the virus strain is capable of efficiently infecting macrophages[1,3–6]. These studies provide evidence that Vpr enhances infection of macrophages and increases viral burden in tissues containing macrophages. Vpr localizes to the nucleus[7], where it is known to induce G2 cell cycle arrest by targeting a variety of host factors involved in post-replication DNA repair[8–15]. To target these factors, Vpr requires the host protein DCAF1 (also known as Vpr binding protein (VprBP))[16]. Through DCAF1, Vpr interacts with Damaged DNA binding protein 1 (DDB1) as part of the Cul4A E3 ubiquitin ligase complex where host proteins recruited by Vpr can be ubiquitylated, then degraded[17–22]. However, the effects of Vpr on DNA repair and cell cycle arrest do not provide a clear explanation for how Vpr enhances HIV infection of macrophages.

Macrophages are terminally differentiated antigen presenting cells that are critical for many immune functions, including antiviral innate immune responses[23,24]. Macrophage identity is tightly controlled through the timed expression of myeloid transcription factors, particularly PU.1 (Reviewed in [25]). PU.1 is a hematopoietic-specific and ETS family transcription factor that is essential for lymphoid and myeloid development [25–28]. ETS family proteins bind to purine rich DNA domains with a central GGAA/T core consensus[29]. Macrophages require early and continuously high levels of PU.1 expression (Reviewed in [25,26]) to maintain normal functionality. PU.1 regulates many essential macrophage genes including those for cytokine receptors such as M-CSF and GM-CSF and the adhesion molecule CD11b. PU.1 also coordinates with other transcription factors such as IRF4 and C/EBPα, and TET methyl cytosine dioxygenase 2 (TET2; also known as ten-eleven translocation 2) to regulate gene expression[25,30–32].

We previously reported that Vpr counteracts accelerated degradation of the HIV Env protein, which occurs in HIV infected macrophages but not T cells[33,34]. We hypothesized that Vpr disables a macrophage-specific restriction factor that detects and degrades HIV Env. We

recently identified this factor as the macrophage mannose receptor (MR)[35], which is highly

expressed in macrophages but not T cells[36,37]. MR senses HIV Env via densely packed high

mannose residues that serve as pathogen-associated molecular patterns (PAMPs) because

they are normally absent from host cellular proteins. MR recognition of HIV Env disrupts

infection by promoting lysosomal degradation of Env and Env-containing viral particles[33,35].

Although the canonical targeting of host factors by Vpr involves the binding of Vpr to DCAF1,

leading to ubiquitylation and degradation via the proteosome, Vpr does not directly interact with

MR[35]. Instead, Vpr suppresses expression of the MR gene, *MRC1*[35].

In this study, we examine the mechanism by which Vpr suppresses the expression of *MRC1*

and demonstrate that Vpr additionally exerts suppressive effects on other important genes in

macrophages to enhance virion assembly and spread. Single-cell RNA sequencing (scRNA-

seq) of primary human macrophages infected with infectious wild type or Vpr-null HIV allowed

us to distinguish the effects of Vpr on both infected and virally exposed, uninfected cells in the

same culture. Within infected cells, Vpr selectively downregulated genes controlled by the

macrophage-selective transcription factor, PU.1. By targeting PU.1, Vpr systemically disrupted

several antiviral factors, including *MRC1,* and *IFITM3,* a previously reported target of Vpr that is

also capable of disrupting Env function[38]. Consistent with this, silencing PU.1 in macrophages

infected with Vpr-null virus rescued Env production. We found Vpr caused a systemic reduction

of genes implicated in Toll-like receptor (TLR) and Type I interferon (IFN-I) signaling that

affected all the cells in the culture. Thus, we provide a systems level explanation for the positive

effect of Vpr on HIV spread in macrophages. Finally, we found that Vpr-mediated

downmodulation of PU.1-regulated gene expression is mediated by protein-protein interaction

between Vpr and PU.1, resulting in accelerated proteasomal degradation of PU.1. This activity

of Vpr is conserved among all HIV-1 molecular clones tested as well as HIV-2 and SIV.

Remarkably, both the interaction between PU.1 and Vpr as well PU.1's subsequent

degradation, require the Vpr interacting protein, DCAF1. The PU.1 transcriptional co-factor,

TET2, is co-recruited with PU.1 to DCAF1 by Vpr, which aligns our results with other reports

that Vpr targets TET2[38,39]. Altogether, our data support a model in which Vpr promotes HIV

spread via systemic detrimental effects on the host innate antiviral response to infection.

RESULTS

Single-cell RNA sequencing of HIV-1 infected MDMs reveals Vpr-dependent transcriptional changes. Vpr is known to counteract a number of host factors, primarily those implicated in DNA repair and restriction of HIV Env [35,38]. Recently we demonstrated Vpr decreases transcriptional expression of the host restriction factor, mannose receptor (*MRC1*), which we hypothesize results from Vpr targeting a macrophage-specific transcription factor (Figure 1A). To identify the transcription factor targeted by Vpr, we undertook an unbiased approach to characterize the genome-wide effects HIV-1 Vpr in monocyte derived macrophages (MDMs). For these studies, we used an infectious HIV molecular clone (89.6) plus or minus an intact *vpr* gene (Figure 1B). We infected primary human CD14$^+$ MDMs from three independent healthy donors and allowed the infection to spread for 10 days before harvesting cells (uninfected, 89.6^{wt} or 89.6$^{\Delta vpr}$ infected) and prepared them for single-cell gene expression analysis (Figure 1C). LIGER (Linked Inference of Genomic Experimental Relationships) [40] was used to integrate all nine data sets. This analysis facilitated alignment despite significant transcriptomic differences between donors and sample types. Uniform Manifold Approximation and Projection (UMAP) visualization of all donor sets showed no significant donor or batch-specific differences amongst the samples (Figure 1D). Based on gene expression patterns in each of the clusters, we determined the main cluster to be pro-inflammatory macrophages (Cluster 0) and we identified a minor population of cycling cells (Cluster 1). These two clusters do not appear to have an infection phenotype because cells from cultures exposed to HIV were distributed across both the clusters (Figure 1E and F).

To separate the bona fide infected cells from bystander cells, we computationally segregated cells into a subset that expressed both *tat* and *gag* transcripts from those that expressed neither (Figure 1G, Supplementary Table 1). From a total of 13,639 WT-Vpr and 35,780 Vpr-*null* exposed MDMs, 6156 and 8699 were identified as *gag$^+$/tat$^+$*, respectively. The proportion of infected cells identified by donor using this analysis was similar to that found by the standard method of identifying infected cells by intracellular Gag staining and detection by flow cytometry (Supplementary Table 2). A comparison of gene expression profiles in cells with and without Vpr revealed that Vpr expression boosted HIV gene expression as previously reported[41] and caused a significant transcriptional shift in host gene expression (Figure 1H). Using a Wilcoxon rank-sum test and false discovery rate correction, we identified 3150 genes with statistically significant two-fold or greater differential expression between 89.6^{wt} and 89.6$^{\Delta vpr}$-infected cells.

Vpr-repressed genes in MDMs include targets of the transcription factor, PU.1.

To identify the transcription factor(s) targeted by Vpr, we used the DNA motif identification software, HOMER (Hypergeometric Optimization of Motif EnRichment), to scan for common transcription factor binding motifs in the promoters of genes downregulated in the presence of Vpr. Our analysis revealed several candidate transcription factors (TFs) and TF-binding families, including many members of the ETS family of TFs (Figure 2A; Supplementary Figure 1A). However, only PU.1 (encoded by the gene *SPI1*) was highly expressed in our data sets (Figure 2B). Moreover, PU.1 appeared twice in the list – as both a member of the ETS family and as a co-factor in the ETS/IRF (Interferon Regulatory Factor) family of transcription factors, a relationship that is well documented[30] (Figure 2A). PU.1 is the master myeloid transcription factor (Reviewed in [27,42,43]). It is required for terminal differentiation of macrophages and is necessary to maintain macrophage immune function[25]. From the list of most downregulated genes in Vpr-expressing MDMs, HOMER identified 316 genes with a PU.1 binding motif in their promoter and 670 PU.1-IRF co-regulated genes, for a combined 840 distinct PU.1-regulated genes identified (Figure 2C, E). One of the genes identified with a PU.1 binding motif was interferon induced transmembrane protein 3 (*IFITM3).* IFITM3 was previously identified as an important host restriction factor that is similar to mannose receptor (MR), in that IFITM3 targets HIV Env and has reduced gene expression (by reverse transcription quantitative PCR (RT-qPCR)) in the presence of Vpr in HIV-infected MDMs[38]. Gene ontology analysis of the 840 genes targeted by PU.1 revealed that they are enriched for regulation of the immune system and defense response (Figure 2D).

Because we had previously demonstrated that the gene encoding mannose receptor (*MRC1*) was transcriptionally downmodulated approximately two-fold by Vpr in HIV infected primary human macrophages using RT-qPCR[35], we examined our data set to specifically identify *MRC1* transcripts. In the scRNA-seq data set shown here, we identified a Vpr-dependent change in *MRC1* transcripts (log2FC = 0.932) that is statistically significant (false-discovery rate corrected two-sided Wilcoxon rank-sum p-value = 0.002) but fell just below our arbitrary cutoff of a two-fold change (Figures 1H, 2C). Consistent with the pattern we observed for other Vpr-suppressed genes, we identified several PU.1 binding sites in the promoter region of *MRC1* (Supplementary Figure 2A) when we manually scanned for PU.1 binding motifs, some of which have been previously described[44]. Furthermore, MDMs transduced with an shRNA-expressing lentivirus targeting *SPI1* transcripts, reduced both PU.1 and MR protein (Supplementary Figure 2B).

While the effect of Vpr on *MRC1* transcriptional expression is modest, it combines

synergistically with HIV Nef-dependent disruption of MR trafficking to reverse mannose

receptor-dependent lysosomal degradation of Env in HIV infected primary macrophages[35].

Vpr suppresses PU.1 regulated genes implicated in Toll-like receptor and IFN-I

responses.

PU.1 regulated genes that are downmodulated in Vpr expressing cells include several factors

implicated in Toll-like receptor (TLR) signaling pathways (Figure 3A). TLRs are evolutionarily

conserved, pattern recognition receptors (PRRs) that recognize PAMPs ([45–47]), including HIV. To

determine whether these effects of Vpr are restricted to infected cells within the culture, we

compared gene expression profiles amongst all the sample types collected from our single-cell

analysis: uninfected (virus naive), 89.6^{wt} and 89.6$^{\Delta vpr}$-infected, and 89.6^{wt} and 89.6$^{\Delta vpr}$ virally

exposed but uninfected (bystander) cells. This was accomplished by computationally

segregating cells into subsets that expressed both *tat* and *gag* transcripts from those that were

virally exposed but appeared uninfected (*tat* and *gag*-negative) (Figure 3B).

As expected, expression of genes involved in the TLR-mediated IFN-I response to infection,

including Interferon Stimulated Gene 15 (*ISG15),* which is upregulated in response to IFN-I and

TLR-signaling (Reviewed in [48,49]), *LY96, and* interferon inducible protein-6 (*IFI6*) (PU.1-IRF

regulated genes) and *IFITM3* were very low in uninfected, unexposed cells (Figure 3C). In

contrast, each of these genes was upregulated in cells from HIV-treated primary macrophage

cultures (Figure 3C). As discussed above, Vpr counteracted upregulation of this anti-viral

response in infected cells (Figure 2C, E and Figure 3C). In addition, we were surprised to find

many of these genes upregulated in bystander cells as well (Figure 3C) and that Vpr limited

their induction. For example, we found that *ISG15* was more highly expressed in 89.6$^{\Delta vpr}$

bystander cells compared to 89.6^{wt}-exposed bystander cells. Similar observations were made

for *LY96*, *IFI6*, and *IFITM3*. Interestingly, the protein product for *LY96* is myeloid differentiation 2

(MD2), which is required for TLR4 ligand-induced activation at the cell surface[50,51]. By

comparison, other subsets of genes including *SPI1* and *MRC1* were primarily downmodulated in

the infected subset and either not at all in the bystander cells or to a lesser extent than their

infected counterparts (Figure 3B).

Next, we investigated whether we could observe a Vpr-dependent suppression of PU.1

regulated gene-products involved in the TLR and IFN-I signaling pathways using confocal

fluorescent microscopy, which allowed us to distinguish between infected and uninfected cells

within the same culture of MDMs. Quantification of ISG15 and IFITM3 in Gag+ cells showed

significant reduction of both PU.1 gene products in HIV-infected MDMs expressing Vpr relative

to cells infected with a Vpr-*null* HIV (Figure 3 E, H). Strikingly, we also observed a similar

pattern in virus-exposed but uninfected MDMs (Figure 3 F, I), where uninfected cells exposed to

wild type HIV had lower levels of PU.1 gene products compared to uninfected cells exposed to

Vpr-*null* HIV. These results are consistent with the scRNA-seq data from Figure 3C, confirming

Vpr-dependent downmodulation of PU.1 regulated genes and their protein products in wild type

infected MDMs as well as bystander cells in the same culture.

PU.1 protein levels decrease in the presence of Vpr.

Vpr is known to act as an adaptor protein that links host proteins to the Cul4A E3 ubiquitin

ligase complex for ubiquitylation and degradation[17,18]. Thus, we hypothesized that Vpr

downregulates PU.1-regulated genes in infected cells by targeting PU.1 for proteasomal

degradation. To test our hypothesis, we first investigated whether Vpr reduced PU.1 protein

levels in infected MDMs. We treated macrophages with a VSV-G envelope-pseudotyped

replication-defective clone of HIV NL4-3 that expresses GFP in the *env* reading frame (Figure

4A). We transduced MDMs with this virus containing all intact accessory proteins, or with

additional mutations in the open reading frames of either *vpr, nef*, or both. PU.1 levels were

measured by flow cytometry at 5-, 7-, and 10-days post infection. Notably, we found that Vpr

expression resulted in lower PU.1 protein levels in infected (GFP$^+$) cells whether Nef was

expressed or not (Figure 4B, C). These changes in PU.1 were consistently observed in MDMs

from four independent donors (Figure 4C). Based on these results, we concluded that Vpr

downmodulates PU.1 in HIV-infected primary human MDMs. Under the conditions of this assay,

in which MDMs were treated with a replication-defective virus and analyzed at least five days

post infection, we did not observe a significant Vpr-dependent downmodulation of PU.1 in

uninfected (GFP$^-$) bystander cells (Supplementary Figure 3A).

To determine whether Vpr was sufficient for PU.1 downmodulation, we tested the ability of

several 3xFLAG-tagged Vpr proteins derived from a panel of HIV molecular clones [89.6, NL4-3,

AD8 and YU2-Vpr (Figure 4D)] to reduce exogenous PU.1 expression in transfected HEK 293T

cells. Because the gene that encodes PU.1 (*SPI1*) contains PU.1 binding sites and is

downmodulated by Vpr expression (Figure 2B), for these experiments, PU.1 was expressed

under the control of a heterologous promoter that allowed separation of transcriptional and post-

transcriptional changes (Figure 4E). For comparison, we used the evolutionarily related protein,

Vpx, from the HIV2$_{ROD}$ molecular clone. We found that all Vpr-expression constructs tested

resulted in a notably lower level of PU.1 protein compared to PU.1 protein with HIV-2$_{ROD}$Vpx

(Figure 4F). Vpr-dependent reduced levels of PU.1 were consistently observed in four

independent experiments and were statistically significant (Figure 4G).

After determining that Vpr was sufficient to downmodulate PU.1 in HEK 293T cells, we

confirmed these results in a second cell line, the lymphoblastic chronic myelogenous leukemia

cell line K562, which endogenously expresses PU.1. For these experiments Vpr-expression was

achieved using VSV-G envelope pseudotyped lentiviruses. Like the results obtained in HEK

293T cells, infections with viruses expressing all Vprs tested (89.6, NL4-3, AD8, YU2) had lower

endogenous PU.1 levels compared to mock infection and HIV-2$_{ROD}$Vpx infection (Figure 4H).

Because the Vpr-dependent decrease of PU.1 occurred both for endogenous PU.1 expressed

from its native promoter in K562 cells as well as in HEK 293T cells, when PU.1 was expressed

from a heterologous promoter, we hypothesized that Vpr was affecting PU.1 protein levels post-

transcriptionally. To rule out transcriptional effects on the heterologous promoter, we quantified

PU.1 mRNA in HEK 293T cells transfected with a PU.1 expression plasmid and HIV constructs

derived from 89.6 that had or lacked the Vpr gene (Supplementary Figure 4A). As shown in

Figure 4I, Vpr did not reduce expression of PU.1 mRNA when expressed from a heterologous

promoter in transfected cells. Thus, we concluded that the mechanisms by which Vpr reduced

PU.1 protein in HEK 293T cells must be post-transcriptional. Moreover, in macrophages, Vpr

likely exerts negative effects directly on PU.1 protein and indirectly on PU.1 RNA through PU.1

binding sites in the *SPI1* promoter [52].

Prior reports indicated that PU.1 suppresses HIV gene expression in a Tat-reversible manner[53].

If true, Vpr-targeting of PU.1 could explain the upregulation of HIV genes observed in Vpr-

expressing HIV infected macrophages (Figure 1H). To attempt to confirm these results, we co-

transfected HEK 293T cells with a PU.1 expression plasmid (Figure 4E) and an 89.6-derived

HIV genome that expresses GFP constitutively and mCherry upon HIV activation, allowing us to

distinguish transfected cells through GFP expression and HIV-LTR activity through mCherry

expression (Supplementary Figure 4A). To eliminate any complications from Vpr-mediated

reduction of PU.1, we used an HIV genome that did not express *vpr*. Transfection of increasing

amounts of PU.1 plasmid with the same amount of HIV in all conditions, resulted in increasing

amounts of PU.1 protein as measured by flow cytometry (Supplementary Figure 4B, C).

However, we failed to confirm a PU.1-dependent suppression of HIV-LTR activity

(Supplementary Figure 4D).

PU.1 downmodulation is a conserved activity of Vprs from HIV-2 and closely related SIV

molecular clones. Vpr is highly conserved in lentiviruses, including HIV-2 and all SIV strains

[54,55]. Interestingly, HIV-2 and certain SIVs contain both Vpr and Vpx, whereas HIV-1 and some

SIVs contain only Vpr (Figure 5A). The extent to which these two proteins harbor overlapping

functions has been the subject of a number of research studies [18,56,57]. Thus, we first examined

whether Vprs from viruses containing both Vpr and Vpx would reduce PU.1 levels. To do this,

we used the HIV-2$_{ROD}$ molecular clone from which our Vpx control is derived. Transient

transfection of HEK 293T cells with a PU.1 expression vector and either HIV-2$_{ROD}$Vpx, HIV-

2$_{ROD}$Vpr, or HIV-1$_{89.6}$Vpr confirmed a significant decrease in PU.1 for cells expressing Vpr^{ROD} or

Vpr$^{89.6}$ but not Vpx^{ROD} (Figure 5B, C). This indicates that Vpx and Vpr from the same molecular

clone have divergent functions with respect to PU.1 downmodulation.

We next tested the ability of Vpr from six isolates evolutionarily similar and disparate to HIV-1 to

determine whether the PU.1 targeting function of Vpr is evolutionarily conserved. Vpr

sequences from the indicated SIV molecular clones from separate clades, including two from

chimpanzee and one from gorilla (the direct evolutionary relatives of HIV-1), were indeed able to

mediate the degradation of PU.1 in HEK 293T cells (Figures 5D, E). Thus, Vpr-mediated

reduction of PU.1 was consistently observed across all HIV-1, HIV-2, and SIV molecular clones

tested, indicating a strong selective pressure for HIV-related viruses to downmodulate PU.1 in

infected cells.

Both PU.1 and DCAF1 form a complex with Vpr. To determine whether Vpr recruits PU.1

similarly to other host proteins for proteasomal degradation via the CRL4-DCAF1 ubiquitin

ligase complex (Reviewed in [54]), we first assessed whether Vpr and PU.1 formed a complex in

transfected HEK 293T cells. Remarkably, we found that PU.1 efficiently co-precipitated with

FLAG-tagged 89.6-Vpr. In addition, and as expected[16,17], DCAF1 also co-precipitated with Vpr

(Figure 6A).

We next examined whether Vpr co-precipitated with endogenous PU.1 in the K562 cell line

originally derived from a patient with CML[58]. To accomplish this, we stably expressed FLAG-

tagged HIV-1$_{89.6}$Vpr or HIV-2$_{ROD}$Vpx in K562 cells using lentiviral vectors and repeated co-

immunoprecipitation experiments. We found that both PU.1 and DCAF1 co-precipitated with Vpr

to a greater extent than Vpx, again suggesting PU.1 binding is specific to Vpr and not to the

similar accessory protein Vpx (Figure 6B). Finally, to determine if Vpr and DCAF1 interact with

PU.1 in macrophages, we expressed FLAG-tagged HIV-1$_{89.6}$Vpr in MDMs using a lentiviral

vector. As shown in Figure 6C, we indeed determined that both endogenous PU.1 and DCAF1

interact with Vpr in this physiologically relevant target of HIV-1. Collectively, these data indicate

that Vpr selectively forms complexes with both PU.1 and DCAF1 in cells endogenously

expressing PU.1.

Interactions between PU.1 and Vpr require DCAF1. To better understand the mechanism

through which Vpr downmodulates PU.1, we assessed a potential requirement for the host

protein DCAF1. DCAF1 (also known as VprBP[16]) is the Vpr binding partner in the CRL4

ubiquitin ligase complex, necessary for proteasomal degradation of Vpr-recruited host proteins

[10,11,14,17,18]. This was tested using a K562 cell line expressing a lentiviral vector containing either

a non-targeting shRNA (shScramble) or a DCAF1-targeting shRNA cassette (shDCAF1). We

found that the amount of PU.1 that co-precipitated with Vpr was decreased with DCAF1

silencing (Figure 6D), suggesting a role for DCAF1 in the interaction between Vpr and PU.1.

Consistent with this, endogenously expressed PU.1 from K562 cells did not coprecipitate with a

Vpr mutant (89.6-Vpr^{Q65R}) protein that is defective for DCAF1 interactions[59,60] (Figure 6E). These

interesting results suggest the unexpected conclusion that PU.1, Vpr, and DCAF1 may form a

trimeric complex and PU.1 binding in the complex is dependent on both Vpr and DCAF1, though

Vpr can associate with DCAF1 independently of PU.1 (Figures 6A, B).

To validate this physical interaction, we performed the immunoprecipitation (IP) in reverse using

HEK 293T cells that overexpressed FLAG-tagged-PU.1. Because these experiments utilized un-

tagged Vpr, they were limited by the availability of antibodies, which do not recognize all Vpr

allotypes equally. To overcome this limitation, we performed FLAG-tagged PU.1 pull-down

experiments with NL4-3 Vpr, which was efficiently recognized by the available antibodies

whereas 89.6 Vpr was not. As we observed in K562 cells using 89.6 FLAG-tagged Vpr, the

amount of NL4-3 Vpr that co-precipitated with FLAG-tagged PU.1 in HEK 293T cells was

decreased with DCAF1 silencing (Figure 6F). In addition, the Vpr mutant that is defective at

interactions with DCAF-1 (NL4-3-Vpr^{Q65R}) was also defective at co-precipitating PU.1 (Figure

6F), further confirming a role for DCAF1 in the interaction between Vpr and PU.1.

TET2, a PU.1 cofactor and Vpr target, coprecipitates with PU.1, Vpr, and DCAF1. The
PU.1-regulated antiviral factor, *IFITM3* (Figure 2C, E) is also controlled by TET2, another target
of Vpr [38,39]. TET2 is a DNA dioxygenase that de-methylates the *IFITM3* promoter during viral
infection, inhibiting HIV Env trafficking and reducing viral spread[61,62]. To prevent this, Vpr
mediates the ubiquitylation and degradation of TET2, which in turn inhibits *IFITM3* expression.
Interestingly, PU.1 and TET2 have been reported to form a complex to co-regulate myeloid
specific genes [25,26,32]. We therefore hypothesized that TET2 could be co-recruited with PU.1 to
DCAF1 by Vpr for degradation. Indeed, we found that TET2 co-precipitated with PU.1 in HEK
293T cells with Vpr and DCAF1 (Figure 6G).

Vpr downmodulates PU.1 via a pathway that depends on proteasome activity.
Having shown that Vpr requires DCAF1 to form a stable complex with PU.1, we next asked
whether interactions with DCAF1 were also important for promoting PU.1 degradation. As
compared to the flow cytometric approaches shown in Figures 4 and 5, detection of degradation
by western blot is more challenging. The model systems described in Figure 7 did not reliably
detect PU.1 degradation, most likely because PU.1 was overexpressed and/or Vpr was not
expressed in a sufficient number of PU.1 expressing cells. Therefore, to further study PU.1
degradation, we employed a previously reported approach in which virion-associated Vpr is
delivered to primary MDMs resulting in Vpr-dependent proteasomal degradation of targets
within five hours of viral treatment[34]. Using this approach, we observed Vpr-dependent
degradation of PU.1 treated with wild type HIV-1 but not an HIV harboring a Vpr mutant
defective at interacting with DCAF1 (89.6-Vpr^{Q65R}, Figure 7A, B). These findings support the
conclusion that interactions between PU.1, DCAF1, and Vpr are needed for efficient PU.1
degradation. Additionally, this experiment suggests that it is possible for virion-associated Vpr to
act on uninfected bystander macrophages, potentially explaining some of the results from
Figure 3 showing that Vpr can suppress innate immune response in uninfected bystander cells.

Because the Vpr/DCAF1 complex is known to promote ubiquitylation and proteasomal
degradation of Vpr-bound host proteins, we investigated whether inhibition of the proteosome
could restore PU.1 levels. As shown in Figure 7C, where we optimized PU.1 expression in HEK
293T cells expressing Vpr, PU.1 levels were restored with MG132, a specific inhibitor of the
proteasome. Furthermore, MG132 treatment of MDMs transduced with 89.6^{wt} (as for Figure 7A)
prevented virion-associated Vpr-mediated degradation of endogenous PU.1 (Figure D, E).

The results from the immunoprecipitation and degradation assays indicate a requirement for

DCAF1 for Vpr to bind PU.1 and promote its degradation, suggesting all three molecules

interact as modeled in Figure 7F. While other explanations remain possible, our data suggest

PU.1 binding in the complex is dependent on both Vpr and DCAF1. Vpr can associate with

DCAF1 independently of PU.1 and PU.1 can associate with TET2 independently of Vpr.

Reducing PU.1 enhances HIV-1 Env production in MDMs

Based on scRNA-seq data (Figure 1), Vpr reduces several antiviral factors regulated by the

transcription factor, PU.1 (Figure 2). Two of these genes, *MRC1* and *IFITM3*, have been

previously shown to inhibit the spread of HIV in MDMs by targeting HIV-1 Env[35,38]. Therefore,

we hypothesized that reduction of PU.1 in MDMs would restore Env production in Vpr-*null*-HIV-

infected primary macrophages. To test this hypothesis, we modified the 89.6 genome to remove

vpr and replace it with either an shScramble sequence, or one of three different shRNAs

targeting the PU.1 gene, *SPI1* (Figure 8A). All three PU.1-targeting cassettes were

independently validated in K562 cells to confirm their ability to reduce endogenous PU.1

expression (Figure 8B). Consistent with our hypothesis, we found that MDMs from two

independent donors infected with all three shSPI1-containing viruses showed a marked

increase in Env compared to MDMs infected with virus expressing shScramble (Figure 8C).

Overall, our findings support a model in which Vpr dramatically alters the transcriptional

landscape in macrophages by targeting myeloid-specific transcription factors that are required

for the expression of key antiviral restriction factors, including those that target HIV-1 Env.

(Figure 8D).

DISCUSSION

Although HIV accessory proteins have been widely studied, the critical function that drives

evolutionary conservation of Vpr has remained largely enigmatic. While Vpr significantly

enhances infection of macrophage-containing cultures, primarily by enhancing cell-to-cell

transmission, it does not substantially affect HIV infection in cultures of CD4[+] T cells that lack

macrophages[1,3–5,33–35]. While a number of Vpr targets implicated in post-replication DNA repair

have been identified, a compelling explanation for the dramatic selective effects of Vpr on

infection and spread in macrophage-containing cultures has remained elusive. In this work, we

identified a macrophage-specific target of Vpr, the myeloid transcription factor PU.1 and its

associated co-factors, which Vpr targets, averting antiviral effects (Figure 8D). Remarkably, by

utilizing single-cell RNA sequencing of MDMs treated with replication competent virus with and

without Vpr we could distinguish effects of Vpr on cells harboring bona fide infections as well as

bystander cells.

In infected cells, we found that Vpr reduces the transcription of hundreds of genes regulated

through PU.1 and its cofactors. Several PU.1 regulated genes we identified as being impacted

by Vpr are involved in TLR signaling. TLRs are highly conserved PRRs that help cells identify

PAMPs and respond quickly to infection. Activation of TLRs through binding of a PAMP, initiates

a cascade of intracellular signaling that results in the release of inflammatory cytokines and

upregulation of Type I interferon response genes[47,49,63]. The products of these genes have

known antiviral effects on HIV, and include *ISG15*, *STMN1, IFI6, LY96*, *TREM2*, *FABP4*,

IFITM3, and *MRC1*. *STMN1* is thought to play a role in the establishment of HIV latency, and

depletion leads to higher expression of HIV-1[64]. *MRC1* and *IFITM3* interrupt Env trafficking,

reducing viral spread. *IFI6*, *LY96, and ISG15* are members of the Type I IFN response to

infection[48,65,66] and *ISG15* has also been shown to inhibit HIV spread by disrupting Gag

polymerization[67].

Unexpectedly, we found that bystander cells within HIV infected cultures also responded to HIV

infection by upregulating a subset of these antiviral genes, including *ISG15*, *LY96*, *IFI6*, and

IFITM3. Moreover, we found that upregulation of these factors was reduced in bystander cells

from Vpr-positive versus Vpr-negative HIV-infected primary macrophage cultures. Because Vpr

is efficiently packaged into the virus particle through specific interactions with p6Gag[2], the Vpr

phenotype we observed in bystander cells could be due to a low and/or transient presence of

Vpr in MDMs that have taken up viral particles, but remained uninfected. This could occur if a

subset of Vpr-containing viral particles were defective and/or if innate immune responses

blocked completion of reverse transcription and/or integration. We did not consistently detect

effects of Vpr in uninfected bystander cells exposed to VSV-G pseudotyped replication defective

virus five days post-infection, but this is not surprising given the likely turnover of the viral

protein and the lack of continuous exposure to virions by MDMs treated at a single time point

with replication defective viruses. In contrast, we did find that MDMs exposed to wild type HIV-1

had low PU.1 levels at short time points (five hours) following exposure to virus. Thus, the Vpr-

dependent bystander phenotype we identified in MDMs continuously exposed to wild type

spreading HIV is most likely due to Vpr delivered by wild type viral particles. However alternative

and additional explanations remain possible. These include the possibility that there is

differential antiviral cytokine production by wild type versus Vpr-mutant infected macrophages

that acts on bystander cells. Regardless of the precise mechanism, our results indicate that Vpr

can exert systemic antiviral affects that favor virus infection and spread.

We have previously shown that mannose receptor is a host restriction factor that reduces HIV

spread in macrophages by binding to mannose residues on Env and directing Env to the

lysosome for degradation[35]. However, our previous data lacked a mechanism for Vpr-mediated

transcriptional reduction of *MRC1* in macrophages. The data here confirm findings from other

groups that PU.1 regulates *MRC1* expression[44,53], and we provide new evidence that Vpr

reduces the transcriptional expression of *MRC1* in macrophages via PU.1 degradation.

In agreement with other studies[38,39], our unbiased survey of Vpr's effect on expression of the

host transcriptome showed that Vpr reduced *IFITM3* gene expression substantially. IFITM

proteins are broad antiviral factors that inhibit viral entry and exit for HIV-1, SIV, MLV, VSV,

EBOV, WNV, among other viruses [68]. Of the IFITM proteins, IFITM3 is well documented as an

HIV-1 antagonist. Like mannose receptor, IFITM3 interacts with Env in infected cells to inhibit

Env processing and virion incorporation and strongly inhibits cell-to-cell spread. Interruption of

Env processing by IFITM3 has been demonstrated for several HIV-1 molecular clones[69] many of

which were included within our own study (AD8, YU2, and NL4-3, HIV-2$_{ROD}$, SIVagm, and

SIVcpz). *IFITM3* expression is regulated by the DNA dioxygenase, TET2[38]. During HIV-1

infection of macrophages, in the absence of Vpr, TET2 demethylates the *IFITM3* promoter,

relieving suppression, contributing to the anti-viral response. When Vpr is present, *IFITM3*

expression is reduced after Vpr recruitment of TET2 to DCAF1 for polyubiquitylation and

degradation of TET2[39]. TET2 is ubiquitously expressed in the nuclei of all cells[70], thus it was

unclear how Vpr targeting of TET2 could result in a macrophage-specific Env phenotype.

Interestingly, in monocytes, TET2 interacts with PU.1, leading to interaction with genetic

targets[32]. Therefore, we hypothesized that Vpr exerts a macrophage-specific effect on the

antiviral response by targeting PU.1 and by extension, PU.1-associated proteins. Consistent

with this hypothesis, we were able to demonstrate TET2 immunoprecipitating with PU.1, Vpr,

and DCAF1, however confirmation of this interaction in MDMs has not yet been achieved.

Altogether, this provides evidence that Vpr can re-program cellular transcription in macrophages

by targeting myeloid-specific transcription factors.

Vpr-mediated reduction of PU.1 occurred in all cell types tested in our study including MDM,

K562, and HEK 293T cells. The Vpr-dependent reduced PU.1 levels were observed regardless

of whether PU.1 was expressed from its native promoter or a heterologous promoter. Reversal

of PU.1 downmodulation with proteasome inhibitors in both HEK 293T cells and MDMs support

the conclusion that Vpr directly reduces PU.1 protein by promoting its degradation. Vpr-

mediated degradation of PU.1 was consistent for all HIV-2, SIV, and HIV-1 Group M isolates

tested - Group M being largely responsible for the global HIV pandemic. In macrophages, *SPI1*

mRNA levels encoding PU.1 protein were also lower in the presence of Vpr, indicating that

downmodulation of PU.1 can occur at both the transcriptional and the post-transcriptional level

in macrophages.

It is important to note that we failed to confirm prior observations that PU.1 suppresses HIV-LTR

activity in a Tat reversible manner[53]. While it is possible that expression of Tat by our construct

prevented our ability to detect an effect of PU.1 on the HIV-1 LTR, these results nevertheless

indicate that downmodulation of PU.1 by Vpr is unlikely to impact HIV-1 gene expression in

infected macrophages that also express Tat. Thus, in primary HIV infected macrophages, Vpr

mainly targets PU.1 to counteract its anti-viral defense response rather than to counteract an

inhibitory effect on HIV gene expression. However, it is possible that the higher expression of

HIV genes we observed in Vpr-containing cells may result from Vpr counteracting a PU.1-

regulated factor that inhibits HIV gene transcription.

Vpr is highly conserved amongst primate lentiviruses and promotes infection of nondividing

cells, especially macrophages[34]. In addition, a requirement for Vpr to achieve maximal

replication and persistence in vivo was first discovered using an SIV molecular clone in rhesus

monkeys[71]. We therefore speculated that the ability of Vpr to degrade PU.1 is an important

evolutionary function of Vpr. HIV-2 differs from HIV-1 in that it contains Vpx, an accessory

protein that shares a common genetic ancestor with Vpr[55] but is lacking from all HIV-1

genomes. Thus, we tested the relative ability of Vpx and Vpr from the same molecular clone to

promote PU.1 degradation[55]. We were able to identify a unique function of HIV-2 Vpr to degrade

PU.1 that was not shared by Vpx. Similarly, both SIVcpz and SIVgor, the evolutionary

precursors to HIV-1, and Vpr proteins from more evolutionarily distant viruses were able to

consistently result in lower PU.1 levels. However, the greatest decrease in PU.1 levels were

achieved with HIV-1 derived molecular clones. We therefore speculate that the strength of Vpr-

mediated degradation of PU.1 plays an important role in driving spread in human pandemic

strains of HIV-1, and PU.1 may be a critical restriction factor, limiting spread between species

and within populations. However, more extensive studies comparing the relationship between

Vpr and PU.1 across SIV, HIV-2, and HIV-1 isolates is necessary and part of our ongoing

efforts.

The relationship between DCAF1 and Vpr is well documented[16–18]. We confirmed the binding of

Vpr to DCAF1 and identified PU.1 as a new Vpr-binding factor. We demonstrated that all three

components co-precipitate under many different cellular conditions regardless of whether we

pull-down with Vpr or PU.1. To our surprise, mutation of the glutamine residue at position 65 of

Vpr (Vpr^{Q65R}) not only resulted in the loss of DCAF1 association but also disrupted formation of

the Vpr-PU.1 complex. Additionally, when we reduced DCAF1 or used Vpr^{Q65R}, the amount of

both Vpr and DCAF1 precipitating with PU.1 was reduced. These results indicate that

interactions amongst all three proteins are necessary for stable complex formation, although

further studies are necessary to understand the detailed protein-protein interactions. Based on

these findings, we propose a model in which interactions amongst PU.1, Vpr, and DCAF1

promote the ubiquitylation of PU.1 via the associated CUL4A ubiquitin ligase complex with

resultant proteasomal degradation. Consistent with this model, PU.1 was not degraded in the

presence of proteasome inhibitors or by a Vpr mutant defective at interacting with DCAF1 in

primary MDMs. Thus, our working model is that PU.1 is poly-ubiquitylated following its

interaction with Vpr and DCAF1. However complete confirmation of this model has not yet been

achieved because we have not yet directly detected ubiquitylated intermediates of PU.1 in Vpr-

expressing cells.

Although the role of Vpr in HIV infection has remained largely undefined, we provide evidence

that the primary selective pressure for Vpr in lentiviruses is to disrupt the macrophage innate

antiviral response to infection, which is achieved by reducing PU.1 levels in infected cells. The

ability of Vpr to degrade PU.1 is highly conserved among all HIV-1, HIV-2, and SIV isolates

tested, and degradation of PU.1 relies on both Vpr and DCAF1. Reducing PU.1 in HIV-infected

macrophages lacking Vpr rescued macrophage-dependent restriction of HIV-1 Env, helping to

explain the requirement for Vpr in macrophage spreading infections. We are continuing to

investigate the transcriptional consequences of PU.1 degradation in macrophages. In addition to

TET2, PU.1 associates with other transcription factors in cells, potentially piggybacking other

secondary targets to the DCAF1-Cul4A E3 Ubiquitin ligase complex via Vpr. Because PU.1 is

necessary to maintain macrophage function, future studies should address the greater impact of
Vpr-mediated reduction of PU.1 on the infected macrophage.
540

ACKNOWLEDGMENTS

This work was supported by National Institutes of Health grants R01AI149669 and
1R61DA059916-01 to K.L.C. and J.D.W., F31AI15504 to M.C.V. and F31AI125090-01 to J.L.,
R21AI32379 to K.L.C, training grants T32GM007315 to M.C.V. and 5T32GM008353-27 to J.L.,
and additional funding provided by the Rackham Regents Fellowship to M.C.V. We thank Chen
Li from the Welch lab for his invaluable advice and aid with single-cell data analysis. We thank
the University of Michigan Flow Cytometry Core and Advanced Genomics Core for assistance
with experiments, particularly Wenpu Trim and Tricia Tamsen. Thank you to all members of the
Collins lab for their helpful discussions.

AUTHOR CONTRIBUTIONS

552 Conceptualization, M.C.V, J.L., and K.L.C.; Methodology, M.C.V, J.L, J.D.W., and K.L.C.;
553 Software, M.C.V. and J.D.W.; Validation, M.C.V.; Formal Analysis, M.C.V.; Investigation,
554 M.C.V., B.R., T.C., W.M.D., and J.L.; Resources, K.L.C.; Writing – Original Draft, M.C.V.;
555 Writing – Review & Editing, M.C.V., J.D.W., and K.L.C.; Visualization, M.C.V.; Supervision,
556 J.D.W. and K.L.C., Funding Acquisition, M.C.V., J.D.W., and K.L.C.
557

DECLARATION OF INTEREST

The authors declare no competing interest.

METHODS

Cell culture and preparation of human MDMs

All cell cultures were maintained at 37°C in a 5% CO_2 humidified atmosphere. HEK 293Ts

(CRL-3216) and K562 (CCL-243) cells were obtained from ATCC and independently

authenticated by STR profiling. HEK 293Ts were maintained in DMEM medium (Gibco)

supplemented with 100 U/mL penicillin, 100μg/mL streptomycin, 2mM glutamine (Pen-Strep-

Glutamine, Invitrogen), and 10% fetal bovine serum (Invitrogen). K562 cells were maintained in

IMDM (Gibco) and supplemented as HEK293Ts. Monocyte-derived macrophages (MDMs) were

isolated and cultured as previously described[35] and maintained in R10 (RPMI, supplemented as

HEK 293Ts). Monocytes were isolated from anonymous donors by apheresis and obtained from

the New York Blood Center Component Laboratory. The use of human blood from anonymous,

de-identified donors was classified as non-human subject research in accordance with federal

regulations and therefore not subject to formal IRB review.

Viruses, viral vectors, and expression plasmids

The molecular clone p89.6 (cat# 3552 from Dr. Ronald G. Collman) was obtained from the AIDS

Reagent Program. The *vpr*-null version was generated as previously described[34]. The human,

full length PU.1 (hPU.1) expression vector was a gift of Dr. Gregory M. K. Poon and generated

as previously described[72]. A triple N-FLAG-tagged version of PU.1 was generated by using PCR

amplification of hPU.1 to add a KpnI site to the 5' end and EcoRI to the 3' end (5'

GTAGGTACCGCCACCATGGAAGGGTT primer and 3'

GTAGAATTCCACCACACTGGACTAGTG primer). The new product replaced an existing gene

when inserted into pcDNA3.1 containing a triple N-FLAG-tag between KpnI and EcoRI

(Addgene plasmid #67788). The GFP transfection control plasmid, pcDNA3-EGFP, was a gift

from Doug Golenbock (Addgene plasmid # 13031; http://n2t.net/addgene:13031; RRID:

Addgene_13031). The pNL4-3 ΔGPE-GFP plasmid used for single round infection of

macrophages was previously described[73], as were the *vpr*-null, *nef*-null, and *vpr-nef* null

versions[35]. pUC19[74] was used as control DNA to adjust transfection samples to the same final

DNA concentration. The p89.6-ΔGPEN-mCherry-pSFFV-EGFP single-round infection plasmid

was previously described[75] and further modified by replacing *gag* and *pol* with mCherry.

Deletion of *vpr* was achieved using Q5 Site-Directed Mutagenesis kit (New England Biolabs)

where the majority of the *vpr* coding sequence was deleted using PCR exclusion (forward -

CAGAATTGGGTGTCGACATAG, reverse - TCACAGCTTCATTCTTAAGC).

Triple FLAG-tagged Vpr and Vpx lentiviral expression vectors for HIV-2$_{ROD}$Vpx (Addgene

plasmid #115816), SIV$_{SAB-92018}$Vpr (Addgene plasmid #115822), SIV$_{AGM-MAL}$Vpr (Addgene

plasmid #115828), SIV$_{CPZ-TAN3}$Vpr (Addgene plasmid #115833), SIV$_{CPZ-LB7}$Vpr (Addgene plasmid

#115834), SIV$_{gorCP684con}$Vpr (Addgene plasmid #115835), and SIV$_{rcm02CM8081}$Vpr (Addgene

plasmid #115838) were a gift from Jeremy Luban[57]. Similar vectors for HIV-2$_{ROD}$Vpr, HIV-

1$_{89.6}$Vpr, HIV-1$_{89.6}$Vpr^{Q65R}, HIV-1$_{NL4-3}$Vpr, HIV-1$_{AD8}$Vpr, HIV-1$_{YU2}$Vpr were generated by

synthesizing the gene as a gBlock (IDT, Coralville, Iowa, USA) between NotI and either EcoRI

or AflIII in the same lentiviral expression plasmid. Untagged expression vectors for HIV-1$_{NL4-3}$Vpr

and HIV-1$_{NL4-3}$Vpr^{Q65R} were generated by synthesizing the genes as a gBlocks (IDT, Coralville,

Iowa, USA) and inserting them between SbfI and NotI in LeGO-IV, a gift from Boris Fehse

(Addgene plasmid #27360)[76].

The short hairpin RNAs targeting *DCAF1* (target sequence: CCTCCCATTCTTCTGCCTTTA)

and *SPI1* (target sequence 1: GCCCTATGACACGGATCTATA, target sequence 2:

CGGATCTATACCAACGCCAAA, and target sequence 3: CCGTATGTAAATCAGATCTCC) were

designed using Genetic Perturbation Platform (Broad institute) and cloned into pLKO.1 – TRC

cloning vector, a gift from David Root (Addgene plasmid # 10878 ; http://n2t.net/addgene:10878

; RRID:Addgene_10878)[77]. The control shRNA, scramble shRNA was a gift from David Sabatini

(Addgene plasmid # 1864 ; http://n2t.net/addgene:1864 ; RRID:Addgene_1864)[78]. For shRNA

expression from full-length HIV-1-89.6 virus, the *vpr*-ORF was first disrupted by the insertion of

a U6-promoter followed by multiple unique restriction enzyme sequences generated by

synthesizing the segment as a gBlock (IDT, Coralville, Iowa, USA). The segment was inserted

between XcmI and SalI without disrupting the *vif* or *tat* ORFs. The same shRNAs as above were

then cloned into HIV-1-89.6 after the U6-promoter.

Co-transfections

Co-transfections of 3xFLAG-Vpx/Vpr and hPU.1 were performed in HEK 293T cells. Cells were

plated at 1.6 x 10^5 per well in a 12-well dish. 24hrs after plating, 1ng of hPU.1, 500ng of

3xFLAG-Vpx/Vpr, and pUC19 to a total of 1μg of DNA per well were combined with 4μg of PEI,

mixed and added to each well. 48hrs later, cells were harvested for flow cytometry or

immunoblotting. For co-immunoprecipitations, transfection experiments were scaled to achieve

60 x 10^6 cells per condition. Co-transfections with p89.6-ΔGPERN-mCherry-pSFFV-EGFP and

hPU.1 were performed as described above and with DNA amounts described in the legend.

Transduction of MDM, K562, and HEK 293T

All transductions were performed via spin inoculation at 1050 x g for 2 hr at 25°C with equal

virus amounts determined by Gag p24 mass in medium containing 4µg/mL polybrene (Sigma).

MDMs were inoculated with 10µg p24 mass equivalents of NL4-3 ∆GPE-GFP virus or 20µg p24

mass equivalents of 3xFLAG-89.6-Vpr. K562 and HEK 293Ts were spin inoculated with 10µg

p24 mass equivalents of shScramble and shDCAF1 viruses. K562 cells were inoculated with

varying amounts of 3xFLAG-tagged Vpr/Vpx expression viruses to achieve equal FLAG

expression. After infection, viral medium was removed and replaced with fresh medium.

HIV infection of MDM

Prior to infection, half of the medium was removed from each well of MDMs and saved to make

diluted conditioned media post-infection. MDM were infected with 5µg, 10µg, and 20µg (scRNA-

seq) or 20µg and 50µg (immunofluorescence) equivalents of Gag p24 mass diluted in R10 for 6

hr at 37°C. After the 6 hr infection, media was removed and replaced with conditioned media

diluted 1:2 in R10. Half-media changes were performed every four days. For assessment of

virion associated impact on PU.1 (including MG132 treatment), MDMs were infected with 200-

300µg of virus in R10 of either 89.6^{wt}, 89.6$^{\Delta vpr}$, or 89.6$^{\Delta vpr-Q65R}$ for 5hrs.

Single-cell RNA sequencing

At 10 days post-infection, uninfected, 89.6^{WT}, and 89.6$^{\Delta vpr}$ infected MDMs were lifted using

enzyme free cell dissociation buffer (ThermoFisher). Replicate samples were fixed with

paraformaldehyde and stained for Gag to assess viral spread by flow cytometry. The resulting

flow cytometry data was used to select 89.6^{WT} and 89.6$^{\Delta vpr}$ conditions with similar percentages

of infected cells. Wells of the selected conditions were harvested, counted, and prepared

according to manufacturer instructions for 10X Chromium Next GEM Single Cell 3' v3 Gene

Expression (10X Genomics).

Single-cell data analysis

10X filtered expression matrices were generated from CellRanger version 3.0.0 (10X

Genomics). We analyzed all single-cell gene expression data using the standard LIGER[40]

(https://github.com/welch-lab/liger) data integration pipeline. All WT and Vpr-null infected MDM

raw data expression matrices from each donor were combined before merging the data. We

used a value of k = 20 during joint matrix factorization, of resolution of 0.05 for Louvain

clustering, and nearest neighbor = 30 with a minimum distance = 0.3 for UMAP visualization.

Infected cells were identified by sub-setting cells with a non-zero expression value for both *gag*

and *tat* transcripts. Differential gene expression between WT and Vpr-null infected MDMs was

determined using the Wilcoxin rank-sum test. Volcano plots of differentially expressed genes

were generated using ggplot2[79] (https://ggplot2.tidyverse.org/). Downregulated genes in the

presence of Vpr with a log2FC > 1 and adjusted p-value > 0.05 were used as input for the

HOMER[80] (http://homer.ucsd.edu/homer/motif/) 'findMotifs' function using the human reference

set. PU.1-motif associated genes were identified using the 'find' function in 'findMotifs' from

HOMER. Gene Ontology analysis for biological processes for PU.1 motif-containing genes was

determined using GOrilla[81,82] (http://cbl-gorilla.cs.technion.ac.il/) with PU.1 regulated genes as

target genes and all expressed genes in our dataset as background. Biological processes were

plotted using REVIGO[83] (http://revigo.irb.hr/). The -500bp sequence for MRC1 used for PU.1

motif scanning was obtained from UCSC Genome Browser[84] (https://genome.ucsc.edu/) using

Human reference genome GRCh38/hg38. The PU.1 binding motif probability matrix was

obtained from HOMER and used with FIMO[85] (https://meme-suite.org/meme/doc/fimo.html) to

scan the *MRC1* input sequence. Violin plots were generated by importing our LIGER generated

dataset into Seurat[86] (https://satijalab.org/seurat/) and running the VlnPlot function. All single-

cell data analysis and plots were done using RStudio[87] (http://www.rstudio.com/) except for

HOMER-predicted motifs.

Virus production

Virus stocks were produced by transfected HEK 293T cells (ATCC, Manassas, Virginia) with

viral DNA and polyethylenimine (PEI) as previously described[35]. For replication defective

constructs, cells were plated 24hrs before transfection with a DNA ratio of 1:1:1 with pCMV-HIV-

1[88], pHCMV-V (VSV-G expression plasmid) (ATCC 75497), and lentiviral expression plasmid.

Viral supernatant was collected two days post-transfection and stored at -80°C. For infectious

virus, pCMV-HIV and pHCMV-V were omitted.

Virion quantification

Viral supernatants were quantified by ELISA as previously described[35] with the following

modification: lysed samples were captured at 4 □C overnight.

Immunoblot

Western blots: cells were lysed in Blue Loading Buffer (cat# 7722, Cell Signaling Technology),

sonicated with a Misonix sonicator (Qsonica, LLC. Newtown, CT), boiled for 10min at 95°C

before loading, and analyzed by SDS-PAGE immunoblot.

Co-immunoprecipitation: cells were lysed in Pierce IP Lysis Buffer (Thermo Fisher Scientific)

and 1x Halt Protease Inhibitor Cocktail (Thermo Fisher Scientific). Lysates were incubated with

Anti-FLAG M2 Magnetic Beads (MilliporeSigma, Darmstadt, Germany) according to

manufacturer instructions. Proteins were eluted using 3xFLAG peptide (MilliporeSigma,

Darmstadt, Germany) and analyzed by SDS-PAGE immunoblot. FLAG-tagged proteins were

visualized using Pierce ECL (Thermo Scientific) after treatment with an HRP-conjugated primary

antibody directed against the FLAG epitope (Millipore Sigma). HRP-conjugated secondary

antibodies against murine and rabbit antibodies to other targets (see below) plus ECL Prime

reagent (Cytiva Amersham) were used to visualize all other proteins.

Antibodies

Antibodies to Vinculin (cat# V9131, Millipore Sigma), DCAF1 (cat# 11612-1AP, Proteintech),

PU.1 (cat# 2266S, Cell Signalling Technology), FLAG (cat# F1804, Millipore Sigma), TET2

(cat#MABE462, EMD Millipore), Vpr (AIDS Reagent Program cat# 3951 from Dr. Jeffrey Kopp),

pr55 and p24 (AIDS Reagent Program cat# 3957), gp120 (AIDS Reagent Program cat# 288),

GFP (cat# ab13970, Abcam) were used for immunoblot analysis. Secondary HRP conjugated

antibodies against murine (rat anti-mouse IgG1, eBioscience), rabbit (goat anti-rabbit IgG,

Invitrogen), sheep (rabbit anti-sheep IgG, Dako), and human (goat anti-human IgG,

ThermoFisher) were also used. Antibodies to PU.1 (clone 7C6B05, BioLegend), FLAG (cat#

637324, BioLegend) were used for flow cytometry. Antibodies to ISG15 (cat# 15981-1-AP,

Proteintech), IFITM3 (cat# 11714-1-AP, Proteintech), and AlexaFluor 647 (A21244, Fisher

Scientific) secondary antibody were used for immunofluorescence. CAp24 (clone KC57-PE cat#

6604667, Beckman Coulter) was used for both flow cytometry and immunofluorescence.

Immunofluorescence

MDMs were generated as described above in μ-slide glass-bottomed cell chambers (Ibidi,

Gräfelfing Germany) and infected as described above. Cells were fixed by adding 4%

paraformaldehyde (PFA) and permeabilized by adding 0.1% Triton X in PBS. Cells were then

blocked by incubating with 5% goat serum (Millipore Sigma) and 1% bovine serum albumin

(BSA) in PBS for 30mins at room temperature. Primary antibodies against ISG15 (Proteintech)

or IFITM3 (Proteintech) were diluted 1:450 or 1:400 respectively in 1% BSA in PBS and were

incubated with cells for 90mins at room temperature. Goat anti-rabbit AlexaFluor 647 secondary

antibody (Fisher Scientific) was diluted 1:200 in 1% BSA and incubated with the cells for 30mins

at room temperature, protected from light. Cells were incubated with Anti-PE conjugated Gag

(1:400) antibody for 30mins at room temperature. Cells were washed three times with PBS after

each step. Nuclei were stained by diluting a 1 mg/mL stock of DAPI (ThermoFisher Scientific)

1:1,000 in PBS and incubating with the cells for 5 minutes at room temperature. Cells were

imaged with a Nikon N-SIM + A1R confocal microscope. Identical laser and gain settings were

used across all images for each individual replicate of the experiment. Images were processed

using NIS viewer imaging software and corrected total cell fluorescence (CTCF) was calculated

using image J software. CTCF = Integrated Density – (Area of Selected Cell x Mean

Fluorescence of Background readings).

Quantitative RT PCR

HEK 293T cells sorted as described in 'Flow Cytometry' below were counted using the

Countess II Cell Counter (Thermo Fisher Scientific) and cell samples were diluted such that all

conditions contained the same cell numbers as input. RNA was isolated using the Zymo

DirectZol RNA MiniPrep Plus extraction kit with an on-column DNaseI digestion. RNA was

reverse transcribed using the High-Capacity cDNA Reverse Transcriptase kit (Applied

Biosystems). Quantitative PCR was performed using SYBR green qPCR Master Mix (Applied

Biosystems) on a QuantStudio 3 Real-Time PCR System (Applied Biosystems) with

ReadyMade PrimeTime primers for *SPI1* (Integrated DNA Technologies Inc, USA) and RT2

qPCR Primer Assay for Human GAPDH (cat# PPH00150F-200, Qiagen). Expression was

quantified using ABI Sequence Detection software compared to serial dilutions of an SPI1 or

GAPDH synthetic sequence gBlock (Integrated DNA Technologies Inc, USA). Measured values

for *SPI1* were normalized to measured values of *GAPDH*.

Flow cytometry

For cells requiring intracellular staining using antibodies directed against HIV Gag p24, FLAG-

Vpx and -Vpr, and PU.1, paraformaldehyde-fixed cells were permeabilized with 0.1% Triton-X in

PBS for 2min followed by incubation with antibody for 30min at room temperature. In all

experiments, cells were gated sequentially by forward scatter vs. side scatter for cells and then

by forward scatter area vs. height to exclude doublets. The gating strategy is shown in

Supplementary Figure 3B. All transduced MDMs and transiently transfected HEK 293T cells

were assessed for protein expression on the Cytek Aurora. GFP+ HEK 293T cells in the MG132

treatment experiments and for qPCR were sorted on the Sony SH800 cell sorter into R10.

Untreated, GFP- cells were also sorted. All flow cytometry data was analyzed using FlowJo

software (TreeStar, Ashland, Oregon).

Proteasome inhibition

Lyophilized MG132 was purchased from MilliporeSigma (cat# M8699) and dissolved in DMSO.

For HEK 293Ts, MG132 was added to cellular medium to achieve a final concentration of 10µM

at varying timepoints. Cells were harvested from replicate wells such that all drug treatment

timepoints were collected at once. DMSO only control treatment wells (Vehicle) were treated

with the same volume of DMSO as contained in the MG132 treatment conditions. For MDMs,

cells were pre-treated with 2.5µM MG132 for 2hrs prior to infection, then maintained in 2.5µM

MG132 throughout the infection.

Statistical Analysis

All non-single cell statistical analyses were performed using GraphPad Prism software as

described in figure legends for each experiment.

Data Availability

CellRanger version 3.0.0 processed data generated in the manuscript have been deposited in

GEO (GSE220574). Raw sequencing data are deposited in NCBI dbGAP: phs002915.v2.p1. TF

motif data and analysis from HOMER[80] (http://homer.ucsd.edu/homer/) are described above.

Code Availability

We analyzed all single-cell gene expression data using the standard LIGER[40]

(https://github.com/welch-lab/liger) data integration pipeline. TF binding motif data was

generated using the HOMER[80] (http://homer.ucsd.edu/homer/motif/) 'findMotifs' function using

the human reference set. PU.1-motif associated genes were identified using the 'find' function in

'findMotifs' from HOMER.

 FIGURE LEGENDS

Figure 1. Single-cell RNA sequencing of HIV-1 infected MDMs reveals Vpr-dependent

transcriptional changes. (A) Schematic diagram illustrating the objective of single-cell RNA

sequencing; the identification of Vpr-targeted transcription factor(s) in HIV infected primary

macrophages. (B) Genome maps for full-length 89.6^{wt} HIV-1 (top) and the same viral genome

with a premature stop-codon in *vpr* (bottom). (C) Experimental setup for the generation of the

scRNA-seq datasets. (D) UMAP representation of LIGER-integrated scRNA-seq data from

MDM samples treated as shown in (C) and listed in Dataset. (E, F, G) UMAP representations of

LIGER-integrated scRNA-seq data from MDMs treated as indicated. (E) Colors indicate

individual clusters. Cluster 0 = pro-inflammatory macrophages, Cluster 1 = cycling cells. (F, G)

cells colored according to whether they were exposed to 89.6 WT-Vpr virus (blue) or Vpr-null

virus (pink). (G) Bona fide infected cells were identified based on expression of HIV *tat* and *gag.*

(H) Volcano plot of differentially expressed genes from HIV 89.6^{wt} verses 89.6$^{\Delta vpr}$ infected MDMs

from (G) as determined by Wilcoxin Rank Sum. Significance determined as greater than 1 log2

fold change and false discovery rate adjusted p-value of $p < 0.05$ (red-bars). Blue colored genes

indicate genes less highly expressed in HIV 89.6^{wt} verses 89.6$^{\Delta vpr}$ infected MDMs, red colored

genes indicate genes more highly expressed, and black colored genes indicate no significant

difference between datasets.

Figure 2. Vpr downmodulates PU.1-dependent transcription. (A) PU.1 motifs identified by

HOMER as present in the promoters of Vpr-downmodulated genes (Figure 1H, Blue). (B) Violin

plots displaying RNA abundance of the indicated transcription factor genes in MDMs infected

with the indicated virus. (C) Volcano plot as in Figure 1H except that genes containing a PU.1 or

PU.1-IRF binding motif in their promoter region are highlighted in green (See also

Supplementary Figure 2). (D) Biological processes associated with the PU.1 targeted genes

from (C). Size of circle indicates the relative number of GO terms associated with the process.

FDR adjusted q-values associated with GO terms as indicated by the color. Bolded rings are

associated with biological processes listed. (E) Violin plots displaying RNA abundance of the

indicated genes in MDMs infected with the indicated virus. HIV-1 89.6^{wt} (WT); HIV-1 89.6$^{\Delta vpr}$

(ΔVpr). False-discovery rate corrected Wilcoxon rank-sum p-values are shown above the

conditions being compared.

Figure 3. Vpr counteracts the innate immune response to HIV infection. (A) Selected

biological processes associated with infection and inflammation (selected from GO terms

represented in Figure 2D). Pathways were identified as associated with PU.1 regulated genes

downmodulated in MDMs infected with 89.6^{wt} or 89.6$^{\Delta vpr}$ virus as determined by expression of

gag and *tat* transcripts (highlighted in Figure 2C). The -log10 FDR-adjusted p-values (q-values)

are plotted for each gene ontology term. Blue bars represent terms associated with TLR

signaling; Black bars represent related gene ontology terms similar, but not directly associated

with TLR signaling. (B) Violin plots summarizing single-cell RNA transcripts expressed by

primary human macrophages treated with the indicated virus and cultured for 10 days.

Gag+/tat+ are the subset of cells expressing HIV genes within in each culture. *SPI1* is the gene

that codes for PU.1. *MRC1* is the gene that codes for mannose receptor. (C) Violin plots

summarizing single-cell RNA transcripts expressed by primary human macrophages as in (B).

False-discovery rate corrected Wilcoxon rank-sum p-values are shown above the conditions

being compared. (D, G) Representative immunofluorescent images of MDMs from a single

donor infected with either 89.6^{wt} or 89.6$^{\Delta vpr}$ (MDMs from n=3 independent donors). HIV-infected

cells identified by Gag staining. (E, H) Quantification of ISG15 or IFITM3 corrected total cell

fluorescence in Gag+ cells, or (F, I) Gag- cells. The number of cells quantified for each condition

as indicated. Error bars represent standard error mean, n = the number of cells quantified. P

values were determined using an analysis of variance (ANOVA) with Tukey's multiple

comparisons test. ** indicates $p < 0.01$, *** indicates $p < 0.001$, **** indicates $p < 0.0001$.

Figure 4. PU.1 levels decrease in the presence of Vpr. (A) NL4-3 ΔGPE-GFP viral genome

map. (B) Representative flow cytometry histogram of PU.1 expression in infected (GFP$^+$) or

uninfected bystander (GFP$^-$) MDMs infected with NL4-3 ΔGPE-GFP with or without *vpr* and

collected on day 7 post infection. (C) Summary graph showing the percentage of infected

(GFP$^+$) cells that do not express PU.1 as determined by flow cytometry as depicted in Figure

4B. The mean +/- standard deviation from n=4 independent donors is shown for each time point.

P values were determined using an analysis of variance (ANOVA) with Tukey's multiple

comparisons test. *** indicates $p = 0.001$, **** indicates $p < 0.0001$. (D) Lentiviral map of

vectors encoding 3xFLAG-encoding *vpr* or *vpx* genes. (E) PU.1 expression plasmid map for full-

length, human PU.1. (F) Representative flow cytometric plots of HEK 293T cells transiently

transfected with PU.1 and the indicated FLAG-tagged viral protein. (G) Summary graph showing

PU.1$^+$ expression in transfected (FLAG$^+$) cells. Each point represents the mean of three

technical replicates. The mean +/- standard deviation of four independent experiments is

shown. P values were determined using one-way ANOVA compared to control, **** indicates p < 0.0001. (H) Immunoblot analysis of PU.1 in K562 cells transduced with the indicated lentivirus expressing 3xFLAG-tagged viral proteins. Results are representative of those from three independent experiments. (I) Summary graph of *SPI1* (the gene encoding PU.1) expression in HEK 293T cells co-transfected with PU.1 and an 89.6 expression vector with or without an intact *vpr* open reading frame. *SPI1* levels were assessed from purified RNA using RT quantitative real time PCR. Results represent the mean fold change compared with wild type +/- standard deviation for samples performed in triplicate for two independent experiments.

Figure 5. Vpr-mediated reduction of PU.1 is conserved in HIV-2 and SIV molecular clones. (A) Genomic maps for HIV-1, HIV-2, and select SIV genomes. HIV-1, SIVcpz, and SIVgor genomes contain *vpr* and *vpu* genes but not *vpx*. HIV-2, SIVrcm, and SIVsm contain *vpx* and *vpr*. SIVagm contains *vpr*. (B and D) Representative flow cytometric plots of HEK 293T cells transiently transfected with expression plasmids for PU.1 and the indicated FLAG-tagged viral protein. (C and E) Summary graph of data from B and D, respectively. The percentage of PU.1$^+$ cells per transfected (FLAG$^+$) cells is shown. Each point represents the average of three technical replicates. The mean +/- standard deviation is shown for n=6 or 11 independent experiments, respectively. Part E was additionally normalized to HIV-2$_{ROD}$Vpx for each experiment. **** indicates p < 0.0001 using one-way ANOVA compared to control.

Figure 6. Vpr forms a DCAF-1-dependent complex with PU.1. (A) Western blot analysis of lysates from HEK 293T cells co-transfected with the indicated expression constructs and immunoprecipitated with an antibody directed against the FLAG epitope. (B) Western blot analysis of lysates from K562 cells transduced with lentiviruses and treated as in part A. (C) Western blot analysis of lysates from MDMs transduced with lentivirus and treated as in part A. For parts A, B, and C, results are representative of three independent experiments. (D) Western blot analysis of lysates from K562 cells stably expressing either control, non-targeting shRNA (shScramble) or an shRNA targeting DCAF1 (shDCAF1), then transduced with FLAG-tagged 89.6-Vpr expression lentivirus and immunoprecipitated using an antibody directed against the FLAG epitope. (E) Western blot analysis of lysates from K562 cells transduced with FLAG-tagged 89.6-Vpr^{WT} or 89.6-Vpr^{Q65R} expressing lentiviruses and immunoprecipitated using an antibody directed against the FLAG epitope. (F) Western blot of lysates from HEK 293T cells stably transduced with virus expressing a non-targeting shRNA (shScramble) or one targeting DCAF1 (shDCAF1) and immunoprecipitated with an antibody directed against the FLAG

epitope. (G) Western blot analysis of lysates from HEK 293T cells co-transfected with the

indicated expression constructs and immunoprecipitated with an antibody directed against the

FLAG epitope. Results are representative of three independent experiments.

Figure 7. An intact DCAF-1 interaction domain is required for Vpr and DCAF1 to degrade

PU.1. (A) Immunoblot analysis of lysates from MDMs treated for five hours with the indicated

virions. (B) Summary graph of PU.1 protein normalized to vinculin from MDMs incubated for five

hours with the indicated viruses from (A). Each point is representative of an independent donor.

Statistical significance was determined using a mixed-effects analysis with Tukey's multiple

comparisons test. * indicates $p < 0.05$. (C) Western blot analysis of lysates from HEK 293T cells

transfected with the indicated expression construct and treatment as indicated with 10µM

MG132 or vehicle (Veh) control (DMSO). A GFP-expressing plasmid was included where

indicated as control for transfection efficiency. (D) Immunoblot analysis of lysates from MDMs

preincubated for two hours with vehicle (Veh) or MG132 as indicated and then treated for five

hours with the indicated virus as in part (A). (E) Summary graph of PU.1 protein normalized to

vinculin from MDMs treated for five hours with the indicated viruses from (D) and quantified as

for part (B). Each point is representative of an independent donor. (F) Working model of PU.1

and TET2 interacting with Vpr and DCAF1 in macrophages.

Figure 8. Reducing PU.1 increases Env output in HIV-1 infected MDMs. (A) Genome map

for full-length 89.6^{wt} HIV-1 modified to replace the *vpr* ORF with a U6-promoter followed by

either a non-targeting shRNA control (shScramble) or an shRNA targeting the PU.1 transcripts

(shSPI1). (B) Immunoblot from K562 cells stably expressing the shRNAs from (A). (C)

Immunoblot analysis from two independent MDM donors. Lysates collected from MDMs infected

with the virus from (A) expressing the indicated shRNA. White bands represent the removal of

irrelevant conditions from the same blot. (D) Proposed model of the PU.1-mediated antiviral

response disrupted by HIV-Vpr. HIV (-) Vpr indicates an infection of primary macrophages with

HIV that does not express Vpr. PU.1 protein is maintained and available to regulate anti-viral

response genes with co-factors such as IRF or TET2. HIV (+) Vpr indicates an infection with

HIV that expresses Vpr. PU.1 protein is less available to regulate anti-viral response genes,

contributing to innate immune evasion.

REFERENCES

1. Balliet, J. W. *et al.* Distinct Effects in Primary Macrophages and Lymphocytes of the Human Immunodeficiency Virus Type 1 Accessory Genes vpr, vpu, and nef: Mutational Analysis of a Primary HIV-1 Isolate. *Virology* **200**, 623–631 (1994).

2. YUH-LING LU, ROBERT P. BENNETT, JOHN W. WILLS, ROBERT GORELICK, & LEE RATNER. A leucine triplet repeat sequence (LXX)4 in p6gag is important for Vpr incorporation into human immunodeficiency virus type 1 particles. *J. Virol.* **69**, 6873–6879 (1995).

3. Eckstein, D. A. *et al.* HIV-1 Vpr enhances viral burden by facilitating infection of tissue macrophages but not nondividing CD4+ T cells. *J. Exp. Med.* **194**, 1407–1419 (2001).

4. Connor, R. I., Chen, B. K., Choe, S. & Landau, N. R. Vpr Is Required for Efficient Replication of Human Immunodeficiency Virus Type-1 in Mononuclear Phagocytes. *Virology* **206**, 935–944 (1995).

5. Dedera, D., Hu, W., Vander Heyden, N. & Ratner, L. Viral protein R of human immunodeficiency virus types 1 and 2 is dispensable for replication and cytopathogenicity in lymphoid cells. *J. Virol.* **63**, 3205–3208 (1989).

6. Rücker, E., Grivel, J.-C., Münch, J., Kirchhoff, F. & Margolis, L. Vpr and Vpu Are Important for Efficient Human Immunodeficiency Virus Type 1 Replication and CD4+ T-Cell Depletion in Human Lymphoid Tissue Ex Vivo. *J. Virol.* **78**, 12689–12693 (2004).

7. Lu, Y. L., Spearman, P. & Ratner, L. Human immunodeficiency virus type 1 viral protein R localization in infected cells and virions. *J. Virol.* **67**, 6542–6550 (1993).

8. Laguette, N. *et al.* Premature Activation of the SLX4 Complex by Vpr Promotes G2/M Arrest and Escape from Innate Immune Sensing. *Cell* **156**, 134–145 (2014).

9. Yan, J., Shun, M. C., Zhang, Y., Hao, C. & Skowronski, J. HIV-1 Vpr counteracts HLTF-mediated restriction of HIV-1 infection in T cells. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 9568–9577 (2019).

10. Ahn, J. *et al.* HIV-1 Vpr Loads Uracil DNA Glycosylase-2 onto DCAF1, a Substrate Recognition Subunit of a Cullin 4A-RING E3 Ubiquitin Ligase for Proteasome-dependent Degradation *. *J. Biol. Chem.* **285**, 37333–37341 (2010).

11. Lahouassa, H. *et al.* HIV-1 Vpr degrades the HLTF DNA translocase in T cells and macrophages. *Proc. Natl. Acad. Sci.* **113**, 5311–5316 (2016).

12. Withers-Ward, E. S., Mueller, T. D., Chen, I. S. Y. & Feigon, J. Biochemical and Structural
Analysis of the Interaction between the UBA(2) Domain of the DNA Repair Protein HHR23A
and HIV-1 Vpr. *Biochemistry* **39**, 14103–14112 (2000).

13. Wang, X. *et al.* HIV-1 Vpr Protein Inhibits Telomerase Activity via the EDD-DDB1-VPRBP
E3 Ligase Complex *. *J. Biol. Chem.* **288**, 15474–15480 (2013).

14. Romani, B., Baygloo, N. S., Aghasadeghi, M. R. & Allahbakhshi, E. HIV-1 Vpr Protein
Enhances Proteasomal Degradation of MCM10 DNA Replication Factor through the Cul4-
DDB1[VprBP] E3 Ubiquitin Ligase to Induce G2/M Cell Cycle Arrest *. *J. Biol. Chem.* **290**,
17380–17389 (2015).

15. Yan, J. *et al.* HIV-1 Vpr Reprograms CLR4DCAF1 E3 Ubiquitin Ligase to Antagonize
Exonuclease 1-Mediated Restriction of HIV-1 Infection. *mBio* **9**, e01732-18 (2018).

16. Zhang, S., Feng, Y., Narayan, O. & Zhao, L.-J. Cytoplasmic retention of HIV-1 regulatory
protein Vpr by protein-protein interaction with a novel human cytoplasmic protein VprBP.
Gene **263**, 131–140 (2001).

17. McCall, C. M. *et al.* Human Immunodeficiency Virus Type 1 Vpr-Binding Protein VprBP, a
WD40 Protein Associated with the DDB1-CUL4 E3 Ubiquitin Ligase, Is Essential for DNA
Replication and Embryonic Development. *Mol. Cell. Biol.* **28**, 5621–5633 (2008).

18. Romani, B. & Cohen, É. A. Lentivirus Vpr and Vpx accessory proteins usurp the cullin4–
DDB1 (DCAF1) E3 ubiquitin ligase. *Curr. Opin. Virol.* **2**, 755–763 (2012).

19. Hakata, Y., Miyazawa, M. & Landau, N. R. Interactions with DCAF1 and DDB1 in the CRL4
E3 ubiquitin ligase are required for Vpr-mediated G2 arrest. *Virol. J.* **11**, 108 (2014).

20. Schröfelbauer, B., Hakata, Y. & Landau, N. R. HIV-1 Vpr function is mediated by interaction
with the damage-specific DNA-binding protein DDB1. *Proc. Natl. Acad. Sci. U. S. A.* **104**,
4130–4135 (2007).

21. Belzile, J.-P. *et al.* HIV-1 Vpr-Mediated G2 Arrest Involves the DDB1-CUL4AVPRBP E3
Ubiquitin Ligase. *PLoS Pathog.* **3**, e85 (2007).

22. Hrecka, K. *et al.* HIV-1 and HIV-2 exhibit divergent interactions with HLTF and UNG2 DNA
repair proteins. *Proc. Natl. Acad. Sci. U. S. A.* **113**, E3921–E3930 (2016).

23. Lavin, Y., Mortha, A., Rahman, A. & Merad, M. Regulation of macrophage development and
function in peripheral tissues. *Nat. Rev. Immunol.* **15**, 731–744 (2015).

24. Silvin, A. & Manel, N. Innate immune sensing of HIV infection. *Curr. Opin. Immunol.* **32**, 54–
60 (2015).

25. Turkistany, S. A., Dekoter, R. P. & Hirszfeld, Ó. L. The Transcription Factor PU.1 is a
Critical Regulator of Cellular Communication in the Immune System. doi:10.1007/s00005-
011-0147-9.

26. Fisher, R. C. & Scott, E. W. Role of PU.1 in Hematopoiesis. *Stem Cells* **16**, 25–37 (1998).

27. Dakic, A. *et al.* PU.1 regulates the commitment of adult hematopoietic progenitors and
restricts granulopoiesis. *J. Exp. Med.* **201**, 1487–1502 (2005).

28. Gupta, P., Gurudutta, G. U., Saluja, D. & Tripathi, R. P. PU.1 and partners: regulation of
haematopoietic stem cell fate in normal and malignant haematopoiesis. *J. Cell. Mol. Med.*
13, 4349–4363 (2009).

29. Graves, B. J. & Petersen, J. M. Specificity within the ets Family of Transcription Factors. in
Advances in Cancer Research (eds. Vande Woude, G. F. & Klein, G.) vol. 75 1–57
(Academic Press, 1998).

30. Marecki, S. & Fenton, M. J. PU.1/interferon regulatory factor interactions. *Cell Biochem.*
Biophys. **33**, 127–148 (2000).

31. Imperato, M. R., Cauchy, P., Obier, N. & Bonifer, C. The RUNX1–PU.1 axis in the control of
hematopoiesis. *Int. J. Hematol.* **101**, 319–329 (2015).

32. de la Rica, L. *et al.* PU.1 target genes undergo Tet2-coupled demethylation and DNMT3b-
mediated methylation in monocyte-to-osteoclast differentiation. *Genome Biol.* **14**, 1–21
(2013).

33. Collins, D. R., Lubow, J., Lukic, Z., Mashiba, M. & Collins, K. L. Vpr Promotes Macrophage-
Dependent HIV-1 Infection of CD4+ T Lymphocytes. *PLoS Pathog.* **11**, e1005054 (2015).

34. Mashiba, M., Collins, D. R., Terry, V. H. & Collins, K. L. Vpr overcomes macrophage-
specific restriction of HIV-1 Env expression and virion production. *Cell Host Microbe* **16**,
722–735 (2014).

35. Lubow, J. *et al.* Mannose receptor is an HIV restriction factor counteracted by Vpr in
macrophages. *eLife* **9**, (2020).

36. Liang, G. *et al.* Membrane metalloprotease TRABD2A restricts HIV-1 progeny production in
resting CD4 + T cells by degrading viral Gag polyprotein. *Nat. Immunol.* (2019)
doi:10.1038/s41590-019-0385-2.

37. Linehan, S. A., Martínez-Pomares, L., Stahl, P. D. & Gordon, S. Mannose Receptor and Its
Putative Ligands in Normal Murine Lymphoid and Nonlymphoid Organs: In Situ Expression
of Mannose Receptor by Selected Macrophages, Endothelial Cells, Perivascular Microglia,
and Mesangial Cells, but not Dendritic Cells. *J. Exp. Med.* **189**, 1961–1972 (1999).

38. Wang, Q. & Su, L. Vpr Enhances HIV-1 Env Processing and Virion Infectivity in
Macrophages by Modulating TET2-Dependent IFITM3 Expression. *mBio* **10**, e01344-19
(2019).

39. Lv, L. *et al.* Vpr Targets TET2 for Degradation by CRL4VprBP E3 Ligase to Sustain IL-6
Expression and Enhance HIV-1 Replication. *Mol. Cell* **70**, 961-970.e5 (2018).

40. Welch, J. D. *et al.* Single-Cell Multi-omic Integration Compares and Contrasts Features of
Brain Cell Identity. *Cell* **177**, 1873–1887 (2019).

41. Zhang, F. & Bieniasz, P. D. HIV-1 Vpr induces cell cycle arrest and enhances viral gene
expression by depleting CCDC137. *eLife* **9**, e55806 (2020).

42. Pham, T. H. *et al.* Mechanisms of in vivo binding site selection of the hematopoietic master
transcription factor PU.1. *Nucleic Acids Res.* **41**, 6391–6402 (2013).

43. Buckland, J. Haematopoiesis: PU.1 pinned down. *Nat. Rev. Immunol.* **2**, 149 (2002).

44. Caldwell, R. L., Egan, B. S. & Shepherd, V. L. HIV-1 Tat Represses Transcription from the
Mannose Receptor Promoter. *J. Immunol.* **165**, 7035–7041 (2000).

45. Medzhitov, R., Preston-Hurlburt, P. & Janeway, C. A. A human homologue of the Drosophila
Toll protein signals activation of adaptive immunity. *Nature* **388**, 394–397 (1997).

46. Molteni, M., Gemma, S. & Rossetti, C. The Role of Toll-Like Receptor 4 in Infectious and
Noninfectious Inflammation. *Mediators Inflamm.* **2016**, 6978936 (2016).

47. Beutler, B. A. TLRs and innate immunity. *Blood* **113**, 1399–1407 (2009).

48. Perng, Y. C. & Lenschow, D. J. ISG15 in antiviral immunity and beyond. *Nat. Rev. Microbiol.*
16, 423–439 (2018).

49. Schneider, W. M., Chevillotte, M. D. & Rice, C. M. Interferon-Stimulated Genes: A Complex
Web of Host Defenses. *Annu. Rev. Immunol.* **32**, 513–545 (2014).

50. Shimazu, R. *et al.* MD-2, a Molecule that Confers Lipopolysaccharide Responsiveness on
Toll-like Receptor 4. *J. Exp. Med.* **189**, 1777–1782 (1999).

51. Nagai, Y. *et al.* Essential role of MD-2 in LPS responsiveness and TLR4 distribution. *Nat.*
Immunol. **3**, 667–672 (2002).

52. Chen, H. *et al.* PU.1 (Spi-1) autoregulates its expression in myeloid cells. *Oncogene* **11**,
1549–1560 (1995).

53. Kao, S. *et al.* The Myeloid-Specific Transcription Factor PU.1 Upregulates Mannose
Receptor Expression but Represses Basal Activity of the HIV-LTR Promoter. *J. Virol.* **96**,
e00652-22 (2022).

54. Collins, D. R. & Collins, K. L. HIV-1 accessory proteins adapt cellular adaptors to facilitate
immune evasion. *PLoS Pathog.* **10**, e1003851 (2014).

55. Sharp, P. M. & Hahn, B. H. Origins of HIV and the AIDS Pandemic. *Cold Spring Harb.*
Perspect. Med. **1**, a006841 (2011).

56. Sakai, Y. *et al.* Expression Profiles of Vpx/Vpr Proteins Are Co-related with the Primate
Lentiviral Lineage. *Front. Microbiol.* **7**, (2016).

57. Yurkovetskiy, L. *et al.* Primate immunodeficiency virus proteins Vpx and Vpr counteract
transcriptional repression of proviruses by the HUSH complex. *Nat. Microbiol.* **3**, 1354–1361
(2018).

58. Lozzio, C. B. & Lozzio, B. B. Human Chronic Myelogenous Leukemia Cell-Line With
Positive Philadelphia Chromosome. *Blood* **45**, 321–334 (1975).

59. Zhao, L.-J., Mukherjee, S. & Narayan, O. Biochemical mechanism of HIV-I Vpr function.
Specific interaction with a cellular protein. *J. Biol. Chem.* **269**, 15577–15582 (1994).

60. DeHart, J. L. *et al.* HIV-1 Vpr activates the G2 checkpoint through manipulation of the
ubiquitin proteasome system. *Virol. J.* **4**, 57 (2007).

61. Lu, J., Pan, Q., Rong, L., Liu, S.-L. & Liang, C. The IFITM Proteins Inhibit HIV-1 Infection. *J.*
Virol. **85**, 2126–2137 (2011).

62. Compton, A. A. *et al.* IFITM Proteins Incorporated into HIV-1 Virions Impair Viral Fusion and
Spread. *Cell Host Microbe* **16**, 736–747 (2014).

63. Kumar, H., Kawai, T. & Akira, S. Toll-like receptors and innate immunity. *Biochem. Biophys.*
Res. Commun. **388**, 621–625 (2009).

64. Deletsu, S. D. *et al.* Identification and characterization of Stathmin 1 as a host factor
involved in HIV-1 latency. *Biochem. Biophys. Res. Commun.* **567**, 106–111 (2021).

65. Park, G.-H. *et al.* Association between Interferon-Inducible Protein 6 (IFI6) Polymorphisms
and Hepatitis B Virus Clearance. *Genomics Inform.* **11**, 15–23 (2013).

66. Del Cornò, M. *et al.* HIV-1 gp120 signaling through TLR4 modulates innate immune
activation in human macrophages and the biology of hepatic stellate cells. *J. Leukoc. Biol.*
100, 599–606 (2016).

67. Okumura, A., Lu, G., Pitha-Rowe, I. & Pitha, P. M. Innate antiviral response targets HIV-1
release by the induction of ubiquitin-like protein ISG15. *Proc. Natl. Acad. Sci.* **103**, 1440–
1445 (2006).

68. Tartour, K. *et al.* Interference with the production of infectious viral particles and bimodal
inhibition of replication are broadly conserved antiviral properties of IFITMs. *PLOS Pathog.*
13, e1006610 (2017).

69. Yu, J. *et al.* IFITM Proteins Restrict HIV-1 Infection by Antagonizing the Envelope
Glycoprotein. *Cell Rep.* **13**, 145–156 (2015).

70. Lorsbach, R. B. *et al.* TET1, a member of a novel protein family, is fused to MLL in acute
myeloid leukemia containing the t(10;11)(q22;q23). *Leukemia* **17**, 637–641 (2003).

71. Lang, S. M. *et al.* Importance of vpr for infection of rhesus monkeys with simian
immunodeficiency virus. *J. Virol.* **67**, 902–912 (1993).

72. Munde, M. *et al.* Structure-dependent inhibition of the ETS-family transcription factor PU.1
by novel heterocyclic diamidines. *Nucleic Acids Res.* **42**, 1379–1390 (2014).

73. McNamara, L. A., Ganesh, J. A. & Collins, K. L. Latent HIV-1 infection occurs in multiple
subsets of hematopoietic progenitor cells and is reversed by NF-κB activation. *J. Virol.* **86**,
9337–50 (2012).

74. Norrander, J., Kempe, T. & Messing, J. Construction of improved M13 vectors using
oligodeoxynucleotide-directed mutagenesis. *Gene* **26**, 101–106 (1983).

75. Carter, C. C. *et al.* HIV-1 infects multipotent progenitor cells causing cell death and
establishing latent cellular reservoirs. *Nat. Med.* **16**, 446–451 (2010).

76. Weber, K., Bartsch, U., Stocking, C. & Fehse, B. A multicolor panel of novel lentiviral 'gene
ontology' (LeGO) vectors for functional gene analysis. *Mol. Ther. J. Am. Soc. Gene Ther.*
16, 698–706 (2008).

77. Moffat, J. *et al.* A Lentiviral RNAi Library for Human and Mouse Genes Applied to an
Arrayed Viral High-Content Screen. *Cell* **124**, 1283–1298 (2006).

78. Sarbassov, D. D., Guertin, D. A., Ali, S. M. & Sabatini, D. M. Phosphorylation and
Regulation of Akt/PKB by the Rictor-mTOR Complex. *Science* **307**, 1098–1101 (2005).

79. Wickham, H. *Ggplot2.* (Springer, New York, NY, 2009). doi:10.1007/978-0-387-98141-3.

80. Heinz, S. *et al.* Simple Combinations of Lineage-Determining Transcription Factors Prime
cis-Regulatory Elements Required for Macrophage and B Cell Identities. *Mol. Cell* **38**, 576–
589 (2010).

81. Eden, E., Navon, R., Steinfeld, I., Lipson, D. & Yakhini, Z. GOrilla: a tool for discovery and
visualization of enriched GO terms in ranked gene lists. *BMC Bioinformatics* **10**, 48 (2009).

82. Eden, E., Lipson, D., Yogev, S. & Yakhini, Z. Discovering Motifs in Ranked Lists of DNA
Sequences. *PLOS Comput. Biol.* **3**, e39 (2007).

83. Supek, F., Bošnjak, M., Škunca, N. & Šmuc, T. REVIGO Summarizes and Visualizes Long
Lists of Gene Ontology Terms. *PLOS ONE* **6**, e21800 (2011).

84. Kent, W. J. *et al.* The Human Genome Browser at UCSC. *Genome Res.* **12**, 996–1006
(2002).

85. Grant, C. E., Bailey, T. L. & Noble, W. S. FIMO: scanning for occurrences of a given motif.
Bioinformatics **27**, 1017–1018 (2011).

86. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29

(2021).

87. RStudio Team. RStudio: Integrated Development for R. RStudio, PBC (2020).

88. Gasmi, M. *et al.* Requirements for Efficient Production and Transduction of Human

Immunodeficiency Virus Type 1-Based Vectors. *J. Virol.* **73**, 1828–1834 (1999).

1129

Figure 1

Figure 2

A

ETS Family Transcription Factors

Motif	TF Name	q-value	# Targets w/ Sequence
AGAGGAAGTG	PU.1	0.0005	316.0

ETS/IRF Family Transcription Factors

Motif	TF Name	q-value	# Targets w/ Sequence
CGGAAGTGAAAC	PU.1-IRF	0.0004	670.0

B HOMER predicted transcription factor expression

SPI1 KLF14 NFYA SP2

IRF8 ETV4 NRF1 CLOCK

C HOMER-identified targets of PU.1 binding

Expression
- DOWN
- NO CHANGE
- UP
- PU.1(SPI1) Targets

D Biological processes associated with PU.1 targeted genes

E PU.1 regulated genes

TREM2 p = 3.13x10^-244
FABP4 p = 5.15x10^-166
LY96   p = 6.24x10^-163
STMN1 p = 3.12x10^-144
IFI6   p < 1x10^-287
IFITM3 p = 1.96x10^-190
ISG15 p = 8.53x10^-25
MRC1   p = 0.002

A Selected biological processes associated with PU.1 targeted genes

B Expression of HIV and macrophage markers

C PU.1 regulated genes involved in TLR and IFN-I signalling

D DAPI | Gag | ISG15 | Merge

E Gag-positive MDM

F Gag-negative MDM

G DAPI | Gag | IFITM3 | Merge

H Gag-positive MDM

I Gag-negative MDM

A HIV-1 NL4-3 ΔGPE-GFP

B MDMs

C MDMs

D 3xFLAG-Vpx/Vpr

E PU.1 Expression Vector

F HEK 293T

G HEK 293T

H K562

I HEK 293T

A

HIV-1/SIVcpz/SIVgor

HIV-2/SIVrcm/SIVsm

SIVagm

B HEK 293T

C HEK 293T

D HEK 293T

E HEK 293T

Figure 6

A HIV-treated primary macrophages

B HIV-treated primary macrophages

C HEK 293T FLAG-Vpr

D HIV-treated primary macrophages

E HIV-treated primary macrophages

F Working model of PU.1 and TET2 interacting with Vpr and DCAF1 in macrophages

Figure 8

A 89.6^Δvpr-U6-shRNA

B K562

C

D