TITLE & AUTHORS

HIV-1 Vpr combats the PU.1-driven antiviral response in primary human macrophages.

Maria C. Virgilio[1,2,3], W. Miguel Disbennett[4,5,a], Thomas Chen[5], Jay Lubow[5,b], Joshua D. Welch[2],

Kathleen L. Collins[1,3,5,*]

[1]Cellular and Molecular Biology Program, University of Michigan, Ann Arbor, MI, USA

[2]Department of Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor,

MI, USA

[3]Department of Internal Medicine, University of Michigan, Ann Arbor, MI, USA\

[4]Post-Baccalaureate Research Education Program (PREP), University of Michigan, Ann Arbor,

MI, USA

[5]Department of Microbiology and Immunology, University of Michigan, Ann Arbor, MI, USA

[a]Present location: University of Pennsylvania, Philadelphia, PA, USA

[b]Present location: Fred Hutchinson Cancer Research Center, Seattle, WA, USA

[*] Correspondence: klcollin@umich.edu

SUMMARY

The HIV-1 accessory protein, Vpr, is an enigmatic protein required for efficient spread of HIV

from macrophages to T cells, a necessary step for propagation of infection. To illuminate the

role of Vpr in HIV-infection of primary macrophages, we used single-cell RNA sequencing to

capture the transcriptional changes during an HIV-1 spreading infection plus and minus Vpr. We

found that Vpr reprogramed HIV-infected macrophage gene expression by targeting the master

transcriptional regulator, PU.1. PU.1 was required for efficient induction of the host innate

immune response to HIV, including upregulation of *ISG15*, *LY96,* and *IFI6*. In contrast, we did

not observe direct effects of PU.1 on HIV gene transcription. Single cell gene expression

analysis also revealed Vpr countered an innate immune response to HIV-infection within

bystander macrophages via a PU.1-independent mechanism. The capacity of Vpr to target PU.1

and disrupt the anti-viral response was highly conserved across primate lentiviruses including

HIV-2 and several SIVs. By demonstrating how Vpr overcomes a critical early warning system

of infection, we identify a crucial reason why Vpr is necessary for HIV infection and spread.

KEYWORDS

Vpr, HIV-1, HIV-2, SIV, PU.1, macrophages, scRNA-seq, interferon, ISG15, TLR

INTRODUCTION

The HIV-1 genome encodes several accessory proteins that counteract innate and adaptive antiviral responses. Although HIV accessory proteins have been widely studied, the role of the Vpr accessory protein has remained enigmatic. Vpr is highly conserved amongst lentiviruses and is necessary for optimal replication in macrophages[1]. Studies using human lymphoid tissue, which is rich in both T cells and macrophages, have found that loss of Vpr decreases virus production but only when the virus strain is capable of efficiently infecting macrophages[1–5]. These studies provide evidence that Vpr enhances infection of macrophages and increases viral burden in tissues containing macrophages. Vpr localizes to the nucleus[6], where it is known to induce G2 cell cycle arrest by targeting a variety of host factors involved in post-replication DNA repair[7–14]. To target these factors, Vpr requires the host protein DCAF1 (also known as Vpr binding protein (VprBP))[15]. Through DCAF1, Vpr interacts with Damaged DNA binding protein 1 (DDB1) as part of the Cul4A E3 ubiquitin ligase complex where host proteins recruited by Vpr can be ubiquitinated, then degraded[16–21]. However, the effects of Vpr on DNA repair and cell cycle arrest do not provide a clear explanation for how Vpr enhances HIV infection of macrophages.

Macrophages are terminally differentiated antigen presenting cells that are critical for many immune functions, including antiviral innate immune responses[22,23]. Macrophage identity is tightly controlled through the timed expression of myeloid transcription factors, particularly PU.1 (Reviewed in [24]). PU.1 is a hematopoietic-specific and ETS family transcription factor that is essential for lymphoid and myeloid development [24–27]. ETS family proteins bind to purine rich DNA domains with a central GGAA/T core consensus[28]. Macrophages require early and continuously high levels of PU.1 expression (Reviewed in [24,25]) to maintain normal functionality. PU.1 regulates many essential macrophage genes including those for cytokine receptors including M-CSF and GM-CSF and the adhesion molecule CD11b. PU.1 also coordinates with other transcription factors such as IRF4 and C/EBPα, and TET methyl cytosine dioxygenase 2 (TET2; also known as ten-eleven translocation 2) to regulate gene expression[24,29–31].

We previously reported that Vpr counteracts accelerated degradation of the HIV Env protein, which occurs in HIV infected macrophages but not T cells. We hypothesized that Vpr disables a macrophage-specific restriction factor that detects and degrades HIV Env. We recently identified this factor as the macrophage mannose receptor (MR), which is highly expressed in

macrophages but not T cells[32,33]. MR senses HIV Env via densely packed high mannose

residues that serve as pathogen-associated molecular patterns (PAMPs) because they are

normally absent from host cellular proteins. MR recognition of HIV Env disrupts infection by

promoting lysosomal degradation of Env and Env-containing viral particles [34,35]. Although the

canonical targeting of host factors by Vpr involves the binding of Vpr to DCAF1, leading to

ubiquitination and degradation via the proteosome, Vpr does not directly interact with MR[34].

Instead, Vpr suppresses expression of the MR gene, *MRC1*[34]. Here, we examine the

mechanism by which Vpr suppresses the expression of *MRC1* and demonstrate that Vpr

additionally exerts suppressive effects on other important genes in macrophages to enhance

virion assembly and spread.

To accomplish this goal, we employed single-cell RNA sequencing (scRNA-seq) of primary

human macrophages infected with infectious wild type or Vpr-null HIV. Single-cell resolution

allowed us to distinguish the effects of Vpr on both infected and virally exposed, uninfected cells

in the same culture. Within infected cells, Vpr selectively downregulated genes controlled by the

macrophage-selective transcription factor, PU.1. By targeting PU.1, Vpr systemically disrupted

several antiviral factors, including *MRC1,* that would otherwise restrict HIV infection and spread.

Interestingly, we also identified PU.1 as the transcription factor responsible for partnering with

TET2 to control expression of *IFITM3,* a previously reported target of Vpr that is also capable of

disrupting Env function[36]. Comparing the expression of antiviral-response genes between

infected cells and virus-exposed uninfected bystander cells in the same culture, we discovered

that Vpr caused a systemic reduction in genes implicated in Toll-like receptor and Type I

interferon (IFN-I) signaling that affected all the cells in the culture. Thus, we provide a systems

level explanation for the positive effect of Vpr on HIV spread in macrophages.

Finally, we found that Vpr-mediated downmodulation of PU.1-regulated gene expression is

mediated by direct protein-protein interaction between Vpr and PU.1, resulting in accelerated

proteasomal degradation of PU.1. This activity of Vpr is conserved among all HIV-1 molecular

clones tested as well as HIV-2 and SIV. Remarkably, both the interaction between PU.1 and Vpr

as well PU.1's subsequent degradation, require the Vpr interacting protein, DCAF1. Intriguingly,

the PU.1 transcriptional co-factor, TET2, is co-recruited with PU.1 to DCAF1 by Vpr, which

aligns our results with other reports that Vpr targets TET2[36,37]. Altogether, our data support a

model in which Vpr promotes HIV spread via systemic detrimental effects on the host innate

antiviral response to infection.

RESULTS

Single-cell RNA sequencing of HIV-1 infected MDMs reveals Vpr-dependent transcriptional changes. Vpr is known to counteract a number of host factors, primarily those implicated in DNA repair and restriction of HIV Env [34,36]. Recently we demonstrated Vpr decreases transcriptional expression of the host restriction factor, mannose receptor (*MRC1*), which we hypothesize results from Vpr targeting a macrophage-specific transcription factor (Figure 1A). To identify the transcription factor targeted by Vpr, we undertook an unbiased approach to characterize the genome-wide effects of HIV-1 Vpr in monocyte derived macrophages (MDMs). For these studies, we used an infectious HIV molecular clone (89.6) plus or minus an intact *vpr* gene (Figure 1B). We infected primary human CD14$^+$ MDMs from three independent healthy donors and allowed the infection to spread for 10 days before harvesting cells (uninfected, 89.6^{wt} or 89.6$^{\Delta vpr}$ infected) and prepared them for single-cell gene expression analysis (Figure 1C). LIGER (Linked Inference of Genomic Experimental Relationships) [38] was used to integrate all nine data sets. LIGER enabled us to identify shared cell profiles across donors while reducing sample-specific differences. Uniform Manifold Approximation and Projection (UMAP) visualization of all donor sets showed no significant donor or batch-specific differences amongst the samples (Figure 1D). Integration of virus-treated cells revealed a relatively homogeneous population (Figure 1E, left), with 89.6^{wt} and 89.6$^{\Delta vpr}$-exposed cells evenly dispersed throughout the clusters (Figure 1E, center).

To separate the bona fide infected cells from bystander cells, we computationally segregated cells into a subset that expressed both *tat* and *gag* transcripts from those that expressed neither (Figure 1E, right, designated "Infected"). The proportion of infected cells identified by this analysis was like that found by the standard method of identifying infected cells by intracellular Gag staining and detection by flow cytometry (Supplemental Table 1). A comparison of gene expression profiles in cells with and without Vpr revealed that Vpr expression boosted HIV gene expression as previously reported[39] and caused a significant transcriptional shift in host gene expression (Figure 1F). Using a Wilcoxon rank-sum test and false discovery rate correction, we identified 3150 genes with statistically significant two-fold or greater differential expression between 89.6^{wt} and 89.6$^{\Delta vpr}$-infected cells.

Figure 1. Single-cell RNA sequencing of HIV-1 infected MDMs reveals Vpr-dependent transcriptional changes. (A) Schematic diagram illustrating the objective of single-cell RNA sequencing; the identification of Vpr-targeted transcription factor(s) in HIV infected primary macrophages. (B) Genome maps for full-length 89.6^{wt} HIV-1 (top) and the same viral genome with a premature stop-codon in *vpr* (bottom). (C) Experimental setup for the generation of the scRNA-seq datasets. (D) UMAP representation of LIGER-integrated scRNA-seq data from MDM samples treated as shown in (C) and listed in Dataset. (E, Right) UMAP representations of LIGER-integrated scRNA-seq data from MDMs treated as indicated. (E, Right) Bona fide infected cells were identified based on expression of HIV *tat* and *gag*. (E, left) Colors indicate individual clusters. Cluster 0 = pro-inflammatory macrophages, Cluster 1 = cycling cells, Cluster 2 = dying cells, Cluster 3 = pro-inflammatory macrophages. (E, center and right) cells colored according to whether they were exposed to 89.6 WT-Vpr virus (blue) or Vpr-*null* virus (pink). (F) Volcano plot of differentially expressed genes from HIV 89.6^{wt} verses 89.6$^{\Delta vpr}$ infected MDMs from (E) as determined by Wilcoxin Rank Sum. Significance determined as greater than 1 log2 fold change and false discovery rate adjusted p-value of p < 0.05 (red-bars). Blue colored genes indicate genes less highly expressed in HIV 89.6^{wt} verses 89.6$^{\Delta vpr}$ infected MDMs, Red colored genes indicate genes more highly expressed, and black colored genes indicate no significant difference between datasets.

135 **Vpr-repressed genes in MDMs include targets of the transcription factor, PU.1.**

136 To identify the transcription factor(s) targeted by Vpr, we used the DNA motif identification

137 software, HOMER (Hypergeometric Optimization of Motif EnRichment), to scan for common

138 transcription factor binding motifs in the promoters of genes downregulated in the presence of

139 Vpr. Our analysis revealed several candidate transcription factors (TFs) and TF-binding families,

140 including many members of the ETS family of TFs (Supplemental Figure 1A). However, only

141 PU.1 (encoded by the gene *SPI1*) was highly expressed in our data sets (Figure 2B). Moreover,

142 PU.1 appeared twice in the list – as both a member of the ETS family and as a co-factor in the

143 ETS/IRF (Interferon Regulatory Factor) family of transcription factors, a relationship that is well

144 documented[29] (Figure 2A). PU.1 is the master myeloid transcription factor (Reviewed in [26,40,41]).

145 It is required for terminal differentiation of macrophages and is necessary to maintain

146 macrophage immune function [24]. From the list of most downregulated genes in Vpr-expressing

147 MDMs, HOMER identified 316 genes with a PU.1 binding motif in their promoter and 670 PU.1-

148 IRF co-regulated genes, for a combined 840 distinct PU.1-regulated genes identified (Figure 2C,

149 E). One of the genes identified with a PU.1 binding motif was interferon induced transmembrane

150 protein 3 (*IFITM3)*. IFITM3 was previously identified as an important host restriction factor that is

151 similar to mannose receptor, in that IFITM3 targets HIV Env and has reduced gene expression

152 (by reverse transcription quantitative PCR (RT-qPCR)) in the presence of Vpr in HIV-infected

153 MDMs[36]. Gene ontology analysis of the 840 genes targeted by PU.1 revealed that they are

154 enriched for regulation of the immune system and defense response (Figure 2D).

155

156 Because we had previously demonstrated that the gene encoding mannose receptor (*MRC1*)

157 was transcriptionally downmodulated approximately two-fold by Vpr in HIV infected primary

158 human macrophages using RT-qPCR[34], we examined our data set to specifically identify *MRC1*

159 transcripts. In the scRNA-seq data set shown here, we identified a Vpr-dependent change in

160 *MRC1* transcripts (log2FC = 0.932) that is statistically significant (false-discovery rate corrected

161 Wilcoxon rank-sum p-value = 0.002) but fell just below our arbitrary cutoff of a two-fold change

162 (Figures 1F, 2C). Consistent with the pattern we observed for other Vpr-suppressed genes, we

163 identified several PU.1 binding sites in the promoter region of *MRC1* (Supplemental Figure 2)

164 when we manually scanned for PU.1 binding motifs, some of which have been previously

165 described[42]. While the effect of Vpr on *MRC1* expression is relatively small, it combines

166 synergistically with HIV Nef-dependent disruption of MR trafficking to reverse mannose

167 receptor-dependent lysosomal degradation of Env in HIV infected primary macrophages[34].

168

A

ETS Family Transcription Factors			
Motif	TF Name	q-value	# Targets w/ Sequence
AGAGGAAGTG	PU.1	0.0005	316.0

ETS/IRF Family Transcription Factors			
Motif	TF Name	q-value	# Targets w/ Sequence
CGGAAGTGAAAC	PU.1-IRF	0.0004	670.0

B HOMER predicted transcription factor expression

C HOMER-identified targets of PU.1 binding

D Biological processes associated with PU.1 targeted genes

E PU.1 regulated genes

Figure 2. Vpr downmodulates PU.1-dependent transcription. (A) PU.1 motifs identified by HOMER as present in the promoters of Vpr-downmodulated genes (Figure 1E, Blue). (B) Violin plots displaying RNA abundance of the indicated transcription factor genes in MDMs infected with the indicated virus. (C) Volcano plot as in Figure 1E except that genes containing a PU.1 or PU.1-IRF binding motif in their promoter region are highlighted in green (See also Supplemental Figure 2). (D) Biological processes associated with the PU.1 targeted genes from (B). Size of circle indicates the relative number of GO terms with the process. FDR adjusted q-values associated with GO terms as indicated by the color. (E) Violin plots displaying RNA abundance of the indicated genes in MDMs infected with the indicated virus. HIV-1 89.6^{wt} (WT); HIV-1 89.6$^{\Delta vpr}$ (ΔVpr).

Vpr suppresses PU.1 regulated genes implicated in Toll-like receptor and IFN-I

responses.

PU.1 regulated genes that are downmodulated in Vpr expressing cells include several factors

implicated in Toll-like receptor (TLR) signaling pathways (Figure 3A). TLRs are evolutionarily

conserved, pattern recognition receptors (PRRs) that recognize PAMPs ([43–45]), including HIV. To

determine whether these effects of Vpr are restricted to infected cells within the culture, we

compared gene expression profiles amongst all the sample types collected from our single-cell

analysis: uninfected (virus naive), 89.6[wt] and 89.6[Δvpr]-infected, and 89.6[wt] and 89.6[Δvpr] virally

exposed but uninfected (bystander) cells. This was accomplished by computationally

segregating cells into subsets that expressed both *tat* and *gag* transcripts from those that were

virally exposed but appeared uninfected (*tat* and *gag*-negative) (Figure 3B).

As expected, expression of genes involved in the TLR-mediated IFN-I response to infection,

including Interferon Stimulated Gene 15 (*ISG15),* which is upregulated in response to IFN-I and

TLR-signaling (Reviewed in [46,47]), *LY96, and* interferon inducible protein-6 (*IFI6)* (PU.1-IRF

regulated genes) and *IFITM3* was very low in uninfected, unexposed cells (Figure 3C). In

contrast, each of these genes was upregulated in cells from HIV treated primary macrophage

cultures (Figure 3C). As discussed above, Vpr counteracted upregulation of this anti-viral

response in infected cells (Figure 2C, E and Figure 3C). In addition, we were surprised to find

many of these genes upregulated in bystander cells as well (Figure 3C) and that Vpr limited

their induction. For example, we found that *ISG15*, was more highly expressed in 89.6[Δvpr]

bystander cells compared to 89.6[wt]-exposed bystander cells. Similar observations were made

for *LY96*, *IFI6*, and *IFITM3*. Interestingly, the protein product for *LY96* is myeloid differentiation 2

(MD2) which is required for TLR4 ligand-induced activation at the cell surface[48,49]. By

comparison, other subsets of genes, including *SPI1* and *MRC1* were primarily downmodulated

in the infected subset and either not at all in the bystander cells or to a lesser extent than their

infected counterparts (Figure 3B).

PU.1 protein levels decrease in the presence of Vpr.

Vpr is known to act as an adaptor protein that links host proteins to the Cul4A E3 ubiquitin

ligase complex for ubiquitylation and degradation[16,17]. Thus, we hypothesized that Vpr

downregulates PU.1-regulated genes in infected cells by targeting PU.1 for proteasomal

degradation. To test our hypothesis, we first investigated whether Vpr reduced PU.1 protein

levels in infected MDMs. We treated macrophages with a VSV-G envelope-pseudotyped

A **Selected biological processes associated with PU.1 targeted genes**

B **Expression of HIV and macrophage markers**

C **PU.1 regulated genes involved in TLR and IFN-I signalling**

Figure 3. Vpr counteracts the innate immune response to HIV infection. (A) Selected biological processes associated with infection and inflammation (selected from GO terms represented in Figure 2D). Pathways were identified as associated with PU.1 regulated genes downmodulated in MDMs infected with 89.6^{wt} or 89.6$^{\Delta vpr}$ virus as determined by expression of *gag* and *tat* transcripts (highlighted in Figure 2C). The -log10 FDR-adjusted p-values (q-values) are plotted for each gene ontology term. Blue bars represent terms associated with TLR signaling; Black bars represent related gene ontology terms similar, but not directly associated with TLR signaling. (B) Violin plots summarizing single-cell RNA transcripts expressed by primary human macrophages treated with the indicated virus and cultured for 10 days. *Gag+/tat+* are the subset of cells expressing HIV genes within in each culture. *SPI1* is the gene that codes for PU.1. *MRC1* is the gene that codes for mannose receptor. (C) Violin plots summarizing single-cell RNA transcripts expressed by primary human macrophages as in (B). False-discovery rate corrected Wilcoxon rank-sum p-values are shown above the conditions being compared.

replication-defective clone of HIV NL4-3 that expresses GFP in the *env* reading frame (Figure

4A). We transduced MDMs with this virus containing all intact accessory proteins, or with

additional mutations in the open reading frames of either *vpr, nef,* or both. PU.1 levels were

measured by flow cytometry at 5-, 7-, and 10-days post infection. Notably, we found that Vpr

expression resulted in lower PU.1 protein levels in infected (GFP$^+$) cells but not in virus-exposed

(GFP$^-$) bystander cells, whether Nef was expressed or not (Figure 4B, C). These changes in

PU.1 were consistently observed in MDMs from four independent donors (Figure 4C). Based on

these results, we concluded that Vpr downmodulates PU.1 in HIV-infected primary human

MDMs.

To determine whether Vpr was sufficient for PU.1 downmodulation, we tested the ability of

several 3xFLAG-tagged Vpr proteins derived from a panel of HIV molecular clones [89.6, NL4-3,

AD8 and YU2-Vpr (Figure 4D)] to reduce exogenous PU.1 expression in transfected HEK 293T

cells. Because the gene that encodes PU.1 (*SPI1*) contains PU.1 binding sites and is

downmodulated by Vpr expression (Figure 1B), for these experiments, PU.1 was expressed

under the control of a heterologous promoter that allowed separation of transcriptional and post-

transcriptional changes (Figure 4E). For comparison, we used the evolutionarily related protein,

Vpx, from the HIV2$_{ROD}$ molecular clone. We found that all Vpr-expression constructs tested

resulted in a notably lower level of PU.1 protein compared to PU.1 protein with HIV-2$_{ROD}$Vpx

(Figure 4F). Vpr-dependent reduced levels of PU.1 were consistently observed in four

independent experiments and were statistically significant (Figure 4G).

After determining that Vpr was sufficient to downmodulate PU.1 in HEK 293T cells, we

confirmed these results in a second cell line, the lymphoblastic chronic myelogenous leukemia

cell line K562, which endogenously expresses PU.1. For these experiments Vpr-expression was

achieved using VSV-G envelope pseudotyped lentiviruses. Like the results obtained in HEK

293T cells, infections with viruses expressing all Vprs tested (89.6, NL4-3, AD8, YU2) had lower

endogenous PU.1 levels compared to mock infection and HIV-2$_{ROD}$Vpx infection (Figure 4H).

Because the Vpr-dependent decrease of PU.1 occurred both for endogenous PU.1 expressed

from its native promoter in K562 cells as well as in HEK 293T cells, when PU.1 was expressed

from a heterologous promoter, we hypothesized that Vpr was affecting PU.1 protein levels post-

transcriptionally. To rule out transcriptional effects on the heterologous promoter, we quantified

PU.1 mRNA in HEK 293T cells transfected with a PU.1 expression plasmid and HIV constructs

Figure 4. PU.1 levels decrease in the presence of Vpr. (A) NL4-3 ΔGPE-GFP viral genome map. (B) Representative flow cytometry histogram of PU.1 expression in infected (GFP+) or uninfected bystander (GFP-) MDMs infected with NL4-3 ΔGPE-GFP with or without *vpr* and collected on day 7 post infection. (C) Summary graph showing the percentage of infected (GFP+) cells that do not express PU.1 as determined by flow cytometry as depicted in Figure 4B. The mean +/- standard deviation from n=4 independent donors is shown for each time point. P values were determined using an analysis of variance (ANOVA) with Tukey's multiple comparisons test. *** indicates p = 0.001, **** indicates p < 0.0001. (D) Lentiviral map of vectors encoding 3xFLAG-encoding *vpr* or *vpx* genes. (E) PU.1 expression plasmid map for full-length, human PU.1. (F) Representative flow cytometric plots of HEK 293T cells transiently transfected with PU.1 and the indicated FLAG-tagged viral protein. (G) Summary graph showing PU.1+ expression in transfected (FLAG+) cells. Each point represents the mean of three technical replicates. The mean +/- standard deviation of four independent experiments is shown. P values were determined using one-way ANOVA compared to control, **** indicates p < 0.0001. (H) Immunoblot analysis of PU.1 in K562 cells transduced with the indicated lentivirus expressing 3xFLAG-tagged viral proteins. Results are representative of those from three independent experiments. (I) Summary graph of *SPI1* (the gene encoding PU.1) expression in HEK 293T cells co-transfected with PU.1 and an 89.6 expression vector with or without an intact *vpr* open reading frame. *SPI1* levels were assessed from purified RNA using RT quantitative real time PCR. Results represent the mean fold change compared with wild type +/- standard deviation for samples performed in triplicate for two independent experiments.

derived from 89.6 that had or lacked the Vpr gene (Supplemental Figure 3A). As shown in

Figure 4I, Vpr did not reduce expression of PU.1 mRNA when expressed from a heterologous

promoter in transfected cells. Thus, we concluded that the mechanisms by which Vpr reduced

PU.1 protein in HEK 293T cells must be post-transcriptional. Moreover, in macrophages, Vpr

likely exerts negative effects directly on PU.1 protein and indirectly on PU.1 RNA through PU.1

binding sites in the *SPI1* promoter [50].

Prior reports indicated that PU.1 suppresses HIV gene expression in a Tat-reversible manner[51].

If true, Vpr-targeting of PU.1 could explain the upregulation of HIV genes observed in Vpr-

expressing HIV infected macrophages (Figure 1F). To attempt to confirm these results, we co-

transfected HEK 293T cells with a PU.1 expression plasmid (Figure 4E) and an 89.6-derived

HIV genome that expresses GFP constitutively and mCherry upon HIV activation, allowing us to

distinguish transfected cells through GFP expression and HIV-LTR activity through mCherry

expression (Supplemental Figure 3A). To eliminate any complications from Vpr-mediated

reduction of PU.1, we used an HIV genome that did not express *vpr*. Transfection of increasing

amounts of PU.1 plasmid with the same amount of HIV in all conditions, resulted in increasing

amounts of PU.1 protein as measured by flow cytometry (Supplemental Figure 3B, Left).

However, we failed to confirm a PU.1-dependent suppression of HIV-LTR activity (Supplemental

Figure 3B, Right).

PU.1 downmodulation is a conserved activity of Vprs from HIV-2 and closely related SIV

molecular clones. Vpr is highly conserved in lentiviruses, including HIV-2 and all SIV strains

[52,53]. Interestingly, HIV-2 and certain SIVs contain both Vpr and Vpx, whereas HIV-1 and some

SIVs contain only Vpr (Figure 5A). The extent to which these two proteins harbor overlapping

functions has been a subject of a number of research studies [17,54,55]. Thus, we first examined

whether Vprs from viruses containing both Vpr and Vpx would reduce PU.1 levels. To do this,

we used the HIV-2$_{ROD}$ molecular clone from which our Vpx control is derived. Transient

transfection of HEK 293T cells with our PU.1 expression vector and either HIV-2$_{ROD}$Vpx, HIV-

2$_{ROD}$Vpr, or HIV-1$_{89.6}$Vpr confirmed a significant decrease in PU.1 for cells expressing Vpr^{ROD} or

Vpr$^{89.6}$ but not Vpx^{ROD} (Figure 5B, C). This indicates that Vpx and Vpr from the same molecular

clone have divergent functions with respect to PU.1 downmodulation.

We next tested the ability of Vpr from six isolates evolutionarily similar and disparate to HIV-1 to

determine whether the PU.1 targeting function of Vpr is evolutionarily conserved. Vpr

Figure 5. Vpr-mediated reduction of PU.1 is conserved in HIV-2 and SIV molecular clones. (A) Genomic maps for HIV-1, HIV-2, and select SIV genomes. HIV-1, SIVcpz, and SIVgor genomes contain *vpr* and *vpu* genes but not *vpx*. HIV-2, SIVrcm, and SIVsm contain *vpx* and *vpr*. SIVagm contains *vpr*. (B and D) Representative flow cytometric plots of HEK 293T cells transiently transfected with expression plasmids for PU.1 and the indicated FLAG-tagged viral protein. (C and E) Summary graph of data from B and D respectively. The percentage of PU.1$^+$ cells per transfected (FLAG$^+$) cells is shown. Each point represents the average of three technical replicates. The mean +/- standard deviation is shown for n=6 or 11 independent experiments, respectively. Part E was additionally normalized to HIV-2$_{ROD}$Vpx for each experiment. **** indicates p < 0.0001 using one-way ANOVA compared to control.

sequences from several SIV molecular clones from separate clades, including two from

chimpanzee and one from gorilla (the direct evolutionary relatives of HIV-1), were indeed able to

mediate the degradation of PU.1 in HEK 293T cells (Figures 5D, E). Thus, Vpr-mediated

reduction of PU.1 was consistently observed across all HIV-1, HIV-2, and SIV molecular clones

tested, indicating a strong selective pressure for HIV-related viruses to downmodulate PU.1 in

infected cells.

Both PU.1 and DCAF1 form a complex with Vpr. To determine whether Vpr recruits PU.1

similarly to other host proteins for proteasomal degradation via the CRL4-DCAF1 ubiquitin

ligase complex (Reviewed in [52]), we first assessed whether Vpr and PU.1 formed a complex in

transfected HEK 293T cells. Remarkably, we found that exogenously expressed PU.1 (Figure

4E) efficiently co-precipitated with FLAG-tagged 89.6-Vpr. In addition, and as expected [15,16],

DCAF1 also co-precipitated with Vpr (Figure 6A).

We next examined

whether Vpr co-

precipitated with

endogenous PU.1 in the

K562 cell line originally

derived from a patient

with CML[56]. To

accomplish this, we stably

expressed FLAG-tagged

HIV-1$_{89.6}$Vpr or HIV-

2$_{ROD}$Vpx in K562 cells

using lentiviral vectors

and repeated co-

immunoprecipitation

experiments. We found

that both PU.1 and

Figure 6. PU.1, and DCAF1 co-precipitate with Vpr but not Vpx. (A) Western blot analysis of lysates from HEK 293T cells co-transfected with the indicated expression constructs and immunoprecipitated with an antibody directed against the FLAG epitope. (B) Western blot analysis of lysates from K562 cells transduced with lentiviruses and treated as in part A. For parts A and B, results are representative of three independent experiments.

301 DCAF1 co-precipitated with Vpr to a greater extent than Vpx, again suggesting PU.1 binding is

302 specific to Vpr and not to the similar accessory protein Vpx (Figure 6B). Collectively, these data

303 indicate that Vpr selectively forms complexes with both PU.1 and DCAF1 in cells expressing

304 PU.1 at normal levels from an endogenous promoter.

Interactions between PU.1 and Vpr require DCAF1. To better understand the mechanism

through which Vpr downmodulates PU.1, we assessed a potential requirement for the host

protein DCAF1. DCAF1 (also known as VprBP[15]) is the Vpr binding partner in the CRL4

ubiquitin ligase complex, necessary for proteasomal degradation of Vpr recruited host proteins

(Refs). We generated an HEK 293T cell line with reduced DCAF1 expression. The cell line was

generated using a lentiviral vector stably expressing either a non-targeting shRNA (shScramble)

or a DCAF1-targeting shRNA cassette (shDCAF1) (Supplemental Figure 4; (Figure 7A, Lanes

1-6 compared to 7-12 and lanes 13-18 compared to 19-24, respectively).

Having previously demonstrated that PU.1 and DCAF1 co-precipitate with Vpr (Figure 6), we

performed the IP in reverse and found that Vpr and DCAF1 coprecipitated with a FLAG-tagged-

PU.1 (Figure 7A, Lanes 4 and 10). Because the antibody directed against Vpr does not

recognize 89.6-Vpr, these experiments utilizing un-tagged Vpr were performed using NL4-3,

whose Vpr is recognized by the antibody.

We found that the amount of Vpr that co-precipitated with PU.1 was decreased with DCAF1

silencing (Figure 7A, upper panel, Lane 10 compared to 4), suggesting a role for DCAF1 in the

interaction between Vpr and PU.1. Consistent with this, NL4-3-Vpr^{Q65R} protein, which is

defective for DCAF1 interactions[57,58], did not coprecipitate with PU.1 (Figure 7A, upper panel

Lanes 6 and 12 compared to 4 and 10). These interesting results suggest the unexpected

conclusion that PU.1, Vpr, and DCAF1 may form a trimeric complex and PU.1 binding in the

complex is dependent on both Vpr and DCAF1 (Figure 7D), though Vpr can associate with

DCAF1 independently of PU.1 (Figures 6A, B).

Vpr downmodulates PU.1 via a pathway that depends on proteasome activity.
To further test the relationship between PU.1 and the Vpr/DCAF1 complex, which together are

known to promote ubiquitylation and proteasomal degradation of Vpr-bound host proteins, we

investigated whether inhibition of the proteosome could restore PU.1 levels. As shown in Figure

7B, we were indeed able to restore PU.1 levels in transfected HEK 293T cells expressing Vpr

using MG132, as a specific inhibitor of the proteasome.

**An intact DCAF-1 interaction domain is required for Vpr and DCAF1 to co-precipitate with
TET2 and PU.1.** The PU.1-regulated antiviral factor, *IFITM3* (Figure 2C and E) is also controlled

A HEK 293T Lysates

B HEK 293T Lysates

C HEK 293T Lysates

D Working model of PU.1 and TET2 interacting with Vpr and DCAF1 in macrophages

Figure 7. An intact DCAF-1 interaction domain is required for Vpr and DCAF1 to co-precipitate with TET2 and PU.1. (A) Western blot analysis of lysates from HEK 293T cells stably expressing either control, non-targeting shRNA (scramble) or an shRNA targeting DCAF1 (shDCAF1), transfected with the indicated expression plasmid and immunoprecipitated using an antibody directed against the FLAG epitope. (B) Western blot analysis of lysates from HEK 293T cells transfected with the indicated expression construct and treatment as indicated with 10μM MG132 or vehicle (Veh) control (DMSO). A GFP-expressing plasmid was included where indicated as control for transfection efficiency. (C) Western blot analysis of lysates from HEK 293T cells co-transfected with the indicated expression constructs and immunoprecipitated with an antibody directed against the FLAG epitope. For parts A, B, and C, results are representative of three independent experiments. (D) Working model of PU.1 and TET2 interacting with Vpr and DCAF1 in macrophages.

by TET2, another target of Vpr [36,37]. TET2 is a DNA dioxygenase that de-methylates the *IFITM3*

promoter during viral infection, inhibiting HIV Env trafficking and reducing viral spread[59,60]. To

prevent this, Vpr mediates the ubiquitination and degradation of TET2, which in turn inhibits

IFITM3 expression. PU.1 also regulates myeloid specific gene expression by interacting with

other transcriptional regulatory proteins, including TET2[24,25,31]. We therefore hypothesized that

TET2 could be co-recruited with PU.1 to DCAF1 by Vpr for degradation (Figure 7D). Indeed, we

observed TET2 co-precipitating in HEK 293T cells with FLAG-tagged-PU.1, Vpr, and DCAF1

(Figure 7C). Our findings support a model in which Vpr can dramatically alter the transcriptional

landscape in macrophages by targeting myeloid-specific transcription factors – like PU.1 and its

co-factors – that are required for the expression of key antiviral restriction factors (Figure 8).

Figure 8. Proposed model of the PU.1-mediated antiviral response disrupted by HIV-Vpr. HIV (-) Vpr indicates an infection of primary macrophages with HIV that does not express Vpr. PU.1 protein is maintained and available to regulate anti-viral response genes with co-factors such as IRF or TET2. HIV (+) Vpr indicates an infection with HIV that expresses Vpr. PU.1 protein is less available to regulate anti-viral response genes, contributing to innate immune evasion.

DISCUSSION

Although HIV accessory proteins have been widely studied the critical function that drives

evolutionary conservation of Vpr has remained largely enigmatic. While Vpr significantly

enhances infection of macrophage-containing cultures, primarily by enhancing cell-to-cell

transmission, it does not substantially affect HIV infection in cultures of CD4[+] T cells that lack

macrophages [1–4,34,35,61]. While a number of Vpr targets implicated in post-replication DNA repair

have been identified, a compelling explanation for the dramatic selective effects of Vpr on

infection and spread in macrophage-containing cultures has remained elusive. In this work, we

identified a macrophage-specific target of Vpr, the myeloid transcription factor PU.1 and its

associated co-factors, which Vpr targets, averting antiviral effects (Figure 8). Remarkably, by

utilizing single-cell RNA sequencing of MDMs treated with replication competent virus with and

without Vpr we could distinguish effects of Vpr on cells harboring bona fide infections as well as

bystander cells. This allowed us to detect both cell autonomous and non-autonomous effects

of Vpr.

In infected cells, we found that Vpr reduces the transcription of hundreds of genes regulated

through PU.1 and its cofactors. Several of the PU.1 regulated genes we identified as being

impacted by Vpr are involved in TLR signaling. TLRs are highly conserved PRRs that help cells

identify PAMPs and respond quickly to infection. Activation of TLRs through binding of a PAMP,

initiates a cascade of intracellular signaling that results in the release of inflammatory cytokines

and upregulation of Type I interferon response genes[45,47,62]. The products of these genes have

known antiviral effects on in HIV, and include *ISG15, STMN1, IFI6, LY96, TREM2, FABP4,*

IFITM3, and *MRC1*. *STMN1* is thought to play a role in the establishment of HIV latency, and

depletion leads to higher expression of HIV-1[63]. *MRC1* and *IFITM3* interrupt Env trafficking,

reducing viral spread. *IFI6*, *LY96, and ISG15* are members of the Type I IFN response to

infection[46,64,65] and *ISG15* has also been shown to inhibit HIV spread by disrupting Gag

polymerization[66].

Unexpectedly, we found that bystander cells within HIV infected cultures also responded to HIV

infection by upregulating a subset of these antiviral genes, including *ISG15*, *LY96*, *IFI6*, and

IFITM3. Moreover, we found that upregulation of these factors was reduced in bystander cells

from Vpr-positive versus Vpr-negative HIV-infected primary macrophage cultures. By

comparison, other subsets of genes, including *SPI1* and *MRC1* were primarily

downmodulated in the infected subset and either not at all in the bystander cells or to a

lesser extent than their infected counterparts. Thus, we observed both cell autonomous and

non-autonomous effects of Vpr. Because we only detected significant downmodulation of PU.1

protein in the infected cell subset in our flow cytometric analyses (Figure 2B), Vpr does not

appear to enter and directly affect bystander cells in our cultures. Instead, the detrimental effect

of Vpr on antiviral gene expression in bystander cells likely occurs by an indirect mechanism.

We propose that the inhibitory effects of Vpr on PU.1/IRF gene regulation reduces the

expression of paracrine factors that alert bystander macrophages to the presence of the virus.

Likely candidate genes that could act in this way include type I interferons. Unfortunately, we

were unable to efficiently detect *IFN* RNA expression via single cell RNA sequencing.

Nevertheless, our results indicate that Vpr can exert systemic antiviral affects that favor virus

infection and spread.

We have previously shown that mannose receptor is a host restriction factor that reduces HIV

spread in macrophages by binding to mannose residues on Env and directing Env to the

lysosome for degradation[34]. However, our previous data lacked a mechanism for Vpr-mediated

transcriptional reduction of *MRC1* in macrophages. The data here confirm findings from other

groups that PU.1 regulates *MRC1* expression[42,51], and we provide new evidence that Vpr

reduces the transcriptional expression of *MRC1* in macrophages via PU.1 degradation.

In agreement with other studies[36,37], our unbiased survey of Vpr's effect on expression of host

transcriptome showed that Vpr reduced *IFITM3* gene expression substantially. IFITM proteins

are broad antiviral factors that inhibit viral entry and exit for HIV-1, SIV, MLV, VSV, EBOV,

WNV, among other viruses [67]. Of the IFITM proteins, IFITM3 is well documented as an HIV-1

antagonist. Like mannose receptor, IFITM3 interacts with Env in infected cells to inhibit Env

processing and virion incorporation and strongly inhibits cell-to-cell spread. Interruption of Env

processing by IFITM3 has been demonstrated for several HIV-1 molecular clones[68] many of

which were included within our own study (AD8, YU2, and NL4-3, HIV-2$_{ROD}$, SIVagm, and

SIVcpz). *IFITM3* expression is regulated by the DNA dioxygenase, TET2[36]. During HIV-1

infection of macrophages, in the absence of Vpr, TET2 demethylates the *IFITM3* promoter,

relieving suppression, contributing to the anti-viral response. When Vpr is present, *IFITM3*

expression is reduced after Vpr recruitment of TET2 to DCAF1 for polyubiquitination and

degradation of TET2[37]. Although TET2 is highly expressed in myeloid cells[69], it is ubiquitously

expressed in the nuclei of all cells[70]. How TET2 is specifically recruited to *IFITM3* to regulate

expression of *IFITM3* in macrophages remains unknown. Interestingly, in monocytes, TET2

interacts with PU.1, leading to interaction with genetic targets[31]. Therefore, we hypothesized

that Vpr exerts a macrophage-specific effect on the antiviral response by targeting PU.1 and by

extension, PU.1-associated proteins. Consistent with this hypothesis, we were able to

demonstrate TET2 immunoprecipitating with PU.1, Vpr, and DCAF1 (Figure 7). This provides

evidence that Vpr can re-program cellular transcription in macrophages by targeting myeloid-

specific transcription factors (Figure 1).

Vpr-mediated reduction of PU.1 occurred in all cell types tested in our study including MDM,

K562, and HEK 293T cells. The Vpr-dependent reduced PU.1 levels were observed regardless

of whether PU.1 was expressed from its native promoter or a heterologous promoter. Reversal

of PU.1 downmodulation with proteasome inhibitors in HEK 293T cells supports the conclusion

that Vpr directly reduces PU.1 protein by promoting its degradation. Vpr-mediated degradation

of PU.1 was consistent for all HIV-2, SIV, and HIV-1 Group M isolates tested - Group M being

largely responsible for the global HIV pandemic. In macrophages, *SPI1* mRNA levels encoding

PU.1 protein were also lower in the presence of Vpr, indicating that downmodulation of PU.1

can occur at both the transcriptional and the post-transcriptional level in macrophages.

It is important to note that we failed to confirm prior observations that PU.1 suppresses HIV-LTR

activity in a Tat reversible manner[51]. While it is possible that expression of Tat by our construct

prevented our ability to detect an effect of PU.1 on the HIV-1 LTR, these results nevertheless

indicate that downmodulation of PU.1 by Vpr is unlikely to impact HIV-1 gene expression in

infected macrophages that also express Tat. Thus, in primary HIV infected macrophages, Vpr

mainly targets PU.1 to counteract its anti-viral defense response rather than to counteract an

inhibitory effect on HIV gene expression. However, it is possible that the higher expression of

HIV genes we observed in Vpr-containing cells may result from Vpr counteracting a PU.1-

regulated factor that inhibits HIV gene transcription.

Vpr is highly conserved amongst primate lentiviruses and promotes infection of nondividing

cells, especially macrophages[61]. In addition, a requirement for Vpr to achieve maximal

replication and persistence in vivo was first discovered using an SIV molecular clone in rhesus

monkeys[71]. We therefore speculated that the ability of Vpr to degrade PU.1 is an important

evolutionary function of Vpr. HIV-2 differs from HIV-1 in that it contains Vpx, an accessory

protein that shares a common genetic ancestor with Vpr[53] but is lacking from all HIV-1

genomes. Thus, we tested the relative ability of Vpx and Vpr from the same molecular clone to

promote PU.1 degradation[53]. We were able to identify a unique function of HIV-2 Vpr to degrade

PU.1 that was not shared by Vpx. Similarly, both SIVcpz and SIVgor, the evolutionary

precursors to HIV-1, and Vpr proteins from more evolutionarily distant viruses were able to

consistently result in lower PU.1 levels. However, the greatest decrease in PU.1 levels was

achieved with HIV-1 derived molecular clones. We therefore speculate that the strength of Vpr-

mediated degradation of PU.1 plays an important role in driving spread in human pandemic

strains of HIV-1, and PU.1 may be a critical restriction factor, limiting spread between species

and within populations. However, more extensive studies comparing the relationship between

Vpr and PU.1 across SIV, HIV-2, and HIV-1 isolates is necessary and part of our ongoing

efforts.

The relationship between DCAF1 and Vpr is well documented[15–17]. We confirmed the binding of

Vpr to DCAF1 and identified PU.1 as a new Vpr-binding factor. We demonstrated that all three

components co-precipitate under many different cellular conditions regardless of whether we

pull-down with Vpr or PU.1. To our surprise, mutation of the glutamine residue at position 65 of

Vpr not only resulted in the loss of DCAF1 association but also disrupted formation of the Vpr-

PU.1 complex. Additionally, when we reduced DCAF1 levels, the amount of both Vpr and

DCAF1 precipitating with PU.1 was reduced. These results indicate that interactions amongst all

three proteins are necessary for stable complex formation, although further studies are

necessary to understand the detailed protein-protein interactions. Based on these findings, we

propose a model in which interactions amongst PU.1, Vpr, and DCAF1 promote the

ubiquitylation of PU.1 via the associated CUL4A ubiquitin ligase complex with resultant

proteasomal degradation (Figure 7D). Consistent with this model, inhibition of the proteosome

with MG132 treatment rescued PU.1 levels in the presence of Vpr.

Although the role of Vpr in HIV infection has remained largely undefined, we provide evidence

that the primary selective pressure for Vpr in lentiviruses is to disrupt the macrophage innate

antiviral response to infection, which is achieved by reducing PU.1 levels in infected cells. The

ability of Vpr to degrade PU.1 is highly conserved among all HIV-1, HIV-2, and SIV isolates

tested, and degradation of PU.1 relies on both Vpr and DCAF1. We are continuing to investigate

the transcriptional consequences of PU.1 degradation in macrophages. In addition to TET2,

PU.1 associates with other transcription factors in cells, potentially piggybacking other

secondary targets to the DCAF1-Cul4A E3 Ubiquitin ligase complex via Vpr. Because PU.1 is

necessary to maintain macrophage function, future studies should address the greater impact of

Vpr-mediated reduction of PU.1 on the infected macrophage.

ACKNOWLEDGMENTS

This work was supported by National Institutes of Health grants R01AI149669 to K.L.C. and

J.D.W., F31AI15504 to M.C.V. and F31AI125090-01 to J.L., R21AI32379 to K.L.C, training

grants T32GM007315 to M.C.V. and 5T32GM008353-27 to J.L., and additional funding provided

by the Rackham Regents Fellowship to M.C.V. We thank Chen Li from the Welch lab for his

invaluable advice and aid with single-cell data analysis. We thank the University of Michigan

Flow Cytometry Core and Advanced Genomics Core for assistance with experiments,

particularly Wenpu Trim and Tricia Tamsen. Thank you to all members of the Collins lab for their

helpful discussions.

AUTHOR CONTRIBUTIONS

Conceptualization, M.C.V, J.L., and K.L.C.; Methodology, M.C.V, J.L, J.D.W., and K.L.C.; Software, M.C.V. and J.D.W.; Validation, M.C.V.; Formal Analysis, M.C.V.; Investigation, M.C.V., W.M.D., T.C., and J.L.; Resources, K.L.C.; Writing – Original Draft, M.C.V.; Writing – Review & Editing, M.C.V., J.D.W., and K.L.C.; Visualization, M.C.V.; Supervision, J.D.W. and K.L.C., Funding Acquisition, M.C.V., J.D.W., and K.L.C.

DECLARATION OF INTEREST

The authors declare no competing interest.

METHODS

Cell culture and preparation of human MDMs

All cell cultures were maintained at 37°C in a 5% CO_2 humidified atmosphere. HEK 293Ts (CRL-3216) and K562 (CCL-243) cells were obtained from ATCC and independently authenticated by STR profiling. HEK 293Ts were maintained in DMEM medium (Gibco) supplemented with 100 U/mL penicillin, 100μg/mL streptomycin, 2mM glutamine (Pen-Strep-Glutamine, Invitrogen), and 10% fetal bovine serum (Invitrogen). K562 cell were maintained in IMDM (Gibco) and supplemented as HEK293Ts. Monocyte-derived macrophages (MDMs) were isolated and cultured as previously described[34] and maintained in R10 (RPMI, supplemented as HEK 293Ts). Monocytes were isolated from anonymous donors by apheresis and obtained from the New York Blood Center Component Laboratory. The use of human blood from anonymous, de-identified donors was classified as non-human subject research in accordance with federal regulations and therefore not subject to formal IRB review.

Viruses, viral vectors, and expression plasmids

The molecular clone p89.6 (cat# 3552 from Dr. Ronald G. Collman) was obtained from the AIDS Reagent Program. The *vpr*-null version was generated as previously described[61]. The human, full length PU.1 (hPU.1) expression vector was a gift of Dr. Gregory M. K. Poon and generated as previously described[72]. A triple N-FLAG-tagged version of PU.1 was generated by using PCR amplification of hPU.1 to add a KpnI site to the 5' end and EcoRI to the 3' end (5' GTAGGTACCGCCACCATGGAAGGGTT primer and 3' GTAGAATTCCACCACACTGGACTAGTG primer). The new product replaced an existing gene when inserted into pcDNA3.1 containing a triple N-FLAG-tag between KpnI and EcoRI (Addgene plasmid #67788). The GFP transfection control plasmid, pcDNA3-EGFP, was a gift from Doug Golenbock (Addgene plasmid # 13031; http://n2t.net/addgene:13031; RRID: Addgene_13031). The pNL4-3 ΔGPE-GFP plasmid used for single round infection of macrophages was previously described[73], as were the vpr-null, nef-null, and vpr-nef null versions[34]. pUC19[74] was used as control DNA to adjust transfection samples to the same final DNA concentration. The p89.6-ΔGPEN-mCherry-pSFFV-EGFP single-round infection plasmid was previously described[75] and further modified by replacing *gag* and *pol* with mCherry. Deletion of *vpr* was achieved using Q5 Site-Directed Mutagenesis kit (New England Biolabs) where the majority of the *vpr* coding sequence was deleted using PCR exclusion (forward - CAGAATTGGGTGTCGACATAG, reverse - TCACAGCTTCATTCTTAAGC).

Triple FLAG-tagged Vpr and Vpx lentiviral expression vectors for HIV-2$_{ROD}$Vpx (Addgene

plasmid #115816), SIV$_{SAB-92018}$Vpr (Addgene plasmid #115822), SIV$_{AGM-MAL}$Vpr (Addgene

plasmid #115828), SIV$_{CPZ-TAN3}$Vpr (Addgene plasmid #115833), SIV$_{CPZ-LB7}$Vpr (Addgene plasmid

#115834), SIV$_{gorCP684con}$Vpr (Addgene plasmid #115835), and SIV$_{rcm02CM8081}$Vpr (Addgene

plasmid #115838) were a gift from Jeremy Luban[55]. Similar vectors for HIV-2$_{ROD}$Vpr, HIV-

1$_{89.6}$Vpr, HIV-1$_{NL4-3}$Vpr, HIV-1$_{AD8}$Vpr, HIV-1$_{YU2}$Vpr were generated by synthesizing the gene as a

gBlock (IDT, Coralville, Iowa, USA) between NotI and either EcoRI or AflIII in the same lentiviral

expression plasmid. Untagged expression vectors for HIV-1$_{NL4-3}$Vpr and HIV-1$_{NL4-3}$Vpr^{Q65R} were

generated by synthesizing the genes as a gBlocks (IDT, Coralville, Iowa, USA) and inserting

them between SbfI and NotI in LeGO-IV, a gift from Boris Fehse (Addgene plasmid #27360)[76].

The short hairpin RNA targeting DCAF1 (target sequence: CCTCCCATTCTTCTGCCTTTA) was

designed using Genetic Perturbation Platform (Broad institute) and cloned into pLKO.1 – TRC

cloning vector, a gift from David Root (Addgene plasmid # 10878 ; http://n2t.net/addgene:10878

; RRID:Addgene_10878)[77]. The control shRNA, scramble shRNA was a gift from David Sabatini

(Addgene plasmid # 1864 ; http://n2t.net/addgene:1864 ; RRID:Addgene_1864)[78].

Co-transfections

Co-transfections of 3xFLAG-Vpx/Vpr and hPU.1 were performed in HEK 293T cells. Cells were

plated at 1.6 x 10^5 per well in a 12-well dish. 24hrs after plating, 1ng of hPU.1, 500ng of

3xFLAG-Vpx/Vpr, and pUC19 to a total of 1µg of DNA per well were combined with 4µg of PEI,

mixed and added to each well. 48hrs later, cells were harvested for flow cytometry or

immunoblotting. For co-immunoprecipitations, transfection experiments were scaled to achieve

60 x 10^6 cells per condition. Co-transfections with p89.6-ΔGPERN-mCherry-pSFFV-EGFP and

hPU.1 were performed as described above and with DNA amounts described in the legend.

Transduction of MDM, K562, and HEK 293T

All transductions were performed via spin inoculation at 1050 x g for 2 hr at 25°C with equal

virus amounts determined by Gag p24 mass in medium containing 4µg/mL polybrene (Sigma).

MDMs were inoculated with 10µg p24 mass equivalents of NL4-3 ΔGPE-GFP virus. K562 and

HEK 293Ts were spin inoculated with 10µg p24 mass equivalents of shScramble and shDCAF1

viruses. K562 cells were inoculated with varying amounts of 3xFLAG-tagged Vpr/Vpx

expression viruses to achieve equal FLAG expression. After infection, viral medium was
removed and replaced with fresh medium.

HIV infection of MDM

Prior to infection, 500µL of medium was removed from each well of MDMs and saved to make
diluted conditioned media post-infection. MDM were infected with 5µg, 10µg, and 20µg
equivalents of Gag p24 mass in 500µL total of R10 6 hr at 37°C. After the 6 hr infection, media
was removed and replaced with conditioned media diluted 1:2 in R10. Half-media changes were
performed every three days.

Single-cell RNA sequencing

At 10 days post-infection, uninfected, 89.6^{WT}, and 89.6$^{\Delta vpr}$ infected MDMs were lifted using
enzyme free cell dissociation buffer (ThermoFisher). Replicate samples were fixed with
paraformaldehyde and stained for Gag to assess viral spread by flow cytometry. The resulting
flow cytometry data was used to select 89.6^{WT} and 89.6$^{\Delta vpr}$ conditions with similar percentages
of infected cells. Wells of the selected conditions were harvested, counted, and prepared
according to manufacturer instructions for 10X Chromium Next GEM Single Cell 3' v3 Gene
Expression (10X Genomics).

Single-cell data analysis

10X filtered expression matrices were generated from CellRanger version 3.0.0 (10X
Genomics). We analyzed all single-cell gene expression data using the standard LIGER[38]
(https://github.com/welch-lab/liger) data integration pipeline. All WT and Vpr-null infected MDM
raw data expression matrices from each donor were combined before merging the data. We
used a value of k = 20 during joint matrix factorization, of resolution of 0.05 for Louvain
clustering, and nearest neighbor = 30 with a minimum distance = 0.3 for UMAP visualization.
Infected cells were identified by sub-setting cells with a non-zero expression value for both *gag*
and *tat* transcripts. Differential gene expression between WT and Vpr-null infected MDMs was
determined using the Wilcoxin rank-sum test. Volcano plots of differentially expressed genes
was generated using ggplot2[79] (https://ggplot2.tidyverse.org/). Downregulated genes in the
presence of Vpr with a log2FC > 1 and adjusted p-value > 0.05 were used as input for the
HOMER[80] (http://homer.ucsd.edu/homer/motif/) 'findMotifs' function using the human reference
set. PU.1-motif associated genes were identified using the 'find' function in 'findMotifs' from
HOMER. Gene Ontology analysis for biological processes for PU.1 motif-containing genes was

determined using GOrilla[81,82] (http://cbl-gorilla.cs.technion.ac.il/) with PU.1 regulated genes as

target genes and all expressed genes in our dataset as background. Biological processes were

plotted using REVIGO[83] (http://revigo.irb.hr/). The -500bp sequence for MRC1 used for PU.1

motif scanning was obtained from UCSC Genome Browser[84] (https://genome.ucsc.edu/) using

Human reference genome GRCh38/hg38. The PU.1 binding motif probability matrix was

obtained from HOMER and used with FIMO[85] (https://meme-suite.org/meme/doc/fimo.html) to

scan the *MRC1* input sequence. Violin plots were generated by importing our LIGER generated

dataset into Seurat[86] (https://satijalab.org/seurat/) and running the VlnPlot function. All single-

cell data analysis and plots were done using RStudio[87] (http://www.rstudio.com/) except for

HOMER-predicted motifs.

Virus production

Virus stocks were produced by transfected HEK 293T cells (ATCC, Manassas, Virginia) with

viral DNA and polyethylenimine (PEI) as previously described[34]. For replication defective

constructs, cells were plated 24hrs before transfection with a DNA ratio of 1:1:1 with pCMV-HIV-

1[88], pHCMV-V (VSV-G expression plasmid) (ATCC 75497), and lentiviral expression plasmid.

Viral supernatant was collected two days post-transfection and stored at -80°C. For infectious

virus, pCMV-HIV and pHCMV-V were omitted.

Virion quantification

Viral supernatants were quantified by ELISA as previously described[34] with the following

modification: lysed samples were captured at 4°C overnight.

Immunoblot

Western blots: cells were lysed in Blue Loading Buffer (cat# 7722, Cell Signaling Technology),

sonicated with a Misonix sonicator (Qsonica, LLC. Newtown, CT), boiled for 10min at 95°C

before loading, and analyzed by SDS-PAGE immunoblot.

Co-immunoprecipitation: cells were lysed in Pierce IP Lysis Buffer (Thermo Fisher Scientific)

and 1x Halt Protease Inhibitor Cocktail (Thermo Fisher Scientific). Lysates were incubated with

Anti-FLAG M2 Magnetic Beads (MilliporeSigma, Darmstadt, Germany) according to

manufacturer instructions. Proteins were eluted using 3xFLAG peptide (MilliporeSigma,

Darmstadt, Germany) and analyzed by SDS-PAGE immunoblot. FLAG-tagged proteins were

visualized using Pierce ECL (Thermo Scientific) after treatment with an HRP-conjugated primary

antibody directed against the FLAG epitope (Millipore Sigma). HRP-conjugated secondary

antibodies against murine and rabbit antibodies to other targets (see below) plus ECL Prime

reagent (Cytiva Amersham) were used to visualize all other proteins.

Antibodies

Antibodies to Vinculin (cat# V9131, Millipore Sigma), DCAF1 (cat# 11612-1AP, Proteintech),

PU.1 (cat# 2266S, Cell Signalling Technology), FLAG (cat# F1804, Millipore Sigma), TET2

(cat#MABE462, EMD Millipore), CAp24 (clone KC57-PE cat# 6604667, Beckman Coulter), Vpr

(AIDS Reagent Program cat# 3951 from Dr. Jeffrey Kopp), GFP (cat# ab13970, Abcam) were

used for immunoblot analysis. Secondary HRP conjugated antibodies against murine (rat anti-

mouse IgG1, eBioscience) and rabbit (goat anti-rabbit IgG, Invitrogen) were also used.

Antibodies to PU.1 (clone 7C6B05, BioLegend), FLAG (cat# 637324, BioLegend) were used for

flow cytometry.

Quantitative RT PCR

HEK 293T cells sorted as described in 'Flow Cytometry' below were counted using the

Countess II Cell Counter (Thermo Fisher Scientific) and cell samples were diluted such that all

conditions contained the same cell numbers as input. RNA was isolated using the Zymo

DirectZol RNA MiniPrep Plus extraction kit with an on-column DNaseI digestion. RNA was

reverse transcribed using the High-Capacity cDNA Reverse Transcriptase kit (Applied

Biosystems). Quantitative PCR was performed using SYBR green qPCR Master Mix (Applied

Biosystems) on a QuantStudio 3 Real-Time PCR System (Applied Biosystems) with

ReadyMade PrimeTime primers for *SPI1* (Integrated DNA Technologies Inc, USA) and RT2

qPCR Primer Assay for Human GAPDH (cat# PPH00150F-200, Qiagen). Expression was

quantified using ABI Sequence Detection software compared to serial dilutions of an SPI1 or

GAPDH synthetic sequence gBlock (Integrated DNA Technologies Inc, USA). Measured values

for *SPI1* were normalized to measured values of *GAPDH*.

Flow cytometry

For cells requiring intracellular staining using antibodies directed against HIV Gag p24, FLAG-

Vpx and -Vpr, and PU.1, paraformaldehyde-fixed cells were permeabilized with 0.1% Triton-X in

PBS for 2min followed by incubation with antibody for 30min at room temperature.

In all experiments, cells were gated sequentially by forward scatter vs. side scatter for cells and

then by forward scatter area vs. height to exclude doublets. All transduced MDMs and

transiently transfected HEK 293T cells were assessed for protein expression on the Cytek

Aurora. GFP+ HEK 293T cells in the MG132 treatment experiments and for qPCR were sorted

on the Sony SH800 cell sorter into R10. Untreated, GFP- cells were also sorted. All flow

cytometry data was analyzed using FlowJo software (TreeStar, Ashland, Oregon).

Proteasome inhibition

Lyophilized MG132 was purchased from MilliporeSigma (cat# M8699) and dissolved in DMSO.

MG132 was added to cellular medium to achieve a final concentration of 10µM at varying

timepoints. Cells were harvested from replicate wells such that all drug treatment timepoints

were collected at once. DMSO only control treatment wells (Vehicle) were treated with the same

volume of DMSO as contained in the MG132 treatment conditions.

Statistical Analysis

All non-single cell statistical analyses were performed using GraphPad Prism software as

described in figure legends for each experiment.

 FIGURE LEGENDS

Figure 1. Single-cell RNA sequencing of HIV-1 infected MDMs reveals Vpr-dependent

transcriptional changes. (A) Schematic diagram illustrating the objective of single-cell RNA

sequencing; the identification of Vpr-targeted transcription factor(s) in HIV infected primary

macrophages. (B) Genome maps for full-length 89.6^{wt} HIV-1 (top) and the same viral genome

with a premature stop-codon in *vpr* (bottom). (C) Experimental setup for the generation of the

scRNA-seq datasets. (D) UMAP representation of LIGER-integrated scRNA-seq data from

MDM samples treated as shown in (C) and listed in Dataset. (E) UMAP representations of

LIGER-integrated scRNA-seq data from MDMs treated as indicated. (E, Right) Bona fide

infected cells were identified based on expression of HIV *tat* and *gag*. (E, left) Colors indicate

individual clusters. Cluster 0 = pro-inflammatory macrophages, Cluster 1 = cycling cells, Cluster

2 = dying cells, Cluster 3 = pro-inflammatory macrophages. (E, center and right) cells colored

according to whether they were exposed to 89.6 WT-Vpr virus (blue) or Vpr-null virus (pink). (F)

Volcano plot of differentially expressed genes from HIV 89.6^{wt} verses 89.6$^{\Delta vpr}$ infected MDMs

from (E) as determined by Wilcoxin Rank Sum. Significance determined as greater than 1 log2

fold change and false discovery rate adjusted p-value of p < 0.05 (red-bars). Blue colored genes

indicate genes less highly expressed in HIV 89.6^{wt} verses 89.6$^{\Delta vpr}$ infected MDMs, Red colored

genes indicate genes more highly expressed, and black colored genes indicate no significant

difference between datasets.

Figure 2. Vpr downmodulates PU.1-dependent transcription. (A) PU.1 motifs identified by

HOMER as present in the promoters of Vpr-downmodulated genes (Figure 1E, Blue). (B) Violin

plots displaying RNA abundance of the indicated transcription factor genes in MDMs infected

with the indicated virus. (C) Volcano plot as in Figure 1E except that genes containing a PU.1 or

PU.1-IRF binding motif in their promoter region are highlighted in green (See also Supplemental

Figure 2). (D) Biological processes associated with the PU.1 targeted genes from (B). Size of

circle indicates the relative number of GO terms with the process. FDR adjusted q-values

associated with GO terms as indicated by the color. (E) Violin plots displaying RNA abundance

of the indicated genes in MDMs infected with the indicated virus. HIV-1 89.6^{wt} (WT); HIV-1

89.6$^{\Delta vpr}$ (ΔVpr).

Figure 3. Vpr counteracts the innate immune response to HIV infection. (A) Selected

biological processes associated with infection and inflammation (selected from GO terms

represented in Figure 2D). Pathways were identified as associated with PU.1 regulated genes

downmodulated in MDMs infected with 89.6^{wt} or 89.6$^{\Delta vpr}$ virus as determined by expression of

gag and *tat* transcripts (highlighted in Figure 2C). The -log10 FDR-adjusted p-values (q-values)

are plotted for each gene ontology term. Blue bars represent terms associated with TLR

signaling; Black bars represent related gene ontology terms similar, but not directly associated

with TLR signaling. (B) Violin plots summarizing single-cell RNA transcripts expressed by

primary human macrophages treated with the indicated virus and cultured for 10 days.

Gag+/tat+ are the subset of cells expressing HIV genes within in each culture. *SPI1* is the gene

that codes for PU.1. *MRC1* is the gene that codes for mannose receptor. (C) Violin plots

summarizing single-cell RNA transcripts expressed by primary human macrophages as in (B).

False-discovery rate corrected Wilcoxon rank-sum p-values are shown above the conditions

being compared.

Figure 4. PU.1 levels decrease in the presence of Vpr. (A) NL4-3 ΔGPE-GFP viral genome

map. (B) Representative flow cytometry histogram of PU.1 expression in infected (GFP$^+$) or

uninfected bystander (GFP$^-$) MDMs infected with NL4-3 ΔGPE-GFP with or without *vpr* and

collected on day 7 post infection. (C) Summary graph showing the percentage of infected

(GFP$^+$) cells that do not express PU.1 as determined by flow cytometry as depicted in Figure

4B. The mean +/- standard deviation from n=4 independent donors is shown for each time point.

P values were determined using an analysis of variance (ANOVA) with Tukey's multiple

comparisons test. *** indicates p = 0.001, **** indicates p < 0.0001. (D) Lentiviral map of

vectors encoding 3xFLAG-encoding *vpr* or *vpx* genes. (E) PU.1 expression plasmid map for full-

length, human PU.1. (F) Representative flow cytometric plots of HEK 293T cells transiently

transfected with PU.1 and the indicated FLAG-tagged viral protein. (G) Summary graph showing

PU.1$^+$ expression in transfected (FLAG$^+$) cells. Each point represents the mean of three

technical replicates. The mean +/- standard deviation of four independent experiments is

shown. P values were determined using one-way ANOVA compared to control, **** indicates p

< 0.0001. (H) Immunoblot analysis of PU.1 in K562 cells transduced with the indicated lentivirus

expressing 3xFLAG-tagged viral proteins. Results are representative of those from three

independent experiments. (I) Summary graph of *SPI1* (the gene encoding PU.1) expression in

HEK 293T cells co-transfected with PU.1 and an 89.6 expression vector with or without an intact

vpr open reading frame. *SPI1* levels were assessed from purified RNA using RT quantitative

real time PCR. Results represent the mean fold change compared with wild type +/- standard

deviation for samples performed in triplicate for two independent experiments.

Figure 5. Vpr-mediated reduction of PU.1 is conserved in HIV-2 and SIV molecular clones.

(A) Genomic maps for HIV-1, HIV-2, and select SIV genomes. HIV-1, SIVcpz, and SIVgor

genomes contain *vpr* and *vpu* genes but not *vpx*. HIV-2, SIVrcm, and SIVsm contain *vpx* and

vpr. SIVagm contains *vpr*. (B and D) Representative flow cytometric plots of HEK 293T cells

transiently transfected with expression plasmids for PU.1 and the indicated FLAG-tagged viral

protein. (C and E) Summary graph of data from B and D respectively. The percentage of PU.1$^+$

cells per transfected (FLAG$^+$) cells is shown. Each point represents the average of three

technical replicates. The mean +/- standard deviation is shown for n=6 or 11 independent

experiments, respectively. Part E was additionally normalized to HIV-2$_{ROD}$Vpx for each

experiment. **** indicates p < 0.0001 using one-way ANOVA compared to control.

Figure 6. PU.1, and DCAF1 co-precipitate with Vpr but not Vpx. (A) Western blot analysis of

lysates from HEK 293 T cells co-transfected with the indicated expression constructs and

immunoprecipitated with an antibody directed against the FLAG epitope. (B) Western blot

analysis of lysates from K562 cells transduced with lentiviruses and treated as in part A. For

parts A and B, results are representative of three independent experiments.

Figure 7. An intact DCAF-1 interaction domain is required for Vpr and DCAF1 to co-

precipitate with TET2 and PU.1. (A) Western blot analysis of lysates from HEK 293T cells

stably expressing either control, non-targeting shRNA (scramble) or an shRNA targeting DCAF1

(shDCAF1), transfected with the indicated expression plasmid and immunoprecipitated using an

antibody directed against the FLAG epitope. (B) Western blot analysis of lysates from HEK

293T cells transfected with the indicated expression construct and treatment as indicated with

10μM MG132 or vehicle (Veh) control (DMSO). A GFP-expressing plasmid was included where

indicated as control for transfection efficiency. (C) Western blot analysis of lysates from HEK

293T cells co-transfected with the indicated expression constructs and immunoprecipitated with

an antibody directed against the FLAG epitope. For parts A, B, and C, results are representative

of three independent experiments. (D) Working model of PU.1 and TET2 interacting with Vpr

and DCAF1 in macrophages.

Figure 8. Proposed model of the PU.1-mediated antiviral response disrupted by HIV-Vpr.

HIV (-) Vpr indicates an infection of primary macrophages with HIV that does not express Vpr.

PU.1 protein is maintained and available to regulate anti-viral response genes with co-factors

such as IRF or TET2. HIV (+) Vpr indicates an infection with HIV that expresses Vpr. PU.1

protein is less available to regulate anti-viral response genes, contributing to innate immune

evasion.

Supplemental Figure legends

Supplemental Table 1. MDM infection rates measured with Gag. Percent infection of 89.6^{wt}

and 89.6$^{\Delta vpr}$ infected MDMs from each of three donors was determined by quantifying the

percent Gag$^+$ cells by either flow cytometry (protein) or gene expression levels from scRNA-seq

data (mRNA), as indicated by the experiment type and molecule, over the total number of cells

analyzed.

Supplemental Figure 1. Transcription factors and their predicted binding motifs in

downregulated genes in MDMs expressing Vpr. Motifs, transcription factor names,

significance (FDR q-value), and number of target genes identified by HOMER analysis of genes

downregulated in the presence of Vpr (Figure 1E) from several transcription factor families.

Supplemental Figure 2. The *MRC1* promoter contains multiple PU.1 binding motifs. The

first 500 bp upstream of the start codon in *MRC1*, the gene that codes for mannose receptor.

Previously reported PU.1 binding motifs are outlined in red. The double-box PU.1 motif was

identified through inputting the HOMER generated PU.1 motif parameters into FIMO (Find

Individual Motif Occurrences). The TATA box is outlined in blue, the 5'UTR in yellow, and the

start codon in green.

Supplemental Figure 3. PU.1 does not alter HIV-LTR activity. (A) Genomic map for 89.6-

derived HIV-1 fluorescent reporter virus (89.6-ΔGPERN-pSFFV-EGFP). (B) Representative flow

plots and bar graph assessing transfection rate via GFP expression in HEK 293T cells

transfected with the indicated amount of PU.1 construct from Figure 2E plus 500ng of 89.6-

ΔGPERN-pSFFV-EGFP from (A). (C) Summary graph of PU.1 expression in GFP$^+$ cells from

(B). PU.1 levels were assessed using intracellular staining as described in Methods and

measured by flow cytometry. (D) Summary graph of HIV-LTR activity as assessed by mCherry

MFI from the same cells as in (C), measured flow cytometrically. All conditions were transfected

with 500ng of HIV expression plasmid plus 0, 1, 10, 100, or 500ng of PU.1 plasmid. MFI = mean

fluorescence intensity.

Supplemental Figure 4. DCAF1 protein is reduced in K562 cells via shRNA. Immunoblot of

lysates from K562 cells stably transduced with virus expressing a non-targeting shRNA

(shScramble) or one targeting DCAF1 (shDCAF1) compared to un-transduced cells (Mock).

REFERENCES

1. Balliet, J. W. *et al.* Distinct Effects in Primary Macrophages and Lymphocytes of the Human

Immunodeficiency Virus Type 1 Accessory Genes vpr, vpu, and nef: Mutational Analysis of a

Primary HIV-1 Isolate. *Virology* **200**, 623–631 (1994).

2. Eckstein, D. A. *et al.* HIV-1 Vpr enhances viral burden by facilitating infection of tissue

macrophages but not nondividing CD4+ T cells. *J. Exp. Med.* **194**, 1407–1419 (2001).

3. Connor, R. I., Chen, B. K., Choe, S. & Landau, N. R. Vpr Is Required for Efficient Replication of

Human Immunodeficiency Virus Type-1 in Mononuclear Phagocytes. *Virology* **206**, 935–944

(1995).

4. Dedera, D., Hu, W., Vander Heyden, N. & Ratner, L. Viral protein R of human

immunodeficiency virus types 1 and 2 is dispensable for replication and cytopathogenicity

in lymphoid cells. *J. Virol.* **63**, 3205–3208 (1989).

5. Rücker, E., Grivel, J.-C., Münch, J., Kirchhoff, F. & Margolis, L. Vpr and Vpu Are Important for

Efficient Human Immunodeficiency Virus Type 1 Replication and CD4+ T-Cell Depletion in

Human Lymphoid Tissue Ex Vivo. *J. Virol.* **78**, 12689–12693 (2004).

6. Lu, Y. L., Spearman, P. & Ratner, L. Human immunodeficiency virus type 1 viral protein R

localization in infected cells and virions. *J. Virol.* **67**, 6542–6550 (1993).

7. Laguette, N. *et al.* Premature Activation of the SLX4 Complex by Vpr Promotes G2/M Arrest

and Escape from Innate Immune Sensing. *Cell* **156**, 134–145 (2014).

8. Yan, J., Shun, M. C., Zhang, Y., Hao, C. & Skowronski, J. HIV-1 Vpr counteracts HLTF-

mediated restriction of HIV-1 infection in T cells. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 9568–

9577 (2019).

9. Ahn, J. *et al.* HIV-1 Vpr Loads Uracil DNA Glycosylase-2 onto DCAF1, a Substrate Recognition

Subunit of a Cullin 4A-RING E3 Ubiquitin Ligase for Proteasome-dependent Degradation *. *J.

Biol. Chem.* **285**, 37333–37341 (2010).

10. Lahouassa, H. *et al.* HIV-1 Vpr degrades the HLTF DNA translocase in T cells and

macrophages. *Proc. Natl. Acad. Sci.* **113**, 5311–5316 (2016).

11. Withers-Ward, E. S., Mueller, T. D., Chen, I. S. Y. & Feigon, J. Biochemical and Structural

Analysis of the Interaction between the UBA(2) Domain of the DNA Repair Protein HHR23A

and HIV-1 Vpr. *Biochemistry* **39**, 14103–14112 (2000).

12. Wang, X. *et al.* HIV-1 Vpr Protein Inhibits Telomerase Activity via the EDD-DDB1-VPRBP E3

Ligase Complex *. *J. Biol. Chem.* **288**, 15474–15480 (2013).

13. Romani, B., Baygloo, N. S., Aghasadeghi, M. R. & Allahbakhshi, E. HIV-1 Vpr Protein

Enhances Proteasomal Degradation of MCM10 DNA Replication Factor through the Cul4-

DDB1[VprBP] E3 Ubiquitin Ligase to Induce G2/M Cell Cycle Arrest *. *J. Biol. Chem.* **290**,

17380–17389 (2015).

14. Yan, J. *et al.* HIV-1 Vpr Reprograms CLR4DCAF1 E3 Ubiquitin Ligase to Antagonize

Exonuclease 1-Mediated Restriction of HIV-1 Infection. *mBio* **9**, e01732-18 (2018).

15. Zhang, S., Feng, Y., Narayan, O. & Zhao, L.-J. Cytoplasmic retention of HIV-1 regulatory

protein Vpr by protein-protein interaction with a novel human cytoplasmic protein VprBP.

Gene **263**, 131–140 (2001).

16. McCall, C. M. *et al.* Human Immunodeficiency Virus Type 1 Vpr-Binding Protein VprBP, a

WD40 Protein Associated with the DDB1-CUL4 E3 Ubiquitin Ligase, Is Essential for DNA

Replication and Embryonic Development. *Mol. Cell. Biol.* **28**, 5621–5633 (2008).

17. Romani, B. & Cohen, É. A. Lentivirus Vpr and Vpx accessory proteins usurp the cullin4–DDB1

(DCAF1) E3 ubiquitin ligase. *Curr. Opin. Virol.* **2**, 755–763 (2012).

18. Hakata, Y., Miyazawa, M. & Landau, N. R. Interactions with DCAF1 and DDB1 in the CRL4 E3

ubiquitin ligase are required for Vpr-mediated G2 arrest. *Virol. J.* **11**, 108 (2014).

19. Schröfelbauer, B., Hakata, Y. & Landau, N. R. HIV-1 Vpr function is mediated by interaction

with the damage-specific DNA-binding protein DDB1. *Proc. Natl. Acad. Sci. U. S. A.* **104**,

4130–4135 (2007).

20. Belzile, J.-P. *et al.* HIV-1 Vpr-Mediated G2 Arrest Involves the DDB1-CUL4AVPRBP E3

Ubiquitin Ligase. *PLoS Pathog.* **3**, e85 (2007).

21. Hrecka, K. *et al.* HIV-1 and HIV-2 exhibit divergent interactions with HLTF and UNG2 DNA

repair proteins. *Proc. Natl. Acad. Sci. U. S. A.* **113**, E3921–E3930 (2016).

22. Lavin, Y., Mortha, A., Rahman, A. & Merad, M. Regulation of macrophage development and

function in peripheral tissues. *Nat. Rev. Immunol.* **15**, 731–744 (2015).

23. Silvin, A. & Manel, N. Innate immune sensing of HIV infection. *Curr. Opin. Immunol.* **32**, 54–

60 (2015).

24. Turkistany, S. A., Dekoter, R. P. & Hirszfeld, Ó. L. The Transcription Factor PU.1 is a Critical

Regulator of Cellular Communication in the Immune System. doi:10.1007/s00005-011-

0147-9.

25. Fisher, R. C. & Scott, E. W. Role of PU.1 in Hematopoiesis. *Stem Cells* **16**, 25–37 (1998).

26. Dakic, A. *et al.* PU.1 regulates the commitment of adult hematopoietic progenitors and

restricts granulopoiesis. *J. Exp. Med.* **201**, 1487–1502 (2005).

27. Gupta, P., Gurudutta, G. U., Saluja, D. & Tripathi, R. P. PU.1 and partners: regulation of

haematopoietic stem cell fate in normal and malignant haematopoiesis. *J. Cell. Mol. Med.*

13, 4349–4363 (2009).

28. Graves, B. J. & Petersen, J. M. Specificity within the ets Family of Transcription Factors. in

Advances in Cancer Research (eds. Vande Woude, G. F. & Klein, G.) vol. 75 1–57 (Academic

Press, 1998).

29. Marecki, S. & Fenton, M. J. PU.1/interferon regulatory factor interactions. *Cell Biochem.*

Biophys. **33**, 127–148 (2000).

30. Imperato, M. R., Cauchy, P., Obier, N. & Bonifer, C. The RUNX1–PU.1 axis in the control of

hematopoiesis. *Int. J. Hematol.* **101**, 319–329 (2015).

31. de la Rica, L. *et al.* PU.1 target genes undergo Tet2-coupled demethylation and DNMT3b-

mediated methylation in monocyte-to-osteoclast differentiation. *Genome Biol.* **14**, 1–21

(2013).

32. Liang, G. *et al.* Membrane metalloprotease TRABD2A restricts HIV-1 progeny production in

resting CD4 + T cells by degrading viral Gag polyprotein. *Nat. Immunol.* (2019)

doi:10.1038/s41590-019-0385-2.

33. Linehan, S. A., Martínez-Pomares, L., Stahl, P. D. & Gordon, S. Mannose Receptor and Its

Putative Ligands in Normal Murine Lymphoid and Nonlymphoid Organs: In Situ Expression

of Mannose Receptor by Selected Macrophages, Endothelial Cells, Perivascular Microglia,

and Mesangial Cells, but not Dendritic Cells. *J. Exp. Med.* **189**, 1961–1972 (1999).

34. Lubow, J. *et al.* Mannose receptor is an HIV restriction factor counteracted by Vpr in

macrophages. *eLife* **9**, (2020).

35. Collins, D. R., Lubow, J., Lukic, Z., Mashiba, M. & Collins, K. L. Vpr Promotes Macrophage-

Dependent HIV-1 Infection of CD4+ T Lymphocytes. *PLoS Pathog.* **11**, e1005054 (2015).

36. Wang, Q. & Su, L. Vpr Enhances HIV-1 Env Processing and Virion Infectivity in Macrophages

by Modulating TET2-Dependent IFITM3 Expression. *mBio* **10**, e01344-19 (2019).

37. Lv, L. *et al.* Vpr Targets TET2 for Degradation by CRL4VprBP E3 Ligase to Sustain IL-6

Expression and Enhance HIV-1 Replication. *Mol. Cell* **70**, 961-970.e5 (2018).

38. Welch, J. D. *et al.* Single-Cell Multi-omic Integration Compares and Contrasts Features of

Brain Cell Identity. *Cell* **177**, 1873–1887 (2019).

39. Zhang, F. & Bieniasz, P. D. HIV-1 Vpr induces cell cycle arrest and enhances viral gene

expression by depleting CCDC137. *eLife* **9**, e55806 (2020).

40. Pham, T. H. *et al.* Mechanisms of in vivo binding site selection of the hematopoietic master

transcription factor PU.1. *Nucleic Acids Res.* **41**, 6391–6402 (2013).

41. Buckland, J. Haematopoiesis: PU.1 pinned down. *Nat. Rev. Immunol.* **2**, 149 (2002).

42. Caldwell, R. L., Egan, B. S. & Shepherd, V. L. HIV-1 Tat Represses Transcription from the

Mannose Receptor Promoter. *J. Immunol.* **165**, 7035–7041 (2000).

43. Medzhitov, R., Preston-Hurlburt, P. & Janeway, C. A. A human homologue of the Drosophila

Toll protein signals activation of adaptive immunity. *Nature* **388**, 394–397 (1997).

44. Molteni, M., Gemma, S. & Rossetti, C. The Role of Toll-Like Receptor 4 in Infectious and

Noninfectious Inflammation. *Mediators Inflamm.* **2016**, 6978936 (2016).

45. Beutler, B. A. TLRs and innate immunity. *Blood* **113**, 1399–1407 (2009).

46. Perng, Y. C. & Lenschow, D. J. ISG15 in antiviral immunity and beyond. *Nat. Rev. Microbiol.*

16, 423–439 (2018).

47. Schneider, W. M., Chevillotte, M. D. & Rice, C. M. Interferon-Stimulated Genes: A Complex

Web of Host Defenses. *Annu. Rev. Immunol.* **32**, 513–545 (2014).

48. Shimazu, R. *et al.* MD-2, a Molecule that Confers Lipopolysaccharide Responsiveness on

Toll-like Receptor 4. *J. Exp. Med.* **189**, 1777–1782 (1999).

49. Nagai, Y. *et al.* Essential role of MD-2 in LPS responsiveness and TLR4 distribution. *Nat.*

Immunol. **3**, 667–672 (2002).

50. Chen, H. *et al.* PU.1 (Spi-1) autoregulates its expression in myeloid cells. *Oncogene* **11**,

1549–1560 (1995).

51. Kao, S. *et al.* The Myeloid-Specific Transcription Factor PU.1 Upregulates Mannose Receptor

Expression but Represses Basal Activity of the HIV-LTR Promoter. *J. Virol.* **96**, e00652-22

(2022).

52. Collins, D. R. & Collins, K. L. HIV-1 accessory proteins adapt cellular adaptors to facilitate

immune evasion. *PLoS Pathog.* **10**, e1003851 (2014).

53. Sharp, P. M. & Hahn, B. H. Origins of HIV and the AIDS Pandemic. *Cold Spring Harb.*

Perspect. Med. **1**, a006841 (2011).

54. Sakai, Y. *et al.* Expression Profiles of Vpx/Vpr Proteins Are Co-related with the Primate

Lentiviral Lineage. *Front. Microbiol.* **7**, (2016).

55. Yurkovetskiy, L. *et al.* Primate immunodeficiency virus proteins Vpx and Vpr counteract

transcriptional repression of proviruses by the HUSH complex. *Nat. Microbiol.* **3**, 1354–1361

(2018).

56. Lozzio, C. B. & Lozzio, B. B. Human Chronic Myelogenous Leukemia Cell-Line With Positive

Philadelphia Chromosome. *Blood* **45**, 321–334 (1975).

57. Zhao, L.-J., Mukherjee, S. & Narayan, O. Biochemical mechanism of HIV-I Vpr function.

Specific interaction with a cellular protein. *J. Biol. Chem.* **269**, 15577–15582 (1994).

58. DeHart, J. L. *et al.* HIV-1 Vpr activates the G2 checkpoint through manipulation of the

ubiquitin proteasome system. *Virol. J.* **4**, 57 (2007).

59. Lu, J., Pan, Q., Rong, L., Liu, S.-L. & Liang, C. The IFITM Proteins Inhibit HIV-1 Infection. *J.*

Virol. **85**, 2126–2137 (2011).

60. Compton, A. A. *et al.* IFITM Proteins Incorporated into HIV-1 Virions Impair Viral Fusion and

Spread. *Cell Host Microbe* **16**, 736–747 (2014).

61. Mashiba, M., Collins, D. R., Terry, V. H. & Collins, K. L. Vpr overcomes macrophage-specific

restriction of HIV-1 Env expression and virion production. *Cell Host Microbe* **16**, 722–735

(2014).

62. Kumar, H., Kawai, T. & Akira, S. Toll-like receptors and innate immunity. *Biochem. Biophys.*

Res. Commun. **388**, 621–625 (2009).

63. Deletsu, S. D. *et al.* Identification and characterization of Stathmin 1 as a host factor

involved in HIV-1 latency. *Biochem. Biophys. Res. Commun.* **567**, 106–111 (2021).

64. Park, G.-H. *et al.* Association between Interferon-Inducible Protein 6 (IFI6) Polymorphisms

and Hepatitis B Virus Clearance. *Genomics Inform.* **11**, 15–23 (2013).

65. Del Cornò, M. *et al.* HIV-1 gp120 signaling through TLR4 modulates innate immune

activation in human macrophages and the biology of hepatic stellate cells. *J. Leukoc. Biol.*

100, 599–606 (2016).

66. Okumura, A., Lu, G., Pitha-Rowe, I. & Pitha, P. M. Innate antiviral response targets HIV-1

release by the induction of ubiquitin-like protein ISG15. *Proc. Natl. Acad. Sci.* **103**, 1440–

1445 (2006).

67. Tartour, K. *et al.* Interference with the production of infectious viral particles and bimodal

inhibition of replication are broadly conserved antiviral properties of IFITMs. *PLOS Pathog.*

13, e1006610 (2017).

68. Yu, J. *et al.* IFITM Proteins Restrict HIV-1 Infection by Antagonizing the Envelope

Glycoprotein. *Cell Rep.* **13**, 145–156 (2015).

69. Cull, A. H., Snetsinger, B., Buckstein, R., Wells, R. A. & Rauh, M. J. Tet2 restrains

inflammatory gene expression in macrophages. *Exp. Hematol.* **55**, 56-70.e13 (2017).

70. Lorsbach, R. B. *et al.* TET1, a member of a novel protein family, is fused to MLL in acute

myeloid leukemia containing the t(10;11)(q22;q23). *Leukemia* **17**, 637–641 (2003).

71. Lang, S. M. *et al.* Importance of vpr for infection of rhesus monkeys with simian

immunodeficiency virus. *J. Virol.* **67**, 902–912 (1993).

72. Munde, M. *et al.* Structure-dependent inhibition of the ETS-family transcription factor PU.1

by novel heterocyclic diamidines. *Nucleic Acids Res.* **42**, 1379–1390 (2014).

73. McNamara, L. A., Ganesh, J. A. & Collins, K. L. Latent HIV-1 infection occurs in multiple

1002   subsets of hematopoietic progenitor cells and is reversed by NF-κB activation. *J. Virol.* **86**,

1003   9337–50 (2012).

1004   74. Norrander, J., Kempe, T. & Messing, J. Construction of improved M13 vectors using

1005   oligodeoxynucleotide-directed mutagenesis. *Gene* **26**, 101–106 (1983).

1006   75. Carter, C. C. *et al.* HIV-1 infects multipotent progenitor cells causing cell death and

1007   establishing latent cellular reservoirs. *Nat. Med.* **16**, 446–451 (2010).

1008   76. Weber, K., Bartsch, U., Stocking, C. & Fehse, B. A multicolor panel of novel lentiviral 'gene

1009   ontology' (LeGO) vectors for functional gene analysis. *Mol. Ther. J. Am. Soc. Gene Ther.* **16**,

1010   698–706 (2008).

1011   77. Moffat, J. *et al.* A Lentiviral RNAi Library for Human and Mouse Genes Applied to an Arrayed

1012   Viral High-Content Screen. *Cell* **124**, 1283–1298 (2006).

1013   78. Sarbassov, D. D., Guertin, D. A., Ali, S. M. & Sabatini, D. M. Phosphorylation and Regulation

1014   of Akt/PKB by the Rictor-mTOR Complex. *Science* **307**, 1098–1101 (2005).

1015   79. Wickham, H. *ggplot2*. (Springer, 2009). doi:10.1007/978-0-387-98141-3.

1016   80. Heinz, S. *et al.* Simple Combinations of Lineage-Determining Transcription Factors Prime cis-

1017   Regulatory Elements Required for Macrophage and B Cell Identities. *Mol. Cell* **38**, 576–589

1018   (2010).

1019   81. Eden, E., Navon, R., Steinfeld, I., Lipson, D. & Yakhini, Z. GOrilla: a tool for discovery and

1020   visualization of enriched GO terms in ranked gene lists. *BMC Bioinformatics* **10**, 48 (2009).

1021   82. Eden, E., Lipson, D., Yogev, S. & Yakhini, Z. Discovering Motifs in Ranked Lists of DNA

1022   Sequences. *PLOS Comput. Biol.* **3**, e39 (2007).

1023   83. Supek, F., Bošnjak, M., Škunca, N. & Šmuc, T. REVIGO Summarizes and Visualizes Long Lists

1024   of Gene Ontology Terms. *PLOS ONE* **6**, e21800 (2011).

1025   84. Kent, W. J. *et al.* The Human Genome Browser at UCSC. *Genome Res.* **12**, 996–1006 (2002).

1026   85. Grant, C. E., Bailey, T. L. & Noble, W. S. FIMO: scanning for occurrences of a given motif.

1027   *Bioinformatics* **27**, 1017–1018 (2011).

1028   86. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29

1029   (2021).

1030   87. RStudio Team. RStudio: Integrated Development for R. (2020).

1031   88. Gasmi, M. *et al.* Requirements for Efficient Production and Transduction of Human

1032   Immunodeficiency Virus Type 1-Based Vectors. *J. Virol.* **73**, 1828–1834 (1999).

1033

Donor	Experiment Type	Molecule	Percent Infected	
			89.6 WT vpr	89.6 vpr-null
1	Flow Cytometry	Protein	65%	50%
1	scRNA-seq	mRNA	71%	48%
2	Flow Cytometry	Protein	45%	31%
2	scRNA-seq	mRNA	53%	28%
3	Flow Cytometry	Protein	52%	35%
3	scRNA-seq	mRNA	N/A	(a) 42%
3	scRNA-seq	mRNA	N/A	(b) 54%

Supplemental Table 1. MDM infection rates measured with Gag. Percent infection of 89.6[wt] and 89.6[Δvpr] infected MDMs from each of three donors was determined by quantifying the percent Gag+ cells by either flow cytometry (protein) or gene expression levels from scRNA-seq data (mRNA), as indicated by the experiment type and molecule, over the total number of cells analyzed.

Supplemental Figure 1

ETS Family Transcription Factors

Motif	TF Name	q-value	# Targets w/ Sequence
	ETV4	0.0000	1270
	ETV1	0.0000	1228
	ETS1	0.0000	987
	Fli1	0.0000	1229
	Elk4	0.0000	1038
	Elf4	0.0000	930
	Elk1	0.0000	1013
	ERG	0.0000	1135
	GABPA	0.0000	995
	ELF1	0.0000	925
	EHF	0.0000	876
	ETS	0.0000	674
	ELF5	0.0000	557
	Etv2	0.0000	768
	EWS	0.0001	327
	PU.1	0.0005	316
	ELF3	0.0011	417
	SpiB	0.0038	180
	EWS:FLI1-fusion	0.0063	534
	ETS:RUNX	0.0082	121
	SPDEF	0.0121	624

ETS/IRF Family Transcription Factors

Motif	TF Name	q-value	# Targets w/ Sequence
	PU.1-IRF	0.0004	670

IRF Family Transcription Factors

Motif	TF Name	q-value	# Targets w/ Sequence
	IRF8	0.0006	209
	IRF2	0.0049	75
	IRF1	0.0453	68
	T1ISRE	0.0504	8

ZF Transcription Factors

Motif	TF Name	q-value	# Targets w/ Sequence
	Sp5	0.0000	1786
	Sp2	0.0000	2047
	KLF1	0.0000	1607
	Sp1	0.0000	1125
	KLF5	0.0000	1787
	KLF3	0.0000	1089
	YY1	0.0000	203
	KLF6	0.0000	1497
	KLF14	0.0000	2044
	Klf9	0.0001	780
	Maz	0.0012	1644
	Klf4	0.0056	531
	GFY	0.0183	200
	ZNF143|ST	0.0567	275

bHLH Family Transcription Factors

Motif	TF Name	q-value	# Targets w/ Sequence
	E-box	0.0000	212
	Usf2	0.0000	290
	MITF	0.0000	505
	TFE3	0.0000	160
	CLOCK	0.0000	402
	USF1	0.0002	371
	c-Myc	0.0006	495
	bHLHE41	0.0036	888
	bHLHE40	0.0041	280
	BMAL1	0.0043	679
	n-Myc	0.0150	432
	Max	0.0228	347
	NPAS	0.0254	612
	HIF2a	0.0636	246

Homeobox Family Transcription Factors

Motif	TF Name	q-value	# Targets w/ Sequence
	Hoxa13	0.0043	647
	HOXB13	0.0183	292
	Hoxd13	0.0183	1
	Nkx3.1	0.0255	746
	Hoxa9	0.0355	796

bZIP Family Transcription Factors

Motif	TF Name	q-value	# Targets w/ Sequence
	CRE	0.0000	385
	Atf1	0.0043	434
	CREB5	0.0632	203

NRF Family Transcription Factors

Motif	TF Name	q-value	# Targets w/ Sequence
	NRF1	0.0000	544
	NRF	0.0003	1

CCAAT Family Transcription Factors

Motif	TF Name	q-value	# Targets w/ Sequence
	NFY	0.0000	782

THAP Family Transcription Factors

Motif	TF Name	q-value	# Targets w/ Sequence
	Ronin	0.0227	185

Supplemental Figure 1. Transcription factors and their predicted binding motifs in downregulated genes in MDMs expressing Vpr. Motifs, transcription factor names, significance (FDR q-value), and number of target genes identified by HOMER analysis of genes downregulated in the presence of Vpr (Figure 1E) from several transcription factor families.

MCR1 promoter has both predicted and known PU.1 binding motifs

-500

AGAAAGGCTCTAAGCACTGAATGTGGAAACTGAAGGGGATGAGCTTCAACTCTGAAGTGTTTCCAGCGTAAAACT
GTCCTTTCCAGGGCCCGTGTGGCTGTCACTTCAGAGTGGAGGTTGTCTGCTGAGGGACCCCTGACTCAGCTGC
TTCCCAGGGGAAGCTCCGTCTTCCGGCACAGGTAATGGCCTGCAGCTTGATCTCCACCCAGCCCCATCTGAGCA
GGCCGGGAGCTCCCAGGCTGTTTCACTTCTCTCCTTCCTGACTCCTCACCATCACCATCGCCCTCTCTCCTCCC
CACCCCGCCACTCCTCTCCCACACGTGTCCCTTTCTCCCCTTCCTCTGCGTCTGCTCTTCTCAGAAGTTAGCTTA
CGAAGCAAAGTTGTTACTTTGAATTCCTGTTTTTCCAGCCACCCTCATGTGACAGGATGTCTCCTCAGTAGAGGCT
TTCCCTAAATTCAGGAGCCCTTTAAAAGGGAGGGCTTCCTCTGTAGTTCTTTTCAGCTGGGCAGCTCTGGGAACT
TGGATTAGGTGGAGAGGCAGTTGGGGGGCCTCGTTGTTTTGCGTCTTAGTTCCGCCCTCCTGTCCATCAGGAGA
AGGAAAGGATAAACCCTGGGCCATG
　　　　　　　　　　　+1

PU.1 Binding Motif　TATA Box　5'UTR　Start Codon

Supplemental Figure 2. The *MRC1* promoter contains multiple PU.1 binding motifs. The first 500 bp upstream of the start codon in *MRC1*, the gene that codes for mannose receptor. Previously reported PU.1 binding motifs are outlined in red. The double-box PU.1 motif was identified through inputting the HOMER generated PU.1 motif parameters into FIMO (Find Individual Motif Occurrences). The TATA box is outlined in blue, the 5'UTR in yellow, and the start codon in green.

A Genomic map of p89.6-ΔGPERN-pSFFV-EGFP

Supplemental Figure 3. PU.1 does not alter HIV-LTR activity. (A) Genomic map for 89.6-derived HIV-1 fluorescent reporter virus (89.6-ΔGPERN-pSFFV-EGFP). (B) Representative flow plots and bar graph assessing transfection rate via GFP expression in HEK 293T cells transfected with the indicated amount of PU.1 construct from Figure 2E plus 500ng of 89.6-ΔGPERN-pSFFV-EGFP from (A). (C) Summary graph of PU.1 expression in GFP+ cells from (B). PU.1 levels were assessed using intracellular staining as described in Methods and measured by flow cytometry. (D) Summary graph of HIV-LTR activity as assessed by mCherry MFI from the same cells as in (C), measured flow cytometrically. All conditions were transfected with 500ng of HIV expression plasmid plus 0, 1, 10, 100, or 500ng of PU.1 plasmid. MFI = mean fluorescence intensity.

K562 Lysates

shRNA

Vinculin

DCAF1

Supplemental Figure 4. DCAF1 protein is reduced in K562 cells via shRNA. Immunoblot of lysates from K562 cells stably transduced with virus expressing a non-targeting shRNA (shScramble) or one targeting DCAF1 (shDCAF1) compared to un-transduced cells (Mock).