Mixing selects for predation resistance in lab-evolved communities of bacterial prey and

social predator *Myxococcus xanthus*

Saheli Saha[1], Bhoomika Bhat[1,2], Jessica Mellicent Laloo[1], Akshit Goyal[3], Samay Pande[1]

[1]Bacterial Ecology and Evolution Group, Department of Microbiology and Cell Biology, Indian Institute

of Science, Bangalore, India.

[2]Undergraduate program, Indian Institute of Science, Bangalore, India.

[3]International Centre for Theoretical Sciences (ICTS), Tata Institute of Fundamental Research,

Bangalore, India.

Corresponding author: Samay Pande

13   Address: Bacterial Ecology and Evolution Group, Department of Microbiology

14   and Cell Biology, Indian Institute of Science, Bangalore, India

15   Email: samayrp@gmail.com

17   Saheli Saha

18   Address: Bacterial Ecology and Evolution Group, Department of Microbiology

19   and Cell Biology, Indian Institute of Science, Bangalore, India

20   Email: sahelisaha@iisc.ac.in

22   Keywords: *M. xanthus*, bacterial predation, coevolution, predator-prey, population-mixing, Red-

23   Queen hypothesis, evolutionary arms race.

Abstract

Population mixing, and transmission modes strongly influence social interactions. However, the impact of repeated mixing on the evolutionary dynamics of microbial predator-prey interactions remains underexplored[1,2]. Hence, we conducted a laboratory evolution experiment with bacterial predator-prey communities under two transfer regimens: repeated mixing (horizontal transfer) versus no mixing (vertical transfer). For this, *Myxococcus xanthus* served as the generalist predator[3,4] and *Escherichia* coli as prey. We show that prey populations from vertical regimen were less resistant to predation than the ones from horizontal regimen. This was because prey isolates in the vertical treatment showed varying resistance levels relative to ancestors, while those in the horizontal treatment demonstrated less variation. Moreover, analysis of prey populations over evolutionary time revealed that the populations from horizontal treatment showed increasing levels of resistance to predation over time, whereas the ones from vertical treatment did not show similar trend. The differences in the outcomes of the two treatments was because the variants better at intraspecies competition, can only be maintained in the vertical treatment, whereas in horizontal treatment the benefits of superior intraspecies competitive fitness are nullified because of population mixing, as predicted by mathematical modeling approach. These predictions were empirically confirmed. Moreover, contrary to our expectations, predatory efficiency of evolved *M. xanthus* isolates was either the same or lesser than the ancestor. Together, we demonstrate that mixing affects the evolution of prey bacteria but has little effect on the hunting ability of the social predator *M. xanthus*.

Introduction:

The influence of mixing on the evolution of microbial interactions and their effects on community

assembly has garnered significant interest in recent times [1,2]. This interest seems justified, given that

the mixing has been shown to strongly affect the evolution of antagonistic as well as synergistic

interactions within populations (i.e., groups of individuals of single species) [1,2,5-9]. Microbial

communities in nature are frequently mixed because of human activities, abiotic, and biotic factors [10-

12]. However, despite the prevalence of mixing events, we have limited understanding of the effects of

repeated mixing on the evolution of microbial interactions both in the context of simple communities

such as predator-prey interaction, and complex multispecies communities [1,2,13].

Evolutionary trajectories of antagonistically interacting species, like host-parasite and predator-

prey systems follow the Red Queen (RQ) dynamics in which interacting partners coevolve in an

evolutionary arms-race [14-16]. The RQ hypothesis has gained significant prominence over the years and

has been proposed to explain the evolution of sex, antagonist-driven emergence of diversity,

symbiosis, and antimicrobial resistance [8,17,18]. Originally proposed by Van Valen (1973) [19], the Red

Queen hypothesis is intentionally focused only on the influence of the biotic factor. However, in

nature, a combination of both biotic and abiotic factors will likely influence the outcome of

antagonistic evolution. Therefore, extensions of this theory that incorporate both the effects of biotic

and abiotic factors, such as diversity, migration, and changes in environmental conditions have

attracted considerable attention[7]. For example, studies using host-parasite interactions suggest a

significant effect of population mixing on the Red Queen dynamics between co-evolving antagonists

[18,20]. While predator-prey coevolutionary dynamics has been the focus of multiple studies in the past

[21-24], surprisingly the influence of repeated mixing, across habitats or even ecosystem, on the predator-

prey coevolutionary dynamics is relatively less explored among microbes [1].

Here we asked whether repeated mixing of distinct microbial predator-prey consortia will affect

their coevolutionary dynamics. For within-species interactions, theory and empirical evidence suggest

that the coevolutionary dynamics without mixing events are likely to promote synergistic interactions

[25]. In contrast, repeated mixing of microbial consortia, that are growing independent of each other

are likely to promote antagonism [25]. Hence, we predicted that for interspecies interactions (such as

predator-prey) too, repeated mixing will result in the evolution of predators that show better

predatory performance relative to the ones that evolved in communities that did not experience

repeated mixing events. Similarly, prey populations that experience repeated mixing events will

evolve to be more resistant to predation than when predator-prey communities are propagated

independent of mixing events. This is because of differences in the partner-fidelity between the two

treatments. We also hypothesized that the populations from mixed (horizontal) and unmixed (vertical)

treatments would follow distinct evolutionary strategies. This is because the mixing events will
increase the supply of mutation which might affect the fixation probabilities (by selecting larger effect
mutations) because of changes in the clonal interference. To test these hypotheses, bacterial
predator-prey communities of *M. xanthus* and *E. coli* were propagated in a lab evolution experiment.

M. xanthus is a generalist bacterial predator that kills prey bacteria by both contact-dependent and
contact-independent mechanisms [4,26-30]. This includes the production of enzymes [27,31], antibiotics [3],
and secretion systems [28,30]. In the presence of *M. xanthus,* coevolving *E. coli* populations have been
shown to diversify and maintain predation-resistant mucoid variants along with the predation-
sensitive types over the evolutionary timescale [24,32]. Further, as expected in a winnerless Red Queen
dynamic, *M. xanthus* evolves higher predatory performance[24].

Here, we demonstrate that coevolution with a social microbial-predator results in an asymmetric
arms-race where only the prey continues to evolve better survival strategies over evolutionary time
in response to predation. Further, we show that the mixing of predator-prey populations during each
transfer results in the evolution of resistance to predation. Whereas prey populations that do not
experience mixing events maintained greater diversity in prey phenotype based on their varying
degree of resistance to predation, indicating maintenance of diverse strategies. Most interestingly,
evolved populations of predators from both regimens show the emergence of within-population
diversity which harbors genotypes with similar predatory performance as their ancestors and the
genotypes with reduced predatory performance relative to their ancestors. The mathematical
modelling approach revealed that the differences in the outcomes between the two treatments are
because of changes in the clonal interference caused by mixing events. These predictions were further
confirmed by performing a laboratory evolution experiment.

Results:
Repeated mixing results in increased resistance to predation in the evolved prey population
The evolution experiment designed to test the effect of mixing on the evolution of prey and
predator populations involved repeated mixing of predator-prey consortia every fourth day
(Supplementary figure S1) for eighty days. Whereas control populations were propagated without
mixing. To perform these experiments glucose was used as a carbon source. Since *E. coli* can use
glucose as a carbon source but *M. xanthus* cannot, predation on *E. coli* population was the only source
of nutrients for *M. xanthus*. The evolved communities after sixty-four days of serial transfers were
analysed for the changes in the resistance to predation in the evolved *E. coli* and changes in the
predatory performance of evolved *M. xanthus*.

To test the effect of repeated mixing, we measured the growth of evolved (after sixteen rounds of
propagation in the lab evolution experiment) and ancestral *E. coli* populations when cocultured with
ancestral *M. xanthus* in conditions identical to the ones during the evolution experiment. This measure
accounts for the growth and survival of prey in the presence of *M. xanthus*. In line with our
expectations, *E. coli* populations from the horizontal regimen on average exhibit better growth in the
presence of predators than their ancestors (1.09 log-fold relative to ancestor, one sample t-test, $p^{[FDR}$
$^{corrected]}$ < 0.001). Whereas growth of populations from vertical treatment did not show a significant
difference relative to the ancestor (1.03 log-fold relative to ancestor, one sample t-test, $p^{[FDR\ corrected]}$ >
0.05). Thus, *E. coli* population from vertical treatment were in general less resistant to predation than
the ones from horizontal treatment (Figure 1A, independent sample t-test, p = 0.057). However, one
of the four vertically coevolved populations (population 3) did evolve higher levels of resistance to
predation, which was comparable to horizontally transmitted lines (Figure 1A, Supplementary figure
S2). Together, vertically propagated prey populations seem to evolve different mechanisms of survival
in the presence of predators, which in most cases increases the sensitivity of the prey populations to
predators.
Differences between the vertically coevolved and horizontally coevolved prey populations can
emerge in response to the presence of predators or as an outcome of adaptation to abiotic factors.
However, since mono-evolved *E. coli* populations (propagated in the absence of *M. xanthus,*
Supplementary figure S1C and S1D) from the vertical as well as horizontal regimen showed no
difference in the overall levels of resistance to predation (Figure 1B and Supplementary figure S2, one
sample Wilcoxon test, p > 0.05), we concluded that the differences between vertical and horizontal *E.*
coli populations propagated along with *M. xanthus* were not the result of adaptation to abiotic factors.
Together, these results demonstrate a strong effect of the transfer regimen on the evolution of the
prey population.
138
Evolution of reduced predatory performance in co-evolved *M. xanthus* populations
It is difficult to reliably measure the growth of *M. xanthus* in our experimental setup (See methods).
Hence, to assess changes in the evolved predator populations, the growth of ancestral prey was used
as a measure of the predatory performance of *M. xanthus*. This measure of predation is a combination
of growth and death of *E. coli* in the presence of the predator. Contrary to our expectations and
previously reported findings [24], we observed that the ancestral *E. coli* was able to grow to a similar
density in the presence of evolved as well as ancestral *M. xanthus* populations (Figure 1C, and
Supplementary figure S3, one sample t-test p > 0.05). Further, no significant differences between
coevolved *M. xanthus* populations (after sixteen transfers) from horizontal and vertical regimens

(Figure 1C, independent sample t-test for differences between prey growth in the presence of evolved
predators, $p > 0.3$) were observed. Together, the results suggest that the predator populations were
probably under relatively weaker selection to evolve enhanced hunting ability. Thus, adaptive
response other than the increased predatory performance as measured in our experiments might
have been selected in our experiments.

It is expected that the evolved populations are internally diverse and hence population-level data
might not be reflective of the intra-population diversity. Hence, we measured the predatory
performance of distinct clones isolated from evolved *M. xanthus* populations. To do so, four *M.*
xanthus clones were randomly isolated from each of the replicate lines from the vertical regimen and
sixteen clones were isolated from the horizontal regimen. Surprisingly, a minority of the clones from
both the transfer regimens showed reduced hunting performance relative to the ancestors
(Supplementary figure S3, ANOVA and SNK *post hoc* test). Whereas the majority showed no
difference.

However, genomic analysis of the evolved clones revealed mutations specific to the co-evolved
clones that were absent in the ancestors. Most of these mutations were either located in the genes
that putatively code for non-ribosomal peptide synthetases (NRPS) or transporter proteins
(Supplementary table T4). Previous reports have demonstrated that NRPS is involved in the synthesis
of antimicrobial peptides and hence is a major component of predatory behavior of *M. xanthus* [33,34].
Thus, mutation in this locus is in line with the selection pressure during the evolution experiment.
While in the bad predators (clones with lower predatory efficiency) most consistent mutations were
in the genes that putatively code for chemotaxis proteins (Supplementary table T4)**.** Mutations in the
chemotactic pathways like in the CheA gene, which is a part of the Che3 cluster of chemotaxis proteins
in *M. xanthus*, also explain reduced swarming efficiencies of the *M. xanthus* clones that have evolved
reduced predatory efficiency (Supplementary table T4). The evolution experiments were conducted
in a spatially structured environment, thus *M. xanthus* cells might need the ability to find the next
patch of prey populations. Thus, chemotactic mutants that cannot swarm towards the prey show
lower prey-killing efficiencies. Interestingly, despite the similarities in the predatory efficiencies of the
predators from both regimens, the overall mutations were more similar in the horizontally coevolved
M. xanthus isolates, compared to the vertically coevolved ones (Supplementary table T4). These
results therefore strongly suggest that the predators' populations have also evolved differentially.
However, the differential evolution has not affected the prey-killing efficiency of the predator
population**.**

Coevolved *E. coli* clones from vertical regimen exhibit large within-population differences in their sensitivity to *M. xanthus* predation than the ones from horizontal treatment

Since evolved prey populations from the two treatments were significantly different from each other, we further tested whether the population level differences were representative of the phenotypes of distinct clones within each population. To do so, four *E. coli* clones were isolated from each of the vertically propagated lines and sixteen clones were isolated from the horizontally evolved population. We observed significant variation among the isolates derived from vertically coevolved populations than between clones isolated from horizontal transfer regimen (Supplementary figure S4, average variance among vertical clones = 0.414, average variance across horizontal clones = 0.130, independent sample t-test for differences between variance $p < 0.02$). Importantly, out of sixteen clones from the vertical regimen, five clones seem to have higher resistance and four clones seem to show lower resistance to predation relative to ancestors (Figure 2 and Supplementary figure S4). Whereas all clones from the horizontal regimen were more resistant to predation than the ancestors.

The evolution of *E. coli* strains sensitive to *M. xanthus* predation in vertically evolved populations suggested that the evolution of resistance to predation might not be the only adaptive strategy. This was indeed the case, as pairwise competition experiments between evolved *E. coli* clones and their respective Kanamycin-marked fitness-neutral ancestors (Supplementary figure S8) revealed that irrespective of their sensitivity to *M. xanthus* predation, most coevolved clones showed higher competitive fitness than ancestors in the presence of *M. xanthus* (Figure 3, one sample Wilcoxon test $p^{[FDR\ corrected]} < 0.0005$). Moreover, the fitness of co-evolved clones was in general significantly higher in the presence of *M. xanthus* than in the absence of it (Figure 3, Kruskal Wallis and Dunn's *post hoc* test).

Similar experiments with strains from mono-evolved lines of *E. coli* showed that all the clones had lower resistance against *M. xanthus* than their ancestors (Figure 2, an average of prey survival of mono-evolved clones was 1.03-fold lower than ancestors, one sample Wilcoxon test, p < 0.0003). Further, we did not detect differences in the relative fitness of mono-evolved *E. coli* in the presence of *M. xanthus* than in its absence (Figure 3, Kruskal Wallis, Dunn's *post hoc*).

Further analysis of the data from predation experiments (Figure 2) and competition experiments (Figure 3) revealed a positive correlation between resistance to predation and relative fitness for vertically coevolved clones of *E. coli* (Supplementary figure S5, Pearson's correlation coefficient (r) = 0.6, p < 0.019), but not for *E. coli* coevolved in the horizontal regimen. Thus, though the evolution of high levels of resistance to predation is not essential for competitive advantage against ancestors, strains with higher resistance will generally have higher fitness than the ones with lower levels of resistance to predation.

Genomic analysis of the evolved and ancestral prey clones from both coevolved and mono-evolved treatments revealed several mutations that are specific to coevolved prey clones (Supplementary table T3). Amongst these, insertions at different locations in and after fliPQR operon in RcsB binding site were most common [35]. This binding site is important for the regulation of RcsA gene that is involved in Rcs-phosphorelay[35]. Previous studies have demonstrated a diverse set of functions for Rcs signal transduction system that include regulation of motility, capsule formation and virulence [35-37]. A detailed understanding of how these mutations affect the sensitivity of prey bacteria in our experiments will require exhaustive analysis that is not part of the current manuscript. Further, a higher genetic diversity was observed in the vertically coevolved prey, that is consistent with their phenotypic diversity. Whereas the horizontally coevolved prey was more genetically similar with comparable mutations in the majority of the clones.

Mathematical modelling reveals prey evolution in horizontal treatment is mainly driven by adaptation to the presence of predator rather than intraspecific competition between prey

Together, our experiments demonstrate that horizontally propagated populations, which were mixed, only evolved predator avoidance, while vertically propagated ones, with no mixing, evolved both predator avoidance and prey competitive ability (Figure 2 and Figure 3). To explain why mixing suppressed prey competitive ability and amplified predator avoidance, we formulated a simple mathematical model of our communities. Briefly, our model partitioned each growth cycle into two phases: an initial period of prey growth, and a later period of predator growth and prey death (Figure 4A), consistent with experimental data of the dynamics within a growth cycle (Supplementary figure S6).

The relative fitness of prey in each of the two phases operated along two distinct measures of fitness in our model. During prey growth, prey fitness was determined primarily by prey competition, i.e., the growth rate of each prey subpopulation relative to the others (Figure 4B). During predator growth, prey fitness was set primarily by predator avoidance, i.e., the death rate of each prey subpopulation (Fig. 4B and Methods for details of the model). Thus, the fitness for each prey subpopulation was determined by two parameters. Our model simulated population dynamics over these growth cycles for a population comprising several prey subpopulations whose prey competition (growth rate) and predator avoidance (death rate) were randomly picked from an appropriate distribution (details in Methods). To simulate mixing (horizontal) and no mixing (vertical) regimens, we mimicked our experimental protocol (Methods).

Simulations of our model reproduced our experimental observations: mixed populations showed greater predator avoidance and lower relative prey competitive ability (Figure 4C and 4D).

Further, the model explained why mixing suppressed prey competition. The key idea is that mixing
increases the number of competing prey subpopulations, which increases clonal interference (Figure
4D). This slows down the takeover of prey with a growth advantage and favors prey with improved
predator avoidance. Thus, our model suggests that mixing increases clonal interference among
competing prey, potentially explaining the different experimental outcomes in vertical and horizontal
regimens.

Prey populations from vertical and horizontal treatment show distinct evolutionary dynamics

Together, empirical observations and theoretical predations suggest that the prey populations in
the vertical transfer regimen are under fluctuating selection. This was because during each growth
cycle interference competition against conspecifics was under selection during the first growth phase
and resistance to predation was under selection in the second growth phase (Figure 4A). This resulted
in the emergence of distinct genotypes with different degrees of resistance to *M. xanthus* predation
in vertical treatment. Since the benefits of better intra-prey population competitive ability during the
growth phase were limited in the horizontal treatment, prey populations from the horizontal regimen
were probably under directional selection, which selects for lesser phenotypic diversity. If true,
horizontally evolved prey populations should over evolutionary time keep on either increasing or
maintaining high levels of resistance against *M. xanthus*. While vertically co-evolved prey populations
are likely to not show a trend of increasing resistance to predation over evolutionary time.

To test this hypothesis, we isolated *M. xanthus* populations after twenty rounds of coevolution.
These *M. xanthus* populations were then used to test the resistance of evolved *E. coli* populations
from three distinct evolutionary time points (i.e., after ten, sixteen and twentieth transfer). In line
with our hypothesis, we observed that after sixteen transfers, populations of horizontally coevolved
prey were on average 1.06 log-fold more resistant to predators than the ones from the tenth transfer.
Further, though not statistically significant, differences were observed between prey at the sixteenth
and twentieth transfer, prey at the sixteenth transfer on average seemed more sensitive to predation
than the ones after twenty transfers (Figure 5A, ANOVA and Tukey's post-hoc). Whereas the vertically
coevolved prey did not show differences in the levels of resistance to predation from transfer-ten to
transfer-twenty (Figure 5A, ANOVA and Tukey's post-hoc). These data, coupled with our observations
that within-population variation among clones from the horizontally coevolved prey populations is
less (Figure 2), and that vertically evolved prey populations are internally heterogeneous (i.e., show
diversity in predation resistance phenotypes) (Figure 2), supports our hypothesis that the horizontally
co-evolved prey populations tend to either continuously evolve higher level of resistance of maintain

the levels of resistance over evolutionary time. Whereas vertically co-evolved prey populations evolve

two strategies, i.e. evolution of superior intraspecies competitive fitness and resistance to predation.

However, when *M. xanthus* population isolated from different evolutionary time points (i.e., tenth,

sixteenth and twentieth) were cultured with *E. coli* population isolated after the twentieth transfer,

no significant differences were observed in the predatory ability of the predators over evolutionary

time from both the treatments. Here, the survival of terminally coevolved *E. coli* (CFU/mL) in the

presence of predators from distinct evolutionary timepoints within respective treatments, was taken

as a read-out of the predatory abilities of respective predator population (Figure 5B, ANOVA and

Tukey's post-hoc)

Switching the mixing regimen between the terminally evolved horizontal and vertical populations

Predictions from our model suggest that the distinct outcomes in the two selection regimens

depend on the continuation of the selection regime. Thus, if mixing indeed resulted in directional

selection in prey survival strategies, repeating the same evolution experiment again with the

diversified populations from the vertical regimen should result in one or both of the following

possibilities. First, reduced variation among clones after being propagated in the mixing treatment.

And second, increased resistance to predation both at the population and isolate levels. To test these

hypotheses, we propagated a terminally coevolved population (i.e., after twenty transfers) by

Switching the transfer regimen - i.e., populations from the horizontal regimen were propagated

vertically, and the vertically evolved populations were propagated in the horizontal regimen -

(Supplementary figure S7). Similar to the experiments above, prey populations were isolated from the

evolved prey-predator consortia (See methods) after they were propagated for sixteen transfers. On

testing the ability to survive predation in the evolved preys against mixing, not only evolved a

significantly higher ability to survive predation than their ancestors (Figure 6A, ANOVA and Tukey's

post-hoc) but also exhibited a lower within-population diversity than their ancestors (Figure 6B,

ANOVA and Tukey's post-hoc for difference between groups and also for differences within each of

the groups). Moreover, consistent with our previous results, while the overall resistance to predation

did not change much at a population level (Figure 6A, ANOVA and Tukey's Post-hoc), in the prey

population that was propagated under vertical regimen exhibited a greater degree of within-

population variation than their ancestors (Figure 6B, ANOVA and Tukey's post-hoc for difference

between groups and also for differences within each of the groups). Together, these experiments

reveal that irrespective of preexisting standing diversity, prey populations that experience repeated

mixing will evolve to have reduced within-population diversity and higher resistance than the ones

that evolved in the absence of mixing events.

Discussion

Discussion

Events that affect the mixing of communities are common in nature [1,9,13,38-40]. However, their

influence on the evolution of bacterial predator-prey interaction remains less explored [1]. Here we

used model predator-prey communities of bacteria to show that repeated mixing of predator-prey

communities can result in the evolution of enhanced prey survival ability in the presence of the

predator, relative to when communities are allowed to evolve in the absence of repeated mixing

events. Surprisingly though, we could not detect increased predatory performance in the evolved

populations of predators. Moreover, contrary to expectations we observed the evolution of reduced

predatory performances in certain predator isolates. Hence the observations reported in our study do

not follow the coevolutionary pattern of the evolutionary arms-race as expected from host-parasite,

and host-pathogen interactions [16,20,41]. Several factors such as social predation by *M. xanthus*, inter-

vs intra-species competition, differences in the effective mutation supply in the two selection

regimens, and asymmetry in the rate of evolution of prey and predator can explain the observed

outcomes.

Two aspects of *M. xanthus* biology can introduce intra-species competition among predators. First,

M. xanthus is a social predator that uses diffusible compounds to kill prey [3,4,29,42]. Moreover, digestion

of the prey is done extracellularly [27,43]. Hence, both the killing factors, as well as the digested prey, are

available as public goods to the individuals in the neighborhood. In our experiments, we find that most

of the individual isolates from both treatments show similar predatory performance as their

ancestors, except for a few clones, that perform worse than their ancestors. Hence, it seems possible

that the apparent decrease in population productivity is mediated by the exploitation of public goods

by the non-producers. If this is true, the use of non-social predators in a similar experimental setup

should result in the evolution of increased predatory performance. Indeed, this is the case, predator-

prey coevolution experiments using non-social *M. xanthus* isolates have resulted in enhanced

predatory performance relative to the ancestral variants [24]. Though not the focus of the current

manuscript, it is an interesting opportunity to test the interaction dynamics between the coevolved

efficient predator and coevolved less-efficient predator. Such population studies coupled with

mechanistic understanding might reveal ecological and evolutionary mechanisms that lead to the

evolution of scavenging strategies. That is the evolution of reduced prey-killing ability coupled with a

strong dependence on the presence of efficient predators for better survival.

Microbial antagonism is pervasive in complex microbial communities [44,45]. Hence, studies of such

interactions have attracted tremendous interest in recent years to combat microbial infections and

contaminations [46-48], and to better the understanding of microbial ecology [24,32,49,50]. To our surprise,

studies have used the terms host-parasite and predator-prey interactions interchangeably for

identical interactions (for example bacteria and phage) [38,51,52]. However, predator-prey interactions

are distinct from host-parasite interactions in many ways [51,54-58] (for example: relationship intimacy,

relationship duration, generation time, population size, and trophic level). Among them we feel two

are relatively more significant than others. First, the population size of the exploiter and exploited is

inverse for predator-prey interactions relative to that of the host-parasite interactions. Second, the

generation time of the predator and prey is similar, whereas the generation time of parasites is

significantly smaller than their hosts. Thus, the exploited partner has a faster rate of evolution in

predator-prey interactions, which is the opposite for host-parasite interactions where the exploiter

has a faster rate of evolution relative to the exploited partner. Here, we did not observe the evolution

of increased predation efficiency among *M. xanthus* populations, whereas bacteria-phage (host-

parasite) coevolution experiments have repeatedly shown the evolution of increased virulence [1,20, 53].

The differences we find here relative to the ones reported in the previous studies [1,19,52] in which the

exploiter (parasite) evolves faster might be attributed to the differences in the relative rate of

evolution of exploiter to the exploited partner.

The effect of population mixing has been studied in multiple instances [1,59]. Whereas the effect of

mixing of predator-prey communities has not been studied in as much detail. Interestingly, we

observed that the evolution of prey bacteria followed a similar fate to previous reports in which the

effect of population mixing was studied on the evolution of bacterial host-parasite interaction [1]. Albeit

with one key difference, i.e., unlike in the host-parasite evolution experiment in which vertical

propagation either selected for resistance to phage in some replicate lineages or intraspecies

competitive fitness in others, we observed that both of these strategies simultaneously evolved within

each of the replicate population. Further, we observed that repeated mixing resulted in directional

selection on the evolution of resistance against the exploiting partner. And unmixed communities, like

in previous report [24,32], selected for the coexistence of distinct survival strategies that include the

evolution of resistance and sensitivity to the exploiter, which is suggestive of fluctuating selection. If

these trends observed in our experiments are indeed solely a result of the repeated mixing of

communities, it is expected that when the diversified vertically coevolved communities are

propagated under horizontal treatment regimen, homogenization of the survival strategies should be

observed conferring overall increased resistance to predation. When the transfer regimen was flipped

from horizontal to vertical, diverse survival strategies should emerge from a more homogeneously

distributed population. Consistent with our expectation we observe an increased resistance to

predation in our flipped horizontal treatment and diversification in our vertical treatments. Thus,

supporting our previous observation and further establishing that mixing predominantly will result in

an increased resistance to predation in prey irrespective of whether the ancestral population is
isogenic or has a standing diversity.

Though the experimental approaches to studying the effect of population mixing are numerous,
subtle differences between them introduce differential importance to the spatial structure. For
example, population mixing within a single spatially structured arena will allow the mixing of distinct
genotypes but will not affect the overall availability of diffused shared resources such as public-good
[60-62]. Whereas mixing of populations as done in this study will result in the mixing of distinct lineages
repeatedly relative to the vertically propagated communities in which identical genotypes interact
with each other over evolutionary time. Separation of populations during the growth phase and
mixing at each transfer will result in increased mutational supply even after segregation of the
populations. Thus, the design of the evolution experiment allows the horizontal population to behave
as a single large metapopulation. Thus, we used a mathematical modelling approach to demonstrate
that the mixing events, as expected, will effectively allow the horizontal populations (using the growth
phase) to sample a larger number of mutations. The model further shows that this results in changes
in the dynamics of interference competition. These changes in the dynamics of interference
competition resulted in the selection for resistance to predation. This was because of the diminishing
adaptive benefits of resource competition. The model further predicted that the outcomes of the
evolution experiment depend on the continuation of regimens. That is, if the populations evolved in
mixed treatment are propagated as unmixed, then it is expected that the populations will evolve
diverse adaptive strategies like in the vertical treatment. We demonstrate that this outcome is indeed
true by performing additional evolution experiments in which the selection regimens were switched.

The logistics of our evolution experiment involved aspects of evolution in the spatially structured
as well as unstructured environment. For example: limited diffusion of resources is an important
distinguisher between structured and unstructured environments, which has been shown to strongly
affect the evolution of synergistic [25,63] as well as antagonistic interactions [64]. The transfer protocol used
in this study involved population mixing even in the vertically transmitted regimen, thus limiting the
evolution of interaction networks. However, the probability of the evolution of synergistic interactions
was most likely higher in the vertically propagated regimen than in the horizontal regimen. This is
because of the mixing of multiple lineages before each transfer in the latter.

Interestingly, one of the four vertically propagated coevolving prey populations also demonstrated
high levels of resistance to predation, which was comparable to horizontally propagated populations.
Thus, vertically propagated prey populations might explore distinct evolutionary fates that range from
the evolution of higher levels of resistance to *M. xanthus*, to the evolution of within-population
diversity with both resistant and sensitive individuals. This finding is not surprising as similar effects

of differential effects of bacterial predators, protists, and phages on bacteria have been demonstrated

previously [32,65-67]. Nonetheless, unlike prey populations that evolved in horizontal regimen, the vertical

population with high levels of resistance to predation harboured intraspecies diversity (prey with low

and high resistance to predation) which was similar to the other vertically propagated coevolving *E.*

coli populations. Together, it seems that the evolution of the homogenous predation-resistant-prey

population in horizontally propagated lines indicates that the predation-resistant individuals will

outcompete predation-sensitive ones when communities are repeatedly mixed.

In recent years the call for harnessing the potential of microbial predators to control microbial

contaminations and infections has generated tremendous interest. Thus, it is important to understand

the effect of microbial predation on the evolution of prey species. In this context, it is unavoidable

that distinct predator-prey populations might experience mixing events which might change the

effective supply of mutations. Here we demonstrate that because of mixing events microbial predator-

prey interactions might have a strong influence on the evolutionary outcomes for prey populations

and that mixing will most likely have a stronger influence on the evolution of prey bacteria but not the

predator.

Acknowledgements

The authors thank Deepa Agashe and Gregory Velicer for providing *E. coli* (MG1655) and *M. xanthus*

(GJV1) strains, members of the Bacterial Ecology and Evolution lab at the Indian Institute of Science

for helpful discussion. We also thank DST-FIST for support to the Department of Microbiology and Cell

Biology, UGC Centre for advanced study, and the DBT-IISc partnership. AG acknowledges support from

the Ashok and Gita Vaish Junior Researcher Award, as well as the Government of India's DBT

Ramalingaswami Fellowship. This study was mainly supported by the funds from India Alliance

Intermediate fellowship to SP (IA/I/20/1/504921) and HGK-IYBA grant (BT/13/IYBA/2020/15) to SP.

Author contribution

SP conceived the project. SS, BB, and SP designed experiments. SS, BB, and JL performed experiments.

SS, BB and JL analyzed the data. AG performed the mathematical modelling. SS wrote the first draft of

the manuscript. SP edited the manuscript.

446
447

 Figures and figure legends

Figure 1: Effect of selection regimen on the evolution of resistance to predation in prey and predatory performance of *M. xanthus* populations. (A and B) Growth of evolved *E. coli* relative to the growth of ancestral *E. coli* in the presence of ancestral predator is shown. (A) Prey populations coevolved in the vertical transfer regimen were more susceptible to predation than the populations that evolved in the horizontal transfer regimen (paired t-test for differences between vertically evolved *E. coli* relative to horizontally co-evolved populations, p*** < 0.001). (B) Whereas mono-evolved populations showed no significant differences both between treatments as well as relative to the ancestors (Wilcoxon-test for differences between vertically and horizontally mono-evolved populations, *p* > 0.05, n = 4). Color of dots in the vertical regimen represents distinct lineages. (C) Predator populations coevolved in vertical and horizontal treatment exhibited similar predatory performance, which was not significantly different from their respective ancestors. Relative numbers (evolved/ancestor) are log(CFU/ml) of ancestral *E. coli* when cocultured with evolved predator population relative to their respective ancestor (one sample T-test for differences between coevolved predator population and their ancestors, *p* > 0.05, n=4). (D) Mono-evolved predator populations showed no significant differences both between treatment as well as from ancestors *p* > 0.05, n=4). Relative numbers (evolved/ancestor) in log (CFU/ml) of ancestral *E. coli* when cocultured with evolved predator population relative to their respective ancestor. Different color of dots in the vertical regimen represents distinct lineages.

Figure 2: Effect of the selection regimens on the evolution of resistance to predation in prey clones.

Survival of evolved and ancestor *E. coli* clones is shown as log(CFU/ml) of prey numbers when

cocultured with the ancestral *M. xanthus.* Letters indicate significant difference between groups

(Kruskal-Wallis, Dunn's post hoc test, n =3 for each clone studied). Different color of dots in the vertical

and ancestral regimen represents a distinct lineage that started from the ancestor shown in the same

color.

482
483

Figure 3: Effect of the selection regimens on the competitive fitness of prey clones relative to their ancestors. Relative fitness of evolved *E. coli* clones (dots) is shown. All groups are significantly different from 1.0 (one sample Wilcoxon test for differences between average relative-fitness of all clones relative to 1.0, $p^{[FDR\ corrected]} < 0.01$, n = 3), and letters indicate significant differences between groups (Kruskal-Wallis, Dunn's post hoc test for differences among four treatments). Different color of dots in the vertical treatment represents distinct lineages.

Figure 4: Mathematical model of population dynamics suggests that mixing increases clonal interference and preferentially selects for predator avoidance. A) Schematic illustrating the two dynamic phases of the prey and predator in each growth cycle. The first phase, lasting time T_1, exclusively involves exponential prey growth (linear on a log axis), while the second phase, lasting time T_2, involves predator growth and consequent prey death (Methods). B) In each dynamic phase, different aspects of the prey affect its fitness. In the prey growth phase, prey competition dominates, while during prey death, predator avoidance dominates. C) Schematic illustrating how mixing increases clonal interference and slows the evolution of prey competition. Shown are hypothetical growth rate distributions for prey subpopulations. Vertical lines show hypothetical prey variants. Mixing increases the diversity of competing prey variants, lowering the gaps between their growth rates and lowering the relative competitive advantage of prey variants with large growth rates. Thus, prey variants with decreased death rates have a larger selective advantage. D) Results from simulations of our model (for details see Methods) suggest that mixing selects for variants with increased predator avoidance over prey competition, providing an explanation for our observations.

510
511
Figure 5: Prey from the horizontal selection regimen become increasingly more resistant over
evolutionary time whereas the ones from the vertical regimen do not. A) Coevolved *E. coli*
populations after ten (T10), sixteen (T16), and twenty (T20) transfers from vertical (green line) and
horizontal (red line) regimens were analysed for changes in the resistance to predation against
terminally evolved (after twenty cycles) respective *M. xanthus* populations. B) Coevolved *M. xanthus*
populations after ten (T10), sixteen (T16) and twenty (T20) transfers from vertical (green line) and
horizontal (red line) regimens were analysed for the change in the predatory performance against
terminally evolved (after twenty cycles) respective *E. coli* populations from either vertical or horizontal
regimen. Data shown is the average log(CFU/ml) for prey survival against respective *M. xanthus* from
both regimens. Error bars are 95% confidence interval, letters indicate significant differences within
the regimen (ANOVA, Tukey's post hoc test, n = 3).

525
526
Figure 6: Mixing communities always results in evolution of higher resistance to predation while
isolated communities evolve diversified survival strategies in prey population. A) Growth of
ancestral vertically coevolved and horizontally coevolved *E. coli* population (T20 ancestors) along with
the *E. coli* population after sixteen transfers in flipped regimen (Flipped: where vertically coevolved
communities were further propagated under horizontal regimen and horizontally coevolved clones
were further propagated under vertical treatment regimen), in presence of ancestral *M. xanthus*
(ANOVA, Tukey's post hoc test, n = 4). B) Growth of *E. coli* clones isolated from ancestral vertically
coevolved and horizontally coevolved *E. coli* population (T20 ancestors) along with the *E. coli* clones
isolated after sixteen transfers in flipped regimen (Flipped) in presence of ancestral *M. xanthus*
(ANOVA, Tukey's post hoc test, n = 4). Different color of dots in the vertical treatment represents
distinct lineages.

Methods

Semantics

We define predation-prey interaction as an antagonist interaction in which the exploiting partner kills the exploited partner to acquire nutrients. Predator-prey interactions are distinct from host-parasite interactions in many ways [51,54-58] (for example: relationship intimacy, relationship duration, generation time, population size, and trophic level). Importantly, we argue that two factors most likely to differentiate the evolutionary dynamics of host-parasite interactions relative to that of predator-prey interactions are.

1)	The overall population size of the predator is similar or lower relative to the prey. Thus, prey is generally present in higher numbers than predators.

2)	Unlike host-parasite interactions in which parasites have smaller generation time, predator and prey organisms generally have similar generation times.

Thus, unlike host-parasite or host-pathogen interactions, the exploited organisms evolve faster than the exploiters in predator-prey communities.

Strains used

M. xanthus (GJV1) and *E. coli* (MG1655) were used as predator and prey respectively. *M. xanthus* strain used in this study was provided by Dr. Gregory Velicer's laboratory at ETH Zurich, Zurich, Switzerland. Whereas *E. coli* (MG1655) strain was provided by Dr. Deepa Agashe's laboratory at the National Centre for Biological Sciences, Bangalore, India.

Culture condition

All plates were incubated at 32 °C, all liquid cultures were incubated at 32 °C, 200 RPM. *M. xanthus* strains were stored as frozen glycerol (20%) stocks in CTT [68] medium, and *E. coli* strains were stored in frozen glycerol (20%) stocks in LB medium (HIMedia Laboratories Pvt. Ltd, India).

To initiate the evolution experiment *M. xanthus* glycerol stock was inoculated on CTT medium with 1.5 % agar, incubated for 3-5 days, inoculated from colony edge into 8 mL liquid CTT medium, and incubated for 24 h at 200 RPM. Grown liquid culture of *M. xanthus* was dilution plated on CTT 0.5 % agar plates and colonies (subclones) were isolated, stored as freezer stock and distinct subclones were used as biological replicates to initiate the evolution experiment.

Similarly, *E. coli* strains were inoculated on LB medium with 1.5 % agar, incubated for 1-day, individual colonies were inoculated into 8 mL liquid LB medium, and incubated for 24 h at 200 RPM. Each colony (subclone) of *E. coli* served as an ancestor for an independent replicate line.

For experiments performed to analyse evolved population/clones of *M. xanthus*, glycerol stock of

M. xanthus was spotted on CTT medium with 1.5% agar plates and incubated at 32°C for 3-4 days.

Once the swarm appeared, cells from the outer edge (actively growing and swarming) were inoculated

in CTT liquid and incubated under shaking conditions at 32 °C for 20 h. For *E. coli*, cultures were grown

in M9 (MP Biomedicals, United States) + 0.2% glucose (ThermoFisher Scientific, India) liquid, prior to

setting all experiments.

Evolution experiment 1:

Protocol for the evolution experiment is demonstrated in Figure S1. Evolution experiment was

performed in 50 mL flasks containing 10 mL M9 medium + 1.5 % agar medium (M9 salts (MP

Biomedicals) + 0.2 % glucose + 0.01mM $CaCl_2$ (ThermoFisher Scientific, India) + 0.2mM $MgSO_4$ + 1.5%

bacto agar (DIFCO, BD Biosciences, India)). Glucose in the M9 medium supports the growth of *E. coli*

whereas *M. xanthus* cannot utilise glucose as a carbon source, and hence depend on their predatory

ability to feed on the growing population of *E. coli.*

To initiate the evolution experiment, the densities of *E. coli* and *M. xanthus* were adjusted to ~10^6

cells/mL and ~$2*10^{10}$ cells/mL respectively by resuspending in M9 buffer (M9 salts (MP Biomedicals)

+ 0.01mM $CaCl_2$ + 0.2mM $MgSO_4$). 50 µL culture of prey and predator was spread across the agar bed

with 9-12 sterile glass beads. For the *E. coli* only control 50 µL of *E. coli* culture was spread on the agar

beds along with 50 µL of M9 buffer (M9 salts (MP Biomedicals, United States) + 0.01mM $CaCl_2$ + 0.2mM

$MgSO_4$). For predator-only control, agar beds were supplemented with 5% dead *E. coli.* Dead *E. coli*

were obtained by autoclaving 10 mL of 1.0 OD culture of *E. coli* for 20 minutes. All agar beds were

prepared 1 day prior to the transfers. After 96 h of incubation at 32°C the predator-prey communities

were harvested by adding 4 mL of M9 buffer followed by shaking for ~30 mins at 200 RPM to

resuspend the growing culture. For the vertical transfer regimen, 200 µL (5 %) of the resuspended

culture from each flask was transferred to a fresh flask and spread using glass beads. In horizontal

transfer regimen, all four replicate cultures of the respective treatment were pooled in a single 15 mL

conical centrifuge tube, mixed, 200 µL of these pooled cultures were redistributed in each of the four

replicate flasks with fresh medium and spread with the help of 9-12 glass-beads. The evolution

experiment was carried out for 20 such transfer cycles with populations being stored at regular

intervals in 20 % glycerol stocks at -80 °C.

Isolation of clones from evolved populations of *M. xanthus* and *E. coli*

To isolate the evolved population of *E. coli*, a fraction of frozen glycerol stock was inoculated in 5 mL LB liquid medium (with 1 % NaCl instead of 0.5 %), incubated at 32 °C at 200 RPM for 7-8 h, and frozen as 20 % glycerol stock. *M. xanthus* cells are sensitive to the presence of salt and hence do not grow in these conditions, whereas cells of *E. coli* do.

To isolate *M. xanthus* population, frozen glycerol stock was inoculated in 8 mL CTT liquid with Gentamicin (Sisco Research Laboratories Pvt. Ltd, India) (10 µg/mL), incubated at 32 °C at 200 RPM for 18-20 h, and frozen as 20 % glycerol stock. Being naturally resistant to Gentamicin, *M. xanthus* can grow in the presence of Gentamicin.

Next, to isolate *E. coli* clones glycerol stocks of frozen *E. coli* populations were dilution plated on LB + 0.5 % agar (Agar agar, ThermoFisher Scientific, India), incubated at 32 °C for 18 h. Distinct colonies were isolated as clones, inoculated in liquid LB medium, incubated at 32 °C at 200 RPM for 7-8 h, and stored as glycerol (20 %) stocks at -80 °C.

To isolate *M. xanthus* clones glycerol stocks of frozen *M. xanthus* populations were pour-platted in CTT + 0.5 % agar, and incubated at 32 °C for 96 h. Distinct colonies were isolated as subclones, inoculated in liquid CTT medium, incubated at 32 °C at 200 RPM for 20 h, and stored at glycerol (20 %) stocks at -80 °C.

Predation assay

Predatory performance can be measured as the growth of M. *xanthus* when prey is the only source of nutrient, or growth of *E. coli* in the presence of *M. xanthus* relative to its absence. In our study, we used growth of *E. coli* as a measure of predation by *M. xanthus*. This is because *M. xanthus* is a sticky biofilm forming bacterium. Therefore, estimates of Colony Forming Units (CFU) are generally unreliable, and a single colony might emerge from a group of cells resulting in significant underestimation of *M. xanthus* cell numbers.

All predation assays were performed on 10 mL M9 + 1.5 % agar beds (M9 salts (MP Biomedicals) + 0.2 % glucose + 0.01 mM $CaCl_2$ + 0.2 mM $MgSO_4$ +1.5 % Bacto agar) in 50 mL conical flasks with 9-12 sterile glass beads. Media was poured into the flask one day prior to the experiment. All *E. coli* cultures were grown to log phase in liquid M9 media (M9 salts (MP Biomedicals) + 0.2 % glucose + 0.01 mM $CaCl_2$ + 0.2 mM $MgSO_4$) prior to setting up the experiments. The *M. xanthus* cultures were grown to log phase in liquid CTT media. The prey cell density was adjusted to 0.1 OD (~10^6 cells/mL) and predator cell density was adjusted to $5*10^9$ cells/mL in TPM buffer (10 mM Tris pH 8.0, 8 mM $MgSO_4$, 1 mM K_2HPO_4 – KH_2PO_4), 50µL of these cultures were then inoculated on the M9+1.5 % agar beds and spread with sterile glass beads. For monoculture of *E. coli,* 50 µL of cell density adjusted culture was

inoculated along with 50 µL TPM buffer and spread on the M9+ 1.5 % agar medium. The cultures were

incubated for 96 h at 32 °C. Following the incubation period, cells were harvested by adding 4 mL of

TPM buffer with 9-12 sterile glass beads followed by ~30 min shaking at 200 RPM. Viable number of

E. coli was determined by dilution plating the resuspended cell suspension in LB + 0.5 % agar.

Competition assay

Competition between ancestors with and without Kanamycin marker

E. coli ancestral strains with and without Kanamycin marker were grown to log phase in LB liquid

medium and 1:1 ratio of 0.1 OD adjusted culture of each strain was co-inoculated to initiate the

experiment. The competition experiment was performed on 10 mL M9 + 1.5 % agar beds (M9 salts

(MP Biomedicals) + 0.2 % glucose + 0.01 mM $CaCl_2$ + 0.2 mM $MgSO_4$ + 1.5 % Bacto agar) in 50 mL

conical flasks with 9-12 sterile glass beads. The culture was spread evenly on the agar-beds using glass

beads and incubated for 24 h at 32 °C. Post incubation, the cells were harvested by adding 4 mL TPM

buffer with ~30 min shaking at 200 RPM. Viable counts of Kanamycin sensitive and resistant *E. coli*

were determined by dilution plating on LB + 0.5 % agar and LB + 0.5 % agar with Kanamycin (40 µg /

mL) (Merck Germany).

Competition between evolved clones and Kanamycin marked ancestors

Four *E. coli* clones were isolated from each of the vertically evolved population, sixteen from each

from horizontally co-evolved and mono-evolved populations. Each of these clones were competed

against their respective ancestors, in the presence or absence of ancestral *M. xanthus.* For these

experiments we used Kanamycin-resistant ancestor *E. coli*, which allowed us to distinguish them from

the evolved clones. Competition experiments were performed on 10 mL M9 + 1.5 % agar beds (M9

salts (MP Biomedicals) + 0.2 % glucose + 0.01 mM $CaCl_2$ + 0.2 mM $MgSO_4$ + 1.5 % Bacto agar) in 50 mL

conical flasks with 9-12 sterile glass beads. All *E. coli* cultures were grown to log phase in M9 liquid

media, 25 µL of 0.1 OD (~10^6 cells/mL) adjusted culture of each evolved and ancestor *E. coli* were co-

inoculated, either with (50 µL of $5*10^9$ cells/mL) or without (50 µL TPM buffer) ancestral *M. xanthus*,

spread on the agar medium with sterile glass beads. These were incubated for 96 h at 32 °C and

harvested by adding in 4 mL TPM buffer with ~30 min shaking at 200 RPM. Viable counts of ancestor

and evolved *E. coli* were determined by dilution plating on LB + 0.5 % agar and LB+0.5% agar with

Kanamycin (40 µg/mL) (Merck Germany).

Determination of prey-predator growth dynamics in co-cultures experiments

To demonstrate the relative growth dynamics of the prey and predator when co-cultured together in

media containing only glucose as a carbon source, the prey numbers and predator densities were

tracked for 4 days. Since only *E. coli* can utilise glucose and *M. xanthus* cannot. The only available

source of nutrition for *M. xanthus* was the prey population. Additionally, given the differences in

generation times of *E. coli* and *M. xanthus*, their growth in the coculture is expected to be temporally

segregated. Hence, it is expected that *E. coli* will have distinct growth and death phases. Thus, to

determine the changes in the population size of prey bacteria during the four-day transfer cycle, prey

counts were obtained every 24 h from 0 to 96 h. To do so, 50 µL of 0.1 OD (~10^6 cells/mL) adjusted

culture of ancestor *E. coli* was co-inoculated, either with (50 µL of $5*10^9$ cells/mL) or without (50 µL

TPM buffer) ancestral *M. xanthus*, and spread on 10 mL of M9 + 1.5 % agar beds (M9 salts (MP

Biomedicals) + 0.2 % glucose + 0.01 mM $CaCl_2$ + 0.2 mM $MgSO_4$ + 1.5 % Bacto agar). The cultures were

spread on the agar medium with the help of sterile glass beads, incubated at 32 °C, and harvested by

adding in 4 mL TPM buffer with ~30 min shaking at 200 RPM. Viable counts of *E. coli* were determined

by dilution plating on LB + 0.5 % agar. Because of the destructive sampling approach, distinct prey-

predator microcosms were set up for each time point.

Since exopolysaccharides secreted by *M. xanthus* result in the formation of biofilms that are difficult

to disrupt, it reduces the reliability of the obtained *M. xanthus* viable counts, using similar methods

as for *E. coli*. Therefore, we used the swarm sizes as a proxy to determine the relative densities of *M.*

xanthus in the cocultures. Swarming of *M. xanthus* on hard agar surfaces is known to be a positive

density-dependent trait. Hence, it is expected that as the relative numbers increase, the swarm sizes

will also increase. To estimate the swarm sizes, post harvesting of the cocultures in TPM buffer (as

described above), 10 µL of the culture was spotted on CTT hard agar (1.5 Bacto agar) surface in 90 mm

plates. The initial swarm was marked after 24 h and the final swarms were marked after 120 h. The

average distance between the initial and final swarms was recorded and presented as the swarm sizes.

Mathematical model of evolution in predator-prey communities

To explain our surprising experimental observation that variants with superior competitive fitness that

did not show any increase in their ability to resist predation evolved only in the vertical treatment and

not in the horizontal treatment, we developed a mathematical model of population dynamics. Our

model followed the dynamics of prey subpopulations explicitly and the predator populations

implicitly. Specifically, we assumed each growth cycle (which lasted time T) had two distinct periods:

the first where prey subpopulations grew (lasting time T_1) and the second where they died due to

predation (lasting time T_2). In each period, each subpopulation (labeled α) had an exponential growth

rate g_α and death rate d_α respectively, both drawn randomly and independently from a normal

distribution with means 0.7 hr^{-1} and 0.2 hr^{-1} respectively, and standard deviations 0.3 hr^{-1} and 0.2 hr^{-1}

respectively (disallowing negative rates). With these the growth of each subpopulation was described

by the following equation:

$$N\alpha(T) = N_\alpha(0) \exp\left(g_\alpha T_1 - d_\alpha T_2\right)$$

Inspired by evolutionary ecology, our model implemented evolution by introducing distinct

"mutants" as prey subpopulations at the beginning of population dynamics and following their fate

over time. We modeled 4 distinct communities, each seeded with 5 subpopulations of prey (chosen

arbitrarily). At the beginning of each simulation, each subpopulation started with equal abundance

0.1 (arbitrary units) and grew according to the dynamics described above. After each growth cycle

(time T), all subpopulations in each population were diluted by a factor $D = 20$ (5% bottleneck) and

grown again in a new cycle. In the vertical regimen, different communities were not mixed, while in

the horizontal one, they were mixed and redistributed equally, mimicking our experiment. We

continued these dynamics across multiple growth-dilution cycles till communities reached a steady

state, after which we compared the mean ratio of the relative growth rates (quantifying prey

competition) and death rates (quantifying predator avoidance) of each surviving subpopulation. We

repeated and averaged our results over such 100 simulations with randomly chosen parameters and

plotted the results (Fig. 4D).

Time-course assay

Predation assays (were performed with populations of coevolved prey and predator from both the

transfer regimes (vertical and horizontal), respectively isolated from different evolutionary time. The

populations of both prey and predator were isolated from the glycerol stocks stored at different

transfers, namely, after transfer 10 (T10), transfer 16 (T16) and transfer 20 (T20). The prey, and the

predator populations were isolated using a similar protocol as described earlier (See isolation of

population and clones). To determine the change in prey survival over evolutionary time against the

terminally coevolved predator, *E. coli* populations from different evolutionary times (T10, T16 and

T20) were inoculated with the terminally evolved *M. xanthus* (after 20 transfers). The predation assay

for each combination of prey and predator population was set up as discussed earlier (See predation

assay). The change in prey survival over time was recorded for prey population by dilution plating on

LB + 0.5 % agar. Similarly, the *M. xanthus* populations from all time points (T10, T16 and T20) were

pitched against the terminally co-evolved *E. coli* population, to determine the changes in the

predatory ability over evolutionary time. The survival of the terminally coevolved prey against *M.*

xanthus from different evolutionary time-point was used as a measure of change in the resistance to

predation over evolutionary time. Here, a lower prey survival implies better predatory ability, and a
higher prey survival indicates worse predatory ability.

Flipped-regime evolution experiment (Evolution experiment 2)

Experimental protocol for the second evolution experiment is demonstrated in Supplementary

figure S7. Evolution experiment was performed in 50 mL flasks containing 10 mL M9 + 1.5 % agar

medium (M9 salts (MP Biomedicals) + 0.2 % glucose+ 0.01mM CaCl2 + 0.2mM MgSO4 + 1.5 % bacto

agar). To initiate the second evolution experiment, glycerol stocks of the T20 population from

evolution experiment 1 were thawed on ice and 5% of the culture was washed with M9 buffer two

times and then resuspended in 200 μL M9 buffer. These cultures were then spread on fresh M9 + 1.5

% agar beds with the help of 9-12 sterile glass beads. Similar treatment was done for all populations

from both mixed communities of prey and predator and monoculture prey (control) treatments. The

mixed communities of prey and predator (coevolved) /prey only (mono-evolved) population were

incubated in fresh media at 32 ^{O}C for 96 h. This initial 96 h incubation was done to allow the

stabilization of the communities after revival from freezer stocks. Following this both the mixed

communities of prey-predator and prey only treatments were harvested in 4 mL of M9 buffer and

shaking at ~200 RPM for 30 mins to resuspend the growing cultures. 5 % of the resuspended culture

was transferred to fresh M9 medium + 1.5 % agar medium under two different transfer regimens:

Vertical transfer regimen and Horizontal transfer regimen (similar to the first experiment). However,

in this experiment the transfer regimens of the evolved population from evolution experiment-1 were

interchanged. The vertically coevolved/mono-evolved population were propagated under the

horizontal transfer regimen, i.e., the population from replicate lines were pooled in a 15 mL centrifuge

tube and then 5% of the resuspended culture was transferred to each of the 4 flasks with fresh media,

at every transfer (ensuring a periodic mixing of population). Whereas the horizontally

coevolved/mono-evolved population were propagated under the vertical transfer regimen, where

suspended cultures from the 4 replicate lines were serially transferred to respective fresh M9 + 1.5 %

agar medium. All the cultures were spread on the agar beds using the glass beads. The experiment

was carried out for 20 such cycles with population being stored at regular intervals in the form of 20

% glycerol stocks in -80 °C. Evolved population of prey and predator was isolated after 16 transfer

cycles using similar protocols as used for the isolation of population in the previous experiments.

Genomic analysis

For the isolation of genomic DNA, *E. coli* and *M. xanthus* clones were grown in M9 + 0.2 % glucose

and CTT liquid medium respectively at 32 ^{o}C, 200 RPM till they achieved optical density between 0.2

to 0.8 at 600 nm. These cultures were used for the extraction of genomic DNA using the QIAGEN

genomic DNA isolation kit using 20/G column. Manufacturer's instructions were followed for the

isolation of genomic DNA. Next steps for sequencing were performed at Macrogen Inc (Korea). Before

TrueSeq library preparation the quality control of the samples was done using Agilent TapeStation[R]

DNA Screen Tape (Agilent). The DNA samples were sequenced on HiSeq platform, for 150 bp paired-

end sequencing. Ambiguous low- quality sequences were trimmed using Trimmomatic v0.39. The

Genomic libraries were analysed using breseq[69] against reference genome of *E. coli* MG1655 (GenBank

Accession No. GCA_000005845.2) or *M. xanthus* DK1622 (GenBank Accession No.GCA_000012685.1)

from NCBI genomic data bank. The breseq pipeline was used with all default parameters.

Statistical analysis

All statistical analysis were performed in R (version 4.0.4). Levene's test was performed with each

set of data to check for the homogeneity of variance. For a $p > 0.05$ for Levene's test, either an

independent sample t-test or one sample t-test was performed wherever applicable, and the

Benjamini-Hochberg method was used for correcting for multiple testing. Significant ANOVA in all

analysis was followed by an SNK *post hoc* test. However, for a $p < 0.05$ for Levene's test nonparametric

tests were used for data analysis. The significant Kruskal-Wallis test was followed by Dunn's *post hoc*

test. The statistical test for the respective data set is mentioned in the figure legends.

References

1.   Brockhurst, M. A., Morgan, A. D., Rainey, P. B., & Buckling, A. (2003). Population
mixing accelerates coevolution. Ecol. Lett., *6*(11), 975-979.

2.   Benmayor, R., Hodgson, D. J., Perron, G. G., & Buckling, A. (2009). Host mixing and
disease emergence. *Curr. Biol.*, *19*(9), 764-767.

3.   Xiao, Y., Wei, X., Ebright, R., and Wall, D. (2011). Antibiotic production by
myxobacteria plays a role in predation. J. Bacteriol. *193*, 4626-4633.

4.   Berleman, J.E., Allen, S., Danielewicz, M.A., Remis, J.P., Gorur, A., Cunha, J., Hadi, M.Z.,
Zusman, D.R., Northen, T.R., Witkowska, H.E., and Auer, M. (2014). The lethal cargo of
Myxococcus xanthus outer membrane vesicles. Front. Microbiol. *5*. ARTN 474.

5.   Song, H., Payne, S., Gray, M., and You, L.C. (2009). Spatiotemporal modulation of
biodiversity in a synthetic chemical-mediated ecosystem. Nat. Chem. Biol. *5*, 929-935.

6.   Sicardi, E.A., Fort, H., Vainstein, M.H., and Arenzon, J.J. (2009). Random mobility and
spatial structure often enhance cooperation. J. Theor. Biol. *256*, 240-246.

7.   Vigneux, F., Bashey, F., Sicard, M., and Lively, C.M. (2008). Low migration decreases
interference competition among parasites and increases virulence. J. Evol. Biol. *21*, 1245-
1251.

8.   Jokela, J., Dybdahl, M., and Lively, C. (2009). The maintenance of sex, clonal dynamics,
and host-parasite coevolution in a mixed population of sexual and asexual snails. Am. Nat.
174, S43-S53.

9.   Furey, N.B., Armstrong, J.B., Beauchamp, D.A., and Hinch, S.G. (2018). Migratory
coupling between predators and prey. Nat. Ecol. Evol. *2*, 1846-1853.

10.   Hughes, K.A., Convey, P., Maslen, N.R., and Smith, R.I.L. (2010). Accidental transfer of
non-native soil organisms into Antarctica on construction vehicles. Biol. Invasions *12*, 875-
891.

11.   Gilbert, J.A., Field, D., Swift, P., Newbold, L., Oliver, A., Smyth, T., Somerfield, P.J.,
Huse, S., and Joint, I. (2009). The seasonal structure of microbial communities in the Western
English Channel. Environ. Microbiol. *11*, 3132-3139.

12.   Bottos, E.M., Laughlin, D.C., Herbold, C.W., Lee, C.K., McDonald, I.R., and Cary, S.C.
(2020). Abiotic factors influence patterns of bacterial diversity and community composition
in the dry valleys of Antarctica. FEMS Microbiol. Ecol. *96*.

13.   Rillig, M.C., Antonovics, J., Caruso, T., Lehmann, A., Powell, J.R., Veresoglou, S.D., and
Verbruggen, E. (2015). Interchange of entire communities: microbial community coalescence.
Trends Ecol. Evol. *30*, 470-476.

14.   Lipsitch, M., Nowak, M.A., Ebert, D., and May, R.M. (1995). The population dynamics
of vertically and horizontally transmitted parasites. Proc. Biol. Sci. *260*, 321-327.

15.   Reger, J., Lind, M.I., Robinson, M.R., and Beckerman, A.P. (2018). Predation drives
local adaptation of phenotypic plasticity. Nat. Ecol. Evol. *2*, 100-107.

16.   Decaestecker, E., Gaba, S., Raeymaekers, J.A.M., Stoks, R., Van Kerckhoven, L., Ebert,
D., and De Meester, L. (2007). Host-parasite 'Red Queen' dynamics archived in pond
sediment. Nature *450*, 870-873.

17.   Larsen, J., Raisen, C.L., Ba, X.L., Sadgrove, N.J., Padilla-Gonzalez, G.F., Simmonds,
M.S.J., Loncaric, I., Kerschner, H., Apfalter, P., Hartl, R., et al. (2022). Emergence of methicillin
resistance predates the clinical use of antibiotics. Nature *602*, 135-141.

18.   Lively, C.M., Clay, K., Wade, M.J., and Fuqua, C. (2005). Competitive co-existence of
vertically and horizontally transmitted parasites. Evol. Ecol. Res. *7*, 1183-1190.

19.   Leigh, V.V. (1973). A new evolutionary law. Evol Theory *1*, 30.

20.   Buckling, A., and Rainey, P.B. (2002). Antagonistic coevolution between a bacterium
and a bacteriophage. P. Roy. Soc. B-Biol. Sci. *269*, 931-936.

21.   Lively, C.M. (1986). Predator-induceds shell dimorphism in the acorn barnacle
Chthamalus-anisopoma. Evolution *40*, 232-242.

22.   Baum, J.K., and Worm, B. (2009). Cascading top-down effects of changing oceanic
predator abundances. J. Anim. Ecol. *78*, 699-714.

23.   Gurevitch, J., Morrison, J.A., and Hedges, L.V. (2000). The interaction between
competition and predation: A meta-analysis of field experiments. Am. Nat. *155*, 435-453.

24.   Nair, R.R., Vasse, M., Wielgoss, S., Sun, L., Yu, Y.N., and Velicer, G.J. (2019). Bacterial
predator-prey coevolution accelerates genome evolution and selects on virulence-associated
prey defences. Nat. Commun. *10*, 4301.

25.   Pande, S., Kaftan, F., Lang, S., Svatos, A., Germerodt, S., and Kost, C. (2016).
Privatization of cooperative benefits stabilizes mutualistic cross-feeding interactions in
spatially structured environments. ISME J *10*, 1413-1423.

26.   Rosenberg, E., Vaks, B., and Zuckerberg, A. (1973). Bactericidal action of an antibiotic
produced by *Myxococcus xanthus*. Antimicrob. Agents Ch *4*, 507-513.

27.   Sudo, S., and Dworkin, M. (1972). Bacteriolytic enzymes produced by *Myxococcus*
xanthus. J. Bacteriol. *110*, 236-245.

28.   Zhang, W., Wang, Y., Lu, H., Liu, Q., Wang, C., Hu, W., and Zhao, K. (2020). Dynamics
of solitary predation by *Myxococcus xanthus* on *Escherichia coli* observed at the single-cell
level. Appl. Environ. Microbiol. *86*. ARTN e02286-19.

29.   Pan, H.W., He, X.S., Lux, R., Luan, J., and Shi, W.Y. (2013). Killing of *Escherichia coli by*
Myxococcus xanthus in aqueous environments requires exopolysaccharide-dependent
physical contact. Microb. Ecol. *66*, 630-638.

30.   Seef, S., Herrou, J., de Boissier, P., My, L., Brasseur, G., Robert, D., Jain, R., Mercier, R.,
Cascales, E., Habermann, B.H., and Mignot, T. (2021). A Tad-like apparatus is required for
contact-dependent prey killing in predatory social bacteria. Elife *10*. ARTN e72409

31.   Evans, A.G.L., Davey, H.M., Cookson, A., Currinn, H., Cooke-Fox, G., Stanczyk, P.J., and
Whitworth, D.E. (2012). Predatory activity of *Myxococcus xanthus* outer-membrane vesicles
and properties of their hydrolase cargo. Microbiol-SGM *158*, 2742-2752.

32.   Nair, R.R., and Velicer, G.J. (2021). Predatory bacteria select for sustained prey
diversity. Microorganisms *9*. ARTN 2079
878

33.   Cortina, N.S., Krug, D., Plaza, A., Revermann, O., and Muller, R. (2012).
Myxoprincomide: a natural product from *Myxococcus xanthus* discovered by comprehensive
analysis of the secondary metabolome. Angew. Chem. Int. Ed. Engl. *51*, 811-816.

34.   Simunovic, V., Zapp, J., Rachid, S., Krug, D., Meiser, P., and Muller, R. (2006).
Myxovirescin A biosynthesis is directed by hybrid polyketide synthases/nonribosomal peptide
synthetase, 3-hydroxy-3-methylglutaryl-CoA synthases, and trans-acting acyltransferases.
ChembioChem. *7*, 1206-1220.

35.   Wehland, M., and Bernhard, F. (2000). The RcsAB box - Characterization of a new
operator essential for the regulation of exopolysaccharide biosynthesis in enteric bacteria. J.
Biol. Chem. *275*, 7013-7020.

36.   Howery, K.E., Clemmer, K.M., and Rather, P.N. (2016). The Rcs regulon in Proteus
mirabilis: implications for motility, biofilm formation, and virulence. Curr. Genet. *62*, 775-789.

37.   Wang, Q.F., Zhao, Y.F., McClelland, M., and Harshey, R.A. (2007). The RcsCDB signaling
system and swarming motility in *Salmonella enterica* serovar *Typhimurium*: Dual regulation
of flagellar and SPI-2 virulence genes. J. Bacteriol. *189*, 8447-8457.

38.   Morgan, A.D., Gandon, S., and Buckling, A. (2005). The effect of migration on local
adaptation in a coevolving host-parasite system. Nature *437*, 253-256

39.   Mansour, I., Heppell, C.M., Ryo, M., and Rillig, M.C. (2018). Application of the
microbial community coalescence concept to riverine networks. Biol. Rev. Camb. Philos. Soc.
93, 1832-1845.

40.   Durham, W.M., Climent, E., Barry, M., De Lillo, F., Boffetta, G., Cencini, M., and
Stocker, R. (2013). Turbulence drives microscale patches of motile phytoplankton. Nat.
Commun. *4*, 2148.

41.   Ignacio-Espinoza, J.C., Ahlgren, N.A., and Fuhrman, J.A. (2020). Long-term stability and
Red Queen-like strain dynamics in marine viruses. Nat. Microbiol. *5*, 265-271.

42.   Berleman, J.E., and Kirby, J.R. (2009). Deciphering the hunting strategy of a bacterial
wolfpack. FEMS Microbiol. Rev. *33*, 942-957.

43.   Hart, B.A., and Zahler, S.A. (1966). Lytic Enzyme Produced by *Myxococcus xanthus*. J.
Bacteriol. *92*, 1632-1637.

44.   Foster, K.R., and Bell, T. (2012). Competition, not cooperation, dominates interactions
among culturable microbial species. Curr. Biol. *22*, 1845-1850.

45.   Xavier, J.B., and Foster, K.R. (2007). Cooperation and conflict in microbial biofilms. P.
Nat. Acad. Sci. USA *104*, 876-881.

46.   Sockett, R.E., and Lambert, C. (2004). Bdellovibrio as therapeutic agents: a predatory
renaissance? Nat. Rev. Microbiol. *2*, 669-675.

47.   Hobley, L., Summers, J.K., Till, R., Milner, D.S., Atterbury, R.J., Stroud, A., Capeness,
M.J., Gray, S., Leidenroth, A., Lambert, C., et al. (2020). Dual predation by bacteriophage and
Bdellovibrio bacteriovorus can eradicate *Escherichia coli* prey in situations where single
predation cannot. J. Bacteriol. *202*. ARTN e00629-19

48.   Kadouri, D.E., To, K., Shanks, R.M.Q., and Doi, Y. (2013). Predatory bacteria: A potential
ally against multidrug-resistant Gram-negative pathogens. PLoS One *8*. ARTN e63397
920
49.   Morgan, A.D., MacLean, R.C., Hillesland, K.L., and Velicer, G.J. (2010). Comparative
analysis of *Myxococcus* predation on soil bacteria. Appl. Environ. Microb. *76*, 6920-6927.

50.   Matz, C., Webb, J.S., Schupp, P.J., Phang, S.Y., Penesyan, A., Egan, S., Steinberg, P., and
Kjelleberg, S. (2008). Marine biofilm bacteria evade eukaryotic predation by targeted
chemical defense. PLoS One *3*. ARTN e2744.

51.   Raffel, T.R., Martin, L.B., and Rohr, J.R. (2008). Parasites as predators: unifying natural
enemy ecology. Trends Ecol. Evol. *23*, 610-618.

52.   Cowley, L.A., Dallman, T.J., Jenkins, C., and Sheppard, S.K. (2019). Phage predation
shapes the population structure of shiga-toxigenic *Escherichia coli* O157:H7 in the UK: An
evolutionary perspective. Front. Genet. *10*. ARTN 763.

53.   Morgan, A.D., Brockhurst, M.A., Lopez-Pascua, L.D.C., Pal, C., and Buckling, A. (2007).
Differential impact of simultaneous migration on coevolving hosts and parasites. BMC Evol.
Biol. 7, ARTN 1.

54.   Combes, C. (2001). Parasitism: the ecology and evolution of intimate interactions. U
of Chicago P.

55.   Antia, R., Levin, B. R., & May, R. M. (1994). Within-host population dynamics and the
evolution and maintenance of microparasite virulence. Am. Nat., 144(3), 457-472.

56.   Martin, L. B., Weil, Z. M., & Nelson, R. J. (2008). Seasonal changes in vertebrate
immune activity: mediation by physiological trade-offs. Philo. Trans. R. Soc. B: Biol.
Sci., 363(1490), 321-339.

57. Lafferty, K. D., Allesina, S., Arim, M., Briggs, C. J., De Leo, G., Dobson, A. P., … &
Thieltges, D. W. (2008). Parasites in food webs: the ultimate missing links. Ecol. Lett., 11(6),
533-546.

58. Lafferty, K. D., & Kuris, A. M. (2002). Trophic strategies, animal diversity and body
size. Trends Ecol. Evol., 17(11), 507-513.

59. Livingston, G., Jiang, Y.X., Fox, J.W., and Leibold, M.A. (2013). The dynamics of
community assembly under sudden mixing in experimental microcosms. Ecology *94*, 2898-
2906.

60. Popat, R., Crusz, S.A., Messina, M., Williams, P., West, S.A., and Diggle, S.P. (2012).
Quorum-sensing and cheating in bacterial biofilms. P. Roy. Soc. B-Biol. Sci. *279*, 4765-4771.

61. Butaite, E., Baumgartner, M., Wyder, S., and Kummerli, R. (2017). Siderophore
cheating and cheating resistance shape competition for iron in soil and freshwater
Pseudomonas communities. Nat. Comm. *8*. ARTN 414.

62. Drescher, K., Nadell, C.D., Stone, H.A., Wingreen, N.S., and Bassler, B.L. (2014).
Solutions to the public goods dilemma in bacterial biofilms. Curr. Biol. *24*, 50-55.

63. Estrela, S., Morris, J.J., and Kerr, B. (2015). Private benefits and metabolic conflicts
shape the emergence of microbial interdependencies. Environ. Microbiol. 10.1111/1462-
2920.13028.

64. Kerr, B., Riley, M.A., Feldman, M.W., and Bohannan, B.J.M. (2002). Local dispersal
promotes biodiversity in a real-life game of rock-paper-scissors. Nature *418*, 171-174

65. Johnke, J., Baron, M., de Leeuw, M., Kushmaro, A., Jurkevitch, E., Harms, H., and
Chatzinotas, A. (2017). A generalist protist predator enables coexistence in multitrophic
predator-prey systems containing a phage and the bacterial predator *Bdellovibrio.* Front. Ecol.
Evol. *5*. ARTN 124.

66. Bell, T., Bonsall, M.B., Buckling, A., Whiteley, A.S., Goodall, T., and Griffiths, R.I. (2010).
Protists have divergent effects on bacterial diversity along a productivity gradient. Biol. Lett.
6, 639-642.

67. Pyenson, N.C., and Marraffini, L.A. (2020). Co-evolution within structured bacterial
communities results in multiple expansion of CRISPR loci and enhanced immunity. Elife *9*.
ARTN e53078.

68. Hodgkin, J., and Kaiser, D. (1977). Cell-to-Cell Stimulation of movement in nonmotile
mutants of *Myxococcus.* P. Natl. Acad. Sci. USA *74*, 2938-2942.

69. Deatherage D, Barrick J (2014). Identification of mutations in laboratory-evolved
microbes from next-generation sequencing data using breseq. Engineering and Analyzing
Multicellular Systems (*pp. 165-188*). Humana Press, New York, NY.