Microsaccades track location-based object rehearsal in visual working memory

Eelke de Vries (e3.de.vries@vu.nl) & Freek van Ede (freek.van.ede@vu.nl)

Institute for Brain and Behavior Amsterdam, Department of Experimental and Applied Psychology, Vrije Universiteit Amsterdam, The Netherlands

ABSTRACT

Besides controlling eye movements, the brain's oculomotor system has been implicated in the control of covert spatial attention and the rehearsal of spatial information in working memory. We investigated whether the oculomotor system also contributes to rehearsing visual objects in working memory when object location is never asked about. To address this, we tracked the incidental use of locations for mnemonic rehearsal via directional biases in microsaccades while participants maintained two visual objects (coloured oriented gratings) in working memory. By varying the stimulus configuration (horizontal, diagonal, and vertical) at encoding, we could quantify whether microsaccades were more aligned with the configurational axis of the memory contents, as opposed to the orthogonal axis. Experiment 1 revealed that microsaccades continued to be biased along the axis of the memory content several seconds into the working-memory delay. In Experiment 2, we confirmed that this directional microsaccade bias was specific to memory demands, ruling out lingering effects from passive and attentive encoding of the same visual objects in the same configurations. Thus, by studying microsaccade directions, we uncover oculomotor-driven rehearsal of visual objects in working memory through their associated locations.

INTRODUCTION

The brain's oculomotor system, which controls our eye movements, has also been linked to the regulation of covert cognitive processes. To date, the cognitive involvement of the oculomotor system has been established most clearly in the studies of covert visual-spatial attention and the rehearsal of spatial information in working memory (e.g., Corbetta et al., 1998; Engbert & Kliegl, 2003; Hafed et al., 2015; Jonikaitis & Moore, 2019; Krauzlis et al., 2013; Moore et al., 2003; Nobre et al., 2000; Rizzolatti et al., 1987; Theeuwes et al., 2009). In the current study, we tested whether the oculomotor system also contributes to the location-based rehearsal of visual objects in working memory when object locations are never explicitly asked about. To address this, we here studied microsaccades as a direct time-resolved output from the oculomotor system.

Studies in which participants are explicitly tasked with memorising the locations of visual stimuli have revealed that the oculomotor system plays a role in the mnemonic rehearsal of these locations. Most directly, spatial rehearsal of visual locations has been shown to trigger eye movements to these locations (Olsen et al., 2014; Wynn et al., 2018). Complementary evidence comes from studies showing better visual-spatial memory when eye movements are allowed throughout the delay interval (McAteer et al., 2023; Pearson et al., 2014; Souza et al., 2020; Williams et al., 2013; Postle et al., 2006; but see also Godijn & Theeuwes, 2012; Loaiza & Souza, 2022), and worse memory when eye movements are prompted to other locations (Lawrence et al., 2004; Ohl & Rolfs, 2020; Postle et al., 2006; Tremblay et al., 2006). Neuroscience studies in humans (Courtney et al., 1998; Curtis et al., 2004; Jerde et al., 2012; Postle et al., 2000) and non-human primates (Armstrong et al., 2009; Merrikhi et al., 2017) have further revealed how spatial locations in working memory are represented in the frontal eye fields – a key cortical node in the brain's oculomotor system (e.g., Marshall et al., 2015; Moore & Armstrong, 2003). These studies collectively point to an important role for the oculomotor system in the rehearsal of spatial-location information in working memory.

Besides the retention of locations, working memory enables us to retain object-specific visual features, such as object colour and shape. The aforementioned oculomotor involvement in visual working memory may be specific to the rehearsal of locations (as suggested, for example, in McAteer et al., 2023; Postle et al., 2006). At the same time, we know from ample prior studies that space remains a profound organizing principle for object retention in working memory, even when the location of memoranda is never asked about. For example, spatial locations may help to keep memory objects separate, and facilitate the binding of features belonging to the same memory object (Abrahamse et al., 2014; Hollingworth, 2007; Manohar et al., 2017; Pertzov & Husain, 2014; Schneegans & Bays, 2017; Treisman & Zhang, 2006). Likewise, spatial locations may act as a medium for object selection and prioritisation from working memory (Griffin & Nobre, 2003; Kuo et al., 2009; Theeuwes et al., 2011; van Ede & Nobre, 2023), and may serve 'spontaneous' object rehearsal in working memory (Foster et al., 2017). It has remained unclear, however, whether location-based rehearsal of visual object features other than location relies on the brain's oculomotor system.

The aim of the current study was to address whether the oculomotor system contributes to working memory through location-based rehearsal of visual objects, even when object location is never asked about. To address this, we capitalised on directional biases in microsaccades (cf. Engbert & Kliegl, 2003; Hafed & Clark, 2002; Liu et al., 2022) as a direct and time-resolved read-out of oculomotor engagement during working-memory rehearsal of two visual objects. By presenting memory objects in different spatial configurations at encoding, we were able to isolate and track object rehearsal through the incidental locations of the memoranda. Furthermore, by comparing directional microsaccade biases across different task demands with the same spatial configurations (Experiment 2), we could disentangle the contribution of working-memory rehearsal from passive and attentive encoding of the same visual objects in the same configurations. Doing so, we reveal oculomotor rehearsal of visual objects in working memory through their associated – but in our case always task-irrelevant – locations.

METHODS
Participants
We conducted two experiments with 25 healthy human volunteers in each (Experiment 1: mean age = 21.96 years, age range = 18-28 years, 13 women; 1 left-handed; Experiment 2, mean age = 21.76 years, age range = 18-37 years, 17 women; 1 left-handed). Sample size of 25 per experiment was determined a-priori based on the use of the same sample size in relevant prior studies from the lab that used the same outcome measure

to target complementary questions (de Vries et al., 2023; Liu et al., 2022; van Ede et al., 2019, 2020, 2021). The recruitment of participants for the two experiments was conducted independently. Data from all participants were included in the analysis. All participants received course credit or monetary compensation (€10 per hour), provided written informed consent prior to participation, and reported normal or corrected-to-normal vision.

Task and procedure Experiment 1

Experiment 1 used a within-subject design in which we manipulated stimulus configuration at encoding. Participants performed a visual working memory task (see **Fig. 1a**), in which they memorised the orientation and colour of two objects presented on opposite points around a central fixation dot. The key manipulation was the variation of the stimulus configuration at encoding (horizontal, vertical, or either of two diagonal configurations; see **Fig. 2a**, top). This manipulation enabled us to investigate whether microsaccades during the subsequent delay-interval were more closely aligned with the configurational axes of the memory content, as opposed to the orthogonal axes – even if object location was incidental to the task because a location was never asked about.

Each trial started with a one-second intertrial interval that featured a small fixation dot, followed by the simultaneous presentation of two memory objects for 200 ms and a delay-interval of 2800 ms. The colour of the fixation dot then changed for 1000 ms serving as a retrocue to instruct (with 100% validity) which visual memory object would be probed after the second delay interval (1000 ms). The probe was always tilted at an angle of ±20 degrees relative to the cued memory target. The probe was always black and remained on the center of the screen until a response was provided. Participants had to match the orientation of the probe with that of the cued memory object by pressing the F key (counterclockwise) or the J key (clockwise).

The experimental stimuli were generated and presented on an LCD monitor with a solid grey background (ASUS ROG Strix XG248; 1920 x 1080 pixels; 240 Hz refresh rate) using PsychToolbox (Brainard, 1997) for MATLAB. Participants were seated in a dimly lit room with their heads resting on a chin rest at approximately 70 cm from the display. The central fixation dot was black and approximately 0.3 degrees visual angle. The memoranda had a diameter of 1.6 degrees, and were presented with a horizontal and vertical displacement of 6.4 degrees. The orientation of each stimulus was chosen randomly. The configuration (horizontal, vertical, and diagonal), colour (red and blue; 21/165/234; 234/74/21), and target position of the memoranda were varied pseudo-randomly to ensure that all these features were distributed equally within each block. The duration of each experimental session, which had 12 blocks of 36 trials each (totaling 432 trials), was roughly 70 minutes. Participants prepared for the experiment by practising 36 trials.

Task and procedure Experiment 2

Experiment 2 was designed to determine whether the directional microsaccade bias observed in Experiment 1 reflected the active location-based rehearsal of memory content, or instead reflected a lingering trace from encoding. To this end, Experiment 2 followed the same overall set-up as Experiment 1 but introduced different task demands, that were blocked. Moreover, to reduce factors, in Experiment 2, we always presented objects in either of the two diagonal encoding configurations.

In the passive viewing blocks (**Fig. 1b**), participants were instructed to ignore the stimuli, while maintaining central fixation. In the perceptual-judgement blocks (**Fig. 1c**), participants had to indicate whether the orientations of the two objects that were concurrently presented in any given display were identical (~7% of encoding displays) by pressing the spacebar. The task in the working-memory blocks (**Fig. 1d**) was the same as Experiment 1, only now the colour cue (informing the target memory object) was integrated into the probe by making the probe itself match the colour of either memory object. The orientation of each stimulus was chosen randomly, but, unlike Experiment 1, target and non-target objects always had a difference in orientation of at least 20 degrees (except in identical-target trials that served as targets in the perception task). This separation was necessary to account for the fact that, in perception blocks, participants had to be able to differentiate between targets (identical orientations) from non-target (non-identical orientations). Participants completed 24 blocks, each with 48 trials (totalling 1152 trials). The order of the blocks was randomised in sets of three (i.e. passive, perception, memory). Each block began with a brief overview of the instructions

for the upcoming task. Participants practised 32 trials of each of the three task types in preparation for the
experiment.

Figure 1. Task schematics of Experiment 1 (panel a) and Experiment 2 (panels b-d). In every case, trials started with the
bilateral presentation of two sample gratings, followed by a delay interval. **a)** In Experiment 1, participants indicated whether
a probe was oriented clockwise or counterclockwise with respect to the target memory object that was indicated by a colour
retrocue presented during the delay. In Experiment 1, the two visual objects could occupy one of four configurations at
encoding (horizontal, vertical, and either of two diagonal configurations). **b)** In the passive viewing condition of Experiment
2, participants were instructed to ignore the stimuli, while maintaining central fixation. **c)** In the perceptual-judgement
condition of Experiment 2, participants had to indicate whether the orientations of the two simultaneously presented gratings
in a given display were identical. **d)** The task in the memory condition of Experiment 2 was the same as in the first
experiment, only now the cue was integrated into the probe, by virtue of the probe's colour. In Experiment 2, the two visual
objects always occupied either of the two diagonal configurations at encoding.

Data acquisition & preprocessing

131
The right eye position was continuously recorded with an EyeLink 1000 Plus (SR Research, Ltd.) at a
sampling rate of 1000 Hz. The eye-tracking data were converted into ASCII format and analysed with
Matlab and Fieldtrip. Eye blinks were detected with an algorithm based on fluctuations in the pupil data
(Hershman et al., 2018) and linearly interpolated. The data were then segmented into epochs from -500 ms to
3000 ms relative to encoding onset. We estimated participants' attentiveness to the memory task by
measuring reaction times and excluding trials with responses slower than the mean reaction time +4 standard
deviations (following an iterative procedure until no more outliers remained). We also did not analyse trials
from perception blocks that contained targets (identical orientations) or were misidentified as such (false
detection). Following these criteria, only a small percentage of trials were eliminated in each experiment
(Experiment 1: $M = 0.015\%$, SD = 0.008%; Experiment 2: $M = 0.36\%$, $SD = 0.008\%$).

Eye-tracking analysis

142
To detect gaze shifts (saccades) we build on an existing velocity-based detection approach that has been
successfully employed and validated in prior studies from our lab (de Vries et al., 2023; Liu et al., 2022,
2023). Saccade detection was achieved by calculating the euclidean distance between successive gaze points
in the horizontal and vertical planes, smoothing the resulting velocity vector (using the built-in "smoothdata"
function in MATLAB, with a 7 ms sliding window), and identifying the onset of a saccade as the instant that
the velocity exceeds a predetermined threshold. A trial-based threshold of 5 times the median velocity was

employed to determine saccade onset, and a minimum interval of 100 milliseconds was used between
saccades to prevent multiple detections of the same saccade.

To index the incidental use of locations for working memory, we tracked the direction (angle) of saccades
and classified them as "aligned" when their direction was closer to the configurational axis of the visual
objects during encoding, and "misaligned" when their direction was closer to the orthogonal axis. Every
detected saccade was thus either classified as aligned or as misaligned. The resulting aligned and misaligned
saccade-rate time courses (expressed in Hz) were then smoothed (using the built-in function "smoothdata" in
MATLAB) using a moving average with a 100 ms sliding window. Crucially, directions classified as aligned
in one configuration (e.g., vertical stimulus arrangement) would be classified as misaligned in the
complementary configuration (e.g., horizontal stimulus arrangement). Accordingly, by collapsing across
configurations, we were able to isolate the effect of stimulus configuration by subtracting out any
background preferred saccade directions that would be common across all configurations.

Statistical analyses
To assess whether the oculomotor system is involved in the rehearsal of visual objects in working memory,
we compared the time courses of aligned and misaligned saccade rates. This was statistically evaluated using
a cluster-based permutation approach (Maris & Oostenveld, 2007), which effectively avoids the issue of
multiple comparisons by evaluating the full dataspace under a single permutation distribution of the largest
cluster. It groups neighbouring time points that show significant differences between the two time-courses
into a cluster and then randomly permutes the condition labels to create a null distribution of the largest
cluster that would be observed by chance. If the observed cluster size is larger than what would be observed
by chance, the two time-courses differ statistically. We used the Fieldtrip toolbox (Oostenveld et al., 2010) to
identify significant clusters using the default configuration settings (10,000 permutations; alpha level of
0.025 per side).

In Experiment 2, we additionally compared the directional microsaccade biases from our working-memory
condition of interest to each of the control conditions (passive, perception) using paired sample t-tests. For
this comparison, we extracted the effect of interest as the difference in aligned vs. misaligned saccades rates,
averaged across the full delay interval, as determined a-priori.

Data availability
Raw data and relevant code will be made publicly available upon publication.

RESULTS
Experiment 1: Microsaccades reveal location-based oculomotor rehearsal in visual working memory
The purpose of Experiment 1 (**Fig. 1a**) was to investigate whether the oculomotor system is involved in the
maintenance of visual objects in working memory, even when object location is never asked about. The
incidental use of location for working memory was tracked by classifying saccades as "aligned" when their
direction was closer to the configurational axis of the visual objects at encoding, and "misaligned" when
their direction was closer to the orthogonal axis.

Figure 2a depicts the distribution of saccade sizes across the full 2.8-second delay period for trials with
cardinal and diagonal stimulus configurations. The majority of saccades in the delay period were less than 1
degree visual angle (i.e. microsaccades; Engbert, 2006; Hafed & Clark, 2002; Martinez-Conde et al., 2004;
Nyström et al., 2016; Poletti & Rucci, 2016; Rolfs, 2009) and did not revisit the initially encoded locations
of the memory objects (which were presented at a distance of 6.4 degrees).

Our main result concerns the direction of the identified saccades as a function of memory-object
configurations at encoding. **Figure 2c** depicts the polar distribution of all saccades detected during the delay
period. The coloured bins correspond to saccades along the aligned axis, while the grey bins correspond to
saccades along the misaligned axis. Taking all of the saccades during the memory delay into account reveals
how saccade directions were generally biased along the axis of the visual objects at encoding (i.e. the aligned
axis). While there is generally a horizontal bias in cardinal configurations (**Fig. 2c**, left panels; consistent
with e.g., Engbert & Kliegl, 2003; Hermens & Walker, 2010), we found a configural memory bias on top of

this general preference for horizontal saccades. This memory-configuration-specific bias is especially evident when comparing the two diagonal configurations, where the data is less distorted by the background horizontal bias (**Fig. 2c**, right panels).

Figure 2. A directional microsaccade bias during working-memory maintenance reveals location-based object rehearsal. a) Spatial lay-out of the four stimulus configurations that were used in Experiment 1, with two solid black circles marking the locations of each stimulus pair at encoding **b)** Saccade-size distributions of delay-period saccades reveal that eye movements did not revisit the encoding locations of the memoranda. **c)** Polar distributions of delay-period saccades, with the coloured bins corresponding to saccades along the aligned axis, and the grey bins corresponding to saccades along the misaligned axis. **d)** The time courses of the saccade rates (in Hz) for saccades along the aligned and misaligned axes. The horizontal lines at the bottom highlight statistically significant clusters. Shaded lines represent ±1 standard error of the mean. The two vertical dotted lines mark stimulus onset and stimulus offset.

To quantify this bias and track its evolution over time, we collapsed across horizontal/vertical and across the two diagonal configurations and counted saccades along aligned versus misaligned axes (relative to encoding configurations) as a function of time. Crucially, directions that were classified as aligned in one configuration would be classified as misaligned in the complementary configuration. Accordingly, by collapsing configurations, we could isolate condition-specific saccadic biases by subtracting out any 'background' saccade-direction preferences that would be common to all configurations. Further, note how the probe was always presented centrally, so any directional-microsaccade bias could not reflect probe anticipation.

Figure 2d shows the saccade rates across time for the aligned (coloured) and misaligned (grey) axes. A cluster-based permutation comparison of the time courses confirmed significantly higher saccade rates on the aligned than on the misaligned axis (indicated by the horizontal lines in **Fig. 2d**, cluster P's < 0.001, 0.01, and 0.012 for the cardinal-configuration data; cluster P's < 0.001 and 0.003 for the diagonal-configuration

data). Importantly, this was not only evident during encoding, but lasted for a considerable period of time
well into the working-memory delay interval.

Thus, these data show that during working-memory maintenance of visual objects, saccades – particularly
microsaccades – occur more frequently along the spatial axis where memoranda were incidentally encoded
than along the orthogonal axis (where no stimuli were encoded). This is consistent with object rehearsal by
the oculomotor system through location, even when object location is never asked about.

Experiment 2: The microsaccade bias during the delay reflects working-memory demands
In Experiment 1, microsaccades continued to be biased along the aligned axis well into working-memory
delay. However, merely demonstrating that microsaccades remained biased in their direction during the delay
does not necessarily imply that they reflect active memory maintenance, as such biases could also reflect
residual, lingering consequences from prior sensory encoding.

Figure 3. Directional bias in microsaccades is driven by working-memory demands. Spatial lay-out of the stimulus
configurations that were used in Experiment 2, with the locations of each stimulus pair indicated by the two solid black
circles. **b)** Saccade-size distributions of delay-period saccades. **c)** Polar distributions of delay-period saccades, with coloured
bins representing saccades along the aligned axis and grey bins representing saccades along the misaligned axis. **d)** The time
courses of the saccade rates (in Hz) for saccades along the aligned and misaligned axes. The horizontal lines at the bottom
highlight clusters that are statistically significant. Shaded lines represent ±1 standard error of the mean. The two vertical
dashed lines denote the onset and offset of the stimulus.

Experiment 2 (**Fig. 1b-d**) sought to replicate the microsaccade bias during the memory interval (this time
using only the two diagonal conditions; **Fig 3a**) and to compare it to two control conditions in which
participants either (1) passively viewed the same objects, or (2) were required to make a perceptual
judgement on the two objects (detect instances where both simultaneously presented objects had identical
orientations) without subsequent working-memory demands. If the bias truly reflects active working-
memory maintenance (rehearsal) processes, it should only be profound in our memory task, and not after

passive presentation, or perceptual evaluation of the same objects, that we presented in the same
configurations, followed by the same delay.

Figure 3b shows the distribution of saccade magnitudes for passive, perceptual, and memory trials. As in
Experiment 1, the vast majority of saccades occurred within the microsaccade range and again this was the
case for all conditions. **Figure 3c** depicts the polar distribution of saccades during the delay period, again
showing a clear bias on memory trials (in blue, on the right), but little evidence of a similar bias for the two
control conditions with passive object presentation (in red, left) or the perceptual judgement task (in green,
middle).

To quantify the bias and track its temporal profile, we again counted saccades along the aligned and
misaligned axes, as defined by the incidental stimulus configurations at encoding. **Figure 3d** shows that
saccades in passive blocks are equally likely to occur along the aligned and misaligned axes (cluster $P > 0.3$),
indicating that stimulus presentation alone cannot account for the reported directional saccade bias in
working-memory trials. In perception blocks, we discovered a slight bias for saccades to occur more along
the aligned axis (cluster P's = 0.01 and 0.019), but this bias was relatively weak and appeared to persist for
only about 1 second after stimulus onset, which was also the time limit for participants to indicate that the
two objects were identical.

In comparison, and replicating the results of Experiment 1, the memory condition again reveals a profound
directional bias that persisted well into the working-memory delay period (cluster P's < 0.001, 0.007, and
0.01). A direct comparison of the memory condition with both control conditions confirmed that the memory
condition had a significantly stronger bias in microsaccade direction: the saccade rate effect (aligned minus
misaligned saccades rates, averaged across the full a-priori defined delay period) was significantly larger in
memory blocks ($M = 0.19$, $SD = 0.11$), than in passive blocks ($M = 0.01$, $SD = 0.03$), $t(24) = 7.99$, $p < 0.001$,
and perception blocks ($M = 0.05$, $SD = 0.06$), $t(24) = 6.07$, $p < 0.001$.

These findings rule out that the observed saccade bias during working-memory maintenance is due to
lingering processing of the passive viewing or attentive encoding of salient visual objects per se. Instead, the
prolonged directional saccade bias in the memory task must be attributed to working-memory demands,
which is consistent with an oculomotor-driven rehearsal of memorised visual object features (colour and
orientation) through their memorised locations.

DISCUSSION
It has been established that the brain's oculomotor system not only controls where we look, but also
participates in the rehearsal of spatial-location information in working memory (as reviewed in: Awh et al.,
2006; Jonikaitis & Moore, 2019; Olsen et al., 2014; Souza et al., 2020; Theeuwes et al., 2009; Wynn et al.,
2018). It had been called into question, however, whether this role extends to visual working memory of
objects whose location is never asked about (see e.g., McAteer et al., 2023; Postle et al., 2006). By analysing
microsaccades as a direct and time-resolved output from the oculomotor system, we provide unique evidence
that the brain's oculomotor system also participates in the rehearsal of visual objects in working memory,
even when object location is never asked about.

Experiment 1 revealed how microsaccade directions remained biased during the working-memory delay
along the axis at which objects were incidentally presented at encoding. Experiment 2 replicated this central
finding and ruled out lingering effects from passive or attentive encoding. By demonstrating directional
biases in microsaccades during working-memory maintenance, we thus reveal rehearsal of memorised
visual-object features through their incidentally encoded memorised locations – and implicate the
oculomotor system in such rehearsal. In the following, we consider and discuss three central questions raised
by these findings.

What may the observed microsaccade bias during working-memory maintenance reflect?
Spatial biases in microsaccades have previously been linked to covert shifts of attention to peripheral
locations (Engbert & Kliegl, 2003; Hafed & Clark, 2002; Yuval-Greenberg et al., 2014), including when
directing attention within the spatial-layout of visual working memory (e.g., de Vries et al., 2023; Liu et al.,

2022; van Ede et al., 2019). Accordingly, our findings of directional microsaccade biases during working-memory maintenance may reflect attentional rehearsal of memory representations. Though the location was never asked about, it may still facilitate rehearsal. This is consistent with prior research demonstrating that space remains important for organising working-memory content, even when location is never asked about (e.g., de Vries et al., 2023; Kuo et al., 2009; Liu et al., 2022; Theeuwes et al., 2011; van Ede et al., 2019). Our data go beyond this existing body of work by implicating the brain's oculomotor system in such incidental use of location, and by demonstrating this role during working-memory maintenance (extending prior work implicating the oculomotor system in mnemonic selection; see e.g., de Vries et al., 2023; Liu et al., 2022; van Ede et al., 2019).

By focusing on object rehearsal through location, our findings are also distinct from complementary recent studies that have linked directional biases in microsaccades to the rehearsal of visual shape and orientation information (Dotson et al., 2018; Linde-Domingo & Spitzer, 2022; Mostert et al., 2018; Quax et al., 2019; Thielen et al., 2019). In contrast to these studies where microsaccades directly tracked the target-relevant memory feature (shape or orientation), we used microsaccades to track incidental object-locations that were never queried for report.

Consistent with a role of the oculomotor system in object rehearsal, complementary dual-task experiments have revealed enhanced memory for objects whose incidental locations became saccade targets during the working-memory delay (Hanning et al., 2016; Ohl & Rolfs, 2020). This suggests some type of "refreshing" of memory objects when they become the target of a saccade. Our microsaccade observations were made without asking participants to make any eye movements, and persisted for several seconds after encoding. Accordingly, our microsaccade observations may signal serial refreshing by the oculomotor system that may serve as a mechanism supporting object maintenance. In future studies, it will be interesting to address whether and how these microsaccade findings relate to recent complementary findings of serial object-sampling in working memory (e.g., Chota et al., 2022; Peters et al., 2021; Pomper & Ansorge, 2021).

Why does the observed microsaccade-maintenance bias diminish with time?
Experiment 2 demonstrated that the directional microsaccade bias was specific to memory demands, ruling out lingering effects from passive and attentive encoding. At the same time, despite ongoing memory demands, the bias diminished with time in the delay interval, and this was the case in both experiments. Interestingly, however, this appears not to be unique to the working-memory signature that we uncovered here. For example, the Contralateral Delay Activity (an EEG signature frequently implicated in working-memory maintenance; e.g., Luria et al., 2016; McCollough et al., 2007; Vogel & Machizawa, 2004), similarly decays during the memory delay (e.g., Hakim et al., 2020; Jin et al., 2020; Roy & Faubert, 2023).

Even so, the observed decay raises the intriguing question of why this may occur and what it may reflect. One possible explanation is that rehearsal demands become less urgent as time passes. Early after encoding, active rehearsal may facilitate consolidation, while later in the delay active rehearsal may be less critical. For example, with time, working-memory representations may become increasingly reliant on activity-silent mechanisms (through temporary changes in synaptic connectivity; Mongillo et al., 2008; Stokes, 2015), rather than states of active rehearsal. Another possibility is that over time, location information simply becomes less relevant for working-memory maintenance, as objects are gradually recoded to a less visual format, or shifted from peripheral to more foveal (e.g., Chambers et al., 2013; Fan et al., 2016) or global (Ester et al., 2009) cortical representations. Finally, it is conceivable that the anticipation of the probe stimulus itself may have affected our ability to measure robust microsaccade biases, since stimulus expectation is known to reduce overall microsaccade rates (Amit et al., 2019; Denison et al., 2019; Olmos-Solis et al., 2017; Tal-Perry & Yuval-Greenberg, 2021). It is worth noting that these explanations are not mutually exclusive, as multiple (as well as additional) factors may contribute jointly; providing a relevant avenue for future research.

What opportunities and challenges do these findings bring?
Our findings demonstrate how microsaccades can be used to track working-memory rehearsal through spatial location over time. This may open new avenues for characterising the spatial-temporal trajectories of working-memory maintenance in future studies with complementary manipulations of spatial and/or

temporal working-memory demands. We have previously demonstrated that microsaccades are a valuable tool for studying attentional selection and prioritisation of objects in visual working memory (e.g., de Vries et al., 2023; Liu et al., 2022; van Ede et al., 2019). Here, we extend this involvement to the maintenance period. Zooming out, these findings add to a growing body of work showing the utility of using microsaccades to track cognition at large, including for tracking complementary cognitive functions such as temporal expectations (Amit et al., 2019; Denison et al., 2019; Olmos-Solis et al., 2017; Tal-Perry & Yuval-Greenberg, 2021), perceptual learning (Hung & Carrasco, 2022) and decision making (Loughnane et al., 2018).

At the same time, our data bring a practical implication to the fore that presents a challenge, rather than an opportunity. They show how microsaccades cannot be neglected in neuroscience studies of visual working memory. Several recent reports already made clear how microsaccades may confound neural decoding of memory representations, by showing how microsaccades can be systematically biased by task-relevant object-features such as shape and orientation (Dotson et al., 2018; Linde-Domingo & Spitzer, 2022; Mostert et al., 2018; Quax et al., 2019; Thielen et al., 2019). Building on this work, our data make clear how systematic biases in microsaccades can be a concern also for arbitrarily chosen spatial stimulus-configurations at encoding, even if stimulus configurations are never explicitly relevant for the participant.

Conclusion

In conclusion, by studying microsaccades, we provide unique evidence that the brain's oculomotor system participates in rehearsing visual objects in working memory through their incidental locations. The microsaccade-maintenance bias we observed may signal an attentional refreshing mechanism by the oculomotor system that supports visual object maintenance. At the same time, the observed decay in the observed microsaccade-maintenance bias raises intriguing questions about the dynamic nature of working-memory rehearsal across time.

ACKNOWLEDGEMENTS

This research was supported by an ERC Starting Grant from the European Research Council (MEMTICIPATION, 850636) and an NWO Vidi grant by the Dutch Research Council (grant number 14721) to F.v.E. The authors thank Anna van Harmelen and Baiwei Liu for their valuable comments on the manuscript.

REFERENCES

Abrahamse, E., van Dijck, J.-P., Majerus, S., & Fias, W. (2014). Finding the answer in space: The mental whiteboard hypothesis on serial order in working memory. *Frontiers in Human Neuroscience*, *8*. https://doi.org/10.3389/fnhum.2014.00932

Amit, R., Abeles, D., Carrasco, M., & Yuval-Greenberg, S. (2019). Oculomotor inhibition reflects temporal expectations. *NeuroImage*, *184*, 279–292. https://doi.org/10.1016/j.neuroimage.2018.09.026

Armstrong, K. M., Chang, M. H., & Moore, T. (2009). Selection and Maintenance of Spatial Information by Frontal Eye Field Neurons. *Journal of Neuroscience*, *29*(50), 15621–15629. https://doi.org/10.1523/JNEUROSCI.4465-09.2009

Awh, E., Armstrong, K. M., & Moore, T. (2006). Visual and oculomotor selection: Links, causes and implications for spatial attention. *Trends in Cognitive Sciences*, *10*(3), 124–130. https://doi.org/10.1016/j.tics.2006.01.001

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436. https://doi.org/10.1163/156856897X00357

Chambers, C. D., Allen, C. P., Maizey, L., & Williams, M. A. (2013). Is delayed foveal feedback critical for extra-foveal perception? *Cortex*, *49*(1), 327–335.

Chota, S., Leto, C., van Zantwijk, L., & Van der Stigchel, S. (2022). Attention rhythmically samples multi-feature objects in working memory. *Scientific Reports*, *12*(1), Article 1. https://doi.org/10.1038/s41598-022-18819-z

Corbetta, M., Akbudak, E., Conturo, T. E., Snyder, A. Z., Ollinger, J. M., Drury, H. A., Linenweber, M. R., Petersen, S. E., Raichle, M. E., Van Essen, D. C., & Shulman, G. L. (1998). A Common Network of Functional Areas for Attention and Eye Movements. *Neuron*, *21*(4), 761–773. https://doi.org/10.1016/S0896-6273(00)80593-0

Courtney, S. M., Petit, L., Maisog, J. Ma., Ungerleider, L. G., & Haxby, J. V. (1998). An Area Specialized for Spatial Working Memory in Human Frontal Cortex. *Science*, *279*(5355), 1347–1351. https://doi.org/10.1126/science.279.5355.1347

Curtis, C. E., Rao, V. Y., & D'Esposito, M. (2004). Maintenance of Spatial and Motor Codes during Oculomotor Delayed Response Tasks. *Journal of Neuroscience*, *24*(16), 3944–3952. https://doi.org/10.1523/JNEUROSCI.5640-03.2004

de Vries, E., Fejer, G., & van Ede, F. (2023). *No trade-off between the use of space and time for working memory* (p. 2023.01.20.524861). bioRxiv. https://doi.org/10.1101/2023.01.20.524861

Denison, R. N., Yuval-Greenberg, S., & Carrasco, M. (2019). Directing Voluntary Temporal Attention Increases Fixational Stability. *Journal of Neuroscience*, *39*(2), 353–363. https://doi.org/10.1523/JNEUROSCI.1926-18.2018

Dotson, N. M., Hoffman, S. J., Goodell, B., & Gray, C. M. (2018). Feature-Based Visual Short-Term Memory Is Widely Distributed and Hierarchically Organized. *Neuron*, *99*(1), 215-226.e4. https://doi.org/10.1016/j.neuron.2018.05.026

Engbert, R. (2006). Microsaccades: A microcosm for research on oculomotor control, attention, and visual perception. In *Progress in Brain Research* (Vol. 154, pp. 177–192). Elsevier. https://doi.org/10.1016/S0079-6123(06)54009-9

Engbert, R., & Kliegl, R. (2003). Microsaccades uncover the orientation of covert attention. *Vision Research*, *43*(9), 1035–1045. https://doi.org/10.1016/S0042-6989(03)00084-1

Ester, E. F., Serences, J. T., & Awh, E. (2009). Spatially Global Representations in Human Primary Visual Cortex during Working Memory Maintenance. *Journal of Neuroscience*, *29*(48), 15258–15265. https://doi.org/10.1523/JNEUROSCI.4388-09.2009

Fan, X., Wang, L., Shao, H., Kersten, D., & He, S. (2016). Temporally flexible feedback signal to foveal cortex for peripheral object recognition. *Proceedings of the National Academy of Sciences*, *113*(41), 11627–11632. https://doi.org/10.1073/pnas.1606137113

Foster, J. J., Bsales, E. M., Jaffe, R. J., & Awh, E. (2017). Alpha-Band Activity Reveals Spontaneous Representations of Spatial Position in Visual Working Memory. *Current Biology*, *27*(20), 3216-3223.e6. https://doi.org/10.1016/j.cub.2017.09.031

Godijn, R., & Theeuwes, J. (2012). Overt is no better than covert when rehearsing visuo-spatial information in working memory. *Memory & Cognition*, *40*(1), 52–61. https://doi.org/10.3758/s13421-011-0132-x

Griffin, I. C., & Nobre, A. C. (2003). Orienting attention to locations in internal representations. *Journal of Cognitive Neuroscience*, *15*(8), 1176–1194.

Hafed, Z. M., Chen, C.-Y., & Tian, X. (2015). Vision, Perception, and Attention through the Lens of Microsaccades: Mechanisms and Implications. *Frontiers in Systems Neuroscience*, *9*, 167. https://doi.org/10.3389/fnsys.2015.00167

Hafed, Z. M., & Clark, J. J. (2002). Microsaccades as an overt measure of covert attention shifts. *Vision Research*, *42*(22), 2533–2545. https://doi.org/10.1016/S0042-6989(02)00263-8

Hakim, N., Feldmann-Wüstefeld, T., Awh, E., & Vogel, E. K. (2020). Perturbing Neural Representations of Working Memory with Task-irrelevant Interruption. *Journal of Cognitive Neuroscience*, *32*(3), 558–569. https://doi.org/10.1162/jocn_a_01481

Hanning, N. M., Jonikaitis, D., Deubel, H., & Szinte, M. (2016). Oculomotor selection underlies feature retention in visual working memory. *Journal of Neurophysiology*, *115*(2), 1071–1076. https://doi.org/10.1152/jn.00927.2015

Hermens, F., & Walker, R. (2010). What determines the direction of microsaccades? *Journal of Eye Movement Research*, *3*(4), 1–20.

Hershman, R., Henik, A., & Cohen, N. (2018). A novel blink detection method based on pupillometry noise. *Behavior Research Methods*, *50*(1), 107–114. https://doi.org/10.3758/s13428-017-1008-1

Hollingworth, A. (2007). Object-position binding in visual memory for natural scenes and object arrays. *Journal of Experimental Psychology: Human Perception and Performance*, *33*(1), 31.

Hung, S.-C., & Carrasco, M. (2022). Microsaccades as a long-term oculomotor correlate in visual perceptual learning. *Psychonomic Bulletin & Review*. https://doi.org/10.3758/s13423-022-02151-8

Jerde, T. A., Merriam, E. P., Riggall, A. C., Hedges, J. H., & Curtis, C. E. (2012). Prioritized Maps of Space in Human Frontoparietal Cortex. *Journal of Neuroscience*, *32*(48), 17382–17390. https://doi.org/10.1523/JNEUROSCI.3810-12.2012

Jin, W., Nobre, A. C., & van Ede, F. (2020). Temporal Expectations Prepare Visual Working Memory for Behavior. *Journal of Cognitive Neuroscience*, *32*(12), 2320–2332. https://doi.org/10.1162/jocn_a_01626

Jonikaitis, D., & Moore, T. (2019). The interdependence of attention, working memory and gaze control: Behavior and neural circuitry. *Current Opinion in Psychology*, *29*, 126–134. https://doi.org/10.1016/j.copsyc.2019.01.012

Krauzlis, R. J., Lovejoy, L. P., & Zénon, A. (2013). Superior colliculus and visual spatial attention. *Annual Review of Neuroscience*, *36*, 165–182. https://doi.org/10.1146/annurev-neuro-062012-170249

Kuo, B.-C., Rao, A., Lepsien, J., & Nobre, A. C. (2009). Searching for targets within the spatial layout of visual short-term memory. *Journal of Neuroscience*, *29*(25), 8032–8038.

Lawrence, B. M., Myerson, J., & Abrams, R. A. (2004). Interference with spatial working memory: An eye movement is more than a shift of attention. *Psychonomic Bulletin & Review*, *11*(3), 488–494. https://doi.org/10.3758/BF03196600

Linde-Domingo, J., & Spitzer, B. (2022). *Geometry of visual working memory information in human gaze patterns*. bioRxiv. https://doi.org/10.1101/2022.11.17.516917

Liu, B., Alexopoulou, Z.-S., & Ede, F. van. (2023). *Jointly looking to the past and the future in visual working memory* (p. 2023.01.30.526235). bioRxiv. https://doi.org/10.1101/2023.01.30.526235

Liu, B., Nobre, A. C., & van Ede, F. (2022). Functional but not obligatory link between microsaccades and neural modulation by covert spatial attention. *Nature Communications*, *13*(1), Article 1. https://doi.org/10.1038/s41467-022-31217-3

Loaiza, V. M., & Souza, A. S. (2022). The eyes don't have it: Eye movements are unlikely to reflect refreshing in working memory. *PLOS ONE*, *17*(7), e0271116. https://doi.org/10.1371/journal.pone.0271116

Loughnane, G. M., Newman, D. P., Tamang, S., Kelly, S. P., & O'Connell, R. G. (2018). Antagonistic Interactions Between Microsaccades and Evidence Accumulation Processes During Decision Formation. *Journal of Neuroscience*, *38*(9), 2163–2176. https://doi.org/10.1523/JNEUROSCI.2340-17.2018

Luria, R., Balaban, H., Awh, E., & Vogel, E. K. (2016). The contralateral delay activity as a neural measure of visual working memory. *Neuroscience & Biobehavioral Reviews*, *62*, 100–108. https://doi.org/10.1016/j.neubiorev.2016.01.003

Manohar, S. G., Pertzov, Y., & Husain, M. (2017). Short-term memory for spatial, sequential and duration information. *Current Opinion in Behavioral Sciences*, *17*, 20–26. https://doi.org/10.1016/j.cobeha.2017.05.023

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-data. *Journal of Neuroscience Methods*, *164*(1), 177–190. https://doi.org/10.1016/j.jneumeth.2007.03.024

Marshall, T. R., O'Shea, J., Jensen, O., & Bergmann, T. O. (2015). Frontal eye fields control attentional modulation of alpha and gamma oscillations in contralateral occipitoparietal cortex. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *35*(4), 1638–1647. https://doi.org/10.1523/JNEUROSCI.3116-14.2015

Martinez-Conde, S., Macknik, S. L., & Hubel, D. H. (2004). The role of fixational eye movements in visual perception. *Nature Reviews Neuroscience*, *5*(3), 229–240. https://doi.org/10.1038/nrn1348

McAteer, S. M., McGregor, A., & Smith, D. T. (2023). Oculomotor rehearsal in visuospatial working memory. *Attention, Perception & Psychophysics*, *85*(1), 261–275. https://doi.org/10.3758/s13414-022-02601-4

McCollough, A. W., Machizawa, M. G., & Vogel, E. K. (2007). Electrophysiological Measures of Maintaining Representations in Visual Working Memory. *Cortex*, *43*(1), 77–94. https://doi.org/10.1016/S0010-9452(08)70447-7

Merrikhi, Y., Clark, K., Albarran, E., Parsa, M., Zirnsak, M., Moore, T., & Noudoost, B. (2017). Spatial working memory alters the efficacy of input to visual cortex. *Nature Communications*, *8*(1), 15041. https://doi.org/10.1038/ncomms15041

Mongillo, G., Barak, O., & Tsodyks, M. (2008). Synaptic Theory of Working Memory. *Science*, *319*(5869), 1543–1546. https://doi.org/10.1126/science.1150769

Moore, T., & Armstrong, K. M. (2003). Selective gating of visual signals by microstimulation of frontal cortex. *Nature*, *421*(6921), Article 6921. https://doi.org/10.1038/nature01341

Moore, T., Armstrong, K. M., & Fallah, M. (2003). Visuomotor Origins of Covert Spatial Attention. *Neuron*, *40*(4), 671–683. https://doi.org/10.1016/S0896-6273(03)00716-5

Mostert, P., Albers, A. M., Brinkman, L., Todorova, L., Kok, P., & de Lange, F. P. (2018). Eye Movement-Related Confounds in Neural Decoding of Visual Working Memory Representations. *Eneuro*, *5*(4), ENEURO.0401-17.2018. https://doi.org/10.1523/ENEURO.0401-17.2018

Nobre, A. C., Gitelman, D. R., Dias, E. C., & Mesulam, M. M. (2000). Covert Visual Spatial Orienting and Saccades: Overlapping Neural Systems. *NeuroImage*, *11*(3), 210–216. https://doi.org/10.1006/nimg.2000.0539

Nyström, M., Hansen, D. W., Andersson, R., & Hooge, I. (2016). Why have microsaccades become larger? Investigating eye deformations and detection algorithms. *Vision Research*, *118*, 17–24. https://doi.org/10.1016/j.visres.2014.11.007

Ohl, S., & Rolfs, M. (2020). Bold moves: Inevitable saccadic selection in visual short-term memory. *Journal of Vision*, *20*(2), 11–11. https://doi.org/10.1167/jov.20.2.11

Olmos-Solis, K., van Loon, A. M., Los, S. A., & Olivers, C. N. (2017). Oculomotor measures reveal the temporal dynamics of preparing for search. *Progress in Brain Research*, *236*, 1–23. https://doi.org/10.1016/bs.pbr.2017.07.003

Olsen, R. K., Chiew, M., Buchsbaum, B. R., & Ryan, J. D. (2014). The relationship between delay period eye movements and visuospatial memory. *Journal of Vision*, *14*(1), 8–8. https://doi.org/10.1167/14.1.8

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2010). FieldTrip: Open Source Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data. *Computational Intelligence and Neuroscience*, *2011*, e156869. https://doi.org/10.1155/2011/156869

Pearson, D. G., Ball, K., & Smith, D. T. (2014). Oculomotor preparation as a rehearsal mechanism in spatial working memory. *Cognition*, *132*(3), 416–428. https://doi.org/10.1016/j.cognition.2014.05.006

Pertzov, Y., & Husain, M. (2014). The privileged role of location in visual working memory. *Attention, Perception, & Psychophysics*, *76*(7), 1914–1924. https://doi.org/10.3758/s13414-013-0541-y

Peters, B., Kaiser, J., Rahm, B., & Bledowski, C. (2021). Object-based attention prioritizes working memory contents at a theta rhythm. *Journal of Experimental Psychology: General*, *150*, 1250–1256. https://doi.org/10.1037/xge0000994

Poletti, M., & Rucci, M. (2016). A compact field guide to the study of microsaccades: Challenges and functions. *Vision Research*, *118*, 83–97. https://doi.org/10.1016/j.visres.2015.01.018

Pomper, U., & Ansorge, U. (2021). Theta-Rhythmic Oscillation of Working Memory Performance. *Psychological Science*, *32*(11), 1801–1810. https://doi.org/10.1177/09567976211013045

Postle, B. R., Berger, J. S., Taich, A. M., & D'esposito, M. (2000). Activity in human frontal cortex associated with spatial working memory and saccadic behavior. *Journal of Cognitive Neuroscience*, *12*(Supplement 2), 2–14.

Postle, B. R., Idzikowski, C., Sala, S. D., Logie, R. H., & Baddeley, A. D. (2006). The selective disruption of spatial working memory by eye movements. *Quarterly Journal of Experimental Psychology*, *59*(1), 100–120. https://doi.org/10.1080/17470210500151410

Quax, S. C., Dijkstra, N., van Staveren, M. J., Bosch, S. E., & van Gerven, M. A. J. (2019). Eye movements explain decodability during perception and cued attention in MEG. *NeuroImage*, *195*, 444–453. https://doi.org/10.1016/j.neuroimage.2019.03.069

Rizzolatti, G., Riggio, L., Dascola, I., & Umiltá, C. (1987). Reorienting attention across the horizontal and vertical meridians: Evidence in favor of a premotor theory of attention. *Neuropsychologia*, *25*(1, Part 1), 31–40. https://doi.org/10.1016/0028-3932(87)90041-8

Rolfs, M. (2009). Microsaccades: Small steps on a long way. *Vision Research*, *49*(20), 2415–2441. https://doi.org/10.1016/j.visres.2009.08.010

Roy, Y., & Faubert, J. (2023). Is the Contralateral Delay Activity (CDA) a robust neural correlate for Visual Working Memory (VWM) tasks? A reproducibility study. *Psychophysiology*, *60*(2), e14180. https://doi.org/10.1111/psyp.14180

Schneegans, S., & Bays, P. M. (2017). Neural Architecture for Feature Binding in Visual Working Memory. *Journal of Neuroscience*, *37*(14), 3913–3925. https://doi.org/10.1523/JNEUROSCI.3493-16.2017

Souza, A. S., Czoschke, S., & Lange, E. B. (2020). Gaze-based and attention-based rehearsal in spatial working memory. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *46*(5), 980–1003. https://doi.org/10.1037/xlm0000771

Stokes, M. G. (2015). 'Activity-silent' working memory in prefrontal cortex: A dynamic coding framework. *Trends in Cognitive Sciences*, *19*(7), 394–405. https://doi.org/10.1016/j.tics.2015.05.004

Tal-Perry, N., & Yuval-Greenberg, S. (2021). Prestimulus inhibition of eye movements reflects temporal expectation rather than time estimation. *Attention, Perception, & Psychophysics*, *83*(6), 2473–2485. https://doi.org/10.3758/s13414-021-02319-9

Theeuwes, J., Belopolsky, A., & Olivers, C. N. L. (2009). Interactions between working memory, attention and eye movements. *Acta Psychologica*, *132*(2), 106–114. https://doi.org/10.1016/j.actpsy.2009.01.005

Theeuwes, J., Kramer, A. F., & Irwin, D. E. (2011). Attention on our mind: The role of spatial attention in visual working memory. *Acta Psychologica*, *137*(2), 248–251. https://doi.org/10.1016/j.actpsy.2010.06.011

Thielen, J., Bosch, S. E., Leeuwen, T. M. van, Gerven, M. A. J. van, & Lier, R. van. (2019). Evidence for confounding eye movements under attempted fixation and active viewing in cognitive neuroscience. *Scientific Reports*, *9*(1), 1–8. https://doi.org/10.1038/s41598-019-54018-z

Treisman, A., & Zhang, W. (2006). Location and binding in visual working memory. *Memory & Cognition*, *34*(8), 1704–1719. https://doi.org/10.3758/BF03195932

Tremblay, S., Saint-Aubin, J., & Jalbert, A. (2006). Rehearsal in serial memory for visual-spatial information: Evidence from eye movements. *Psychonomic Bulletin & Review*, *13*(3), 452–457. https://doi.org/10.3758/bf03193869

van Ede, F., Board, A. G., & Nobre, A. C. (2020). Goal-directed and stimulus-driven selection of internal representations. *Proceedings of the National Academy of Sciences*, *117*(39), 24590–24598. https://doi.org/10.1073/pnas.2013432117

van Ede, F., Chekroud, S. R., & Nobre, A. C. (2019). Human gaze tracks attentional focusing in memorized visual space. *Nature Human Behaviour*, *3*(5), 462–470. https://doi.org/10.1038/s41562-019-0549-y

van Ede, F., Deden, J., & Nobre, A. C. (2021). Looking ahead in working memory to guide sequential behaviour. *Current Biology*, *31*(12), R779–R780. https://doi.org/10.1016/j.cub.2021.04.063

van Ede, F., & Nobre, A. C. (2023). Turning Attention Inside Out: How Working Memory Serves Behavior. *Annual Review of Psychology*, *74*(1), 137–165. https://doi.org/10.1146/annurev-psych-021422-041757

Vogel, E. K., & Machizawa, M. G. (2004). Neural activity predicts individual differences in visual working memory capacity. *Nature*, *428*(6984), Article 6984. https://doi.org/10.1038/nature02447

Williams, M., Pouget, P., Boucher, L., & Woodman, G. F. (2013). Visual–spatial attention aids the maintenance of object representations in visual working memory. *Memory & Cognition*, *41*(5), 698–715. https://doi.org/10.3758/s13421-013-0296-7

Wynn, J. S., Olsen, R. K., Binns, M. A., Buchsbaum, B. R., & Ryan, J. D. (2018). Fixation reinstatement supports visuospatial memory in older adults. *Journal of Experimental Psychology: Human Perception and Performance*, *44*(7), 1119–1127. https://doi.org/10.1037/xhp0000522

Yuval-Greenberg, S., Merriam, E. P., & Heeger, D. J. (2014). Spontaneous Microsaccades Reflect Shifts in Covert Attention. *The Journal of Neuroscience*, *34*(41), 13693–13700. https://doi.org/10.1523/JNEUROSCI.0582-14.2014