# Title: Microbially-derived indole-3-acetate alleviates diet induced steatosis and inflammation in mice

Yufang Ding[1], Karin Yanagi[2], Fang Yang[1], Evelyn Callaway[1], Clint Cheng[1], Martha E Hensel[3], Rani Menon[1], Robert C. Alaniz[4], Kyongbum Lee[2*], Arul Jayaraman[1,4*]

[1] Artie McFerrin Department of Chemical Engineering, Texas A&M University; College Station, TX 77843, USA

[2] Department of Chemical and Biological Engineering, Tufts University; Medford, MA 02155, USA

[3] Department of Pathobiology, College of Veterinary Medicine and Biomedical Sciences, Texas A&M University; College Station, TX, USA

[4] Department of Microbial Pathogenesis and Immunology, College of Medicine, Texas Health Science Center, Texas A&M University; Bryan, TX, USA

*Corresponding author. Kyongbum.Lee@tufts.edu, arulj@tamu.edu

**Abstract:** Non-alcoholic fatty liver disease (NAFLD) is the most common chronic liver disease in Western countries. There is growing evidence that dysbiosis of the intestinal microbiota and disruption of microbiota-host interactions contribute to the pathology of NAFLD. We previously demonstrated that gut microbiota derived tryptophan metabolite indole-3-acetate (I3A) was decreased in both cecum and liver of high-fat diet-fed mice and attenuated the expression of inflammatory cytokines in macrophages and TNF-a and fatty acid induced inflammatory responses in an aryl-hydrocarbon receptor (AhR) dependent manner in hepatocytes. In this study, we investigated the effect of orally administered I3A in a mouse model of diet induced NAFLD. Western diet (WD)-fed mice given sugar water (SW) with I3A showed dramatically decreased serum ALT, hepatic TG, liver steatosis, hepatocyte ballooning, lobular inflammation, and hepatic production of inflammatory cytokines, compared to WD-fed mice given only SW. Metagenomic analysis show that I3A administration did not significantly modify the intestinal microbiome, suggesting that I3A's beneficial effects likely reflect the metabolite's direct actions on the liver. Administration of I3A partially reversed WD induced alterations of liver metabolome and proteome, notably, decreasing expression of several enzymes in hepatic lipogenesis and β-oxidation. Mechanistically, we also show that AMP-activated protein kinase (AMPK) mediates the anti-inflammatory effects of I3A in macrophages. The potency of I3A in alleviating liver steatosis and inflammation clearly demonstrates its potential as a therapeutic modality for preventing the progression of steatosis to NASH.

## INTRODUCTION

Non-alcoholic fatty liver disease (NAFLD) is the most common chronic liver disease in Western countries, with a prevalence rate of 21-25% in North American and Europe [1]. It is a multi-stage disease [2] that can progress from liver steatosis (characterized by macrovesicular fat deposition), which is benign and reversible [2,3], to more severe forms of the disease such as non-alcoholic steatohepatitis (NASH) and fibrosis. Approximately 25% of individuals having liver steatosis develop NASH, which is characterized by liver inflammation, dysregulated lipid metabolism, and cell damage [3]. A subset of NASH patients develops cirrhosis and even hepatocellular carcinoma [4]. Although significant progress has been made in understanding the pathogenesis of NAFLD, factors leading to the progression from steatosis to NASH remain poorly understood, and there are currently no pharmacological treatments available for NASH.

There is growing evidence that dysbiosis of the intestinal microbiota and disruption of microbiota-host interactions contribute to the pathology of NASH [5-8]. Certain shifts in the intestinal microbiota community composition, e.g., expansion of the phyla *Verrucomicrobia* and *Proteobacteria,* correlate with NASH in both human and animal studies [9,10]. One potential mechanism linking intestinal microbiota dysbiosis and NASH is compromised intestinal barrier integrity, which promotes the translocation of bacterial products from the lumen to circulation. This can contribute directly and indirectly (through intestinal inflammation) to liver inflammation [11-13]. Additionally, microbial dysbiosis alters the balance of bioactive metabolites produced by gut bacteria such as bile acids, short chain fatty acids and aromatic amino acid derivatives, which have been shown to impact liver metabolism and inflammation in NAFLD through host receptor mediated pathways [13].

In our previous study [14], we demonstrated that gut microbiota derived tryptophan metabolites indole-3-acetate (I3A) and tryptamine (TA) were decreased in both cecum and liver of high-fat

diet (HFD)-fed mice compared to low-fat diet (LFD)-fed control mice. *In vitro*, both I3A and TA
attenuated the expression of inflammatory cytokines (*Tnfα*, *Il-1β* and *Mcp-1*) in macrophages
exposed to palmitate and LPS. In hepatocytes, I3A significantly attenuated TNF-α and fatty acid
induced inflammatory responses in an aryl-hydrocarbon receptor (AhR) dependent manner.
Based on these findings, we hypothesized that I3A could protect against NAFLD progression *in*
*vivo*. We show that supplementation of I3A in drinking water alleviates liver steatosis and
inflammation even when mice are continued on the NAFLD inducing diet, and that these effects
correlate with a decrease in both lipogenesis and β-oxidation in the liver. We also demonstrate
that the anti-inflammatory effects of I3A in macrophages can be mediated by AMP-activated
protein kinase (AMPK).

**RESULTS**

**Oral administration of I3A alleviates diet induced hepatic steatosis and inflammation**

We utilized a mouse model of diet-induced fatty liver disease [15] to investigate the effect of I3A
administration through drinking water (Fig. S1A) on liver steatosis and inflammation. Similar to
our previous finding in HFD fed mice [14], the levels of I3A were significantly reduced in fecal
material (50% and 64% decrease at week 8 and 16, respectively) and the liver (70% decrease at
week 16) of Western diet (WD) fed mice compared to control mice (CN) fed a low-fat diet. Mice
that were given WD and sugar water (SW) containing I3A showed increased I3A levels in fecal
material, the liver and serum compared to mice given the same WD and SW without I3A (Fig.
S1A and 1B, Fig. S1C). Next, we sought to determine if the I3A administration impacted WD
induced features of NAFLD. Compared to CN, WD fed mice had elevated serum alanine
aminotransferase (ALT) levels, indicating liver injury. Treatment of WD fed mice with the high
dose of I3A (WD-100) significantly reduced serum ALT levels two weeks after beginning the
I3A treatment (week 10 time point, Fig. 1C) and beyond. Additionally, I3A treatment reduced
liver TG in a dose-dependent manner at week 16 (Fig. 1D). Scoring of H&E-stained liver
sections indicated that the I3A treatment improved liver steatosis, hepatocyte ballooning, and
lobular inflammation in a dose-dependent manner (Fig. 1E and F). Treatment with I3A didn't
alter weight increase nor food intake compared to WD fed mice (Fig. S1D).

We also investigated whether I3A treatment modulated the levels of inflammatory cytokines in
WD-fed mice. All pro-inflammatory cytokines in the panel (e.g., *Tnfα*, *Il-6, Mcp-1*) were
significantly elevated in livers of WD mice compared to CN (Fig. 2A and Fig. S2A). Treatment
with I3A reduced the expression of these cytokines in a dose-dependent manner. Interestingly,
WD upregulated IL-10, considered an anti-inflammatory cytokine, and I3A treatment
downregulated this cytokine (Fig. S2A). Taken together, these results demonstrate that I3A,
provided via drinking water, attenuates diet-induced hepatic steatosis, cellular injury, and
inflammation in mice.

**I3A reverses diet induced alterations in liver bile acids and free fatty acids**

Studies in mice and humans showed that NAFLD is associated with alterations in liver bile acids.
Previously, we found that the ratio of chenodeoxycholic acid (CDCA) to cholic acid (CD)
greatly decreased when HepG12 and AML12 cells were treated with I3A, indicating that I3A can
directly alter host cell bile acid metabolism *in vitro* [14]. In the present study, mice in the WD
group showed reduced liver total bile acids and CDCA-derived bile acids compared to CN (Fig.
2B). Treatment with I3A led to a further decrease in the CDCA-derived bile acids, consistent
with our previous observation of I3A's effect on hepatocytes *in vitro*. We also measured the

concentrations of nine major free fatty acids (FFAs, Table S2). In the liver, WD significantly
increased the total concentration of these FFAs by 2.7-fold compared to CN (Fig. S2B).
Treatment with I3A had no significant effect, although the total FFA concentration trended lower
in the WD-100 group (36% reduction, $p = 0.11$, Fig. S2B). The total FFA concentrations in
serum were similar across all four groups (Fig. S2B).

**I3A administration does not significantly alter the fecal microbial community as well as**
**fecal metabolome**

Next, we investigated whether the hepatoprotective effects of I3A were due to a modification of
the gut microbiome. The fecal microbial communities at week 8 of WD (prior to I3A treatment)
and week 16 (at termination) were analyzed using 16S rRNA sequencing. Fecal microbiome of
WD fed mice showed reduced α-diversity compared to CN, which was not further altered upon
I3A treatment (Fig. S3A). Similarity analysis using the Bray-Curtis dissimilarity metric showed
that the microbiome compositions of CN and WD-fed mice were significantly different (Fig.
S3B). The fecal microbiome compositions of WD-fed mice did not change significantly upon
I3A treatment (Fig. S3B). Analysis of fecal microbiota taxonomic profiles at the phylum and
genus level (Fig. S3C and S3D) showed major shifts in bacterial abundance of fecal microbiota
from WD-fed mice relative to CN. At week 16, the relative abundance of phylum
*Verrucomicrobia* increased, whereas *Bacteroidetes* decreased. LEfSe detected 18 taxa showing
significant differences in relative abundance between the WD group and CN, including an
expansion of the genus *Akkermansia* and reduction of family *Muribaculaceae*. In contrast, only 3
genera showed significant shifts in WD fed mice upon I3A treatment (Fig. S3E).

We also investigated whether administration of I3A altered the metabolic profile of the gut
microbiota. Principal component analysis (PCA) of untargeted LC-MS data showed that the fecal
metabolome of WD fed mice was significantly different from CN, whereas the metabolomes of
WD-100 and WD groups largely overlapped (Fig. S4A). Although permutational multivariate
analysis of variance (PERMANOVA) suggested a modest difference in the variance of
metabolite levels between WD-50 and WD groups, Hotelling's $T^2$ test found that the means
(centroids) were not significantly different. Clustering analysis using the k-means showed that
fecal metabolites could be assigned into 4 groups (Fig. S4B). We did not observe any clusters
that showed a reversal of WD induced changes in fecal metabolome by I3A treatment. Taken
together, these results indicate that treatment with I3A had minimal effects on the fecal
microbiota composition and metabolome of WD fed mice.

**High-dose I3A administration partially reverses diet induced metabolome alterations in the**
**liver**

We next investigated the impact of I3A treatment on the liver metabolome using untargeted LC-
MS. Projections of all annotated features onto latent variables from PLS-DA showed that the
liver metabolome of WD-fed mice was significantly different from CN (Fig. 3A). The low dose
I3A (WD-50) group completely overlapped with the WD group, whereas the high dose (WD-
100) group was significantly different from both WD and CN groups (Fig. 3A). To further
determine if the high dose of I3A was able to reverse the diet-induced alterations in liver
metabolome, we performed a clustering analysis on all 30 annotated features that were
significantly (FDR<0.1) different in WD mice compared to CN. This analysis placed the
significant features into 4 groups. Group 1 comprised metabolites that were further increased or
depleted by the high dose of I3A, whereas groups 3 and 4 comprised metabolites having WD-

induced changes that were reversed by I3A treatment (except for vigabatrin) (Fig. 3B).
Enrichment analysis based on KEGG pathways (Fig. 3C) showed that nicotinate and
nicotinamide metabolism and steroid hormone biosynthesis were significantly altered by WD
(CN vs WD), which were reversed by the high dose I3A treatment (WD vs WD-100).

Because nicotinamide and nicotinate are products of tryptophan metabolism, we manually
integrated the peak areas for all features we identified as tryptophan metabolites (Fig. 3D). This
added formylkynurenine, kynurenic acid and xanthurenic acid to the list of annotated metabolites
comprising the kynurenine branch of tryptophan metabolism. The high dose of I3A restored
WD-induced depletion of metabolites in the upper kynurenine branch, and ameliorated WD-
induced increases in niacin and nicotinamide in the lower branch. These effects were dose-
dependent, as they were not present in the WD-50 group.

Acyl-carnitines comprise another class of metabolites having abundances that were returned to
CN levels in WD-fed mice by the high dose of I3A (Fig. 3E). Compared to CN, mice in the WD
group showed elevated levels of long-chain 3-hydroxylacyl-carnitines. These acyl-carnitines are
closely linked to 3-hydroxylacyl-CoAs, which are intermediates of beta-oxidation, and their
accumulations suggest incomplete beta-oxidation [16,17]. Treatment with the high dose of I3A
significantly reversed these accumulations. As was the case for tryptophan metabolites, the effect
of I3A on acyl-carnitines was dose-dependent.

**I3A administration partially reverses diet induced proteome alterations in the liver**

We next investigated if I3A also changed the expression levels of liver proteins altered by the
WD. Latent variable projections from PLS-DA on liver proteomics data (Fig. 4A) showed that
the WD group had a protein abundance profile significantly different from CN ($p < 10^{-6}$,
Hotelling's $T^2$). The effect of I3A was dose-dependent; whereas the WD-50 group was not
significantly different from the WD group, the WD-100 group was significantly different from
the WD group (p < 0.001). The PLS-DA projections for WD-100 were closer to CN than WD,
but still significantly different from CN. Principal component analysis (PCA) of the proteomics
data showed similar trends (Fig. S5).

Clustering and silhouette analysis on significant (differentially abundant) proteins with VIP
scores > 1.2 in the PLS-DA projections identified three clusters of proteins based on their
abundance profiles. Proteins in cluster 1 had lower abundance in the WD group compared to CN.
This trend was reversed in the WD-100 group (Fig. 4B). Proteins in cluster 2 had lower
abundance in the WD group compared to CN, but this trend was not reversed in the WD-100
group. Proteins in cluster 3 had higher abundance in the WD group compared to CN. These
proteins had similar abundance in the WD-100 group and CN. The protein abundance profiles
were similar between WD-50 and WD groups in all three clusters. These results suggest that the
I3A treatment partially reversed the diet-induced alterations in the liver proteome in a dose-
dependent manner, similar to the trend observed for the liver metabolome.

We performed an enrichment analysis to identify liver biochemical pathways significantly
altered by diet and I3A treatment. In total, 454 quantified proteins were associated with 276
KEGG pathways. Of these, 22 proteins were differentially abundant in WD mice compared to
CN. These proteins were associated with 6 significantly enriched pathways (Fig. 4C). A
comparison of the WD-100 and WD groups identified 52 differentially abundant proteins that
were associated with 5 significantly enriched pathways. One metabolic pathway, biosynthesis of
unsaturated fatty acids, was common to both sets of enriched pathways.

We used targeted proteomics to quantify fatty acid metabolism enzymes identified in the untargeted proteomics data (Fig. 5). Platelet glycoprotein 4 (CD36), a major fatty acid uptake protein, was elevated in the WD group compared to CN. The abundance of CD36 was decreased, but not significantly, in the WD-100 group compared to WD (Fig. 5A). The abundance of fatty acid synthase (Fasn) was not significantly different between the WD group and CN but was decreased in the WD-100 group compared to the WD group (Fig. 5A). Acetyl-CoA carboxylase-2 (Acab), which regulates mitochondrial β-oxidation, was decreased in the WD group compared to CN, but was not significantly different between the WD-100 group compared to WD (Fig. 5B). Enzymes catalyzing mitochondrial β-oxidation, including long-chain acyl-CoA dehydrogenase (Acadl), medium-chain acyl-CoA dehydrogenase (Acadm), short-chain acyl-CoA dehydrogenase (Acads) and hydroxyacyl-CoA dehydrogenase (Hadh) trended higher in the WD group compared to CN, although only Hadh showed a statistically significant difference (Fig. 5B). All four enzymes were significantly decreased in the WD-100 group compared to WD. A similar trend was observed for peroxisomal β-oxidation enzymes acyl-CoA oxidase 1 (Acox1) and peroxisomal 3-ketoacyl-CoA thiolases (Acaa1a and Acaa1b, Fig. 5C). These results suggest that WD feeding led to increased cellular uptake, transport into mitochondria, and β-oxidation of fatty acids, while administration of I3A reduced fatty acid synthesis and β-oxidation. As β-oxidation generates reactive oxygen species (ROS), we also performed a targeted analysis of two antioxidant enzymes detected by the untargeted proteomics experiments, catalase (CatB) and glutathione peroxidase 1 (Gpx1), and administration of I3A reduced the levels of both enzymes compared to WD (Fig. S6A). Previous *in vitro* studies in hepatocytes have shown that AhR activation upregulates hepcidin (Hamp) [18,19], a secreted liver protein that regulates iron absorption. Mice fed the WD showed significantly reduced Hamp abundance in the liver, which was dose-dependently increased by I3A treatment (Fig. S6B). Another iron carrier protein, serotransferrin, showed an opposite trend; its abundance significantly increased in the WD group compared to CN, and dose-dependently decreased by I3A treatment (Fig. S6B).

**I3A suppresses macrophage inflammation in an AMPK but not AhR dependent manner**

Since our previous study found that the metabolic effects of I3A in hepatocytes depend on the AhR, we tested if this was also the case in macrophages. The expression of AhR in RAW 264.7 macrophages is very low compared to murine AML12 hepatocytes (Fig. S7A). Treatment with AhR inhibitor CH223191 did not alter I3A's effect on reducing palmitate and LPS induced *Tnfα* and *Il-β* expression in RAW 264.7 macrophages (Fig. S7B and S7C), indicating that the anti-inflammatory effect of I3A in macrophages likely does not require AhR activity.

Given that AMPK is a well-known signaling mediator in lipid metabolism and inflammation [20-22], and has been shown to play a role in the progression of NAFLD [23,24], we investigated if the levels of AMPK and p-AMPK were altered in the different treatment groups. Both p-AMPK and AMPK were significantly downregulated in livers of the WD group compared to CN (Fig. 6A). Administration of I3A reversed this effect in a dose-dependent manner and increased both AMPK as well as p-AMPK (Fig. 6B).

We tested the role of AMPK in RAW 264.7 macrophages *in vitro*. Palmitate and LPS treatment significantly reduced p-AMPK levels by 50% compared to vehicle control but did not affect total AMPK (Fig. 6C and 6D). Treatment with I3A upregulated p-AMPK to baseline levels without altering total AMPK (Fig. 6C and 6D). Palmitate and LPS also led to an increase in *Tnfα* and *Il-1β* expression (Fig. 6E), while activation of AMPK with AICAR attenuated the palmitate and LPS stimulated increase in *Tnfα* and *Il-1β* expression by 30% and 80%, respectively (Fig. 6E).

These results suggest that upregulation of p-AMPK modulates the expression of *Tnfα* and *Il-1β*
in RAW 264.7 macrophages.

We next investigated if the anti-inflammatory effect of I3A in RAW macrophages that we
previously reported [14] depends on AMPK activation. We used siRNA to knock down prkaa1, the
gene encoding AMPKα1, the main form of AMPKα in murine macrophages [25], and measured its
effect on pro-inflammatory cytokine expression. The knockdown efficiency was ~50% for
mRNA (Fig. S8A), ~ 40-50% for AMPK protein, and ~50 – 60% for p-AMPK (Figs. S8B and
S8C), and the knockdown was stable for up to 96 h at both mRNA and protein levels.  In cells
transfected with the non-targeted control siRNA (Fig. 6F left panel), palmitate and LPS
increased tnfα and il-1β expression by ~10- and 250-fold, respectively (Fig. 6F), and I3A
significantly downregulated this by ~30% and 50% for *Tnfα* and *Il-1β*, respectively, relative to
the DMF control. In contrast, I3A did not significantly modulate *Tnfα* expression in prkaa1
siRNA transfected cells stimulated with palmitate and LPS, and only reduced *Il-1β* expression by
~20 % (Fig. 6F) relative to the DMF control. These results suggest that I3A's anti-inflammatory
effect in RAW 264.7 macrophages depend on AMPK activation.

**DISCUSSION**

Based on our previous finding that I3A modulates inflammatory cytokine expression in
macrophages and fatty acid metabolism in hepatocytes [14], we hypothesized that I3A could
mitigate liver steatosis and other NAFLD features *in vivo*. Using a mouse model of WD-induced
NAFLD [15], we show that oral administration of I3A protects against liver injury in a dose-
dependent manner (Fig. 1C), attenuates liver TG accumulation (Fig. 1D), and reduces steatosis,
hepatocyte ballooning, and lobular inflammation (Figs. 1E and 1F), suggesting that
administration of I3A altered both metabolic and inflammation pathways in the liver. As
expected, WD induced significant additional weight gain. Treatment with I3A had negligible
impact on the weight gain and did not reduce food intake.

Production of pro-inflammatory cytokines such as TNFα, IL-1β, MCP-1 and IL-6 by resident
macrophages (Kupffer cells) signals recruitment and infiltration of monocytes into the liver.
These cytokines also promote the differentiation of monocytes into pro-inflammatory
macrophages, which in turn exacerbates dysregulation of hepatic lipid metabolism [26,27].
Consistent with the lobular inflammation score, there was a significant increase in the expression
of inflammatory cytokines in livers of the WD group (Fig. 2A). Remarkably, treatment of WD
fed mice with I3A dose-dependently reduced the expression of every cytokine in the 13-member
panel, including IL-10. The downregulation of IL-10 in I3A treated mice could be a result of
reduced hepatic inflammation (i.e., feedback regulation of IL-10 production). Whether the
decrease in inflammatory cytokine production is due to I3A's anti-inflammatory effects in liver
immune cells [14] or due to modulation of lipid metabolism in hepatocytes warrants further
investigation.

Studies have shown that the alteration of bile acid metabolism could be a biomarker for NAFLD
[28]. Mice in the WD group had a lower total concentration of liver bile acids and higher
proportion of primary bile acids compared to CN (Fig. 2B). The latter trend is consistent with
previous studies comparing bile acid profiles between NASH patients and healthy controls [29].
Compared to our previous *in vitro* results, the effect of I3A treatment on liver bile acids was
modest, possibly due to regulatory mechanisms present *in vivo*. Moreover, CDCA is metabolized
into muricholic acids in mice which are absent in humans and have different affinities for bile

acid receptors compared to CDCA and lithocholic acid, a CDCA-derived secondary bile acid.
Knockout mouse models that lack the bile acid metabolizing enzymes Cyp2c70/Cyp2a12 have
bile acid pools closer to humans [30,31]. Future studies utilizing these models could help elucidate
the physiological impact of I3A on liver bile acids and their role in fatty liver disease.

The observation that the fecal microbial community composition was not significantly altered by
I3A administration suggests that the hepatoprotective effects of I3A are likely due to the
metabolite's direct action on the host, rather than the gut microbiota composition. One caveat to
this interpretation is the shallow resolution of 16S rRNA sequencing. It is possible that I3A
administration altered the microbiota composition at the species level. It is also possible that I3A
could affect the metabolism of the microbiota without altering its composition. A recent study [32]
showed that a diet containing high levels of methionine and cysteine protected mice against
chronic kidney disease by reducing uremic toxins through post-translational modification of gut
bacterial enzymes. To test the possibility that I3A administration altered the intestinal
metabolome, we performed untargeted metabolomics experiments on fecal samples. The results
showed no significant impact of I3A on the fecal metabolomes of WD mice. While further
studies are warranted to fully elucidate the impact of I3A administration on the microbiome and
microbial gene expression in the intestine, these results suggest that the partial reversal of WD-
induced changes in liver metabolome and proteome by I3A treatment is due to the metabolite's
direct action on the liver.

Analysis of enzymes in lipid transport, synthesis, and mobilization showed that WD and I3A
impacted different facets of lipid metabolism. Targeted proteomics detected a 6- fold increase in
fatty acid translocase CD36 in the WD group compared to CN, suggesting that the elevation of
liver FFAs in the WD group (Fig. S2B) could be due to increased uptake. I3A did not affect
CD36 abundance, but significantly decreased Fasn by 86%, the rate limiting enzyme in *de novo*
lipogenesis. This suggests that the significant decrease in liver TG in the WD-100 group could
reflect a downregulation of *de novo* lipid synthesis in the liver.

Liver fatty acid oxidation (FAO) enzymes also showed different responses to WD and I3A.
Acetyl-CoA carboxylase 2 (Acab) is a mitochondrial regulatory enzyme that produces malonyl-
CoA, which inhibits carnitine palmitoyltransferase 1 (CPT1) [33]. Downregulation of Acab in the
WD group compared to CN suggests increased CPT1 activity and hence elevated long-chain FA
(LCFA) transport into the mitochondria [34]. This interpretation is also supported by the
observation that long-chain 3-hydoxy-acyl carnitines were elevated in the WD group compared
to CN, which is consistent with a recent study reporting elevated serum C14 hydroxy-
acylcarnitine in NAFLD patients [35]. Other mitochondrial fatty acid oxidation (FAO) enzymes
also showed a trend towards upregulation in the WD group compared to CN. Upregulation of
FAO by WD was significant in the peroxisomes, the initial site of very long-chain FA oxidation.
The literature is conflicted on FAO in human subjects with NAFLD or NASH. Depending on the
study, enhanced [36-38], unchanged [39,40] or decreased [41] FAO has been reported. The different
results may reflect varying severity of the disease (degree of steatosis or steatosis vs. NASH) and
variations in FAO capacity across individual subjects. One study [42] showed that the expression of
genes in peroxisomal and mitochondria β-oxidation was higher in patients with more severe
steatosis compared to patients with less sever steatosis or healthy control. Administration of I3A
significantly reduced the abundance of both mitochondrial and peroxisomal FAO enzymes.
Increased FAO and concomitant ROS generation can overwhelm cellular antioxidant defenses,
inducing oxidative stress. Although there is a lack of consensus regarding FAO, studies

consistently report elevated markers of oxidative stress in human subjects with steatosis [43-45]. In
this regard, I3A may protect from oxidative stress by reducing FAO. This is consistent with our
observations that I3A administration reversed the WD-induced accumulation of long-chain 3-
hydroxy-acylcarnitines and that the changes in abundance of FAO enzymes were upstream of 3-
hydroxy-acyl-CoAs. Oxidative stress induced enzymes CAT and GPx were upregulated in the
WD group compared to CN and downregulated in the WD-100 group compared to the WD
group. It should be noted that all the biochemical parameters related to lipid metabolism were
assessed at termination of the mouse experiments. It is possible that the downregulation of FAO
enzymes in I3A treated mice reflects a response to reduced lipid accumulation. Further
longitudinal studies are warranted to determine the (likely) dynamic effects of I3A on the liver
lipid metabolism.

Recent studies have demonstrated that tryptophan derived gut bacterial metabolites, including
I3A, are AhR agonists in intestinal epithelial cells [46] hepatocytes [14] and immune cells [28]. In our
previous study [14], we observed that I3A decreased the expression of Fasn and its transcriptional
regulator SREBP-1c in hepatocytes in an AhR dependent manner. In the current study, we again
found that Fasn was downregulated by I3A treatment. Additionally, we observed a dose-
dependent upregulation of hepcidin in mice treated with I3A. Previously, tryptophan derived
metabolites indoxyl sulfate and kynurenine have been shown to induce hepcidin expression in
HepG2 cells in an AhR-dependent manner [18,19]. Hamp is a master regulator of systemic iron
homeostasis that binds to ferroportin and inhibits the absorption of dietary iron and efflux of iron
from liver resident macrophages. Iron overload is common among NAFLD patients [47], and the
excess iron along with inflammation can induce hepcidin in these patients [48-51]. However,
hepcidin gene expression decreases when NAFLD progresses to NASH [51], suggesting that
disease progression is linked to impaired regulation of iron metabolism. Hepcidin knockout in a
mouse model of NAFLD ameliorated steatosis while exacerbating fibrosis [52]. In contrast,
hepcidin overexpression decreased choline-deficient diet induced steatosis, inflammation and
fibrosis in a mouse model of NASH [53]. Whether I3A induction of hepcidin ameliorates iron
overload and how this could contribute to I3A's protective role needs to be investigated.

Another set of WD-induced metabolic alterations that were reversed by I3A treatment and likely
mediated through the AhR are restorations of tryptophan metabolite levels. Studies in
hepatocytes have shown that AhR knockdown down-regulates tryptophan 2,3-dioxygenase,
kynurenine 3-monooxygenase, and kynureninase, enzymes that direct flux into the kynurenine
branch of tryptophan metabolism. Another study reported that kynurenine activation of the AhR
upregulated expression of indoleamine 2,3-dioxygenase 2 in dendritic cells [54]. In the present
study, I3A treatment of WD-fed mice directed flux of tryptophan into kynurenic, anthranilic, and
xanthurenic acids, reducing flux towards niacin and nicotinamide. Given that kynurenine,
kynurenic acid, anthranilic acid, and xanthurenic acid are all AhR agonists, this could have
further enhanced AhR activation by I3A treatment.

In macrophages, however, I3A's effects were independent of AhR activity, as addition of the
AhR inhibitor did not alter the effects (Fig. S7). Studies in HFD-fed mice showed dysregulation
of liver AMPK activity and its association with increased lipid accumulation [20,21,55]. *In vitro*, LPS
activated macrophage showed upregulation of heme oxygenase-1 (HO-1), an AMPK regulated
protein, upon I3A exposure [56]. Based on these reports, we investigated if AMPK was involved in
mediating the response to I3A. We show that both liver AMPK expression and phosphorylation
were reduced in WD-fed mice relative to CN, and that administration of I3A rescued both

AMPK expression and phosphorylation in a dose-dependent manner (Fig. 6). Using an *in vitro* model, we demonstrate a similar AMPK dependence of I3A's anti-inflammatory effect in murine macrophages (Fig. 6). Whether AMPK signaling plays a role in mediating I3A's effects in hepatocytes *in vivo* and whether this is important relative to I3A's activation of the AhR warrants further investigation.

In summary, we have shown that oral administration of a microbiota derived tryptophan metabolite, I3A, in WD-fed mice alleviates diet-induced liver steatosis and inflammation, even when the mice were continued on WD. These hepatoprotective effects occurred without significant alterations in the gut microbiome composition and metabolome profiles, suggesting that I3A acted directly on host cells. Untargeted LC-MS experiments showed activation of several AhR-regulated pathways, suggesting that I3A's effects are partially mediated through this nuclear receptor. Additionally, *in vivo* results show a correlation between AMPK phosphorylation and the efficacy of I3A, while our *in vitro* studies with RAW macrophages show that AMPK mediates I3A's attenuation of pro-inflammatory cytokine expression. While our results do not rule out the involvement of additional signaling pathways in both hepatocytes and macrophages, or interactions between the liver and other tissues (e.g., adipose tissue), the data nevertheless strongly point to I3A directly modulating lipid metabolism and inflammatory cytokine production in the liver.

Recently, several other studies reported on the protective effects of I3A in various mouse models [57,58]. However, there are significant differences between the other studies and our work. The prior studies modeled steatosis using short-term (less than 4 weeks) HFD exposure, whereas we modeled the later stages of NAFLD (i.e., with steatosis and inflammation) utilizing long-term (16 weeks) exposure to WD and sugar water. Moreover, in contrast to the other studies, I3A was administered concurrently with WD and sugar water for the last eight weeks of the study. Despite the differences in the study design, the common conclusion from the above studies and our work is that I3A administration decreases hepatic TG. In addition, our study shows that administration of I3A significantly decreases hepatic inflammation and lipid metabolism with minimal modulation of the gut microbiome composition and metabolic function. Mechanistically, we also show that I3A's anti-inflammatory effects in RAW 264.7 macrophages are mediated through activation of AMPK, whereas its effects in hepatocytes are mediated through the AhR [14]. Thus, demonstrating that I3A elicits protective effects in NAFLD through two signaling pathways in different cell types is a novel and significant finding from our study.

Based on our data, we propose a model describing I3A's effects in the liver (Fig. S9). While additional studies are warranted to investigate the above-mentioned possibilities, the remarkable potency of I3A in alleviating steatosis and inflammation in a therapeutic model clearly demonstrates the potential for developing I3A as an inherently safe treatment option for NAFLD. To this end, preclinical studies should be conducted to evaluate delivery options such as botanical extracts and probiotics, while human subject studies should be performed to determine safety, tolerability, and side effects of I3A. Prospectively, other tryptophan derived microbiota metabolites as well as diets that result in enhanced production of these metabolites in the GI tract could also be investigated for protection against WD-induced hepatic steatosis and inflammation.

**MATERIALS AND METHODS**

**Study design**

The overall objective of the study was to investigate if I3A can alleviate diet induced liver steatosis and inflammation *in vivo*. We induced these NAFLD features in mice by feeding the animals a WD and SW for 8 weeks. The drinking water for these mice was then supplemented with low and high doses of I3A, while continuing the WD for another 8 weeks. The phenotypic changes were assessed by measuring body weight gain, serum ALT, hepatic inflammatory cytokines, and histopathological examination of liver tissue. To investigate potential mechanisms of action, we performed a series of omics analyses on fecal material (metagenomics and metabolomics) and liver tissue (metabolomics and proteomics). To determine cell-type specific signaling mediating I3A effects in macrophages, we used an *in vitro* cell culture model. The histopathological scoring of liver sections was blinded; all other analyses were not blinded. Sample processing and statistical analysis were performed concurrently on treatment and control groups using identical methods. Numbers of replicates and outcomes of statistical tests are indicated in the figure legends.

**Materials**

RAW 264.7 cells were purchased from ATCC (Manassas, MA). Dulbecco's Modified Eagle Medium (DMEM), penicillin/streptomycin, and LPS (from *Salmonella minnesota*) were purchased from Invitrogen (Carlsbad, CA). Fetal bovine serum (FBS) was purchased from Atlanta Biologicals (Flowery Branch, GA). All free fatty acid chemicals, 5-aminoimidazole-4-carboxamide ribonucleotide (AICAR) and the AhR inhibitor CH-223191 were purchased from Millipore Sigma (St. Louis, MO). Indole-3-acetate sodium salt (I3A) was purchased from Cayman chemicals (Ann Arbor, MI). All other chemicals were purchased from VWR (Radnor, PA) or Millipore Sigma unless otherwise specified.

**Animal experiments**

Male B6 129SF1/J mice at 6 weeks of age were obtained from Jackson Laboratories (Bar Harbor, ME). Mice were acclimatized to the animal facility for one week. At the start of the experiment, mice were randomly divided into four groups (n=10 for each group). Three of the four groups were fed *ad libitum* a Western diet (WD) with 40% Kcal from fat and containing 0.15% cholesterol (D12079B, Research Diets) and a sugar water (SW) solution (23.1g/L d-fructose + 18.9g/L d-glucose) as previously described (DIAMOND model [15]). After 8 weeks, the three groups of WD-fed mice were randomly selected for treatment with vehicle (WD group) or low (WD-50 group) or high dose (WD-100 group) of I3A. The WD group was fed WD and drank sugar water. The WD-50 and WD-100 groups were fed WD and drank sugar water containing, respectively, 50 or 100 mg I3A per kg body weight. The treatments were continued for another 8 weeks. Water bottles containing SW and I3A or only SW were changed every other day. A fourth group of mice was fed a low-fat control diet (D12450B, Research Diets) and normal water for 16 weeks (CN group) (Fig. S1A). Mice belonging to the same treatment group were housed together (5 mice/cage). Mice were maintained on 12:12-h light-dark cycles. All procedures were performed in accordance with Texas A&M University Health Sciences Center Institutional Animal Care and Use Committee guidelines under an approved animal use protocol (AUP #2017-0145).

**Sample collection**

Fecal pellets were collected every other week prior to I3A treatment, right before I3A treatment, every week during I3A treatment, and on the last day of the experiments (Fig. S1B). All fecal pellets were flash frozen in liquid nitrogen after collection. Blood samples from the submandibular vein were taken right prior to I3A treatment, every other week during I3A

treatment and on the last day of the experiments. The blood samples were centrifuged at 4,000g for 15 min at 4°C to obtain serum. At the end of the experiment, the mice were sacrificed by euthanasia. The liver was quickly excised and rinsed with 10x volume of ice-cold PBS. The right medium lobe of the liver was fixed in 10% neutral formalin for histological analysis. A small piece from right lateral lobe was stored in RNAlater (Sigma Aldrich, St. Louis, MO) using the manufacturer's protocol. The remaining liver tissue samples were flash frozen in liquid nitrogen, homogenized in HPLC-grade water, lyophilized to a dry powder, and stored at -80°C until further processing.

**Histological analysis**

Formalin fixed liver were embedded in paraffin, sectioned (5µm) and stained with hematoxylin and eosin (H&E) through VWR histological services (Radnor, PA). Liver histology sections were evaluated by an expert pathologist at Texas A&M University who was blinded to the treatment conditions. Histology was assessed using the NASH CRN [59] and fatty liver inhibition of progression (FLIP) consortia criteria [60].

**Serum ALT and hepatic TG measurement**

Alanine aminotransferase (ALT) was measured in serum samples using a commercial ELISA kit (Cayman Chemical Company). Liver triglycerides (TG) were quantified using a commercial colorimetric assay kit (Cayman Chemical Company). Briefly, lyophilized liver samples were weighted and lysed in diluted NP-40 buffer. After centrifugation at 10,000g and 4°C for 15 min, the supernatant was stored on ice for quantification of TG. A small amount of lyophilized liver was used for DNA isolation using a DNA miniprep kit (Zymo Research, Irvine, CA). The TG concentrations (mg/dl) were normalized to tissue DNA contents (µg).

**Fecal microbiome analysis**

Microbial DNA was extracted from homogenized fecal material using the Power soil DNA extraction kit (Qiagen, Carlsbad, CA). The V4 region of 16S rRNA was sequenced on a MiSeq Illumina platform [61] at the Microbial Analysis, Resources, and Services (MARS) core facility (University of Connecticut). Illumina sequence reads were quality filtered, denoised, joined, chimera filtered, aligned, and classified using mothur (v 1.40.4) following the Miseq SOP pipeline. The SILVA database [62] was used for alignment and classification of the operational taxonomic units (OTU) at 97% similarity. The taxonomic dissimilarities between different treatment groups were calculated using the Bray-Curtis dissimilarity metric and visualized on non-metric multidimensional scaling (NMDS) plots. Analysis of similarities (ANOSIM) was used for statistical comparison of microbiome compositions between treatment groups. The differential abundances of OTUs between two different groups were determined using Linear discriminant analysis Effect Size (LEfSe) [63]. Fisher's index was calculated to estimate the alpha diversity.

**Fecal and serum metabolite extraction**

Metabolites were extracted from fecal samples as described previously [64]. Briefly, fecal material was weighed, homogenized, and extracted twice with chloroform/methanol/water. The aqueous phase from the two extractions were combined, lyophilized, and stored at -80°C. Samples were reconstituted in 100 µl methanol/water (1:1, v/v) prior to LC-MS analysis. Serum metabolites were extracted with ice-cold methanol (1:4 serum:methanol). Samples were centrifuged twice at 15,000 x $g$ for 5 min at 4°C. The supernatant was stored at -80°C until LC-MS analysis.

**Liver metabolite and protein extraction**

Lyophilized liver samples were weighted and homogenized using soft tissue homogenization beads (Omni International) on a bead beater (VWR) with 1 ml ice- cold methanol/ water (91:9, v/v). The samples were homogenized for one min, incubated on ice for 5 min, and centrifuged at 12,000 x $g$ for 10 min at 4°C. The supernatant was transferred into a new sample tube through a 70 μm cell strainer. Ice-cold chloroform was added into the tube to obtain a final solvent ratio of 47.6/47.6/4.8 % methanol/chloroform/water [65]. After vigorous mixing, the samples were frozen in liquid nitrogen and thawed at room temperature. The freeze-thaw cycle was carried out three times. Samples were centrifuged at 15,000 x $g$ for 5 min at 4°C. The supernatant and pellet were each transferred into a new sample tube for metabolite and protein analysis, respectively.

For metabolite analysis, 1 ml of HPLC water was added to the supernatant and centrifuged at 10,000 x $g$ for 5 min at 4°C to obtain phase separation. The upper and lower phases were collected separately and filtered through 0.2 μm filters into new sample tubes. The filtered samples were dried to pellets with a lyophilizer and stored at -80°C until further analysis. The upper and lower phases were reconstituted in 100 μl methanol/water (1:1, v/v) and 200 μl methanol, respectively, prior to LC-MS analysis. For protein analysis, the pellet was solubilized in 650 μl extraction buffer (0.5% SDS, 1% v/v β-mercaptoethanol and 1M Tris-HCl, pH=7.6) and 650 μl TRIzol reagent (Thermo Fisher Scientific, Waltham, MA) and incubated at 37°C for one hour. The sample was centrifuged at 14,000 x $g$ for 15 min at 4°C to obtain phase separation. One ml of ice-cold acetone was mixed into the bottom protein layer. Following an overnight incubation at -20°C, the samples were centrifuged at 14,000 x $g$ for 15 min at 4°C. The supernatant was discarded, and the pellet was washed 3× with 1 ml ethanol. The protein pellet was then lyophilized and stored at -80°C until further analysis.

**Untargeted metabolomics**

Untargeted LC-MS experiments were performed on a Q Exactive Plus orbitrap mass spectrometer (Thermo Fisher Scientific, Waltham, MA) coupled to a binary pump HPLC system (UltiMate 3000, Thermo Fisher Scientific, Waltham, MA) at the Integrated Metabolomics Analysis Core (IMAC) facility of Texas A&M university as previously described [64]. Chromatographic separation was achieved on a reverse phase (RP) column (Synergi Fusion 4μm, 150 mm x 2 mm, Phenomenex, Torrance, CA) using a gradient method (Table S3). Sample acquisition was performed Xcalibur (Thermo Fisher Scientific, Waltham, MA). Raw data files were processed in Compound Discoverer (version 3.0, Thermo Fisher Scientific, Waltham, MA). Metabolite identification was performed by searching the features against mzCloud and ChemSpider. For comparing the levels of metabolites between samples, the area under the curve (AUC) determined in Compound Discoverer for each feature was normalized to the sum of the AUCs for all features. If there're multiple features were annotated as the same metabolite, the feature with highest confidence score and highest AUC was selected to represent the metabolite.

**Liver bile acid analysis**

Lyophilized liver samples were homogenized in 10 mM phosphate buffer at pH 6 (28 mg tissue per ml buffer) on a bead beater (VWR, Radnor, PA) for 1 min. Samples were then centrifuged at 15,000 x $g$ for 5 min at 4°C. The supernatant (200 μl) was mixed vigorously with 100 ng of d4-glycohenodeoxycholic acid internal standard, 20 μl of saturated ammonium sulfate and 800μl of acetonitrile, and then centrifuged at 15,000 x $g$ for 5 min at 4°C. The supernatant was transferred into a new sample tube and dried with a vacufuge (Eppendorf, Hauppauge, NY).  Pellets were

resuspended in 100 µl methanol/water (1:1, v/v), vortexed for 30 sec, sonicated for 1 min, and
then centrifuged at 15,000 g and 4°C for 1 min. The supernatant was transferred into a new
sample tube and analyzed by LC-MS.

Targeted LC-MS experiments were performed on a TSQ Altis triple quadrupole mass
spectrometer (Thermo Fisher Scientific, Waltham, MA) coupled to a binary pump UHPLC
(Vanquish, Thermo Scientific) at the TAMU IMAC core facility. Chromatographic separation
was achieved on a Kinetex 2.6 µm, 100 mm x 2.1 mm polar C18 column (Phenomenex,
Torrance, CA) using a gradient method (Table S4). Scan parameters for target ions are listed in
Table S1. Sample acquisition and data analysis were performed with Trace Finder 4.1 (Thermo
Fisher Scientific, Waltham, MA). The calculated bile acid concentrations were normalized to the
level of spiked internal standard for each sample.

### Quantification of free fatty acids

575  Serum and liver FFAs were analyzed using targeted LC-MS experiments performed on a
576  quadrupole-time of flight instrument (TripleTOF 5600+, AB Sciex, Framingham, MA) as
577  previously described [14] with minor modifications (Tables S2 and S5).

### Untargeted proteomics

Proteins were reduced, alkylated, and digested into peptides using information dependent
acquisition (IDA) on a quadrupole-time of flight instrument (TripleTOF 5600+) as previously
described [66] with minor modifications (Supplemental Information, Table S6). Peptide ions
detected in the IDA experiment were annotated using ProteinPilot (v.5.0, AB Sciex) and further
processed using an inhouse script written in MATLAB to associate each protein identified in the
sample with a unique high-quality peptide (Fig. S9). The relative abundance of a protein was
determined by quantifying the corresponding peptide peak's AUC in Multi-Quant (v2.0, AB) and
normalizing the value by the sample's sum of all peptide AUCs.

### Targeted proteomics

588  A panel of 15 proteins having differential abundance in the untargeted proteomics data were
589  selected for targeted measurements using product ion scans for representative peptides (Table
590  S7). The relative abundance of a target protein was determined by quantifying the corresponding
591  peptide peak's AUCs in MultiQuant and normalizing the value by the sample protein weight.

### Liver cytokine measurements

Lyophilized liver samples were weighted and homogenized with 500 µl lysis buffer (50mM Tris,
150mM NaCl, 1% Triton X-100, 1mM EDTA and 1% protease inhibitor cocktail, PH=7.4) for 1
min on a bead beater (VWR, Radnor, PA). Samples were centrifuged at 20,000 x $g$ for 15min at
4°C. The lipid layer was removed by pipetting and the supernatant was transferred into a new
tube. The centrifugation and lipid removal steps were repeated three times. The supernatant was
transferred into a new tube and the total protein concentration was measured with the BCA
protein assay (Thermo Fisher Scientific, Waltham, MA). A panel of 13 cytokines was quantified
with a bead-based ELISA kit (BioLegend, San Diego, CA) following the manufacturer's
protocol. Cytokine concentrations (pg/ml) were normalized to the corresponding sample total
protein concentration (mg protein/mg tissue).

### RAW 264.7 macrophage culture

RAW 264.7 murine macrophages were cultured in a humidified incubator at 37°C and 5% $CO_2$ using DMEM supplemented with 10% heat inactivated FBS, penicillin (200 U/mL) and streptomycin (200 µg/mL). Cells were passaged every 2-3 days and used within 10 passages after thawing. For the two-hit model experiment, RAW 264.7 cells were seeded into 24-well plates at a density of 2 x $10^5$ cells/ml and then treated with 1 mM I3A, followed by palmitate and LPS as reported before [14]. For the p-AMPK activation experiment, RAW 264.7 cells were treated with 1 mM AICAR for 4 h, followed by addition of 300 µM palmitate complexed with BSA. Following an 18 h incubation, the cells were treated with 10 ng/ml LPS for another 6 hours.

**RNA extraction and qRT-PCR**

Total RNA was extracted from RAW 264.7 cell pellets using the EZNA Total RNA kit (Omega Bio-Tek, Norcross, GA). Purity of isolated RNA was confirmed by A260/280 ratio. qRT-PCR analysis was carried out using the qScript One-Step PCR kit (Quanta Biosciences, Gaithersburg, MD) on a LightCycler 96 System (Roche, Indianapolis, IN). Fold-change values were calculated using the $2^{-\Delta\Delta Ct}$ method, with β-actin as the housekeeping gene. The primer sequences are listed in Table S8.

**AMPK Western blot analysis**

Cell pellets or lyophilized liver samples were lysed with modified RIPA buffer (50 mM Tris-HCl, PH 7.4, 1% Triton X-100, 150 mM NaCl, 1 mM EDTA, and 0.5% Sodium deoxycholate) supplemented with a protease inhibitor cocktail (Sigma), 10 mM NaF, and 1 mM $Na_3VO_4$. The protein concentration was determined using the BCA protein assay kit (Pierce). Protein samples were denatured with SDS and ~10 µg of protein was separated on a 10% SDS-PAGE gel. Proteins were transferred to a PVDF membrane (Thermo Scientific, Waltham, MA) by wet-transfer electrophoresis. Non-fat milk (5%) in TBST solution was used to block non-specific binding. The blots were probed with appropriate primary antibodies (p-AMPK: 2535, total AMPK: 2603, β-actin: 12620, Cell Signaling Technology) and secondary antibody (Anti-rabbit horseradish peroxidase-conjugated secondary antibody, 7074, Cell Signaling Technology). Proteins bound by both primary and secondary antibodies were visualized by chemiluminescence after incubating the blot with Clarity Max Western ECL Blotting Substrate (Bio-Rad, Hercules, CA). Blot images were acquired on a ChemiDoc gel imaging system (Bio-Rad, Hercules, CA). Proteins were quantified by normalizing the intensity of the protein band of interest to the intensity of the β-actin band in the same lane using the Image Lab software (Bio-Rad, Hercules, CA).

**Small interfering RNA transfection**

Raw 264.7 cells were seeded into 6-well or 24-well plates at ~ 30% confluence and cultured for 24 h prior to transfection. Cells were transfected with ON-TARGETplus mouse prkaa1 siRNA (Dharmacon, Lafayette, CO) or ON-TARGETplus non-targeting pool (negative control, Dharmacon, Lafayette, CO) using the GenMute siRNA transfection reagent (SignaGen Laboratories, Rockville, MD) according to manufacturer's instruction. After 24 h, the medium was replaced with siRNA-free growth medium and incubated for an additional 24 to 72 h. The transfection efficiency was determined by monitoring the AMPK mRNA and protein levels using qRT-PCR and Western blot, respectively.

**Statistical analysis**

Determination of significant difference in the level of a metabolite between two experimental groups used the Student's t-test with FDR correction. KEGG pathway enrichment analysis of metabolomics data was performed using MetaboAnalyst 5.0 [67]. Principal component analysis (PCA) and PLS-DA of the metabolomics and proteomics data were conducted using the mixOmics R package (v6.10.9). Ellipses drawn to represent 95% confidence regions assumed Gaussian distribution of latent variables (for PLS-DA) or scores (for PCA). Significance of separation between treatment groups was determined by calculating a standardized Euclidean distance matrix on the coordinates of latent variables or scores and performing a pairwise permutational multivariate analysis of variance (PERMANOVA) with 999 permutations on the distance matrix [68] and Hotelling's T2 tests. A p-value of 0.05 was set as the significance threshold for all statistical comparisons. Heatmaps for liver metabolome (only significantly changed features), fecal metabolome and liver proteome were generated with auto-scaled data. The features were clustered using the k-means method with Pearson correlation as the similarity metric. For protein analysis, a Variable Importance in Projection (VIP) score was calculated for each protein based on the PLS-DA result, and proteins with a VIP score < 1.2 were excluded from the clustering analysis to avoid overfitting. KEGG pathway enrichment analysis of proteomics data was performed using clusterProfiler R package (3.14.3).

**List of Supplementary Materials**

Materials and Methods

Tables S1 to S8

Figures S1 to S10

**References and Notes**

Loomba, R. & Sanyal, A. J. The global NAFLD epidemic. *Nat Rev Gastroenterol Hepatol* **10**, 686-690 (2013). https://doi.org:10.1038/nrgastro.2013.171

Day, C. P. Non-alcoholic fatty liver disease: a massive problem. *Clinical medicine* **11**, 176-178 (2011).

Tiniakos, D. G., Vos, M. B. & Brunt, E. M. Nonalcoholic fatty liver disease: pathology and pathogenesis. *Annual review of pathology* **5**, 145-171 (2010). https://doi.org:10.1146/annurev-pathol-121808-102132

Calzadilla Bertot, L. & Adams, L. A. The Natural Course of Non-Alcoholic Fatty Liver Disease. *International journal of molecular sciences* **17** (2016). https://doi.org:10.3390/ijms17050774

Brandl, K. & Schnabl, B. Intestinal microbiota and nonalcoholic steatohepatitis. *Curr Opin Gastroenterol* **33**, 128-133 (2017). https://doi.org:10.1097/MOG.0000000000000349

Boursier, J. *et al.* The Severity of Nonalcoholic Fatty Liver Disease Is Associated With Gut Dysbiosis and Shift in the Metabolic Function of the Gut Microbiota. *Hepatology* **63**, 764-775 (2016). https://doi.org:10.1002/hep.28356

Loomba, R. *et al.* Gut Microbiome-Based Metagenomic Signature for Non-invasive Detection of Advanced Fibrosis in Human Nonalcoholic Fatty Liver Disease. *Cell Metab* **25**, 1054-1062 e1055 (2017). https://doi.org:10.1016/j.cmet.2017.04.001

Aron-Wisnewsky, J. *et al.* Gut microbiota and human NAFLD: disentangling microbial signatures from metabolic disorders. *Nat Rev Gastroenterol Hepatol* **17**, 279-297 (2020). https://doi.org:10.1038/s41575-020-0269-9

9     Hoyles, L. *et al.* Molecular phenomics and metagenomics of hepatic steatosis in non-
diabetic obese women. *Nat Med* **24**, 1070-1080 (2018). https://doi.org:10.1038/s41591-
018-0061-3
10    Le Roy, T. *et al.* Intestinal microbiota determines development of non-alcoholic fatty
liver disease in mice. *Gut* **62**, 1787-1794 (2013). https://doi.org:10.1136/gutjnl-2012-
303816
11    Saltzman, E. T., Palacios, T., Thomsen, M. & Vitetta, L. Intestinal Microbiome Shifts,
Dysbiosis, Inflammation, and Non-alcoholic Fatty Liver Disease. *Front Microbiol* **9**, 61
(2018). https://doi.org:10.3389/fmicb.2018.00061
12    Bibbo, S. *et al.* Gut Microbiota as a Driver of Inflammation in Nonalcoholic Fatty Liver
Disease. *Mediat Inflamm* (2018). https://doi.org:Artn 9321643
10.1155/2018/9321643
13    Ding, Y. *et al.* Interactions between gut microbiota and non-alcoholic liver disease: The
role of microbiota-derived metabolites. *Pharmacol Res* **141**, 521-529 (2019).
https://doi.org:10.1016/j.phrs.2019.01.029
14    Krishnan, S. *et al.* Gut Microbiota-Derived Tryptophan Metabolites Modulate
Inflammatory Response in Hepatocytes and Macrophages. *Cell reports* **23**, 1099-1111
(2018). https://doi.org:10.1016/j.celrep.2018.03.109
15    Asgharpour, A. *et al.* A diet-induced animal model of non-alcoholic fatty liver disease
and hepatocellular cancer. *Journal of hepatology* **65**, 579-588 (2016).
https://doi.org:10.1016/j.jhep.2016.05.005
16    Su, X., Han, X., Mancuso, D. J., Abendschein, D. R. & Gross, R. W. Accumulation of
long-chain acylcarnitine and 3-hydroxy acylcarnitine molecular species in diabetic
myocardium: identification of alterations in mitochondrial fatty acid processing in
diabetic myocardium by shotgun lipidomics. *Biochemistry* **44**, 5234-5245 (2005).
https://doi.org:10.1021/bi047773a
17    Natarajan, S. K. & Ibdah, J. A. Role of 3-Hydroxy Fatty Acid-Induced Hepatic
Lipotoxicity in Acute Fatty Liver of Pregnancy. *International journal of molecular*
*sciences* **19** (2018). https://doi.org:10.3390/ijms19010322
18    Hamano, H. *et al.* The uremic toxin indoxyl sulfate interferes with iron metabolism by
regulating hepcidin in chronic kidney disease. *Nephrol Dial Transplant* **33**, 586-597
(2018). https://doi.org:10.1093/ndt/gfx252
19    Eleftheriadis, T., Pissas, G., Antoniadi, G., Liakopoulos, V. & Stefanidis, I. Kynurenine,
by activating aryl hydrocarbon receptor, decreases erythropoietin and increases hepcidin
production in HepG2 cells: A new mechanism for anemia of inflammation. *Exp Hematol*
**44**, 60-67 e61 (2016). https://doi.org:10.1016/j.exphem.2015.08.010
20    Muse, E. D. *et al.* Role of resistin in diet-induced hepatic insulin resistance. *The Journal*
*of clinical investigation* **114**, 232-239 (2004). https://doi.org:10.1172/JCI21270
21    Sag, D., Carling, D., Stout, R. D. & Suttles, J. Adenosine 5'-monophosphate-activated
protein kinase promotes macrophage polarization to an anti-inflammatory functional
phenotype. *Journal of immunology* **181**, 8633-8641 (2008).
22    Day, E. A., Ford, R. J. & Steinberg, G. R. AMPK as a Therapeutic Target for Treating
Metabolic Diseases. *Trends Endocrinol Metab* **28**, 545-560 (2017).
https://doi.org:10.1016/j.tem.2017.05.004

23 Smith, B. K. *et al.* Treatment of nonalcoholic fatty liver disease: role of AMPK.
*American journal of physiology. Endocrinology and metabolism* **311**, E730-E740 (2016).
https://doi.org:10.1152/ajpendo.00225.2016
24 Herzig, S. & Shaw, R. J. AMPK: guardian of metabolism and mitochondrial homeostasis.
*Nat Rev Mol Cell Biol* **19**, 121-135 (2018). https://doi.org:10.1038/nrm.2017.95
25 Yang, Z., Kahn, B. B., Shi, H. & Xue, B. Z. Macrophage alpha1 AMP-activated protein
kinase (alpha1AMPK) antagonizes fatty acid-induced inflammation through SIRT1. *J*
*Biol Chem* **285**, 19051-19059 (2010). https://doi.org:10.1074/jbc.M110.123620
26 Seki, E. & Schwabe, R. F. Hepatic inflammation and fibrosis: functional links and key
pathways. *Hepatology* **61**, 1066-1079 (2015). https://doi.org:10.1002/hep.27332
27 Braunersreuther, V., Viviani, G. L., Mach, F. & Montecucco, F. Role of cytokines and
chemokines in non-alcoholic fatty liver disease. *World J Gastroenterol* **18**, 727-735
(2012). https://doi.org:10.3748/wjg.v18.i8.727
28 De Juan, A. & Segura, E. Modulation of Immune Responses by Nutritional Ligands of
Aryl Hydrocarbon Receptor. *Front Immunol* **12**, 645168 (2021).
https://doi.org:10.3389/fimmu.2021.645168
29 Puri, P. *et al.* The presence and severity of nonalcoholic steatohepatitis is associated with
specific changes in circulating bile acids. *Hepatology* **67**, 534-548 (2018).
https://doi.org:10.1002/hep.29359
30 de Boer, J. F. *et al.* A human-like bile acid pool induced by deletion of hepatic Cyp2c70
modulates effects of FXR activation in mice. *J Lipid Res* **61**, 291-305 (2020).
https://doi.org:10.1194/jlr.RA119000243
31 Ueda, H. *et al.* Sex-, age-, and organ-dependent improvement of bile acid hydrophobicity
by ursodeoxycholic acid treatment: A study using a mouse model with human-like bile
acid composition. *PLoS One* **17**, e0271308 (2022).
https://doi.org:10.1371/journal.pone.0271308
32 Lobel, L., Cao, Y. G., Fenn, K., Glickman, J. N. & Garrett, W. S. Diet posttranslationally
modifies the mouse gut microbial proteome to modulate renal function. *Science* **369**,
1518-1524 (2020). https://doi.org:10.1126/science.abb3763
33 Abu-Elheiga, L. *et al.* The subcellular localization of acetyl-CoA carboxylase 2. *Proc*
*Natl Acad Sci U S A* **97**, 1444-1449 (2000). https://doi.org:10.1073/pnas.97.4.1444
34 Reddy, J. K. & Hashimoto, T. Peroxisomal beta-oxidation and peroxisome proliferator-
activated receptor alpha: an adaptive metabolic system. *Annu Rev Nutr* **21**, 193-230
(2001). https://doi.org:10.1146/annurev.nutr.21.1.193
35 Enooku, K. *et al.* Altered serum acylcarnitine profile is associated with the status of
nonalcoholic fatty liver disease (NAFLD) and NAFLD-related hepatocellular carcinoma.
*Sci Rep* **9**, 10663 (2019). https://doi.org:10.1038/s41598-019-47216-2
36 Sunny, N. E., Parks, E. J., Browning, J. D. & Burgess, S. C. Excessive hepatic
mitochondrial TCA cycle and gluconeogenesis in humans with nonalcoholic fatty liver
disease. *Cell Metab* **14**, 804-810 (2011). https://doi.org:10.1016/j.cmet.2011.11.004
37 Dasarathy, S. *et al.* Elevated hepatic fatty acid oxidation, high plasma fibroblast growth
factor 21, and fasting bile acids in nonalcoholic steatohepatitis. *Eur J Gastroenterol*
*Hepatol* **23**, 382-388 (2011). https://doi.org:10.1097/MEG.0b013e328345c8c7
38 Miele, L. *et al.* Hepatic mitochondrial beta-oxidation in patients with nonalcoholic
steatohepatitis assessed by 13C-octanoate breath test. *Am J Gastroenterol* **98**, 2335-2336
(2003). https://doi.org:10.1111/j.1572-0241.2003.07725.x

39 Petersen, K. F., Befroy, D. E., Dufour, S., Rothman, D. L. & Shulman, G. I. Assessment
of Hepatic Mitochondrial Oxidation and Pyruvate Cycling in NAFLD by (13)C Magnetic
Resonance Spectroscopy. *Cell Metab* **24**, 167-171 (2016).
https://doi.org:10.1016/j.cmet.2016.06.005
40 Kotronen, A. *et al.* Liver fat and lipid oxidation in humans. *Liver Int* **29**, 1439-1446
(2009). https://doi.org:10.1111/j.1478-3231.2009.02076.x
41 Croci, I. *et al.* Whole-body substrate metabolism is associated with disease severity in
patients with non-alcoholic fatty liver disease. *Gut* **62**, 1625-1633 (2013).
https://doi.org:10.1136/gutjnl-2012-302789
42 von Loeffelholz, C. *et al.* Increased lipogenesis in spite of upregulated hepatic 5'AMP-
activated protein kinase in human non-alcoholic fatty liver. *Hepatol Res* **47**, 890-901
(2017). https://doi.org:10.1111/hepr.12825
43 Palmieri, V. O., Grattagliano, I., Portincasa, P. & Palasciano, G. Systemic oxidative
alterations are associated with visceral adiposity and liver steatosis in patients with
metabolic syndrome. *J Nutr* **136**, 3022-3026 (2006).
https://doi.org:10.1093/jn/136.12.3022
44 Del Ben, M. *et al.* Serum Cytokeratin-18 Is Associated with NOX2-Generated Oxidative
Stress in Patients with Nonalcoholic Fatty Liver. *Int J Hepatol* **2014**, 784985 (2014).
https://doi.org:10.1155/2014/784985
45 Del Ben, M. *et al.* NOX2-generated oxidative stress is associated with severity of
ultrasound liver steatosis in patients with non-alcoholic fatty liver disease. *BMC*
*Gastroenterol* **14**, 81 (2014). https://doi.org:10.1186/1471-230X-14-81
46 Jin, U. H. *et al.* Microbiome-derived tryptophan metabolites and their aryl hydrocarbon
receptor-dependent agonist and antagonist activities. *Molecular pharmacology* **85**, 777-
788 (2014). https://doi.org:10.1124/mol.113.091165
47 Datz, C., Muller, E. & Aigner, E. Iron overload and non-alcoholic fatty liver disease.
*Minerva Endocrinol* **42**, 173-183 (2017). https://doi.org:10.23736/S0391-1977.16.02565-
7
48 Vela, D. Low hepcidin in liver fibrosis and cirrhosis; a tale of progressive disorder and a
case for a new biochemical marker. *Mol Med* **24**, 5 (2018).
https://doi.org:10.1186/s10020-018-0008-7
49 Zhou, W. & Qiu, K. The correlation between lncRNA NEAT1 and serum hepcidin in the
peripheral blood of non-alcoholic fatty liver disease patients. *Am J Transl Res* **14**, 2593-
2599 (2022).
50 Senates, E. *et al.* Serum levels of hepcidin in patients with biopsy-proven nonalcoholic
fatty liver disease. *Metab Syndr Relat Disord* **9**, 287-290 (2011).
https://doi.org:10.1089/met.2010.0121
51 Mitsuyoshi, H. *et al.* Analysis of hepatic genes involved in the metabolism of fatty acids
and iron in nonalcoholic fatty liver disease. *Hepatol Res* **39**, 366-373 (2009).
https://doi.org:10.1111/j.1872-034X.2008.00464.x
52 Lu, S., Bennett, R. G., Kharbanda, K. K. & Harrison-Findik, D. D. Lack of hepcidin
expression attenuates steatosis and causes fibrosis in the liver. *World J Hepatol* **8**, 211-
225 (2016). https://doi.org:10.4254/wjh.v8.i4.211
53 Chen, H., Zhao, W., Yan, X., Huang, T. & Yang, A. Overexpression of Hepcidin
Alleviates Steatohepatitis and Fibrosis in a Diet-induced Nonalcoholic Steatohepatitis. *J*
*Clin Transl Hepatol* **10**, 577-588 (2022). https://doi.org:10.14218/JCTH.2021.00289

54  Li, Q., Harden, J. L., Anderson, C. D. & Egilmez, N. K. Tolerogenic Phenotype of IFN-
gamma-Induced IDO+ Dendritic Cells Is Maintained via an Autocrine IDO-
Kynurenine/AhR-IDO Loop. *Journal of immunology* **197**, 962-970 (2016).
https://doi.org:10.4049/jimmunol.1502615
55  Wu, Y., Song, P., Xu, J., Zhang, M. & Zou, M. H. Activation of protein phosphatase 2A
by palmitate inhibits AMP-activated protein kinase. *J Biol Chem* **282**, 9777-9788 (2007).
https://doi.org:10.1074/jbc.M608310200
56  Ji, Y. *et al.* Anti-Inflammatory and Anti-Oxidative Activity of Indole-3-Acetic Acid
Involves Induction of HO-1 and Neutralization of Free Radicals in RAW264.7 Cells.
*International journal of molecular sciences* **21** (2020).
https://doi.org:10.3390/ijms21051579
57  Zhang, C. *et al.* Indole-3-acetic acid improves the hepatic mitochondrial respiration
defects by PGC1a up-regulation. *Cell Signal* **99**, 110442 (2022).
https://doi.org:10.1016/j.cellsig.2022.110442
58  Ji, Y., Gao, Y., Chen, H., Yin, Y. & Zhang, W. Indole-3-Acetic Acid Alleviates
Nonalcoholic Fatty Liver Disease in Mice via Attenuation of Hepatic Lipogenesis, and
Oxidative and Inflammatory Stress. *Nutrients* **11** (2019).
https://doi.org:10.3390/nu11092062
59  Kleiner, D. E. *et al.* Design and validation of a histological scoring system for
nonalcoholic fatty liver disease. *Hepatology* **41**, 1313-1321 (2005).
https://doi.org:10.1002/hep.20701
60  Bedossa, P. & Consortium, F. P. Utility and appropriateness of the fatty liver inhibition
of progression (FLIP) algorithm and steatosis, activity, and fibrosis (SAF) score in the
evaluation of biopsies of nonalcoholic fatty liver disease. *Hepatology* **60**, 565-575 (2014).
https://doi.org:10.1002/hep.27173
61  Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K. & Schloss, P. D.
Development of a dual-index sequencing strategy and curation pipeline for analyzing
amplicon sequence data on the MiSeq Illumina sequencing platform. *Appl Environ*
*Microbiol* **79**, 5112-5120 (2013). https://doi.org:10.1128/AEM.01043-13
62  Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data
processing and web-based tools. *Nucleic Acids Res* **41**, D590-596 (2013).
https://doi.org:10.1093/nar/gks1219
63  Segata, N. *et al.* Metagenomic biomarker discovery and explanation. *Genome Biol* **12**,
R60 (2011). https://doi.org:10.1186/gb-2011-12-6-r60
64  Yang, F. *et al.* Effect of diet and intestinal AhR expression on fecal microbiome and
metabolomic profiles. *Microb Cell Fact* **19**, 219 (2020). https://doi.org:10.1186/s12934-
020-01463-5
65  Dettmer, K. *et al.* Metabolite extraction from adherently growing mammalian cells for
metabolomics studies: optimization of harvesting and extraction protocols. *Anal Bioanal*
*Chem* **399**, 1127-1139 (2011). https://doi.org:10.1007/s00216-010-4425-x
66  Manteiga, S. & Lee, K. Monoethylhexyl Phthalate Elicits an Inflammatory Response in
Adipocytes Characterized by Alterations in Lipid and Cytokine Pathways. *Environ*
*Health Perspect* **125**, 615-622 (2017). https://doi.org:10.1289/EHP464
67  Pang, Z. *et al.* Using MetaboAnalyst 5.0 for LC-HRMS spectra processing, multi-omics
integration and covariate adjustment of global metabolomics data. *Nat Protoc* **17**, 1735-
1761 (2022). https://doi.org:10.1038/s41596-022-00710-w

68      Anderson, M. J. in *Wiley StatsRef: Statistics Reference Online*     1-15.

**Acknowledgments:**

We gratefully acknowledge the use of facilities at the Microbial Analysis, Resources, and Services, (MARS) Center for Open Research Resources and Equipment at the University of Connecticut, Texas A&M High Performance Research Computing, Integrated Metabolomics Analysis Core (IMAC) at Texas A&M University, and Mass Spectrometry Core Lab at Tufts University.

**Funding:**

Ray B. Nesbitt Chair endowment, Texas A&M (AJ)
Karol Family Professorship (KL)

**Author contributions:**
>
> Conceptualization: KL, AJ
> Methodology: YD, KY, FY, EC, CC, MEH, RM, RCA
> Investigation: YD, KY, FY, EC, CC, MEH, RM, KL, AJ
> Visualization: YD, KY
> Funding acquisition: KL, AJ
> Project administration: KL, AJ
> Supervision: RCA, KL, AJ
> Writing – original draft: YD, KY, EC
> Writing – review & editing: YD, KY, KL, AJ

**Competing interests:** Authors declare they have no competing interests

**Data and materials availability:** All data are available in the main text or Supplemental Information.

897
898
**Fig. 1. Oral administration of I3A alleviates diet induce hepatic steatosis and inflammation.**

**(A)** Fecal (at weeks 8 and 16) and **(B)** liver (at week 16) concentrations of I3A in male B6 129SF1/J

mice in control low-fat diet (CN), Western diet (WD), Western diet with low-dose I3A (WD-50),

and Western diet with high-dose I3A (WD-100). **(C)** Serum alanine aminotransferase (ALT) levels

in mice at weeks 10 and 16. **(D)** Liver triglyceride (TG) levels at week 16. Data shown are TG

concentrations (mg/dl) normalized to corresponding tissue DNA contents (µg DNA). **(E)**

905     Representative liver sections stained with hematoxylin-eosin (H&E). **(F)** Histology score for

906     steatosis, hepatocyte ballooning and lobular inflammation. H&E-stained liver sections were

907     evaluated by an expert pathologist using the NASH CRN and fatty liver inhibition of progression

908     (FLIP) consortia criteria. Data shown are mean ± SEM (n = 10 per group). *: $p < 0.05$, **: $p < 0.01$,

909     ***: $p < 0.001$ using Wilcoxon rank sum test.

915
916
917

**Fig. 2. I3A administration reverses WD induced alterations in liver inflammatory cytokines**

**and bile acids. (A)** Inflammatory cytokines in liver tissue at week 16. **(B)** Liver total bile acid

concentration (left panel), and abundance of CDCA branch bile acids relative to total bile acids

pool (right panel). Data shown are mean ± SEM. *: $p<0.05$, **: $p<0.01$, ***: $p<0.001$ using

Wilcoxon rank sum test.

**Fig. 3. I3A administration partially reverses diet-induced metabolome alterations in the**

**liver. (A)** Scatter plots of latent variable projections from PLS-DA of untargeted metabolomics

data features. Comparison of all four experimental groups (left panel), CN vs. WD group (middle

panel), and WD vs. WD-50 and WD-100 groups (right panel). **(B)** Heatmap of significant

metabolite features (FDR<0.05) based on statistical comparisons of treatment groups (CN vs.

WD). (C) KEGG pathway enrichment analysis of the metabolites. Number in the parenthesis

represents the metabolites detected in the pathway. (D) Schematic for tryptophan metabolism (left

panel).  Tryptophan metabolism metabolites fold changes of WD-50, WD-100, CN relative to WD

(right panel). (E) Acyl-Carnitine fold change of WD-50, WD-100, CN relative to WD. p-values

were calculated using Student *t*-test and corrected by FDR.

946
947
**Fig. 4. I3A administration partially reverses diet-induced proteome alterations in the liver.**

**(A)** Scatter plots of latent variable projections from PLS-DA of confidently identified proteins.

Comparison of all four experimental groups (left panel), CN vs WD group (middle panel), and

WD vs. WD-50 and WD-100 groups (right panel). **(B)** Heatmap of significant proteins having

Variable Importance in Projection score >1.2. The proteins were clustered using k-means. **(C)**

Pathway enrichment analysis of significant proteins differentially abundant in CN vs. WD

comparison (upper panel) and WD-100 vs. WD comparison (lower panel). GeneRatio divides the

number of significantly altered proteins that are in the pathway by the total number of significantly

altered proteins. BGRatio divides the number of proteins that are in the pathway by the number of

all detected proteins. The $p$-value was calculated using Fisher's exact test.

**Fig. 5. I3A administration reduces the levels of enzymes in fatty acid transport,** *de novo*

**lipogenesis and β-oxidation. (A)** Abundance of fatty acid translocase (CD36) and fatty acid

synthase. **(B)** Mitochondrial and **(C)** peroxisomal fatty acid oxidation enzymes. Data shown are

mean ± SEM. *: p<0.05, **: p<0.01 using Wilcoxon rank sum test

**Fig. 6. I3A suppresses macrophage inflammation *in vitro* in an AMPK dependent manner**.

**(A)** and **(B)** I3A administration reverses WD-induced reduction in liver p-AMPK and AMPK. **(A)**

Levels of p-AMPK and AMPK in liver tissue at week 16 as determined by Western blot analysis.

**(B)** Ratios of p-AMPK (left panel) and AMPK (right panel) to β-actin. The ratios were determined

based on the p-AMPK and AMPK band intensities quantified using Image Lab (Bio-Rad) and

normalized to the loading control (β-actin). Data shown are mean ± SEM. *: $p<0.05$, **: $p<0.01$

using Wilcoxon rank sum test. **(C)** Expression levels of p-AMPK and total AMPK in RAW 264.7

macrophages pre-treated with either I3A or vehicle (DMF) control followed by stimulation with

palmitate and LPS, determined by Western blot analysis. **(D)** Fold-changes in p-AMPK and total

AMPK. Fold-changes were calculated relative to the DMF and no palmitate and LPS stimulation

condition. The band intensities were quantified and normalized to loading control (β-actin) by

using Image Lab (Bio-Rad). **(E)** Expression levels of *tnfα* and *il-1β* in RAW 264.7 cells treated

with p-AMPK activator AICAR, followed by stimulation with palmitate and LPS. **(F)** Expression

levels of *tnfα* (top row) and *il-1β* (bottom row) in RAW 264.7 cells transduced with non-targeted

control siRNA (left panels) or *Prkaa1* siRNA (middle panels), pre-treated with I3A, and then

stimulated with palmitate and LPS. Data shown are mean ± SEM from three independent cultures

with three biological replicates. *: $p<0.05$, **: $p<0.01$, ***: $p<0.001$ using Student's *t*- test.

**Microbially-derived indole-3-acetate alleviates diet induced steatosis and inflammation in mice**

Yufang Ding[1], Karin Yanagi[2], Fang Yang[1], Evelyn Callaway[1], Clint Cheng[1], Martha E Hensel[3], Rani Menon[1], Robert Alaniz[4], Kyongbum Lee[2*], Arul Jayaraman[1,4*]

[1]Artie McFerrin Department of Chemical Engineering, Texas A&M University, College Station, TX 77843, USA

[2]Department of Chemical and Biological Engineering, Tufts University, Medford, MA 02155, USA

[3]Department of Pathobiology, College of Veterinary Medicine and Biomedical Sciences, Texas A&M University, College Station, TX, USA

[4] Department of Microbial Pathogenesis and Immunology, College of Medicine, Texas Health Science Center, Texas A&M University, Bryan, TX, USA

**Material and methods**

**Free fatty acid analysis**

Free fatty acids (FFAs) in the liver and serum were quantified using product ion scan experiments (TripleTOF 5600+, AB Sciex, Framingham, MA). Chromatographic separation was achieved on a C18 column (Gemini 5 μm C18 110 Å, LC Column 250 x 2 mm, 5μm, Phenomenex) using the solvent gradient described in Table S5. Solvent A was acetonitrile/water (3/2, v/v) containing 10 mM ammonium acetate. Solvent B was acetonitrile/isopropanol (1/1, v/v). The injection volume was 5 μL, and the oven temperature was set to 55℃. The flow rate was held at 1 mL/min. The mass spectrometer was operated in negative electrospray ionization (ESI-) mode with $MS^2$ scan. The experimental parameters are described in Table S2. "Light" experiments with low collision energy were used for quantification. "Heavy" experiments with high collision energy were used to confirm peak identity. The sample AUCs were converted into absolute concentrations using calibration curves from chemical standards. The FFA concentrations were then normalized by the corresponding sample DNA concentration.

**Untargeted proteomics**

Proteins extracted from homogenized liver tissue samples were reduced at 37 °C for 30 min using a 50 mM dithiothreitol solution in 50 mM Tris-HCl containing 8 M urea. The reduced proteins were alkylated by incubating the sample in the dark for 15 min with 45.5 mM iodoacetamide. The reduced and alkylated proteins were digested using trypsin. After incubating the sample overnight at 37°C, formic acid was added to lower the pH to 2 and terminate the digest. The sample solution was centrifuged for 5 min at 14,000×g and desalted using spin columns (Pierce, Thermo Fisher). The desalting procedure followed the manufacturer's instruction except that 0.1% (v/v) formic acid was used instead of 0.1% (v/v) trifluoroacetic acid.

Chromatographic separation was achieved on a reverse phase (RP) column (Ascentis Express C18, 2.7 μm 100Å 15cm x 2.1 mm, Sigma) using a gradient method (Table S6). Solvent A was a 0.1% formic acid solution in water, and solvent B was a 0.1% formic acid solution in acetonitrile. The mobile phase flow rate was held constant at 200 μL/min. The IDA method comprised a TOF MS (survey) scan and (triggered) high-resolution MS/MS (product ion) scans monitoring up to 25 candidate ions per cycle. The dependent scans were triggered whenever the survey scan detected a precursor ion matching the following criteria: mass range of 300-1250 m/z, charge state within +2 to +5, mass tolerance of 50 mDa, ion peak is not an isotope within 6 Da, and ion count exceeds 100 cps. Previously

fragmented target ions were excluded for 15 sec to improve the variety of fragmented ions. Collision energy was set

to 10 eV.

Ions detected in the IDA experiment were annotated using ProteinPilot (v.5.0, AB Sciex) against mouse

proteins in the SwissProt database (Figure S8, Step A). Here, an ion refers to a charged peptide comprising a sequence

of amino acids with or without modifications. Ions annotated as the same peptide were collected into one record (Step

B). For each record, the average chromatographic retention time (RT), RT variation (relative to average), median

peptide confidence score was calculated from the associated ion data. The peptides collected into records were filtered

using an automated procedure implemented as a script in MATLAB (R2019b, MathWorks, Natick, MA) based on the

following criteria (Step C). For each peptide, average RT and accurate mass in the record were compared with the

ions of every other peptide having a different predicted sequence and modification. If a match with another peptide

ion was found within 0.1 m/z and 0.5 min, the peptide was excluded from quantification since the overlap decreases

confidence in quantitation. All comparisons were performed with ions instead of records because the average RT of a

record can be unreliable due to multiple elution of peptides belonging to the same record. The data were analyzed to

ensure that a peptide belongs to only one protein, is detected more than once, i.e., at least two ions are included in one

record, the median confidence score of a record is greater than 95%, and the maximal difference between RTs of

peptide ions within a single record is less than 0.5 min.

Peptides meeting the above criteria were analyzed in MultiQuant (v2.0, AB Sciex) to quantify the areas-

under-the curve (AUCs) of corresponding extracted ion chromatograms (XICs) (Step D). The peptide AUCs were

then normalized by the corresponding sample's sum of all peptide AUCs to control for sample-to-sample variability

in total protein loaded onto the HPLC column (Step E). An additional filter was applied to exclude peptides that did

not have AUCs above blank in at least 80% of the samples (Step F). Peptides with RTs that deviate more than 0.5 min

among the samples were also excluded from further analysis. To select a representative peptide for protein

quantification, peptides having post-translational modification sites were excluded, since the modified peptides can at

most represent a fraction of the protein (Step G). If peptides having the same sequence and different charge states

were identified for a protein, then the representative peptide was selected from this pool. Otherwise, the representative

peptide was selected from all peptides belonging to the protein. In either case, the peptide with the largest average

normalized AUC across all samples was selected to represent the corresponding protein's relative abundance.

1069

1070

1071

1072 Table S1. LC-MS parameters for bile acid analysis
1073

| Bile acid | RT (min) | Polarity | Precursor (m/z) | Product (m/z) | CE (V) | RF Lens (V) |
|---|---|---|---|---|---|---|
| Muricholic acid | 1.39 | Negative | 407.3 | 371.3 | 31.7 | 249 |
| Tauromuricholic acid | 0.9 | Negative | 514.5 | 79.9 | 55 | 185 |
| Tauro-β-muricholic acid | 1.87 | Negative | 514.5 | 80 | 55 | 248 |
| Taurohyodeoxycholic acid | 1.58 | Negative | 498.6 | 79.9 | 55 | 137 |
| Murideoxycholic acid | 2.54 | Negative | 391.5 | 327.3 | 34.3 | 170 |
| Glycocholic acid | 1.76 | Negative | 464.5 | 74 | 36.8 | 111 |
| Taurocholic acid | 1.87 | Negative | 514.5 | 79.9 | 55 | 137 |
| Ursodeoxycholic acid | 2.2 | Negative | 391.4 | 355.2 | 32.4 | 178 |
| Cholic acid | 2.64 | Negative | 407.8 | 290.3 | 37.5 | 131 |
| Hyodeoxycholic acid | 2.54 | Negative | 391.5 | 353.2 | 35.7 | 173 |
| Taurodeoxycholic acid | 3.05 | Negative | 498.4 | 79.9 | 54.6 | 198 |
| Deoxycholic acid | 4.83 | Negative | 391.4 | 327.2 | 34.6 | 146 |
| Chenodeoxycholic acid | 4.6 | Negative | 391.4 | 355.1 | 34.8 | 154 |

1074
1075 Table S2. LC-MS parameters for free fatty acid analysis
1076

| Free fatty acids | RT (min) | [M-H]- (m/z) | TOF MS$^2$ (m/z) | CE (V) | DE |
|---|---|---|---|---|---|
| Light Experiment for quantification | | | | | |
| Lauric acid | 1.94 | 199.17 | 199.17 | -8 | -100 |
| Myristic acid | 3.28 | 227.2 | 227.2 | -15 | -150 |
| Palmitic acid | 6.02 | 255.23 | 255.23 | -15 | -150 |
| Palmitoleic acid | 3.91 | 253.22 | 253.22 | -15 | -150 |
| Stearic acid | 10.81 | 283.26 | 283.26 | -15 | -150 |
| Oleic acid | 7.01 | 281.25 | 281.25 | -15 | -150 |
| Linoleic acid | 4.79 | 279.22 | 279.22 | -17 | -150 |
| Linolenic acid | 3.38 | 277.22 | 277.22 | -15 | -150 |
| Arachidonic acid | 4.57 | 303.23 | 303.23 | -10 | -175 |
| Heavy Experiment for confirmation | | | | | |
| Lauric acid | 1.94 | 199.17 | 199.17 | -21 | -100 |
| Myristic acid | 3.28 | 227.20 | 227.20 | -23 | -150 |
| Palmitic acid | 6.02 | 255.23 | 255.23 | -26 | -150 |
| Palmitoleic acid | 3.91 | 253.22 | 253.22 | -25 | -150 |
| Stearic acid | 10.81 | 283.26 | 283.26 | -25 | -150 |
| Oleic acid | 7.01 | 281.25 | 281.25 | -26 | -150 |
| Linoleic acid | 4.79 | 279.22 | 279.22 | -25 | -150 |
| Linolenic acid | 3.38 | 277.22 | 277.22 | -23 | -150 |
| Arachidonic acid | 4.57 | 303.23 | 303.23 | -20 | -175 |

1077
1078

1079    Table S3. Gradient method for untargeted metabolomics
1080

| Time (min) | % Solvent A | % Solvent B |
|---|---|---|
| 0 | 90 | 10 |
| 5 | 60 | 40 |
| 7 | 5 | 95 |
| 9 | 5 | 95 |
| 9.1 | 90 | 10 |
| 13 | 90 | 10 |

1081
1082    Solvent A: 0.1% (v/v) formic acid solution in water. Solvent B was a 0.1% (v/v) formic acid solution in methanol.
1083    The flow rate was 0.4 mL/min.
1084
1085
1086    Table S4. Gradient method for bile acid analysis
1087

| Time (min) | % Solvent A | % Solvent B |
|---|---|---|
| 0 | 55 | 45 |
| 9 | 30 | 70 |
| 9.5 | 30 | 70 |
| 9.51 | 55 | 45 |
| 12 | 60 | 40 |

1088
1089    Solvent A: 2 mM ammonium acetate in water. Solvent B: methanol:acetonitrile (50:50, V:V). The flow rate was 0.4
1090    mL/min.
1091
1092
1093    Table S5. Gradient method for free fatty acid analysis
1094

| Time (min) | % Solvent A | % Solvent B |
|---|---|---|
| 0 | 90 | 10 |
| 1.7 | 90 | 10 |
| 11.9 | 65 | 35 |
| 14.9 | 0 | 100 |
| 17.4 | 0 | 100 |
| 17.9 | 90 | 10 |
| 20 | 90 | 10 |

1095
1096
1097    Table S6. Gradient method for untargeted proteomics
1098

| Time (min) | % Solvent A | % Solvent B |
|---|---|---|
| 0 | 98 | 2 |
| 15 | 98 | 2 |
| 50 | 55 | 45 |
| 60 | 55 | 45 |
| 62 | 5 | 95 |
| 75 | 5 | 95 |
| 75.5 | 98 | 2 |
| 80 | 98 | 2 |

1099    Table S7. LC-MS parameters for targeted proteomics
1100

| Protein | Selected Peptide | CE (V) | m/z | Charge | RT (min) | Product ion(s) |
|---|---|---|---|---|---|---|

| | | | | | | |
|---|---|---|---|---|---|---|
| CD36 | VAIIESYK | 21.6 | 461.8 | 2 | 32.8 | 752.4 |
| Acox1 | TSNHAIVLAQLITR | 22.6 | 513 | 3 | 37.8 | 701.4 |
| Acaa1a | NTTPDELLSAVLTAVLQDVR | 32.5 | 719.1 | 3 | 53.3 | Sum of 901.5, 1014.6, 1113.7, 800.5, 729.4, 630.4, 517,3 |
| Acaa1b | DTTPDELLSAVLTAVLQDVK | 32.1 | 710.1 | 3 | 54.2 | 873.5 |
| Acads | IGIASQALGIAQASLDCAVK_Carbamidomethyl(C)@17 | 29.8 | 662.7 | 3 | 42.8 | 792.4 |
| Acadm | LLVEHQGVSFLLAEMAMK | 30.3 | 672.7 | 3 | 42.6 | 793.4 |
| Acadl | SPAHGISLFLVENGMK | 25.2 | 567.3 | 3 | 39 | 677.3 |
| Hadh | LLVPYLIEAVR | 30.5 | 643.4 | 2 | 43.1 | 960.6 |
| Acaa2 | TNVSGGAIALGHPLGGSGSR | 27 | 603.3 | 3 | 33.5 | 747.4 |
| Fasn | VLEALLPLK | 23.4 | 498.3 | 2 | 39.7 | 783.5 |
| Acab | VEANLLSSEESLSASELSGEQLQEHGDHSCLSYR_Carbamidomethyl(C)@30 | 45.1 | 941.2 | 4 | 38.1 | 1001.9 |
| Cat | GAGAFGYFEVTHDITR | 25.9 | 580.9 | 3 | 38.1 | 742.7 |
| Gpx1 | YVRPGGGFEPNFTLFEK | 29.4 | 653.3 | 3 | 39.5 | 498.3 |
| Hamp | DTNFPICIFCCK_Carbamidomethyl(C)@7; Carbamidomethyl(C)@10; Carbamidomethyl(C)@11 | 37.6 | 787.8 | 2 | 41.6 | 1097.5 |
| Tf | SAGWVIPIGLLFCK_Carbamidomethyl(C)@13 | 37.3 | 780.9 | 2 | 47 | 947.5 |

1101

1102 CD36: platelet glycoprotein 4; Acox1: peroxisomal acyl-coenzyme A oxidase 1; Acaa1a: peroxisomal 3-ketoacyl-
1103 CoA thiolase A; Acaa1b: peroxisomal 3-ketoacyl-CoA thiolase B; Acads: short-chain acyl-CoA dehydrogenase;
1104 Acadm: medium-chain acyl-CoA dehydrogenase; Acadl: long-chain acyl-CoA dehydrogenase; Hadh: hydroxyacyl-
1105 CoA dehydrogenase; Acaa2: mitochondrial 3-ketoacyl-CoA thiolase; Acab: acetyl-CoA carboxylase 2; Cat:
1106 catalase; Gpx1: Glutathione peroxidase 1; Hamp: hepcidin; Tf: serotransferrin

1107
1108
1109 Table S8. Primer sequences
1110

| Gene | Forward Primer Sequence | Reverse Primer Sequence |
|---|---|---|
| TNFα | TCTCATGCACCACCATCAAGGACT | TGACCACTCTCCCTTTGCAGAACT |
| IL-1β | TCCAGGATGAGGACATGAGCAC | GAACGTCACACACCAGCAGGTTA |
| β-actin | GGCTGTATTCCCCTCCATCG | CCAGTTGGTAACAATGCCATGT |

1111
1112

1113

1114

**Fig. S1. Study design**. **(A)** Three groups of male B6 129SF1/J mice (n=10 for each group) were fed *ad libitum* a

Western diet (WD) and a sugar water (SW) solution while a fourth group was given normal chow diet. After 8 weeks,

the three groups of WD-fed mice were randomly selected for treatment with vehicle (WD group) or low (WD-50

group) or high dose (WD-100 group) of I3A for an additional 8 weeks. The fourth group (CN) was continued on low-

fat diet calorie matched with the WD. **(B)** Sampling scheme. **(C)** Quantification of I3A in serum. **(D)** Body weight

increase and food intake of the four groups. Body weights were normalized to week 8 body weight when I3A

administration was started. Food intakes were measured per cage and average food intake was calculated as per gram

food intake per mouse per day. Data shown are mean ± SEM. *: $p < 0.05$, **: $p < 0.01$, WD group compared to CN

group using Wilcoxon rank sum test.

1124

1125

1126

1127

1128

**A**

**B**

1129

1130 **Fig. S2. I3A reduces liver inflammatory cytokine expression and total FFA concentration. (A)** Inflammatory

1131 cytokines in liver tissue at week 16. **(B)** FFAs in diet, serum and liver samples. Data shown are mean ± SEM. *:

1132 p<0.05, **: p<0.01, ***: p<0.001 using Wilcoxon rank sum test.

1133

1134 **Fig. S3. I3A administration does not significantly alter the fecal microbial community**. **(A)** Alpha diversity of the

1135 fecal microbiome from CN, WD, WD-50, and WD-100 groups. **(B)** Analysis of Similarities (ANOSIM) comparison

1136 for CN vs. WD group and WD vs. WD-50 vs. WD-100 groups. **(C)** Phylum and **(D)** genus level relative abundance

1137 of the fecal microbial community members. **(E)** LEfSe results at the genus level. *: p<0.05, ***: p<0.001 using

1138 Wilcoxon rank sum test.

1139

**Fig. S4. I3A administration does not significantly alter the fecal metabolome. (A)** Score plots show the first two

principal components for all four experimental groups (left panel), CN vs. WD group (middle panel), and WD vs.

WD-50 and WD-100 groups (right panel). Numbers in the parentheses of axis titles show percent of variance explained

by the corresponding principal component. Ellipses circumscribe 95% confidence regions for the experimental groups

assuming Gaussian distribution of component scores. **(B)** Heatmap of fecal microbiome metabolite features detected

in all treatment groups. Rows and columns are features and treatment groups, respectively. The features were clustered

using k-means.

1147

1148

1149 **Fig. S5. Principal Component Analysis of liver proteome after I3A administration. (A)** Score plots of the first

1150 two principal components for all four experimental groups (left panel), CN vs. WD group (middle panel), and WD vs.

1151 WD-50 and WD-100 groups (right panel). Numbers in the parentheses of axis titles show percent of variance explained

1152 by the corresponding principal component. Ellipses represent 95% confidence regions for the experimental groups

1153 assuming Gaussian distribution of component scores.

1154

1155

1156

1157 **Fig. S6. I3A administration reduces the levels of antioxidant enzymes and affects iron metabolism related**

1158 **proteins. (A)** Catalase and glutathione peroxidase-1 abundance for the four experimental groups. **(B)** Hepcidin and

1159 Serotransferrin abundance. Data shown are mean ± SEM. *: p<0.05, **: p<0.01 using Wilcoxon rank sum test.

Fig. S7. I3A's anti-inflammatory effects in macrophages are independent of AhR activation. A. Expression

level of AhR in RAW264.7 and AML12 cells determined by Western blot analysis. B. Raw 264.7 cells were treated

with 1mM I3A (or DMF solvent control) for 4h, then stimulated with 300μM Palmitate for 18h and 10ng/ml LPS for

6h (two-hits model). The AhR inhibitor CH223191 (5μM) or DMSO control were added 10min before I3A

1165    treatment. Total RNA were isolated from the cells and the expression of *tnfα* and *il-1β* were measured with qRT-

1166    PCR. C. *tnfα* and *il-1β* expression were plotted as fold change normalized to the DMF control group. Data presented

1167    as the mean ± SEM. ***: p<0.001 using Student's *t*-test.

1168

1169

**Fig. S8. Prkaa1 siRNA reduces AMPK expression in macrophages. (A)** Levels of prkaa1 mRNA in RAW 264.7

cells transfected with prkaa1 or non-targeted control siRNA for 24 h, followed by incubation for an additional 24, 48,

and 72 h. The expression level of prkaa1 is normalized to that of the housekeeping gene β-actin. **(B)** Western blot

analysis of p-AMPK and total AMPK from cells treated with the different siRNA. A representative blot is shown. **(C)**

Quantified intensities of p-AMPK and total AMPK bands normalized to loading control (β-actin).

1175

Western Diet

Western Diet + I3A

1176

**Fig. S9. Proposed model for the effects of I3A on hepatic lipid metabolism and inflammation.** When mice are

fed with a WD (top panel), TG and FFAs accumulate in the liver due to increased uptake of fatty acids. This also leads

to increased β-oxidation in the mitochondria and peroxisomes. In liver macrophages, the increase in FFAs, possibly

in conjunction with circulating endotoxins (e.g., LPS) stimulate production of inflammatory cytokines. When mice

fed the WD are treated with I3A (bottom panel), both TG and FFAs decrease in the liver. Rather than impact fatty

acid uptake, I3A treatment reduces *de novo* lipogenesis through a downregulation of Fasn, while also reducing both

mitochondrial and peroxisomal β-oxidation. In macrophages, I3A attenuates fatty acid and LPS stimulated production

of inflammatory cytokines through activation of AMPK.

1185

1186

## Proteomic Data processing workflow

Raw LC-MS Data

**Step A**
Peptide* prediction (identification and annotation with protein name) of ion library using biological reference protein sequence set

ProteinPilot™

**Step B**
Generate records by combining information (RT, m/z, charge, etc.) from ions with same prediction

**Step C**
Pick high-quality peptides for quantification based on information in the records

**Step D**
Quantify each peptide's area under curve (AUC) for each sample.

MQ MultiQuant™ Software

**Step E**
Normalize each peptide AUC with sum of all peptide AUCs for each sample

**Step F**
Filter for peptides with AUCs above baseline and consistent RT across samples

**Step G**
Select the peptide to represent protein

*Peptide refers to ion comprising an amino acid sequence with certain modification and charge, unless otherwise noted

1187

1188 **Fig. S10. Untargeted proteomic data analysis workflow.** The individual steps (A - G) are described in the untargeted

1189 proteomics section of Supplemental Information.

1190

1191