Type of article: Research in Context

Seed functional ecology in Brazilian rock outcrop vegetation: an integrative synthesis

Carlos A. Ordóñez-Parra[1,2*], Natália F. Medeiros[2,3], Roberta L.C. Dayrell[4], Soizig Le Stradic[5],

Daniel Negreiros[6], Tatiana Cornelissen[2], Fernando A. O. Silveira[2]

1. Programa de Pós-Graduação em Biologia Vegetal, Departamento de Botânica, Instituto de

Ciências Biológicas, Universidade Federal de Minas Gerais, Belo Horizonte, Brazil.

2. Centro de Síntese Ecológica e Conservação, Departamento de Genética, Ecologia e

Evolução, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Belo

Horizonte, Brazil.

3. Programa de Pós-Graduação em Ecologia, Conservação e Manejo da Vida Silvestre,

Departamento de Genética, Ecologia e Evolução, Instituto de Ciências Biológicas,

Universidade Federal de Minas Gerais, Belo Horizonte, Brazil.

4. Royal Botanic Gardens, Kew, Wakehurst, Ardingly, Haywards Heath, West Sussex,

RH17 6TN, UK.

5. UMR BIOGECO, INRAE and Université de Bordeaux, Pessac, France.

6. Laboratório de Ecologia Evolutiva e Biodiversidade, Departamento de Genética, Ecologia

e Evolução, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Belo

Horizonte, Brazil

* Corresponding author. E-mail: carlos.ordonez.parra@gmail.com

Running Title: Seed ecology in Brazilian rock outcrop vegetation

ABSTRACT

Background and Aims

Rock outcrop vegetation is distributed worldwide and hosts a diverse and unique flora that evolved under harsh environmental conditions. Unfortunately, seed ecology in such ecosystems has received little attention, especially regarding seed traits, germination responses to abiotic factors and the potential role of phylogenetic relatedness on such features Here, we provide the first quantitative and phylogenetically-informed synthesis of the seed functional ecology of Brazilian rock outcrop vegetation, with a particular focus on quartzitic and ironstone *campo rupestre*.

Methods

Using a database of functional trait data, we calculated the phylogenetic signal of seven seed traits for 371 taxa and tested whether they varied among growth forms, geographic distribution, and microhabitats. We also conducted meta-analyses that included 4,252 germination records for 102 taxa to assess the effects of light, temperature, and fire-related cues on the germination of *campo rupestre* species and explored how the aforementioned ecological groups and seed traits modulate germination responses.

Key Results

All traits and germination responses showed a moderate-to-strong phylogenetic signal. *Campo rupestre* species responded positively to light and had maximum germination between 20-25 ºC. The effect of temperatures beyond this range was moderated by growth form, species geographic distribution, and microhabitat. Seeds exposed to heat shocks above 80 °C lost viability, but smoke accelerated germination. We found a moderating effect of seed mass

for in responses to light and heat shocks, with larger, dormant seeds tolerating heat better but

less sensitive to light. Species from xeric habitats evolved phenological strategies to

synchronise germination during periods of increased soil water availability.

Conclusions

Phylogenetic relatedness plays a major role in shaping seed ecology of Brazilian rock outcrop

vegetation. Nevertheless, seed traits and germination responses varied significantly between

growth forms, species geographic distribution and microhabitats, providing support to the

regeneration niche hypothesis and the role of functional traits in shaping germination in these

ecosystems.

Keywords

Campo de altitude; campo rupestre; *canga*; germination requirements; inselberg; regeneration

niche; seed dispersal; seed dormancy; seed mass; seed viability.

INTRODUCTION

Rock outcrops are outstanding geological features where the bedrock protrudes above the

land's surface due to the erosion of softer parts of the landscape (Fitzsimons and Michael

2017). They offer a unique habitat that drastically contrasts with the neighbouring vegetation

(Porembski 2007). Notably, they experience severe surface temperatures and have shallow,

poorly-developed soils with low water retention capacity –a combination of abiotic factors

that has driven the evolution of distinctive traits that allow plants to establish and survive in

such harsh environments (Kluge and Brulfert 2000; Escudero *et al.* 2015; Oliveira *et al.* 2016;

Bondi *et al.* 2023). Most studies, however, have focused on the ecophysiology of adult plants,

and only few comprehensive reviews on germination ecology have been produced to date

(e.g., Wyatt 1997; Biedinger et al. 2000). As a result, the potential role of regeneration traits

(Donohue *et al.* 2010; Larson and Funk 2016) and regeneration niche (Grubb 1977) in

shaping ecological processes in rock outcrops has been largely overlooked.

In Brazil, there are four main vegetation types associated with rock outcrops: *campo*

rupestre, *canga*, *campo de altitude*, and inselbergs (Martinelli 2007). *Campo rupestre* occurs

on quartzite, sandstone, and ironstone outcrops throughout the country, but it is most common

at high elevations in the Espinhaço Range, eastern Brazil (Miola *et al.* 2021). This vegetation

is characterised by sandy and rocky grasslands dominated by grasses, sedges and forbs

(mostly Xyridaceae and Eriocaulaceae) with isolated patches of shrubs (mainly

Melastomataceae, Fabaceae, Asteraceae, Myrtaceae, Vochysiaceae and Clusiaceae) and

desiccation-tolerant and epilithic species establishing directly in the rock (Silveira *et al.*

2016). The *campo rupestre* that develops on ironstone outcrops receives the name *canga* and

is functionally analogous to the quartzitic outcrops (Jacobi *et al.* 2007). The *campo de altitude*

is found on granite and gneissic outcrops within the Atlantic Forest and consists of a mosaic

of shrubs (mostly Asteraceae, Melastomataceae, and Myrtaceae) and small patches of dwarf

forests amidst a matrix of grasses (Safford 1999). In contrast, inselbergs are isolated, dome-

shaped granitic outcrops present across the country. In such geological formations, the mat-

like vegetation –dominated mostly by Bromeliaceae, Cactaceae, Orchidaceae, Velloziaceae

and shrubs from several families– is notably patchy, occurring only in depressions in the rock

surface that accumulate soil (Safford and Martinelli 2000). All these open, grassy-shrubby,

fire-prone and highly heterogeneous vegetation types are centres of species richness,

phylogenetic diversity and endemism (Porembski 2007; Silveira *et al.* 2016; Campos *et al.*

2018).

The germination ecology of Brazilian rock outcrop vegetation has been previously

reviewed (Garcia and Oliveira 2007; Nunes *et al.* 2016; Garcia *et al.* 2020), but these

syntheses were restricted to the *campo rupestre* and focused on the differences in germination

responses in a few emblematic families (e.g., Asteraceae, Bromeliaceae, Cactaceae,

Eriocaulaceae, Melastomataceae, Velloziaceae, and Xyridaceae). Still, based on seminal

studies on seed functional ecology, these syntheses put forward various promising hypotheses

on the role of seed and germination traits in the ecological dynamics of plant communities. A

first hypothesis poses that germination responses are shaped by seed traits, especially seed

mass. For example, Nunes *et al.* (2016) pointed out that *campo rupestre* species with lighter

seeds were more dependent on light for germination than larger ones, as expected under the

trade-off between light requirements and seed mass (Milberg *et al.* 2000; Pearson *et al.* 2003).

Still, such an association between seed traits and germination responses remains to be

formally tested for this ecosystem. The second hypothesis is that contrasting microhabitat

preferences and geographical ranges derive from distinct regeneration traits, as proposed by

the regeneration niche hypothesis (Grubb 1977; Saatkamp *et al.* 2019). For instance, species

from xeric microhabitats in the *campo rupestre* have been shown to germinate at higher

temperatures than those from mesic environments (Oliveira and Garcia 2011), and broadly

distributed species have broader germination niches than those restricted to a single substrate

(Marques *et al.* 2014). Species with a broader distribution range have also been shown to have

a wider germination temperature range (Ranieri *et al.* 2012). Similar studies, however, have

failed to find such a significant association between microhabitat preferences or geographic

distributions in the *campo rupestre* (Oliveira and Garcia 2011; Silveira et al. 2012; Giorni et

al. 2018), suggesting the patterns might be more complicated than previously thought.

Moreover, these studies have been limited to a handful of species within single families,

making it impossible to draw broad conclusions for the entire rock outcrop community. A

new opportunity to address these questions at the community level has emerged owing to a

comprehensive database recently published (Ordóñez-Parra *et al.* 2023). Another limitation of

previous reviews is that they did not explicitly address the potential role of shared

evolutionary history in seed ecology. Local and global scale studies have highlighted that

several seed traits, such as seed mass (Moles *et al.* 2005; Zanetti *et al.* 2020), dormancy

(Willis *et al.* 2014; Dayrell *et al.* 2017) and germination requirements (Arène *et al.* 2017), are

phylogenetically clustered, potentially due to a combination of adaptive evolution and

phylogenetic conservatism (Vandelook *et al.* 2012; Carta, Mattana, *et al.* 2022). As a result, a

proper vegetation-wide assessment of ecological hypotheses regarding seed ecology must

incorporate the relative importance of phylogenetic relatedness in seed functional traits and

germination responses.

Here, we conducted the first integrative synthesis of seed functional ecology of

Brazilian rock outcrop vegetation to address the two previously described long-standing

hypotheses and the potential role of phylogenetic relatedness on the seed ecology of these

ecosystems. First, we assessed whether seed functional traits associated with different

dimensions of the seed ecological spectrum exhibit a significant phylogenetic signal and

evaluated whether they differ between growth forms, species geographic distribution and

microhabitats. Second, we assessed the effect of light, temperature, and fire-related cues on

the germination of plant species in these communities through phylogenetically informed

meta-analyses. Such abiotic factors are established germination cues in these ecosystems and

are the most studied in Brazilian rock outcrop vegetation, allowing us to provide more robust

conclusions about their effects on germination (Figure S1). We then contrasted germination

responses between growth forms, species geographic distribution and microhabitats to test the

regeneration niche hypothesis (Grubb 1977). Finally, we tested if such responses were

moderated by seed mass and dormancy –central traits associated with multiple seed functions

(Saatkamp *et al.* 2019). Because rock outcrops occur in all continents and biomes, and their

associated vegetation is quite similar in structure and function (Barthlott and Porembski

2000), we hope this study provides a solid starting point towards a global synthesis of the

germination ecology of rock outcrop vegetation.

MATERIALS AND METHODS

Data sources

Seed functional traits and germination experiments

We retrieved seed trait and germination records from *Rock n' Seeds* (Ordóñez-Parra *et al.*

2023), a database of seed functional traits and germination experiments of plant species from

Brazilian rock outcrop vegetation (Figure 1). This database contains information on 16

functional traits in 383 taxa and 10,187 germination records for 281 taxa. We focused on

seven functional traits associated with regeneration from seeds and major seed functions

(Grubb 1977; Saatkamp *et al.* 2019): seed dry mass (mg), seed water content (%), percentage

of embryoless seeds (%), percentage of viable seeds (%), seed dispersal syndrome

(anemochory, autochory and zoochory), seed dispersal season (early-dry, late-dry, early-rain

and late-rainy season) and primary dormancy [dormant (D) and non-dormant (ND)]. This

subset of traits included data from 103 studies on 371 species. We also retrieved information

from experiments assessing the effects of light availability, constant and alternate

temperatures, and fire-related cues (i.e., heat shocks and smoke) from 22 studies and 102

species, totalising 4,252 germination records. Beyond seed and germination traits, the

database also provides species-level data on growth form, microhabitat ('xeric' substrates

experiencing pronounced seasonal drought, or otherwise 'mesic'), and geographic distribution

('restricted' to rock outcrop vegetation, and 'widespread' if found in other vegetation types in

addition to rock outcrops) (Ordóñez-Parra *et al.* 2023).

Phylogenetic tree

We reconstructed the phylogenetic tree of the studied species using the package

V.PhyloMaker2 (Jin and Qian 2022). First, species names were checked and updated

following The Leipzig Catalogue of Vascular Plants using the R package *lcvplants* (Freiberg

et al. 2020). We kept taxa identified at the species level, and all subspecies and varieties (18

cases) were upgraded to the species level. A phylogenetic tree for the resulting 370 species

was generated, based on the GBOTB phylogeny (Smith and Brown 2018), which was updated

and standardised following Freiberg *et al*. (2020). Taxa absent from this phylogeny were

bound to their designated relatives using the bind.relative function of *V.PhyloMaker2* based

on different sources (Almeda *et al.* 2016; Rivera *et al.* 2016). Species with no known relatives

in the phylogeny (*Austrocritonia velutina*, *Cavalcantia glomerata*, *Cavalcantia percymosa*

and *Parapiqueria cavalcantei*) were added using the at.node function from the *ape* package

(Paradis and Schliep 2019). All these species belong to the Eupatorieae (Asteraceae), a tribe

where infrageneric relationships remain unresolved (Rivera *et al.* 2016). Therefore, they were

added to the base of the tribe as polytomies. Most infrageneric relationships in the phylogeny

remained unresolved, appearing as polytomies, partially because the relationships within

some genera of the highly diverse families in our database have low support (Alcantara *et al.*

2018; Guimarães *et al.* 2019). Nevertheless, phylogenetic metrics estimated from trees

resolved to the genus level have shown to be highly correlated with those derived from fully

resolved trees, suggesting that these polytomies do not weaken the results (Qian and Jin

2021).

Statistical analyses

All analyses were made using R v. 4.3.2 (R Core Team 2023), and the code prepared will be

provided as Supplementary Material and uploaded to GitHub upon acceptance (see Open

Data).

Variation and phylogenetic signal of seed functional traits

To test for the phylogenetic signal in the quantitative seed traits (i.e., dry mass, water

content, percentage of viable and embryoless seeds), we calculated Pagel's λ (Pagel 1999)

using the phylosig function from the package *phytools* (Revell 2012). λ values range from

zero to one, with $\lambda = 0$ indicating that there is no phylogenetic signal; thus, related taxa are

not more similar than expected by chance, and $\lambda = 1$ implying a strong phylogenetic signal, so

the traits of related taxa are more similar than expected by chance (Pagel 1999). Pagel's λ was

selected among the available phylogenetic signal indices given its robustness in measuring

phylogenetic signals based on incompletely resolved phylogenies (Molina-Venegas and

Rodríguez 2017). We also calculated Moran's I coefficient to assess the phylogenetic

autocorrelation of traits at the genus, family and order levels using the correlogram.formula

function from *ape* (Paradis and Schliep 2019). Values of Moran's I range from -1 to 1, with I = 0 indicating that species resemble each other as expected by the Brownian motion model. Values above and below zero imply that related species are more or less similar than expected by such model, respectively (Moran 1950; Gittleman and Kot 1990). All these tests were carried out using log-transformed seed mass values and logit-transformed water content, percentage of embryoless and viable seed values to reduce the skewness of the data. For species with more than one record (25 species), we carried out the analyses using the average species value.

For the qualitative seed traits (i.e., seed dormancy, seed dispersal syndrome and seed dispersal season), the phylogenetic signal was tested by implementing the rtestdeciv function from the *adiv* package using 9,999 permutations (Pavoine 2020). This method decomposes trait diversity among the nodes of the phylogenetic tree and assesses whether it is skewed towards the tree's root or tips, with significant root skewness implying the presence of a significant phylogenetic signal. Seed dormancy and dispersal season were treated as multichoice variables, as 11 species had records of both D and ND seeds, and 18 species reported more than one dispersal season.

To test for differences in quantitative traits between ecological groups (growth form, microhabitat, and geographic distribution), we used phylogenetic generalised least square models as implemented in the package *caper* (Orme *et al.* 2018). Likewise, qualitative trait variation was assessed using a phylogenetic logistic regression (Ives and Garland 2010) implemented in the *phylolm* package (Ho and Ané 2014). Because dispersal season and syndrome had more than two possible states, individual models were run to assess the probability of each season and syndrome. Species with records for more than one distribution class were classified as "widespread", and species with occurrence in both microhabitats were coded as "mesic/xeric". For this analysis, species with records from more than one

combination of qualitative traits were included as different populations, using the

add.taxa.phylo function from the *daee* package (Debastiani 2021) and keeping the tree

ultrametric.

Preliminary tests showed that the interaction between our qualitative predictors did not

significantly affect any of the tested variables. Moreover, the models that included such

interactions were outperformed by simple additive models based on Akaike Information

Criteria (see code provided). As a result, our final models did not include these interactions.

Growth form comparisons were made between herbs and shrubs only due to the low sample

size of trees (26 species), succulents (nine species) and lianas (six species).

Meta-analyses of germination responses to abiotic factors

We estimated the overall effects of i) light, ii) constant and alternate temperatures, and

iii) fire-related cues on germination through meta-analyses. We achieved this by fitting

phylogenetic generalised linear mix models with Bayesian estimations using the Markov

chain Monte Carlo (MCMC) method –an approach that has been previously used in meta-

analyses of primary germination data (Vandelook et al. 2018; Fernández-Pascual et al. 2021;

Fernández-Pascual et al. 2022; Carta, Fernández-Pascual, et al. 2022). This analysis was

restricted to *campo rupestre* and *canga* species, given the lack of data from *campo de altitude*

and the small sample size of independent observations for inselbergs (two studies, three

species) for these abiotic factors. Germination records under complete darkness were used as

controls to test the effects of light. Also, given the overall positive effect of light on

germination (see Results), only experiments carried out under light conditions were used to

assess the effects of different temperatures and fire-related cues on seed germination. We used

25 ºC as a control for constant and alternate temperatures because it is considered the

optimum temperature for most species in this ecosystem (Garcia and Oliveira 2007; Nunes *et*

al. 2016; Garcia *et al.* 2020). Finally, we used untreated seeds (i.e., seeds not exposed to

either heat shock or smoke) as controls for fire-related cues.

We assessed the effects of these abiotic factors on the proportion of germinated seeds

and median germination time (t_{50}). The number of germinated and ungerminated seeds was

taken from the raw germination data deposited in *Rock n' Seeds*. To avoid confounding

effects from treatments with extremely low germination, we only included observations where

either the control or treatment had ≥10% germination. The t_{50} (Coolbear *et al.* 1984 as

modified by Farooq *et al.* 2005) was calculated for the experiments that included raw daily

germination data using the *germinationmetrics* package (Aravind *et al.* 2022). We calculated

this index only for observations where both the control and treatment conditions had ≥10%

germination. Moreover, we assessed the effect of light, constant and alternate temperatures as

germination cues in ND seeds or in seeds where dormancy has been experimentally

alleviated. Alternate temperatures are known to break physical dormancy (Baskin and Baskin

2014), but none of the studies in the dataset have tested this. In contrast, we included D

species in the meta-analyses of fire-related cues because these are known to break the seed

dormancy in some species in fire-prone ecosystems (Baskin and Baskin 2014). Species with

non-conclusive dormancy records were excluded.

We evaluated the effect of each abiotic factor (i.e. light, temperature and fire) on

germination proportions and t_{50} on the whole dataset and separately for each growth form

(herbs and shrubs), geographic distribution (widespread and restricted) and microhabitat

(mesic, xeric, and mesic/xeric). We also tested the interaction of seed mass and seed

dormancy on germination responses in the whole dataset, and the interaction effect of seed

dormancy only for responses to fire-related cues. Phylogeny, species identity, study and

observations were included as random variables. Because seed mass exhibited a significant,

strong phylogenetic signal (see Results), the missing seed mass values for 54 species were

inputted using average values for the genera from *Rock n' Seeds* (Ordóñez-Parra *et al.* 2023)

and the Seed Information Database (https://ser-sid.org/). The analyses were performed with

the MCMCglmm package (Hadfield 2010). All models were run with weakly informative

priors, with parameter-expanded priors for the random effects. Each model was run for

500,000 MCMC steps, with an initial burn-in phase of 50,000 and a thinning interval of 50

(de Villemereuil and Nakagawa 2014), resulting in an average of 9000 posterior distributions.

We calculated mean parameter estimates and 95% credible intervals (CIs) from the resulting

posterior distributions. CIs overlapping zero were considered non-significant. To estimate the

phylogenetic signal of germination responses, we also estimated the Pagel's λ of each model,

as described by de Villemereuil and Nakagawa (2014).

RESULTS

Phylogenetic signal and variation of seed functional traits

All examined seed traits exhibited a significant phylogenetic signal (Figure 2, Table 1,

$p < 0.001$). For all quantitative traits (i.e., seed mass, water content, percentage of embryoless

and percentage of viable seeds), the phylogenetic signal was moderate to strong (λ between

0.57-0.90), meaning that closely related taxa strongly resembling each other both at the genus

and family level (Table S1). At the order level, we only found a significant and negative

phylogenetic autocorrelation for seed mass (I = -0.39, $p < 0.001$, Table S1). Our data did not

suggest the presence of phylogenetic autocorrelation for seed water content and the

percentage of embryoless and viable seeds at the order level ($p > 0.05$). A summary of family-

level variation in quantitative and qualitative traits is presented in Figure S2.

Regarding the variation of quantitative traits, seed mass variation differed between

growth forms and microhabitats, with shrubs (t = 2.19, p = 0.030) and species inhabiting both

mesic and xeric microhabitats (t = 2.01, p = 0.045) producing heavier seeds than herbs and

species restricted to xeric microhabitats, respectively. Seed water content, percentage of

embryoless and viable seeds did not vary between any of the groups, and seed mass did not

differ between species with distinct geographic distributions (Table 2).

For qualitative traits, we found that shrubs had a higher probability of dispersing seeds

during the late-dry season (z = 3.43, p < 0.001), while dispersal during the late-rainy season

was more likely in herbs (z = -2.01, p = 0.044). Primary dormancy and dispersal season varied

significantly between microhabitats, with species from xeric environments having higher

probabilities of producing dormant seeds (z = 2.52, p = 0.012) and dispersing their seeds

during the late-rainy season than species from mesic microhabitats (z = 2.15, p = 0.031).

Similarly, species from both mesic and xeric microhabitats also had a higher probability of

dispersing seeds during the late-rainy season (z = 2.03, p = 0.042). We found no significant

differences in the probability of any of the dispersal syndromes or the other dispersal seasons

(Table 2).

In summary, shrubs tended to produce heavier seeds and to have higher probabilities

of dispersal during the late-dry season. In contrast, herbs produced relatively smaller seeds

with higher probabilities of dispersal during the late-rainy season. Species from mesic/xeric

microhabitats tended to produce heavier seeds and disperse them during the late-rainy season

compared to those exclusively from mesic microhabitats. Moreover, species restricted to xeric

environments had higher probabilities of producing dormant seeds during late-rainy dispersal

than those from mesic environments. Finally, species restricted to outcrop vegetation and

widespread ones did not differ in any seed trait (Figure 3).

Meta-analyses of germination responses

Light positively and strongly affected the final germination proportion regardless of

growth form, distribution, or microhabitat. Still, light effects were higher for shrubs and

species from mesic environments compared to herbs and species from other microhabitats,

respectively. Seed responses to light did not vary between species with different geographic

distributions. Seed mass had a negative interactive effect with responses to light (Figure 4).

Regarding the overall effect of constant temperatures on germination, temperatures

below 20 °C had no significant effect on germination proportions when compared to the

control. However, temperatures below and above 20 °C had a consistent negative effect on

germination proportion and a positive effect on t_{50}. Herbs and shrubs exhibited differential

responses to temperature regimes: germination proportion decreased more with decreasing

temperature for shrubs, whereas the final germination proportion of herbs increased at 20 °C

and remained relatively constant at 30 °C. For instance, the t_{50} of herbs was not significantly

affected by 30 °C, contrasting with shrubs for which germination was slowed at this

temperature. Differential responses were also found for species with contrasting geographic

distributions, with species with widespread distribution showing a more severe reduction in

the germination proportion at 10 °C as well as a significant negative reduction at 30 °C.

Similarly, the germination proportion of species from mesic microhabitats was less negatively

affected by 15 °C conditions but was more inhibited at 35 and 40 °C than species from xeric

microhabitats. Finally, seed mass exhibited a significant interaction with germination

responses to temperature, being positive for germination proportion at 10 and 35 °C and

negative at 20 and 40 °C. Still, we did not find a significant interaction between seed mass

and germination responses at 15 and 30 °C (Figure 5A).

When evaluating the effects of alternate temperatures, we found that 25/15 and 30/20

°C regimes had an overall negative and a non-significant effect on germination proportion,

respectively. Moreover, both treatments significantly increased t_{50}. 30/20 °C regime also had

a negative effect on the germination proportion in shrubs and increased t_{50} in herbs. Seed

mass had a negative interaction with germination proportion at 25/15 °C and a positive

interaction with t_{50} at both treatments (Figure 5B).

Finally, we found that fire-related cues reduced germination proportion while also

affecting t_{50}. The effect varied according to the treatment applied, with heat shocks –alone or

in combination with smoke– reducing germination proportion but not germination time.

Instead, smoke alone or in combination with heat did not significantly affect germination

proportion but significantly reduced t_{50} (Figure 6A). When comparing heat shock treatments,

we found that treatments heat shocks of 100 (both of 1 and 5 minutes) and 200 °C for 1

minute significantly reduced germination proportion, although they did not affect t_{50} (Figure

6B). This negative effect of heat shocks on germination proportion varied between growth

forms, species distribution and microhabitats, with only shrubs, restricted species and species

from xeric microhabitats being negatively affected by heat shocks. Instead, heat shocks did

not significantly affect herbs, widespread species, and species from mesic microhabitats

(Figure 6C). The negative effect of smoke on t_{50} remained significant for shrubs but was non-

significant in herbs. Our analysis did not indicate that smoke's effect was modulated by seed

mass (Figure 6D). Contrastingly, primary dormancy and seed mass altered germination

responses to heat shocks, with ND seeds exhibiting a reduction in germination proportion and

seed mass holding a positive effect (Figure 6C).

Overall, phylogeny played a major role in germination responses to all these abiotic

factors, with most of our models having a λ significantly different form zero. Moreover, in

most cases, species identity had a stronger effect on germination responses variability than

study or observations within studies (Table S2 and S3).

DISCUSSION

Seed mass shapes germination responses in **campo rupestre**

Our analysis points out a significant interaction between seed mass and germination

responses to light, constant temperature and heat shocks, supporting the long-standing first

hypothesis that seed mass modulates germination responses. Regarding light, we found a

significant and negative interaction between seed mass and germination responses to light,

with larger seeds being less responsive to light availability. Such results support the

hypothesis raised by Nunes *et al.* (2016) on the existence of this trade-off in the *campo*

rupestre (Milberg *et al.* 2000; Pearson *et al.* 2003).

We also found that seed mass had a significant interaction with response to some

temperatures. Nevertheless, there was no overall pattern between germination responses to

temperature and seed mass, with a positive interaction with changes in germination proportion

at 10 and 30 °C, a negative one at 20 and 40 °C and non-significant interactions at 15 and 35

°C. Despite being significant, the effect of such interactions is rather small (see Table S2 and

S3), which agrees with the results of Arène *et al.* (2017), who found a weak association

between cardinal temperatures and seed mass that disappeared when accounting for

phylogenetic relatedness. Therefore, it seems reasonable to conclude that variation in

germination responses in *campo rupestre* is probably better explained by phylogeny rather

than by seed mass, an observation supported by the high λ values found in our models.

Moreover, the small seed mass variation in our dataset (86% of records < 1 mg) could prevent

us from finding a more consistent pattern for the role of seed mass in this ecosystem.

We also found a significant interaction between seed mass and germination responses

to heat shocks that indicate that *campo rupestre* species with larger seeds tend to have

increased and faster germination after a heat shock, a result that is consistent with a positive

correlation between seed mass and heat tolerance reported in the Cerrado (Ribeiro *et al.* 2015;

Daibes *et al.* 2019). Such association can be explained by decreasing surface-to-volume ratio

with increasing seed mass, indicating that larger, heavier seeds have higher insulation against

higher, lethal temperatures (Ruprecht *et al.* 2015).

Finally, the lack of a significant effect of alternate temperatures or an interaction of

such responses with seed mass could be explained by the fact that most *campo rupestre*

species produce small, lighter seeds, which are not expected to benefit from alternate

temperature regimes (Pearson *et al.* 2003).

Seed functional traits are phylogenetically clustered in Brazilian rock outcrop vegetation

Our results indicate that all seed traits showed moderate-to-strong phylogenetic

signals, which is consistent with global studies showing phylogenetic clustering for multiple

seed traits, such as seed mass (Moles *et al.* 2005) and dormancy (Willis *et al.* 2014).

However, our results partially contrast with local studies in similar ecosystems. First, Zanetti

et al. (2020) did not find a phylogenetic signal for seed water content in their study of 48

species from the *cangas* of Carajás (Eastern Brazilian Amazon). This discrepancy between

this and our study might arise from differences in sample size and geographic extent or the

distinct phylogenetic structure of each rock outcrop vegetation type (Massante *et al.* 2023).

Second, we found a significant phylogenetic signal for seed dispersal season, while seed

dispersal season in the Cerrado (Escobar *et al.* 2021) and other phenological events in the

campo rupestre (Zanetti *et al.* 2020; Oliveira *et al.* 2021) have not shown such signal. As a

result, seed dispersal season is the only phenophase that has shown a significant phylogenetic

signal in the Brazilian rock outcrop vegetation, potentially because of an intense evolutionary

pressure towards germination timing in these harsh ecosystems, where water is even scarcer

than in the Cerrado (Oliveira et al. 2021), thus may create stronger selective pressures for

establishment.

Seed mass and seed dispersal season differed between herbs and shrubs, with shrubs

tending to produce relatively larger seeds during the late-dry season and herbs producing

lighter seeds during the late-rain season. Differences in seed mass between herbs and shrubs

have been attributed to the distinct strategies employed by such growth forms: while shrubs

invest in fewer, larger seeds that cope better with environmental hazards, herbs invest in

numerous, lighter seeds to increase their establishment opportunities (Westoby *et al.* 2002;

Moles *et al.* 2005). This relationship is also supported by the global spectrum of plant form

and function (Díaz *et al.* 2016), suggesting that this global pattern also holds at small scales,

such as for Brazilian rock outcrop vegetation. On the other hand, differences in seed dispersal

season suggest differences in phenological strategies to cope with precipitation seasonality

(see below).

13 *Species from distinct geographic distributions and microhabitats exhibit different*
14 *germination responses (and seed traits)*

Rock outcrops and the surrounding vegetation differ in several aspects that should

impact germination, such as nutrient and water availability, irradiance, and daily temperature

fluctuations (Oliveira *et al.* 2016). While we did not find significant differences in seed traits

between restricted and widespread species, their germination responses to temperature and

heat shocks differed. First, the germination proportion of widespread species was reduced or

delayed at relatively low and high temperatures, implying they are more susceptible to these

conditions than species restricted to outcrop vegetation. Second, species restricted to outcrop

vegetation were more sensitive to heat shocks than widespread ones. The rocky substrate of

the *campo rupestre* does not favour fire spread (Conceição *et al.* 2016), so natural fire events

in these ecosystems are unlikely to reach the vegetation establishing directly on rock outcrops,

reducing evolutionary pressures towards heat tolerance. In contrast, widespread species

present in our dataset include species from the Cerrado, where fire events are relatively more frequent, and seeds exhibit relatively higher heat tolerance (Daibes *et al.* 2022).

Contrastingly, species from mesic and xeric microhabitats were found to differ both in terms of seed traits and germination responses, agreeing with local studies showing a correspondence between seed traits and species microhabitats (Oliveira and Garcia 2011; Marques et al. 2014; Ranieri et al. 2012). These results support our second hypothesis that predicts distinct germination requirements expected between species with contrasting microhabitat preferences and geographical ranges, as expected under the regeneration niche hypothesis (Grubb 1977). Interestingly, these results contrast with large-scale studies carried out in temperate ecosystems where an association between ecological preferences and germination responses has been found (Fernández-Pascual *et al.* 2021; Fernández-Pascual *et al.* 2022), suggesting that the extreme abiotic conditions of both the *campo rupestre* and rock outcrop vegetation as a whole impose a stronger selective pressure to germination traits. However, seed traits did not differ in species with distinct geographic ranges suggesting that multiple pressures operate on the selection of seed traits at larger scales.

Regardless of microhabitat, most species in our dataset dispersed their seeds during the late- or the early-dry season. However, species from xeric habitats had higher probabilities of producing dormant seeds and dispersing their seeds during the late-rain season –two strategies presumably arising due to evolutionary pressures towards strategies to synchronise germination with optimum conditions for seedling establishment, that is under higher soil water availability (Jurado and Flores 2005; Silveira, Ribeiro, *et al.* 2012; Garcia *et al.* 2020). Nevertheless, only 36% of our species produce dormant seeds, and *campo rupestre* is known to have the highest ND:D ratio globally (Dayrell *et al.* 2017), implying that seed dormancy might not be the main driver of seedling establishment in our study system. Additional strategies to control germination timing include germination requirements and the acquisition

of secondary dormancy. A mismatch between temperature germination requirements and

environmental conditions at the moment of dispersal might provide an alternative strategy to

prevent germination in the absence of dormancy (Escobar *et al.* 2021) –for example, seeds

dispersed during the dry season have evolved to germinate under relatively higher

temperatures of the rainy season. Furthermore, secondary dormancy, as reported in a few

Eriocaulaceae and Xyridaceae species from *campo rupestre* (see Garcia *et al.* 2020), can

contribute to fine-tuning seedling establishment with optimum conditions. All these species

disperse their non-dormant seeds between the early-dry and the early-rain season and become

increasingly dormant as the rainy season advances. When the dry season starts, dormancy is

progressively alleviated by reduced temperatures and low water availability, allowing

germination at the onset of the rainy season (Duarte and Garcia 2015).

Species from mesic microhabitats also exhibit distinct responses to light and

temperature, with stronger positive responses to light and being more tolerant to low (below

20 °C) but more sensitive to high temperatures (above 30 °C). As a result, a stronger response

to light could ensure that germination under light conditions occurs as quickly as possible.

Moreover, high soil humidity in mesic habitats buffers these habitats from wide temperature

fluctuations and contributes to maintaining relatively low soil temperatures (Oliveira and

Garcia, 2011).

Campo rupestre *species depend on light for germination*

Our meta-analysis showed that light positively affects seed germination across all

ecological groups, supporting previous assessments about its importance in *campo rupestre*

germination ecology. Interestingly, we also found that germination responses to light were

stronger in shrubs, a result that could be attributed to the fact that species with the lightest

seeds in our dataset are shrubs from the Melastomataceae (see Figure S2), which is both the

best-represented family in our dataset (Ordóñez-Parra *et al.* 2023) and a family characterised

by an absolute light requirement for germination (Ordóñez-Parra *et al.* 2022).

Small-seeded species from our study are expected to have narrowly defined microsite

requirements due to their limited internal resources, so further aspects of the light

environment, such as spectral quality and the interaction between light and temperature, are

expected to control their germination (Pearson *et al.* 2003; Pons 2014). Still, studies assessing

such aspects in our study system are scarce and have only involved very few species (Pereira

et al. 2009; Hmeljevski *et al.* 2014; Vieira *et al.* 2018; Garcia *et al.* 2020), preventing robust

inferences about the functional relevance of these germination cues.

Herbs and shrubs respond differently to low and high temperatures

We found that the germination of *campo rupestre* species was maximised between 20-

25 ºC, supporting previous qualitative reviews (Nunes *et al.* 2016; Garcia *et al.* 2020).

Decreased germination proportion below this range could be a putative mechanism to avoid

germination during the dry season when temperatures decrease and soil water potential does

not support seedling establishment (Garcia *et al.* 2020). The negative effect of low

temperatures was stronger in shrubs, suggesting that these require higher temperatures to

germinate. In contrast, herbs had increased germination at 20 °C but were unaffected by 30 °C

conditions. These results suggest that herbs dominate microsites experiencing lower soil

temperatures.

Alternate temperature regimes had either a negative or null effect on the germination.

The reduction of germination proportion and the increase of germination time at 25/15 ºC

regime is probably due to part of these temperatures being suboptimal. Conversely, studies in

the *campo de altitude* have shown that several species benefit from alternate temperature

regimes, suggesting this cue could be more important in this particular vegetation that occurs

on higher elevation belts (Andrade *et al.* 2021). Therefore, additional studies are required to

elucidate the functional relevance of alternate temperatures and whether this varies between

vegetation types.

Heat kills small, non-dormant seeds, but smoke accelerates germination

Seeds exposed to 100 or 200 ºC heat shocks had their germination proportion reduced,

while milder heat shocks (≤ 80 ºC) did not affect germination proportion or time, supporting

studies showing heat tolerance in the Cerrado (Daibes *et al.* 2022). In addition to seed mass,

responses to heat shocks were also moderated by seed dormancy, with ND seeds being

relatively less tolerant to heat shocks, as previously described for Cerrado species (Ramos *et*

al. 2016; Daibes *et al.* 2019). Dormant species were mostly unaffected by heat shocks,

supporting the notion that fire-mediated dormancy alleviation is not expected in rocky areas

or savannas (Pausas and Lamont 2022). The lack of detrimental effects on dormant species

also implies that traits associated with seed dormancy promote heat tolerance, such as the

accumulation of heat shock proteins or the presence of hard and water-impermeable coats

(Tweddle *et al.* 2003; Ramos *et al.* 2016).

Smoke did not affect germination proportion but accelerated germination, supporting

similar results found for Cerrado vegetation (Motta *et al.* 2024) and the notion that smoke-

derived compounds stimulate germination of ND seeds or those where dormancy has been

alleviated (Mackenzie *et al.* 2021). Smoke-stimulated germination in *campo rupestre* is

thought to promote the germination of species resprouting and shedding seeds after fire

(Fernandes *et al.* 2021) –a usual phenological syndrome in this vegetation (Figueira *et al.*

2016)– allowing recently-dispersed seeds to exploit the post-fire environment where

competition is relaxed (Le Stradic *et al.* 2015; Fernandes *et al.* 2021). While our meta-

analysis supports this hypothesis, we found that the hastening effect of smoke is minor and

only significant for shrubs. Moreover, such an effect has only been tested on a handful of species, and further studies are needed to rigorously test it.

Conclusion

Seed germination ecology in rock outcrop vegetation in Brazil is strongly shaped by species phylogenetic relatedness, as illustrated by the moderate-to-strong phylogenetic signal exhibited across seed traits and germination responses. As a result, knowledge about one species can provide highly relevant information about the germination ecology of other closely related species. Despite that, our analyses also point out that seed mass can shape germination responses that may differ between species with different growth form, species geographic distribution and microhabitats. Our integrative synthesis provides robust support for long-standing hypotheses in the seed ecology of Brazilian rock outcrop vegetation. We hope that our synthesis paves the way for future research about the functional role of seed traits in these megadiverse ecosystems and a solid baseline for using seeds in ecological restoration and biological conservation programs, such as direct seeding and seed banking.

ACKNOWLEDGEMENTS

The first author dedicates this article to the memory of Martha Galeano (1964-2022), who was fascinated with *tepuis* and whose Vegetation Studies lecture inspired him to pursue a career in plant ecology. This study is part of the first author's Master Dissertation at the Plant Biology Program at Universidade Federal de Minas Gerais. CAO-P and NFM were supported by a scholarship from CAPES and CNPq, respectively. FAOS acknowledges support from FAPEMIG. We also thank James Dalling and Sergey Rosbakh, who provided valuable comments to an earlier version of the manuscript, and Rohit Goswami, who helped in elaboration of the figures and the organization of the GitHub repository.

OPEN DATA

The R code for the analysis and creation of the figures will be provided as Supplementary

Material and uploaded to GitHub upon acceptance.

LITERATURE CITED

Alcantara S, Ree RH, Mello-Silva R. **2018**. Accelerated diversification and functional trait
evolution in Velloziaceae reveal new insights into the origins of the *campos rupestres*'
exceptional floristic richness. *Annals of Botany* **122**: 165–180.
Almeda F, Michelangeli FA, Viana PL. **2016**. *Brasilianthus* (Melastomataceae), a new
monotypic genus endemic to ironstone outcrops in the Brazilian Amazon. *Phytotaxa* **273**:
269–282.
Aravind J, Vimala Devi S, Radhamani J, Jacob SR, Srinivasan K. **2022**.
germinationmetrics: seed germination indices and curve fitting. R package version 0.1.8.9,
https://cran.r-project.org/web/packages/germinationmetrics/.
Arène F, Affre L, Doxa A, Saatkamp A. **2017**. Temperature but not moisture response of
germination shows phylogenetic constraints while both interact with seed mass and lifespan.
Seed Science Research **27**: 110–120.
Barthlott W, Porembski S. **2000**. Why study inselbergs? In: Porembski S, Barthlott W, eds.
Inselbergs. Berlin: Springer, 1–6.
Baskin CC, Baskin JM. **2014**. *Seeds: ecology, biogeography, and evolution of dormancy*
and germination. San Diego: Elsevier Inc.
Biedinger N, Porembski S, Barthlott W. **2000**. Vascular plants on inselbergs: vegetative
and reproductive strategies In: Porembski S, Barthlott W, eds. *Inselbergs*. Berlin, Heidelberg:
Springer, 117–142.
Bondi L, de Paula LFA, Rosado BHP, Porembski S. **2023**. Demystifying the convergent
ecological specialization of desiccation-tolerant vascular plants for water deficit. *Annals of*
Botany **131**: 521–531.
Campitelli E. **2022**. ggnewscale: multiple fill and colour scales in "ggplot2." R package
version 0.4.10, https://cran.r-project.org/package=ggnewscale
Campos P V., Villa PM, Nunes JA, Schaefer CEGR, Porembski S, Neri A V. **2018**. Plant
diversity and community structure of Brazilian Páramos. *Journal of Mountain Science* **15**:
1186–1198.
Carta A, Fernández-Pascual E, Gioria M, *et al*. **2022**. Climate shapes the seed germination
niche of temperate flowering plants: a meta-analysis of European seed conservation data.
Annals of Botany **129**: 775–786.
Carta A, Mattana E, Dickie J, Vandelook F. **2022**. Correlated evolution of seed mass and
genome size varies among life forms in flowering plants. *Seed Science Research* **32**: 46–52.
Conceição AA, Rapini A, do Carmo FF, *et al*. **2016**. Rupestrian Grassland Vegetation,
Diversity, and Origin In: *Ecology and Conservation of Mountaintop grasslands in Brazil*.
Cham: Springer International Publishing, 105–127.
Coolbear P, Francis A, Grierson D. **1984**. The effect of low temperature pre-sowing
treatment on the germination performance and membrane integrity of artificially aged tomato
seeds. *Journal of Experimental Botany* **35**: 1609–1617.

Daibes LF, Ordóñez-Parra CA, Dayrell RLC, Silveira FAO. 2022. Regeneration from
seeds in South American savannas, in particular the Brazilian Cerrado In: Baskin CC, Baskin
JM, eds. *Plant regeneration from seeds: a global warming perspective*. Elsevier, 183–197.
Daibes LF, Pausas JG, Bonani N, Nunes J, Silveira FAO, Fidelis A. 2019. Fire and legume
germination in a tropical savanna: ecological and historical factors. *Annals of Botany* **123**:
1219–1229.
Dayrell RLC, Garcia QS, Negreiros D, Baskin CC, Baskin JM, Silveira FAO. 2017.
Phylogeny strongly drives seed dormancy and quality in a climatically buffered hotspot for
plant endemism. *Annals of Botany* **119**: 267–277.
Debastiani VJ. 2021. Data Analysis for Ecology and Evolution. R package version
0.1.7, https://github.com/vanderleidebastiani/daee (accessed November 2021).
Díaz S, Kattge J, Cornelissen JHC, *et al.* 2016. The global spectrum of plant form and
function. *Nature* **529**: 167–171.
Donohue K, Rubio De Casas R, Burghardt L, Kovach K, Willis CG. 2010. Germination,
postgermination adaptation, and species ecological ranges. *Annual Review of Ecology,*
Evolution, and Systematics **41**: 293–319.
Duarte DM, Garcia QS. 2015. Interactions between substrate temperature and humidity in
signalling cyclical dormancy in seeds of two perennial tropical species. *Seed Science*
Research **25**: 170–178.
Escobar DFE, Rubio de Casas R, Morellato LPC. 2021. Many roads to success: different
combinations of life-history traits provide accurate germination timing in seasonally dry
environments. *Oikos* **130**: 1865–1879.
Escudero A, Palacio S, Maestre FT, Luzuriaga AL. 2015. Plant life on gypsum: a review
of its multiple facets. *Biological Reviews* **90**: 1–18.
Farooq M, Basra SMA, Ahmad N, Hafeez K. 2005. Thermal hardening: a new seed vigor
enhancement tool in rice. *Journal of Integrative Plant Biology* **47**: 187–193.
Fernandes AF, Oki Y, Fernandes GW, Moreira B. 2021. The effect of fire on seed
germination of *campo rupestre* species in the South American Cerrado. *Plant Ecology* **222**:
45–55.
Fernández-Pascual E, Carta A, Mondoni A, *et al.* 2021. The seed germination spectrum of
alpine plants: a global meta-analysis. *New Phytologist* **229**: 3573–3586.
Fernández-Pascual E, Vaz M, Morais B, *et al.* 2022. Seed ecology of European mesic
meadows. *Annals of botany* **129**: 121–134.
Figueira JEC, Ribeiro KT, Ribeiro MC, *et al.* 2016. Fire in rupestrian grasslands: plant
response and management In: *Ecology and conservation of mountaintop grasslands in Brazil*.
Cham: Springer International Publishing, 415–448.
Fitzsimons JA, Michael DR. 2017. Rocky outcrops: a hard road in the conservation of
critical habitats. *Biological Conservation* **211**: 36–44.
Freiberg M, Winter M, Gentile A, *et al.* 2020. LCVP, The Leipzig catalogue of vascular
plants, a new taxonomic reference list for all known vascular plants. *Scientific Data* **7**: 1–7.
Garcia QS, Barreto LC, Bicalho EM. 2020. Environmental factors driving seed dormancy
and germination in tropical ecosystems: a perspective from *campo rupestre* species.
Environmental and Experimental Botany **178**: 104164.
Garcia QS, Oliveira PG. 2007. Germination patterns and seed longevity of monocotyledons
from the Brazilian *campos rupestres*. *Seed Science and Biotechnology* **1**: 35–41.
Giorni VT, Bicalho EM, Garcia QS. 2018. Seed germination of *Xyris* spp. from Brazilian
campo rupestre is not associated to geographic distribution and microhabitat. *Flora* **238**: 102–
109.
Gittleman JL, Kot M. 1990. Adaptation: statistics and a null model for estimating
phylogenetic effects. *Systematic Zoology* **39**: 227.

Grubb PJ. **1977**. The maintenance of species-richness in plant communities: the importance of the regeneration niche. *Biological Reviews* **52**: 107–145.

Guimarães PJF, Michelangeli FA, Sosa K, Santiago Gómez JR. **2019**. Systematics of *Tibouchina* and allies (Melastomataceae: Melastomateae): A new taxonomic classification. *Taxon* **68**: 937–1002.

Hadfield JD. **2010**. MCMC Methods for Multi-Response Generalized Linear Mixed Models: The **MCMCglmm** *R* Package. *Journal of Statistical Software* **33**.

Hmeljevski K V., Freitas L, Domingues R, *et al.* 2014. Conservation assessment of an extremely restricted bromeliad highlights the need for population-based conservation on granitic inselbergs of the Brazilian Atlantic Forest. *Flora* **209**: 250–259.

Ho LST, Ané C. **2014**. A linear-time algorithm for gaussian and non-gaussian trait evolution models. *Systematic Biology* **63**: 397–408.

Ives AR, Garland T. **2010**. Phylogenetic logistic regression for binary dependent variables. *Systematic Biology* **59**: 9–26.

Jacobi CM, Carmo FF, Vincent RC, Stehmann JR. **2007**. Plant communities on ironstone outcrops: a diverse and endangered Brazilian ecosystem. *Biodiversity and Conservation* **16**: 2185–2200.

Jin Y, Qian H. **2022**. V.PhyloMaker2: An updated and enlarged R package that can generate very large phylogenies for vascular plants. *Plant Diversity*: 0–4.

Jurado E, Flores J. **2005**. Is seed dormancy under environmental control or bound to plant traits? *Journal of Vegetation Science* **16**: 559–564.

Kluge M, Brulfert J. **2000**. Ecophysiology of vascular plants on inselbergs In: *Inselbergs*. Berlin, Heidelberg: Springer, 143–174.

Larson JE, Funk JL. **2016**. Regeneration: an overlooked aspect of trait-based plant community assembly models. *Journal of Ecology* **104**: 1284–1298.

Mackenzie BDE, Auld TD, Keith DA, Ooi MKJ. **2021**. Fire seasonality, seasonal temperature cues, dormancy cycling, and moisture availability mediate post-fire germination of species with physiological dormancy. *Frontiers in Plant Science* **12**.

Marques AR, Atman APFF, Silveira FAOO, Lemos-Filho JP. **2014**. Are seed germination and ecological breadth associated? Testing the regeneration niche hypothesis with bromeliads in a heterogeneous neotropical montane vegetation. *Plant Ecology* **215**: 517–529.

Martinelli G. **2007**. Mountain biodiversity in Brazil. *Revista Brasileira de Botanica* **30**: 587–597.

Massante JC, Neri AV, Villa PM, *et al.* 2023. Looking similar but all different: Phylogenetic signature of Brazilian rocky outcrops and the influence of temperature variability on their phylogenetic structure. *Journal of Ecology* **111**: 1905–1920.

Milberg P, Andersson L, Thompson K. **2000**. Large-seeded species are less dependent on light for germination than small-seeded ones. *Seed Science Research* **10**: 99–104.

Miola DTB, Ramos VDV, Silveira FAO. **2021**. A brief history of research in *campo rupestre*: identifying research priorities and revisiting the geographical distribution of an ancient, widespread Neotropical biome. *Biological Journal of the Linnean Society* **133**: 464–480.

Moles AT, Ackerly DD, Webb CO, *et al.* 2005. Factors that shape seed mass evolution. *Proceedings of the National Academy of Sciences of the United States of America* **102**: 10540–10544.

Molina-Venegas R, Rodríguez MÁ. **2017**. Revisiting phylogenetic signal; strong or negligible impacts of polytomies and branch length information? *BMC Evolutionary Biology* **17**: 53.

Moran PAP. **1950**. Notes on continuous stochastic phenomena. *Biometrika* **37**: 17.

Motta GST, Pilon N, Fidelis A, Kolb RM. **2024**. Smoke effects on the germination of Cerrado species. *Plant Ecology* **225**: 685–693.

Nunes FP, Dayrell RLC, Silveira FAO, *et al.* 2016. Seed germination ecology in rupestrian grasslands In: Fernandes GW, ed. *Ecology and conservation of mountaintop grasslands in Brazil*. Springer International Publishing, 207–225.

Oliveira RS, Abrahão A, Pereira C, *et al.* 2016. Ecophysiology of *campos rupestres* plants In: *Ecology and conservation of mountaintop grasslands in Brazil*. Cham: Springer International Publishing, 227–272.

Oliveira PG, Garcia QS. **2011**. Germination characteristics of *Syngonanthus seeds* (Eriocaulaceae) in *campos rupestres* vegetation in south-eastern Brazil. *Seed Science Research* **21**: 39–45.

Oliveira CS, Messeder JVS, Teixido AL, Arantes MRR, Silveira FAO. **2021**. Vegetative and reproductive phenology in a tropical grassland–savanna–forest gradient. *Journal of Vegetation Science* **32**: 1–16.

Ordóñez-Parra CA, Dayrell RLC, Negreiros D, *et al.* 2023. Rock n' Seeds: A database of seed functional traits and germination experiments from Brazilian rock outcrop vegetation. *Ecology* **104**: e3852.

Ordóñez-Parra CA, Messeder JVS, Mancipe-Murillo C, Calderón-Hernández M, Silveira FAO. 2022. Seed Germination Ecology in Neotropical Melastomataceae: Past, Present, and Future In: *Systematics, Evolution, and Ecology of Melastomataceae*. Cham: Springer International Publishing, 707–733.

Orme D, Freckleton R, Thomas G, *et al.* 2018. Comparative analyses of phylogenetics and evolution in R. R package version 1.0.3, https://cran.r-project.org/web/packages/caper/index.html.

Pagel M. **1999**. Inferring the historical patterns of biological evolution. *Nature* **401**: 877–884.

Paradis E, Schliep K. **2019**. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* **35**: 526–528.

Pausas JG, Lamont BB. **2022**. Fire-released seed dormancy - a global synthesis. *Biological Reviews* **97**: 1612-1639.

Pavoine S. **2020**. adiv: an R package to analyse biodiversity in ecology (S Goslee, Ed.). *Methods in Ecology and Evolution* **11**: 1106–1112.

Pearson TRH, Burslem DFRP, Mullins CE, Dalling JW. **2003**. Functional significance of photoblastic germination in neotropical pioneer trees: a seed's eye view. *Functional Ecology* **17**: 394–402.

Pereira AR, Andrade ACS, Pereira TS, Forzza RC, Rodrigues AS. **2009**. Comportamento germinativo de espécies epífitas e rupícolas de bromeliaceae do Parque Estadual do Ibitipoca, Minas Gerais, Brasil. *Revista Brasileira de Botanica* **32**: 827–838.

Pons TL. **2014**. Light-mediated germination. In: Gallagher RS, ed. *Seeds: the ecology of regeneration in plant communities*. UK: CABI, 111–134.

Porembski S. **2007**. Tropical inselbergs: habitat types, adaptive strategies and diversity patterns. *Revista Brasileira de Botanica* **30**: 579–586.

Qian H, Jin Y. **2021**. Are phylogenies resolved at the genus level appropriate for studies on phylogenetic structure of species assemblages? *Plant Diversity* **43**: 255–263.

R Core Team. **2023**. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/

Ramos DM, Liaffa ABS, Diniz P, *et al.* 2016. Seed tolerance to heating is better predicted by seed dormancy than by habitat type in Neotropical savanna grasses. *International Journal of Wildland Fire* **25**: 1273–1280.

Ranieri BD, Pezzini FF, Garcia QS, Chautems A. **2012**. Testing the regeneration niche
hypothesis with Gesneriaceae (tribe Sinningiae) in Brazil: Implications for the conservation of
rare species. *Austral Ecology* **37**: 125–133.
Revell LJ. **2012**. phytools: an R package for phylogenetic comparative biology (and other
things). *Methods in Ecology and Evolution* **3**: 217–223.
Ribeiro LC, Barbosa ERM, van Langevelde F, Borghetti F. **2015**. The importance of seed
mass for the tolerance to heat shocks of savanna and forest tree species. *Journal of Vegetation*
Science **26**: 1102–1111.
Rivera VL, Panero JL, Schilling EE, Crozier BS, Moraes MD. **2016**. Origins and recent
radiation of Brazilian Eupatorieae (Asteraceae) in the eastern Cerrado and Atlantic Forest.
Molecular Phylogenetics and Evolution **97**: 90–100.
Ruprecht E, Fenesi A, Fodor EI, Kuhn T, Tökölyi J. **2015**. Shape determines fire tolerance
of seeds in temperate grasslands that are not prone to fire. *Perspectives in Plant Ecology,*
Evolution and Systematics **17**: 397–404.
Saatkamp A, Cochrane A, Commander L, *et al.* 2019. A research agenda for seed-trait
functional ecology. *New Phytologist* **221**: 1764–1775.
Safford HD. **1999**. Brazilian Páramos I. An introduction to the physical environment and
vegetation of the campos de altitude. *Journal of Biogeography* **26**: 693–712.
Safford HD, Martinelli Gustavo. **2000**. Southeast Brazil In: DeForest Safford H, Martinelli
G., eds. *Inselbergs*. Berlin: Springer, 339–389.
Silveira FAO, Negreiros D, Araújo LM, Fernandes GW. **2012**. Does seed germination
contribute to ecological breadth and geographic range? A test with sympatric *Diplusodon*
(Lythraceae) species from rupestrian fields. *Plant Species Biology* **27**: 170–173.
Silveira FAO, Negreiros D, Barbosa NPU, *et al.* 2016. Ecology and evolution of plant
diversity in the endangered *campo rupestre*: a neglected conservation priority. *Plant and Soil*
403: 129–152.
Silveira FAO, Ribeiro RC, Oliveira DMT, Fernandes GW, Lemos-Filho JP. **2012**.
Evolution of physiological dormancy multiple times in Melastomataceae from Neotropical
montane vegetation. *Seed Science Research* **22**: 37–44.
Smith SA, Brown JW. **2018**. Constructing a broadly inclusive seed plant phylogeny.
American Journal of Botany **105**: 302–314.
Le Stradic S, Silveira FAO, Buisson E, Cazelles K, Carvalho V, Fernandes GW. **2015**.
Diversity of germination strategies and seed dormancy in herbaceous species of *campo*
rupestre grasslands. *Austral Ecology* **40**: 537–546.
Tweddle JC, Dickie JB, Baskin CC, Baskin JM. **2003**. Ecological aspects of seed
desiccation sensitivity. *Journal of Ecology* **91**: 294–304.
Vandelook F, Janssens SB, Probert RJ. **2012**. Relative embryo length as an adaptation to
habitat and life cycle in Apiaceae. *New Phytologist* **195**: 479–487.
Vandelook F, Newton RJ, Carta A. **2018**. Photophobia in Lilioid monocots: photoinhibition
of seed germination explained by seed traits, habitat adaptation and phylogenetic inertia.
Annals of Botany **121**: 405–413.
Vieira BC, Rodrigues BMA, Garcia QS. **2018**. Light exposure time and light quality on
seed germination of *Vellozia* species (Velloziaceae) from Brazilian *campo rupestre*. *Flora*
238: 94–101.
de Villemereuil P, Nakagawa S. **2014**. General Quantitative Genetic Methods for
Comparative Biology In: *Modern Phylogenetic Comparative Methods and Their Application*
in Evolutionary Biology. Berlin, Heidelberg: Springer Berlin Heidelberg, 287–303.
Westoby M, Falster DS, Moles AT, Vesk PA, Wright IJ. **2002**. Plant ecological strategies:
some leading dimensions of variation between species. *Annual Review of Ecology and*
Systematics **33**: 125–159.

Willis CG, Baskin CC, Baskin JM, *et al.* **2014**. The evolution of seed dormancy:
environmental cues, evolutionary hubs, and diversification of the seed plants. *New Phytologist*
203: 300–309.
Wyatt R. **1997**. Reproductive ecology of granite outcrop plants from the south-eastern United
States. *Journal of the Royal Society of Western Australia* **80**: 123–129.
Xu S, Dai Z, Guo P, *et al.* **2021**. ggtreeExtra: compact visualization of richly annotated
phylogenetic data. *Molecular Biology and Evolution* **38**: 4039–4042.
Yu G, Smith DK, Zhu H, Guan Y, Lam TT. **2017**. ggtree: an R package for visualization
and annotation of phylogenetic trees with their covariates and other associated data. *Methods*
in Ecology and Evolution **8**: 28–36.
Zanetti M, Dayrell RLC, Wardil M V., *et al.* **2020**. Seed functional traits provide support
for ecological restoration and ex situ conservation in the threatened Amazon ironstone
outcrop flora. *Frontiers in Plant Science* **11**: 599496.
14
15

Figure 1. The four main vegetation types associated with rock outcrops in Brazil and its representation in the *Rock n' Seeds* database (Ordóñez-Parra *et al.* 2023). A. A typical *Campo rupestre* landscape at Serra do Cipó. B. *Canga* vegetation at Parque Estadual da Serra do Rola Moça. C. *Campo de altitude* at Parque Nacional do Caparaó. D. Inselberg vegetation at Teófilo Otoni municipality. E. Percentage of studies and taxa evaluated on each vegetation type. All locations are in the State of Minas Gerais. Photos by Carlos A. Ordóñez-Parra (A), Roberta L. C. Dayrell (B), Daniela Calaça (C) and Fernando A. O. Silveira (D).

Figure 2. Phylogeny of studied species with available information on seed functional traits.

Bars represent trait values for each species. The ten families with the largest number of

1   species in the dataset are labelled. Figure elaborated with the R packages ggtree (Yu *et al.*

2   2017), ggtreeExtra (Xu *et al.* 2021) and ggnewscale (Campitelli 2022).

1   **Figure 3.** Summary of seed trait differences among growth forms, species geographic

2   distribution and microhabitat, based on results from Table 2. Notes: 1. When compared to

3   species form mesic/xeric microhabitats. 2. When compared to species from xeric

4   microhabitats.

Figure 4. Effect of light availability on germination proportion in *campo rupestre* species

(Control: total darkness). Squares indicate the posterior means of the effect for each

moderator (growth form, distribution and microhabitat) and whiskers the 95 % credible

intervals of the effect size. Numbers in parentheses indicate the number of species and

observations (each observation is a replicate of a given germination experiment) employed for

each model.

2 **Figure 5.** Germination responses to (A) constant and (B) alternate temperatures in *campo*

3 *rupestre* species (Control: 25 ºC). Squares indicate the standardised mean effect size for each

moderator (growth form, distribution and microhabitat) and whiskers the 95 % confidence

interval of the effect size. Coloured estimates indicate significant effects (i.e., where

confidence intervals do not overlap zero). Numbers in parentheses indicate the number of

species and observations (each observation is a replicate of a given germination experiment)

employed for each model. The empty panel for germination time at 10 and 40 ºC refers to a

lack of data.

Figure 6. Germination responses to fire-related cues in *campo rupestre* species (Control:

untreated seeds). (A) Overall effects. (B) Heat shocks temperatures. (C) Heat shocks without

the 200 ºC for one minute. (D) Smoke. Squares indicate the standardised mean effect size for

each moderator (growth form, distribution and microhabitat) and whiskers the 95 %

confidence interval of the effect size. Coloured estimates indicate significant effects (i.e.,

where confidence intervals do not overlap zero). Numbers in parentheses indicate the number

of species and observations (each observation is a replicate of a given germination

experiment) employed for each model.

1   **Table 1.** Results of phylogenetic signal test for seed traits in Brazilian rocky outcrop vegetation. P-values for

2   quantitative traits come from the likelihood ratio test performed by phylosig function, while it corresponds to the

3   root-to-tip skewness test performed by the rtestdecdiv function for qualitative traits.

Quantitative traits	Lambda	p-value
Seed mass	0.90	< 0.001
Water content	0.72	< 0.001
Percentage of embryoless seeds	0.75	< 0.001
Percentage of viable seeds	0.57	< 0.001
Qualitative traits	p-value	
Primary dormancy	< 0.001	
Dispersal syndrome	< 0.001	
Dispersal season	< 0.001	

Table 2. Differences in seed functional traits between growth forms (herbs vs. shrubs), and species distribution (restricted vs. widespread) and microhabitat (mesic vs. mesic/xeric vs. xeric) in Brazilian rocky outcrop vegetation. Bold values indicate p-values < 0.05. t and z correspond to the values obtained in the t- and z-tests that are part of the implementation of phylogenetic least square models and phylogenetic logistic regressions, respectively.

Quantitative traits	Growth form: Shrub		Distribution: Widespread		Microhabitat: Mesic/Xeric		Microhabitat: Xeric	
	t	p-value	t	p-value	t	p-value	t	p-value
Seed mass	**2.1872**	**0.0298**	-0.5986	0.5501	**2.0141**	**0.0453**	1.6115	0.1086
Water content	0.2789	0.7811	-0.0770	0.9388	0.6075	0.5453	0.7834	0.4358
Embryoless seeds	1.9274	0.0572	-0.2368	0.8134	0.9752	0.3322	1.2967	0.1982
Viable seeds	-1.0221	0.3085	0.2381	0.8122	-0.3040	0.7616	0.7006	0.4847
Qualitative traits	Z	p-value	z	p-value	z	p-value	z	p-value
Dormancy	1.4125	0.1578	1.5052	0.1323	1.5266	0.1268	**2.5218**	**0.0117**
Anemochory	-1.2154	0.2242	1.3943	0.1632	1.0047	0.3150	1.1497	0.2503
Autochory	0.0387	0.9691	-0.0710	0.9434	-0.0792	0.9368	-0.0998	0.9205
Zoochory	0.3290	0.7422	0.0048	0.9962	0.6844	0.4937	-0.0040	0.9968
Early-Dry	-0.9086	0.3636	-0.1266	0.8993	-1.0427	0.2971	0.4504	0.6524
Late-Dry	**3.4323**	**0.0006**	-0.7443	0.4567	0.7421	0.4580	-1.3673	0.1713
Early-Rain	0.1036	0.9175	-0.7541	0.4508	-0.4839	0.6284	0.0445	0.9645
Late-Rain	**-2.0137**	**0.0440**	-0.1826	0.8551	**2.0319**	**0.0422**	**2.1507**	**0.0315**

6
7

2 **Figure S1.** State of the art of the germination ecology of Brazilian rock outcrop vegetation. A. Percentage of

3 studies for each germination ecology topic. B. Number of species studied each major topic and their

1   combinations. C. Number of species studied for each abiotic factor and their combination. Figure made using

2   RAWGraphs (Mauri *et al*. 2017) and the venn R package (Dusa 2022).

Figure S2. Variation in seven functional seed traits among the most represented families in the database. A. Variation in quantitative traits, showing the variation across the whole dataset and the top three families with the highest and lowest values on each trait. B. Variation in qualitative traits, shown as the proportion of species presenting each trait state across the ten most represented families in the database.

1 **Table S1.** Moran's I value for the quantitative seed traits across different taxonomic levels. Bold values indicate

2 values significantly different from zero.

Seed trait	Genus		Family		Order	
	I	p.value	I	p.value	I	p.value
Seed mass	**0.7035**	**< 0.001**	**0.6545**	**< 0.001**	**-0.3912**	**< 0.001**
Water content	**0.4456**	**< 0.001**	**0.2368**	**0.0287**	-0.0623	0.4592
Percentage of embryoless seeds	**0.6269**	**< 0.001**	**0.4324**	**< 0.001**	-0.0668	0.5038
Percentage of viable seeds	**0.4808**	**< 0.001**	**0.2176**	**< 0.001**	-0.0998	0.1337

1 **Table S2.** Summary of phylogenetic mixed models with Bayesian estimation (MCMCglmms) examining the effect of the different abiotic factors, seed mass and dormancy

2 on the final seed germination proportions. Fixed and random effects (±95% CI) and phylogenetic signal λ (±95% CI) are presented. Significant effects are in bold. On

3 average, models were the results of 9000 effect samples.

Abiotic factor / Predictor	Fixed effects			Phylogenetic signal (λ)	Random effects		
	Posterior mean	95% C.I.	pMCMC		Species	Study	Observation:Studies
Light							
All	3.412	(3.215 ; 3.613)	**< 0.001**	0.884 (0.788 – 0.931)	7.11 (3.807 – 10.73)	2.963 (0.456 – 6.542)	2.114 (1.568 – 2.628)
Herb	2.489	(2.216 ; 2.758)	**< 0.001**	0.939 (0.843 – 0.973)	12.99 (4.508 – 24.3)	10.01 (0.361 – 32.32)	2.559 (1.925 – 3.223)
Shrub	4.400	(4.111 ; 4.676)	**< 0.001**	0.859 (0.739 – 0.929)	6.208 (2.731 – 10.2)	0.772 (0 – 2.855)	1.11 (0.53 – 1.71)
Restricted	3.581	(3.226 ; 3.914)	**< 0.001**	0.893 (0.802 – 0.968)	9.931 (3.277 – 18.81)	4.374 (0 – 11.9)	1.508 (0.79 – 2.193)
Widespread	3.318	(3.075 ; 3.56)	**< 0.001**	0.869 (0.761 – 0.934)	6.44 (3.061 – 10.35)	2.509 (0.185 – 7.074)	2.327 (1.745 – 2.948)
Mesic	5.282	(4.906 ; 5.701)	**< 0.001**	0.831 (0.498 – 0.937)	3.956 (0.457 – 9.199)	22.59 (0.348 – 71.42)	1.745 (1.081 – 2.47)
Mesic/Xeric	0.844	(0.518 ; 1.136)	**< 0.001**	0.946 (0.733 – 0.996)	16.55 (1.311 – 50.46)	26.24 (0.033 – 92.11)	0.944 (0.28 – 1.572)
Xeric	3.448	(3.159 ; 3.733)	**< 0.001**	0.893 (0.797 – 0.95)	8.204 (3.61 – 13.74)	3.652 (0.002 – 9.413)	1.829 (1.226 – 2.483)
Dry mass	-0.437	(-0.63 ; -0.256)	**< 0.001**	0.846 (0.739 – 0.924)	5.636 (2.83 – 9.193)	4.472 (0.614 – 10.45)	2.099 (1.551 – 2.638)
25/15 °C							
Overall	-1.384	(-1.852 ; -0.87)	**< 0.001**	0.917 (0.269 – 0.99)	5.582 (0.004 – 18.78)	9.212 (0 – 33.01)	0.088 (0 – 0.321)
Seed mass	-1.259	(-2.337 ; -0.138)	0.140	0.968 (0.48 – 0.999)	14.43 (0.049 – 49.4)	1094 (0 – 1690)	0.064 (0 – 0.245)
30/20 °C							
Overall	-0.379	(-0.828 ; 0.06)	0.090	0.829 (0.491 – 0.983)	4.018 (0.22 – 11.2)	8.492 (0 – 29.25)	0.035 (0 – 0.136)
Herb	0.057	(-0.746 ; 0.908)	0.893	0.994 (0.007 – 1)	58.14 (0 – 185.6)	682.7 (0 – 1462)	0.093 (0 – 0.359)
Shurb	-0.617	(-1.179 ; -0.038)	**0.036**	0.931 (0.621 – 0.997)	13.3 (0.297 – 44.31)	90.82 (0 – 245.7)	0.061 (0 – 0.23)
Seed mass	0.322	(-0.229 ; 0.862)	0.250	0.98 (0.096 – 1)	16.63 (0 – 60.55)	776.3 (0 – 1480)	0.043 (0 – 0.165)
10 °C							
All	-4.295	(-4.96 ; -3.613)	**< 0.001**	0.003 (0 – 0.744)	0.726 (0 – 2.951)	95.73 (0 – 290.8)	0.36 (0 – 1.018)
Restricted	-2.142	(-3.194 ; -1.145)	**< 0.001**	0.996 (0.17 – 1)	183.1 (0 – 640.3)	140 (0 – 567.9)	0.35 (0 – 1.314)
Widespread	-5.505	(-6.393 ; -4.691)	**< 0.001**	0.004 (0 – 0.844)	1.469 (0 – 5.12)	0 (0 – 0)	0.187 (0 – 0.645)
Dry mass	1.242	(0.631 ; 1.843)	**< 0.001**	0.004 (0 – 0.725)	0.686 (0 – 2.645)	208.9 (0 – 767.8)	0.255 (0 – 0.791)
15 °C							
All	-1.919	(-2.176 ; -1.663)	**< 0.001**	0.93 (0.876 – 0.969)	14.12 (6.587 – 22.23)	1.742 (0.062 – 4.906)	0.366 (0 – 0.777)
Herb	-1.119	(-1.475 ; -0.786)	**< 0.001**	0.934 (0.852 – 0.98)	15.2 (3.604 – 31.35)	13.16 (0.09 – 48.44)	0.167 (0 – 0.493)
Shrub	-2.661	(-3.045 ; -2.261)	**< 0.001**	0.94 (0.883 – 0.977)	17.55 (6.621 – 30.81)	1.407 (0 – 5.84)	0.506 (0 – 1.047)
Restricted	-1.808	(-2.267 ; -1.363)	**< 0.001**	0.955 (0.849 – 0.986)	17.2 (3.531 – 38.63)	2.111 (0 – 8.446)	0.216 (0 – 0.652)
Widespread	-1.975	(-2.313 ; -1.667)	**< 0.001**	0.943 (0.87 – 0.972)	15.14 (5.954 – 26.12)	5.065 (0.089 – 16.77)	0.507 (0 – 0.993)

Mesic	-1.612	(-1.988 ; -1.262)	**< 0.001**	0.954 (0.868 – 0.983)	17.18 (4.223 – 34.99)	16.03 (0.006 – 58.79)	0.187 (0 – 0.548)
Mesic/Xeric	-0.683	(-1.25 ; -0.159)	**0.015**	0.973 (0.699 – 0.999)	20.47 (0.114 – 65.64)	28.41 (0.017 – 107)	0.22 (0 – 0.703)
Xeric	-2.877	(-3.346 ; -2.43)	**< 0.001**	0.932 (0.817 – 0.974)	12.25 (3.305 – 24.24)	1.601 (0 – 6.554)	0.583 (0 – 1.223)
Dry mass	-0.023	(-0.309 ; 0.27)	0.886	0.937 (0.872 – 0.968)	14.15 (6.584 – 23.48)	2.484 (0.105 – 7.488)	0.304 (0 – 0.696)
20 °C							
All	-0.019	(-0.195 ; 0.165)	0.838	0.843 (0.719 – 0.904)	4.586 (2.058 – 7.342)	2.565 (0.15 – 6.241)	0.119 (0 – 0.348)
Herb	0.666	(0.395 ; 0.937)	**< 0.001**	0.798 (0.606 – 0.923)	3.953 (0.952 – 7.842)	7.608 (0.084 – 24.08)	0.12 (0 – 0.38)
Shrub	-0.572	(-0.799 ; -0.34)	**< 0.001**	0.902 (0.789 – 0.953)	7.058 (2.502 – 12.73)	2.736 (0 – 8.938)	0.072 (0 – 0.242)
Restricted	0.238	(-0.088 ; 0.551)	0.135	0.905 (0.699 – 0.962)	6.047 (1.354 – 13.35)	3.92 (0 – 10.87)	0.093 (0 – 0.313)
Widespread	-0.148	(-0.378 ; 0.073)	0.196	0.818 (0.684 – 0.91)	4.458 (1.82 – 7.669)	2.979 (0.084 – 8.612)	0.154 (0 – 0.431)
Mesic	0.095	(-0.18 ; 0.363)	0.499	0.809 (0.608 – 0.934)	3.226 (0.803 – 6.923)	6.351 (0 – 20.27)	0.035 (0 – 0.129)
Mesic/Xeric	0.463	(0.038 ; 0.912)	**0.038**	0.786 (0.486 – 0.956)	4.477 (0.352 – 11.2)	6.456 (0 – 21.13)	0.456 (0 – 1.016)
Xeric	-0.386	(-0.679 ; -0.089)	**0.015**	0.885 (0.737 – 0.953)	7.27 (1.926 – 14.07)	3.559 (0 – 10.35)	0.205 (0 – 0.561)
Dry mass	-0.341	(-0.534 ; -0.148)	**0.001**	0.853 (0.741 – 0.916)	5.059 (2.226 – 8.205)	3.05 (0.171 – 7.698)	0.098 (0 – 0.304)
30 °C							
All	-0.528	(-0.71 ; -0.342)	**< 0.001**	0.886 (0.816 – 0.937)	7.243 (3.766 – 11.29)	1.852 (0.207 – 4.696)	0.099 (0 – 0.309)
Herb	0.092	(-0.205 ; 0.357)	0.518	0.685 (0.465 – 0.885)	2.563 (0.542 – 5.404)	6.221 (0.138 – 19.95)	0.167 (0 – 0.476)
Shrub	-1.054	(-1.286 ; -0.819)	**< 0.001**	0.948 (0.893 – 0.971)	12.4 (5.656 – 20.78)	1.224 (0 – 4.99)	0.046 (0 – 0.17)
Restricted	-0.264	(-0.538 ; 0.019)	0.064	0.896 (0.766 – 0.967)	6.223 (1.507 – 12.92)	2.957 (0 – 8.832)	0.03 (0 – 0.117)
Widespread	-0.663	(-0.92 ; -0.418)	0.196	0.904 (0.823 – 0.957)	9.655 (4.027 – 16.34)	3.798 (0.112 – 11.08)	0.223 (0 – 0.563)
Mesic	-0.720	(-0.995 ; -0.435)	**< 0.001**	0.909 (0.802 – 0.971)	8.003 (2.23 – 16.42)	9.546 (0 – 34.04)	0.045 (0 – 0.166)
Mesic/Xeric	0.270	(-0.206 ; 0.747)	0.267	0.746 (0.393 – 0.962)	3.905 (0.263 – 10.9)	9.202 (0.02 – 31.67)	0.599 (0 – 1.24)
Xeric	-0.756	(-1.032 ; -0.482)	**0.015**	8.702 (3.05 – 15.57)	1.852 (0 – 6.214)	1.852 (0 – 6.214)	0.085 (0 – 0.289)
Dry mass	-0.188	(-0.388 ; 0.002)	0.062	0.896 (0.815 – 0.941)	7.612 (3.713 – 12.08)	2.138 (0.235 – 5.423)	0.109 (0 – 0.329)
35 °C							
All	-2.486	(-2.793 ; -2.196)	**< 0.001**	0.876 (0.717 – 0.928)	5.625 (2.135 – 9.439)	4.358 (0.186 – 10.91)	1.547 (0.903 – 2.191)
Herb	-2.153	(-2.626 ; -1.684)	**< 0.001**	0.845 (0.594 – 0.94)	4.69 (0.925 – 9.977)	4.553 (0.067 – 14.7)	2.243 (1.354 – 3.185)
Shrub	-2.725	(-3.114 ; -2.367)	**< 0.001**	0.858 (0.711 – 0.955)	7.014 (1.709 – 13.72)	8.105 (0 – 23.78)	0.912 (0.142 – 1.585)
Restricted	-2.132	(-2.617 ; -1.673)	**< 0.001**	0.887 (0.686 – 0.966)	7.348 (1.292 – 15.73)	2.644 (0 – 8.515)	0.935 (0 – 1.636)
Widespread	-2.660	(-3.04 ; -2.271)	**< 0.001**	0.854 (0.678 – 0.938)	5.621 (1.662 – 10.61)	7.73 (0.086 – 22.35)	1.891 (1.169 – 2.705)
Mesic	-4.019	(-4.537 ; -3.524)	**< 0.001**	0.784 (0.294 – 0.944)	3.024 (0 – 7.521)	31.31 (2.09 – 90.38)	1.012 (0.173 – 1.904)
Mesic/Xeric	-0.929	(-1.425 ; -0.416)	**< 0.001**	0.889 (0.697 – 0.976)	8.994 (1.407 – 21.31)	7.368 (0 – 28.39)	0.817 (0 – 1.512)
Xeric	-2.202	(-2.661 ; -1.738)	**< 0.001**	0.85 (0.576 – 0.948)	4.891 (0.716 – 10.62)	3.076 (0 – 10.06)	1.597 (0.73 – 2.458)
Dry mass	0.455	(0.149 ; 0.75)	**0.003**	0.855 (0.715 – 0.928)	5.918 (2.302 – 10.32)	3.504 (0.184 – 9.553)	1.519 (0.895 – 2.188)
40 °C							
All	-1.249	(-1.784 ; -0.708)	**< 0.001**	0.719 (0.398 – 0.87)	2.332 (0.436 – 5.065)	0 (0 – 0)	1.505 (0.538 – 2.446)

Herb	-1.188	(-1.893 ; -0.454)	**0.002**	0.509 (0.167 – 0.908)	1.937 (0 – 5.676)	0 (0 – 0)	1.224 (0 – 2.392)
Shrub	-1.317	(-2.122 ; -0.524)	**0.001**	0.798 (0.315 – 0.955)	3.256 (0.139 – 9.99)	770.8 (0 – 1331)	1.845 (0.573 – 3.308)
Restricted	-0.915	(-1.487 ; -0.351)	**0.001**	0.731 (0.34 – 0.96)	3.738 (0.138 – 11.52)	0 (0 – 0)	0.263 (0 – 0.785)
Widespread	-1.629	(-2.629 ; -0.628)	**0.002**	0.789 (0.394 – 0.961)	3.837 (0.107 – 10.45)	491.2 (0 – 1288)	3.625 (1.643 – 5.935)
Mesic	-3.678	(-4.699 ; -2.687)	**< 0.001**	0.993 (0.102 – 1)	31.33 (0 – 76.51)	0 (0 – 0)	0.242 (0 – 0.908)
Mesic/Xeric	-1.304	(-2.238 ; -0.368)	**0.005**	0.798 (0.413 – 0.992)	6.474 (0.122 – 20.61)	509.6 (0 – 1281)	2.053 (0.399 – 3.835)
Xeric	-0.085	(-0.804 ; 0.657)	0.809	0.744 (0.346 – 0.974)	3.797 (0.138 – 11.37)	0 (0 – 0)	0.938 (0 – 2)
Dry mass	-1.149	(-1.672 ; -0.605)	**< 0.001**	0.738 (0.436 – 0.902)	2.708 (0.539 – 6.155)	579.9 (0 – 1055)	1.156 (0.277 – 2.051)
Fire-related cues							
All	-0.641	(-0.876 ; -0.388)	**< 0.001**	0.959 (0.887 – 0.982)	19.54 (6.709 – 36.63)	1.062 (0 – 3.823)	0.523 (0 – 0.953)
Combined	-2.143	(-3.46 ; -0.794)	**0.003**	0.992 (0.832 – 0.999)	60.61 (0.658 – 200.7)	0 (0 – 0)	2.608 (0.458 – 5.382)
Single	-0.548	(-0.784 ; -0.314)	**< 0.001**	0.952 (0.887 – 0.98)	18.59 (6.779 – 35.54)	1.021 (0 – 3.556)	0.253 (0 – 0.608)
Heat	-0.831	(-1.176 ; -0.485)	**< 0.001**	0.958 (0.894 – 0.986)	22.35 (6.755 – 43.45)	1.455 (0 – 5.276)	0.496 (0 – 0.992)
Smoke	0.057	(-0.237 ; 0.342)	0.697	15.44 (0.903 – 0.989)	15.44 (3.908 – 30.98)	1.439 (0 – 4.972)	0.023 (0 – 0.087)
Dormant	0.218	(-0.248 ; 0.7)	0.359	0.97 (0.487 – 0.999)	18.91 (0 – 61.13)	39.82 (0 – 151.1)	0.059 (0 – 0.223)
Non-dormant	-0.814	(-1.077 ; -0.52)	**< 0.001**	0.969 (0.901 – 0.992)	26.69 (6.285 – 56.7)	2.055 (0 – 6.22)	0.745 (0 – 1.204)
Seed mass	0.179	(-0.093 ; 0.454)	0.198	0.958 (0.895 – 0.984)	21.79 (7.043 – 41.86)	1.113 (0 – 3.797)	0.603 (0 – 1.053)
Heat shock temperature							
80 °C for 2'	0.179	(-0.543 ; 0.903)	0.616	0.964 (0.896 – 0.989)	18.13 (3.958 – 37.41)	0.88 (0 – 3.235)	0.039 (0 – 0.147)
80 °C for 5'	0.109	(-0.567 ; 0.798)	0.755	0.953 (0.857 – 0.987)	15.32 (2.624 – 32.7)	0.933 (0 – 3.074)	0.05 (0 – 0.186)
100 °C for 1'	-0.545	(-1.039 ; -0.089)	**0.026**	0.963 (0.916 – 0.988)	19.55 (5.07 – 37.53)	0.492 (0 – 1.806)	0.039 (0 – 0.148)
100 °C for 5'	-1.122	(-1.6 ; -0.663)	**< 0.001**	0.957 (0.841 – 0.986)	17.69 (2.814 – 38.09)	2.62 (0.036 – 8.24)	0.222 (0 – 0.629)
200 °C for 1'	-2.485	(-3.096 ; -1.889)	**< 0.001**	0.964 (0.905 – 0.989)	21.26 (5.75 – 41.66)	1.231 (0 – 4.557)	0.094 (0 – 0.335)
Heat shocks							
All	-0.500	(-0.814 ; -0.188)	**0.003**	0.965 (0.904 – 0.986)	22.69 (7.102 – 45.39)	1.13 (0 – 3.766)	0.17 (0 – 0.494)
Herb	-0.373	(-0.864 ; 0.111)	0.134	0.974 (0.815 – 0.997)	21.75 (1.228 – 65.16)	27.61 (0 – 95.44)	0.064 (0 – 0.245)
Shrub	-0.566	(-0.975 ; -0.157)	**0.006**	0.98 (0.932 – 0.996)	42.58 (9.428 – 98.09)	3.572 (0.041 – 11.95)	0.312 (0 – 0.788)
Restricted	-0.645	(-1.151 ; -0.177)	**0.007**	0.925 (0.718 – 0.992)	12.45 (0.952 – 34.1)	4.567 (0 – 12.98)	0.315 (0 – 0.843)
Widespread	-0.366	(-0.789 ; 0.064)	0.088	0.98 (0.929 – 0.995)	40 (7.305 – 92.64)	2.755 (0 – 9.573)	0.126 (0 – 0.416)
Mesic/Xeric	-0.072	(-0.629 ; 0.472)	0.794	0.979 (0.709 – 0.999)	21.32 (0.231 – 69.83)	48.92 (0 – 165.4)	0.045 (0 – 0.173)
Xeric	-0.561	(-0.978 ; -0.148)	**0.007**	0.974 (0.915 – 0.995)	37.06 (5.006 – 84.35)	2.851 (0.001 – 9.76)	0.383 (0 – 0.895)
Dormant	0.256	(-0.328 ; 0.898)	0.406	0.983 (0.392 – 1)	30.67 (0 – 105.5)	91.08 (0 – 304.2)	0.08 (0 – 0.305)
Non-dormant	-0.722	(-1.108 ; -0.345)	**< 0.001**	0.966 (0.914 – 0.993)	33.35 (6.73 – 74.78)	1.502 (0 – 4.826)	0.28 (0 – 0.716)
Dry mass	0.399	(0.054 ; 0.74)	0.020	0.958 (0.91 – 0.988)	25.1 (7.582 – 52.74)	1.321 (0 – 4.325)	0.18 (0 – 0.52)
Smoke							
All	0.057	(-0.237 ; 0.342)	0.697	0.96 (0.903 – 0.989)	15.44 (3.908 – 30.98)	1.439 (0 – 4.972)	0.023 (0 – 0.087)

Herb	0.446	(-0.213 ; 1.117)	0.186	0.988 (0.746 – 1)	35.48 (0.756 – 108.6)	835.2 (0.135 – 1691)	0.081 (0 – 0.315)
Shrub	-0.025	(-0.368 ; 0.308)	0.885	0.975 (0.928 – 0.995)	24.68 (4.67 – 55.35)	2.393 (0 – 8.081)	0.023 (0 – 0.087)
Restricted	0.181	(-0.294 ; 0.662)	0.468	0.924 (0.699 – 0.993)	10.47 (0.717 – 30.87)	12.97 (0 – 40.24)	0.073 (0 – 0.276)
Widespread	-0.021	(-0.398 ; 0.399)	0.922	0.983 (0.928 – 0.998)	34.03 (3.48 – 86.99)	2.347 (0 – 8.289)	0.027 (0 – 0.107)
Mesic/Xeric	0.243	(-0.556 ; 1.058)	0.548	0.997 (0.56 – 1)	97.85 (0.046 – 300.5)	0 (0 – 0)	0.1 (0 – 0.382)
Xeric	0.018	(-0.328 ; 0.339)	0.913	0.973 (0.911 – 0.993)	20.47 (3.877 – 46.81)	2.13 (0 – 6.92)	0.024 (0 – 0.092)
Dormant	0.499	(-0.342 ; 1.37)	0.254	0.995 (0.195 – 1)	114.7 (0 – 331.1)	1085 (0 – 1737)	0.139 (0 – 0.543)
Non-dormant	-0.021	(-0.331 ; 0.299)	0.895	0.972 (0.904 – 0.993)	19.07 (4.138 – 42.55)	1.601 (0 – 5.143)	0.027 (0 – 0.105)
Dry mass	0.000	(-0.317 ; 0.332)	0.996	0.973 (0.915 – 0.992)	19.81 (4.747 – 42.28)	1.802 (0 – 6.374)	0.027 (0 – 0.104)

Table S3. Summary of phylogenetic mixed models with Bayesian estimation (MCMCglmms) examining the effect of the different abiotic factors, seed mass and dormancy

on median germination time (t_{50}). Fixed and random effects ($\pm$95% CI) and phylogenetic signal λ ($\pm$95% CI) are presented. Significant effects are in bold. On average, models

were the results of 9000 effect samples.

Abiotic factor / Predictor	Fixed effects			Phylogenetic signal (λ)	Random effects		
	Posterior mean	95% C.I.	pMCMC		Species	Study	Observation:Studies
25/15 °C							
Overall	6.620	(5.385 ; 7.917)	**0.001**	0.998 (0.993 – 1)	426.1 (85.92 – 976.1)	157.4 (0 – 662.7)	6.288 (3.909 – 9.031)
Seed mass	5.732	(3.162 ; 8.24)	**< 0.001**	0.997 (0.975 – 1)	265.4 (13.37 – 803)	2758 (0 – 5097)	4.239 (2.078 – 6.755)
30/20 °C							
Overall	1.234	(0.524 ; 1.956)	**0.002**	0.993 (0.974 – 0.999)	132.9 (22.82 – 317.9)	104.5 (0 – 415.7)	1.772 (0.76 – 2.842)
Herb	2.241	(1.196 ; 3.298)	**< 0.001**	0.998 (0.852 – 1)	284.8 (1.744 – 844.1)	1672 (0 – 2427)	0.906 (0 – 2.209)
Shurb	0.734	(-0.216 ; 1.679)	0.123	0.996 (0.98 – 1)	235 (19.35 – 644.4)	341.9 (0 – 1163)	2.117 (0.781 – 3.554)
Seed mass	1.003	(0.149 ; 1.862)	**0.027**	0.997 (0.98 – 1)	236.5 (23.78 – 632.5)	3957 (0 – 2924)	1.657 (0.537 – 2.916)
15 °C							
All	5.419	(4.691 ; 6.15)	**< 0.001**	0.99 (0.976 – 0.997)	98.21 (32.69 – 185.4)	19.19 (0 – 61.37)	6.87 (5.289 – 8.482)
Herb	5.125	(4.385 ; 5.875)	**< 0.001**	0.982 (0.949 – 0.994)	140 (22 – 337.4)	253.9 (0.009 – 625.7)	4.116 (2.859 – 5.552)
Shrub	5.903	(4.399 ; 7.479)	**< 0.001**	0.992 (0.973 – 0.998)	116.2 (24.07 – 262.7)	18.35 (0 – 75.27)	11.99 (8.108 – 16.47)
Restricted	7.089	(5.617 ; 8.432)	**< 0.001**	0.995 (0.92 – 1)	114.9 (0 – 357)	51 (0 – 186.6)	7.286 (4.296 – 10.57)
Widespread	4.752	(3.89 ; 5.598)	**< 0.001**	0.992 (0.973 – 0.998)	115.1 (32.65 – 232.5)	28.61 (0 – 114.8)	6.52 (4.81 – 8.425)
Mesic	5.018	(3.351 ; 6.654)	**< 0.001**	0.995 (0.969 – 1)	174.8 (11.05 – 491.2)	98.63 (0 – 388.8)	10.7 (6.57 – 15.07)
Mesic/Xeric	6.509	(5.503 ; 7.565)	**< 0.001**	0.993 (0.941 – 0.999)	106.8 (5.34 – 320.2)	108.5 (0 – 329)	4.348 (2.577 – 6.308)
Xeric	4.674	(3.482 ; 5.873)	**< 0.001**	0.99 (0.94 – 0.999)	92.89 (2.359 – 262.6)	50.7 (0 – 167.2)	6.444 (4.055 – 9.138)
Dry mass	-1.109	(-1.842 ; -0.311)	0.886	0.992 (0.974 – 0.997)	102.4 (27.17 – 206)	30.66 (0 – 99.35)	6.684 (5.052 – 8.398)
20 °C							
All	1.037	(0.512 ; 1.549)	**< 0.001**	0.997 (0.994 – 0.999)	343.7 (171.4 – 542.8)	11.86 (0.001 – 41.13)	6.5 (5.318 – 7.678)
Herb	0.835	(0.279 ; 1.395)	**0.003**	0.982 (0.949 – 0.994)	49.8 (13.3 – 101.3)	51.14 (0.206 – 165.7)	2.709 (1.792 – 3.618)
Shrub	1.200	(0.362 ; 2.029)	**0.005**	0.982 (0.998 – 0.999)	495.2 (192.7 – 855.8)	32.46 (0 – 125.6)	9.707 (7.555 – 11.98)
Restricted	0.796	(-0.067 ; 1.644)	0.066	0.994 (0.985 – 0.998)	165 (56.15 – 314.2)	18.19 (0 – 65.24)	6.619 (4.831 – 8.509)
Widespread	1.187	(0.529 ; 1.86)	**0.002**	0.998 (0.995 – 0.999)	495.3 (170.9 – 892.5)	39.31 (0 – 157.9)	6.446 (5.038 – 7.993)
Mesic	1.185	(0.579 ; 1.82)	**< 0.001**	0.999 (0.998 – 1)	1622 (491.7 – 3199)	181.6 (0.087 – 732.7)	2.188 (1.203 – 3.156)
Mesic/Xeric	1.494	(0.196 ; 2.805)	**0.029**	0.984 (0.92 – 0.998)	52.5 (4.734 – 137)	61.51 (0 – 209.5)	9.704 (6.574 – 12.96)

Herb	-0.410	(-0.869 ; 0.052)	0.082	0.982 (0.949 – 0.994)	43.9 (10.39 – 93.8)	82.06 (0.165 – 287.6)	1.497 (0.834 – 2.277)
Shrub	1.690	(0.515 ; 2.841)	**0.005**	0.995 (0.986 – 0.998)	179.9 (54.14 – 349.5)	37.17 (0 – 137.5)	16.38 (12.76 – 20.3)
Restricted	-0.955	(-2.023 ; 0.132)	0.083	0.993 (0.974 – 0.998)	106.3 (24.7 – 219.6)	27.19 (0 – 92.75)	8.628 (6.18 – 11.32)
Widespread	1.432	(0.635 ; 2.177)	**< 0.001**	0.994 (0.984 – 0.998)	154.8 (48.1 – 291.4)	44.67 (0 – 181.8)	8.959 (7.125 – 10.89)
Mesic	2.930	(1.457 ; 4.459)	**< 0.001**	0.996 (0.975 – 1)	194.4 (16.49 – 519.9)	96.69 (0 – 394.6)	15.14 (10.62 – 20.12)
Mesic/Xeric	-1.394	(-2.409 ; -0.455)	**0.008**	0.962 (0.732 – 0.998)	20.24 (0.168 – 60.27)	23.59 (0 – 95.38)	4.378 (2.775 – 6.149)
Xeric	0.315	(-0.489 ; 1.169)	**0.471**	0.995 (0.987 – 0.998)	180.3 (60.2 – 343.4)	35.32 (0 – 119.1)	6.96 (5.31 – 8.946)
Dry mass	-0.624	(-1.535 ; 0.254)	0.062	0.995 (0.988 – 0.998)	181.4 (74.25 – 324.6)	28.59 (0 – 96.32)	10.05 (8.306 – 11.97)
35 °C							
All	2.151	(1.417 ; 2.943)	**< 0.001**	0.994 (0.985 – 0.998)	148.8 (58.67 – 264)	28.29 (0 – 90.39)	9.471 (7.506 – 11.32)
Herb	2.239	(1.472 ; 2.951)	**< 0.001**	0.982 (0.949 – 0.994)	114.9 (24.03 – 246.9)	54.54 (0.001 – 195.5)	5.052 (3.67 – 6.525)
Shrub	2.027	(0.408 ; 3.629)	**< 0.001**	0.992 (0.966 – 0.997)	94.29 (17.9 – 215.4)	73.75 (0 – 230.6)	16.98 (12.32 – 22.44)
Restricted	0.405	(-0.893 ; 1.686)	0.530	0.997 (0.988 – 0.999)	258.3 (50.35 – 589.7)	54.97 (0 – 213.3)	8.923 (5.986 – 12.15)
Widespread	3.085	(2.121 ; 4.017)	**< 0.001**	0.991 (0.976 – 0.998)	111.7 (26.43 – 230.7)	67.71 (0 – 211)	9.349 (7.114 – 11.57)
Mesic	2.665	(1.06 ; 4.247)	**< 0.001**	0.998 (0.985 – 1)	360.7 (25.81 – 1005)	98.31 (0 – 385.8)	7.729 (4.36 – 11.41)
Mesic/Xeric	1.924	(0.231 ; 3.553)	**0.022**	0.985 (0.859 – 0.999)	47.28 (0.001 – 138.1)	42.77 (0 – 175.6)	15.76 (11.24 – 21.18)
Xeric	2.173	(1.262 ; 3.119)	**< 0.001**	0.988 (0.955 – 0.997)	74.39 (12.88 – 178.8)	93.63 (0 – 296.2)	6.013 (4.089 – 7.908)
Dry mass	0.613	(-0.193 ; 1.48)	0.156	0.993 (0.983 – 0.998)	139.4 (47.69 – 259.6)	41.18 (0 – 125.8)	10.09 (7.994 – 12.24)
Fire-related cues							
All	-0.283	(-0.898 ; 0.296)	0.349	0.994 (0.983 – 0.998)	159.6 (41.53 – 333.9)	28.57 (0.296 – 112)	5.565 (4.357 – 6.777)
Combined	-1.885	(-3.496 ; -0.171)	**0.035**	0.998 (0.981 – 1)	720.4 (0.109 – 2327)	0 (0 – 0)	1.686 (0 – 3.876)
Single	-0.260	(-0.851 ; 0.402)	0.413	0.993 (0.983 – 0.998)	158.2 (41.7 – 327.9)	33.63 (0.288 – 127.9)	5.82 (4.543 – 7.083)
Heat	0.012	(-0.925 ; 0.922)	0.971	0.992 (0.973 – 0.999)	114.6 (19.27 – 250.5)	45.23 (0.317 – 160.5)	7.243 (5.256 – 9.338)
Smoke	-0.974	(-1.67 ; -0.31)	**0.004**	0.997 (0.988 – 0.999)	274.3 (50.7 – 653.4)	76.75 (0.141 – 240)	2.187 (1.189 – 3.27)
Dormant	-1.494	(-3.551 ; 0.586)	0.157	0.996 (0.921 – 1)	207.3 (207.3 – 0)	1348 (0 – 3135)	9.817 (5.092 – 15.53)
Non-dormant	-0.088	(-0.728 ; 0.513)	0.777	0.995 (0.985 – 1)	232.8 (40.24 – 574)	77.03 (0.319 – 259.1)	5.053 (3.916 – 6.274)
Seed mass	-0.608	(-1.239 ; 0.039)	0.063	0.994 (0.981 – 0.999)	160.2 (35.3 – 353.6)	34.6 (0.207 – 131.6)	5.881 (4.627 – 7.205)
Heat shock temperature							
80 °C for 2'	-1.463	(-2.994 ; 0.075)	0.061	0.995 (0.986 – 0.999)	196.5 (53.11 – 415.4)	25.71 (0 – 100.5)	2.91 (1.529 – 4.515)
80 °C for 5'	-0.466	(-1.765 ; 0.91)	0.493	0.996 (0.988 – 0.999)	225.5 (58.69 – 483.5)	36 (0.002 – 141.9)	3.112 (1.724 – 4.683)
100 °C for 5'	0.944	(-0.519 ; 2.229)	0.173	0.993 (0.968 – 0.999)	123.1 (11.64 – 287.8)	10.93 (7.717 – 14.52)	1.046 (0.671 – 1.486)
Heat shocks							
All	-0.066	(-1.029 ; 0.836)	0.893	0.992 (0.972 – 0.998)	110.9 (19.99 – 243)	44 (0.485 – 162.5)	7.255 (5.32 – 9.377)
Herb	-1.184	(-2.448 ; 0.111)	0.070	0.993 (0.965 – 0.999)	130.4 (11.62 – 350.5)	199.8 (0.342 – 603.7)	5.474 (3.118 – 8.125)
Shrub	0.716	(-0.555 ; 2.074)	0.285	0.996 (0.949 – 1)	0.716 (-0.555 – 2.074)	190 (0.001 – 609.1)	66.62 (0 – 221.2)
Restricted	-0.900	(-1.862 ; 0.086)	0.067	0.995 (0.979 – 0.999)	183.8 (23.73 – 463.2)	171 (0.296 – 616.9)	3.893 (2.22 – 5.462)

Widespread	1.017	(-0.756 ; 2.855)	0.272	0.994 (0.153 – 1)	87.79 (0 – 346.7)	41.93 (0.001 – 143)	12.54 (7.75 – 18.28)
Mesic/Xeric	-2.855	(-4.804 ; -1.027)	**0.005**	0.997 (0.934 – 1)	187.6 (0.01 – 607.7)	1163 (0 – 2761)	6.932 (2.923 – 11.87)
Xeric	1.043	(-0.139 ; 2.109)	0.073	0.004 (0 – 0.963)	6.092 (0 – 25.93)	27.71 (0.374 – 96.94)	6.755 (4.596 – 9.351)
Dormant	-1.983	(-3.955 ; 0.006)	0.052	0.996 (0.95 – 1)	218.2 (0 – 689.2)	0 (0 – 0)	7.042 (2.952 – 11.98)
Non-dormant	-0.723	(-1.108 ; -0.345)	**< 0.001**	0.995 (0.97 – 0.999)	33.35 (6.73 – 74.78)	1.502 (0 – 4.826)	0.28 (0 – 0.716)
Dry mass	-0.964	(-1.987 ; -0.033)	0.053	0.993 (0.96 – 0.998)	97.35 (10.7 – 233.3)	51.02 (0.14 – 189.8)	7.646 (5.45 – 9.844)
Smoke							
All	-0.974	(-1.67 ; -0.31)	**0.004**	0.997 (0.988 – 0.999)	274.3 (50.7 – 653.4)	76.75 (0.141 – 240)	2.187 (1.189 – 3.27)
Herb	-1.560	(-3.548 ; 0.411)	0.120	0.999 (0.976 – 1)	600.5 (6.639 – 1777)	1176 (0 – 1618)	7.226 (3.348 – 12.26)
Shrub	-0.785	(-1.294 ; -0.298)	**< 0.001**	0.998 (0.99 – 1)	414.3 (42.7 – 1113)	58.16 (0.183 – 207)	0.19 (0 – 0.575)
Restricted	-1.447	(-2.409 ; -0.417)	**0.007**	0.997 (0.988 – 1)	359.8 (44.06 – 950.3)	699.9 (0 – 1312)	3.595 (1.971 – 5.405)
Widespread	-0.071	(-0.655 ; 0.528)	0.819	0.999 (0.461 – 1)	907.6 (0.001 – 1348)	160.5 (0.272 – 243.6)	0.051 (0 – 0.196)
Non-dormant	-1.090	(-1.665 ; -0.541)	**< 0.001**	0.997 (0.988 – 0.999)	295.1 (50.39 – 711.9)	65.08 (0.299 – 225.1)	1.026 (0.248 – 1.858)
Dry mass	-0.131	(-0.848 ; 0.622)	0.724	0.997 (0.984 – 0.999)	265.1 (37.71 – 699.9)	86.51 (0.11 – 231.4)	2.349 (1.276 – 3.48)

1