Type of article: Research in Context

Seed functional ecology in Brazilian rock outcrop vegetation: an integrative synthesis

Carlos A. Ordóñez-Parra[1,2*], Natália F. Medeiros[2,3], Roberta L.C. Dayrell[4], Soizig Le Stradic[5],

Daniel Negreiros[6], Tatiana Cornelissen[2], Fernando A. O. Silveira[2]

1. Programa de Pós-Graduação em Biologia Vegetal, Departamento de Botânica, Instituto de

Ciências Biológicas, Universidade Federal de Minas Gerais, Belo Horizonte, Brasil.

2. Centro de Síntese Ecológica e Conservação, Departamento de Genética, Ecologia e

Evolução, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Belo

Horizonte, Brasil.

3. Programa de Pós-Graduação em Ecologia, Conservação e Manejo da Vida Silvestre,

Departamento de Genética, Ecologia e Evolução, Instituto de Ciências Biológicas,

Universidade Federal de Minas Gerais, Belo Horizonte, Brazil.

4. Faculty of Biology and Preclinical Medicine, University of Regensburg, Regensburg,

Germany.

5. UMR BIOGECO, Institut National de la Recherche Agronomique – INRAE and

Université de Bordeaux, Bordeaux, France.

6. Laboratório de Ecologia Evolutiva e Biodiversidade, Departamento de Genética, Ecologia

e Evolução, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Belo

Horizonte, Brazil

* Corresponding author. E-mail: carlos.ordonez.parra@gmail.com

Running Title: Seed ecology in Brazilian rock outcrop vegetation

ABSTRACT

Background and Aims

Rock outcrop vegetation is distributed worldwide and hosts a diverse, specialised, and unique

flora that evolved under extremely harsh environmental conditions. The germination ecology

in such ecosystems has received little attention, especially regarding the association between

seed traits, germination responses and adult plants ecology. Here, we provide a quantitative

and meta-analytical review of the seed functional ecology of Brazilian rocky outcrop

vegetation, focusing on four vegetation types: *campo rupestre*, *canga*, *campo de altitude* and

inselbergs.

Methods

Using a database with functional trait data for 383 taxa and 10,187 germination records for

281 taxa, we calculated the phylogenetic signal of seven seed traits and tested whether they

varied among growth forms, geographical distributions, and microhabitats. We also

conducted a meta-analysis to understand the effects of light, temperature, and fire-related cues

on the germination of *campo rupestre* species and how the beforementioned ecological groups

and seed mass affect such responses.

Key Results

All traits showed a strong phylogenetic signal. *Campo rupestre* species responded positively

to light and had optimal germination between 20-30ºC. The effect of temperatures below and

above this range was modulated by growth form, with shrubs requiring and tolerating higher

temperatures to germinate. We only found evidence of a moderating effect of seed mass for

responses to heat shocks, with larger, dormant seeds tolerating heat better. Heat shocks above

200°C killed seeds, but smoke accelerated germination. No consistent differences in germination responses were found between restricted and widespread species or microhabitats. Still, species from xeric habitats evolved phenological strategies to synchronise germination with higher soil water availability.

Conclusions

Evolutionary history plays a major role in the seed ecology of Brazilian rock outcrop vegetation. However, seed traits and germination responses did not explain species' geographic distribution and microhabitats, suggesting other traits are more likely to explain such differences.

Keywords

INTRODUCTION

Rock outcrops are outstanding geological features where the bedrock protrudes above the land's surface due to the erosion of softer parts of the landscape (Fitzsimons and Michael 2017). They offer a unique habitat that drastically contrasts with the neighbouring vegetation (Porembski 2007). Notably, they experience severe surface temperatures and have extremely nutrient-poor soils with low water retention capacity, a combination of abiotic factors that has driven the evolution of distinctive traits that allow plants to establish and survive in such harsh environments (Kluge and Brulfert 2000; Escudero *et al.* 2015; Oliveira *et al.* 2016). Most attention, however, has been given to the ecophysiology of adult plants, overlooking the potential role of regeneration traits and regeneration niche, two aspects that are expected to drive species distribution and community assembly (Grubb 1977; Donohue *et al.* 2010; Larson and Funk 2016). Up to date, only a few qualitative reviews of germination ecology in these ecosystems have been conducted (Wyatt 1997; Biedinger *et al.* 2000).

In Brazil, there are four main vegetation types associated with rock outcrops: *campo rupestre*, *canga*, *campo de altitude* and inselbergs (Martinelli 2007). *Campo rupestre* occurs on quartzite, sandstone, and ironstone outcrops throughout the country, but it is most common at the high elevations of the Espinhaço Range in eastern Brazil (Miola *et al.* 2021). The *campo rupestre* that develops on ironstone outcrops receives the name *canga* (Jacobi *et al.* 2007; Skirycz *et al.* 2014). The *campo de altitude* is found on granite and gneissic outcrops within the Atlantic Forest (Vasconcelos 2011), whereas inselbergs are isolated, dome-shaped granitic outcrops present across the country (Safford and Martinelli 2000). All these open, grassy-shrubby, fire-prone vegetations are centres of species richness, phylogenetic diversity and endemism (Porembski 2007; Silveira *et al.* 2016; Campos *et al.* 2018).

The germination ecology of Brazilian rock outcrop vegetations has been previously

reviewed (Garcia and Oliveira 2007; Nunes *et al.* 2016; Garcia *et al.* 2020), but these

syntheses were restricted to the *campo rupestre* and focused on the differences in germination

responses among some of its most emblematic families (i.e., Asteraceae, Bromeliaceae,

Cactaceae, Eriocaulaceae, Melastomataceae, Velloziaceae and Xyridaceae). Still, these

syntheses put forward promising hypotheses about the role of seed and germination traits in

the ecological dynamics of these plant communities by pointing out that 1) seed traits shape

germination responses to abiotic factors and 2) that contrasting microhabitat preferences and

geographical ranges derive from distinct germination requirements, as assumed under the

regeneration niche hypothesis and the seed ecological spectrum framework (Grubb 1977;

Saatkamp *et al.* 2019). Nevertheless, they have only been assessed with a handful of species

from a single family or could not be formally tested without large-scale databases made

available only recently (Ordóñez-Parra *et al.* 2023). Furthermore, these qualitative reviews do

not explicitly address the potential role of shared evolutionary history in species seed ecology.

Studies carried out both at the local and global scale have highlighted that several key seed

traits are highly conserved across the phylogeny, including seed mass (Moles *et al.* 2005;

Zanetti *et al.* 2020), dormancy (Willis *et al.* 2014; Dayrell *et al.* 2017) and germination

requirements (Arène *et al.* 2017). As a result, a proper assessment of the ecological

hypotheses raised by these reviews must incorporate this phylogenetic component.

In this research, we test these two long-standing hypotheses using a phylogenetically-

controlled approach with the ultimate goal of providing an integrative synthesis of seed

functional ecology in Brazilian rock outcrop vegetation. First, we assessed whether seed

functional traits associated with different dimensions of the seed ecological spectrum are

phylogenetically conserved and evaluated whether they differed between growth-forms,

species geographic distribution and microhabitats. We then assessed the effect of light,

temperature and fire-related cues on the germination of plant species in these communities. These abiotic factors are established germination cues in these ecosystems and are the most studied cues in Brazilian rock outcrop vegetation, allowing us to provide more robust conclusions about their effects on germination (Figure S1). We then contrasted germination responses between the before mentioned ecological groups and tested if such responses were moderated by seed mass –a central trait that is considered to control various aspects of germination (Saatkamp *et al.* 2019). Finally, given that seed traits data provide critical information for biodiversity conservation and ecological restoration (Mattana *et al.* 2022) and rock outcrop vegetation is increasingly exposed to several threats –both in Brazil and worldwide (Fernandes 2016; Porembski *et al.* 2016; Fitzsimons and Michael 2017)– we discuss how the available information on seed traits and germination responses support and inform conservation and restoration programs. Since rock outcrops occur in all continents and biomes, and their associated vegetation is quite similar in structure and function (Barthlott and Porembski 2000), we hope this study provides a solid starting point towards a global synthesis of the germination ecology of rock outcrop vegetation.

MATERIALS AND METHODS

Data sources

Seed functional traits and germination experiments

We retrieved seed trait and germination records from *Rock n' Seeds* (Ordóñez-Parra *et al.* 2023), a database of seed functional traits and germination experiments of plant species from Brazilian rock outcrop vegetation (Figure 1). We focused on seven functional traits associated with the investment in regeneration via seeds and seeds main functions (Grubb 1977;

Saatkamp *et al.* 2019) [i.e., seed dry mass (mg), seed water content (%), percentage of

embryoless seeds (%), percentage of viable seeds (%), seed dispersal syndrome

(anemochoroy, autochory and zoochory), seed dispersal season (early-dry, late-dry, early-rain

and late-rain) and primary dormancy (dormant and non-dormant)]; and experiments assessing

the effects of light availability, constant and alternate temperatures, and fire-related cues (i.e.,

heat shocks and smoke). Beyond seed and germination traits, the database also provides

information on species growth form, microhabitat ('xeric' when substrate experiences

pronounced seasonal drought, or otherwise 'mesic'), and geographic distribution ('restricted'

to rock outcrop vegetation, and otherwise 'widespread') of each species (see Ordóñez-Parra *et*

al. 2023 for a full description of data collection).

Phylogenetic tree

We reconstructed the phylogenetic tree of the study species using the package *V.PhyloMaker2*

(Jin and Qian 2022). First, species names were checked and updated following The Leipzig

Catalogue of Vascular Plants using the R package *lcvplants* (Freiberg *et al.* 2020). We kept

taxa identified at the species level, and all subspecies and varieties (18 cases) were upgraded

to the species level. A phylogenetic tree for the resulting 370 species was generated, based on

the GBOTB phylogeny (Smith and Brown 2018) updated and standardised following Freiberg

et al. (2020). Taxa absent from this phylogeny were bound to their designated relatives using

the bind.relative function of *V.PhyloMaker2* based on different sources (Almeda *et al.* 2016;

Rivera *et al.* 2016). Species with no known relatives in the phylogeny (*Austrocritonia*

velutina, Cavalcantia glomerata, Cavalcantia percymosa and *Parapiqueria cavalcantei*) were

added using the at.node function from the *ape* package (Paradis and Schliep 2019). All these

species belong to the Eupatorieae (Asteraceae), a tribe where infrageneric relationships are yet

to be resolved (Rivera *et al.* 2016). Therefore, they were added to the base of the tribe as

polytomies.

Most infrageneric relationships in the phylogeny remained unresolved, appearing as

polytomies, partially because the relationships within some genera of the highly diverse

families in our database have low support (Alcantara *et al.* 2018; Guimarães *et al.* 2019).

Nevertheless, phylogenetic metrics estimated from trees resolved to the genus-level have

shown to be highly correlated with those derived from fully-resolved trees, suggesting that

these polytomies do not impair the results (Qian and Jin 2021).

Statistical analyses

All analyses were made using R v. 4.2.0 (R Core Team 2022), and the code prepared will be

provided as Supplementary Material and uploaded to GitHub upon acceptance (see Open

Data). Results are presented using a gradual language of evidence, which allows

communicating scientific findings in a more nuanced way (Muff *et al.* 2022).

Variation and phylogenetic signal of seed functional traits

To test for the phylogenetic signal in the quantitative seed traits (i.e., dry mass, water content,

percentage of viable and embryoless seeds), we calculated Pagel's λ (Pagel 1999) using the

phylosig function from the package *phytools* (Revell 2012). λ values range from zero to one,

with $\lambda = 0$ indicating that related taxa are not more similar than expected by chance (i.e., no

phylogenetic signal) and $\lambda = 1$ implying that a given trait evolves under a Brownian motion

model (Pagel 1999). Pagel's λ was selected among the available phylogenetic signal indices

given its robustness in measuring phylogenetic signals based on incompletely resolved

phylogenies (Molina-Venegas and Rodríguez 2017). We further explored this phylogenetic

signal by calculating Moran's I at the Genus, Family and Order levels using the

correlogram.formula function from *ape* (Paradis and Schliep 2019). Values of Moran's I range from -1 to 1, with I = 0 indicating that species resemble each other as expected by Brownian motion model. Values above and below zero imply that related species are more or less similar than expected by such model, respectively (Moran 1950; Gittleman and Kot 1990). All these tests were carried out using log-transformed seed mass values and logit-transformed water content, and percentage of embryoless and viable seed values, to reduce the skewness of the data. For species with more than one record, we carried out the analyses using the mean species value.

For the qualitative seed traits (i.e., seed dormancy, seed dispersal syndrome and seed dispersal season), the phylogenetic signal was tested by implementing the rtestdeciv function from the *adiv* package using 9,999 permutations (Pavoine 2020). This method decomposes trait diversity among the nodes of the phylogenetic tree and assesses whether it is skewed towards the tree's root or tips, with significant root skewness implying the presence of a significant phylogenetic signal. Seed dormancy and dispersal season were treated as multichoice variables as 11 species had records of both dormant and non-dormant seeds, and 18 species reported more than one dispersal season.

To test for differences in quantitative seed traits between ecological groups (growth-form, microhabitat, and geographic distribution), we used phylogenetic generalised least square models as implemented in the package *caper* (Orme *et al.* 2018). Likewise, variation in qualitative seed traits were assessed using a phylogenetic logistic regression (Ives and Garland 2010) implemented in the *phylolm* package (Ho and Ané 2014). Because the dispersal season and syndrome had more than two possible states, individual models were run to assess the probability of each season and syndrome. Species with records for more than one distribution records were classified as "Widespread" and species with occurrence in both microhabitats were classified as "Mesic/Xeric". For this analysis, species with records from

more than one combination of qualitative traits were included as different populations, using

the add.taxa.phylo function from the *daee* package (Debastiani 2021) and keeping the tree

ultrametric.

Preliminary tests showed that the interaction between our qualitative predictors did not

significantly affect any of the tested variables. Moreover, the model that included such were

always outperformed by simple additive models based on Akaike Information Criteria (see

code provided). As a result, our final models did not include these interactions. Growth-form

comparisons were made between herbs and shrubs due to the low sample size of trees (26

species), succulents (nine species) and lianas (six species).

Meta-analyses of germination responses to abiotic factors

We estimated the effects of light, constant and temperatures, and fire-related cues on

germination only for *campo rupestre* species, given the lack of data from *campo de altitude*

and the small sample size of independent observations for inselbergs (two studies, three

species) for these abiotic factors. We used the standardised mean difference Hedge's g

(Hedges 1981) using the escalc function from the package *metafor* (Viechtbauer 2010). This

metric measures the standardized change in a given response variable in treatment groups

compared to control groups and includes a correction by sample size and variance

(Rosenberg *et al.* 2013). Experiments under total darkness were used as controls to test the

effects of light, with results from different temperatures being pooled into a single effect for

each plant species in each study. Given the overall positive effect of light on germination (see

below), only experiments carried out under light conditions were used to assess the effects of

different temperatures and fire-related cues on seed germination. We used 25ºC as a control

for constant and alternate temperatures since it is considered the optimal temperature for most

species in this ecosystem (Garcia and Oliveira 2007; Nunes *et al.* 2016; Garcia *et al.* 2020).

Finally, we used untreated seeds (i.e., seeds not exposed to either heat shocks or smoke) as controls for fire-related cues.

We assessed the effect of these abiotic factors on seed germination percentage and median germination time (t_{50}). Mean germination percentage values were taken from the raw germination data deposited in *Rock n' Seeds* database and were complemented with data from thirteen experiments recovered from the database literature survey whose raw data was not already available. To be included in our meta-analysis, the latter studies should present the mean germination percentage and its standard deviation (or any other statistical measure that could be converted into standard deviation units) in either the text, tables figures or supplementary materials or provide enough information for calculations. Data presented in figures were extracted using ImageJ (Schneider *et al.* 2012). To avoid confounding effects from treatments with extremely low germination, we only included observations where either the control or treatment had ≥10% germination. The t_{50} (Coolbear *et al.* 1984 as modified by Farooq *et al.* 2005) was calculated for the experiments deposited in the database *Rock n' Seeds* that include raw daily germination data using the *germinationmetrics* package (Aravind *et al.* 2022). To reduce the effect of low germinability in our analysis, we calculated t_{50} for observations where both the control and treatment conditions had ≥10% germination. Moreover, we assessed the effect of light, constant and alternate temperatures as germination cues in non-dormant seeds or in seeds where dormancy has been alleviated. Alternate temperatures are known to break physical dormancy (Baskin and Baskin 2014), but none of the studies in the dataset have tested this. In contrast, since fire-related cues are known to shape seed dormancy states (Baskin and Baskin 2014), we included dormant species in the meta-analysis of fire-related cues. Species with non-conclusive records regarding seed dormancy were excluded.

We assessed the overall effect of each abiotic factor (i.e. light, temperature and fire)

on germination percentage and time by implementing mixed effect models using the rma.mv

function of *metafor* (Viechtbauer 2010), with effect sizes as response variables and

observations, study and species phylogeny as random variables. Species appearing in more

than one study were included in the phylogeny as distinct populations using the

add.taxa.phylo function from *daee* (Debastiani 2021) with a branch length of zero (Lajeunesse

et al. 2013). A covariance matrix for each species set was built using the vcv function from

ape (Paradis and Schliep 2019). After assessing the overall effect of each abiotic factor, we

tested the moderating effects of growth-forms, distribution, microhabitats, seed mass and

dormancy, but the last one only for fire-related cues. Because seed mass exhibited a

significant, strong phylogenetic signal (see below), the missing seed mass values for some 54

species were inputted using average values for the genera from *Rock n' Seeds* database

(Ordóñez-Parra *et al.* 2023) and the Seed Information Database (https://data.kew.org/sid/).

Ninety-five percent confidence intervals around the effect sizes were calculated. Effect size

was considered significant if the confidence intervals did not overlap zero (Hedges *et al.*

1999). Comparisons within predictors were carried out using the function linearHypothesis

from the *car* package (Fox and Weisberg 2019).

RESULTS

Variation and phylogenetic signal of seed functional traits

We found very strong evidence that all seed traits exhibited a significant phylogenetic signal

(Table 1, $p < 0.001$). For all quantitative traits, the phylogenetic signal was moderate to strong

(λ between 0.57-0.90), with closely related taxa strongly resembling each other at both the

Genus and Family level (Table S1). At the Order level, we only found (very strong) evidence

of a negative phylogenetic correlation for the seed mass (I = -0.39, p < 0.001, Table S1).

Seed mass values ranged from 0.00035 to 175.15 mg (Median: 0.245 mg). Among the

most well-represented families in the database, Fabaceae exhibited the heaviest seeds

(Median: 11.57 mg), whereas Melastomataceae (0.007 mg), Xyridaceae (0.016 mg) and

Eriocaulaceae (0.044 mg) exhibited the lighter ones (Figure 2). There was moderate evidence

that shrubs (t = 2.19, p = 0.030) and species occurring at both mesic and xeric microhabitats (t

= 2.01, p = 0.045) produced heavier seeds than herbs and species from mesic microhabitats,

respectively (Table 2).

Seed water content was less variable than seed mass, with values ranging from 3.9% to

28.3% (Median: 11.7%). Cyperaceae (7.7%), Asteraceae (9.2%) and Fabaceae (9.4%) had the

seeds with the lowest water content, whereas Melastomataceae (12.2%), Bromeliaceae

(13.6%) and Velloziaceae (14.2%) showed the highest values (Figure 2). There was no

evidence that seed water content varied between growth forms, microhabitats or species

distribution (Table 2).

The percentage of embryoless and viable seeds varied greatly among species. On the

one hand, the percentage of embryoless seeds ranged from 0% (i.e., species producing only

filled seeds) to species producing > 95% of empty seeds (Median: 10.6%). Poaceae and

Asteraceae produced the highest proportion of embryoless seeds (Median: 82% and 66%,

respectively), while Eriocaulaceae (0.9%), Xyridaceae (1.5%) and Fabaceae (4%) produced

few embryoless seeds (Figure 2). Likewise, some species produce large amounts of unviable

seeds while others produce mainly viable seeds (i.e., almost 100% of viable seeds).

Eriocaulaceae (Median: 89%), Xyridaceae (87.7%) and Velloziaceae (86%) produced the

highest proportion of viable seeds, while Poaceae species (11.5%) produced the seeds with

the lowest viability (Figure 2). We found no evidence that the percentage of embryoless and

viable seeds varied between growth forms, microhabitats nor species distributions (Table 2).

Sixty-four percent of species produced non-dormant seeds, whereas 36% produced

dormant seeds (Figure 2). The most common dormancy class was physical dormancy (PY),

being present in 37 Fabaceae, three Convolvulaceae and one Malvaceae species.

Physiological dormancy (PD) was the second common dormancy class and Verbenaceae

(nine species), Cyperaceae (six species) and Melastomataceae (five species) were the families

with the highest number of species presenting it. Eleven species were reported to produce

either mostly dormant or non-dormant seeds. We found moderate evidence that species from

xeric environments had higher probabilities of producing dormant seeds ($z = 2.52$, $p = 0.012$).

Autochory was the most common dispersal syndrome (58.9% of species), which was

present in families such as Melastomataceae (49 species), Fabaceae (38) and Velloziaceae

(36). This dispersal syndrome was followed by anemochory (26.2%) –exhibited by

Asteraceae (29 species), Eriocaulaceae (24) and Bromeliaceae (24)– and zoochory (14.9%,

e.g., Melastomataceae). There was no evidence that the probability of dispersal syndrome was

driven by any of the predictors tested (Table 3). Most species dispersed seeds during the dry

season, either in the late-dry season (38.5%) or early-dry season (26.5%), while only 25.6%

and 17.9% dispersed in the late- and early-rainy season, respectively. There was strong

evidence that shrubs had a higher probability of dispersing seeds during the late-dry season (z

$= 3.43$, $p < 0.001$), and moderate evidence that dispersal during the late rainy season was

more likely in herbs ($z = -2.01$, $p = 0.044$). Additionally, the data exhibited moderate evidence

that species from mesic/xeric microhabitats ($z = 2.03$, $p = 0.042$) and those restricted to xeric

microhabitats ($z = 2.15$, $p = 0.031$) had a higher probability of dispersing seeds during the

late-rain season than species from mesic microhabitats. There was no evidence that the

probability of dispersal during the early-dry or the early-rainy season varied between

predictors (Table 3).

In summary, shrubs tended to produce heavier seeds and had a high probability of

dispersal during the late-dry season. In contrast, herbs produced relatively smaller seeds that

had high probabilities of being dispersed during the late-rainy season. Species from

mesic/xeric microhabitats tended to produce heavy seeds and be dispersed during the late-

rainy season compared to those from exclusively mesic microhabitats. Similarly, species

restricted to xeric environments had a higher probability of producing dormant seeds and

presenting late-rainy dispersal than those from mesic environments (Figure 3). Finally, there

was no evidence that species restricted to outcrop vegetation and widespread ones differed in

any seed trait.

Meta-analyses of germination responses

There was very strong evidence that light had a positive overall effect on germination

percentage (g = 5.48, CI = 3.03 to 7.93, z = 4.39, p < 0.001) regardless of growth-form,

distribution, or microhabitat (Figure 4). The effect was higher in herbs and restricted species

when compared to shrubs (χ^2 = 6.04, p = 0.014) and widespread species (X^2 = 9.33, p =

0.002), respectively. There was no evidence of differences in germination responses to light

between microhabitats or that seed mass moderated the effect of light on germination

percentage (z = -0.71, p = 0.48).

Regarding the effect of constant temperatures on germination, there was no evidence

that 20 and 30 ºC differed from the control treatment (25 ºC), regardless of growth form,

distribution, or microhabitat (Figure 5A). Still, the data showed very strong evidence that

temperatures below 20 ºC (10 and 15 °C) reduced germination percentage (10 ºC: g = -12.53,

CI = -17.62 to -7.45, z = -4.83, p < 0.0001; 15ºC: g = -1.75, CI = -2.78 to -0.72, z = -3.32, p <

0.001) and increased germination time (15ºC: g = 5.09, CI = 3.37 to 6.81, z = 5.80, p <

0.0001). Moreover, there was moderate-to-strong evidence that temperatures above 30 ºC (35

and 40°C) decreased germination percentage (35 ºC: g = -4.57, CI = -9.00 to -0.14, z = -2.02,

p = 0.04; 40ºC: g = -7.85, CI = -12.94 to -2.75, z = -3.02, p = 0.002) but no evidence that they

affected germination time (Figure 5A). Growth-form moderated germination responses to

temperature, with our data showing very strong evidence that shrubs exhibited the most

pronounced decreases in germination percentage at 15ºC (X^2 = 8.35, p = 0.004, Figure 5).

Additionally, there was moderate evidence that germination in herbs was accelerated at 30ºC

(z = -2.23, p = 0.03) but decreased (z = -2.12, p = 0.03) and delayed (z = 2.03, p = 0.04) at

35ºC. No evidence of an effect of these temperatures was found germination of shrubs. Seed

mass modulated germination responses at 20 and 30 ºC, with moderate evidence that

relatively heavier seeds had low germination at 20 ºC (z = -2.20, p = 0.03), but strong

evidence that they had higher germination percentage at 30 °C (z = 2.90, p = 0.003). No

consistent differences were found between microhabitats or distributions.

15   Similar results were found when evaluating the effects of alternate temperatures, with

the data showing moderate evidence that 25/15 ºC regimes reduced germination percentage (g

= -2.58, CI = -4.94 to -0.20, z = -2.13, p = 0.03), but very strong evidence that they increased

germination time (g = 2.72, CI = 1.80 to 3.65, z = 5.78, p < 0.001,). Conversely, there was no

evidence that 30/20 ºC affected germination percentage (g = -0.93, CI = -4.22 to 2.36, z = -

0.66, p = 0.51) or time (g = 0.83, CI = -0.42 to 2.06, z = 1.31, p = 0.19). Moreover, there was

no evidence that seed mass modulated the effect of alternate temperatures on germination

percentage or time (Figure 5B).

23   Finally, we found moderate evidence that fire-related cues reduced germination

24   percentage (g = -0.41, CI = -0.81 to -0.01, z = -2.01, p = 0.04) and time (g = -0.41, CI = -0.92

25   to 0.10, z = -2.68, p = 0.007). This effect varied according to the treatment applied, with heat

shocks –alone or in combination with smoke– reducing germination percentage but not germination time. The data showed moderate evidence that the germination percentage of relatively heavier seeds increased after exposed to fire-related ($z = 2.54$, $p = 0.01$), while no evidence of an effect on germination time was found ($z = 1.41$, $p = 0.16$). Moreover, there was strong evidence that primary dormancy significantly shaped the responses to fire-related cues, with an overall decrease in germination percentage in ND species ($z = -2.75$, $p = 0.006$, Fig 6A). When comparing heat shock treatments, there was only evidence of a negative effect on germination percentage by 200 ºC, 1 minute heat shocks ($z = -5.56$, $p < 0.001$, Figure 6B). Further analysis excluding this treatment showed no evidence that heat shocks affected germination, regardless of growth form, distribution or microhabitat (Figure 6C). Still, there was moderate evidence that ND ($z = -2.16$, $p = 0.03$) and relatively lighter seeds were more sensitive to heat ($z = 2.05$, $p = 0.04$). The data showed no evidence that smoke affected germination percentage, but moderate evidence that it accelerated germination ($g = -1.28$, CI $= -2.51$ to -0.06, $z = -2.05$, $p = 0.04$). However, the available data only provided evidence for such an effect in herbs ($g = -1.57$, CI $= -3.13$ to -0.02, $z = -1.9860$, $p = 0.047$) and restricted species ($g = -1.62$, CI $= -2.87$ to -0.38, $z = -2.55$, $p = 0.01$). There was no evidence that smoke's effect was modulated neither by seed mass nor by dormancy (Figure 6D).

DISCUSSION

Seeds functional traits are phylogenetically conserved in Brazilian rock outcrop vegetation

We found that all the seed functional traits we assessed had a significant and moderate-to-strong phylogenetic signal, suggesting that phylogenetic relatedness plays a pivotal role in shaping seed traits in rock outcrop vegetation as globally reported for seed mass (Moles *et al.* 2005) and dormancy (Willis *et al.* 2014). Such phylogenetic signal could

be traced back up to the Family level, implying that several aspects of a species germination ecology can be inferred from other members of the family. However, our results partially contrast with local studies in other Brazilian open ecosystems. First, Zanetti *et al.* (2020) did not find a phylogenetic signal for seed water content in their study of 48 species from the *cangas* of Carajás. Still, the authors tested and measured the phylogenetic signal of this trait using Bloomberg's K, an index known to be highly sensitive to sample sizes and incomplete phylogenetic background, and that provides divergent results to Pagel's λ (Münkemüller *et al.* 2012). Second, we found a significant phylogenetic signal for seed dispersal season, while seed dispersal season in the Cerrado (Escobar *et al.* 2021) and other phenophases in the *campo rupestre* (Zanetti *et al.* 2020; CS Oliveira *et al.* 2021) have not shown such signal. As a result, seed dispersal season is the only phenophase that has shown a significant phylogenetic signal in the Brazilian rock outcrop vegetation, potentially as a result of an intense evolutionary pressure towards germination timing in these ecosystems, where water is even more scarce than in the Cerrado (CS Oliveira *et al.* 2021).

Seed mass and seed dispersal season differed between herbs and shrubs, with shrubs tending to produce relatively larger seeds and having higher probabilities of dispersing during the late-dry season, whereas herbs produced lighter seeds with higher probabilities of dispersal during the late-rain season. Differences in seed mass between herbs and shrubs have been attributed to the distinct strategies employed by such growth forms: while shrubs invest in fewer, larger seeds that cope better with environmental hazards, herbs invest in numerous, lighter seeds to increase their establishment opportunities (Westoby *et al.* 2002; Moles *et al.* 2005). This relationship is also supported by the global spectrum of plant form and function (Díaz *et al.* 2016), suggesting that this global pattern also stands for Brazilian rock outcrop vegetation. On the other hand, differences in seed dispersal season suggest differences in phenological strategies to cope with precipitation seasonality (see below).

Seed traits do not explain large-scale species distribution, but seed dispersal phenology may

shape microhabitat differences

Rock outcrops and the surrounding vegetation differ in several aspects that should

impact germination, such as nutrient and water availability, irradiance, and daily temperature

fluctuations (Oliveira *et al.* 2016). Despite that, we did not find differences in seed functional

traits or germination requirements between restricted and widespread species, suggesting that

the assessed seed traits do not explain large-scale species distribution, as predicted by the

regeneration niche hypothesis, which suggests that differences in regeneration traits

contribute to species ecological breadth and niche segregation (Grubb 1977). Therefore,

community assembly at this scale is probably shaped by vegetative traits that allow these

plants to cope with the stressful conditions of rock outcrops (Oliveira *et al.* 2016) or by other

seed and germination traits not included here.

Species from mesic and xeric microhabitats did differ in some of the traits we

assessed, but these differences were related to seed dispersal phenology rather than seed

germination requirements. This finding agrees with previous research showing little

correspondence between seed traits and species ecological preferences (Fernández-Pascual *et*

al. 2022) but contrasts with local-scale studies showing small-scale differences in the

germination niche of *campo rupestre* species from distinct microhabitats (Oliveira and Garcia

2011; Ranieri *et al.* 2012; Silveira, Negreiros, *et al.* 2012; Marques *et al.* 2014; Giorni *et al.*

2018). All these local studies were conducted with fewer species from a single family,

suggesting that these differences might dilute at larger scales when several different taxa are

considered. While our results do not support the idea that germination responses to the abiotic

factors explain distributions at the microhabitat level in the *campo rupestre*, a more detailed

analysis within each family could be useful to test the relative importance of germination

requirements for microhabitat differentiation across different scales.

Regardless of microhabitat, most species dispersed their seeds during the late- or the

early-dry season. Therefore, dispersal during the dry season is the prevalent strategy in

Brazilian rock outcrop vegetation. However, we found that species from xeric habitats had

higher probabilities of producing dormant seeds and dispersing their seeds during the late-rain

season –two strategies that presumably arise due to evolutionary pressures towards strategies

to synchronize germination with higher water availability (Silveira, Ribeiro, *et al.* 2012;

Garcia *et al.* 2020). Yet, only 36% of species in our dataset produce dormant seeds, and

campo rupestre is the vegetation type with the highest ND/D ratio worldwide (Dayrell *et al.*

2017), implying that seed dormancy is not the main driver of seedling establishment in rock

outcrop vegetation. Additional strategies to control germination timing might include

germination requirements and the acquisition of secondary dormancy. A mismatch between

temperature germination requirements and environmental conditions at the moment of

dispersal might provides an alternative strategy to prevent germination in the absence of

dormancy (Escobar *et al.* 2021) –for example, seeds dispersed during the dry season have

evolved to germinate under relatively higher temperatures of the rainy season. On the other

hand, secondary dormancy has been reported in various Eriocaulaceae and Xyridaceae species

from *campo rupestre* (see Garcia *et al.* 2020). All these species disperse their non-dormant

seeds between the early-dry and the early-rain season, which become increasingly dormant as

the rainy season advances, and humidity and temperature rise. When the dry season starts,

dormancy is progressively alleviated by reduced temperatures and low water availability

(Duarte and Garcia 2015).

Campo rupestre *species depend on light for germination*

Our meta-analysis showed that light positively affects seed germination across all

ecological groups, supporting previous assessments about its importance in *campo rupestre*

germination ecology. Interestingly, restricted species had a stronger response to light than widespread species, suggesting this kind of response may be particularly prevalent in this ecosystem. We also found that germination responses to light were stronger in herbs, which could be associated to our finding of them having lower seed mass compared to shrubs, as predicted by the well-established trade-off between light requirements and seed mass (Milberg *et al.* 2000). Under such trade-off, lighter seeds are expected have stronger light requirements, given the little amount of internal reserves to support seedling development. Still, our meta-analysis did not find evidence to support such a trade-off in Brazilian rock outcrop vegetation. A possible explanation for this lack of support is the high uniformity in germination responses to light in our species which either responded positively to light (i.e., positive photoblastic) or were light-indifferent (non-photoblastic) (Nunes *et al.* 2016; Garcia *et al.* 2020). Negative photoblastism is prevalent in dark and large-seeded herbaceous species from non-tropical latitudes (Carta *et al.* 2017); thus, it is not expected in Brazilian rock outcrop vegetation. For instance, this germination behaviour has only been reported for one species in our study system (*Lippia filifolia,* Pimenta *et al.* 2007). Another possible explanation is the small variation in seed size in our data, with most species producing small (86% records, < 1 mg) seeds and from lineages known to have positively photoblastic seeds, such as Melastomataceae, Xyridaceae and Velloziaceae (Nunes *et al.* 2016; Garcia *et al.* 2020; Ordóñez-Parra *et al.* 2022). Large-seeded species from *campo rupestre*, mainly from Fabaceae, have shown to be light-indifferent (Nunes *et al.* 2016), so additional studies with other large-seeded species are needed to test the prevalence of the seed mass-light responses trade-off in rock outcrop vegetation.

Small-seeded species from our study are expected to have narrowly defined microsite requirements for successful establishment due to their limited internal resources (Pearson *et al.* 2003). Therefore, futher aspects of the light environment, such as light spectral quality and

the interaction between light and temperature, and are expected to control their germination

(Pons 2014). Spectral quality —measured as the ratio between red (R) and far-red (FR) light;

i.e., R:FR— provides information about overhead foliage and litter on the soil surface, with

high R:FR indicating no cover and, thus, high irradiance (Baskin and Baskin 2014). Still, in

our study system, the effect of different R:FR has only been assessed for a handful

Bromeliaceae (Pereira *et al.* 2009; Hmeljevski *et al.* 2014) and Velloziaceae species (Vieira *et*

al. 2018), preventing us from providing robust inferences about the functional relevance of

light quality as a germination cue in Brazilian rock outcrop vegetation. Globally, responses to

R:FR have been both positively and negatively associated with seed mass (Tiansawat and

Dalling 2013). Jankowska-Blaszczuk and Daws (2007) suggest that tiny seeds should restrict

germination to high R:FR conditions to ensure high irradiance and that for such behaviour to

be successful, seeds must be able to persist for extended periods in the soil. Considering that

the small-seed species from Brazilian rock outcrop vegetation have been shown to persist for

several years in soil seed banks (see Garcia *et al.* 2020), a positive relationship between seed

mass and R:FR requirement would be expected. However, Jankowska-Blaszcuk and Daws

(2007) comment that such germination behaviour implies that small-seeded species would

germinate in high-risky environments where soils dry rapidly. In this case, one should expect

that small-seeded species limit their germination to safer sites, such as those with low R:FR

and likely higher soil moisture. Consequently, a positive relationship between seed mass and

R:FR requirements (Pearson *et al.* 2003) could be adaptive in rock outcrop vegetation, a

hypothesis that is yet to be tested. On the other hand, temperature-dependent germination

responses to light have only been described in some Velloziaceae species in which

germination in the darkness only occurs under high (30-40ºC) temperatures. Still, its

functional or evolutionary relevance, if any, is yet to be described (Garcia *et al.* 2020).

Herbs and shrubs respond differently to low and high temperatures

We found that the optimal temperature for *campo rupestre* species germination ranges

between 20-30ºC, supporting previous studies (Nunes *et al.* 2016; Garcia *et al.* 2020).

Temperatures below this range significantly reduced germination percentage and delayed

germination. This reduction is an arguably a mechanism to avoid germination during the dry

season when temperatures decrease and water is not readily available for seedling

establishment (Garcia *et al.* 2020). The negative effect of low temperatures was stronger in

shrubs, suggesting that these require higher temperatures to germinate. Once again,

differences between growth forms could arise from differences in seed size, with the

relatively larger seeds of shrubs having higher base temperatures for germination (Arène *et al.*

2017). Still, our data only support a positive moderating effect of seed mass on germination

responses to 20 and 30 ºC, both temperatures where no significant differences between herbs

and shrubs were found. Also, the small variation in seed size mentioned earlier and the

limited availability of data from hydrothermal time models our study system (but see Duarte

et al. 2018; Oliveira *et al.* 2021) limit our capacity to test this hypothesis. On the other hand,

temperatures above 30ºC decreased and delayed germination in herbs, while shrubs remained

unaffected, implying that shrub seeds tolerate germinating at higher temperatures.

Accordingly, we should expect increased shrub abundance in microsites that experience

higher temperatures.

Alternate temperature regimes have either a negative or null effect on the germination

in *campo rupestre*. The reduction of germination percentage and the increase of germination

time at 25/15ºC regime is probably due to part of these temperatures being below the optimal

range. Similarly to germination responses to light, the lack of a significant effect of 30/20ºC

and an association between response to alternate temperatures and seed mass could be

attributed to the fact that most *campo rupestre* species produce small, lighter seeds, which are

not expected to benefit from alternate temperature regimes (Pearson *et al.* 2003). Recent

studies in the *campo de altitude* have shown that several species benefit from alternate

temperature regimes, suggesting this cue could be more important in this vegetation (Andrade

et al. 2021). Therefore, additional studies with species with large seeds and from other

Brazilian rock outcrop vegetation are required to elucidate the functional relevance of

alternate temperatures and whether this varies between vegetation types.

Heat kills small, non-dormant seeds, but smoke accelerates germination

Seeds exposed to 200 ºC had their germination percentage reduced, but milder heat

shocks ($\leq$ 100 ºC) did not affect germination percentage or time, agreeing with previous

evidence of high heat tolerance in seeds from Neotropical savanna species (Daibes *et al.*

2022). This effect was moderated by seed mass and seed dormancy, with lighter, non-dormant

seeds being less tolerant to heat shocks. These results are aligned with those described for

Cerrado species (Ramos *et al.* 2016; Daibes *et al.* 2019). Contrastingly, dormant species were

mostly unaffected by heat shocks, supporting the notion that fire-mediated dormancy

alleviation is not expected in rocky areas or savannas (Pausas and Lamont 2022). On the other

hand, the lack of negative effects on germination in dormant species implies that traits

associated with seed dormancy promote heat tolerance. For instance, dormancy acquisition

occurs in parallel with the accumulation of heat shock proteins or the consolidation of water-

impermeable coats, which can offer protection against environmental hazards, including heat

(Tweddle *et al.* 2003; Ramos *et al.* 2016).

Smoke did not affect germination percentage but consistently reduced germination

time. This result agrees with the role of smoke-derived compounds as germination stimulants

of non-dormant seeds or those where dormancy has been alleviated (Mackenzie *et al.* 2021).

Smoke-stimulated germination in the *campo rupestre* is thought to be ecologically relevant as

it promotes the germination of species resprouting and shedding seeds after a fire (Fernandes

et al. 2021) –a usual phenological syndrome in our study system (Figueira *et al.* 2016)–

allowing recently-dispersed seeds to exploit the post-fire environment where competition is

relaxe (Le Stradic *et al.* 2015; Fernandes *et al.* 2021). While our meta-analysis largely

supports this hypothesis, the effect of smoke has only been tested on a handful of species and

further studies are needed to rigorously test it.

Implications for seed banking and seed-based restoration

Seed banking is a promising, low-cost *ex situ* conservation strategy with a high

potential for efficiently conserving considerable amounts of plant material and its associated

genetic diversity (Hay and Probert 2013). Easy-to-measure traits can be a reliable proxy for

assessing whether the seeds of a given species are desiccation tolerant and, thus, suitable for

seed banking. Dry seed mass is tightly linked with this trait, with lighter seeds having more

probabilities of producing desiccation tolerant (Wyse and Dickie 2018). For instance, seeds

with both a thousand seed wight < 500g and a moisture content < 30% are assumed to be

tolerate desiccation (Lan *et al.* 2014). Trait data for 87 species in our dataset suggest that there

is a high potential for storage in *ex-situ* seed banks, as shown by empirical studies in Brazilian

rock outcrop vegetation (Tarré *et al.* 2007; Andrade *et al.* 2021) and similar ecosystems in

Western Africa (Godefroid *et al.* 2013, 2020).

On the other hand, seed-based restoration is increasingly recognized as a highly cost-

effective restoration strategy (Merritt and Dixon 2011), yet many of the seed traits assessed in

this research indicate that it may be challenging to apply in Brazilian rock outcrop vegetation

(Dayrell *et al.* 2016). For example, direct seeding has yielded mixed results in *cangas*: while

it led to a high seedling density during the initial months, species showed high mortality and

slow growth after the first year of seeding (Figueiredo *et al.* 2021b), even after the

incorporation of organic matter, which hampered seedling establishment, especially for species with small, light-dependent seeds species (Figueiredo *et al.* 2021a).

One potential limitation for direct seeding is the prevalence of species that produce many embryoless and non-viable seeds (Figueiredo *et al.* 2021b). Since these traits showed a strong phylogenetic signal in our study, we expect that its relatively easy to identify taxa that require additional processing before broadcasting, such as Poaceae and Asteraceae. Seed-based restoration projects involving these species must employ additional processing techniques to improve seed quality (e.g., Feitosa *et al.* 2009; Melo *et al.* 2009). Restoration practitioners must also consider that the *ex-situ* stored seeds of several species from these ecosystems lose viability quite rapidly after collection, implying they ought to be used as soon as possible (Zanetti *et al.* 2020). Despite all these limitations, direct seeding has an underused potential to help reintroduce species in a cheaper way than seedling planting (Figueiredo *et al.* 2021b).

Alternative seed-based restoration techniques for rock outcrop vegetation include hay transfer and topsoil translocation. Unfortunately, hay transfer is not a useful means to restore degraded areas of *campo rupestre* due to the high amount of non-viable seeds produced by most plants (Le Stradic *et al.* 2014). In contrast, topsoil transfer is a more promising technique to restore rock outcrop vegetation considering that species produce small, light-demanding, and long-lived seeds, suggesting a large potential for persistence in soil (Long *et al.* 2015). Still, topsoil translocation has proven useful for restoration purposes in some (Burke 2008) but not all vegetation associated with rock outcrops (Le Stradic *et al.* 2016). Moreover, studies in the *campo rupestre* have shown impoverished soil seed banks (Medina and Fernandes 2007; Le Stradic *et al.* 2018; Luz *et al.* 2018), hence topsoil transfer (5 cm) has shown little use for ecological restoration (Le Stradic *et al.* 2018). However, transferring deeper topsoil samples in degraded *cangas* showed more promising results with restored areas

recovering a considerable percentage of floristic and functional composition (Onésimo *et al.* 2021; Rezende *et al.* 2021). However, this technique has two major limitations. First, areas where topsoil is removed show little recovery after extraction (Le Stradic *et al.* 2018); thus, topsoil extraction is recommended exclusively for sites targeted for mining operations. Second, soil samples –especially thicker ones– might contain a considerable amount of exotic and non-target species (Le Stradic *et al.* 2018; Rezende *et al.* 2021).

Conclusion

Seed germination ecology in Brazilian rock outcrop vegetation is shaped by species evolutionary history, but we were also able to find differences between ecological groups. Unexpectedly, seed traits did not explain large-scale species distribution, neither germination requirements explained microhabitat preferences, providing low support to the regeneration niche hypothesis (Grubb 1977). As a result, large-scale community assembly might be better explained by vegetative traits or other seed germinaton traits not explored here. Still, species from xeric habitats evolved specific risk-reducing strategies exemplified by seed dormancy and late-rain dispersal, ensuring germination is synchronized with optimum conditions for seedling establishment.

Germination in *campo rupestre* was positively influenced by light, and most species had optimal germination between 20-30 ºC. Temperatures below and above this range had different effects according to growth form, with shrubs requiring higher temperatures than herbs. Therefore, we should expect shrubs to colonize microsites that experience higher soil temperatures better, with likely changes in vegetation physiognomy. Larger, dormant seeds showed higher heat tolerance, but seed mass did not affect germination responses to light or temperatures. Smoke accelerated germination, providing opportunities for establishment under conditions of relaxed competition in post-fire environments.

ACKNOWLEDGEMENTS

This study is part of the first author's Master Dissertation at the Plant Biology Program at Universidade Federal de Minas Gerais. CAO-P and NFM were supported by a scholarship from CAPES and CNPq, respectively. FAOS acknowledges support from FAPEMIG. We also thank James Dalling and Sergey Rosbakh, who provided valuable comments to an earlier version of the manuscript.

OPEN DATA

The R code for the analysis and creation of the figures will be provided as Supplementary Material and uploaded to GitHub upon acceptance.

LITERATURE CITED

Alcantara S, Ree RH, Mello-Silva R. **2018**. Accelerated diversification and functional trait evolution in Velloziaceae reveal new insights into the origins of the campos rupestres' exceptional floristic richness. *Annals of Botany* **122**: 165–180.

Almeda F, Michelangeli FA, Viana PL. **2016**. *Brasilianthus* (Melastomataceae), a new monotypic genus endemic to ironstone outcrops in the Brazilian Amazon. *Phytotaxa* **273**: 269–282.

Andrade LG, Sánchez-Tapia A, Andrade ACS. **2021**. Germination, viability and dormancy of 47 species from threatened tropical montane grassland in southeast Brazil: Implications for ex situ conservation. *Plant Biology* **23**: 735–742.

Aravind J, Devi SV, Radhamani J, Jacob SR, Srinivasan K. **2022**. germinationmetrics: Seed Germination Indices and Curve Fitting. R package version 0.1.7, https://cran.r-project.org/web/packages/germinationmetrics/

Arène F, Affre L, Doxa A, Saatkamp A. **2017**. Temperature but not moisture response of

germination shows phylogenetic constraints while both interact with seed mass and lifespan.

Seed Science Research **27**: 110–120.

Barthlott W, Porembski S. **2000**. Why Study Inselbergs? In: Porembski S, Barthlott W, eds.

Inselbergs. Berlin: Springer, 1–6.

Baskin CC, Baskin JM. **2014**. *Seeds: Ecology, Biogeography, and Evolution of Dormancy*

and Germination. San Diego: Elsevier Inc.

Biedinger N, Porembski S, Barthlott W. **2000**. Vascular Plants on Inselbergs: Vegetative

and Reproductive Strategies In: Porembski S, Barthlott W, eds. *Inselbergs*. Berlin,

Heidelberg: Springer, 117–142.

Burke A. **2008**. The effect of topsoil treatment on the recovery of rocky plain and outcrop

plant communities in Namibia. *Journal of Arid Environments* **72**: 1531–1536.

Campitelli E. **2022**. ggnewscale: Multiple Fill and Colour Scales in "ggplot2." https://cran.r-

project.org/package=ggnewscale

Campos P V., Villa PM, Nunes JA, Schaefer CEGR, Porembski S, Neri AV. 2018. Plant

diversity and community structure of Brazilian Páramos. *Journal of Mountain Science* **15**:

1186–1198.

Carta A, Skourti E, Mattana E, Vandelook F, Thanos CA. **2017**. Photoinhibition of seed

germination: occurrence, ecology and phylogeny. *Seed Science Research* **27**: 131–153.

Coolbear P, Francis A, Grierson D. **1984**. The Effect of Low Temperature Pre-Sowing

Treatment on the Germination Performance and Membrane Integrity of Artificially Aged

Tomato Seeds. *Journal of Experimental Botany* **35**: 1609–1617.

Daibes LF, Ordóñez-Parra CA, Dayrell RLC, Silveira FAO. **2022**. Regeneration from

seeds in South American savannas, in particular the Brazilian Cerrado In: Baskin CC, Baskin

JM, eds. *Plant Regeneration from Seeds*. A Global Warming Perspective. San Diego:

Academic Press, 183–197.

Daibes LF, Pausas JG, Bonani N, Nunes J, Silveira FAO, Fidelis A. **2019**. Fire and legume germination in a tropical savanna: Ecological and historical factors. *Annals of Botany* **123**: 1219–1229.

Dayrell RLC, Arruda AJ, Buisson E, Silveira FAO. **2016**. Overcoming challenges on using native seeds for restoration of megadiverse resource-poor environments: a reply to Madsen et al. *Restoration Ecology* **24**: 710–713.

Dayrell RLC, Garcia QS, Negreiros D, Baskin CC, Baskin JM, Silveira FAO. **2017**. Phylogeny strongly drives seed dormancy and quality in a climatically buffered hotspot for plant endemism. *Annals of Botany* **119**: 267–277.

Debastiani VJ. **2021**. daee: Data Analysis for Ecology and Evolution. version 0.1.7. https://github.com/vanderleidebastiani/daee (accessed November 2021)

Díaz S, Kattge J, Cornelissen JHC, *et al.* 2016. The global spectrum of plant form and function. *Nature* **529**: 167–171.

Donohue K, Rubio De Casas R, Burghardt L, Kovach K, Willis CG. **2010**. Germination, postgermination adaptation, and species ecological ranges. *Annual Review of Ecology, Evolution, and Systematics* **41**: 293–319.

Duarte DM, Garcia QS. **2015**. Interactions between substrate temperature and humidity in signalling cyclical dormancy in seeds of two perennial tropical species. *Seed Science Research* **25**: 170–178.

Duarte AA, Lemos Filho JP, Marques AR. **2018**. Seed germination of bromeliad species from the *campo rupestre*: thermal time requirements and response under predicted climate-change scenarios. *Flora* **238**: 119–128.

Dusa A. **2022**. venn: Draw Venn Diagrams. R Package Version 1.11. https://CRAN.R project.org/package=venn (accessed November 2021)

Escobar DFE, Rubio de Casas R, Morellato LPC. **2021**. Many roads to success: different

combinations of life-history traits provide accurate germination timing in seasonally dry

environments. *Oikos* **130**: 1865–1879.

Escudero A, Palacio S, Maestre FT, Luzuriaga AL. **2015**. Plant life on gypsum: A review

of its multiple facets. *Biological Reviews* **90**: 1–18.

Farooq M, Basra SMA, Ahmad N, Hafeez K. **2005**. Thermal Hardening: A New Seed

Vigor Enhancement Tool in Rice. *Journal of Integrative Plant Biology* **47**: 187–193.

Feitosa SS, Davide AC, Tonetti OAO, Fabricante JR, Lui JJ. **2009**. Estudos de viabilidade

de sementes de candeia *Eremanthus erythropappus* (DC.) MacLeish por meio de testes de

germinação e raios x. *Floresta* **39**: 393–399.

Fernandes GW. **2016**. The Shady Future of the Rupestrian Grassland: Major Threats to

Conservation and Challenges in the Anthropocene In: Fernandes GW, ed. *Ecology and*

Conservation of Mountaintop grasslands in Brazil. Cham: Springer International Publishing,

545–561.

Fernandes AF, Oki Y, Fernandes GW, Moreira B. **2021**. The effect of fire on seed

germination of campo rupestre species in the South American Cerrado. *Plant Ecology* **222**:

45–55.

Fernández-Pascual E, Vaz M, Morais B, *et al.* 2022. Seed ecology of European mesic

meadows. *Annals of Botany* **129**: 121–134.

Figueira JEC, Ribeiro KT, Ribeiro MC, *et al.* 2016. Fire in Rupestrian Grasslands: Plant

Response and Management In: Fernandes GW, ed. *Ecology and Conservation of Mountaintop*

grasslands in Brazil. Cham: Springer International Publishing, 415–448.

Figueiredo MA, Messias MCTB, Leite MGP, Kozovits AR. **2021a**. Seed covering and dry

periods in the rainy season interfere with direct seeding success in the restoration of post-

mined grasslands. *Ecologia Austral* **31**: 444–455.

Figueiredo MA, Messias MCTB, Leite MGP, Kozovits AR. **2021b**. Direct seeding in the

restoration of post-mined campo rupestre: Germination and establishment of 14 native

species. *Flora* **276–277**: 151772.

Fitzsimons JA, Michael DR. **2017**. Rocky outcrops: A hard road in the conservation of

critical habitats. *Biological Conservation* **211**: 36–44.

Fox J, Weisberg S. **2019**. *An R Companion to Applied Regression*. Thousand Oaks, CA:

Sage.

Freiberg M, Winter M, Gentile A, *et al.* 2020. LCVP, The Leipzig catalogue of vascular

plants, a new taxonomic reference list for all known vascular plants. *Scientific Data* **7**: 1–7.

Garcia QS, Barreto LC, Bicalho EM. **2020**. Environmental factors driving seed dormancy

and germination in tropical ecosystems: A perspective from *campo rupestre* species.

Environmental and Experimental Botany **178**: 104164.

Garcia QS, Oliveira PG. **2007**. Germination patterns and seed longevity of monocotyledons

from the Brazilian campos rupestres. *Seed Science and Biotechnology* **1**: 35–41.

Giorni VT, Bicalho EM, Garcia QS. **2018**. Seed germination of *Xyris* spp. from Brazilian

campo rupestre is not associated to geographic distribution and microhabitat. *Flora* **238**: 102–

109.

Gittleman JL, Kot M. **1990**. Adaptation: Statistics and a Null Model for Estimating

Phylogenetic Effects. *Systematic Zoology* **39**: 227.

Godefroid S, Van de Vyver A, Boisson S, Mahy G. **2020**. Seed desiccation-tolerance is a

common feature of threatened taxa in metalliferous tropical grasslands from southeastern DR

Congo. *Journal for Nature Conservation* **56**: 125842.

Godefroid S, Van de Vyver A, Lebrun J, *et al.* 2013. Germination capacity and seed storage

behaviour of threatened metallophytes from the Katanga copper belt (D.R. Congo):

Implications for ex situ conservation. *Plant Ecology and Evolution* **146**: 183–192.

Grubb PJ. **1977**. The Maintenance of Species-Richness in Plant Communities: The

Importance of the Regeneration Niche. *Biological Reviews* **52**: 107–145.

Guimarães PJF, Michelangeli FA, Sosa K, Santiago Gómez JR. **2019**. Systematics of

Tibouchina and allies (Melastomataceae: Melastomateae): A new taxonomic classification.

Taxon **68**: 937–1002.

Hay FR, Probert RJ. **2013**. Advances in seed conservation of wild plant species: a review of

recent research. *Conservation Physiology* **1**: cot030.

Hedges L V. **1981**. Distribution Theory for Glass's Estimator of Effect size and Related

Estimators. *Journal of Educational Statistics* **6**: 107–128.

Hedges L V., Gurevitch J, Curtis PS. **1999**. The meta-analysis of response ratios in

experimental ecology. *Ecology* **80**: 1150–1156.

Hmeljevski K V., Freitas L, Domingues R, *et al*. 2014. Conservation assessment of an

extremely restricted bromeliad highlights the need for population-based conservation on

granitic inselbergs of the Brazilian Atlantic Forest. *Flora* **209**: 250–259.

Ho LST, Ané C. **2014**. A linear-time algorithm for gaussian and non-gaussian trait evolution

models. *Systematic Biology* **63**: 397–408.

Ives AR, Garland T. **2010**. Phylogenetic logistic regression for binary dependent variables.

Systematic Biology **59**: 9–26.

Jacobi CM, Carmo FF, Vincent RC, Stehmann JR. **2007**. Plant communities on ironstone

outcrops: a diverse and endangered Brazilian ecosystem. *Biodiversity and Conservation* **16**:

2185–2200.

Jankowska-Blaszczuk M, Daws MI. **2007**. Impact of red: far-red ratios on germination of

temperate forest herbs in relation to shade tolerance, seed mass and persistence in the soil.

Functional Ecology **21**: 1055–1062.

Jin Y, Qian H. **2022**. V.PhyloMaker2: An updated and enlarged R package that can generate

very large phylogenies for vascular plants. *Plant Diversity* **44**: 335–339.

Kluge M, Brulfert J. **2000**. Ecophysiology of Vascular Plants on Inselbergs In: *Inselbergs*.

Berlin, Heidelberg: Springer, 143–174.

Lajeunesse MJ, Rosenberg MS, Jennions MD. **2013**. Phylogenetic Nonindependence and

Meta-analysis In: Koricheva J, Gurevitch J, Mengersen K, eds. *Handbook of Meta-Analysis in*

Ecology and Evolution. Princeton, New Jersey: Princeton University Press, 284–299.

Lan Q, Xia K, Wang X, Liu J, Zhao J, Tan Y. **2014**. Seed storage behaviour of 101 woody

species from the tropical rainforest of southern China: A test of the seed-coat ratio-seed mass

(SCR-SM) model for determination of desiccation sensitivity. *Australian Journal of Botany*

62: 305–311.

Larson JE, Funk JL. **2016**. Regeneration: an overlooked aspect of trait-based plant

community assembly models. *Journal of Ecology* **104**: 1284–1298.

Le Stradic S, Buisson E, Fernandes GW. **2014**. Restoration of Neotropical grasslands

degraded by quarrying using hay transfer. *Applied Vegetation Science* **17**: 482–492.

Le Stradic S, Fernandes GW, Buisson E. **2018**. No recovery of campo rupestre grasslands

after gravel extraction: implications for conservation and restoration. *Restoration Ecology* **26**:

S151–S159.

Le Stradic S, Séleck M, Lebrun J, *et al.* 2016. Comparison of translocation methods to

conserve metallophyte communities in the Southeastern D.R. Congo. *Environmental Science*

and Pollution Research **23**: 13681–13692.

Le Stradic S, Silveira FAO, Buisson E, Cazelles K, Carvalho V, Fernandes GW. **2015**.

Diversity of germination strategies and seed dormancy in herbaceous species of campo

rupestre grasslands. *Austral Ecology* **40**: 537–546.

Long RL, Gorecki MJ, Renton M, *et al.* 2015. The ecophysiology of seed persistence: A

mechanistic view of the journey to germination or demise. *Biological Reviews* **90**: 31–59.

Luz GR, Mota GS, Spadeto C, Tolentino GS, Fernandes GW, Nunes YRF. **2018**. Regenerative potential of the soil seed bank along an elevation gradient of rupestrian grassland in southeastern Brazil. *Botany* **96**: 281–298.

Mackenzie BDE, Auld TD, Keith DA, Ooi MKJ. **2021**. Fire Seasonality, Seasonal Temperature Cues, Dormancy Cycling, and Moisture Availability Mediate Post-fire Germination of Species With Physiological Dormancy. *Frontiers in Plant Science* **12**: 795711.

Marques AR, Atman APFF, Silveira FAOO, Lemos-Filho JP. **2014**. Are seed germination and ecological breadth associated? Testing the regeneration niche hypothesis with bromeliads in a heterogeneous neotropical montane vegetation. *Plant Ecology* **215**: 517–529.

Martinelli G. **2007**. Mountain biodiversity in Brazil. *Revista Brasileira de Botânica* **30**: 587–597.

Mattana E, Ulian T, Pritchard HW. **2022**. Seeds as natural capital. *Trends in Plant Science* **27**: 139–146.

Mauri M, Elli T, Caviglia G, Uboldi G, Azzi M. **2017**. RAWGraphs: A Visualisation Platform to Create Open Outputs In: *Proceedings of the 12th Biannual Conference on Italian SIGCHI Chapter*. New York, NY, USA: ACM, 1–5.

Medina BMO, Fernandes GW. **2007**. The potential of natural regeneration of rocky outcrop vegetation on rupestrian field soils in "Serra do Cipó", Brazil. *Revista Brasileira de Botânica* **30**: 665–678.

Melo PRB, Oliveira JA, Carvalho MLM, Guimarães RM, Carvalho BO. **2009**. Aplicação do teste de raios x no estudo da morfologia interna e da qualidade fisiológica de aquênios de arnica (*Lychnophora pinaster* Mart.). *Revista Brasileira de Sementes* **31**: 146–154.

Merritt DJ, Dixon KW. **2011**. Restoration Seed Banks—A Matter of Scale. *Science* **332**: 424–425.

Milberg P, Andersson L, Thompson K. 2000. Large-seeded species are less dependent on light for germination than small-seeded ones. *Seed Science Research* **10**: 99–104.

Miola DTB, Ramos VDV, Silveira FAO. 2021. A brief history of research in *campo rupestre*: Identifying research priorities and revisiting the geographical distribution of an ancient, widespread Neotropical biome. *Biological Journal of the Linnean Society* **133**: 464–480.

Moles AT, Ackerly DD, Webb CO, *et al.* 2005. Factors that shape seed mass evolution. *Proceedings of the National Academy of Sciences of the United States of America* **102**: 10540–10544.

Molina-Venegas R, Rodríguez MÁ. 2017. Revisiting phylogenetic signal; strong or negligible impacts of polytomies and branch length information? *BMC Evolutionary Biology* **17**: 53.

Moran PAP. 1950. Notes on Continuous Stochastic Phenomena. *Biometrika* **37**: 17.

Muff S, Nilsen EB, O'Hara RB, Nater CR. 2022. Rewriting results sections in the language of evidence. *Trends in Ecology and Evolution* **37**: 203–210.

Münkemüller T, Lavergne S, Bzeznik B, *et al.* 2012. How to measure and test phylogenetic signal. *Methods in Ecology and Evolution* **3**: 743–756.

Nunes FP, Dayrell RLC, Silveira FAO, *et al.* 2016. Seed Germination Ecology in Rupestrian Grasslands In: Fernandes GW, ed. *Ecology and Conservation of Mountaintop Grasslands in Brazil*. Cham: Springer International Publishing, 207–225.

Oliveira RS, Abrahão A, Pereira C, *et al.* 2016. Ecophysiology of Campos Rupestres Plants In: *Ecology and Conservation of Mountaintop grasslands in Brazil*. Cham: Springer International Publishing, 227–272.

Oliveira TGS, Duarte AA, Diamantino IP, Garcia QS. 2021. Thermal niche for seed germination of *Xyris* species from Brazilian montane vegetation: Implications for climate

change. *Plant Species Biology* **36**: 284–294.

Oliveira RS, Galvão HC, Campos MCR, *et al.* 2015. Mineral nutrition of *campos rupestres* plant species on contrasting nutrient-impoverished soil types. *New Phytologist* **205**: 1183–1194.

Oliveira, R.S., Abrahão, A., Pereira, C., Teodoro, G.S., Brum, M., Alcantara, S. & Lambers, H. 2016. Ecophysiology of Campos Rupestres Plants. In: Fernandes GW, ed. *Ecology and Conservation of Mountaintop grasslands in Brazil*. Cham: Springer International Publishing, 227–272.

Oliveira PG, Garcia QS. 2011. Germination characteristics of *Syngonanthus* seeds (Eriocaulaceae) in *campos rupestres* vegetation in south-eastern Brazil. *Seed Science Research* **21**: 39–45.

Oliveira CS, Messeder JVS, Teixido AL, Arantes MRR, Silveira FAO. 2021. Vegetative and reproductive phenology in a tropical grassland–savanna–forest gradient. *Journal of Vegetation Science* **32**: 1–16.

Onésimo CMG, Dias DD, Vale Beirão M, Kozovits AR, Messias MCTB. 2021. Ecological succession in areas degraded by bauxite mining indicates successful use of topsoil. *Restoration Ecology* **29**: 1–11.

Ordóñez-Parra CA, Messeder JVS, Mancipe-Murillo C, Calderón-Hernández M, Silveira FAO. 2022. Seed Germination Ecology in Neotropical Melastomataceae: Past, Present, and Future In: Goldenberg R, Michelangeli FA, Almeda F, eds. *Systematics, Evolution, and Ecology of Melastomataceae*. Cham: Springer International Publishing, 707–733.

Ordóñez-Parra CA, Dayrell RLC, Negreiros D, *et al.* 2023. Rock n' Seeds: A database of seed functional traits and germination experiments from Brazilian rock outcrop vegetation. *Ecology* **104**: e3852.

Orme D, Freckleton R, Thomas G, *et al.* **2018**. caper: Comparative Analyses of Phylogenetics and Evolution in R. https://cran.r-project.org/web/packages/caper/index.html

Pagel M. **1999**. Inferring the historical patterns of biological evolution. *Nature* **401**: 877–884.

Paradis E, Schliep K. **2019**. Ape 5.0: An environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* **35**: 526–528.

Pausas JG, Lamont BB. **2022**. Fire-released seed dormancy - a global synthesis. *Biological Reviews* **9**.

Pavoine S. **2020**. adiv: An R package to analyse biodiversity in ecology. *Methods in Ecology and Evolution* **11**: 1106–1112.

Pearson TRH, Burslem DFRP, Mullins CE, Dalling JW. **2003**. Functional significance of photoblastic germination in neotropical pioneer trees: A seed's eye view. *Functional Ecology* **17**: 394–402.

Pereira AR, Andrade ACS, Pereira TS, Forzza RC, Rodrigues AS. **2009**. Comportamento germinativo de espécies epífitas e rupícolas de bromeliaceae do parque estadual do Ibitipoca, Minas Gerais, Brasil. *Revista Brasileira de Botânica* **32**: 827–838.

Pimenta MR, Fernandes LS, Pereira UJ, *et al.* **2007**. Floração, germinação e estaquia em espécies de *Lippia* L. (Verbenaceae). *Revista Brasileira de Botânica* **30**: 211–220.

Pons TL. **2014**. Light-mediated germination. In: Gallagher RS, ed. *Seeds: The Ecology of Regeneration in Plant Communities*. UK: CABI, 111–134.

Porembski S. **2007**. Tropical inselbergs: Habitat types, adaptive strategies and diversity patterns. *Revista Brasileira de Botânica* **30**: 579–586.

Porembski S, Silveira FAO, Fiedler PL, *et al.* **2016**. Worldwide destruction of inselbergs and related rock outcrops threatens a unique ecosystem. *Biodiversity and Conservation* **25**: 2827–2830.

Qian H, Jin Y. **2021**. Are phylogenies resolved at the genus level appropriate for studies on

phylogenetic structure of species assemblages? *Plant Diversity* **43**: 255–263.

R Core Team. **2022**. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. http://www. R-project. org/

Ramos DM, Liaffa ABS, Diniz P, *et al.* 2016. Seed tolerance to heating is better predicted by seed dormancy than by habitat type in Neotropical savanna grasses. *International Journal of Wildland Fire* **25**: 1273–1280.

Ranieri BD, Pezzini FF, Garcia QS, Chautems A, França MGC. **2012**. Testing the regeneration niche hypothesis with Gesneriaceae (tribe Sinningiae) in Brazil: Implications for the conservation of rare species. *Austral Ecology* **37**: 125–133.

Revell LJ. **2012**. phytools: An R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution* **3**: 217–223.

Rezende LAL, Fernandes GW, Braga RP, Dias LE, Gomes VM. **2021**. Topsoil depth influences the recovery of rupestrian grasslands degraded by mining. *Revista Brasileira de Ciência do Solo* **45**: e0210056.

Rivera VL, Panero JL, Schilling EE, Crozier BS, Moraes MD. **2016**. Origins and recent radiation of Brazilian Eupatorieae (Asteraceae) in the eastern Cerrado and Atlantic Forest. *Molecular Phylogenetics and Evolution* **97**: 90–100.

Rosenberg MS, Rothstein HR, Gurevitch J. **2013**. Effect Sizes: Conventional Choices and Calculation In: Koricheva J, Gurevitch J, Mengersen K, eds. *Handbook of Meta-Analysis in Ecology and Evolution*. Princeton, New Jersey: Princeton University Press, 61–71.

Saatkamp A, Cochrane A, Commander L, *et al.* 2019. A research agenda for seed-trait functional ecology. *New Phytologist* **221**: 1764–1775.

Safford HD, Martinelli Gustavo. **2000**. Southeast Brazil In: Safford HD, Martinelli G., eds. *Inselbergs*. Berlin: Springer, 339–389.

Schneider CA, Rasband WS, Eliceiri KW. **2012**. NIH Image to ImageJ: 25 years of image

analysis. *Nature Methods* **9**: 671–675.

Silveira FAO, Negreiros D, Araújo LM, Fernandes GW. **2012**. Does seed germination contribute to ecological breadth and geographic range? A test with sympatric *Diplusodon* (Lythraceae) species from rupestrian fields. *Plant Species Biology* **27**: 170–173.

Silveira FAO, Negreiros D, Barbosa NPU, *et al.* 2016. Ecology and evolution of plant diversity in the endangered campo rupestre: a neglected conservation priority. *Plant and Soil* **403**: 129–152.

Silveira FAO, Ribeiro RC, Oliveira DMT, Fernandes GW, Lemos-Filho JP. **2012**. Evolution of physiological dormancy multiple times in Melastomataceae from Neotropical montane vegetation. *Seed Science Research* **22**: 37–44.

Skirycz A, Castilho A, Chaparro C, Carvalho N, Tzotzos G, Siqueira JO. **2014**. Canga biodiversity, a matter of mining. *Frontiers in Plant Science* **5**: 653.

Smith SA, Brown JW. **2018**. Constructing a broadly inclusive seed plant phylogeny. *American Journal of Botany* **105**: 302–314.

Tarré E, Pires BBM, Guimarães APM, Carneiro LA, Forzza RC, Mansur E. **2007**. Germinability after desiccation, storage and cryopreservation of seeds from endemic *Encholirium* Mart. ex Schult. & Schult. f. and *Dyckia* Schult. & Schult. f. species (Bromeliaceae). *Acta Botanica Brasilica* **21**: 777–783.

Tiansawat P, Dalling JW. **2013**. Differential seed germination responses to the ratio of red to far-red light in temperate and tropical species. *Plant Ecology* **214**: 751–764.

Tweddle JC, Dickie JB, Baskin CC, Baskin JM. **2003**. Ecological aspects of seed desiccation sensitivity. *Journal of Ecology* **91**: 294–304.

Vasconcelos MF. **2011**. O que são campos rupestres e campos de altitude nos topos de montanha do leste do Brasil? *Revista Brasileira de Botânica* **34**: 241–246.

Viechtbauer W. **2010**. Conducting Meta-Analyses in R with the metafor Package. *Journal of*

Statistical Software **36**: 1–48.

Vieira BC, Rodrigues BMA, Garcia QS. **2018**. Light exposure time and light quality on

seed germination of *Vellozia* species (Velloziaceae) from Brazilian *campo rupestre*. *Flora*

238: 94–101.

Westoby M, Falster DS, Moles AT, Vesk PA, Wright IJ. **2002**. Plant Ecological Strategies:

Some Leading Dimensions of Variation Between Species. *Annual Review of Ecology and*

Systematics **33**: 125–159.

Willis CG, Baskin CC, Baskin JM, *et al.* 2014. The evolution of seed dormancy:

Environmental cues, evolutionary hubs, and diversification of the seed plants. *New*

Phytologist **203**: 300–309.

Wyatt R. **1997**. Reproductive ecology of granite outcrop plants from the south-eastern United

States. *Journal of the Royal Society of Western Australia* **80**: 123–129.

Wyse S V., Dickie JB. **2018**. Taxonomic affinity, habitat and seed mass strongly predict seed

desiccation response: A boosted regression trees analysis based on 17 539 species. *Annals of*

Botany **121**: 71–83.

Xu S, Dai Z, Guo P, *et al.* 2021. ggtreeExtra: Compact Visualization of Richly Annotated

Phylogenetic Data. *Molecular Biology and Evolution* **38**: 4039–4042.

Yu G, Smith DK, Zhu H, Guan Y, Lam TT. **2017**. ggtree: an R package for visualization

and annotation of phylogenetic trees with their covariates and other associated data. *Methods*

in Ecology and Evolution **8**: 28–36.

Zanetti M, Dayrell RLC, Wardil M V., *et al.* 2020. Seed Functional Traits Provide Support

for Ecological Restoration and ex situ Conservation in the Threatened Amazon Ironstone

Outcrop Flora. *Frontiers in Plant Science* **11**: 599496.

Table 1. Results of phylogenetic signal test for seed traits in Brazilian rocky outcrop vegetation. P.values for

quantitative traits comes from the likelihood ratio test performed by phylosig function, while for qualitative traits

it corresponds to the root-to-tip skewness test performed by the rtestdecdiv function.

Quantitative traits	Lambda	p.value
Seed mass	0.90	< 0.001
Water content	0.72	< 0.001
Percentage of embryoless seeds	0.75	< 0.001
Percentage of viable seeds	0.57	< 0.001
Qualitative traits	p.value	
Primary dormancy	< 0.001	
Dispersal syndrome	< 0.001	
Dispersal season	< 0.001	

1   **Table 2.** Differences in seed functional traits between growth-forms (herbs vs. shrubs), and species distribution

2   (restricted vs. widespread) and microhabitat (mesic vs. mesic/xeric vs. xeric) in Brazilian rocky outcrop

3   vegetation. Bold values indicate p.values < 0.05.

	Growth-form: Shrub		Distribution: Widespread		Microhabitat: Mesic/Xeric		Microhabitat: Xeric	
Quantitative traits	t	p.value	t	p.value	t	p.value	t	p.value
Seed mass	**2.1872**	**0.0298**	-0.5986	0.5501	**2.0141**	**0.0453**	1.6115	0.1086
Water content	0.2789	0.7811	-0.0770	0.9388	0.6075	0.5453	0.7834	0.4358
Embryoless seeds	1.9274	0.0572	-0.2368	0.8134	0.9752	0.3322	1.2967	0.1982
Viable seeds	-1.0221	0.3085	0.2381	0.8122	-0.3040	0.7616	0.7006	0.4847
Qualitative traits	z	p.value	z	p.value	z	p.value	z	p.value
Dormancy	1.4125	0.1578	1.5052	0.1323	1.5266	0.1268	**2.5218**	**0.0117**
Anemochory	-1.2154	0.2242	1.3943	0.1632	1.0047	0.3150	1.1497	0.2503
Autochory	0.0387	0.9691	-0.0710	0.9434	-0.0792	0.9368	-0.0998	0.9205
Zoochory	0.3290	0.7422	0.0048	0.9962	0.6844	0.4937	-0.0040	0.9968
Early-Dry	-0.9086	0.3636	-0.1266	0.8993	-1.0427	0.2971	0.4504	0.6524
Late-Dry	**3.4323**	**0.0006**	-0.7443	0.4567	0.7421	0.4580	-1.3673	0.1713
Early-Rain	0.1036	0.9175	-0.7541	0.4508	-0.4839	0.6284	0.0445	0.9645
Late-Rain	**-2.0137**	**0.0440**	-0.1826	0.8551	**2.0319**	**0.0422**	**2.1507**	**0.0315**

Figure 1. Main vegetation types associated with rock outcrops in Brazil. A. *Campo rupestre* at Serra do Cipó. B. *Canga* at Parque Estadual da Serra do Rola Moça. C. *Campo de altitude* at Parque Nacional do Caparaó. D. Inselberg at Teófilo Otoni municipality. E. Percentage of studies and taxa evaluated on each vegetation type. All locations are in the State of Minas Gerais. Photos by Carlos A. Ordóñez-Parra (A), Roberta L. C. Dayrell (B), Daniela Calaça (C) and Fernando A. O. Silveira (D).

Figure 2. Phylogeny of studied species with available information on seed functional traits. The ten families with the largest number of species in the dataset are labelled. Figure elaborated with the R packages ggtree (Yu *et al.* 2017), ggtreeExtra (Xu *et al.* 2021) and ggnewscale (Campitelli 2022).

Figure 3. Seed trait differences among growth forms and species distribution and microhabitat, based on statistical results from Table 2. Notes: 1. In relation with species form Mesic/Xeric microhabitats. 2. When compared to species from xeric microhabitats. This figure was made using icons from Flaticon.com

2 **Figure 4.** Effect of light availability on germination percentage in *campo rupestre* species (Control: total

3 darkness). Squares indicate the standardised mean effect size for each moderator (growth form, distribution

4 and microhabitat) and whiskers the 95 % confidence interval of the effect size. Coloured estimates indicate

5 significant effects (i.e., where confidence intervals do not overlap zero). Numbers in parentheses indicate

6 the number of observations.

2 **Figure 5.** Germination responses to (A) constant and (B) alternate temperatures in *campo rupestre* species

3 (Control: 25 ºC). Squares indicate the standardised mean effect size for each moderator (growth form,

4 distribution and microhabitat) and whiskers the 95 % confidence interval of the effect size. Coloured

1   estimates indicate significant effects (i.e., where confidence intervals do not overlap zero). Numbers in

2   parenthesis indicate the number of observations.

Figure 6. Germination responses to fire-related cues in *campo rupestre* species (Control: untreated seeds).

(A) Overall effects. (B) Heat shocks temperatures. (C) Heat shocks without the 200 ºC for one minute. (D)

Smoke. Squares indicate the standardised mean effect size for each moderator (growth form, distribution

and microhabitat) and whiskers the 95 % confidence interval of the effect size. Coloured estimates indicate

significant effects (i.e., where confidence intervals do not overlap zero). Numbers in parentheses indicate

the number of observations.

1   **Table S1.** Moran's I value for the quantitative seed traits across different taxonomic levels. Bold values indicate

2   values significantly different from zero.

Seed trait	Genus		Family		Order	
	I	p.value	I	p.value	I	p.value
Seed mass	**0.7035**	**< 0.001**	**0.6545**	**< 0.001**	**-0.3912**	**< 0.001**
Water content	**0.4456**	**< 0.001**	**0.2368**	**0.0287**	-0.0623	0.4592
Percentage of embryoless seeds	**0.6269**	**< 0.001**	**0.4324**	**< 0.001**	-0.0668	0.5038
Percentage of viable seeds	**0.4808**	**< 0.001**	**0.2176**	**< 0.001**	-0.0998	0.1337

1   **Figure S1.** State of the art of the germination ecology of Brazilian rock outcrop vegetation. A. Percentage

2   of studies for each germination ecology topic. B. Number of species studied each major topic and their

3   combinations. C. Number of species studied for each abiotic factor and their combination. Figure made

4   using RAWGraphs (Mauri *et al.* 2017) and the venn R package (Dusa 2022).

5