Running title: An endocytic myosin-1 generates power

Power generation by a myosin-1 essential for endocytosis:

implications for biological mechanism

Ross TA Pedersen[1, 3, 4], Aaron Snoberger[2, 4], Serapion Pyrpassopoulos[2], Daniel Safer[2], David

G Drubin[1*], E Michael Ostap[2*]

[1]Department of Molecular and Cell Biology, University of California, Berkeley, Berkeley, CA

94720

[2]Pennsylvania Muscle Institute, Perelman School of Medicine, University of Pennsylvania,

Philadelphia, PA 19104

[3]Present address: Department of Embryology, Carnegie Institution for Science, Baltimore,

MD 21218

[4]Equal Contribution

*Correspondence: drubin@berkeley.edu, ostap@mail.med.upenn.edu

Summary

Pedersen, Snoberger et al. measure the force-sensitivity of the yeast endocytic the myosin-

1 called Myo5 and find that it is more likely to generate power than to serve as a force-

sensitive anchor in cells. Implications for Myo5's role in clathrin-mediated endocytosis are

discussed.

Abstract

Myosins are required for clathrin-mediated endocytosis, but their precise molecular roles in this process are not known. This is, in part, because the biophysical properties of the relevant motors have not been investigated. Myosins have diverse mechanochemical activities, ranging from powerful contractility against mechanical loads to force-sensitive anchoring. To better understand the essential molecular contribution of myosin to endocytosis, we studied the in vitro force-dependent kinetics of the *Saccharomyces cerevisiae* endocytic type I myosin called Myo5, a motor whose role in clathrin-mediated endocytosis has been meticulously studied in vivo. We report that Myo5 is a low-duty-ratio motor that is activated ~10-fold by phosphorylation, and that its working stroke and actin-detachment kinetics are relatively force-insensitive. Strikingly, the in vitro mechanochemistry of Myo5 is more like that of cardiac myosin than like that of slow anchoring myosin-1s found on endosomal membranes. We therefore propose that Myo5 generates power to augment actin assembly-based forces during endocytosis in cells.

Introduction

During clathrin-mediated endocytosis (CME), the plasma membrane invaginates and undergoes scission to become a cytoplasmic vesicle. Coat proteins like clathrin can deform membranes under low tension (Dannhauser and Ungewickell, 2012; Busch et al., 2015; Cail et al., 2022), but when membrane bending is resisted by membrane tension (Hassinger et al., 2017), the actin cytoskeleton drives plasma membrane invagination (Boulant et al., 2011; Kaplan et al., 2022). In yeasts, including *Saccharomyces cerevisiae* and *Schizosaccharomyces pombe*, turgor pressure opposes plasma membrane invagination, so actin is required at every CME site (Aghamohammadzadeh and Ayscough, 2009; Basu et al., 2013).

The actin cytoskeleton can produce pushing and pulling force, both of which are required for CME in *S. cerevisiae* (Sun et al., 2006). When actin filament ends grow against a surface, they push the surface forward (Mogilner and Oster, 1996, 2003). During CME, actin filaments, bound by coat proteins, grow against the plasma membrane at the base the CME site, driving invagination (Picco et al., 2015; Kaksonen et al., 2005, 2003; Skruzny et al., 2012, Fig. 1). Modeling of the homologous CME machinery in mammalian cells has demonstrated that such actin networks generate sufficient power for CME (Akamatsu et al., 2020), but whether actin assembly alone can overcome turgor pressure in yeast cells is under debate (Nickaeen et al., 2019; Carlsson, 2018).

Additional power may be provided by myosins, which generate tension on actin filaments. The myosins critical for CME, Myo3 and Myo5 in budding yeast, and Myo1e in vertebrates, are type I myosins (Geli and Riezman, 1996; Cheng et al., 2012; Krendel et al.,

2007). Some type I myosins are well-suited to generate power – i.e., they carry out

mechanical work per unit time by consuming ATP to execute a power stroke. Other type I

myosins are ideally suited to serve as force-sensitive anchors – their ATPase cycle is easily

arrested by resistance, locking them in a state that maintains tension without completing the

ATPase cycle or powering movement (Greenberg and Ostap, 2013). The possible roles of

type I myosins in CME depend on whether endocytic myosins are power generators or force-

sensitive anchors.

If endocytic type I myosins are acutely force-sensitive, they might organize the actin

filaments of the endocytic actin network, while if they are less force sensitive, they could

power plasma membrane invagination (Evangelista et al., 2000, Fig. 1). Myosin-1 motors

form a ring at the base of CME sites, where the invaginated membrane meets the plasma

membrane (Mund et al., 2018, Fig. 1). Yeast type I myosins serve at least one organizational

function as a membrane anchor for the actin assembly machinery, a function associated

with the non-motor tail of the molecules (Lewellyn et al., 2015), but motor activity is

required in addition to membrane anchorage (Pedersen and Drubin, 2019). If endocytic

myosin-1s are force sensitive anchors, they may serve a further organizational role by

holding growing filaments in an optimal orientation for force generation (Fig. 1, left). If the

myosins are power-generating motors, they may pull actin filament ends away from the

plasma membrane, deepening the plasma membrane invagination and creating space for

monomer addition and filament elongation (Fig. 1, right), a model supported by the

observation that the actin assembly rate at CME sites depends on type I myosin motors in a

dose-dependent manner (Manenschijn et al., 2019).

To distinguish between these possibilities, we measured the force sensitivity of the

endocytic myosin Myo5 (not to be confused with the vertebrate type V myosin). Myo5 is

insensitive to resistive force compared to related myosins. We therefore propose that Myo5

actively powers CME. Because actin and myosin collaborate in a variety of membrane

remodeling processes, we expect that these results will be instructive beyond CME.

Results and Discussion

Heavy Chain Phosphorylation Activates Myo5 ATPase Activity

To determine how force sensitive Myo5 is, we first needed to measure its unloaded

kinetics. We purified a Myo5 construct containing the motor and lever domains from *S.*

cerevisiae (Fig. 2A). Because phosphorylation of Myo5 at the TEDS site is required for most

CME events and is thought to regulate Myo5's motor activity (Grosshans et al., 2006; Sun et

al., 2006; Bement and Mooseker, 1995), we purified a phosphorylated version and an

unphosphorylated version of the protein separately (See Materials and methods). p21

activated kinase was used to phosphorylate the myosin at the TEDS site (S357), as

determined by control experiments with an S357A mutant (Fig. S1). The phosphorylation

state of preparations was judged to be uniform when ATP-induced actoMyo5 dissociation

transients were well fit by single exponential functions (see below). The yeast light chain

for Myo5, calmodulin (Cmd1, Geli et al., 1998) was purified from *E. coli* and included in

excess in all experiments (Fig. 2A).

We measured the steady-state actin-activated ATPase activities of phosphorylated

and unphosphorylated Myo5 using the NADH-coupled assay (De La Cruz and Ostap, 2009)

in the presence of 0 - 80 μM phalloidin-stabilized actin. Unphosphorylated Myo5 ATPase

activity was largely insensitive to actin: the ATPase rate for at 0 μM actin was 0.14 s^{-1}, while

the maximum ATPase rate measured was 0.39 s^{-1} at 40 μM actin. (Fig. 2B). Phosphorylation

activated Myo5 ATPase activity by about 10-fold (Fig. 2B). The actin concentration

dependence of the phosphorylated Myo5 ATPase rate (k_{obs}) was well fit by:

$$k_{obs} = v_o \frac{V_{\max}[Actin]}{K_{ATPase}+[Actin]}$$ (**Equation 1**)

From the fit, the actin concentration at half-maximum of the ATPase rate (K_{ATPase}) was

determined to be 5.1 ± 0.88 µM, and the maximum ATPase rate (V_{max}) was found to be 3.3 ±

0.15 s^{-1} (Fig, 2B; Table 1).

ATP Binding and ADP Release are Non-Rate Limiting for Myo5 ATPase Activity

Resistive force impacts the rate of myosin detachment from actin, and two biochemical

transitions, ADP release and subsequent ATP binding, determine the detachment rate.

Therefore, we used stopped-flow kinetics to measure ADP release from (Fig. 2C, k_{+5}') and

ATP binding to (Fig. 2C, K_1' and k_{+2}') actoMyo5.

We found that yeast Myo5 does not quench the fluorescence of actin labeled at cys-374

with pyrene iodoacetamide, which is the probe most used to measure actomyosin

attachment and detachment (De La Cruz and Ostap, 2009). Thus, we measured actoMyo5

detachment by monitoring light scattering, which decreases as myosin unbinds actin.

To determine the rate constant for ATP binding, we mixed nucleotide-free actoMyo5

(100 nM) with varying concentrations of ATP and monitored 90° light scattering in the

stopped flow instrument. Light scattering time courses followed single exponential

functions (Fig 2D). For phosphorylated Myo5, the observed rates determined from the fits

increased linearly with ATP concentration (Fig. 2E). At concentrations of > 1 mM ATP, the

actomyosin complex dissociated within the response time of the instrument, precluding

measurement. For unphosphorylated Myo5, the observed rates fit a rectangular hyperbola

with increasing ATP concentration (Fig. 2E).

The mechanism was modeled as (De La Cruz and Ostap, 2009):

131   K_1' k_2' k_{diss}

132            $AM + ATP \rightleftharpoons AM(ATP) \longrightarrow AM.ATP \longrightarrow A + M.ATP$,   (**Scheme 1**)

133   where K_1' is a rapid equilibrium binding step, k_2' is a rate-limiting isomerization to the

134   AM.ATP state, and k_{diss} is the rapid actin dissociation step. The apparent second order rate

135   constant for ATP binding to phosphorylated actoMyo5 was determined by a linear fit to the

136   data ($K_1'k_2' = 0.39 \pm 0.017 \ \mu m^{-1} \ s^{-1}$). The unphosphorylated actoMyo5 data were fit by:

137                    $$k_{obs} = \left[\frac{K_1' \ [ATP]}{1 + K_1' \ [ATP]} \right] k_2' ,$$ (**Equation 2**)

138   and the maximum rate of isomerization ($k_2' = 290 \pm 24 \ s^{-1}$) and ATP affinity ($K_1' = 0.006 \pm$

139    $0.0016 \ \mu M^{-1}$) were determined. The apparent second-order rate constant for ATP binding

140   ($K_1'k_2'$) was determined from a linear fit of the observed rates below 100 μM ATP to be 1.1

141    $\pm 0.28 \ \mu M^{-1} \ s^{-1}$ (Table 1).

142   The rate constant for ADP dissociation (k_{+5}') was measured by preincubating

143   100 μM ADP with 200 nM actoMyo5 and then rapidly mixing with 2.5 mM ATP as shown in

144   **scheme 2**:

145   k_5'

146            $A.M.ADP \rightleftharpoons A.M + ATP \rightleftharpoons A.M(ATP) \longrightarrow A + M.ATP$,   (**Scheme 2**)

When myosin active sites are saturated with ADP, the rate of ATP-induced dissociation of

actomyosin is limited by ADP's slow dissociation. Light scattering transients were fitted by

single exponential functions, yielding rates for ADP release for phosphorylated actoMyo5

($k_{+5}' = 74 \pm 2.0 s^{-1}$) and for unphosphorylated actoMyo5 ($k_{+5}' = 107 \pm 5.9 \ s^{-1}$) (Fig. 2F and

Table 1). The signal-to-noise ratio of the fast light scattering transients is low, resulting in

large uncertainties on these fits. However, these rates are substantially faster than the

steady-state ATPase values, but slower than the maximum rate of ATP-induced actomyosin

dissociation. ADP release for actoMyo5 ADP is much faster than ADP release for vertebrate

Myo1b and Myo1c (Greenberg et al., 2012; Lewis et al., 2006). It is more similar to the

vertebrate endocytic myosin-1, Myo1e (El Mezgueldi et al., 2002). Because ADP release is

rate limiting for detachment of Myo5 and Myo1e from actin, fast ADP release by these

molecules mean that the unloaded actin-attachment lifetimes for endocytic type I myosins

are < 15 ms. This property may make these motors particularly well-suited to function in

dynamic actin networks like those at CME sites, where actin filaments elongate and

"treadmill" into the cytoplasm (Kaksonen et al., 2003, 2005).

Actin gliding is dependent on Myo5 phosphorylation state

Our kinetic results suggest that both phosphorylated and unphosphorylated Myo5 have

low duty ratios (i.e., the motor spends a small fraction of its ATPase cycle bound to actin).

Since ADP release limits the rate of phosphorylated Myo5 detachment from actin at

saturating ATP ($k_{+5}' = 74 \pm 2.0$ s^{-1}) and since we have measured the overall ATPase rate

($V_{max} = 3.3 \pm 0.15$ s^{-1}), we can estimate the duty ratio:

$$Duty\ Ratio = \frac{(\frac{1}{k\prime_5})}{(\frac{1}{Vmax})},$$ (**Equation 3**)

The calculated duty ratio of phosphorylated Myo5 is 0.045. Unphosphorylated Myo5 has a

lower duty ratio (< 0.004).

To assess the effect of phosphorylation on Myo5 motility, we performed in vitro

motility assays at 1 mM ATP. Motors were attached site-specifically to coverslips coated

with anti-His$_6$ antibody. Coverslips were incubated with a range of concentrations of

phosphorylated and unphosphorylated Myo5, creating a titration series of surface

densities. At low Myo5 surface densities (incubation with ≤ 30 nM phosphorylated Myo5,

≤ 150 nM unphosphorylated Myo5), actin filaments failed to bind the coverslip (Fig. 2G and

movies S1 and S2). At higher concentrations, phosphorylated Myo5 moved actin filaments

at velocities ranging from 720 ± 40 nm/s (100 nM phosphorylated Myo5) to 880 ± 90 nm/s

(40 nM) (Fig. 2G and movie S1). These gliding velocities are considerably higher than those

reported by Sun et al., 2006, possibly reflecting differences in the phosphorylation state of

the purified Myo5 protein (see below) or differences in other motility assay conditions,

such as light chain availability. Higher (> 5-fold) surface densities of unphosphorylated

Myo5 were required to achieve smooth motility, but this motility occurred at a

substantially slower speed, ~120 nm/s (Fig. 2G and movie S2). While it is possible that

residual phosphorylated Myo5 in the unphosphorylated prep contributed to this motility,

Sun et al., 2006 similarly reported that Myo5 harboring TEDS site mutations moved actin

filaments much more slowly. The slower actin gliding speed for unphosphorylated myosin

was unexpected given the similar rates of ADP release between phosphorylated and

unphosphorylated Myo5 (Table 1). It is possible that the our kinetics experiments have not

determined the rate limiting step for detachment, but it is more likely that motility of the

unphosphorylated myosin is limited by the slow attachment rate of the motor (Stewart et

al., 2021), as suggested by the slow actin-activated ATPase rate. The activation of Myo5

motility by phosphorylation could explain why fast, cargo-induced endocytosis, which

involves rapid and dynamic actin turnover, requires phosphorylated Myo5, while slower

constitutive endocytosis does not (Grosshans et al., 2006).

198 **Myo5's working stroke comprises two substeps that are consistent with unloaded**

199 **kinetic measurements**

200 The kinetics of actin attachment durations and mechanics of single myosin molecules were

201 measured using an optical trapping instrument that can detect sub-nanometer

202 displacements with millisecond temporal resolution (Woody et al., 2018; Snoberger et al.,

203 2021). We used the three-bead optical trapping geometry in which a biotinylated actin

204 filament is held between two laser-trapped polystyrene beads coated with neutravidin,

205 creating a bead-actin-bead dumbbell (Fig. 3A). Dumbbells were lowered onto pedestal

206 beads that were sparsely coated with phosphorylated Myo5-His$_9$ bound to a surface-

207 adsorbed anti-His$_6$ tag antibody. The positions of trapped beads were detected via

208 quadrant photodiode detectors, and single actomyosin binding events were detected by the

209 decrease in covariance of the positions of the two dumbbell beads (Fig. 3B-D).

210 　　　　Traces acquired at 1, 10, and 1000 μM ATP reveal displacements and drops in

211 covariance during actomyosin binding events. Individual points from covariance traces

212 were fit by double gaussian distributions, with the thresholds for event detection indicated

213 as dotted gray lines in Fig. 3 B-D. Event durations decreased with increasing ATP

214 concentrations (Fig. 3 B-D, blue lines).

215 　　　　The myosin-1 working stroke has been shown to occur in two discrete substeps,

216 with the first substep occurring with actin-activated phosphate release, and the second

217 occurring with ADP release (Jontes et al., 1995; Veigel et al., 1999, Fig. 3E). The substeps

218 can be characterized in optical trapping assays by ensemble averaging single interactions

219 (Veigel et al., 1999; Chen et al., 2012; Laakso et al., 2008), where the detected events are

aligned at their beginnings and forward-averaged in time (Fig 3F-H, left), or aligned at their

ends and reverse-averaged in time (Fig 3F-H, right).

Ensemble averages of Myo5 interactions showed a two-step working stroke at the

three ATP concentrations, but the step-size was most accurately resolved at 10 μM ATP

(see Materials and methods). In this condition, an initial substep of 4.8 nm was followed by

a second substep of 0.2 nm (Fig. 3G). We determined the lifetimes of the substeps by fitting

the ensemble averages with single exponential functions. At 1 μM ATP (Fig. 3F, left trace),

the measured rate (> 30 s^{-1}) of the time-forward average was limited by the covariance

smoothing window, but at 10 and 1000 μM ATP (Fig. 3 G-H, left traces), the rates were 49 ±

1.6 and 50 ± 0.2 s^{-1}, respectively (Fig. 3K) which are similar to the measured ADP release

rate (k_{+5}', 74 ± 2.0 s^{-1}, Table 1) supporting the model that the transition from state-1 to

state-2 accompanies ADP release.

The kinetics of time-reversed averages reveal the lifetime of State 2 (Fig. 3F-H, right

traces). Fitting single exponential functions to these traces reveals rates of 0.59 ± 0.003 and

7.34 ± 0.1 s^{-1} at 1 and 10 μM ATP, respectively (Fig. 3K). At 1000 μM ATP, the observed rate

(> 187 s^{-1}) was limited by the size of the covariance smoothing window (5.25 ms; Fig. 3K).

The observed rates at 1 and 10 μM ATP are consistent with the second order rate constant

for ATP binding of 0.39 ± 0.017 μM^{-1}s^{-1} measured by stopped-flow kinetics ($K_1'k_{+2}'$,

Table 1).

We determined the detachment rates of actomyosin events by plotting the

cumulative frequency of individual attachment durations and fitting a single exponential

function to the data by maximum likelihood estimation (Fig. 3I). Data from 1 and 10 μM

ATP were well fit by single exponentials with rates of 0.88 and 6.87 s^{-1}, respectively (Fig. 3I,

Fig. 3K). These rates match well with the observed rate of ATP binding (Table 1), as well as

the fits for the reverse ensemble averages, indicating that at sub-saturating ATP (1 and 10

μM), detachment is limited by ATP binding (Figure 3J, blue squares & gray diamonds, Fig.

3K). Data from 1000 μM ATP were best described as the sum of 2 exponentials, with the

major rate of 67.8 s^{-1} comprising 92.1% of the total, and a minor rate of 11.6 s^{-1} comprising

7.9% of the total (Fig. 3I, Fig. 3K). The major rate is consistent with both the observed ADP

release rate and the measured forward ensemble average rates, indicating that at

saturating ATP, ADP release limits detachment of actomyosin interactions (Fig. 3J, blue

square and black diamond, Fig. 3K).

Myo5 is a relatively force-insensitive motor

To elucidate the force sensitivity of Myo5, we measured how its actin detachment rate was

affected by mechanical force opposing the power stroke using an isometric feedback

system that maintained the actin filament near its initial position (Takagi et al., 2006). The

initial force applied to Myo5 in this system depends in part on where along the actin

filament Myo5 stochastically binds, so this approach allowed measurement of attachment

durations at a range of resistive forces (Fig. 4A). Plotting attachment durations as a

function of force revealed a general trend of longer attachment durations at higher

resisting forces. At each interaction force, attachment durations are exponentially

distributed and, as expected based on prior isometric feedback experiments, the data

appear noisy when plotted this way (Fig. 4A). Converting these data to detachment rates by

binning them by force at every ten points, averaging, and taking the inverse of the

attachment duration more clearly reveals the trend (Fig. 4B).

The force dependence of the Myo5 detachment rate was fit by the Bell Equation:

$$k(F) = k_0 \cdot e^{\frac{-F \cdot d}{k_B \cdot T}}$$ (**Equation 4**)

where $k(F)$ is the detachment rate at force F, k_0 is the detachment rate in the absence of

load, d is the distance parameter (the distance to the force-dependent transition state and a

measure of force sensitivity), k_B is Boltzmann's constant, and T is the temperature. Best fit

parameters for k_0 and d were determined by maximum likelihood estimation of the

unaveraged data from Fig. 4A, incorporating the instrument response time (15-30 ms,

Woody et al., 2016). The estimated detachment rate in the absence of force is 67.6 s^{-1}, in

close agreement with the measured detachment rate under low load conditions at 1000 μM

(saturating) ATP (67.8 s^{-1}, Fig. 3K). The best fit value for the distance parameter, d, was

1.14 nm.

To put Myo5's force sensitivity in context, we re-plotted the function describing the

force-dependent actin detachment rate of Myo5 alongside the same curves for vertebrate

Myo1b, Myo1c, and β-cardiac myosin, which have previously been determined by the same

experimental approach (Fig. 4C, Laakso et al., 2010; Greenberg et al., 2012; Woody et al.,

2018). The mechanochemistry of Myo5 (d = 1.14 nm) is most like that of β-cardiac (muscle)

myosin (d = 1.3 nm), suggesting that it is well-suited for generating power. The difference

between Myo5 and acutely force-sensitive Myo1b, a tension-sensitive anchor myosin (d =

15 nm), is dramatic. From 0 to 2 pN of resistance, Myo1b attachment lifetimes slow from

~600 ms to ~45 s, resulting in negligible power generation (Fig. 4D). Over the same

interval, Myo5 attachment lifetimes slow very modestly from ~15 ms to ~25 ms, allowing

it to generate considerable power (Fig. 4D). Thus, Myo5 is unlikely to act as a force-

sensitive anchor in cells and is more likely to power movements against a resisting load.

Proposed function of type I myosin in clathrin-mediated endocytosis

Myo5 is one of the best-studied myosin-1 proteins in vivo. Quantitative live cell imaging

and electron microscopy have revealed that it is recruited to CME sites simultaneously with

initiation of actin assembly, where it concentrates at the base of the site as membrane

invagination proceeds (Jonsdottir and Li, 2004; Idrissi et al., 2008). Although it has long

been appreciated that the presence (Geli and Riezman, 1996; Goodson et al., 1996) and

mechanochemical activity (Sun et al., 2006) of type-1 myosins are required for CME, the

mechanistic contribution of motor activity to the dynamic actin network was unknown.

When it was discovered that some type I myosins are acutely force sensitive (Laakso et al.,

2008), it became apparent that these motors could have mechanochemical activities that

range from force-dependent anchoring to power generation during CME. However,

distinguishing among these possibilities has been difficult in cells. Mutant Myo5 molecules

lacking the motor head or bearing mutations intended to lock the ATPase in low and high

actin affinity states each block CME, results that do not reveal the mechanochemical role of

Myo5 (Lewellyn et al., 2015; Idrissi et al., 2012). Perhaps the most informative finding in

cells has been the observation that varying the number of type I myosins at CME sites

results in differences in actin assembly rates (Manenschijn et al., 2019). However, because

resistive load remodels growing branched actin networks in complex ways (Bieling et al.,

2016), even this finding did not clearly differentiate between the possible roles for

endocytic myosin-1s.

Here we have shown that Myo5's motor generates power rather than forming force-

sensitive catch bonds. The overall ATPase rate of Myo5 is slow relative to other power-

generating myosins, but its power stroke and detachment from actin are fast, and they slow

only modestly under load (Fig. 4C). Myo5's relative force insensitivity means it generates

steady power against resistance (Fig. 4D). Because Myo3 and Myo5 can each support CME

in the absence of the other, we suspect that Myo3 is similarly force-insensitive. Given the

structural and functional homology between Myo5 and vertebrate Myo1e, together with

the close agreement of their unloaded kinetics (El Mezgueldi et al., 2002), we also predict

the Myo1e generates biologically relevant power.

Our finding that Myo5's kinetics are relatively force insensitive lead us to interpret

the previously described dose dependence of actin assembly on the number of myosin-1s

at endocytic sites to mean that this motor moves actin filaments at CME sites to power

plasma membrane invagination and create space for new monomers to assemble

(Manenschijn et al., 2019, Fig. 1, right). On the order of 300 myosin molecules (Myo3 and

Myo5 combined) are present at CME sites, mostly where the invaginating membrane meets

the plasma membrane (Mund et al., 2018; Idrissi et al., 2008; Sun et al., 2019; Picco et al.,

2015). A related myosin, Myo1c, also binds membranes and can generate and sustain sub-

piconewton forces parallel to the plane of the membrane, and greater forces when moving

actin filaments away from the membrane or when diffusion within the bilayer is impeded

(Pyrpassopoulos et al., 2016). Myo5's diffusion is likely to be impeded by the many proteins

at the base of CME sites, and it may move actin filaments at an angle to the membrane it is

bound to. Actin subunits "treadmill" towards the cytoplasm in endocytic actin networks at

~50-100 nm/s (Kaksonen et al., 2005, 2003), so Myo5's motility rate of 700-900 nm/s (Fig.

2G), which we would expect resistance to slow only modestly, is fast enough to do work on

the actin network as it assembles. We therefore expect that the myosins power membrane

invagination and relive load to accelerate actin assembly during CME.

Type I myosins are involved in a variety of membrane reshaping events in cells,

where they often interact with growing branched actin networks (Sokac et al., 2006;

Almeida et al., 2011; Joensuu et al., 2014; Krendel et al., 2007; Cheng et al., 2012), but the

relative contributions of myosin motor activity and actin assembly have rarely been

resolved. Here, we demonstrated that a type I myosin critical for CME, a process well-

known to be driven by actin assembly, generates power. Implication of endocytic type I

myosin as a force-insensitive motor suggests that actin assembly and myosin power

generation can be coordinated to do coherent work in membrane remodeling processes.

Materials and methods

Reagents, proteins, and buffers

ATP concentrations were determined spectrophotometrically after each experiment by absorbance at 259 nm, ϵ_{259} = 15,400 M^{-1}cm^{-1}. Rabbit skeletal muscle actin was prepared and gel filtered (Spudich and Watt, 1971). Actin concentrations were determined spectrophotometrically by absorbance at 290 nm, ϵ_{290} = 26,600 M^{-1}cm^{-1}. All actin was stabilized with one molar equivalent of phalloidin (Sigma). Steady state, transient, and single molecule experiments were performed at 20°C in KMg25 buffer (60 mM MOPS pH 7, 25 mM KCl, 1 mM EGTA, 1 mM MgCl$_2$, 1 mM DTT). Apyrase VII was obtained from Sigma. Purity and concentration of purified proteins were determined by comparing in-gel Coomassie blue staining to staining of known amounts of bovine serum albumin (Pierce).

Expression and Purification of Cmd1

The *S. cerevisiae* calmodulin gene *CMD1* was cloned from genomic DNA into a bacterial expression plasmid with a sequence encoding His$_6$-TEV situated at the 5' end to generate pDD2743. pDD2743 was transformed into Rosetta *E. coli*, optimized for expression (Novagen). A saturated overnight culture in LB (10 g/L Bacto tryptone, 5 g/L Bacto yeast extract, 10 g/L NaCl) was used to inoculate a 1 L culture in LB to OD$_{600}$ = 0.1. Cells were grown to OD$_{600}$ = 0.6-1, induced with 0.5 mM IPTG for 5 hours at 37°C, pelleted at 4,225 x g for 20 minutes at 4°C in a Sorvall SLA-3000 (fixed angle) rotor, washed with cold 20 mM HEPES pH 7.5, and repelleted at 2,250 x g for 10 minutes at 4°C in a Jouan CR3i (swinging bucket) centrifuge. Cell pellets were flash frozen in 45 mL lysis buffer (20 mM HEPES pH

7.5, 1 M KCl, 20 mM Imidazole). Upon thawing, cells were lysed by sonication, 2 mg DNase I

(Roche) and triton X-100 to 1% were added, and the resulting lysate was incubated on ice

for 30 minutes, then spun at 92,000 x g for 25 minutes in a Beckman Type 70 Ti rotor. The

supernatant was loaded onto a 1 mL HisTrap HP column (GE healthcare) preequilibrated

with binding buffer (20 mM HEPES pH 7.5, 500 mM KCl, 20 mM imidazole). The column

was washed with 20 mL binding buffer, and Cmd1 was eluted using a 30 mL linear gradient

from 0-100% elution buffer (20 mM HEPES pH 7.5, 500 mM KCl, 500 mM imidazole).

Fractions containing Cmd1 were pooled, Cmd1 was cleaved from His$_6$ with TEV protease

and dialyzed overnight at 4°C into low salt buffer (10 mM Tris pH 7, 25 mM NaCl, 2 mM

MgCl$_2$, 5 mM DTT). Following dialysis, purified, cleaved Cmd1 was bound to a MonoQ

column and eluted using a 10 mL linear gradient from 0-70% high salt buffer (10 mM Tris

pH 7, 1 M NaCl, 2 mM MgCl$_2$, 5 mM DTT). Fractions containing Cmd1 were pooled, dialyzed

into KMg50 buffer (60 mM MOPS pH 7, 50 mM KCl, 1 mM MgCl$_2$, 1 mM EGTA, 1 mM DTT,

5% glycerol), and stored at -80°C.

Expression and Purification of Myo5

Myo5 was coexpressed with the myosin chaperone She4 in *S. cerevisiae.* The *MYO5* open

reading frame (ORF) from *S. cerevisiae* was cloned from genomic DNA and truncated at

Gly[763], generating a construct containing the motor domain and both Cmd1-binding IQ

motifs of the lever arm. The *SHE4* ORF was cloned in its entirety from *S. cerevisiae* genomic

DNA. Both ORFs were ligated into a 2μ expression plasmid with a partially defective *LEU2*

gene (*leu2d*) to ensure high copy number, creating plasmid pDD2744 (parent vector

described in Roy et al., 2011). The *MYO5* ORF was situated with a sequence encoding

AviTag-TEV-His$_9$ at the 3' end. Expression of the *MYO5* and *SHE4* ORFs was driven by a

bidirectional Gal 1/10 promotor.

pDD2744 was transformed into D1074 yeast (Roy et al., 2011). Saturated overnight

cultures in synthetic minimal medium (1.5 g/L Difacto yeast nitrogen base, 5 g/L

ammonium sulfate, supplemented with 2% glucose 20 µg/mL adenine, L-histadine, L-

methionine, and 30 µg/mL L-lysine) were used to inoculate 1.5 L cultures in the same

media with raffinose substituted for glucose to OD$_{600}$ = 0.1. After 18 hours of growth at

30°C, cultures were induced with 2% galactose, Bacto yeast extract was added to 10 g/L,

and Bacto peptone to 20 g/L. After 8 hours of expression, the cells were harvested at 4,225

x g for 20 minutes at 4°C in a Sorvall SLA-3000 rotor, washed with 25 mL cold Milli-Q

water, repelleted at 2,250 x g for 10 minutes at 4°C in a Jouan CR3i centrifuge, resuspended

in 0.2 volumes of cold Milli-Q water, and drop frozen into liquid nitrogen. Lysis was

achieved through cryomilling (10 cycles of 3 minutes grinding with one minute cooldown)

in the large vials of a 6870 freezer/mill (SPEX Sample Prep).

Cell powders were thawed in binding buffer (10 mM Tris pH 7, 500 mM NaCl, 4 mM

MgCl$_2$, 2 mM ATP, 20 mM imidazole, 5 mM DTT) supplemented with 1 mM PMSF, 1 x

cOmplete protease inhibitor cocktail without EDTA (Roche), and 1 µM Cmd1. For

purification of phosphorylated Myo5, 1 µg Pak1 (Sigma, Brzeska et al., 1997, Fig. S1) was

included in the lysis buffer and 10 mM β-glycerophosphate, 5 mM sodium pyrophosphate,

and 50 mM sodium fluoride were included in all purification buffers. For purification of

unphosphorylated Myo5, 4000 units lambda phosphatase (NEB) and 1 mM MnCl$_2$ were

included in the lysis buffer. The lysate was then spun at 345,000 x g for 10 minutes at 4°C

in a Beckman TLA100.3 rotor, filtered through a 0.22 µm filter, and loaded onto a 1 mL

HisTrap HP column. The column was washed with wash buffer (Binding buffer with only

200 mM NaCl), and Myo5 was eluted using a 20 mL linear gradient from 0-100% elution

buffer (wash buffer with 1 M imidazole).

Fractions containing Myo5 were pooled and supplemented with Cmd1 to 1 μM. For

unphosphorylated Myo5 purification, a further 20,000 units lambda phosphatase were

added along with $MnCl_2$ to 1 mM and the fractions were incubated at 30°C for 30 minutes.

Purified protein was dialyzed through a 3.5 KDa MWCO membrane into 1 L storage buffer

(KMg50 with 50% glycerol) overnight at 4°C and again into 500 mL of the same buffer for 2

hours at 4°C, then stored at -20°C.

Kinetic measurements

Kinetic experiments were conducted using an Applied Photophysics (Surrey, UK) SX.18 MV

stopped-flow apparatus. To monitor light scattering at 90˚, 450 nm excitation light was

used with a 400 nm emission filter. Experimental transients were fit by single exponentials

using the software provided with the stopped flow apparatus. 1-7 traces were averaged

together to generate each data point. Steady state actin-activated ATPase activity was

measured using the NADH enzyme-linked assay (De La Cruz and Ostap, 2009). NADH loss

was monitored by absorbance at 340 nm (ϵ_{340} = 6,220 $M^{-1}cm^{-1}$). ATP-induced dissociation

of actoMyo5 was measured and analyzed as described (De La Cruz and Ostap, 2009). 0.04

units/mL apyrase was added to solutions of actoMyo5 before mixing to remove

contaminating ADP and ATP. Unphosphorylated actoMyo5 required prolonged treatment

with apyrase to achieve sufficient signal, presumably because a larger fraction of the

population was bound to ATP left over from purification and because the actin-activated

ATPase rate of unphosphorylated Myo5 is slow. Concentrations reported are post mixing.

Motility assays

Motility assays were performed essentially as described (Lin et al., 2005). Myo5 was

adhered to nitrocellulose coated coverslips via 20 μg anti His$_6$ antibody (Sigma) blocked

with 2 mg/mL casein. Blocking coverslips with bovine serum albumin (BSA) led to inferior

gliding. Motility was recorded in the presence of 1 mM ATP and 5 μM Cmd1. The rate of

actin filament gliding was determined using the manual tracking plugin in Fiji.

Optical trapping

Flow chambers for optical trapping were constructed with double-sided tape and vacuum

grease as previously described (Snoberger et al., 2021; Greenberg et al., 2017). Briefly, the

coverslip was coated with a 0.1% mixture of nitrocellulose and 2.47 μm diameter silica

beads. Coverslips were dried at least 30 minutes and were used within 24 hours of

preparation. To define the walls of the flow cell, 2 strips of double-sided tape were placed

~5 mm apart, a 1 mm thick glass slide was placed on top, and carefully sealed with vacuum

grease after addition of final buffer.

Trapping buffer (KMg25 with 1 mM DTT freshly added) was used for all trapping

assays. A 100x stock of glucose oxidase + catalase (GOC) was freshly prepared by

centrifuging catalase (Sigma, > 30000 U·mL^{-1}) at 15,000 x g for 1 minute, and adding 2 μl of

catalase supernatant to 20 μL of 19.1 U/μL glucose oxidase (Sigma).

0.01 mg/mL anti-His$_6$ antibody (Sigma) was flowed in the chamber and incubated

between 30 seconds and 3 minutes, then immediately blocked with two, 3-minute

incubations of 1 - 2 mg/mL BSA. Stocks of phosphorylated His$_9$-tagged Myo5 were diluted

to 1 nM in trapping buffer with 300 mM added KCl and incubated in the flow cell for 2

minutes. The number of myosins bound to the surface was limited by the surface

concentration of anti-His$_6$ antibody, and the incubation time of anti-His$_6$ antibody was

adjusted daily between 30 seconds and 3 minutes such that 1 of 3 - 5 pedestals tested

showed clear myosin interactions with the actin dumbbell.

Following incubation with Myo5, a second blocking step with two, 3-minute

incubations of 1 - 2 mg/mL BSA was performed. Final buffer added to the flow cell

contained trapping buffer with indicated amount of ATP, 1 µL of GOC added immediately

prior to addition to chamber, and 0.1 - 0.25 nM rabbit skeletal muscle actin polymerized

with 15% biotinylated actin (Cytoskeleton) stabilized by rhodamine-phalloidin (Sigma) at a

1.1-1.2 molar ratio with G-actin concentration. Neutravidin-coated beads were prepared by

incubating 0.4 ng of 0.8 µm diameter polystyrene beads (Polysciences) and coated with 5

mg/mL neutravidin (Thermo Fisher). 3 µL of neutravidin-coated beads were added to one

side of the chamber prior to sealing. All trapping data were acquired within 90 minutes of

addition of the final buffer to the chamber.

Optical trapping experiments were performed at room temperature (20 ± 1 °C)

using a dual beam 1064 nm trapping laser as described in (Woody et al., 2018, 2017). A

single laser beam was split into 2 beams using polarizing beam splitters and steered into a

60x water immersion objective (Nikon). Laser light was projected through an oil

immersion condenser and into quadrant photodiodes (JQ-50P, Electro Optical Components,

Inc.), each of which were conjugate to the back focal plane of the objective. Direct force

detection from the quadrant photodiodes was achieved using a custom-built high-voltage

reverse bias and an amplifier. Data acquisition, beam position control output, and isometric

feedback calculations were controlled with custom-built virtual instruments (Labview,

Matlab).

Individual 0.8 μm diameter neutravidin-coated bead were caught in the two traps

and held approximately 5 μm apart. Trap stiffnesses were adjusted to 0.05 – 0.1 pN/nm for

each trap. A biotinylated actin filament visualized by rhodamine phalloidin was bound to

the two trapped beads, creating a bead-actin-bead dumbbell. The dumbbell was

pretensioned (3-5 pN) by steering one beam using a piezo controlled mirror conjugate to

the back focal plane of the objective, and the surface of pedestal beads were probed for

myosins. Putative myosin interactions were detected via drops in variance of the two

beads, and the 3-dimensional position of the dumbbell relative to the myosin was refined

further by maximizing the rate and size of the observed power stroke deflections. Every 30-

60 s, the dumbbell was moved axially along the actin filament in ~6 nm steps between

trace acquisition to ensure even accessibility of actin-attachment target zones. Stage drift

was corrected via a feedback system using a nano-positioning stage and imaging the

position of the pedestal bead with nm precision (Woody et al., 2017). In experiments using

1 μM ATP, due to the longer actomyosin interactions, stage drift was still observed even

with the stage feedback engaged, leading to a presumed underestimation of the

displacement size. All data were acquired at a sampling rate of 250 kHz.

Isometric optical clamping experiments were performed as described in (Woody et

al., 2018; Takagi et al., 2006) using a digital feedback loop and a 1-D electro-optical

deflector (EOD, LTA4-Crystal, Conoptics) to steer the beam position using input from a high

voltage source (Conoptics, Model 420 Amplifier). Briefly, the position of one bead (the

"transducer" bead) was maintained at a constant position by adjusting the position of the

other bead (referred to as the "motor" bead) during actomyosin interactions. The response

time of the feedback loop during actomyosin interactions was ~15-30 ms.

Optical trap data analysis

Actomyosin interactions for non-isometric optical clamping experiments were detected

using the single-molecule computational tool SPASM (Software for Precise Analysis of

Single Molecules , Blackwell et al., 2021), which uses a calculation of the dumbbell bead

covariances and a change-point algorithm. Data collected at 1000 µM ATP were analyzed at

250 kHz, while data collected at 1 and 10 µM ATP were downsampled to 2 kHz by

averaging every 125 points to enhance analysis speed. Events were detected by calculating

the covariance of the 2 beads using a smoothing window of 33.3, 15, and 5.25 ms and an

averaging window 60, 36, and 12 ms at 1, 10, and 1000 µM ATP, respectively. The

instrument deadtime was calculated to be 2 times the covariance averaging window. For

each 15 s trace, the detected covariance was plotted and fit to double gaussian

distributions, with the smaller mean gaussian corresponding to the actomyosin "bound"

portion and the larger mean gaussian corresponding to the "unbound" portion of events. A

putative event was defined as an event where the covariance starts above the unbound

peak mean, drops below the bound peak mean, and remains below the unbound peak mean

for at least the length of the instrument deadtime prior to returning back above unbound

peak mean. Event starts and ends were further refined using a changepoint algorithm as

described (Blackwell et al., 2021). Attachment durations and ensemble averages of single

events were determined using built-in features in the SPASM software. Exponential fits for

forward and reverse ensemble averages were performed in Origin 2019 graphing &

analysis software (OriginLab).

Events detected in isometric optical clamping experiments were detected as

described in (Takagi et al., 2006) using a zero crossing analysis via custom MATLAB scripts.

Briefly, when a myosin is actively engaged with the dumbbell, force is applied to the

transducer bead, a feedback loop is engaged and opposing force applied to the motor bead

until the position of the transducer bead is restored. Beginnings of events are defined at the

point at which the feedback signal increases from baseline in the motor bead and ends of

events are defined when the feedback signal decreases back below the baseline in the

motor bead.

Online supplemental material

Figure S1 shows the results of a kinase assay demonstrating that Pak1, used to in

purifications of phosphorylated Myo5, specifically phosphorylates Myo5 serine-357.

Acknowledgements

We thank M. Greenberg, E. Lewellyn, and A. Kunibe, each of whom played important roles at the inception of this study. We thank Y.E. Goldman for optical trap instrumentation. We thank M. Ferrin for helpful comments on the manuscript. This work was funded by National Institute of General Medical Sciences (NIGMS) grant R35 GM118149 to D.G.D and grant 5R37GM057247 to E.M.O. R.T.A.P. is currently funded by NIGMS F32 GM142145.

References

Aghamohammadzadeh, S., and K.R. Ayscough. 2009. Differential requirements for actin

during yeast and mammalian endocytosis. *Nat. Cell Biol.* 11:1039–1042.

doi:10.1038/ncb1918.

Akamatsu, M., R. Vasan, D. Serwas, M. Ferrin, P. Rangamani, and D.G. Drubin. 2020.

Principles of self-organization and load adaptation by the actin cytoskeleton during

clathrin-mediated endocytosis. *Elife*. 9:1–40. doi:10.7554/eLife.49840.

Almeida, C.G., A. Yamada, D. Tenza, D. Louvard, G. Raposo, and E. Coudrier. 2011. Myosin 1b

promotes the formation of post-Golgi carriers by regulating actin assembly and

membrane remodelling at the trans-Golgi network. *Nat. Cell Biol.* 13:779–789.

doi:10.1038/ncb2262.

Basu, R., E.L. Munteanu, and F. Chang. 2013. Role of turgor pressure in endocytosis in

fission yeast. *Mol. Biol. Cell*. 25:679–87. doi:10.1091/mbc.E13-10-0618.

Bement, W.M., and M.S. Mooseker. 1995. TEDS rule: A molecular rationale for differential

regulation of myosins by phosphorylation of the heavy chain head. *Cell Motil.*

Cytoskeleton. 31:87–92. doi:10.1002/cm.970310202.

Bieling, P., T. Li, J. Weichsel, R. Mcgorty, P. Jreij, B. Huang, D.A. Fletcher, and R.D. Mullins.

2016. Force Feedback Controls Motor Activity and Mechanical Properties of Self-

Assembling Branched Actin Networks. *Cell*. 164:115–127.

doi:10.1016/j.cell.2015.11.057.

Blackwell, T., W.T. Stump, S.R. Clippinger, and M.J. Greenberg. 2021. Computational Tool for

Ensemble Averaging of Single-Molecule Data. *Biophys. J.* 120:10–20.

doi:10.1016/j.bpj.2020.10.047.

Boulant, S., C. Kural, J.-C. Zeeh, F. Ubelmann, and T. Kirchhausen. 2011. Actin dynamics

counteract membrane tension during clathrin-mediated endocytosis. *Nat. Cell Biol.*

13:1124–31. doi:10.1038/ncb2307.

Brzeska, H., U.G. Knaus, Z.Y. Wang, G.M. Bokoch, and E.D. Korn. 1997. p21-activated kinase

has substrate specificity similar to Acanthamoeba myosin I heavy chain kinase and

activates Acanthamoeba myosin I. *Proc. Natl. Acad. Sci. U. S. A.* 94:1092–1095.

Busch, D.J., J.R. Houser, C.C. Hayden, M.B. Sherman, E.M. Lafer, and J.C. Stachowiak. 2015.

Intrinsically disordered proteins drive membrane curvature. *Nat. Commun.* 6.

doi:10.1038/ncomms8875.

Cail, R.C., C.R. Shirazinejad, and D.G. Drubin. 2022. Induced nanoscale membrane curvature

bypasses the essential endocytic function of clathrin. *J. Cell Biol.* 221.

doi:10.1083/jcb.202109013.

Carlsson, A.E. 2018. Membrane bending by actin polymerization. *Curr. Opin. Cell Biol.* 50:1–

7. doi:10.1016/j.ceb.2017.11.007.

Chen, C., M.J. Greenberg, J.M. Laakso, E.M. Ostap, Y.E. Goldman, and H. Shuman. 2012.

Kinetic schemes for post-synchronized single molecule dynamics. *Biophys. J.* 102.

doi:10.1016/j.bpj.2012.01.054.

Cheng, J., A. Grassart, and D.G. Drubin. 2012. Myosin 1E coordinates actin assembly and

cargo trafficking during clathrin-mediated endocytosis. *Mol. Biol. Cell*. 23:2891–904.

doi:10.1091/mbc.E11-04-0383.

Dannhauser, P.N., and E.J. Ungewickell. 2012. Reconstitution of clathrin-coated bud and

vesicle formation with minimal components. *Nat. Cell Biol.* 14:634–639.

doi:10.1038/ncb2478.

Evangelista, M., B.M. Klebl, A.H. Yong, B.A. Webb, T. Leeuw, E. Leberer, M. Whiteway, D.Y.

Thomas, and C. Boone. 2000. A Role for Myosin-I in Actin Assembly through

Interactions with Vrp1p, Bee1p, and the Arp2/3 Complex. *J. Cell Biol.* 148:353–362.

doi:10.1083/jcb.148.2.353.

Geli, M.I., and H. Riezman. 1996. Role of Type I Myosins in Receptor-Mediated Endocytosis

in Yeast. *Science*. 272:533–535.

Geli, M.I., A. Wesp, and H. Riezman. 1998. Distinct functions of calmodulin are required for

the uptake step of receptor-mediated endocytosis in yeast: The type I myosin Myo5p is

one of the calmodulin targets. *EMBO J.* 17:635–647. doi:10.1093/emboj/17.3.635.

Goodson, H. V, B.L. Anderson, H.M. Warrick, L.A. Pon, and J.A. Spudich. 1996. Synthetic

lethality screen identifies a novel yeast myosin I gene (MYO5): myosin I proteins are

required for polarization of the actin cytoskeleton. *J. Cell Biol.* 133:1277–1291.

doi:10.1083/JCB.133.6.1277.

Greenberg, M.J., T. Lin, Y.E. Goldman, H. Shuman, and E.M. Ostap. 2012. Myosin IC generates

power over a range of loads via a new tension-sensing mechanism. *Proc. Natl. Acad.*

Sci. U. S. A. 109:E2433-40. doi:10.1073/pnas.1207811109.

Greenberg, M.J., and E.M. Ostap. 2013. Regulation and control of myosin-I by the motor and

light chain-binding domains. *Trends Cell Biol.* 23:81–9. doi:10.1016/j.tcb.2012.10.008.

Greenberg, M.J., H. Shuman, and E.M. Ostap. 2017. Measuring the Kinetic and Mechanical

Properties of Non-processive Myosins Using Optical Tweezers. *Methods Mol. Biol.*

1486:483–509. doi:10.1201/b22505.

Grosshans, B.L., H. Grötsch, D. Mukhopadhyay, I.M. Fernández, J. Pfannstiel, F.-Z. Idrissi, J.

Lechner, H. Riezman, and M.I. Geli. 2006. TEDS site phosphorylation of the yeast

myosins I is required for ligand-induced but not for constitutive endocytosis of the G

protein-coupled receptor Ste2p. *J. Biol. Chem.* 281:11104–14.

doi:10.1074/jbc.M508933200.

Hassinger, J.E., G. Oster, D.G. Drubin, and P. Rangamani. 2017. Design principles for robust

vesiculation in clathrin-mediated endocytosis. *Proc. Natl. Acad. Sci. U. S. A.* 114:E1118–

E1127. doi:10.1073/pnas.1617705114.

Idrissi, F.-Z., A. Blasco, A. Espinal, and M.I. Geli. 2012. Ultrastructural dynamics of proteins

involved in endocytic budding. *Proc. Natl. Acad. Sci. U. S. A.* 109:E2587–E2594.

doi:10.1073/pnas.1202789109.

Idrissi, F.Z., H. Grötsch, I.M. Fernández-Golbano, C. Presciatto-Baschong, H. Riezman, and

M.I. Geli. 2008. Distinct acto/myosin-I structures associate with endocytic profiles at

the plasma membrane. *J. Cell Biol.* 180:1219–1232. doi:10.1083/jcb.200708060.

Joensuu, M., I. Belevich, O. Rämö, I. Nevzorov, H. Vihinen, M. Puhka, T.M. Witkos, M. Lowe,

M.K. Vartiainen, and E. Jokitalo. 2014. ER sheet persistence is coupled to myosin 1c-

regulated dynamic actin filament arrays. *Mol. Biol. Cell*. 25:1111–1126.

doi:10.1091/mbc.E13-12-0712.

Jonsdottir, G. a., and R. Li. 2004. Dynamics of Yeast Myosin I: Evidence for a Possible Role in

Scission of Endocytic Vesicles. *Development*. 14:1604–1609. doi:10.1016/j.

Jontes, J.D., E.M. Wilson-Kubalek, and R.A. Milligan. 1995. A 32 degree tail swing in brush

border myosin I on ADP release. *Nature*. 378:751–753. doi:10.1038/378751a0.

Kaksonen, M., Y. Sun, and D.G. Drubin. 2003. A pathway for association of receptors,

adaptors, and actin during endocytic internalization. *Cell*. 115:475–87.

Kaksonen, M., C.P. Toret, and D.G. Drubin. 2005. A modular design for the clathrin- and

actin-mediated endocytosis machinery. *Cell*. 123:305–20.

doi:10.1016/j.cell.2005.09.024.

Kaplan, C., S.J. Kenny, X. Chen, J. Schöneberg, E. Sitarska, A. Diz-Muñoz, M. Akamatsu, K. Xu,

and D.G. Drubin. 2022. Load adaptation by endocytic actin networks. *Mol. Biol. Cell*. 33.

doi:10.1091/mbc.E21-11-0589.

Krendel, M., E.K. Osterweil, and M.S. Mooseker. 2007. Myosin 1E interacts with

synaptojanin-1 and dynamin and is involved in endocytosis. *FEBS Lett.* 581:644–50.

doi:10.1016/j.febslet.2007.01.021.

De La Cruz, E.M., and E.M. Ostap. 2009. Kinetic and Equilibrium Analysis of the Myosin

ATPase. *Methods Enzymol.* 455:157–192. doi:10.1016/S0076-6879(08)04206-7.

Laakso, J.M., J.H. Lewis, H. Shuman, and E.M. Ostap. 2008. Myosin I can act as a molecular

force sensor. *Science*. 321:133–6. doi:10.1126/science.1159419.

Laakso, J.M., J.H. Lewis, H. Shuman, and E.M. Ostap. 2010. Control of myosin-I force sensing

by alternative splicing. *Proc. Natl. Acad. Sci. U. S. A.* 107:698–702.

doi:10.1073/pnas.0911426107.

Lewellyn, E.B., R.T. Pedersen, J. Hong, R. Lu, H.M. Morrison, and D.G. Drubin. 2015. An

Engineered Minimal WASP-Myosin Fusion Protein Reveals Essential Functions for

Endocytosis. *Dev. Cell*. 35:281–294. doi:10.1016/j.devcel.2015.10.007.

Lewis, J.H., T. Lin, D.E. Hokanson, and E.M. Ostap. 2006. Temperature dependence of

nucleotide association and kinetic characterization of Myo1b. *Biochemistry*. 45:11589–

97. doi:10.1021/bi0611917.

Lin, T., N. Tang, and E.M. Ostap. 2005. Biochemical and motile properties of Myo1b splice

isoforms. *J. Biol. Chem.* 280:41562–41567. doi:10.1074/jbc.M508653200.

Manenschijn, H.E., A. Picco, M. Mund, J. Ries, and M. Kaksonen. 2019. Type-I myosins

promote actin polymerization to drive membrane bending in endocytosis. *Elife*.

17:490011. doi:10.1101/490011.

El Mezgueldi, M., N. Tang, S.S. Rosenfeld, and E.M. Ostap. 2002. The Kinetic Mechanism of

Myo1e (Human Myosin-IC). *J. Biol. Chem.* 277. doi:10.1074/jbc.M200713200.

Mogilner, A., and G. Oster. 1996. Cell motility driven by actin polymerization. *Biophys. J.*

71:3030–3045. doi:10.1016/S0006-3495(96)79496-1.

Mogilner, A., and G. Oster. 2003. Force Generation by Actin Polymerization II: The Elastic

Ratchet and Tethered Filaments. *Biophys. J.* 84:1591–1605. doi:10.1016/S0006-

3495(03)74969-8.

Mund, M., J.A. Van Der Beek, J. Deschamps, S. Dmitrieff, P. Hoess, J.L. Monster, A. Picco, F.

Nedelec, M. Kaksonen, and J. Ries. 2018. Systematic Nanoscale Analysis of Endocytosis

Links Efficient Vesicle Formation to Patterned Actin Nucleation. *Cell*. 174:884–896.

doi:10.1016/j.cell.2018.06.032.

Nickaeen, M., J. Berro, T.D. Pollard, and B.M. Slepchenko. 2019. Actin assembly produces

sufficient forces for endocytosis in yeast. *Mol. Biol. Cell*. 30:2014–2024.

doi:10.1091/mbc.E19-01-0059.

Pedersen, R.T., and D.G. Drubin. 2019. Type I myosins anchor actin assembly to the plasma

membrane during clathrin-mediated endocytosis. *J. Cell Biol.* 218:1138–1147.

doi:10.1083/jcb.201810005.

Picco, A., M. Mund, J. Ries, F. Nedelec, and M. Kaksonen. 2015. Visualizing the functional

architecture of the endocytic machinery. *Elife*. 4:1–29. doi:10.7554/eLife.04535.

Pyrpassopoulos, S., G. Arpağ, E.A. Feeser, H. Shuman, E. Tüzel, and E.M. Ostap. 2016. Force

generation by membrane-associated myosin-I. *Sci. Rep.* 6:1–14.

doi:10.1038/srep25524.

Roy, M.A., N. Siddiqui, and D. D'Amours. 2011. Dynamic and selective DNA-binding activity

of Smc5, a core component of the Smc5-Smc6 complex. *Cell Cycle*. 10:690–700.

doi:10.4161/cc.10.4.14860.

Skruzny, M., T. Brach, R. Ciuffa, S. Rybina, M. Wachsmuth, and M. Kaksonen. 2012.

Molecular basis for coupling the plasma membrane to the actin cytoskeleton during

clathrin-mediated endocytosis. *Proc. Natl. Acad. Sci. U. S. A.* 109:E2533–E2542.

doi:10.1073/pnas.1207011109.

Snoberger, A., B. Barua, J.L. Atherton, H. Shuman, E. Forgacs, Y.E. Goldman, D.A.

Winkelmann, and E.M. Ostap. 2021. Myosin with hypertrophic cardiac mutation r712l

has a decreased working stroke which is rescued by omecamtiv mecarbil. *Elife*. 10:1–

24. doi:10.7554/eLife.63691.

Sokac, A.M., C. Schietroma, C.B. Gundersen, and W.M. Bement. 2006. Myosin-1c couples

assembling actin to membranes to drive compensatory endocytosis. *Dev. Cell*. 11:629–

40. doi:10.1016/j.devcel.2006.09.002.

Spudich, J.A., and S. Watt. 1971. The Regulation of Rabbit Skeletal Muscle Contraction. *J.*

Biol. Chem. 246:4866–4871.

Stewart, T.J., V. Murthy, S.P. Dugan, and J.E. Baker. 2021. Velocity of myosin-based actin

sliding depends on attachment and detachment kinetics and reaches a maximum when

myosin-binding sites on actin saturate. *J. Biol. Chem.* 297.

doi:10.1016/j.jbc.2021.101178.

Sun, Y., A.C. Martin, and D.G. Drubin. 2006. Endocytic Internalization in Budding Yeast

Requires Coordinated Actin Nucleation and Myosin Motor Activity. *Dev. Cell*. 11:33–46.

doi:10.1016/j.devcel.2006.05.008.

Sun, Y., J. Schöneberg, X. Chen, T. Jiang, C. Kaplan, K. Xu, T.D. Pollard, and D.G. Drubin. 2019.

Direct comparison of clathrin-mediated endocytosis in budding and fission yeast

reveals conserved and evolvable features. *Elife*. 8. doi:10.7554/eLife.50749.

Takagi, Y., E.E. Homsher, Y.E. Goldman, and H. Shuman. 2006. Force generation in single

conventional actomyosin complexes under high dynamic load. *Biophys. J.* 90:1295–

1307. doi:10.1529/biophysj.105.068429.

Veigel, C., L.M. Coluccio, J.D. Jontes, J.C. Sparrow, R. a Milligan, and J.E. Molloy. 1999. The

motor protein myosin-I produces its working stroke in two steps. *Nature*. 398:530–

533. doi:10.1038/19104.

Woody, M.S., M.J. Greenberg, B. Barua, D.A. Winkelmann, Y.E. Goldman, and E.M. Ostap.

2018. Positive cardiac inotrope omecamtiv mecarbil activates muscle despite

suppressing the myosin working stroke. *Nat. Commun.* 9. doi:10.1038/s41467-018-

06193-2.

Woody, M.S., J.H. Lewis, M.J. Greenberg, Y.E. Goldman, and E.M. Ostap. 2016. MEMLET: An

Easy-to-Use Tool for Data Fitting and Model Comparison Using Maximum-Likelihood

Estimation. *Biophys. J.* 111:273–282. doi:10.1016/j.bpj.2016.06.019.

Woody, M.S., E.M. Ostap, Y.E. Goldman, and M. Capitanio. 2017. An ultra-fast EOD-based

force-clamp detects rapid biomechanical transitions. SPIE-Intl Soc Optical Eng. 26.

Figure 1: Models for the functions of actin assembly and myosin activity during

membrane deformation for clathrin-mediated endocytosis

Cartoon diagram illustrating the organization of actin filaments and Myo5 molecules at

endocytic sites. Actin filaments are bound by coat proteins at the tip of the growing

membrane invagination and oriented with their growing ends toward the plasma

membrane, powering membrane invagination. The type I myosin Myo5 could either anchor

the actin network in a favorable orientation (left) or provide an assisting force (right).

Figure 2: In-solution, population biochemical characterization of Myo5

(A) Coomassie-stained SDS-polyacrylamide gels showing example preparations of the

purified Myo5 motor/lever construct and calmodulin (Cmd1, light chain) used in all

experiments. **(B)** The actin concentration dependence of the steady-state ATPase activity of

100 nM unphosphorylated (grey circles) and phosphorylated Myo5 (black circles). Each

data point represents the average of 6-7 time courses, which were 100 s each. The orange

line is a best fit of the phosphorylated Myo5 data to a rectangular hyperbola. **(C)** Schematic

pathway for the Myo5 ATPase cycle. Blue motors are in tightly bound conformations, green

motors are weakly bound/unbound. **(D)** Example light scattering transients reporting on

ATP-induced dissociation of phosphorylated (left, k_{obs} = 17 s^{-1}) and unphosphorylated

(right, k_{obs} = 64.1 s^{-1}) actoMyo5, obtained by mixing 100 nM actoMyo5 (AM) with 94 µm

and 72 µM ATP, respectively, as shown in the inset schematic. The black line is the fit of a

single exponential function to the data. **(E)** ATP concentration dependence of dissociation

of 100 nM unphosphorylated (grey circles) and phosphorylated actoMyo5 (black circles).

Each data point represents 3-6 time courses averaged and fit to a single exponential decay

function. The orange line is a linear best fit of the phosphorylated Myo5 data. The purple

line is a best fit of the unphosphorylated Myo5 data to a rectangular hyperbola.

(F) Example light scattering transients reporting ATP-induced dissociation of ADP-

saturated phosphorylated (left) and unphosphorylated (right) actoMyo5, obtained by

preincubating 200 nM actoMyo5 (AM) with 100 μM ADP, then mixing rapidly with 2.5 mM

ATP, as shown in the inset schematic. The black line is the fit of a single exponential

function to the data. **(G)** Velocity of actin filament gliding, measured at varying surface

densities of Phospho-Myo5 (black circles, orange line) and unphosphorylated Myo5 (gray

circles, purple line) in in vitro motility assays. Myosin concentrations indicate the quantity

of protein incubated in the flow chamber before washing. Each data point represents the

average velocity of 30 – 60 filaments, and the error bars are standard deviations.

Table 1

Summary of rate and equilibrium constants measured for Myo5 in this study. Errors are

standard errors of the fits.

Figure 3: Single molecule, optical trap analysis of Myo5 step size and kinetics

(A) Cartoon schematic of the 3-bead optical trapping setup. A biotinylated actin filament is

tethered between two neutravidin-coated beads that are trapped in a dual beam optical

trap. This bead-actin-bead "dumbbell" is lowered onto pedestal beads that have been

sparsely coated with His_6 antibody to attach Myo5-motor/lever-Avi-Tev-His_9. **(B-D)** Single

Myo5 displacements of a single bead position and covariance traces, calculated using both

beads, showing single molecule interactions acquired in the presence of 1 μM **(B)** 10 μM

(C) and 1000 μM ATP. **(D)**. Blue bars indicate attachment events as identified by covariance

(gray) decreases. The threshold of event detection by the covariance traces are indicated

by dashed gray lines. **(E)** Schematic of displacement traces depicting the 2-step nature of

actomyosin displacements in the optical trap. **(F-H)** Binding events were synchronized at

their beginnings (left) or ends (right) and averaged forward or backward in time,

respectively. Measured total displacement of Myo5 was 5.0 nm at 10 μM ATP, with the 1st

substep contributing a 4.8 nm displacement (arrow 1. in G) and the 2nd substep

contributing a 0.2 nm displacement (arrow 2. In G). **(F-H, left)** Forward-averaged

ensembles synchronized at the beginnings of events. **(F-H, right)** Reverse-averaged

ensembles synchronized at the ends of events. Black and gray lines are single exponential

fits in the forward and reverse ensembles, respectively. **(I)** Cumulative distributions of

attachment durations for Myo5 at 1, 10, and 1000 μM ATP. Blue lines show cumulative

frequency of attachment durations at the indicated ATP concentrations, and the red,

yellow, and green lines indicate fitted exponential distributions at 1, 10, and 1000 μM ATP,

respectively. 1 and 10 μM ATP were fit well to single exponentials, and the 1000 μM ATP

data were best described by the sum of two exponentials. **(J)** Summary of rates at 1, 10, and

1000 μM ATP calculated from (F-H). Blue boxes are the fitted exponential distributions

from (I), black diamonds are forward ensemble fits from (F-H, left), and gray diamonds are

reverse ensemble fits from (F-H, right). At lower concentrations of ATP (1 and 10 μM), the

rate of detachment is limited by ATP association, corresponding to the reverse ensemble

fits, while at saturating ATP concentration (1000 μM), the detachment rate is limited by the

rate of ADP dissociation, corresponding to the forward ensemble fits. **(K)** Summary of rates

determined via single molecule optical trapping. Errors for detachment rates are 95%

confidence intervals. Errors for forward and reverse ensemble fits are standard errors of

the fits. *Detachment rates at 1000 µM ATP were best fit to the sum of 2 exponents. The

major component of the fit (67.8 s^{-1}) comprises 92.1% of the total with the remaining 7.9%

having a rate of 11.6 s^{-1}.

Figure 4: Myo5 attachment lifetimes are substantially less force-dependent than

other known type I myosins

An isometric optical force clamp was utilized to determine the force-sensitivity of the

detachment of Myo5 from actin. **(A)** Durations of individual actomyosin attachments as a

function of force, plotted on a semi-log scale **(B)** The solid black line shows the force

dependence of the detachment rates determined by MLE fitting of unaveraged points in A.

For illustration purposes, attachment durations from (A) were binned by force at every 10

points, averaged, and converted to rates. Best- fit parameters were determined by MLE

fitting and 95% confidence intervals were calculated via bootstrapping. The solid black line

is calculated from best fit parameters (k = 67.6 s^{-1}, d = 1.14 nm), while the gray shaded

region is the 95% confidence interval (k = 62.4-72.9 s^{-1}, d = 1.03-1.26 nm). All MLE fitting

was performed on unaveraged data and was corrected for instrument deadtime. **(C)** The

force dependent detachment rate of Myo5 (from panel B) plotted alongside the force

dependent detachment rates for Myo1b, Myo1c, and β-cardiac muscle myosin, Myh7. **(D)**

Power output for the same four myosins calculated over a range of forces by multiplying

the functions from (C) by the applied force F, and the step size and duty ratios of each

myosin.

Figure S1: P21 Activated Kinase 1 (Pak1) phosphorylates Myo5 on S357

Crude preparations of wild type and S357A Myo5 motor/lever constructs were mixed with

250 µM ATP including 20 µCi of ATPγP32 in kinase assay buffer (5 mM MOPS pH 7, 2.5 mM

β-glycerophosphate, 5 mM MgCl$_2$, 400 µM EDTA, 1 mM EGTA, 50 µM DTT) in either the

presence or absence of Pak1. Reactions were incubated at 25°C for 60 minutes, then

quenched by adding an equal volume of 2x tris urea sample buffer (125 mM Tris pH 6.8,

6 M urea, 2% SDS, 0.1% bromophenol blue, 10% β-mercaptoethanol) and resolved on a

10% polyacrylamide gel. The gel was stained with Coomassie, then dried onto Whatman

paper and exposed to a storage phosphor screen (Amersham). The Coomassie-stained gel

was imaged on a standard photo scanner and the phosphor screen on a Typhoon gel imager

(Amersham). Note that there are differences in baseline labeling in the absence of added

kinase between the two different protein preps, but addition of Pak1 clearly results in

radiolabeling of wild type but not mutant Myo5.

Movie S1: Motility assays with phosphorylated Myo5

Rhodamine phalloidin-labeled actin filaments gliding over coverslips coated with a

concentration series of phosphorylated Myo5 protein in motility buffer with 1 mM ATP.

Movies were collected at 1 frame per second and are played back at 16 frames per second.

Movie S2: Motility assays with unphosphorylated Myo5

Rhodamine phalloidin-labeled actin filaments gliding over coverslips coated with a

concentration series of unphosphorylated Myo5 protein in motility buffer with 1 mM ATP.

847   Short movies of motility at 100 nM and 150 nM unphosphorylated Myo5 were collected

848   because no motility was observed. Movies at all other concentrations were collected at 1

849   frame every 4 seconds and are played back at 16 frames per second. The playback rate of

850   Movie S2 is four times faster than the playback rate of Movie S1.

Figure 1

Arp2/3

Type I Myosin

Coat protein

G-actin

F-actin

Anchoring of actin by Myo5

Power generation by Myo5

Figure 2

Table 1: Rate and equilibrium constants of the Myo5 ATPase cycle

	Phosphorylated Myo5	Unphosphorylated Myo5
Steady-state actin-activated ATPase		
V_{max} (s^{-1})	3.3 (± 0.15)	ND
K_{ATPase} (µM)	5.1 (± 0.88)	ND
ATP binding		
K_1' (µM^{-1})	ND	0.006 (± 0.0016)
k_{+2}' (s^{-1})	≥ 335	290 (± 24)
$K_1'k_{+2}'$ (µM^{-1} s^{-1})[a]	0.39 (± 0.017)[b]	1.1 (± 0.28)[c]
ADP release		
k_{+5}' (s^{-1})	74 (± 2.0)	107 (± 5.9)

[a]Determined from a linear fit of the unbinding rates. [b]Linear fit of all data for Phosphorylated Myo5 in Fig. 2E. [c]Linear fit of observed rates below 100 µM ATP for Unphosphorylated Myo5 in Fig 2E. ND: Not Determined.

Figure 3

K

Summary of rates determined during single molecule optial trapping

[ATP] μM	Detachment Rate (s⁻¹)	Forward Ensemble Fit (s⁻¹)	Reverse Ensemble Fit (s⁻¹)
1	0.88 (+/- 0.07/0.07)	>30	0.59 (+/- 0.003)
10	6.9 (+/- 0.49/0.46)	49 (+/- 1.6)	7.3 (+/- 0.1)
1000	67.8* (+/- 11.9/8.7)	50 (+/- 0.2)	>187

Figure 4