Running title: Myosins power endocytosis

An endocytic myosin essential for plasma membrane invagination

powers motility against resistance

Ross TA Pedersen[1, 3, 4], Aaron Snoberger[2, 4], Serapion Pyrpassopoulos[2], Daniel Safer[2], David

G Drubin[1]*, E Michael Ostap[2]*

[1]Department of Molecular and Cell Biology, University of California, Berkeley, Berkeley, CA

94720

[2]Pennsylvania Muscle Institute, Perelman School of Medicine, University of Pennsylvania,

Philadelphia, PA 19104

[3]Present address: Department of Embryology, Carnegie Institution for Science, Baltimore,

MD 21218

[4]Equal Contribution

*Correspondence: drubin@berkeley.edu, ostap@mail.med.upenn.edu

Abstract

Assembling actin filaments work together with myosins to accomplish a wide array of biological processes. During clathrin-mediated endocytosis (CME), actin assembly and type I myosin cooperate to bend the plasma membrane into a pit that undergoes scission to internalize a cargo-bearing vesicle. How actin assembly and myosin work together in this process is a critical, unanswered question. Some type I myosins directly power motility, while others act as force-sensitive clamps. The *Saccharomyces cerevisiae* endocytic type I myosin Myo5 has been meticulously studied *in vivo*, yet whether this protein's essential CME function is to power membrane invagination or to bind to force-bearing actin filaments to collect and organize them for optimal force production has not been established. We report that Myo5 is a low-duty-ratio motor with a working stroke that is rapid and force-insensitive compared to related myosins that act as force-sensitive anchors. We therefore propose that Myo5 generates power to augment actin assembly-based forces during endocytosis.

Introduction

During clathrin-mediated endocytosis (CME), the plasma membrane invaginates into a pit and undergoes scission to become a cytoplasmic vesicle. The endocytic coat can deform membranes and stabilize curvature for membranes under low tension (Dannhauser and Ungewickell, 2012; Busch et al., 2015; Cail et al., 2022), but in the cell, bending is resisted by membrane tension (Hassinger et al., 2017). Under conditions of high membrane tension, the actin cytoskeleton assists plasma membrane invagination (Boulant et al., 2011; Kaplan et al., 2022). In yeasts, including *Saccharomyces cerevisiae* and *Schizosaccharomyces pombe*, turgor pressure uniformly opposes plasma membrane invagination, so actin assembly is required at every CME site (Aghamohammadzadeh and Ayscough, 2009; Basu et al., 2013). This consistent requirement for actin assembly has made yeasts particularly useful organisms for studying the mechanism by which actin powers membrane deformation during CME.

The actin cytoskeleton can produce pushing and pulling force, both of which are required for CME in *S. cerevisiae* (Sun et al., 2006). When actin filament ends grow against a surface, they push the surface forward, powering motility (Mogilner and Oster, 1996, 2003). During CME, actin filaments bind to coat proteins at the tip of the endocytic invagination (Skruzny et al., 2012), and the filaments grow when new monomers are added at the base of the endocytic site, causing filaments to push against the plasma membrane (Picco et al., 2015; Kaksonen et al., 2005, 2003, Fig. 1). Modeling of the homologous CME actin machinery in mammalian cells has demonstrated that a growing network with this geometry can generate sufficient power for CME (Akamatsu et al., 2020), but whether the

power of actin assembly alone can overcome turgor pressure in yeast cells is under debate

(Nickaeen et al., 2019; Carlsson, 2018).

Myosins, on the other hand, couple ATP hydrolysis to conformational changes that

pull on actin filaments. The myosins critical for CME, Myo3 and Myo5 in budding yeast, and

Myo1e in mammalian cells, are type I myosins (Geli and Riezman, 1996; Cheng et al., 2012;

Krendel et al., 2007). Proper regulation of endocytic myosins is critical for healthy cell

function, as overexpression of Myo1e has recently been discovered to be correlated with

adverse outcomes in lung adenocarcinoma patients (Jusue-Torres et al., 2023). Within the

type I myosin family, some motors are optimized to generate power – i.e., they carry out

mechanical work per unit time by consuming ATP to carry out a power stroke. Other type I

myosins are more ideally suited to serve as force-sensitive anchors – their ATPase cycle is

easily arrested by resistive force, locking them in a state that maintains static tension

without completing the ATPase cycle or powering movement (Greenberg and Ostap, 2013).

For example, detachment of rat Myo1b from actin filaments slows dramatically under

modest resistive forces, making this motor unlikely to power motility but ideally suited to

hold tension (Laakso et al., 2008). A closely related enzyme, mouse Myo1c, has actin

detachment kinetics that are much less sensitive to resistive forces. It takes $\sim$5 pN of

resistive force to slow Myo1c's detachment to the same degree that $\sim$1 pN of resistive force

slows Myo1b's detachment, making Myo1c more ideally suited to generate power and

transport cellular materials (Greenberg et al., 2012). The possible roles of type I myosins in

CME depend on whether endocytic myosins are power generators or force-sensitive

anchors.

Type I myosins could organize actin filaments within this endocytic actin network or

they could displace filaments, invaginating the plasma membrane and facilitating actin

growth by creating space for monomers to bind onto barbed ends (Evangelista et al., 2000).

Myosin-1 motors are organized in a ring at the base of endocytic sites, where the

invaginated membrane meets the plasma membrane (Mund et al., 2018, Fig. 1). Yeast type I

myosins serve at least one organizational function as a membrane anchor for the actin

assembly machinery, which is associated with the non-motor tail of the molecules

(Pedersen and Drubin, 2019; Lewellyn et al., 2015). A mutant Myo5 protein lacking the

sequence encoding its motor domain can suffice as a membrane anchor for the actin

network, but cannot facilitate endocytic internalization (Pedersen and Drubin, 2019). Thus,

if Myo5 is a force sensitive anchor, its essential role in CME may be to hold growing

filaments in an optimal orientation for force generation (Fig. 1, left). On the other hand, the

rate of actin assembly at CME sites depends on the type I myosin motors in a dose-

dependent manner (Manenschijn et al., 2019). If these Myosins are power-generating

motors, they may pull actin filament ends away from the plasma membrane, deepening the

plasma membrane invagination and creating space for monomer addition and filament

elongation (Fig. 1, right).

To distinguish between these possibilities, we measured the force sensitivity of the

S. cerevisiae endocytic type I myosin Myo5 (not to be confused with the vertebrate type-V

myosin). We find that Myo5 motor kinetics are insensitive to resistive force compared to

related myosins. We therefore propose that Myo5 actively powers CME. Because actin and

myosin contribute to a variety of membrane remodeling processes, we expect that these

results will be instructive beyond CME.

Results and Discussion

Heavy Chain Phosphorylation Activates Myo5 ATPase Activity

To measure the force sensitivity of Myo5 kinetics, we first established a baseline by

measuring Myo5 kinetics in the absence of force. We purified, from *S. cerevisiae*, a Myo5

construct containing the motor and lever arm domains. Because phosphorylation of Myo5

at the TEDS site is required for most CME events and is thought to regulate Myo5's motor

activity (Grosshans et al., 2006; Sun et al., 2006; Bement and Mooseker, 1995), we purified

a phosphorylated version and an unphosphorylated version of the protein separately (See

Materials and methods, Fig 2A). The yeast light chain for Myo5, calmodulin (Cmd1, Geli et

al., 1998) was expressed in *E. coli*, purified separately from Myo5, and included in excess in

all experiments (See Materials and methods, Fig. 2A).

The steady-state actin-activated ATPase activities of phosphorylated and

unphosphorylated Myo5 were determined using the NADH-coupled assay (De La Cruz and

Ostap, 2009) in the presence of 0 - 80 µM phalloidin-stabilized actin. Unphosphorylated

Myo5 ATPase activity was largely insensitive to actin: the ATPase rate for

unphosphorylated Myo5 at 0 µM actin was 0.14 s^{-1}, while the maximum ATPase rate

measured was 0.39 s^{-1} at 40 µM actin. (Fig. 2B). Phosphorylation activated Myo5 ATPase

activity by about 10-fold. The actin concentration dependence of the phosphorylated Myo5

ATPase (k_{obs}) was well fit by:

$$k_{obs} = v_o \frac{V_{\max}[Actin]}{K_{\text{ATPase}} + [Actin]} \qquad \textbf{(Equation 1)}$$

From the fit, the actin concentration at half-maximum of the ATPase rate (K_{ATPase}) was

determined to be 5.1 ± 0.88 µM, and the maximum ATPase rate (V_{max}) was found to be 3.3 ±

0.15 s^{-1} (Fig, 2B; Table 1).

ATP Binding and ADP Release are Non-Rate Limiting for Myo5 ATPase Activity

The motility rate of many myosins is limited by the rate of actin detachment, and two

biochemical transitions, ADP release and subsequent ATP binding, limit the detachment

rate. Therefore, we used stopped-flow kinetics to measure ADP release (Fig. 2C, k_{+5}') from

and ATP binding to (Fig. 2C, K_1' and k_{+2}') actoMyo5.

We found that yeast Myo5 does not quench the fluorescence of actin labeled at cys-374

with pyrene iodoacetamide, which is the probe most used to measure actomyosin

attachment and detachment (De La Cruz and Ostap, 2009). Thus, we measured actoMyo5

detachment by monitoring a decrease in light scattering.

To determine the rate constant for ATP binding, we mixed nucleotide-free actoMyo5

(100 nM) with varying concentrations of ATP and monitored 90° light scattering in the

stopped flow instrument. Time courses of light scattering signal followed single

exponential decay functions (Fig 2D). For phosphorylated Myo5, the observed rates

determined from the exponential fits increased linearly with ATP concentration (Fig. 2E).

At concentrations of > 1 mM ATP, the actomyosin complex dissociated within the response

time of the instrument, precluding measurement. For unphosphorylated Myo5, the

observed rates fit a rectangular hyperbola with increasing ATP concentration (Fig. 2E).

The mechanism was modeled as (De La Cruz and Ostap, 2009):

149   K_1' k_2' k_{diss}

150        $AM + ATP \rightleftharpoons AM(ATP) \longrightarrow AM.ATP \longrightarrow A + M.ATP$,   (**Scheme 1**)

151   where K_1' is a rapid equilibrium binding step, k_2' is a rate-limiting isomerization to the

152   AM.ATP state, and k_{diss} is the rapid actin dissociation step. The apparent second order rate

153   constant for ATP binding to phosphorylated actoMyo5 was determined by a linear fit to the

154   data ($K_1'k_2' = 0.39 \pm 0.017$ $\mu m^{-1} s^{-1}$). The unphosphorylated actoMyo5 data were fitted by:

155                    $$k_{obs} = \left[\frac{K_1'\,[ATP]}{1 + K_1'\,[ATP]}\right] k_2' ,$$ (**Equation 2**)

and the maximum rate of dissociation ($k_2' = 290 \pm 24$ s^{-1}) and ATP affinities, ($K_1' = 0.006 \pm$

0.0016 μM^{-1}) were determined. The apparent second-order rate constant for ATP binding

was determined from a linear fit of the observed rates below 100 μM ATP to be $K_1'k_2' = 1.1 \pm$

0.28 $\mu M^{-1} s^{-1}$ (Table 1).

The rate constant for ADP dissociation (k_{+5}') was measured by preincubating 100

μM ADP with 200 nM actoMyo5 and then rapidly mixing with 2.5 mM ATP as shown in

scheme 2:

163   k_5'

$A.M.ADP \rightleftharpoons A.M + ATP \rightleftharpoons A.M(ATP) \longrightarrow A + M.ATP$, (**Scheme 2**)

When myosin active sites are saturated with ADP, the rate of ATP-induced dissociation of

actomyosin is limited by the slow dissociation of ADP. Light scattering transients were

fitted by single exponential functions, yielding rates for ADP release for phosphorylated

actoMyo5 ($k_{+5}' = 74 \pm 2.0 s^{-1}$) and for unphosphorylated actoMyo5 ($k_{+5}' = 107 \pm 5.9$ s^{-1}) (Fig.

2F and Table 1). The signal-to-noise ratio of the fast light scattering transients is low,

resulting in large uncertainties on these fits. However, these rates are substantially faster

than the steady-state ATPase values, but slower than the maximum rate of ATP-induced

actomyosin dissociation. ADP release for actoMyo5 ADP is much faster than ADP release

for either vertebrate Myo1b or Myo1c (Greenberg et al., 2012; Lewis et al., 2006). It is more

similar to the vertebrate Myo5 homologue, Myo1e, which also functions in clathrin-

mediated endocytosis (Mezgueldi et al., 2002; Krendel et al., 2007; Cheng et al., 2012).

Because ADP release is rate limiting for detachment of Myo5 and Myo1e from actin, these

fast values for ADP release mean that the actin-attachment lifetimes for endocytic type I

myosins are < 15 ms. This property may make these motors particularly well-suited to

function in dynamic actin networks like those at CME sites, where actin filaments elongate

and "treadmill" into the cytoplasm (Kaksonen et al., 2003; Picco et al., 2015).

Actin gliding is dependent on Myo5 phosphorylation state

Our kinetic results suggest that both phosphorylated and unphosphorylated Myo5 have

low duty ratios (i.e., the motor spends most of its ATPase cycle unbound from actin). Since

ADP release limits the rate of phosphorylated Myo5 detachment from actin at saturating

ATP (k_{+5}' = 74 ± 2.0 s^{-1}) and since we have measured the overall ATPase rate (V_{max} = 3.3 ±

0.15 s^{-1}), we can estimate the duty ratio:

$$Duty\ Ratio = \frac{(\frac{1}{k\prime_5})}{(\frac{1}{Vmax})} \qquad \textbf{(Equation 3)}$$

The calculated duty ratios of phosphorylated Myo5 is 0.045, with unphosphorylated Myo5

having a lower duty ratio (< 0.004).

To assess the effect of phosphorylation on Myo5 motility, we performed *in vitro*

motility assays at 1 mM ATP. Motors were attached site-specifically to coverslips coated

with anti-His$_6$ antibody. Coverslips were incubated with a range of concentrations of

phosphorylated and unphosphorylated Myo5, creating a titration series of surface

densities. At low Myo5 surface densities (incubation with ≤ 30 nM phosphorylated Myo5,

≤ 150 nM unphosphorylated Myo5), actin filaments failed to bind the coverslip (Fig. 2G and

movies S1 and S2). At higher concentrations, phosphorylated Myo5 moved actin filaments

at velocities ranging from 720 ± 40 nm/s (100 nM phosphorylated Myo5) to 880 ± 90 nm/s

(40 nM) (Fig. 2G and movie S1). These gliding velocities are considerably higher than those

reported by Sun et al., 2006. This may reflect differences in the phosphorylation state of the

purified Myo5 protein (see below) or differences in other motility assay conditions, such as

light chain availability, which we controlled by including excess Cmd1. Higher (> 5-fold)

surface densities of unphosphorylated Myo5 were required to achieve smooth motility, but

this motility occurred at a substantially slower speed, ~120 nm/s (Fig. 2G and movie S2).

Sun et al., 2006 similarly reported that Myo5 harboring TEDS site mutations moved actin

filaments much more slowly. The slower actin gliding speed for unphosphorylated myosin

was unexpected given the similar rates of ADP release between phosphorylated and

unphosphorylated Myo5 (Table 1). It remains possible that the our kinetics experiments

have not determined the rate limiting step for detachment, or more likely that motility of

the unphosphorylated myosin is limited by the slow attachment rate of the motor (Stewart

et al., 2021), as suggested by the slow actin-activated ATPase rate. The activation of Myo5

motility by phosphorylation could explain why fast, cargo-induced endocytosis, which

involves rapid and dynamic actin turnover, requires phosphorylated Myo5, while slower

constitutive endocytosis does not (Grosshans et al., 2006).

Myo5's working stroke comprises two substeps that are consistent with unloaded

kinetic measurements

The kinetics of actin attachment durations and mechanics of single myosin molecules were

measured using an optical trapping instrument that can detect sub-nanometer

displacements with millisecond temporal resolution (Woody et al., 2018; Snoberger et al.,

2021). We used the three-bead optical trapping geometry in which a biotinylated actin

filament is tethered between two laser-trapped polystyrene beads coated with neutravidin,

creating a bead-actin-bead dumbbell (Fig. 3A). Dumbbells were lowered onto pedestal

beads that were sparsely coated with phosphorylated Myo5-His$_9$ bound to a surface-

adsorbed anti-His$_6$ tag antibody. The positions of trapped beads were detected via

quadrant photodiode detectors, and single actomyosin binding events were detected by the

decrease in covariance of the positions of the two dumbbell beads (Fig. 3B-D).

Traces acquired in the presence of 1, 10, and 1000 μM ATP reveal clear

displacements and drops in covariance during actomyosin binding events. Individual

points from covariance traces were fit by double gaussian distributions, with the

thresholds for event detection indicated as dotted gray lines in Fig. 3 B-D. Event durations

clearly decreased with increasing ATP concentrations (Fig. 3 B-D, blue lines).

The myosin-1 working stroke has been shown to occur in two discrete substeps,

with the first substep occurring with actin-activated phosphate release, and the second

occurring with ADP release (Jontes et al., 1995; Veigel et al., 1999). The substeps can be

characterized in optical trapping assays by ensemble averaging single interactions (Veigel

et al., 1999; Chen et al., 2012; Laakso et al., 2008), where the detected events are aligned at

their beginnings and forward-averaged in time (Fig 3F-H, left), or aligned at their ends and

reverse-averaged in time (Fig 3F-H, right).

Ensemble averages of Myo5 interactions showed a two-step working stroke at the

three ATP concentrations, but the step-size was most accurately resolved at 10 µM ATP

(see Materials and Methods). In this condition, an initial substep of 4.8 nm was followed by

a second substep of 0.2 nm (Fig. 3G). We determined the lifetimes of the substeps by fitting

the ensemble averages with single exponential functions. At 1 µM ATP (Fig. 3F, left trace),

the measured rate (> 30 s^{-1}) of the time-forward average was limited by the covariance

smoothing window, but at 10 and 1000 µM ATP (Fig. 3 G-H, left traces), the rates were 49 ±

1.6 and 50 ± 0.2 s^{-1}, respectively (Fig. 3K) which are similar to the measured ADP release

rate (k_{+5}', 74 ± 2.0 s^{-1}, Table 1) supporting the model that the transition from state-1 to

state-2 accompanies ADP release.

The kinetics of time-reversed averages reveal the lifetime of State 2 (Fig. 3F-H, right

traces). Fitting single exponential functions to these traces reveals rates of 0.59 ± 0.003 and

7.34 ± 0.1 s^{-1} at 1 and 10 µM ATP, respectively (Fig. 3K). At 1000 µM ATP, the observed rate

(> 187 s^{-1}) was limited by the size of the covariance smoothing window (5.25 ms; Fig. 3K).

The observed rates at 1 and 10 µM ATP are consistent with the second order rate constant

for ATP binding of 0.39 ± 0.0017 µM^{-1}s^{-1} measured by stopped-flow kinetics ($K_1'k_{+2}'$,

Table 1).

We determined the detachment rates of actomyosin events by plotting the

cumulative frequency of individual attachment durations and fit a single exponential

function to the data by maximum likelihood estimation (Fig. 3I). Data from 1 and 10 µM

ATP fit well by single exponentials with rates of 0.88 and 6.87 s^{-1}, respectively (Fig. 3I, Fig.

3K). These rates match well with the observed rate of ATP binding (Table 1), as well as the

fits for the reverse ensemble averages, indicating that at sub-saturating ATP (1 and 10 μM),

detachment is limited by ATP binding (Figure 3J, blue squares & gray diamonds, Fig. 3K).

Data from 1000 μM ATP were best described as the sum of 2 exponentials, with the major

rate of 67.8 s^{-1} comprising 92.1% of the total, and a minor rate of 11.6 s^{-1} comprising 7.9%

of the total (Fig. 3I, Fig. 3K). The major rate is consistent with both the observed ADP

release rate and the measured forward ensemble average rates, indicating that at

saturating ATP, ADP release limits detachment of actomyosin interactions (Fig. 3J, blue

square and black diamond, Fig. 3K).

Myo5 is a relatively force-insensitive motor

To elucidate the force sensitivity of Myo5, we measured how its actin detachment rate was

affected by mechanical force opposing the power stroke using an isometric feedback

system that maintained the actin filament near its initial position (Takagi et al., 2006).

Because the initial force applied to the myosin during the interaction in this system

depends in part on the position on the actin filament that Myo5 binds, which is subject to

stochasticity, this approach allowed measurement of Myo5 attachment durations at a range

of resistive forces during a single experiment (Fig. 4A). Plotting the attachment durations

as a function of force revealed a general trend of longer attachment durations at higher

resisting forces. At each given interaction force, the attachment durations are

exponentially distributed and, as expected based on prior isometric feedback experiments,

the data appear noisy when plotted in this manner (Fig. 4A). Converting these data to

detachment rates by binning them by force at every ten points, averaging, and taking the

inverse of the attachment duration more clearly reveals the trend (Fig. 4B).

To compare the force sensitivity of Myo5 directly to that of other myosins, the force

dependence of the Myo5 detachment rate was fit by the Bell Equation:

$$k(F) = k_0 \cdot e^{\frac{-F \cdot d}{k_B \cdot T}}$$ (**Equation 4**)

where $k(F)$ is the detachment rate at force F, k_0 is the detachment rate in the absence of

load, d is the distance parameter (the distance to the force-dependent transition state and a

measure of force sensitivity), k_B is Boltzmann's constant, and T is the temperature. Best fit

parameters for k_0 and d were determined by maximum likelihood estimation of the

unaveraged data from Fig. 4A and incorporating the response time (15-30 ms, Woody et al.,

2016) yielding an estimated detachment rate in the absence of force of 67.6 s^{-1}, in close

agreement with the measured detachment rate under low load conditions at 1000 μM

(saturating) ATP (Fig. 3K). The best fit value for the distance parameter, d, was 1.14 nm.

This distance parameter indicates that Myo5 is much less force sensitive than other

characterized type I myosins; the distance parameter for acutely force sensitive rat Myo1b

has been measured to be as high as 18 nm for certain splice isoforms (Laakso et al., 2010),

and even the moderately force sensitive enzyme Myo1c has a distance parameter of 5.2 nm

(Greenberg et al., 2012). Myo5's force sensitivity is more in line with those measured for

muscle myosins, like smooth muscle myosin (d = 1.3 nm, Veigel et al., 2003) or β-cardiac

myosin (d = 0.97, Greenberg et al., 2014), leading us to characterize Myo5 as a power

generating motor, rather than a force sensitive anchor.

Proposed function of type I myosin in clathrin-mediated endocytosis

Myo5 has long been known to play a role in clathrin-mediated endocytosis, but the precise

mechanistic contribution of the motor has remained obscure (Goodson et al., 1996; Geli

and Riezman, 1996). The discovery that both Myo5 and its paralog Myo3 also regulate actin

assembly by binding nucleation promoting factors on their non-motor tail only complicated

matters (Evangelista et al., 2000; Geli et al., 2000; Lechler et al., 2000). Separating the

contributions of the type I myosin motor domain activity from the actin assembly functions

of the non-motor tails of these molecules has been stymied by a limited understanding of

the motor domain itself.

The data we present here indicate that Myo5's motor domain generates power

rather than forming force-sensitive catch bonds. Although the overall ATPase rate of Myo5

is slow relative to other power-generating myosins, its power stroke and detachment from

actin are fast, and resistive force slows its power stroke modestly. While Myo5 can

complete its working stroke and consume ATP to move actin filaments even when

challenged with > 1 pN resistive force, vertebrate Myo1b, a tension-sensitive anchor

myosin, stays attached to actin for 50-100 s at 1 pN of load and does not move the filament

or consume ATP. Myo5's relative force insensitivity allows it to generate steady power

against a wide range of resisting forces/loads. Because Myo3 and Myo5 can each support

CME in the absence of the other (Manenschijn et al., 2019), we suspect that Myo3 is a

similarly force-insensitive motor. Given the structural and functional homology between

Myo5 and vertebrate Myo1e, together with the close agreement of their unloaded kinetics

(Mezgueldi et al., 2002), we also predict the Myo1e may also generate biologically relevant

power.

Myo5's force insensitive kinetics make it most likely that this motor moves actin

filaments relative to the membrane at CME sites, which would serve to assist both in

plasma membrane invagination and in movement of occluded actin filament barbed ends

away from the plasma membrane, creating space for new monomers to assemble

(Manenschijn et al., 2019, Fig. 1, right). Myo5 is found mostly near the base of endocytic

sites, where the invaginating membrane meets the plasma membrane (Mund et al., 2018;

Idrissi et al., 2008). A population of Myo5 has also been localized near the tips of deeply

invaginated endocytic pits in immunogold EM studies, but this population of Myo5 is

thought to appear only just before scission, after the major work to create the membrane

invagination against membrane tension has occurred (Idrissi and Geli, 2014; Idrissi et al.,

2008). A related myosin, Myo1c, when membrane-bound, can generate and sustain sub-

piconewton forces parallel to the plane of the phospholipid bilayer to which it is bound,

and can generate even greater forces when moving actin filaments away from the

membrane or when its diffusion within the bilayer is impeded, for example, by other

associated proteins (Pyrpassopoulos et al., 2016). Myo5 localized at the base of endocytic

invaginations should therefore be capable of binding and moving actin filaments along the

invagination axis, which would serve to further invaginate the membrane (Fig. 1, right).

Type I myosins are involved in a wide variety of membrane reshaping events in

cells, where they often interact with dynamic branched actin networks (Sokac et al., 2006;

Almeida et al., 2011; Joensuu et al., 2014; Krendel et al., 2007; Cheng et al., 2012), but the

relative contributions of myosin motor activity and actin assembly in these processes have

rarely been resolved. Here, we demonstrated that a type I myosin that functions during

clathrin-mediated endocytosis, a process well-known to be driven by actin assembly, is also

capable of generating power. It might have been the case that this myosin showed force-

sensitive binding to actin filaments rather than power generation, which would have

implicated this protein as a recruiter and organizer of force-bearing actin filaments at

endocytic sites. Our biochemical and biophysical analyses suggest that instead,

recruitment of force-bearing actin filaments must be mediated by other actin-binding

components of the endocytic actin network, such as the actin-binding protein Sla2 in yeast

or Hip1R in mammals, each of which contain a talin-like actin binding domain that may

bind actin filaments in a force-sensitive manner (Owen et al., 2022). Implication of

endocytic type I myosin as a force-generating protein suggests that actin assembly and

myosin power generation can be coordinated to do coherent work in membrane

remodeling processes.

Materials and methods

Reagents, proteins, and buffers

ATP concentrations were determined spectrophotometrically after each experiment by

absorbance at 259 nm, ϵ_{259} = 15,400 $M^{-1}cm^{-1}$. Rabbit skeletal muscle actin was prepared

and gel filtered (Spudich and Watt, 1971). Actin concentrations were determined

spectrophotometrically by absorbance at 290 nm, ϵ_{290} = 26,600 $M^{-1}cm^{-1}$. All actin was

stabilized with one molar equivalent of phalloidin (Sigma). Steady state, transient, and

single molecule experiments were performed at 20°C in KMg25 buffer (60 mM MOPS pH 7,

25 mM KCl, 1 mM EGTA, 1 mM $MgCl_2$, 1 mM DTT). Apyrase VII was obtained from Sigma.

Purity and concentration of purified proteins were determined by comparing in-gel

Coomassie blue staining to staining of known amounts of bovine serum albumin (Pierce).

Expression and Purification of Cmd1

The *S. cerevisiae* calmodulin gene *CMD1* was cloned from genomic DNA into a bacterial

expression plasmid with a sequence encoding His_6-TEV situated at the 5' end to generate

pDD2743. pDD2743 was transformed into Rosetta *E. coli*, optimized for expression

(Novagen). A saturated overnight culture in LB was used to inoculate a 1L culture to OD_{600}

= 0.1. Cells were grown to OD_{600} = 0.6-1, induced with 0.5 mM IPTG for 5 hours at 37°C,

pelleted at 5,000 rpm for 20 minutes at 4°C, washed with cold 20 mM HEPES pH 7.5, and

repelleted at 3,500 rpm for 10 minutes at 4°C in a Jouan CR3i centrifuge. Cell pellets were

flash frozen in 45 mL lysis buffer (20 mM HEPES pH 7.5, 1 M KCl, 20 mM Imidazole). Upon

thawing, cells were lysed by sonication, 2 mg DNase I (Roche) and triton X-100 to 1% were

added, and the resulting lysate was incubated on ice for 30 minutes, then spun at 30,000

rpm for 25 minutes in a 70 TI rotor. The supernatant was loaded onto a 1 mL HisTrap HP

column (GE healthcare) preequilibrated with binding buffer (20 mM HEPES pH 7.5, 500

mM KCl, 20 mM imidazole). The column was washed with 20 mL binding buffer and Cmd1

was eluted over a 30 mL 0-100% gradient of elution buffer (20 mM HEPES pH 7.5, 500 mM

KCl, 500 mM imidazole). Fractions containing Cmd1 were pooled, Cmd1 was cleaved from

His$_6$ with TEV protease and dialyzed overnight at 4°C into low salt buffer (10 mM Tris pH 7,

25 mM NaCl, 2 mM MgCl$_2$, 5 mM DTT). Following dialysis, purified, cleaved Cmd1 was

bound to a MonoQ column and eluted over a 10 mL 0-70% gradient of high salt buffer (10

mM Tris pH 7, 1 M NaCl, 2 mM MgCl$_2$, 5 mM DTT). Fractions containing Cmd1 were pooled,

dialyzed into KMg50 buffer (60 mM MOPS pH 7, 50 mM KCl, 1 mM MgCl$_2$, 1 mM EGTA, 1

mM DTT, 5% glycerol), and stored at -80°C.

Expression and Purification of Myo5

Myo5 was coexpressed with the myosin chaperone She4 in *S. cerevisiae.* The *MYO5* open

reading frame (ORF) from *S. cerevisiae* was cloned from genomic DNA and truncated at

Gly^{763}, generating a construct containing the motor domain and both Cmd1-binding IQ

motifs of the lever arm. The *SHE4* ORF was cloned in its entirety from *S. cerevisiae* genomic

DNA. Both ORFs were ligated into a 2µ expression plasmid with a partially defective *LEU2*

gene (*leu2d*) to ensure high copy number, creating plasmid pDD2744 (Roy et al., 2011). The

MYO5 ORF was situated with a sequence encoding AviTag-TEV-His$_9$ at the 3' end.

Expression of the *MYO5* and *SHE4* ORFs was driven by a bidirectional Gal 1/10 promotor.

pDD2744 was transformed into D1074 yeast (Roy et al., 2011). Saturated overnight

cultures in synthetic minimal medium (1.5 g/L Difacto yeast nitrogen base, 5 g/L

ammonium sulfate) supplemented with 2% glucose and dropout uracil and leucine were

used to inoculate 1.5 L cultures in the same media with raffinose substituted for glucose to

OD_{600} = 0.1. After 18 hours of growth at 30°C, cultures were induced with 2% galactose,

Bacto yeast extract was added to 10 g/L, and Bacto peptone to 20 g/L. After 8 hours of

expression, the cells were harvested at 5,000 rpm for 10 minutes at 4°C in a TLA3000

rotor, washed with 25 mL cold Milli-Q water, repelleted at 3,500 rpm for 10 minutes at 4°C

in a Jouan CR3i centrifuge, resuspended in 0.2 volumes of cold Milli-Q water, and drop

frozen into liquid nitrogen. Lysis was achieved through cryomilling (10 cycles of 3 minutes

grinding with one minute cooldown) in the large vials of a 6870 freezer/mill (SPEX Sample

Prep).

Cell powders were thawed in binding buffer (10 mM Tris pH 7, 500 mM NaCl, 4 mM

$MgCl_2$, 2 mM ATP, 20 mM imidazole, 5 mM DTT) supplemented with 1 mM PMSF, 1 x

cOmplete protease inhibitor cocktail without EDTA (Roche), and 1 μM Cmd1. For

purification of phosphorylated Myo5, 1 μg Pak1 (Sigma) was included in the lysis buffer

and 10 mM β-glycerophosphate, 5 mM sodium pyrophosphate, and 50 mM sodium fluoride

were included in all purification buffers. For purification of unphosphorylated Myo5, 4000

units lambda phosphatase (NEB) and 1 mM $MnCl_2$ were included in the lysis buffer. The

lysate was then spun at 80,000 rpm for 10 minutes at 4°C in a TLA100.3 rotor, filtered

through a 0.22 μm filter, and loaded onto a 1 mL HisTrap HP column. The column was

washed with wash buffer (Binding buffer with only 200 mM NaCl), and eluted over a 20 mL

0-100% gradient of elution buffer (wash buffer with 1 M imidazole).

Fractions containing Myo5 were pooled and supplemented with Cmd1 to 1 µM. For

unphosphorylated Myo5 purification, a further 20,000 units lambda phosphatase were

added along with $MnCl_2$ to 1 mM and the fractions were incubated at 30°C for 30 minutes.

Purified protein was dialyzed through a 3.5 KDa MWCO membrane into 1 L storage buffer

(KMg50 with 50% glycerol) overnight at 4°C and again into 500 mL of the same buffer for 2

hours at 4°C, then stored at -20°C.

Kinetic measurements

Kinetic experiments were conducted using an Applied Photophysics (Surrey, UK) SX.18 MV

stopped-flow apparatus. To monitor light scattering at 90°, 450 nm excitation light was

used with a 400 nm emission filter. Experimental transients were fit by single exponentials

using the software provided with the stopped flow apparatus. 1-7 traces were averaged

together to generate each data point. Steady state actin-activated ATPase activity was

measured using the NADH enzyme-linked assay (De La Cruz and Ostap, 2009). NADH loss

was monitored by absorbance at 340 nm (ϵ_{340} = 6,220 $M^{-1}cm^{-1}$). ATP-induced dissociation

of actoMyo5 was measured and analyzed as described (De La Cruz and Ostap, 2009). 0.04

units/mL apyrase was added to solutions of actoMyo5 before mixing to remove

contaminating ADP and ATP. Unphosphorylated actoMyo5 required prolonged treatment

with apyrase to achieve sufficient signal, presumably because a larger fraction of the

population was bound to ATP left over from purification and because the actin-activated

ATPase rate of unphosphorylated Myo5 is slow. Concentrations reported are post mixing.

Motility assays

Motility assays were performed essentially as described (Lin et al., 2005). Myo5 was

adhered to nitrocellulose coated coverslips via 20 µg anti His$_6$ antibody (Sigma) blocked

with 2 mg/mL casein. Blocking coverslips with bovine serum albumin (BSA) led to inferior

gliding. Motility was recorded in the presence of 1 mM Mg^{2+}ATP and 5 µM Cmd1. The rate

of actin filament gliding was determined using the manual tracking plugin in Fiji.

Optical trapping

Flow chambers for optical trapping were constructed with double-sided tape and vacuum

grease as previously described (Snoberger et al., 2021; Greenberg et al., 2017). Briefly, the

coverslip was coated with a 0.1% mixture of nitrocellulose and 2.47 µm diameter silica

beads. Coverslips were dried at least 30 minutes and were used within 24 hours of

preparation. To define the walls of the flow cell, 2 strips of double-sided tape were placed

~5 mm apart, a 1 mm thick glass slide was placed on top, and carefully sealed with vacuum

grease after addition of final buffer.

Trapping buffer (KMg25 with 1 mM DTT freshly added) was used for all trapping

assays. A 100x stock of glucose oxidase + catalase (GOC) was freshly prepared by

centrifuging catalase (Sigma, > 30000 U·mL^{-1}) at 15,000 x g for 1 minute, and adding 2 µl of

catalase supernatant to 20 µL of 19.1 U·µL^{-1} glucose oxidase (Sigma).

0.01 mg·mL^{-1} anti-His$_6$ antibody (Sigma) was flowed in the chamber and incubated

between 30 seconds and 3 minutes, then immediately blocked with two, 3-minute

incubations of 1 - 2 mg·mL^{-1} bovine serum albumin (BSA). Stocks of phosphorylated His$_9$-

tagged Myo5 were diluted to 1 nm in trapping buffer with 300 mM added KCl and

incubated in the flow cell for 2 minutes. The number of myosins bound to the surface was limited by the surface concentration of anti-His$_6$ antibody, and the incubation time of anti-His$_6$ antibody was adjusted daily between 30 seconds and 3 minutes such that 1 of 3 - 5 pedestals tested showed clear myosin interactions with the actin dumbbell.

Following incubation with Myo5, a second blocking step with two, 3-minute incubations of 1 - 2 mg/mL BSA was performed. Final buffer added to the flow cell contained trapping buffer with indicated amount of ATP, 1 μL of GOC added immediately prior to addition to chamber, and 0.1 - 0.25 nM rabbit skeletal muscle actin polymerized with 15% biotinylated actin (Cytoskeleton) stabilized by rhodamine-phalloidin (Sigma) at a 1.1-1.2 molar ratio with G-actin concentration. Neutravidin-coated beads were prepared by incubating 0.4 ng of 0.8 μm diameter polystyrene beads (Polysciences) and coated with 5 mg/mL neutravidin (Thermo Fisher). 3 μL of neutravidin-coated beads were added to one side of the chamber prior to sealing. All trapping data were acquired within 90 minutes of addition of the final buffer to the chamber.

Optical trapping experiments assays were performed at room temperature (20 ± 1 °C) using a dual beam 1064 nm trapping laser as described in (Woody et al., 2018, 2017). A single laser beam was split into 2 beams using polarizing beam splitters and steered into a 60x water immersion objective (Nikon). Laser light was projected through an oil immersion condenser and into quadrant photodiodes (JQ-50P, Electro Optical Components, Inc.), each of which were conjugate to the back focal plane of the objective. Direct force detection from the quadrant photodiodes was achieved using a custom-built high-voltage reverse bias and an amplifier. Data acquisition, beam position control output, and isometric

feedback calculations were controlled with custom-built virtual instruments (Labview,

Matlab).

Individual 0.8 µm diameter neutravidin-coated bead were caught in the two traps

and held approximately 5 µm apart. Trap stiffnesses were adjusted to 0.05 – 0.1 pN/nm

per trap. A biotinylated actin filament visualized by rhodamine phalloidin was bound to the

two trapped beads, creating a bead-actin-bead dumbbell. The dumbbell was pretensioned

(3-5 pN) by steering one beam using a piezo controlled mirror conjugate to the back focal

plane of the objective, and the surface of pedestal beads were probed for myosins. Putative

myosin interactions were detected via drops in variance of the two beads, and the 3-

dimensional position of the dumbbell relative to the myosin was refined further by

maximizing the rate and size of the observed power stroke deflections. Every 30-60 s, the

dumbbell was moved axially along the actin filament in ∼6 nm steps between trace

acquisition to ensure even accessibility of actin-attachment target zones. Stage drift was

corrected via a feedback system using a nano-positioning stage and imaging the position of

the pedestal bead with nm precision (Woody et al., 2017). In experiments using 1 µM ATP,

due to the longer actomyosin interactions, stage drift was still observed even with the stage

feedback engaged, leading to a presumed underestimation of the displacement size. All

data were acquired at a sampling rate of 250 kHz.

Isometric optical clamping experiments were performed as described in (Woody et

al., 2018; Takagi et al., 2006) using a digital feedback loop and a 1-D electro-optical

deflector (EOD, LTA4-Crystal, Conoptics) to steer the beam position using input from a high

voltage source (Conoptics, Model 420 Amplifier). Briefly, the position of one bead (the

"transducer" bead) was maintained at a constant position by adjusting the position of the

other bead (referred to as the "motor" bead) during actomyosin interactions. The response

time of the feedback loop during actomyosin interactions was ~15-30 ms.

Optical trap data analysis

Actomyosin interactions for non-isometric optical clamping experiments were detected

using the single-molecule computational tool SPASM (Software for Precise Analysis of

Single Molecules , Blackwell et al., 2021), which uses a calculation of the dumbbell bead

covariances and a change-point algorithm. Data collected at 1000 μM ATP were analyzed at

250 kHz, while data collected at 1 and 10 μM ATP were downsampled to 2 kHz by

averaging every 125 points to enhance analysis speed. Events were detected by calculating

the covariance of the 2 beads using a smoothing window of 33.3, 15, and 5.25 ms and an

averaging window 60, 36, and 12 ms at 1, 10, and 1000 μM ATP, respectively. The

instrument deadtime was calculated to be 2 times the covariance averaging window. For

each 15 s trace, the detected covariance was plotted and fit to double gaussian

distributions, with the smaller mean gaussian corresponding to the actomyosin "bound"

portion and the larger mean gaussian corresponding to the "unbound" portion of events. A

putative event was defined as an event where the covariance starts above the unbound

peak mean, drops below the bound peak mean, and remains below the unbound peak mean

for at least the length of the instrument deadtime prior to returning back above unbound

peak mean. Event starts and ends were further refined using a changepoint algorithm as

described (Blackwell et al., 2021). Attachment durations and ensemble averages of single

events were determined using built-in features in the SPASM software. Exponential fits for

forward and reverse ensemble averages were performed in Origin 2019 graphing &

analysis software (OriginLab).

Events detected in isometric optical clamping experiments were detected as

described in (Takagi et al., 2006) using a zero crossing analysis via custom MATLAB scripts.

Briefly, when a myosin is actively engaged with the dumbbell, force is applied to the

transducer bead, a feedback loop is engaged and opposing force applied to the motor bead

until the position of the transducer bead is restored. Beginnings of events are defined at

the point at which the feedback signal increases from baseline in the motor bead, and ends

of events are defined when the feedback signal decreases back below the baseline in the

motor bead.

Acknowledgements

Acknowledgements

We acknowledge Michael Greenberg, Eric Lewellyn, and Akemi Kunibe, each of whom

played important roles at the inception of this study. We thank Yale E. Goldman for optical

trap instrumentation. We thank Max Ferrin for helpful comments on the manuscript. This

work was funded by National Institute of General Medical Sciences (NIGMS) grant R35

GM118149 to D.G.D and grant 5R37GM057247 to E.M.O. R.T.A.P. is currently funded by

NIGMS F32 GM142145.

References

Aghamohammadzadeh, S., and K.R. Ayscough. 2009. Differential requirements for actin

during yeast and mammalian endocytosis. *Nat. Cell Biol.* 11:1039–1042.

doi:10.1038/ncb1918.

Akamatsu, M., R. Vasan, D. Serwas, M. Ferrin, P. Rangamani, and D.G. Drubin. 2020.

Principles of self-organization and load adaptation by the actin cytoskeleton during

clathrin-mediated endocytosis. *Elife.* 9:1–40. doi:10.7554/eLife.49840.

Almeida, C.G., A. Yamada, D. Tenza, D. Louvard, G. Raposo, and E. Coudrier. 2011. Myosin 1b

promotes the formation of post-Golgi carriers by regulating actin assembly and

membrane remodelling at the trans-Golgi network. *Nat. Cell Biol.* 13:779–789.

doi:10.1038/ncb2262.

Basu, R., E.L. Munteanu, and F. Chang. 2013. Role of turgor pressure in endocytosis in

fission yeast. *Mol. Biol. Cell.* 25:679–87. doi:10.1091/mbc.E13-10-0618.

Bement, W.M., and M.S. Mooseker. 1995. TEDS rule: A molecular rationale for differential

regulation of myosins by phosphorylation of the heavy chain head. *Cell Motil.*

Cytoskeleton. 31:87–92. doi:10.1002/cm.970310202.

Blackwell, T., W.T. Stump, S.R. Clippinger, and M.J. Greenberg. 2021. Computational Tool for

Ensemble Averaging of Single-Molecule Data. *Biophys. J.* 120:10–20.

doi:10.1016/j.bpj.2020.10.047.

Boulant, S., C. Kural, J.-C. Zeeh, F. Ubelmann, and T. Kirchhausen. 2011. Actin dynamics

counteract membrane tension during clathrin-mediated endocytosis. *Nat. Cell Biol.*

13:1124–31. doi:10.1038/ncb2307.

Busch, D.J., J.R. Houser, C.C. Hayden, M.B. Sherman, E.M. Lafer, and J.C. Stachowiak. 2015.

Intrinsically disordered proteins drive membrane curvature. *Nat. Commun.* 6.

doi:10.1038/ncomms8875.

Cail, R.C., C.R. Shirazinejad, and D.G. Drubin. 2022. Induced nanoscale membrane curvature

bypasses the essential endocytic function of clathrin. *J. Cell Biol.* 221.

doi:10.1083/jcb.202109013.

Carlsson, A.E. 2018. Membrane bending by actin polymerization. *Curr. Opin. Cell Biol.* 50:1–

7. doi:10.1016/j.ceb.2017.11.007.

Chen, C., M.J. Greenberg, J.M. Laakso, E.M. Ostap, Y.E. Goldman, and H. Shuman. 2012.

Kinetic schemes for post-synchronized single molecule dynamics. *Biophys. J.* 102.

doi:10.1016/j.bpj.2012.01.054.

Cheng, J., A. Grassart, and D.G. Drubin. 2012. Myosin 1E coordinates actin assembly and

cargo trafficking during clathrin-mediated endocytosis. *Mol. Biol. Cell*. 23:2891–904.

doi:10.1091/mbc.E11-04-0383.

Dannhauser, P.N., and E.J. Ungewickell. 2012. Reconstitution of clathrin-coated bud and

vesicle formation with minimal components. *Nat. Cell Biol.* 14:634–639.

doi:10.1038/ncb2478.

Evangelista, M., B.M. Klebl, A.H. Yong, B.A. Webb, T. Leeuw, E. Leberer, M. Whiteway, D.Y.

Thomas, and C. Boone. 2000. A Role for Myosin-I in Actin Assembly through

Interactions with Vrp1p, Bee1p, and the Arp2/3 Complex. *J. Cell Biol.* 148:353–362.

doi:10.1083/jcb.148.2.353.

Geli, M.I., R. Lombardi, B. Schmelzl, and H. Riezman. 2000. An intact SH3 domain is required

for myosin I-induced actin polymerization. *EMBO J.* 19:4281–4291.

doi:10.1093/emboj/19.16.4281.

Geli, M.I., and H. Riezman. 1996. Role of Type I Myosins in Receptor-Mediated Endocytosis

608   in Yeast. *Science*. 272:533–535.

609   Geli, M.I., A. Wesp, and H. Riezman. 1998. Distinct functions of calmodulin are required for

610   the uptake step of receptor-mediated endocytosis in yeast: The type I myosin Myo5p is

611   one of the calmodulin targets. *EMBO J.* 17:635–647. doi:10.1093/emboj/17.3.635.

612   Goodson, H. V, B.L. Anderson, H.M. Warrick, L.A. Pon, and J.A. Spudich. 1996. Synthetic

613   lethality screen identifies a novel yeast myosin I gene (MYO5): myosin I proteins are

614   required for polarization of the actin cytoskeleton. *J. Cell Biol.* 133:1277–1291.

615   doi:10.1083/JCB.133.6.1277.

616   Greenberg, M.J., T. Lin, Y.E. Goldman, H. Shuman, and E.M. Ostap. 2012. Myosin IC generates

617   power over a range of loads via a new tension-sensing mechanism. *Proc. Natl. Acad.*

618   *Sci. U. S. A.* 109:E2433-40. doi:10.1073/pnas.1207811109.

619   Greenberg, M.J., and E.M. Ostap. 2013. Regulation and control of myosin-I by the motor and

620   light chain-binding domains. *Trends Cell Biol.* 23:81–9. doi:10.1016/j.tcb.2012.10.008.

621   Greenberg, M.J., H. Shuman, and E.M. Ostap. 2014. Inherent Force-Dependent Properties of

622   β-Cardiac Myosin Contribute to the Force-Velocity Relationship of Cardiac Muscle.

623   *Biophys. J.* 107:L41–L44. doi:10.1016/j.bpj.2014.11.005.

624   Greenberg, M.J., H. Shuman, and E.M. Ostap. 2017. Measuring the Kinetic and Mechanical

625   Properties of Non-processive Myosins Using Optical Tweezers. *Methods Mol. Biol.*

626   1486:483–509. doi:10.1201/b22505.

627   Grosshans, B.L., H. Grötsch, D. Mukhopadhyay, I.M. Fernández, J. Pfannstiel, F.-Z. Idrissi, J.

628   Lechner, H. Riezman, and M.I. Geli. 2006. TEDS site phosphorylation of the yeast

myosins I is required for ligand-induced but not for constitutive endocytosis of the G

protein-coupled receptor Ste2p. *J. Biol. Chem.* 281:11104–14.

doi:10.1074/jbc.M508933200.

Hassinger, J.E., G. Oster, D.G. Drubin, and P. Rangamani. 2017. Design principles for robust

vesiculation in clathrin-mediated endocytosis. *Proc. Natl. Acad. Sci. U. S. A.* 114:E1118–

E1127. doi:10.1073/pnas.1617705114.

Idrissi, F.-Z., and M.I. Geli. 2014. Zooming in on the molecular mechanisms of endocytic

budding by time-resolved electron microscopy. *Cell. Mol. Life Sci.* 71:641–57.

doi:10.1007/s00018-013-1452-8.

Idrissi, F.Z., H. Grötsch, I.M. Fernández-Golbano, C. Presciatto-Baschong, H. Riezman, and

M.I. Geli. 2008. Distinct acto/myosin-I structures associate with endocytic profiles at

the plasma membrane. *J. Cell Biol.* 180:1219–1232. doi:10.1083/jcb.200708060.

Joensuu, M., I. Belevich, O. Rämö, I. Nevzorov, H. Vihinen, M. Puhka, T.M. Witkos, M. Lowe,

M.K. Vartiainen, and E. Jokitalo. 2014. ER sheet persistence is coupled to myosin 1c-

regulated dynamic actin filament arrays. *Mol. Biol. Cell*. 25:1111–1126.

doi:10.1091/mbc.E13-12-0712.

Jontes, J.D., E.M. Wilson-Kubalek, and R.A. Milligan. 1995. A 32 degree tail swing in brush

border myosin I on ADP release. *Nature*. 378:751–753. doi:10.1038/378751a0.

Jusue-Torres, I., R. Tiv, J.C. Ricarte-Filho, A. Mallisetty, L. Contreras-Vargas, M.J. Godoy-

Calderon, K. Khaddour, K. Kennedy, K. Valyi-Nagy, O. David, M. Menchaca, A. Kottorou,

A. Koutras, F. Dimitrakopoulos, K.M. Abdelhady, M. Massad, I. Rubinstein, L. Feldman, J.

Stewart, T. Shimamura, L. Danilova, and A. Hulbert. 2023. Myo1e overexpression in

lung adenocarcinoma is associated with increased risk of mortality. *Sci. Rep.* 13:4107.

doi:10.1038/s41598-023-30765-y.

Kaksonen, M., Y. Sun, and D.G. Drubin. 2003. A pathway for association of receptors,

adaptors, and actin during endocytic internalization. *Cell*. 115:475–87.

Kaksonen, M., C.P. Toret, and D.G. Drubin. 2005. A modular design for the clathrin- and

actin-mediated endocytosis machinery. *Cell*. 123:305–20.

doi:10.1016/j.cell.2005.09.024.

Kaplan, C., S.J. Kenny, X. Chen, J. Schöneberg, E. Sitarska, A. Diz-Muñoz, M. Akamatsu, K. Xu,

and D.G. Drubin. 2022. Load adaptation by endocytic actin networks. *Mol. Biol. Cell*. 33.

doi:10.1091/mbc.E21-11-0589.

Krendel, M., E.K. Osterweil, and M.S. Mooseker. 2007. Myosin 1E interacts with

synaptojanin-1 and dynamin and is involved in endocytosis. *FEBS Lett.* 581:644–50.

doi:10.1016/j.febslet.2007.01.021.

De La Cruz, E.M., and E.M. Ostap. 2009. Kinetic and Equilibrium Analysis of the Myosin

ATPase. *Methods Enzymol.* 455:157–192. doi:10.1016/S0076-6879(08)04206-7.

Laakso, J.M., J.H. Lewis, H. Shuman, and E.M. Ostap. 2008. Myosin I can act as a molecular

force sensor. *Science*. 321:133–6. doi:10.1126/science.1159419.

Laakso, J.M., J.H. Lewis, H. Shuman, and E.M. Ostap. 2010. Control of myosin-I force sensing

by alternative splicing. *Proc. Natl. Acad. Sci. U. S. A.* 107:698–702.

doi:10.1073/pnas.0911426107.

Lechler, T., A. Shevchenko, A. Shevchenko, and R. Li. 2000. Direct involvement of yeast type

I myosins in Cdc42-dependent actin polymerization. *J. Cell Biol.* 148:363–373.

doi:10.1083/jcb.148.2.363.

Lewellyn, E.B., R.T. Pedersen, J. Hong, R. Lu, H.M. Morrison, and D.G. Drubin. 2015. An

Engineered Minimal WASP-Myosin Fusion Protein Reveals Essential Functions for

Endocytosis. *Dev. Cell*. 35:281–294. doi:10.1016/j.devcel.2015.10.007.

Lewis, J.H., T. Lin, D.E. Hokanson, and E.M. Ostap. 2006. Temperature dependence of

nucleotide association and kinetic characterization of Myo1b. *Biochemistry*. 45:11589–

97. doi:10.1021/bi0611917.

Lin, T., N. Tang, and E.M. Ostap. 2005. Biochemical and motile properties of Myo1b splice

isoforms. *J. Biol. Chem.* 280:41562–41567. doi:10.1074/jbc.M508653200.

Manenschijn, H.E., A. Picco, M. Mund, J. Ries, and M. Kaksonen. 2019. Type-I myosins

promote actin polymerization to drive membrane bending in endocytosis. *Elife*.

17:490011. doi:10.1101/490011.

Mezgueldi, M. El, N. Tang, S.S. Rosenfeld, and E.M. Ostap. 2002. The Kinetic Mechanism of

Myo1e (Human Myosin-IC). *J. Biol. Chem.* 277. doi:10.1074/jbc.M200713200.

Mogilner, A., and G. Oster. 1996. Cell motility driven by actin polymerization. *Biophys. J.*

71:3030–3045. doi:10.1016/S0006-3495(96)79496-1.

Mogilner, A., and G. Oster. 2003. Force Generation by Actin Polymerization II: The Elastic

Ratchet and Tethered Filaments. *Biophys. J.* 84:1591–1605. doi:10.1016/S0006-

3495(03)74969-8.

Mund, M., J.A. Van Der Beek, J. Deschamps, S. Dmitrieff, P. Hoess, J.L. Monster, A. Picco, F.

Nedelec, M. Kaksonen, and J. Ries. 2018. Systematic Nanoscale Analysis of Endocytosis

Links Efficient Vesicle Formation to Patterned Actin Nucleation. *Cell*. 174:884–896.

doi:10.1016/j.cell.2018.06.032.

Nickaeen, M., J. Berro, T.D. Pollard, and B.M. Slepchenko. 2019. Actin assembly produces

sufficient forces for endocytosis in yeast. *Mol. Biol. Cell*. 30:2014–2024.

doi:10.1091/mbc.E19-01-0059.

Owen, L.M., N.A. Bax, W.I. Weis, and A.R. Dunn. 2022. The C-terminal actin-binding domain

of talin forms an asymmetric catch bond with F-actin. *Proc. Natl. Acad. Sci. U. S. A.* 119.

doi:10.1073/pnas.2109329119.

Pedersen, R.T., and D.G. Drubin. 2019. Type I myosins anchor actin assembly to the plasma

membrane during clathrin-mediated endocytosis. *J. Cell Biol.* 218:1138–1147.

doi:10.1083/jcb.201810005.

Picco, A., M. Mund, J. Ries, F. Nedelec, and M. Kaksonen. 2015. Visualizing the functional

architecture of the endocytic machinery. *Elife*. 4:1–29. doi:10.7554/eLife.04535.

Pyrpassopoulos, S., G. Arpağ, E.A. Feeser, H. Shuman, E. Tüzel, and E.M. Ostap. 2016. Force

generation by membrane-associated myosin-I. *Sci. Rep.* 6:1–14.

doi:10.1038/srep25524.

Roy, M.A., N. Siddiqui, and D. D'Amours. 2011. Dynamic and selective DNA-binding activity

of Smc5, a core component of the Smc5-Smc6 complex. *Cell Cycle*. 10:690–700.

doi:10.4161/cc.10.4.14860.

Skruzny, M., T. Brach, R. Ciuffa, S. Rybina, M. Wachsmuth, and M. Kaksonen. 2012.

Molecular basis for coupling the plasma membrane to the actin cytoskeleton during

clathrin-mediated endocytosis. *Proc. Natl. Acad. Sci. U. S. A.* 109:E2533–E2542.

doi:10.1073/pnas.1207011109.

Snoberger, A., B. Barua, J.L. Atherton, H. Shuman, E. Forgacs, Y.E. Goldman, D.A.

Winkelmann, and E.M. Ostap. 2021. Myosin with hypertrophic cardiac mutation r712l

has a decreased working stroke which is rescued by omecamtiv mecarbil. *Elife*. 10:1–

24. doi:10.7554/eLife.63691.

Sokac, A.M., C. Schietroma, C.B. Gundersen, and W.M. Bement. 2006. Myosin-1c couples

assembling actin to membranes to drive compensatory endocytosis. *Dev. Cell*. 11:629–

40. doi:10.1016/j.devcel.2006.09.002.

Spudich, J.A., and S. Watt. 1971. The Regulation of Rabbit Skeletal Muscle Contraction. *J.*

Biol. Chem. 246:4866–4871.

Stewart, T.J., V. Murthy, S.P. Dugan, and J.E. Baker. 2021. Velocity of myosin-based actin

sliding depends on attachment and detachment kinetics and reaches a maximum when

myosin-binding sites on actin saturate. *J. Biol. Chem.* 297.

doi:10.1016/j.jbc.2021.101178.

Sun, Y., A.C. Martin, and D.G. Drubin. 2006. Endocytic Internalization in Budding Yeast

Requires Coordinated Actin Nucleation and Myosin Motor Activity. *Dev. Cell*. 11:33–46.

doi:10.1016/j.devcel.2006.05.008.

Takagi, Y., E.E. Homsher, Y.E. Goldman, and H. Shuman. 2006. Force generation in single

conventional actomyosin complexes under high dynamic load. *Biophys. J.* 90:1295–

1307. doi:10.1529/biophysj.105.068429.

Veigel, C., L.M. Coluccio, J.D. Jontes, J.C. Sparrow, R. a Milligan, and J.E. Molloy. 1999. The

motor protein myosin-I produces its working stroke in two steps. *Nature*. 398:530–

533. doi:10.1038/19104.

Veigel, C., J.E. Molloy, S. Schmitz, and J. Kendrick-Jones. 2003. Load-dependent kinetics of

force production by smooth muscle myosin measured with optical tweezers. *Nat. Cell*

Biol. 5:980–986. doi:10.1038/ncb1060.

Woody, M.S., M.J. Greenberg, B. Barua, D.A. Winkelmann, Y.E. Goldman, and E.M. Ostap.

2018. Positive cardiac inotrope omecamtiv mecarbil activates muscle despite

suppressing the myosin working stroke. *Nat. Commun.* 9. doi:10.1038/s41467-018-

06193-2.

Woody, M.S., J.H. Lewis, M.J. Greenberg, Y.E. Goldman, and E.M. Ostap. 2016. MEMLET: An

Easy-to-Use Tool for Data Fitting and Model Comparison Using Maximum-Likelihood

Estimation. *Biophys. J.* 111:273–282. doi:10.1016/j.bpj.2016.06.019.

Woody, M.S., E.M. Ostap, Y.E. Goldman, and M. Capitanio. 2017. An ultra-fast EOD-based

force-clamp detects rapid biomechanical transitions. SPIE-Intl Soc Optical Eng. 26.

Figure 1: Models for the functions of actin assembly and myosin activity during membrane deformation for clathrin-mediated endocytosis

Cartoon diagram illustrating the organization of actin filaments and Myo5 molecules at endocytic sites. Actin filaments are bound by coat proteins at the tip of the growing membrane invagination and oriented with their growing ends toward the plasma membrane, powering membrane invagination. The type I myosin Myo5 could either anchor the actin network in a favorable orientation (left) or provide an assisting force (right).

Figure 2: In-solution, population biochemical characterization of Myo5

(A) Coomassie-stained SDS-polyacrylamide gels showing example preparations of the purified Myo5 motor/lever construct and calmodulin (Cmd1, light chain) used in all experiments. **(B)** The actin concentration dependence of the steady-state ATPase activity of 100 nM unphosphorylated (grey circles) and phosphorylated Myo5 (black circles). Each data point represents the average of 6-7 time courses, which were 100 s each. The orange line is a best fit of the phosphorylated Myo5 data to a rectangular hyperbola. **(C)** Schematic pathway for the Myo5 ATPase cycle. Blue motors are in tightly bound conformations, green motors are weakly bound/unbound. **(D)** Example light scattering transients reporting on ATP-induced dissociation of phosphorylated (left, k_{obs} = 17 s^{-1}) and unphosphorylated (right, k_{obs} = 64.1 s^{-1}) actoMyo5, obtained by mixing 100 nM actoMyo5 (AM) with 94 μm and 72 μM ATP, respectively, as shown in the inset schematic. The black line is the fit of a single exponential function to the data. **(E)** ATP concentration dependence of ATP-induced dissociation of 100 nM unphosphorylated (grey circles) and phosphorylated Myo5 (black circles). Each data point represents 3-6 time courses averaged and fit to a single

exponential decay function. The orange line is a linear best fit of the phosphorylated Myo5

data. The purple line is a best fit of the unphosphorylated Myo5 data to a rectangular

hyperbola. **(F)** Example light scattering transients reporting ATP-induced dissociation of

ADP-saturated phosphorylated (left) and unphosphorylated (right) actoMyo5, obtained by

preincubating 200 nM actoMyo5 (AM) with 100 µM ADP, then mixing rapidly with 2.5 mM

ATP, as shown in the inset schematic. The black line the fit of a single exponential function

to the data. **(G)** Velocity of actin filament gliding, measured at varying surface densities

Phospho-Myo5 (black circles, orange line) and unphosphorylated Myo5 (gray circles,

purple line) in *in vitro* motility assays. Myosin concentrations indicate the quantity of

protein incubated in the flow chamber before washing. Each data point represents the

average velocity of 30 – 60 filaments, and the error bars are standard deviations.

Table 1

Summary of rate and equilibrium constants measured for Myo5 in this study. Errors are

standard errors of the fits.

Figure 3: Single molecule, optical trap analysis of Myo5 step size and kinetics

(A) Cartoon schematic of the 3-bead optical trapping setup. A biotinylated actin filament is

tethered between two neutravidin-coated beads that are trapped in a dual beam optical

trap. This bead-actin-bead "dumbbell" is lowered onto pedestal beads that have been

sparsely coated with His_6 antibody to attach Myo5-motor/lever-Avi-Tev-His_9. **(B-D)** Single

Myo5 displacements of a single bead and covariance traces, calculated using both beads,

showing single molecule interactions acquired in the presence of 1 µM **(B)** 10 µM **(C)** and

1000 μM ATP. **(D)**. Blue bars indicate attachment events as identified by covariance (gray)

decreases. The threshold of event detection by the covariance traces are indicated by

dashed gray lines. **(E)** Schematic of displacement traces depicting the 2-step nature of

actomyosin displacements in the optical trap. **(F-H)** Binding events were synchronized at

their beginnings (left) or ends (right) and averaged forward or backward in time,

respectively. Measured total displacement of Myo5 was 5.0 nm at 10 μM ATP, with the 1^{st}

substep contributing a 4.8 nm displacement (arrow 1. in G) and the 2^{nd} substep

contributing a 0.2 nm displacement (arrow 2. In G). **(F-H, left)** Forward-averaged

ensembles synchronized at the beginnings of events. **(F-H, right)** Reverse-averaged

ensembles synchronized at the ends of events. Black and gray lines are single exponential

fits in the forward and reverse ensembles, respectively. **(I)** Cumulative distributions of

attachment durations for Myo5 at 1, 10, and 1000 μM ATP. Blue lines show cumulative

frequency of attachment durations at the indicated ATP concentrations, and the red,

yellow, and green lines indicate fitted exponential distributions at 1, 10, and 1000 μM ATP,

respectively. 1 and 10 μM ATP were fit well to single exponentials, and the 1000 μM ATP

data were best described by the sum of two exponentials. **(J)** Summary of rates at 1, 10, and

1000 μM ATP calculated from (F-H). Blue boxes are the fitted exponential distributions

from (I), black diamonds are forward ensemble fits from (F-H, left), and gray diamonds are

reverse ensemble fits from (F-H, right). At lower concentrations of ATP (1 and 10 μM), the

rate of detachment is limited by ATP association, corresponding to the reverse ensemble

fits, while at saturating ATP concentration (1000 μM), the detachment rate is limited by the

rate of ADP dissociation, corresponding to the forward ensemble fits. **(K)** Summary of rates

determined via single molecule optical trapping. Errors for detachment rates are 95%

confidence intervals. Errors for forward and reverse ensemble fits are standard errors of

the fits. *Detachment rates at 1000 µM ATP were best fit to the sum of 2 exponents. The

major component of the fit (67.8 s^{-1}) comprises 92.1% of the total with the remaining 7.9%

having a rate of 11.6 s^{-1}.

Fig 4: Myo5 attachment lifetimes are substantially less force-dependent than

other known type I myosins

An isometric optical force clamp was utilized to determine the force-sensitivity of the

detachment of Myo5 from actin. **(A)** Durations of individual actomyosin attachments as a

function of force, plotted on a semi-log scale **(B)** The solid black line shows the force

dependence of the detachment rates determined by MLE fitting of unaveraged points in A.

For illustration purposes, attachment durations from (A) were binned by force at every 10

points. Best- fit parameters were determined by MLE fitting and 95% confidence intervals

were calculated via bootstrapping. The solid black line is calculated from best fit

parameters (k = 67.6 s^{-1}, d = 1.14 nm), while the gray shaded region is the 95% confidence

interval (k = 62.4-72.9 s^{-1}, d = 1.03-1.26 nm). All MLE fitting was performed on unaveraged

data and was corrected for instrument deadtime.

Movie S1: Motility assays with phosphorylated Myo5

Rhodamine phalloidin-labeled actin filaments gliding over coverslips coated with a

concentration series of phosphorylated Myo5 protein in motility buffer with 1 mM ATP.

Movies were collected at 1 frame per second and are played back at 16 frames per second.

Movie S2: Motility assays with unphosphorylated Myo5

Rhodamine phalloidin-labeled actin filaments gliding over coverslips coated with a

concentration series of unphosphorylated Myo5 protein in motility buffer with 1 mM ATP.

Short movies of motility at 100 nM and 150 nM unphosphorylated Myo5 were collected

because no motility was observed. Movies at all other concentrations were collected at 1

frame every 4 seconds and are played back at 16 frames per second. The playback rate of

Movie S2 is four times faster than the playback rate of Movie S1.

Figure 1

Arp2/3   Type I Myosin   Coat protein

G-actin   F-actin

Anchoring of actin by Myo5

Power generation by Myo5

Figure 2

Table 1: Rate and equilibrium constants of the Myo5 ATPase cycle

	Phosphorylated Myo5	Unphosphorylated Myo5
Steady-state actin-activated ATPase		
V_{max} (s^{-1})	3.3 (± 0.15)	ND
K_{ATPase} (µM)	5.1 (± 0.88)	ND
ATP binding		
K_1' (µM^{-1})	ND	0.006 (± 0.0016)
k_{+2}' (s^{-1})	≥ 335	290 (± 24)
$K_1'k_{+2}'$ (µM^{-1} s^{-1})[a]	0.39 (± 0.017)[b]	1.1 (± 0.28)[c]
ADP release		
k_{+5}' (s^{-1})	74 (± 2.0)	107 (± 5.9)

[a]Determined from a linear fit of the unbinding rates. [b]Linear fit of all data for Phospho Myo5 in Fig. 2E. [c]Linear fit of observed rates below 100 µM ATP in Fig 2E.

Figure 3

K

Summary of rates determined during single molecule optial trapping

[ATP] μM	Detachment Rate (s⁻¹)	Forward Ensemble Fit (s⁻¹)	Reverse Ensemble Fit (s⁻¹)
1	0.88 (+/- 0.07/0.07)	>30	0.59 (+/- 0.003)
10	6.9 (+/- 0.49/0.46)	49 (+/- 1.6)	7.3 (+/- 0.1)
1000	67.8* (+/- 11.9/8.7)	50 (+/- 0.2)	>187

Figure 4