Title

Proximity Proteome Analysis Reveals Novel TREM2 Interactors in the ER-Mitochondria Interface of Human Microglia

Authors

Chulhwan Kwak,[1,2,†‡] Gina M. Finan,[4,†] Yu Rim Park,[5,6] Anjali Garg,[7] Oscar Harari,[7] Ji Young Mun,[5,6]*, Hyun-Woo Rhee,[1,3]* Tae-Wan Kim[4]*

Affiliations

[1]Department of Chemistry, Seoul National University, Seoul 08826, Korea

[2]The Research Institute of Basic Science, Seoul National University, Seoul 08826, Korea

[3]School of Biological Sciences, Seoul National University, Seoul 08826, Korea

[4]Department of Pathology and Cell Biology, Taub Institute for Research on Alzheimer's Disease and the Aging Brain, Columbia University Irving Medical Center, New York, NY 10032, USA

[5]Division of Biomedical science, School of Medicine, Kyungpook National University, Daegu, Republic of Korea

[6]Neural Circuit Research Group, Korea Brain Research Institute, 41062 Daegu, Korea

[7]Department of Psychiatry, Washington University School of Medicine, St. Louis, MO 63108, USA.

*Corresponding author: jymun@kbri.re.kr (J.Y.M), rheehw@snu.ac.kr (H.-W.R.), twk16@cumc.columbia.edu (T.-W.K.)

†These authors contributed equally to this work.

‡Present address: Department of Neurosurgery, Stanford University School of Medicine, Stanford, CA, USA

Abstract

Triggering receptor expressed on myeloid cells 2 (TREM2) plays a central role in microglial biology and the pathogenesis of Alzheimer's disease (AD). Besides DNAX-activating protein 12 (DAP12), a communal adaptor for TREM2 and many other receptors, other cellular interactors of TREM2 remain largely elusive. We employed a 'proximity labeling' approach using a biotin ligase, TurboID, for mapping protein–protein interactions in live mammalian cells. We discovered novel TREM2-proximal proteins with diverse functions, including those localized to the Mitochondria-ER contact sites (MERCs), a dynamic subcellular 'hub' implicated in a number of crucial cell physiology such as lipid metabolism. TREM2 deficiency alters the thickness (inter-organelle distance) of MERCs, a structural parameter of metabolic state, in microglia derived from human induced pluripotent stem cells. Our TurboID-based TREM2 interactome study suggest novel roles for TREM2 in the structural plasticity of the MERCs, raising the possibility that dysregulation of MERC-related TREM2 functions contribute to AD pathobiology.

Introduction

Alzheimer's disease (AD) is the leading cause of dementia in the elderly. In addition to plaques, tangles, and widespread neuronal death, there are profound inflammatory changes in the AD brain, which are increasingly presenting new therapeutic targets (Fu et al., 2019; Wyss-Coray and Mucke, 2002). Microglia act as the brain's immune cells, representing roughly 10% of the cells in the brain, and are becoming the subject of intense investigation in the field of neurodegenerative disease research (Leng and Edison, 2021; Hansen et al., 2018; Sarlus et al., 2017). In AD, microglia are linked closely to all pathological cascades including amyloid, tau and neuroinflammatory changes (McQuade and Blurton-Jones, 2019). Recent genome-wide association studies (GWAS) revealed that a number of AD risk genes are highly (and often exclusively) expressed in microglia, including the gene encoding triggering receptor expressed on myeloid cells-2 (TREM2) (Colonna, 2023; Villegas-Llerena et al., 2016; Griciuc and Tanzi 2021; Jorfi et al., 2023). Notably, a rare R47H hypomorphic mutation in the TREM2 increases the risk of AD several fold (R. Guerreiro et al., 2013; Jonsson et al., 2013). Gene network studies reveal that TREM2 and its adapter protein, DNAX-activation protein 12 (DAP12; also known as TYROBP) strongly associate with AD (Zhang et al., 2013; 2016; Matarin et al., 2015). Homozygous variants in TREM2 or its binding partner DAP12 are also shown to cause Nasu-Hakola disease (NHD) with early-onset dementia (Paloneva et al., 2002; Klunemann et al., 2005). Despite the strong clinical phenotype of human NHD, TREM2-deficient and DAP12-deficient mice do not exert strong NHD-like phenotypes (Zhou et al., 2023).

TREM2 is a transmembrane immune signaling receptor, which serves as a receptor for β-amyloid (Zhao et al., 2018; Lessard et al., 2018; Zhong et al., 2018) and various lipoproteins including high density lipoprotein containing apolipoprotein E (Song et al., 2017; Atagi et al., 2015; Bailey et al., 2015). TREM2 promotes the survival and proliferation of microglia and mediates the clearance of apoptotic neurons via phagocytosis without inducing apparent neuroinflammation (Takahashi et al., 2005). TREM2 is essential for retaining a fully mature disease-associated microglia (DAM) profile and for supporting the microglial response to Aβ-plaque-induced pathology (Keren-Shaul et al., 2017; Krasemann et al., 2017). TREM2-dependent response requires the induction of the phosphorylation of tyrosine residues within the ITAM motifs of DAP12, by Src-family kinases and downstream signaling cascades including AKT and mTOR (Kober et al., 2017; Jay et al., 2017). Despite extensive investigation, we currently do not have a complete mechanistic understanding of microglial TREM2 signaling, due in part to the lack of a comprehensive knowledge of TREM2-bearing molecular complex(es) in microglial cells. Most of the focus has been on the interaction of TREM2 with DAP12, which serves as a communal platform for downstream signaling. In addition to TREM2,

DAP12 has been shown to serve as a common hub that interacts with numerous
receptors (Lanier, 2009; Haure-Mirande 2022). Given the complexity and high disease
relevance of TREM2-associated biological processes, it is conceivable that additional
molecular components may interact with TREM2 and contribute substantially to the
regulation of relevant microglial functions.

We set out to identify novel components of TREM2-harboring protein complex(es), not
yet implicated in TREM2 signaling, using an unbiased *in situ* proteomics approach. To
isolate protein complexes of interest and analyzed the components present in the
complex, conventional procedures involve detergent lysis, subcellular fractionation, and
affinity pull-down followed by mass spectrometry, which often produces false positives or
fails to discover weak or unstable interactions (Wilson and Nairn 2018; Mehta and
Trinkle-Mulcahy 2018; Bludau and Aebersold 2020). We used TurboID-based "proximity
labeling (PL)" technique to investigate the protein-protein interactions in living microglial
cells (Lam et al., 2015; Han and Ting, 2018; Lee et al., 2019; Qin et al., 2021; Choi and
Rhee, 2022). TurboID, an engineered biotin ligase, uses ATP to convert biotin into biotin-
AMP, a reactive intermediate that covalently labels biotin to the proximal proteins (Doerr,
2018; Brandon et al., 2018). Recently, we improved the TurboID-based PL further and
performed 'super resolution PL', which involves the highly selective capture of
biotinylated peptides, allowing the detection the TurboID-mediated biotinylation event at
the single amino acid level (Lee et al., 2023).

Using the human microglial cells stably expressing TurboID-fused to the cytoplasmic
region of TREM2, we performed super resolution proximity labeling experiments followed
by mass spectrometry analysis, and successfully identified and validated 47 putative
TREM2 interacting proteins. Interestingly, among these proteins, 21 proteins are found to
localize to the subcellular compartment known as the mitochondria-ER contacts (MERCs)
(Vance 1990; 2014; Lee et al., 2018). Biochemically isolated MERCs are also referred to
as mitochondrial associated membranes (MAMs; Vance 2014). MERCs have been
implicated in many functional roles associated with TREM2, including lipid transport,
mitochondrial energy metabolism, Ca2+ handling and immune sensing (Rieusset, 2018;
Tubbs and Rieusset, 2017), and have also been suggested to be a pathological hub for
several neurodegenerative diseases, including AD (Area-Gomez et al., 2018; Bose and
Beal, 2016; Paillusson et al., 2016; Johri and Chandra, 2021; Area-Gomez and Schon,
2017). Given the physical and/or functional association of TREM2 to MERC-related
cellular activities, we investigated the effects of TREM2 deficiency on the cell biological
changes of MERCs in the human induced pluripotent stem cells (iPSC)-derived microglia
(iMGL) (Abud et al., 2017; McQuade et al., 2018). TREM2 deficiency leads to the
ultrastructural alteration of MERCs, further supporting a functional association of TREM2
to MERCs. Our finding also raises the possibility that TREM2-mediated microglial
functions relevant to AD may be attributed to the aberrant regulation of MERCs in
microglia.

Results
Proximity labeling activity of TREM2-TurboID in human microglial cell line

For enzyme-based proximity labeling (PL), we selected TurboID, one of the engineered
biotin ligases that is suitable for protein-protein interaction mapping due to the high
catalytic activity as compared to other PL enzymes (Branon 2018). We generated
mammalian expression plasmids encoding human TREM2 fused to a C-terminal V5 tag
and TurboID (TREM2-V5-TurboID) (Fig. 1A). TREM2-V5-TurboID was transfected into
the human microglial cell line, IM-HM (Mazzeo et al., 2022). Western blotting with
antibodies against V5-tag or TREM2 revealed that TREM2-V5-TurboID is expressed with
the predicted size (Fig. 1B). Confocal fluorescent microscopy showed that TREM2-V5-
TurboID localizes mainly to intracellular compartments as previously reported (Figures

1C; Kleinberger 2014; Park 2015; Yin 2016; Zhao 2017). We next performed confocal
microscopy analysis to detect subcellular distribution of TREM2-V5-TurboID-mediated
biotinylation using streptavidin-647 (Fig. 1C). TREM2-V5-TurboID immunoreactivity and
biotin-labeled signals appear to co-localize strongly, suggesting that the biotin labeling
occurs at subcellular compartments in close proximity to TREM2-V5-TurboID. TurboID
conjugated to nuclear export signal and V5 tag (V5-TurboID-NES) was used as a control,
and the biotin signals from V5-TurboID-NES vs TREM2-V5-TurboID occurred in spatially
distinct subcellular sites (fig. S1A). We next biochemically validated PL activity of
TREM2-V5-TurboID in IM-HM cells. Upon incubation with exogenous biotin, a wide range
of biotinylated proteins were detected in TREM2-V5-TurboID-expressing cells (fig. S1B).
The biotin-labeled proteins were then analyzed by Western blotting using streptavidin-
HRP. Distinct patterns of biotinylated proteins were observed in the samples from the V5-
TurboID-NES-expressing IM-HM cells (fig. S1C).

Proximal proteomics of TREM2 in living microglial cells

We established stable IM-HM transfectants expressing TREM2-V5-TurboID and
performed 'super resolution PL' (Fig. 1A; Lee at al. 2023). The samples were subjected to
streptavidin pull-down to enrich biotin-conjugated proteins. We then performed on-bead
trypsin digestion, allowing direct mass spectrometry analysis of the peptide harboring
biotin-conjugated lysine (K+226) (Lee et al., 2023). This approach removes the bulk of
unlabeled peptides, therefore drastically improving detection sensitivity and reducing
false positive results. After verifying the biotin labeling, biotin-labeled peptides were
analyzed by mass spectrometry as previously described (Lee et al., 2019; 2023; Kwak et
al., 2020). By statistically assessing the relative enrichment of the biotinylated peptides
from TREM2-V5-TurboID relative to the control (V5-TurboID-NES), we identified 47
proteins as components of a TREM2 interactome (Fig. 2A).

Analysis and validation of TREM2 interactome

Quantification of biotinylated peptides derived from the TREM2 proximal proteins was
performed as previously described (Lee et al., 2019). Our TREM2 interactome analysis
revealed a total of 47 proteins (Fig. 2A). To our surprise, among the 47 identified proteins,
we found that 21 TREM2-proximal proteins are known components of MERCs (Fig. 2, A
and B). These MERC components were identified in a previous study via ER-OMM
contact-dependent localization studies between ER and the outer mitochondrial
membrane (OMM) (Kwak et al., 2020). Quantitative proteomics allowed us to determine
the MS signal intensity of MERC components vs. non-MERC components (fig. S2). Our
assessment of the TREM2-TurboID interactome reveals novel TREM2 interactors in the
MERCs of a human microglial cell line. We summarized the TREM2 interactome and their
corresponding subcellular sites (Fig. 2C). To validate the physical proximity of TREM2-
V5-TurboID and identified proteins, we performed a proximity-dependent biotinylation
assay using the expression plasmid encoding each "hit" protein (fig. S2A). An expression
construct encoding TREM2-V5-TurboID was co-transfected with each expression "hit"
construct in 293 cells and subjected to in situ biotinylation assay for validation. This assay
successfully validated the interaction between TREM2-V5-TurboID and the well-known
TREM2 interactor, DAP12, as well newly discovered interactors such as CISD2 and
CKAP4 (fig. S1B). PL-mediated biotin incorporation is shown to be quantitative (Han and
Ting, 2018; Choi and Rhee, 2022). DAP12 exhibited the highest levels of biotin
incorporation as compared to other identified components of TREM2 interactome (fig.
S1C). Our 47 confirmed TREM proximal proteins were found to be positive. We need to
add the description for old fig S2 (new fig. S3)!

TREM2 localization to the MERCs

We next investigated the subcellular localizations of TREM2-proximal proteins. To
precisely map the MERCs, we used 'Contact-ID', in which two inactive fragments of BioID
(a PL enzyme), are reconstituted via protein-protein interaction across two distal
membranes and become catalytically active (Fig. 3A; Kwak et al., 2020). The Contact-ID
system consists of the N-G78 (B1) fragment of BioID conjugated to the cytosol-facing N
terminus of SEC61B (B1-SEC61B) at the ER membrane and the G79-C (B2) fragments
of BioID conjugated to the C terminus of TOM20 (TOM20-B2) at the OMM. This approach
allows for precise mapping of the MERC contact sites without overlapping distribution of
virtually any MERC components with either the ER or OMM. We transfected TREM2-V5
without TurboID and examine the subcellular distribution of TREM2 vs. Contact-ID-
dependent signals (Fig. 3B). Confocal microscopy analysis showed that the subcellular
localization pattern of TREM2 overlaps substantially to that of the Contact-ID signal.
Biochemical data showed that TREM2 is indeed proximal to the MERC components (Fig.
3C). Please add additional description on F3C here. These data suggest that TREM2
localizes to the MERCs and biotinylation of TREM2 is mediated by the PL activity of
TREM2-TurboID within the MERCs.

TREM2 deficiency increases the thickness of MERCs in iMGL

Prevalent localization of the TREM2 interactome in the MERCs suggests that TREM2
may be required for the proper structural and/or functional regulation of MERCs in human
microglia. Recently, human induced pluripotent stem cell (iPSC)-derived microglial
models (iMGL) for AD became available (Abud et al., 2017; McQuade et al., 2018). We
therefore investigated the cell biological impact of TREM2 deficiency in human iMGL -a
more physiological model system as compared to a microglial cell line. We performed
electron microscopy analysis to test if TREM2 deficiency affects the ultrastructure of
MERCs in iMGL (Fig. 4). Consistent with the previous mouse *in vivo* studies (Nugent et
al., 2020), there is a marked increase of lipid droplets (LDs) in TREM2-deficient iMGL as
compared to wild-type iMGL (fig. S4). We next performed morphological characterization
of MERCs in wild-type vs. TREM2 deficient iMGL (Fig. 4A). The established key
structural parameter of MERCs includes the width of the gap (10-30 nm distance)
between cytosolic sites of ER membrane and OMM (MERC thickness; Csordás et al.,
2006; Giacomello and Pellegrini, 2016). MERC thickness is a regulated parameter
strongly associated with metabolic state of the cells. Cell function and survival relies on
the maintenance of a proper spacing between the ER and mitochondria (Csordás et al.,
2006; Giacomello and Pellegrini, 2016). We found that overall MERC thickness is greatly
increased in TREM2 knockout iMGL as compared to wild-type (Fig 4, B and C). To
ensure that the observed increases of the MERC thickness in the TREM2-deficient iMGL
is specific to cell biological alterations owing to TREM2, we next tested whether re-
introduction of TREM2 into the TREM2-deficient iMGL could restore the observed MERC
alterations associated with TREM2 deficiency (Fig. 5). Lentiviral delivery of TREM2 in the
TREM2 deficient iMGL leads to a decrease in the MERC thickness as compared to
iMGLs infected with control lentiviral particles (Fig. 5). In addition, the MERC thickness
increased occurrence of LDs, a prominent phenotype of TREM2-deficient iMGL, was also
rescued by the introduction of wild-type human TREM2 (fig. S5). These data suggest that
the increased MERC thickness in the TREM2-deficient cells is a TREM2- specific cell
biological change.

Effects of R47H TREM2 on the MERC thickness

The loss-of-function variants of TREM2, including R47H, are associated with an
increased risk of AD (Guerreiro et al., 2013; Jonsson et al., 2013). To test whether the
altered MERC thickness is associated with hypomorphic TREM2 R47H variant, we next
analyzed the MERC thickness of the iMGL derived from human iPSC carrying the R47H
TREM2 variant (Figure 6A). R47H TREM2 loss of function

We determined the distance of ER-to-mitochondria of iMGL carrying TREM2 R47H
mutant, as compared to isogenic wild-type control. The MERC thickness was significantly
elevated in the TREM2 R47H iMGL relative to control (Fig. 6B). The MERC thickness
phenotype was less profound than that observed in the TREM2 full knockout iMGL,
probable due to the partial loss-of-function characteristics of TREM2 R47H (Cheng-
Hathaway et al., 2018; Sudom et al., 2018). The occurrence of LDs was also increased in
the iMGLs carrying R47H TREM2 as compared to isogenic wild-type (fig. S5).

Enrichment of TREM2 interactome genes across microglia transcriptional states

To assess potential involvement of the TREM2 interactome with AD, we next leveraged
the single-cell RNA-sequencing (scRNA-seq) data from human AD samples. Specifically,
we analyzed whether the transcriptomic profiles of the genes encoding the MERC
TREM2 interactome associate with specific activation state(s) among transcriptionally
distinct sub-populations of human microglia, including homeostatic, activated, IFN and
IL1B (Fig. 7). The scRNA-seq microglia data was collected from human donors (parietal
cortex from 67 donors from the Knight ADRC and DIAN brain banks; N=16,849) (Brase et
al., 2023). Curated marker genes were employed to label the cell types based on the
overall expression profile of the nuclei as previously described (Da Masquita et al., 2021).
Detailed clinical data and genetic characterization of the collected microglia are described
in Table 1. Here, we used four different microglia subtypes based on their transcriptomic
profiles and corresponding functional/activation status: mic-resting (n=7001), mic-
activated (n=3653), mic-reduced (n=2078) and mic-inflammatory (n=1668) (Fig. 7, A and
B). The average expression of the genes encoding the TREM2 interactome in MERC
(MERC genes) into each microglia transcriptional states by Control vs. TREM2 reduced
(R47H, R62H, H157Y variants) genetic status was done (Fig. 7C). A MERC genes score
was also calculated for microglia cell by running Seurat's AddModuleScore. The
enrichment of TREM2 interactome MAM genes into Control and TREM2 reduced
microglia cell was also evaluated (Fig. 7D). Further, the significant pairwise differences
into each Control and TREM2 reduced transcriptional states was calculated using the
linear mixed model. moduleScore ~ Cluster + (1|Subject) (Fig. 7E).

Discussion

Protein interaction networks underlie the biological processes and pathological
progression of human diseases (Rees et al., 2015; Bludau and Aebersold, 2020). Given
the critical role for TREM2 in microglial biology as well as the pathophysiology of AD, a
comprehensive understanding of the protein-protein interaction networks of TREM2
should provide new insights on TREM2 functions in microglia. We utilized an TurboID-
catalyzed PL approach combined with mass spectrometry to determine the TREM
interactome in live microglial cells. The PL approach overcomes some inherent limitations
of the traditional methods associated with affinity pull-down or co-immunoprecipitation,
which often involve a post-lysis manipulation of the starting material (e.g. detergent
extraction, subcellular fractionation). PL can detect native protein-protein interactions in
live cells, and is thereby highly suitable for mapping target protein interactome with a
greater specificity. Among a number of PL enzymes, TurboID was chosen because of its
catalytic efficiency, simplicity and the use non-toxic biotin substrate (Branon et al., 2019).
Here we identified novel proteins proximal to TREM2-TurboID, including those localized
predominantly to MERCs, unveiling the connection between TREM2 and MERCs.

MERC is a dynamic and plastic structure that can undergo remodeling in response to
metabolic change and stress (Rowland 2012; Csordás et al., 2006). The thickness of

MERC, the width of the cleft that separates the ER and OMM, is considered a regulated structural parameter that changes depending on the metabolic state of the cells (Giacomello and Pellegrini, 2016). Changes in the width of the cleft in MERC influence the lipid transfer process across two membranes (Giacomello and Pellegrini, 2016). We have demonstrated that deficiency or hypomorphic R47H mutation of TREM2 lead to increased thickness of MERCs in iMGs (Fig. 4 and 6). In addition to our novel observation of the MERC alterations, our results show the increased occurrence of LDs in TREM2 deficient iMGL (fig. S4). This is consistent with the previous report demonstrating the increased accumulation of cholesterol ester (e.g. CE20:4 and CE22:6), the major cholesterol metabolites stored in LDs, in both mouse brain and cultured human myeloid cells (Nugent et al., 2020). Since MERC has been shown to play an important role in the biogenesis of LD (Liu et al., 2015; Freyre et al., 2019; Choudhary et al., 2021; Guyard et al., 2022), it is conceivable that altered MERC in the TREM2 deficient iMGL may be attributed to the aberrant accumulation of LDs in iMGL. When cell conditions promote active expansion of LDs, the synthesis and trafficking of new phospholipids are regulated in the area between mitochondria and ER (Olzmann et al., 2019). According to a report by Area-Gomez et al., the ACAT protein relating LD generation was more abundant in the mitochondria-associated ER membranes (MAMs) fraction and had approximately 3 times higher enzymatic activity compared to the ER and mitochondrial fractions (Area-Gomez et al., 2012). Collectively, virtually all known TREM2 functions including metabolic regulation of lipids and Ca2+, as well as inflammatory activation, are linked intimately to those of MERC, substantiating the idea that TREM2 is an important factor for proper MERC functions.

Aberrant MERCs have been reported in cellular models of AD. Gene deletion and FAD mutations of familial AD-associated presenilin genes (PSEN1 and PSEN2) greatly increases the MERC thickness (50-200 nm) in fibroblasts (Area-Gomez et al., 2012). Presenilin 2 has been shown to regulate Ca2+ crosstalk across the MERCs (Zampese et al., 2011). PS1 has also been shown to interact with TREM2 (Zhao et al., 2017). In addition to AD, MERCs have been implicated in other neurodegenerative diseases, such as Parkinson's disease (PD) (Area-Gomez et al., 2018; Bose and Beal, 2016; Paillusson et al., 2016; Johri and Chandra, 2021; Area-Gomez and Schon, 2017). Occurrence of the microglial TREM2 interactome in the MERCs is particularly relevant to a known MERC functions involving immune activation. MERC is shown to serve as a signaling hub for NLRP3 inflammasome activation (Horng et al., 2014; Zhou et al., 2020). TREM2 is a direct regulator of the NLRP3 inflammasome and the AD-linked R47H TREM2 variant is not capable of activating the inflammasome (Cosker et al., 2021; Li et al., 2021; Jiang et al., 2022; Wang et al., 2022). These observations further substantiate the idea of functional association of TREM2 with MERC.

In our previous study, we developed Contact-ID, a split-pair BioID system used to identify the protein components of MERCs (Kwak et al., 2020). Since all known MERC markers show overlapping subcellular distribution in either the ER or OMM, the Contact-ID approach allows precise mapping of ER-OMM contact-dependent localization (Fig. 3), indicting co-localization of TREM2 with identified MERC components in the TREM2 interactome. In addition to the components of MERC, our PL interactome study revealed newly discovery TREM2-interacting proteins with distinct subcellular localizations (Fig. 2C), suggesting potential mediators or modulators of cellular functions of TREM2 in microglial cells. For instance, anoctamin 6 encoded by ANO6 is an essential component for Ca^{2+}-dependent exposure of phosphatidylserine (PS) on the cell surface (Yang et al., 2012; Kunzelmann et al., 2014). PS exposure is considered to be one of the "eat-me" signals for the engulfment by immune cells (Lemke et al., 2019; Scott-Hewitt et al., 2020) and ANO6 has been shown to increase microglial phagocytosis of neurons (Batti et al., 2016; Zhang et al., 2020). Interestingly, ANO6 deficiency reduces activation of the microglial NLRP3 inflammasome in the brain of AD mouse model (Cui et al., 2023),

suggesting the functional link of ANO6 with TREM2 in the regulation of NLRP3 inflammasome (Li et al., 2021; Jiang et al., 2022). CDK5 regulatory subunit Associated protein 1-Like 1 (CDKAL1) has been implicated in the pathogenesis of type2 diabetes (Ghosh et al., 2022) and a family-based GWAS meta-analysis revealed genetic association of two SNPs in the CDKAL1 gene (Herold et al., 2016). ITPRIP gene (The gene encoding inositol 1,4,5-trisphosphate receptor interacting protein) has also been suggested as an AD risk gene (Simino et al., 2017). AUP1 and REEP5 are known to play a key role in LD biogenesis (Robichaud et al., 2021; Chen et al., 2022), consistent with the functional role for TREM2 in the LD biogenesis. Further studies are needed for the characterization of the components of the TREM2 interactome in microglial cells, including other non-MERC proteins. Future studies would also need to elaborate if the observed TREM2-associated MERCs occurs in vivo especially in the context of mouse models. Collectively, our current study provides new insights on TREM2 functions and serves as a foundation for further research aimed at deciphering new molecular players in microglial TREM2 functions.

Materials and Methods

Culture of iMGL
The wild type iMGL, TREM2 deficient iMGL, and iMGL derived from human iPSC carrying R47H TREM2 variant (Fujifilm Cellular dynamics iCell microglia, AHN, TREM2 R47H, #11969) were cultured in iCell microglia media kit (Fujifilm cellular dynamics, USA, #R1204) as recommended by a manufacturer. Cells were maintained on 0.07 % Poly ethyleneimine solution coated plate in suspension. For re-introduction of TREM2 into the TREM2-deficient iMGL, TREM2-deficient iMGL was infected with two different lentiviral particles (Control lentiviral particles (CH404_empty_GFP reporter_pDual) and TREM2 delivery lentiviral particles (CH400_ss-V5-TREM2-twin strep tag_GFP reporter_pDual)).

Transmission electron microscopy (TEM) analysis
Cells were pre-fixed with 2% paraformaldehyde/2.5% glutaraldehyde and post-fixed in 2% Osmium tetroxide (OsO4; EMS, USA). After block staining with 1.5% potassium ferrocyanide (sigma, USA), 1% thiocarbohydrazide (TCH; TCI, Japan), 2% OsO4, and 2% uranyl acetate (EMS, USA), dehydration with a graded ethanol series (20%, 50%, 70%, 90%, and 100%) and infiltration with an epoxy resin (30, 50, 70, and 100%) were done. Then, samples were embedded with an EMBed-812 embedding kit (EMS, USA) following the manufacturer's instructions. Cells were sectioned at 60 nm using an ultramicrotome (UC7; Leica Microsystems, Germany) and sections were mounted on copper slot grids with a formva film. The sections were imaged by the Tecnai G2 microscope with a US1000X-P camera 200 (Thermo Fisher Scientific, USA) at an accelerating voltage of 120 kV.

Analysis of MERCs in iMGL
Ultrastructural analysis of MERCs in iMGL was performed as previously described (Giacomello et al., 2016; Raturi et al., 2016; Bartok et al., 2019) The thickness of MERCs with less than 30 nm between the mitochondria and ER was measured in 10 regions of MERCs. MERCs thickness and length in each group were determined in nanometers and averaged. Data were analyzed using a Image J software. Statistical analysis was performed using GraphPad Prism 8 software (GraphPad Software Inc., LaJolla, CA). The statistical significance was $p < 0.001$.

Analysis of lipid droplets in iMGL
To identify the number of lipid droplets (LDs), cells were stained with Lipi-red (Dojindo, Japan; 1 umol/L) and Hoechst solution (Sigma, USA; 1 umol/L). Live cells were imaged

393 under a confocal light microscopy (Ti-RCP, Nikon, Japan). The ultrastructure of LDs was
394 observed under a TEM and the number of LDs was measured by Image J software.
395
396
397

References

Abud EM, Ramirez RN, Martinez ES, Healy LM, Nguyen CHH, Newman SA, Yeromin AV, Scarfone VM, Marsh SE, Fimbres C, Caraway CA, Fote GM, Madany AM, Agrawal A, Kayed R, Gylys KH, Cahalan MD, Cummings BJ, Antel JP, Mortazavi A, Carson MJ, Poon WW, Blurton-Jones M. iPSC-Derived Human Microglia-like Cells to Study Neurological Diseases. Neuron. 2017 Apr 19;94(2):278-293. PMID: 28426964Area-Gomez E, de Groof A, Bonilla E, Montesinos J, Tanji K, Boldogh I, Pon L, Schon EA. A key role for MAM in mediating mitochondrial dysfunction in Alzheimer disease. Cell Death Dis. 2018 Feb 28;9(3):335. PMID: 29491396

Area-Gomez E, Del Carmen Lara Castillo M, Tambini MD, Guardia-Laguarta C, de Groof AJ, Madra M, Ikenouchi J, Umeda M, Bird TD, Sturley SL, Schon EA. Upregulated function of mitochondria-associated ER membranes in Alzheimer disease. EMBO J. 2012 Nov 5;31(21):4106-23. PMID: 22892566

Area-Gomez E, Schon EA. On the Pathogenesis of Alzheimer's Disease: The MAM Hypothesis. FASEB J. 2017 Mar;31(3):864-867. PMID: 28246299

Atagi Y, et al. Apolipoprotein E is a ligand for triggering receptor expressed on myeloid cells 2 (TREM2) J. Biol. Chem. 2015;290:26043–26050.

Bailey CC, DeVaux LB, Farzan M. The triggering receptor expressed on myeloid cells 2 binds apolipoprotein E. J. Biol. Chem. 2015;290:26033–26042.

Bartok, A., et al. IP3 receptor isoforms differently regulate ER-mitochondrial contacts and local calcium transfer. Nat Commun 10, 3726 (2019).

Batti L, Sundukova M, Murana E, Pimpinella S, De Castro Reis F, Pagani F, Wang H, Pellegrino E, Perlas E, Di Angelantonio S, Ragozzino D, Heppenstall PA. TMEM16F Regulates Spinal Microglial Function in Neuropathic Pain States. Cell Rep. 2016 Jun 21;15(12):2608-15. PMID: 27332874

Bludau I, Aebersold R. Proteomic and interactomic insights into the molecular basis of cell functional diversity. Nat Rev Mol Cell Biol. 2020 Jun;21(6):327-340. PMID: 32235894

Bose A, Beal MF. Mitochondrial dysfunction in Parkinson's disease. J Neurochem. 2016 Oct;139 Suppl 1:216-231. PMID: 27546335

Branon TC, Bosch JA, Sanchez AD, Udeshi ND, Svinkina T, Carr SA, Feldman JL, Perrimon N, Ting AY. Efficient proximity labeling in living cells and organisms with TurboID. Nat Biotechnol. 2018 Oct;36(9):880-887. PMID: 30125270

Brase L, You SF, D'Oliveira Albanus R, Del-Aguila JL, Dai Y, Novotny BC, Soriano-Tarraga C, Dykstra T, Fernandez MV, Budde JP, Bergmann K, Morris JC, Bateman RJ, Perrin RJ, McDade E, Xiong C, Goate AM, Farlow M; Dominantly Inherited Alzheimer Network (DIAN); Sutherland GT, Kipnis J, Karch CM, Benitez BA, Harari O. Single-nucleus RNA-sequencing of autosomal dominant Alzheimer disease and risk variant carriers. Nat Commun. 2023 Apr 21;14(1):2314. PMID: 37085492

Chen F, Yan B, Ren J, Lyu R, Wu Y, Guo Y, Li D, Zhang H, Hu J. FIT2 organizes lipid droplet biogenesis with ER tubule-forming proteins and septins. J Cell Biol. 2021 May 3;220(5):e201907183. PMID: 33861319

Cheng-Hathaway PJ, Reed-Geaghan EG, Jay TR, Casali BT, Bemiller SM, Puntambekar SS, von Saucken VE, Williams RY, Karlo JC, Moutinho M, Xu G, Ransohoff RM, Lamb BT, Landreth GE. The Trem2 R47H variant confers loss-of-function-like phenotypes in Alzheimer's disease. Mol Neurodegener. 2018 Jun 1;13(1):29. PMID: 29859094

Choi CR, Rhee HW. Proximity labeling: an enzymatic tool for spatial biology. Trends Biotechnol. 2022 Feb;40(2):145-148. PMID: 34663510

Choudhary V, Schneiter R. A Unique Junctional Interface at Contact Sites Between the Endoplasmic Reticulum and Lipid Droplets. Front Cell Dev Biol. 2021 Apr 8;9:650186. PMID: 33898445

Colonna M. The biology of TREM receptors. Nat Rev Immunol. 2023 Feb 7:1-15. PMID: 36750615

Cosker K, Mallach A, Limaye J, Piers TM, Staddon J, Neame SJ, Hardy J, Pocock JM. Microglial signalling pathway deficits associated with the patient derived R47H TREM2 variants linked to AD indicate inability to activate inflammasome. Sci Rep. 2021 Jun 25;11(1):13316. PMID: 34172778

Csordás G., Renken C., Várnai P., Walter L., Weaver D., Buttle K. F., et al. (2006). Structural and functional features and significance of the physical linkage between ER and mitochondria. J. Cell. Biol. 174, 915–921. PMID: 16982799

Cui ZQ, Hu XY, Yang T, Guan JW, Gu Y, Li HY, Zhang HY, Xiao QH, Sun XH. TMEM16F may be a new therapeutic target for Alzheimer's disease. Neural Regen Res. 2023 Mar;18(3):643-651. PMID: 36018189

Da Mesquita S, Papadopoulos Z, Dykstra T, Brase L, Farias FG, Wall M, Jiang H, Kodira CD, de Lima KA, Herz J, Louveau A, Goldman DH, Salvador AF, Onengut-Gumuscu S, Farber E, Dabhi N, Kennedy T, Milam MG, Baker W, Smirnov I, Rich SS; Dominantly Inherited Alzheimer Network; Benitez BA, Karch CM, Perrin RJ, Farlow M, Chhatwal JP, Holtzman DM, Cruchaga C, Harari O, Kipnis J. Meningeal lymphatics affect microglia responses and anti-Aβ immunotherapy. Nature. 2021 May;593(7858):255-260. PMID: 33911285

Doerr A. Proximity labeling with TurboID. Nat Methods. 2018 Oct;15(10):764. doi: 10.1038/s41592-018-0158-0. PMID: 30275580

Freyre CAC, Rauher PC, Ejsing CS, Klemm RW. MIGA2 Links Mitochondria, the ER, and Lipid Droplets and Promotes De Novo Lipogenesis in Adipocytes. Mol Cell. 2019 Dec 5;76(5):811-825.e14. PMID: 31628041

Fu WY, et al. Targeting Neuroinflammation as a Therapeutic Strategy for Alzheimer's Disease: Mechanisms, Drug Candidates, and New Opportunities. ACS Chem Neurosci. 2019. PMID:30221933

Ghosh C, Das N, Saha S, Kundu T, Sircar D, Roy P. Involvement of Cdkal1 in the etiology of type 2 diabetes mellitus and microvascular diabetic complications: a review. J Diabetes Metab Disord. 2022 Jan 13;21(1):991-1001. PMID: 35673487

Giacomello M, Pellegrini L. The coming of age of the mitochondria-ER contact: a matter of thickness. Cell Death Differ. 2016 Sep 1;23(9):1417-27. PMID: 27341186

Griciuc A, Tanzi RE. The role of innate immune genes in Alzheimer's disease. Curr Opin Neurol. 2021 Apr 1;34(2):228-236. PMID: 33560670

Guerreiro R, Wojtas A, Bras J, Carrasquillo M, Rogaeva E, Majounie E, Cruchaga C, Sassi C, Kauwe JS, Younkin S, Hazrati L, Collinge J, Pocock J, Lashley T, Williams J, Lambert JC, Amouyel P, Goate A, Rademakers R, Morgan K, Powell J, St George-Hyslop P, Singleton A, Hardy J; Alzheimer Genetic Analysis Group. TREM2 variants in Alzheimer's disease. N Engl J Med. 2013 Jan 10;368(2):117-27. PMID: 23150934

Guyard V, Monteiro-Cardoso VF, Omrane M, Sauvanet C, Houcine A, Boulogne C, Ben Mbarek K, Vitale N, Faklaris O, El Khallouki N, Thiam AR, Giordano F. ORP5 and ORP8 orchestrate lipid droplet biogenesis and maintenance at ER-mitochondria contact sites. J Cell Biol. 2022 Sep 5;221(9):e202112107. PMID: 35969857

Han S, Li J, Ting AY. Proximity labeling: spatially resolved proteomic mapping for neurobiology. Curr Opin Neurobiol. 2018 Jun;50:17-23. PMID: 29125959

Hansen DV, Hanson JE, Sheng M. Microglia in Alzheimer's disease. J Cell Biol. 2018 Feb 5;217(2):459-472. PMID: 29196460

Haure-Mirande JV, Audrain M, Ehrlich ME, Gandy S. Microglial TYROBP/DAP12 in Alzheimer's disease: Transduction of physiological and pathological signals across TREM2. Mol Neurodegener. 2022 Aug 24;17(1):55. PMID: 36002854

Herold C, Hooli BV, Mullin K, Liu T, Roehr JT, Mattheisen M, Parrado AR, Bertram L, Lange C, Tanzi RE. Family-based association analyses of imputed genotypes reveal genome-wide significant association of Alzheimer's disease with OSBPL6, PTPRG, and PDCL3. Mol Psychiatry. 2016 Nov;21(11):1608-1612. PMID: 26830138

Horng T. Calcium signaling and mitochondrial destabilization in the triggering of the nlrp3 inflammasome. Trends Immunol. 2014;35:253–261.PMID: 24646829

Jay TR, von Saucken VE, Landreth GE. TREM2 in Neurodegenerative Diseases. Mol Neurodegener. 2017 Aug 2;12(1):56. PMID: 28768545

Jiang W, Liu F, Li H, Wang K, Cao X, Xu X, Zhou Y, Zou J, Zhang X, Cui X. TREM2 ameliorates anesthesia and surgery-induced cognitive impairment by regulating mitophagy and NLRP3 inflammasome in aged C57/BL6 mice. Neurotoxicology. 2022 May;90:216-227. PMID:35447280

Johri A, Chandra A. Connection Lost, MAM: Errors in ER-Mitochondria Connections in Neurodegenerative Diseases. Brain Sci. 2021 Oct 28;11(11):1437. PMID: 34827436

Jonsson T, Stefansson H, Steinberg S, Jonsdottir I, Jonsson PV, Snaedal J, Bjornsson S, Huttenlocher J, Levey AI, Lah JJ, Rujescu D, Hampel H, Giegling I, Andreassen OA, Engedal K, Ulstein I, Djurovic S, Ibrahim-Verbaas C, Hofman A, Ikram MA, van Duijn CM, Thorsteinsdottir U, Kong A, Stefansson K. Variant of TREM2 associated with the risk of Alzheimer's disease. N Engl J Med. 2013 Jan 10;368(2):107-16. PMID: 23150908

Jorfi M, Maaser-Hecker A, Tanzi RE. The neuroimmune axis of Alzheimer's disease. Genome Med. 2023 Jan 26;15(1):6. PMID: 36703235

Keren-Shaul H, et al. A unique microglia type associated with restricting development of Alzheimer's disease. Cell. 2017;169:1276–1290.e17. PMID: 28602351

Kim DI, Roux KJ. Filling the Void: Proximity-Based Labeling of Proteins in Living Cells. Trends Cell Biol. 2016 Nov;26(11):804-817. PMID: 27667171

Kleinberger G, Yamanishi Y, Suarez-Calvet M, Czirr E, Lohmann E, Cuyvers E et al. TREM2 mutations implicated in neurodegeneration impair cell surface transport and phagocytosis. Sci Transl Med 2014; 6: 243ra86. [PubMed] [Google Scholar] PMID: 24990881

Klunemann, H. H. et al. The genetic causes of basal ganglia calcification, dementia, and bone cysts: DAP12 and TREM2. Neurology 64, 1502–1507 (2005).

Kober DL, Brett TJ. TREM2-Ligand Interactions in Health and Disease. J Mol Biol. 2017 Jun 2;429(11):1607-1629. PMID: 28432014

Krasemann S, et al. The TREM2–APOE pathway drives the transcriptional phenotype of dysfunctional microglia in neurodegenerative diseases. Immunity. 2017;47:566–581. PMID: 28930663

Kunzelmann K, Nilius B, Owsianik G, Schreiber R, Ousingsawat J, Sirianant L, Wanitchakool P, Bevers EM, Heemskerk JW. Molecular functions of anoctamin 6 (TMEM16F): a chloride channel, cation channel, or phospholipid scramblase? Pflugers Arch. 2014 Mar;466(3):407-14. doi: 10.1007/s00424-013-1305-1. PMID: 23748496

Kwak C, Shin S, Park JS, Jung M, Nhung TTM, Kang MG, Lee C, Kwon TH, Park SK, Mun JY, Kim JS, Rhee HW. Contact-ID, a tool for profiling organelle contact sites, reveals regulatory proteins of mitochondrial-associated membrane formation. Proc Natl Acad Sci U S A. 2020 Jun 2;117(22):12109-12120. PMID: 32414919

Mehta V, Trinkle-Mulcahy L. Recent advances in large-scale protein interactome mapping. F1000Res. 2016 Apr 29;5. PMID: 27158474

Lam SS, Martell JD, Kamer KJ, Deerinck TJ, Ellisman MH, Mootha VK, Ting AY. Directed evolution of APEX2 for electron microscopy and proximity labeling. Nat Methods. 2015 Jan;12(1):51-4. PMID: 25419960

Lanier LL. DAP10- and DAP12-associated receptors in innate immunity. Immunol Rev. 2009 Jan;227(1):150-60. PMID: 19120482

Lee YB, Jung M, Kim J, Charles A, Christ W, Kang J, Kang MG, Kwak C, Klingström J, Smed-Sörensen A, Kim JS, Mun JY, Rhee HW. Super-resolution proximity labeling reveals anti-viral protein network and its structural changes against SARS-CoV-2 viral proteins. Cell Rep. 2023 Jul 20;42(8):112835. PMID: 37478010

Lee S, Min KT. The Interface Between ER and Mitochondria: Molecular Compositions and Functions. Mol Cells. 2018 Dec 31;41(12):1000-1007. PMID: 30590907

Lee SY, Seo JK, Rhee HW. Direct Identification of Biotinylated Proteins from Proximity Labeling (Spot-BioID). Methods Mol Biol. 2019;2008:97-105. PMID: 31124091

Lemke G. How macrophages deal with death. Nat Rev Immunol. 2019 Sep;19(9):539-549. PMID: 31019284

Leng F, Edison P.Neuroinflammation and microglial activation in Alzheimer disease: where do we go from here? Nat Rev Neurol. 2021 Mar;17(3):157-172. PMID: 33318676

Lessard CB, et al. High-affinity interactions and signal transduction between Aβ oligomers and TREM2. EMBO Mol. Med. 2018;10:e9027. PMID: 30341064

Li Y, Long W, Gao M, Jiao F, Chen Z, Liu M, Yu L. TREM2 Regulates High Glucose-Induced Microglial Inflammation via the NLRP3 Signaling Pathway. Brain Sci. 2021 Jul 7;11(7):896. PMID: 34356130

Liu L, Zhang K, Sandoval H, Yamamoto S, Jaiswal M, Sanz E, Li Z, Hui J, Graham BH, Quintana A, Bellen HJ. Glial lipid droplets and ROS induced by mitochondrial defects promote neurodegeneration. Cell. 2015 Jan 15;160(1-2):177-90. PMID: 25594180

Matarin M, Salih DA, Yasvoina M, Cummings DM, Guelfi S, Liu W, Nahaboo Solim MA, Moens TG, Paublete RM, Ali SS, Perona M, Desai R, Smith KJ, Latcham J, Fulleylove M, Richardson JC, Hardy J, Edwards FA. A genome-wide gene-expression analysis and database in transgenic mice during development of amyloid or tau pathology. Cell Rep. 2015 Feb 3;10(4):633-44.

Mazzeo A, Porta M, Beltramo E. Characterization of an Immortalized Human Microglial Cell Line as a Tool for the Study of Diabetic Retinopathy. Int J Mol Sci. 2022 May 20;23(10):5745. PMID: 35628555

McQuade A, Blurton-Jones M. Microglia in Alzheimer's Disease: Exploring How Genetics and Phenotype Influence Risk. J Mol Biol. 2019 Apr 19;431(9):1805-1817. PMID: 30738892

McQuade A, Coburn M, Tu CH, Hasselmann J, Davtyan H, Blurton-Jones M. Development and validation of a simplified method to generate human microglia from pluripotent stem cells. Mol Neurodegener. 2018 Dec 22;13(1):67. PMID: 30577865

Nugent AA, Lin K, van Lengerich B, Lianoglou S, Przybyla L, Davis SS, Llapashtica C, Wang J, Kim DJ, Xia D, Lucas A, Baskaran S, Haddick PCG, Lenser M, Earr TK, Shi J, Dugas JC, Andreone BJ, Logan T, Solanoy HO, Chen H, Srivastava A, Poda SB, Sanchez PE,

Watts RJ, Sandmann T, Astarita G, Lewcock JW, Monroe KM, Di Paolo G. TREM2 Regulates Microglial Cholesterol Metabolism upon Chronic Phagocytic Challenge. Neuron. 2020 Mar 4;105(5):837-854.e9. PMID: 31902528

Olzmann, J.A.; Carvalho, P. Dynamics and functions of lipid droplets. Nat. Rev. Mol. Cell Biol. 2019, 20, 137–155

Paillusson S, Stoica R, Gomez-Suaga P, Lau DHW, Mueller S, Miller T, Miller CCJ. There's Something Wrong with my MAM; the ER-Mitochondria Axis and Neurodegenerative Diseases. Trends Neurosci. 2016 Mar;39(3):146-157. PMID: 26899735

Paloneva, J. et al. Mutations in two genes encoding different subunits of a receptor signaling complex result in an identical disease phenotype. Am. J. Hum. Genet. 71, 656–662 (2002).

Park JS, Ji IJ, An HJ, Kang MJ, Kang SW, Kim DH et al. Disease-associated mutations of TREM2 alter the processing of N-linked oligosaccharides in the Golgi apparatus. Traffic 2015; 16: 510–518. PMID: 25615530

Qin W, Cho KF, Cavanagh PE, Ting AY. Deciphering molecular interactions by proximity labeling. Nat Methods. 2021 Feb;18(2):133-143. PMID: 33432242

Raturi A., et al. TMX1 determines cancer cell metabolism as a thiol-based modulator of ER-mitochondria Ca2+ flux. J Cell Biol. 2016 Aug 15;214(4):433-44.

Rees JS, Li XW, Perrett S, Lilley KS, Jackson AP. Protein neighbors and proximity proteomics. MCP. 2015;14(11):2848–56. PMID: 26355100

Rieusset J. Mitochondria-associated membranes (MAMs): An emerging platform connecting energy and immune sensing to metabolic flexibility. Biochem Biophys Res Commun. 2018 May 27;500(1):35-44. PMID: 28647358

Robichaud S, Fairman G, Vijithakumar V, Mak E, Cook DP, Pelletier AR, Huard S, Vanderhyden BC, Figeys D, Lavallée-Adam M, Baetz K, Ouimet M. Identification of novel lipid droplet factors that regulate lipophagy and cholesterol efflux in macrophage foam cells. Autophagy. 2021 Nov;17(11):3671-3689. PMID: 33590792

Sarlus H, Heneka MT. Microglia in Alzheimer's disease. J Clin Invest. 2017 Sep 1;127(9):3240-3249. PMID: 28862638

Scott-Hewitt N, Perrucci F, Morini R, Erreni M, Mahoney M, Witkowska A, Carey A, Faggiani E, Schuetz LT, Mason S, Tamborini M, Bizzotto M, Passoni L, Filipello F, Jahn R, Stevens B, Matteoli M. Local externalization of phosphatidylserine mediates developmental synaptic pruning by microglia. EMBO J. 2020 Aug 17;39(16):e105380. PMID: 32657463

Simino J, Wang Z, Bressler J, Chouraki V, Yang Q, Younkin SG, Seshadri S, Fornage M, Boerwinkle E, Mosley TH Jr. Whole exome sequence-based association analyses of plasma amyloid-β in African and European Americans; the Atherosclerosis Risk in Communities-Neurocognitive Study. PLoS One. 2017 Jul 13;12(7):e0180046. PMID: 28704393

Song W, et al. Alzheimer's disease-associated TREM2 variants exhibit either decreased or increased ligand-dependent activation. Alzheimers Dement. 2017;13:381–387. PMID: 27520774

Sudom A, Talreja S, Danao J, Bragg E, Kegel R, Min X, Richardson J, Zhang Z, Sharkov N, Marcora E, Thibault S, Bradley J, Wood S, Lim AC, Chen H, Wang S, Foltz IN, Sambashivan S, Wang Z. Molecular basis for the loss-of-function effects of the Alzheimer's disease-associated R47H variant of the immune receptor TREM2. J Biol Chem. 2018 Aug 10;293(32):12634-12646. PMID: 29794134

Takahashi K, Rochford CD, Neumann H. Clearance of apoptotic neurons without inflammation by microglial triggering receptor expressed on myeloid cells-2. J. Exp. Med. 2005;201:647–657. PMID: 15728241

Tubbs E, Rieusset J. (2017) Metabolic signaling functions of ER-mitochondria contact sites: role in metabolic diseases. J Mol Endocrinol 58, R87-R106. PMID: 27965371

Vance JE. MAM (mitochondria-associated membranes) in mammalian cells: lipids and beyond. Biochim Biophys Acta. 2014 Apr 4;1841(4):595-609. PMID: 24316057

Vance JE. Phospholipid synthesis in a membrane fraction associated with mitochondria. J Biol Chem 1990 May 5;265(13):7248-56. PMID: 2332429

Villegas-Llerena C, Phillips A, Garcia-Reitboeck P, Hardy J, Pocock JM. Microglial genes regulating neuroinflammation in the progression of Alzheimer's disease. Curr Opin Neurobiol. 2016 Feb;36:74-81. PMID: 26517285

Wang Y, Cao C, Zhu Y, Fan H, Liu Q, Liu Y, Chen K, Wu Y, Liang S, Li M, Li L, Liu X, Zhang Y, Wu C, Lu G, Wu M. TREM2/β-catenin attenuates NLRP3 inflammasome-mediated macrophage pyroptosis to promote bacterial clearance of pyogenic bacteria. Cell Death Dis. 2022 Sep 6;13(9):771. PMID: 36068223

Wilson RS, Nairn AC. Cell-Type-Specific Proteomics: A Neuroscience Perspective. Proteomes. 2018 Dec 9;6(4). PMID: 30544872

Wyss-Coray T, Mucke L. Inflammation in neurodegenerative disease--a double-edged sword. Neuron. 2002;35:419–432. doi: 10.1016/S0896-6273(02)00794-8.

Yang H, Kim A, David T, Palmer D, Jin T, Tien J, Huang F, Cheng T, Coughlin SR, Jan YN, Jan LY. TMEM16F forms a Ca2+-activated cation channel required for lipid scrambling in platelets during blood coagulation. Cell. 2012 Sep 28;151(1):111-22. PMID: 23021219

Yin J, Liu X, He Q, Zhou L, Yuan Z, Zhao S. Vps35-dependent recycling of Trem2 regulates microglial function. Traffic. 2016 Dec;17(12):1286-1296. PMID: 27717139

Zampese E, Fasolato C, Kipanyula MJ, Bortolozzi M, Pozzan T, Pizzo P. Presenilin 2 modulates endoplasmic reticulum (ER)-mitochondria interactions and Ca2+ cross-talk. Proc Natl Acad Sci U S A. 2011 Feb 15;108(7):2777-82. PMID: 21285369

Zhang B, Gaiteri C, Bodea LG, Wang Z, McElwee J, Podtelezhnikov AA, Zhang C, Xie T, Tran L, Dobrin R, Fluder E, Clurman B, Melquist S, Narayanan M, Suver C, Shah H, Mahajan M, Gillis T, Mysore J, MacDonald ME, Lamb JR, Bennett DA, Molony C, Stone DJ, Gudnason V, Myers AJ, Schadt EE, Neumann H, Zhu J, Emilsson V. Integrated systems approach identifies genetic nodes and networks in late-onset Alzheimer's disease. Cell. 2013 Apr 25;153(3):707-20. PMID: 23622250

Zhang L, Feng XK, Ng YK, Li SC. Reconstructing directed gene regulatory network by only gene expression data. BMC Genomics. 2016 Aug 18;17 Suppl 4:430. PMID: 27556418

Zhang Y, Li H, Li X, Wu J, Xue T, Wu J, Shen H, Li X, Shen M, Chen G. TMEM16F Aggravates Neuronal Loss by Mediating Microglial Phagocytosis of Neurons in a Rat Experimental Cerebral Ischemia and Reperfusion Model. Front Immunol. 2020 Jul 7;11:1144. PMID: 32733436

Zhao Y, et al. TREM2 is a receptor for beta-amyloid that mediates microglial function. Neuron. 2018;97:1023–1031. PMID: 29518356

Zhao Y, Li X, Huang T, Jiang LL, Tan Z, Zhang M, Cheng IH, Wang X, Bu G, Zhang YW, Wang Q, Xu H. Intracellular trafficking of TREM2 is regulated by presenilin 1. Exp Mol Med. 2017 Dec 1;49(12):e405. PMID: 29611543

Zhong L, et al. Amyloid-beta modulates microglial responses by binding to the triggering receptor expressed on myeloid cells 2 (TREM2) Mol. Neurodegener. 2018;13:15. PMID: 29587871

Zhou Y, Tada M, Cai Z, Andhey PS, Swain A, Miller KR, Gilfillan S, Artyomov MN, Takao M,
Kakita A, Colonna M. Human early-onset dementia caused by DAP12 deficiency reveals
a unique signature of dysregulated microglia. Nat Immunol. 2023 Mar;24(3):545-557.
PMID: 36658241
Zhou Y, Tong Z, Jiang S, Zheng W, Zhao J, Zhou X. The Roles of Endoplasmic Reticulum in
NLRP3 Inflammasome Activation. Cells. 2020 May 14;9(5):1219. PMID: 32423023

685
686
687

688　**Acknowledgments:**
689
690
691　**Funding:** This research was supported by NIH grants R01AG067606 (GMF, TWK).
692
693
694　**Author contributions:** H.-W.R and T.-W.K. conceived the study and wrote the manuscript. C.K.,
695　　　　G.M.F., Y.R.P., J.Y.M., H.-W.R and T.-W.K. designed the experiments. C.K., G.M.F.,
696　　　　Y.R.P. performed experiments and analyzed and interpreted the data. A.G. and O.H.
697　　　　analyzed and interpreted bioinformatics data.
698
699
700　**Competing interests:** TWK is a cofounder of BL Melanis Co. Ltd. The remaining authors have
701　　　　no competing interests related to this study.
702
703
704　**Data and materials availability:** All data need for evaluating the main conclusions of this work
705　　　　are included in the paper and/or Supplementary Materials.

706
707
708

Figures and Tables

Fig. 1. TurboID-based proximity labeling of TREM2 in microglial cells. (**A**) Schematic
workflow of proximity labeling, super-resolution TurboID, mass spectrometry and in situ
TREM2 interactome mapping. (**B**) Western blot analysis of the samples from the IM-HM
microglial cells transfected with the expression constructs: human wild-type or R47H
version of TREM2 with C-terminal V5 tag and TruboID and mCherry-V5. (**C**) Confocal
microscopy analysis of IM-HM cells transfected with TREM2-V5-TurboID using anti-V5
antibodies or with Streptavidin-647 for the detection of biotin-labeled protein.

723
724

Fig. 2. Clustering of TREM2 interactome. (**A**) Volcano plots showing statistically
significant enrichment of biotinylated proteins (group-TREM2-TurboID) over the cytosolic
biotinylated proteins by NES-TurboID (TREM2 interactome). Among the TREM2
interactome, MERC components are also indicated. Enrichment of the Spot-TurboID
detected biotinylated proteins by TREM2-V5-TurboID. x-axis: Log2 value of fold change
(FC) of mass signal intensities. Cut off value is 1.2; y-axis: -Log10 value of p-value, cut
off value is 1.2. (**B**) Segregation of TREM2 interactome vs. previously known MERC
components shown by Contact-ID (Kwak et al., 2020). (**C**) Subcellular distribution of
TREM2 interactome proteins across the MERC (MAM), ER, mitochondria, secretory
vesicles, Golgi and cell surface.
735

Fig. 3. Demonstration of MERC localization of TREM2 using the contact-dependent PL labeling. (**A**) Schematic diagram of Contact-ID for MERC mapping using split BioID. (**B**) Confocal microscopy imaging of MERC biotinylation by Contact-ID in HEK293T cells.

TREM2 was visualized by anti-V5 antibody. Biotinylated proteins were visualized by AF647-conjugated streptavidin. (**C**) Streptavidin-horseradish peroxidase (SA-HRP) Western blot of biotinylated proteins.

Fig. 4. Ultrastructural Analysis of MERCs in wild type iMGL and TREM2 deficient iMGL. (**A**) Transmission Electron microscopy (TEM) images of wild-type and TREM2 deficient iMGL cells. Red boxes are enlarged images of dotted area. Each ultrastructure of intracellular organelles including mitochondria and ER is intact without abnormality. scale bar: 1 um. (**B**) The thickness and length of MERC were analyzed in each group. Red arrows represent MERC. The thickness of MERCs with less than 30 nm between the mitochondria and ER was measured in 10 regions of MERCs. (**C**) The averaged thickness was significantly increased in TREM2 deficient iMGL, and the thickness (nm) was each 14.66 ± 0.5824 and 18.58 ± 0.4943 in wild-type and TREM2 deficient cells. N=50 MERCs. Data are represented as mean ± SEM and the statistical significance was p <0.001.

Fig. 5. The thickness of MERC in TREM2 deficient iMGL and the iMGL infected with TREM2 lentivirus. (**A**) Transmission Electron microscopy (TEM) images of TREM2 deficient iMGL cells with control lentivirus (left), and TREM2 lentivirus (right). An increased number of LDs in TREM2-deficient iMGL was prominent, in contrast, LD was decreased in the cell infected with TREM2 lentivirus. (**B**) Black squares are enlarged images from dotted rectangles showing MERC. Arrows showed MERC. (**C**) Increased thickness of MERC in TREM2 deficient iMGL was rescued by TREM2 encoding vector. Each averaged thickness (nm) of MERC was 19.52 ± 0.5512 and 14.1 ± 0.3741 in TREM2 deficient iMGL and the cell infected with TREM2 encoding vector. N=87 MERCs. Data are represented as mean ± SEM and the statistical significance was p <0.001.

Fig. 6. Increased thickness of MERC in TREM2 R47mutated iMGL. (A) Transmission
Electron microscopy (TEM) images of iMGL cells from wild type and patient with R47H
mutation. Enlarged images from dotted boxes (Right) showed MERCs. (B) The thickness
of MERCs was analyzed in isogenic wild type and TREM2 R47H mutated iMGL. Each
averaged thickness (nm) of MERC was 14.36 ± 0.3996 and 17.96 ± 0.3982 in isogenic
wild type and TREM2 R47H mutated iMGL. N=110. Data are represented as mean ±
SEM and the statistical significance was p <0.001.

Fig. 7. Expression of MERC proteins in human microglia derived from control and TREM-carrier of R47H, R62H, and H157Y mutation. (**A**) UMAP plot showing the microglia nuclei from the Homeostatic (overexpression of TMEM119, p= 3.34 x 10^{-5}), Activated (overexpression of CD9, p = 2.93E-93), IFN (overexpression of IFIT3, p= 3.24 x 10^{-58}), and IL1B (overexpression of IL1B, p= 8.98 x 10$_{-200}$) clusters. (**B**) Expression of MERC gene in microglia transcriptional state. Red line shows the mean expression value. (**C**) Differential gene expression of each MERC genes (scale expression >0.1) in the four microglia transcriptional states. (**D**) UMAP representation of the Control (N = 2251; left) and TREM2-carrier (N = 2136; right) cells. The color code represents the expression profile of the MERC genes (depicted in panel c) in each microglia transcriptional state. (**E**) Comparison of expression profile of MERC gene in Control and AD carrier in TREM2-carrier risk variants.