Arteries are finely tuned thermosensors regulating myogenic tone and blood flow

Thieu X. Phan[1,2], Niaz Sahibzada[1] and Gerard P. Ahern[1*]

[1]Department of Pharmacology and Physiology, Georgetown University, Washington DC, USA

[2]Department of Biology, Vinh University, Vinh City, Vietnam

[*]Address correspondence to:
Gerard Ahern, Department of Pharmacology and Physiology, Georgetown University, Med-Dent Building SW401, 3900 Reservoir Rd, Washington DC 20007, USA, Phone (202) 687-9678; Fax (202) 687-2585; Email: gpa3@georgetown.edu

Subject Areas: Ion Channels/Membrane Transport, Vascular biology

Summary

The smooth muscle cells surrounding small arteries and arterioles sense changes in local blood pressure and reflexively adjust their contractility to regulate vessel caliber. This autoregulatory property, known as myogenic tone, helps maintain a constant capillary pressure, which is important for tissue perfusion, fluid balance and vessel integrity. Here, we show that in addition to pressure, temperature is a critical determinant of myogenic tone. Heating steeply activates tone in mouse skeletal muscle, brain, gut, and skin arteries with a temperature co-efficient (Q_{10}) of ~11-20, reflecting a highly thermosensitive process. Further, the temperatures for half-maximal tone correspond closely to resting tissue temperatures, indicating that arterial tone is sensitive to small thermal fluctuations. In skeletal muscle arteries, we show that temperature and intraluminal pressure are sensed largely independently yet integrated to trigger contractility. Further, we reveal key roles for heat-sensitive channels TRPV1 and TRPM4 in heat-induced myogenic tone in the skeletal muscle, both *in vitro* and *in vivo*. Elevations in temperature are known to increase tissue perfusion by altering blood viscosity; remarkably, we demonstrate that thermosensitive tone counterbalances this effect. Thus, the thermosensitivity of myogenic tone is tuned to tissue temperature and serves as a fundamental mechanism to regulate local blood flow.

Introduction

The cardiovascular system has critical roles both in transporting oxygen and nutrients and in thermoregulation. To achieve these goals the cardiac output must be appropriately distributed among tissues. Several important extrinsic and local mechanisms regulate tissue blood flow including the autonomic nervous stimuli, humoral transmitters, shear stress and metabolic factors (*1, 2*). In addition, small arteries and arterioles possess the intrinsic capacity to constrict or dilate in response to changes in intravascular pressure. This autoregulatory mechanism is known as the vascular myogenic response or myogenic tone (*3, 4*). It is proposed that myogenic tone allows for a highly localized, fine tune control of blood perfusion and protects against fluctuations in capillary pressure that would negatively affect fluid transport or compromise vessel integrity (*5*).

In the prevailing theory, first established by William Bayliss in 1902 (*3*), blood pressure is the primary determinant of myogenic tone. Specifically, blood pressure exerts a circumferential stretch of the surrounding vascular smooth muscle cells (VSMC)(*5-7*) leading to membrane depolarization, the activation of L-type Ca^{2+} channels and a subsequent increase in myoplasmic $[Ca^{2+}]$ necessary to trigger contraction (*8-12*). Recent studies have identified G-protein coupled receptors (GPCR) as key VSMC mechano-sensors (*7, 13-15*). In turn, these GPCRs activate transduction channels, which mediate membrane depolarization. Putative transduction channels include members of the transient receptor potential (TRP) family (*16-19*) and ANO1 (*20*). Intravascular pressure via GPCRs also activates a minor, Rho-kinase mediated, Ca^{2+} independent pathway that promotes VSMC contraction (*21*).

Temperature also strongly affects blood flow. The autonomic nervous system bi-directionally regulates cutaneous perfusion in response to heating or cooling (*22, 23*). In addition, heating reduces blood viscosity (~2-3% per degree Celsius) (*24, 25*) and hence vascular resistance, and this leads to increased tissue perfusion per unit pressure (*23, 26-30*). This is an important and often overlooked consideration in the circulatory system, which carries blood through tissues of varying temperatures. Indeed, the skin and skeletal muscle in the body's thermal shell experience large temperature changes that at distal sites may exceed 10-15°C (*26, 31, 32*). Further, core body temperatures display temporal variations that are amplified by exercise, metabolism, or thermal stress, including significant deviations during torpor (*33*). If unchecked, these fluctuating tissue temperatures would lead to large changes in tissue perfusion independent of blood pressure. In this study, we show that arteries are intrinsically highly thermosensitive and generate myogenic tone by integrating both pressure and temperature stimuli. In this manner, the myogenic response optimally regulates blood flow under diverse physiological conditions.

Results

Myogenic tone is steeply heat-dependent and tuned to tissue temperature
It is well established that myogenic tone is temperature sensitive, in that tone is absent at room temperature and only develops when tissues are maintained at physiologic temperature (*34, 35*). The underlying basis for this temperature dependence is unclear. Intriguingly, several of the candidate transduction channels for myogenic tone are highly thermosensitive, including TRPM4 (*36*), TRPV1 (*37*), and ANO1 (*38*). We therefore hypothesized that myogenic tone is dynamically regulated by temperature. As TRPV1 and TRPM4 play a key role in the myogenic tone of skeletal muscle arteries (*19*), we first explored whether temperature dynamically regulates tone in radial muscle branch arteries (Supp. Fig. 1). In isolated, pressurized (80 mmHg) arteries, myogenic tone was mostly absent below ~28°C, but developed steeply to a maximum of 45% as temperature was incrementally raised to 40°C (**Figs. 1a-d**). The mean temperature for half-maximal activation ($T_{1/2}$), calculated by fitting data from individual arteries, was 33.4 ± 1.8 °C (n=11). Analysis of the temperature dependence using an Arrhenius plot, revealed a steep component with a threshold of ~27-28°C and a mean temperature coefficient (Q_{10}) of 12.6 ± 1.8 (n=6, **Fig. 1d and Supp. Fig. 2**). Further, rapid cooling from 37°C to 32°C abruptly dilated arteries (**Fig. 1c**), supporting a highly temperature-sensitive process.

Next, to test whether the dynamic temperature sensitivity of myogenic tone is specific to skeletal muscle arteries or is a general property of resistance vessels, we studied pressurized arteries isolated from the gut (mesenteric), brain (pial) and skin (hairy and glabrous, **Supp. Fig. 1**). Remarkably, we found that myogenic tone in all these arteries is temperature-dependent over a narrow range; albeit the $T_{1/2}$ was tissue-specific (**Fig. 1e-i**). In mesenteric arteries, heating activated a maximal tone of ~46% with a $T_{1/2}$ of 34.4 ± 1.9°C (n=11) and a Q_{10} of 11.7 ± 1.1 (n=5) **(Figs. 1e,f and Supp. Fig. 2)**. In pial arteries, heating activated 49% tone with a $T_{1/2}$ of 35.7 ± 1.4°C (n=6) and a Q_{10} of 11.3 ± 1.3 (n=4) **(Fig. 1G and Supp. Fig. 2)**. In "hairy" skin arteries, heating activated 44% tone with a $T_{1/2}$ of 30.8 ± 1.4°C (n=10) and a Q_{10} of 20.0 ± 2.4 (n=4) **(Figs. 1e,h and Supp. Fig. 2)**. In glabrous skin (digital) arteries, heating activated a bi-phasic response; an initial component of ~13% tone with a $T_{1/2}$ of 23.7 ± 1.4°C (n=7) and an additional component of 40% with a $T_{1/2}$ of 38.0 ± 1.9°C **(Fig. 1i)**. The respective Q_{10} values were 17.3 ± 3.6 (20-26°C, n=5), and 11.9 ± 5.0 (>35°C, n=5) **(Fig. 1i and Supp. Fig. 2)**. These arteries all reside in tissues with distinct temperatures reflecting their location in the body's thermal core or shell (**Fig. 2a**). Although warm core blood perfuses peripheral tissues, due to heat loss and counter-current heat exchange with veins, the temperature of the blood in small arteries (50-150μM) approximates the tissue temperature (*39, 40*). Thus, VSMCs of small peripheral arteries sense the tissue temperature. The published values for tissue temperatures in mice (ambient temperature of 21°C) are skeletal muscle, 34.5°C (*41*); gut, 36°C (*42*); brain, 37°C (*43*); hairy skin, 32°C (*44, 45*); and glabrous (plantar) skin, 25°C (*46*). Notably, the $T_{1/2}$ for myogenic tone from individual arteries showed a high degree of correlation (r^2=0.93, P<0.0001) with published tissue temperatures (**Fig. 2b**). Thus, the thermosensitivity of myogenic tone (hereafter abbreviated as thermo-tone) is tuned to tissue temperature.

Pressure and temperature are independent sensors for myogenic tone

Next, we explored the underlying basis for thermo-tone. Temperature affects all proteins and enzymatic reactions with a Q_{10} of ~2-3. In contrast, the Q_{10} for myogenic tone of ~11-20 suggests a highly thermosensitive mechanism, consistent with a role for specialized heat-sensitive proteins. To identify the locus for temperature sensitivity, we first tested the interrelationship between intraluminal pressure and temperature in skeletal muscle arteries. **Figures 3a&b** show the effect of varying the pressure (0-120 mmHg) at fixed temperatures. At 32°C, raising the intraluminal pressure evoked a maximal tone of 15% and the half-maximal response ($P_{1/2}$) was obtained at 68 ± 14 mmHg. Increasing the temperature to 35°C and then 39°C significantly augmented the maximal tone to 25% and 40%, respectively, with only a modest reduction in the $P_{1/2}$ to 58 ± 6 and 47 ± 12 mmHg respectively (**Fig. 3b**). Similarly, we measured the temperature sensitivity at different fixed pressures (40, 80 and 120 mmHg). **Figures 3c&d** show that raising the pressure increased the maximal tone without significantly altering the $T_{1/2}$ of ~34°C. Thus, both pressure and temperature are critical stimuli that are sensed largely independently, yet integrated to generate myogenic tone. Further, these results indicate that mechanosensors in VSMCs do not significantly contribute to thermosensing.

To understand how temperature and pressure are integrated, we tested for thermo-sensitivity in the signaling pathways downstream of pressure sensing. Previous studies have shown that stretch-mediated depolarization of vascular smooth muscle, followed by activation of L-type Ca^{2+} channels (Cav1.2), accounts for the majority of tone (*8-12*). Accordingly, we found that the Ca^{2+} channel blocker, nifedipine inhibited ~80% of tone in skeletal muscle arteries (**Figs. 3f&g**). Heating had only a modest effect on the residual nifedipine-insensitive tone (**Fig. 3g**). This is consistent with the minimal temperature sensitivity of smooth muscle contraction (Q_{10} of 1.8) (*47*). Further, the gating of L-type Ca^{2+} channels is only weakly temperature-dependent (*48*). Together, these data suggest that temperature sensing resides upstream of Ca^{2+} entry. Indeed, we found that heating regulated the stretch-induced membrane depolarization of VSMCs; at 28°C, pressurized cells remained hyperpolarized but increasing the temperature to 36°C depolarized the membrane potential by 20 mV (**Fig. 3e**). Thus, these data suggest that the transduction channels, that mediate membrane depolarization, are the locus for thermo-tone.

TRPV1 and TRPM4 mediate thermo-tone in skeletal muscle arteries

TRPV1 and TRPM4 are putative transduction channels for myogenic tone in skeletal muscle arteries, activated via a phospholipase C (PLC) dependent pathway (*19*). To test a role for TRPV1 in thermo-tone, we used BCTC, an antagonist with demonstrated on-target selectivity for TRPV1 in VSMCs (*19*). BCTC inhibited VSMC membrane depolarization in pressurized arteries at 35°C (**Fig. 3e**). Further, BCTC inhibited myogenic tone (80 mmHg) in a temperature dependent manner (**Fig. 3f-g**). BCTC had a greater inhibitory effect between 30°C to 34°C (~60% block) compared with 37°C (~40% block) supporting a larger relative contribution of TRPV1 to tone at these lower temperatures (**Fig. 3g**). Accordingly, BCTC reduced the Q_{10} over the 30-33°C

temperature range (**Fig. 3h)** and produced a right shift in the temperature-myogenic tone relationship (**Supp. Fig. 3)**; the $T_{1/2}$ increased from 33.4 ± 1.8 °C (n=11) to 35.9 ± 0.6 (n=5, P=0.01). Likewise, myogenic tone (80 mmHg) in TRPV1-deficient arteries was significantly depressed at 32°C and 34°C (**Fig. 3i)** but not at 37°C, suggesting compensation for loss of TRPV1 at higher temperatures. To test a role for TRPM4 in thermo-tone, we used the antagonist, 9-phenantrol, and a new, highly selective inhibitor, 4-Chloro-2-(1-naphthyloxyacetamido)benzoic acid (NBA). Both compounds significantly blocked myogenic tone by ~40% at 37°C (**Fig. 3f)**. The inhibitory effect of NBA was reduced at lower temperatures (**Fig. 3g)**; unlike BCTC, NBA did not significantly inhibit tone at 30°C and 32°C suggesting that TRPM4 is activated at a slightly higher temperature range. Further, combining NBA with BCTC, suppressed tone to the same extent as blocking L-type Ca^{2+} channels, suggesting that TRPV1 and TRPM4 are the major contributors to thermo-tone in the 30-37°C temperature range (**Fig. 3f)**. Thus, in skeletal muscle arteries, TRPV1 and TRPM4 appear to confer a largely overlapping thermal sensitivity of myogenic tone with TRPV1 making a greater contribution to thermosensing at 30-34°C.

The activation of TRPV1 at 30°C in VSMCs may seem surprising, since thermal activation of TRPV1 typically occurs at greater than 40°C (*37*). However, allosteric modulators are known to substantially lower the heat threshold. Indeed, PLC-mediated signaling, leading to phosphoinositide depletion, generation of arachidonic acid metabolites or PKC activation (*49*), triggers activation of TRPV1 at <32°C and this may be accompanied by a substantial reduction in the Q_{10} (*50*). Accordingly, we tested whether PLC regulated thermo-tone. **Figures 3j&k** show that the PLC inhibitor, U73122, reduced the maximal tone in pressurized arteries from ~45% to 12% and shifted the temperature threshold from 28°C to 38°C and the $T_{1/2}$ from 33.4°C to 39.1°C. Furthermore, the TRPV1 inhibitor, BCTC significantly inhibited the residual tone in U73122-treated arteries (P<0.001) (**Fig. 3k**). Thus, PLC signaling (with a major contribution of TRPV1) tunes the thermo-tone close to the resting temperature of skeletal muscle.

Next, to assess the role of thermo-tone *in vivo,* we exploited the well-described pharmacologic effects of TRPV1 antagonists. When administered acutely to mice, TRPV1 inhibitors reduce the systemic blood pressure (*19*). This response is attributed to blockade of TRPV1-dependent myogenic tone and is retained after the deletion of TRPV1-positive sensory neurons (*19*). Therefore, we measured the systemic blood pressure response to the TRPV1 antagonist, BCTC, at different core and skeletal muscle temperatures ($T_{Core}/_{Sk.M}$). As previously reported, BCTC (3 mg/kg) administered to mice at a near normal $T_{Core}/_{Sk.M}$ of 36.4/33.0°C produced an immediate decrease in blood pressure of ~10 mm Hg (**Figs. 4b-e**). In contrast, when the $T_{Core}/_{Sk.M}$ was lowered to 32/29.0°C, BCTC failed to change the blood pressure (**Figs. 4b-e**). Hypothermia alone did not significantly affect the resting blood pressure (**Fig. 4d**) and therefore, the results are consistent with a loss of TRPV1-dependent thermo-tone. Further, intravital imaging showed that raising the local temperature surrounding the radial muscle branch arteries from 30°C to 37°C significantly decreased the vessel diameter (**Fig. 4f**). Thus, temperature affects myogenic tone *in vivo*.

Thermo-tone counteracts heat-evoked changes in blood flow

Our results indicate that increasing skeletal muscle temperature over the physiological range increases myogenic tone **(Fig. 4).** This may seem surprising and counterintuitive, since many earlier studies have demonstrated that local heating over this same temperature range increases blood flow. Indeed, heating in various tissues, including skeletal muscle, bone, and isolated liver results in a 50-100% increase in perfusion (*23, 28, 29*). The increased blood flow is attributed to an increase in local vascular conductance explained by reductions in blood viscosity and/or vasodilation (*29*). Therefore, to understand how thermo-tone affects blood perfusion, we performed laser speckle imaging of the skeletal muscle radial branch arteries (**Fig. 5a&b**). To prevent any confounding effects of changes in sympathetic tone in these experiments, we included the α1-adrenergic antagonist, prazosin, in the super-perfusate. **Figure 5b&c** shows that step increases in the local temperature from 28°C to 34°C and 39°C modestly increased perfusion (by 31% and 80%, respectively), in line with previous studies (*23, 28*). To test whether changes in myogenic tone regulate this heat-induced response we blocked tone using a Ca^{2+}-free medium (supplemented with sodium nitroprusside) or nifedipine. After blocking myogenic tone, heating markedly increased perfusion (by 221% and 247%, respectively, **Fig. 5b&c**). Over the 28-34°C range, this represents a ~25% increase in perfusion for each degree. Since vascular tone is essentially eliminated by the Ca^{2+} free/sodium nitroprusside treatment, the heat-induced changes in perfusion must reflect altered blood fluidity rather than vasodilation. Thus, inhibiting myogenic tone unmasks the full effects of temperature on fluidity. Similarly, treatment with BCTC plus NBA increased perfusion at 34°C and 39°C but not 28°C consistent with the relaxation of TRPV1/TRPM4-dependent thermo-tone (**Fig. 5b&c**). Taken together, these data show that thermo-tone counteracts the temperature-induced facilitation of blood perfusion.

Discussion

Our data reveal that vascular myogenic tone is dynamically heat dependent over a narrow temperature range (**Fig. 1**). This temperature dependence for myogenic tone appears to be a ubiquitous feature for all resistance arteries. Although previous studies performed at a fixed temperature, report varying levels of myogenic tone, our analysis shows that the peak myogenic tone at 39-41°C exceeds ~40-45% in most tissues. Notably, we found that the $T_{1/2}$ for tone, residing at the steepest portion of the temperature relationship, closely matches the resting tissue temperature (**Fig. 2**). Thus, the resultant vessel caliber and blood flow are maximally sensitive to temperature changes. While core body temperature is generally maintained as the ambient temperature changes, the skin experiences constant temperature fluctuations. Fittingly, we demonstrate that myogenic tone in skin arteries exhibits the greatest temperature sensitivity with a Q_{10} of ~20. The biphasic response to temperature in glabrous skin (**Fig. 1i)** matches the two distinct states of this tissue: the first operating under high, resting sympathetic tone with minimal blood flow and a skin temperature just above the ambient level, and the second after removal of sympathetic tone accompanied by a large inflow (>500%) of heated blood from the core (*32*).

At first glance, thermo-tone may appear to contradict key features of thermal physiology. For example, local or whole-body heating typically leads to an increase in vascular conductance and a fall in systemic blood pressure (*22, 23*), which seems at odds with thermo-tone. However, myogenic tone is just one component of hemodynamic regulation acting in combination with metabolic, humoral and neurogenic factors as well as blood rheology. Thus, heating reduces blood viscosity (*24, 25*) and triggers a sympathetic nerve-mediated increase in cutaneous blood flow (*22, 23*), and these effects can more than compensate for the effects of thermo-tone on vascular conductance and blood pressure. Similarly, during exercise the temperature of skeletal muscle can increase significantly, and the resultant increase in thermo-tone would appear to imperil the required increase in blood supply. However, metabolic factors produced by contracting muscle are well known to inhibit myogenic tone such that there is an ascending relaxation of the supply vessels (*51*). Indeed, in this context, increased thermo-tone may be beneficial by re-directing blood flow to actively contracting muscle units.

Our findings suggest that VSMCs integrate the output of distinct pressure and temperature sensors to generate myogenic tone (**Fig. 3a-d**). Although the precise molecular mechanisms underlying the myogenic response are still unresolved, there is strong evidence supporting a role for GPCRs as mechano-sensors (*7, 13-15*). Indeed, the Angiotensin II Receptor AT1 plays a key role in the myogenic tone of mesenteric, brain, and skeletal muscle arteries (*52-55*). In turn, PLC signaling downstream of GPCRs, activates transduction channels that depolarize membrane potential (*8, 10, 56*) to activate L-type Ca^{2+} channels and ultimately trigger VSMC contraction. Our results show that stretch-induced depolarization of VSMCs is absolutely heat dependent (**Fig. 3e)**, supporting the hypothesis that transduction channels are the primary locus of thermo-sensitivity. Using a combination of pharmacologic and genetic approaches, we show that TRPV1 mediates a major portion of thermo-tone in skeletal muscle both *in vitro* and *in vivo* (**Fig. 3&4**). TRPV1 makes a greater relative contribution to tone in the 30-34°C temperature range and blockade of TRPV1 increases the $T_{1/2}$ from ~33°C to 36°C. Notably, stretch-induced PLC signaling lowers the temperature threshold for tone from ~38°C to <30°C with a major contribution of TRPV1 (**Fig. 3j,k**). Thus, the integration of pressure (via PLC) and temperature occurs at the transduction channels (**see Fig. 5e**). Additionally, we found that TRPM4, also contributes to thermo-tone in skeletal muscle arteries, particularly at temperatures greater than 34°C (**Fig. 3g**). The molecular basis for thermo-tone in other tissues remains to be determined, but similar to skeletal muscle it is likely that a combination of thermosensitive channels is responsible in each tissue. Indeed, previous studies have implicated TRPM4 as well as the heat-gated channels TRPV2 (*57, 58*) and ANO1 (*20*) in the myogenic tone of retinal, mesenteric and cerebral arteries, and all of these channels could confer thermo-sensitivity. On the other hand, TRP channels with limited thermo-sensitivity, such as TRPC6 (*16*) are also implicated in myogenic tone. It is possible that heat-gated channels provide the initial signaling event, which in turn, recruits thermo-insensitive channels to amplify the membrane depolarization.

Why is myogenic tone temperature sensitive? We propose that thermo-tone compensates for temperature-dependent changes in the rheological properties of blood. Heating enhances blood fluidity (*24, 25*), and unchecked, would lead to over perfusion. This change in fluidity mostly impacts the arterioles, capillaries, and venules where frictional forces of blood are most apparent (low Reynolds number). Indeed, we found that inhibiting myogenic tone resulted in very large increases in heat-stimulated blood flow. Thus, these observations lead to a general principle regarding blood flow regulation: small deviations (positive or negative) from the resting tissue temperature that alter blood fluidity are countered by changes in myogenic tone. Therefore, in our model (**Fig**. **5f**), myogenic tone can be represented by two independent relationships: 1) a traditional pressure-flow relationship in which temperature is constant (36°C) and blood flow is stabilized over a discrete pressure range (40 to 150 mmHg), and 2) a temperature-flow relationship in which pressure is fixed (80 mmHg) and blood flow is stabilized over a discrete temperature range (30-40°C). While this temperature range is specific to skeletal muscle, it can be readily adjusted to describe the thermo-sensitivity in other tissues. Our data therefore suggest a fundamental role for thermo-tone in the homeostatic regulation of blood flow. These findings are also relevant to the regulation of perfusion in pathological states such as hypothermia/hyperthermia.

Methods

Ethical approval: All procedures were approved by Georgetown University, IACUC Protocol Number: 2018-0033.

Animals: Male C57Bl6 wild-type and TRPV1-KO mice (25-30 g) were housed at 21-22°C and had *ad libitum* access to a standard laboratory chow and water.

Ex vivo artery physiology: Mice were euthanized (CO_2/decapitation) and the following arteries were isolated: skeletal muscle (radial artery branch, ~60μm, or subscapular muscle branch, ~120mm subscapular branch), mesenteric (3rd order, ~120μm), pial (~80-120μm) and skin (hairy skin, and glabrous skin ~100-150 μm, see **Supp. Fig. 1**). Arteries were cannulated with glass micropipettes, and secured with monofilament threads (*59*). The pipette and bathing physiological salt solution (PSS) containing (in mM): 125 NaCl, 3 KCl, 26 $NaHCO_3$, 1.25 NaH_2PO_4, 1 $MgCl_2$, 4 D-glucose, and 2 $CaCl_{2,}$) was aerated with a gas mixture consisting of 21% O_2, 5% CO_2 to maintain pH (pH 7.4). To induce maximal dilation, arteries were perfused with a PSS solution containing 0 $CaCl_2$, 0.4 mM EGTA and 100 μM sodium nitroprusside (SNP). Arteries were mounted in a single vessel chamber (Living Systems Instrumentation) and placed on a heated imaging stage (Tokai Hit) to maintain bath temperature between 15-42°C, while intraluminal pressure was maintained by a Pressure Control Station (Stratagene). Arteries were viewed with a 10X objective using a Nikon TE2000 microscope and recorded by a digital camera (Retiga 3000, QImaging). Increases in intraluminal pressure or temperature were maintained for a 3-5 min duration to enable a stable vessel response. The arteriole diameter was measured at several locations along each

arteriole using the NIH-ImageJ software's edge-detection plug-in (Diameter) (*60*). The software automatically detects the distance between edges (by resampling a total of five times from adjacent pixels) yielding a continuous read-out ±SD of a vessel's diameter. Myogenic tone (%) was calculated by: $100 \times (\text{Diameter}_{ZeroCa2+} - \text{Diameter}_{PSS}) / \text{Diameter}_{ZeroCa2+}$. The temperature for half-maximal tone was obtained by fitting a Hill function to data from individual arteries.

Membrane potential measurements: Sharp glass microelectrodes (50 MΩ) filled with 3 M KCl were inserted through the adventitial layer of pressurized (20 or 100 mmHg) arteries. The membrane potential was recorded using a Warner OC-725C amplifier. Data analysis was restricted to impalements that produced a sharp voltage deflection and a subsequent stable potential.

Intravital imaging: Intravital imaging was performed in anesthetized mice (urethane 1.2-1.5 g/kg/IP) as previously described (*19, 59*). Briefly, the brachial-radial artery junction was surgically exposed and visualized with a Zeiss stereomicroscope. The exposed arteries were locally perfused (using a 250 µm cannula connected to a valve-controlled gravity-fed perfusion system) with preheated PSS buffer. The surface tissue temperature was measured via a thermistor (Warner Instruments) that was positioned next to the artery. Measurements were made several minutes after temperature had stabilized.

Laser Speckle Contrast Imaging: LSCI was performed as per intravital imaging except that the forelimb was exteriorized into a PSS-filled clear heating probe (VHP3, Moor Instruments; Devon, UK) with a temperature controller (moorVMS HEAT and LDF2). The exposed tissue was imaged with the RFLSI Ⅲ system (RWD Life Science, Shenzhen, China). Resulting images were digitized and analyzed off-line.

Systemic blood pressure recording: The experiments were performed in anesthetized mice (urethane 1.2-1.5 g/kg/IP) as previously described (*19, 59*). Briefly, the left carotid artery and the right jugular vein were cannulated with a Millar catheter (1F SPR-1000) and a polyethylene tubing (PE-10), respectively, for monitoring arterial blood pressure and for systemic (intravenous) infusion of drugs. Body temperature was maintained using a heating plate and core (T_{Core}) and skeletal muscle (T_{SkM}) temperatures were monitored by a digital rectal thermometer and thermoprobe implanted in deep hindlimb muscle. Stock solutions of BCTC (30 mg/ml in DMSO) were dissolved in physiological saline containing 2-hydroxypropyl-Beta-cyclodextrin (25% w/v). Final DMSO concentration was <33%. Intravenous infusion of drugs was initiated only when a stable baseline of blood pressure and heart rate was present.

Chemicals: 4-(3-Chloro-2-pyridinyl)-*N*-[4-(1,1-dimethylethyl)phenyl]−1-piperazinecarboxamide (BCTC), U73122, 9-phenantrol and nifedipine were purchased from Tocris Bioscience. 4-Chloro-2-(1-naphthyloxyacetamido)benzoic acid ammonium salt (NBA) was purchased from Aobious. Stock solutions were prepared in EtOH or DMSO. Unless otherwise indicated, all other chemicals were obtained from Sigma–Aldrich.

Statistical analysis: Data were analyzed using Prism (GraphPad Software, La Jolla, CA) and are expressed as means ± SD. Unless otherwise stated, statistical significance was evaluated using *t*-test or one-way ANOVA with treatment interactions assessed by Tukey's *post hoc* multiple comparisons test. A *P* value of < 0.05 was considered statistically significant.

Acknowledgments

This study was supported by the National Heart, Lung, and Blood Institute of the National Institutes of Health under Award Number R01 HL155979 (G.A). We thank Attila Toth for helpful suggestions and Rosa Miyares for comments on the manuscript.

References

1. M. Jacob, D. Chappell, B. F. Becker, Regulation of blood flow and volume exchange across the microcirculation. *Critical care (London, England)* **20**, 319 (2016).
2. T. W. Secomb, Theoretical models for regulation of blood flow. *Microcirculation (New York, NY : 1994)* **15**, 765-775 (2008).
3. W. M. Bayliss, On the local reactions of the arterial wall to changes of internal pressure. *The Journal of physiology* **28**, 220-231 (1902).
4. P. C. Johnson, M. Intaglietta, Contributions of pressure and flow sensitivity to autoregulation in mesenteric arterioles. *The American journal of physiology* **231**, 1686-1698 (1976).
5. M. J. Davis, Perspective: Physiological Role(s) of the Vascular Myogenic Response. *Microcirculation* **19**, 99-114 (2012).
6. M. A. Hill, G. A. Meininger, Arteriolar vascular smooth muscle cells: mechanotransducers in a complex environment. *The international journal of biochemistry & cell biology* **44**, 1505-1510 (2012).
7. K.-S. Hong, K. Kim, M. A. Hill, Regulation of blood flow in small arteries: mechanosensory events underlying myogenic vasoconstriction. *Journal of exercise rehabilitation* **16**, 207-215 (2020).
8. D. R. Harder, Pressure-dependent membrane depolarization in cat middle cerebral artery. *Circulation Research* **55**, 197-202 (1984).
9. H. J. Knot, M. T. Nelson, Regulation of arterial diameter and wall [Ca2+] in cerebral arteries of rat by membrane potential and intravascular pressure. *The Journal of Physiology* **508 (Pt 1)**, 199-209 (1998).
10. N. Kotecha, M. A. Hill, Myogenic contraction in rat skeletal muscle arterioles: smooth muscle membrane potential and Ca(2+) signaling. *American journal of physiology Heart and circulatory physiology* **289**, H1326-1334 (2005).
11. J. R. H. Mauban, J. Zacharia, J. Zhang, W. G. Wier, Vascular tone and Ca(2+) signaling in murine cremaster muscle arterioles in vivo. *Microcirculation* **20**, 269-277 (2013).

12. S. Moosmang *et al.*, Dominant role of smooth muscle L-type calcium channel Cav1.2 for blood pressure regulation. *The EMBO journal* **22**, 6027-6034 (2003).
13. Y. Zou *et al.*, Mechanical stress activates angiotensin II type 1 receptor without the involvement of angiotensin II. *Nature cell biology* **6**, 499-506 (2004).
14. M. Mederos y Schnitzler *et al.*, Gq-coupled receptors as mechanosensors mediating myogenic vasoconstriction. *EMBO J* **27**, 3092-3103 (2008).
15. A. L. Gonzales *et al.*, A PLCγ1-dependent, force-sensitive signaling network in the myogenic constriction of cerebral arteries. *Sci Signal* **7**, ra49 (2014).
16. D. G. Welsh, A. D. Morielli, M. T. Nelson, J. E. Brayden, Transient receptor potential channels regulate myogenic tone of resistance arteries. *Circulation Research* **90**, 248-250 (2002).
17. S. Earley, B. J. Waldron, J. E. Brayden, Critical role for transient receptor potential channel TRPM4 in myogenic constriction of cerebral arteries. *Circulation research* **95**, 922-929 (2004).
18. S. A. Reading, J. E. Brayden, Central role of TRPM4 channels in cerebral blood flow regulation. *Stroke; a journal of cerebral circulation* **38**, 2322-2328 (2007).
19. T. X. Phan *et al.*, TRPV1 in arteries enables a rapid myogenic tone. *The Journal of Physiology* **600**, 1651-1666 (2022).
20. S. Bulley *et al.*, TMEM16A/ANO1 channels contribute to the myogenic response in cerebral arteries. *Circulation Research* **111**, 1027-1036 (2012).
21. A. Gohla, G. Schultz, S. Offermanns, Role for G(12)/G(13) in agonist-induced vascular smooth muscle cell contraction. *Circulation research* **87**, 221-227 (2000).
22. L. B. Rowell, Reflex control of the cutaneous vasculature. *The Journal of investigative dermatology* **69**, 154-166 (1977).
23. I. Heinonen *et al.*, Local heating, but not indirect whole body heating, increases human skeletal muscle blood flow. *Journal of applied physiology (Bethesda, Md : 1985)* **111**, 818-824 (2011).
24. P. W. Rand, E. Lacombe, H. E. Hunt, W. H. Austin, Viscosity of normal human blood under normothermic and hypothermic conditions. *Journal of applied physiology* **19**, 117-122 (1964).
25. G. K. Snyder, Influence of temperature and hematocrit on blood viscosity. *The American journal of physiology* **220**, 1667-1672 (1971).
26. H. Barcroft, O. G. Edholm, The effect of temperature on blood flow and deep temperature in the human forearm. *The Journal of Physiology* **102**, 5-20 (1943).
27. D. Akyürekli, L. H. Gerig, G. P. Raaphorst, Changes in muscle blood flow distribution during hyperthermia. *International journal of hyperthermia : the official journal of European Society for Hyperthermic Oncology, North American Hyperthermia Group* **13**, 481-496 (1997).
28. T. Binzoni, D. Tchernin, J. Richiardi, D. Van De Ville, J.-N. Hyacinthe, Haemodynamic responses to temperature changes of human skeletal muscle studied by laser-Doppler flowmetry. *Physiological measurement* **33**, 1181-1197 (2012).
29. N. Koch Esteves, O. R. Gibson, A. W. Khir, J. González-Alonso, Regional thermal hyperemia in the human leg: Evidence of the importance of thermosensitive

mechanisms in the control of the peripheral circulation. *Physiological reports* **9**, e14953 (2021).

30. A. M. Thorne *et al.*, Hyperthermia-induced changes in liver physiology and metabolism: a rationale for hyperthermic machine perfusion. *American journal of physiology Gastrointestinal and liver physiology* **319**, G43-G50 (2020).

31. M. B. Ducharme, W. P. VanHelder, M. W. Radomski, Tissue temperature profile in the human forearm during thermal stress at thermal stability. *Journal of applied physiology (Bethesda, Md : 1985)* **71**, 1973-1978 (1991).

32. A. A. Romanovsky, Skin temperature: its role in thermoregulation. *Acta physiologica (Oxford, England)* **210**, 498-507 (2014).

33. C. J. Gordon, The mouse thermoregulatory system: Its impact on translating biomedical data to humans. *Physiology & behavior* **179**, 55-66 (2017).

34. B. R. Duling, R. W. Gore, R. G. Dacey, D. N. Damon, Methods for isolation, cannulation, and in vitro study of single microvessels. *The American journal of physiology* **241**, H108-116 (1981).

35. M. J. Davis, P. J. Sikes, Myogenic responses of isolated arterioles: test for a rate-sensitive mechanism. *The American journal of physiology* **259**, H1890-1900 (1990).

36. K. Talavera *et al.*, Heat activation of TRPM5 underlies thermal sensitivity of sweet taste. *Nature* **438**, 1022-1025 (2005).

37. M. J. Caterina *et al.*, The capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature* **389**, 816-824 (1997).

38. H. Cho *et al.*, The calcium-activated chloride channel anoctamin 1 acts as a heat sensor in nociceptive neurons. *Nature neuroscience* **15**, 1015-1021 (2012).

39. H. C. Bazett, L. Love, Temperature changes in blood flowing in arteries and veins in man. *Journal of applied physiology* **1**, 3-19 (1948).

40. Q. He, L. Zhu, S. Weinbaum, Effect of blood flow on thermal equilibration and venous rewarming. *Annals of biomedical engineering* **31**, 659-666 (2003).

41. E. Gorrell *et al.*, Skeletal muscle thermogenesis induction by exposure to predator odor. *The Journal of experimental biology* **223**, (2020).

42. V. Škop, N. Liu, J. Guo, O. Gavrilova, M. L. Reitman, The contribution of the mouse tail to thermoregulation is modest. *American journal of physiology Endocrinology and metabolism* **319**, E438-E446 (2020).

43. F. Baracchi, M. R. Opp, Sleep-wake behavior and responses to sleep deprivation of mice lacking both interleukin-1 beta receptor 1 and tumor necrosis factor-alpha receptor 1. *Brain, behavior, and immunity* **22**, 982-993 (2008).

44. J. Mei *et al.*, Body temperature measurement in mice during acute illness: implantable temperature transponder versus surface infrared thermometry. *Scientific Reports* **8**, 3526 (2018).

45. V. van der Vinne *et al.*, Continuous and non-invasive thermography of mouse skin accurately describes core body temperature patterns, but not absolute core temperature. *Scientific Reports* **10**, 20680 (2020).

46. W. Richter, Estimation of vasodilator drug effects in mice by measurements of paw skin temperature. *Acta pharmacologica et toxicologica* **21**, 91-104 (1964).

47. P. Klemt, U. Peiper, The dynamics of cross-bridge movement in vascular smooth muscle estimated from a single isometric contraction of the portal vein: the influence of temperature and calcium. *Pflugers Archiv : European journal of physiology* **378**, 31-36 (1978).

48. R. K. Wang, Edward; Pang, Peter K.P., Temperature dependence of L-type calcium channel currents in isolated smooth muscle cells from the rat tail artery. *Journal of Thermal Biology* **16**, 83-87 (1991).

49. T. Sugiura, M. Tominaga, H. Katsuya, K. Mizumura, Bradykinin lowers the threshold temperature for heat activation of vanilloid receptor 1. *J Neurophysiol* **88**, 544-548 (2002).

50. J. Yao, B. Liu, F. Qin, Kinetic and energetic analysis of thermally activated TRPV1 channels. *Biophysical journal* **99**, 1743-1753 (2010).

51. S. S. Segal, Regulation of blood flow in the microcirculation. *Microcirculation (New York, NY : 1994)* **12**, 33-45 (2005).

52. J. Schleifenbaum *et al.*, Stretch-activation of angiotensin II type 1a receptors contributes to the myogenic response of mouse mesenteric and renal arteries. *Circulation Research* **115**, 263-272 (2014).

53. S. Blodow *et al.*, Novel role of mechanosensitive AT1B receptors in myogenic vasoconstriction. *Pflugers Archiv : European journal of physiology* **466**, 1343-1353 (2014).

54. K. Hong *et al.*, Mechanical activation of angiotensin II type 1 receptors causes actin remodelling and myogenic responsiveness in skeletal muscle arterioles. *The Journal of Physiology* **594**, 7027-7047 (2016).

55. P. W. Pires *et al.*, The angiotensin II receptor type 1b is the primary sensor of intraluminal pressure in cerebral artery smooth muscle cells. *The Journal of Physiology* **595**, 4735-4753 (2017).

56. G. Osol, I. Laher, M. Kelley, Myogenic tone is coupled to phospholipase C and G protein activation in small cerebral arteries. *The American journal of physiology* **265**, H415-420 (1993).

57. K. Muraki *et al.*, TRPV2 is a component of osmotically sensitive cation channels in murine aortic myocytes. *Circulation Research* **93**, 829-838 (2003).

58. M. K. McGahon *et al.*, TRPV2 Channels Contribute to Stretch-Activated Cation Currents and Myogenic Constriction in Retinal Arterioles. *Investigative ophthalmology & visual science* **57**, 5637-5647 (2016).

59. T. X. Phan *et al.*, TRPV1 expressed throughout the arterial circulation regulates vasoconstriction and blood pressure. *The Journal of Physiology* **598**, 5639-5659 (2020).

60. M. J. M. Fischer, S. Uchida, K. Messlinger, Measurement of meningeal blood vessel diameter in vivo with a plug-in for ImageJ. *Microvascular research* **80**, 258-266 (2010).

Figure 1. Myogenic tone is thermosensitive. **a** A representative, pressurized (80 mmHg) skeletal muscle artery at 30, 34 and 37°C. **b&c**, Changes in the diameter of pressurized (80 mmHg) skeletal muscle arteries in response to an increase or decrease in bath temperature. **d-i**. Mean myogenic tone (%) versus temperature and representative Arrhenius plots yielding indicated Q_{10} values and threshold temperatures, for skeletal muscle (n=11), mesenteric (n=11), pial (n=7), hairy skin (n=10) and glabrous skin (n=7) arteries. The mean half maximal temperature ($T_{1/2}$) for tone was derived from fits to individual arteries.

Figure. 2. **The thermosensitivity of myogenic tone correlates with tissue temperature**. **a,** Illustration of thermal tissue variations in mice. Approximate resting tissue temperatures were obtained from published values for brain, gut, hairy skin, glabrous skin (see text) at an ambient T of 21-22°C. **b,** Plot of myogenic tone $T_{1/2}$ for individual arteries from skeletal muscle (n=11), mesenteric (n=11), pial (n=6), hairy skin (n=10) and glabrous skin (n=7) versus the tissue temperature.

Figure. 3. **Distinct pressure and temperature sensors mediate myogenic tone in skeletal muscle. a,** Myogenic tone versus pressure in a single skeletal muscle arteriole at 32°C, 35°C and 39°C (*left*) and the mean responses of arteries from six animals (*right*). **b,** Plot of $P_{1/2}$ versus temperature (n=5, unpaired t test, *P<0.05). **c&d,** Myogenic tone versus temperature, measured at intraluminal pressures of 40, 80 and 120 mmHg and plot of $T_{1/2}$ versus pressure (n=4). **e-k, TRPV1 and TRPM4 mediate thermo-tone. e,** Membrane potential in smooth muscle cells of skeletal muscle arteries pressurized to 10 mmHg (n=25) or 100 mmHg (36°C, n=25; 28°C, n=19; 35°C, n=21; 35°C plus BCTC, n=14; one-way ANOVA, **P<0.01, ***P<0.001). **f,** Summary of myogenic tone at 37°C and 80 mmHg under control conditions (PSS, n=9) or during pharmacological inhibition with the TRPV1 inhibitor, BCTC (3μM, n=4), TRPM4 inhibitors, 9-phenanthrol (9-phe, 30μM, n=5) and NBA (3μM, n=6), 9-phe+BCTC (n=5), NBA+BCTC (n=6) and L-type Ca^{2+} channel inhibitor nifedipine (NIF, 5 μM, n=4) (one-way ANOVA, *P<0.05, ***P<0.001). **g,** Pharmacologic inhibition of thermo-tone at different temperatures (n=4, one-way ANOVA, *P<0.05,**P<0.01, ***P<0.001). **h,** Plots of myogenic tone versus temperature for individual arteries under control conditions or in the presence of BCTC. The mean Q_{10} derived from Arrhenius plots between 30-33°C are shown. **i,** Temperature-dependent myogenic tone (80 mmHg) in WT (n=12) versus TRPV1-null (n=3) arteries (unpaired t test, *P<0.05, **P<0.01). **j,** Temperature-dependent myogenic tone (80 mmHg) in control arteries (PSS) (data re-plotted from Fig.1D, $T_{1/2}$ =33.4°C), and after treatment with the PLC inhibitor, U73122 (10μM, n=10). The $T_{1/2}$ =39.1°C. **k,** The maximal myogenic tone at 41°C for control (PSS, n=10), U73122 (n=10), U73122 + BCTC (n=9) and U73122 +NBA (n=9) (one-way ANOVA, ** P<0.01,***P<0.001).

Figure 4. **Temperature regulates myogenic tone *in vivo*. a,** Anesthetized mice were maintained at a core/skeletal muscle temperature (T_C/$T_{Sk.M}$) of 32°C/29°C or 36°C/33°C. **b,** Representative blood pressure (BP) and heart rate (HR) recording during i.v. administration of the TRPV1 inhibitor, BCTC (3mg/kg, dotted line) **c,** The change in the mean BP evoked by BCTC at different T_C (32°C, n= 4; 36°C, n=6, unpaired t test, *P<0.05) **d&e,** Summary of mean arterial pressure and heart rate (32°C n= 4; 36°C, n=6; one-way ANOVA,, *P<0.05 , **P<0.01). **f&g,** Intravital imaging of mouse radial muscle feed arteriole (green arrow) and veins (yellow arrow) shows constriction in response to local perfusion of heated ringers from at 30, 34 and 37°C (one-way ANOVA, n = 5, **$P < 0.01$, ***$P < 0.001$).

Figure 5. **Thermo-tone opposes heat-evoked increases in blood flow**. **a,** *Top*, illustration of the intravital imaging field (dotted box) in the mouse forearm. *Bottom*, expanded view showing the radial muscle branch artery (red arrows) and a vein (yellow arrows). **b,** Representative laser speckle contract images and **c,** Relative blood perfusion during limb heating from 28 to 39°C with physiological salt solution (PSS), BCTC+NBA, nifedipine and Ca²⁺ free/ sodium nitroprusside PSS. **d,** Mean changes in blood perfusion (20°C, n=5; 28-39°C, n=9, one-way ANOVA, NS=not significant, *P<0.05, **P<0.01, ***P<0.001). Note: all media contained the α1 adrenergic inhibitor, prazosin (10 µM). **e-g,** a model for temperature-dependent myogenic tone. **e,** a proposed signaling pathway for thermo-tone showing that stretch-induced PLC signaling and temperature converge to activate thermo-sensitive ion channels (see text for details). **f,** A conventional blood flow versus pressure relationship in a skeletal muscle artery (with a fixed temperature of 36°C) showing myogenic tone (blue line) operating between 40 and 150 mmHg. The black line indicates zero myogenic tone. **g,** Blood flow versus temperature in a skeletal muscle artery (with a fixed pressure of 80 mmHg) showing myogenic tone acting between 30-40°C.