Transient demographic dynamics of recovering fish stocks shaped by past climate, harvest, and

management

Daisuke Goto[1,2,¶]

[1]Institute of Marine Research, P.O. Box 1870, Nordnes, 5817 Bergen, Norway

[2]Swedish University of Agricultural Sciences, Department of Aquatic Resources, Drottningholm,

Sweden.

[¶]Corresponding author: daisuke.goto2@gmail.com

Abstract

Large-scale commercial harvesting and climate-induced fluctuations in ocean properties shape

the dynamics of marine populations as interdependent drivers at varied timescales. Selective

removals of larger, older members of a population can distort its demographic structure, eroding

resilience to fluctuations in habitat conditions and thus amplifying volatility in transient

population dynamics. Through the implementation of stricter management measures since the

1990s, many historically depleted fish stocks began showing signs of recovery. But these

interventions coincided with accelerated changes in the oceans triggered by increasingly warmer

and more variable climates. Applying multilevel models to annual estimates of demographic

metrics of 38 stocks comprising 11 species across seven ecoregions in the Atlantic Ocean, this

study explores how time-varying local and regional climate conditions contributed to the

transient dynamics of recovering populations exposed to variable fishing pressures moderated by

management actions. Analyses reveal that progressive reductions in fishing pressure and shifting

climate conditions shaped rebuilding patterns of the stocks through restorations of maternal

demographic structure (reversing age truncation) and reproductive capacity. As the survival rates

and demographic structures of reproductive fish improved, transient growths became less

sensitive to variability in recruitment and juvenile survival and more to that in adult survival. As

the biomass of reproductive fish rose, recruitment success rates also became increasingly

regulated by density-dependent intrinsic processes involving higher numbers of older fish. When

reductions in fishing pressure were insufficient or delayed, however, stocks became further

depleted, with more eroded demographic structures. Although warmer climates in spawning

seasons promoted recruitment success in some ecoregions, changing climate conditions over the

past decades began adversely affecting reproductive performance overall. These shared patterns

underscore the value of demographic transients in developing robust strategies for managing

marine resources sustainably. Such strategies could form the foundation for effective

applications of adaptive measures resilient to future environmental changes.

Keywords: age truncation, stock assessment, meta-analysis, demographic buffering, climate

change, fisheries management, matrix population model, life history trait, life table response

experiment, hierarchical modeling

Introduction

Excessive human exploitation continued through the twentieth century has depleted marine

living resources worldwide, threatening biodiversity, food and nutrient security, and other

ecosystem services that the oceans provide (Pauly, Watson & Alder 2005; Crowder *et al.* 2008;

Worm *et al.* 2009). Long-lived, late-maturing vertebrate predators are disproportionately

depleted by industrial fishing as target or incidental catch (Christensen *et al.* 2003; Juan-Jordá *et*

al. 2022). As part of the efforts to halt and reverse the losses, national government and

intergovernmental organizations began applying the precautionary principle to the management

of marine fisheries (Hilborn *et al.* 2001; Punt 2006), following international agreements,

including the Code of Conduct for Responsible Fisheries adopted by the Food and Agriculture

Organization of the United Nations (FAO 1995; Melnychuk *et al.* 2021). Since the

implementation of stricter management measures to regulate fishing operations in the early

1990s, an increasing number of historically overexploited fish stocks began showing signs of

recovery (Hilborn *et al.* 2020; Juan-Jordá *et al.* 2022). These management interventions,

however, coincided with accelerated changes in the oceans (Barange *et al.* 2018). Global climate

is becoming increasingly warmer and more variable, modifying regional and local habitat

conditions that support the biomass production of fish stocks, possibly shifting biologically

sustainable levels of harvesting (Chassot *et al.* 2010; Free *et al.* 2019).

The dynamics of marine resource populations in many systems are shaped by drivers operating

interactively at varied timescales, including high-frequency perturbations like fishing and low-

frequency variability in ocean properties forced by climate (Rouyer *et al.* 2012; Goto *et al.*

2022b). When exposed to such persistent, nonstationary perturbations, near-term (transient)

dynamics of an exploited population, which depend on both variable vital rates (survival, growth,

and fecundity) and demographic (age and stage) structures, may deviate from asymptotic

(equilibrium) dynamics (Hastings 2004; Koons *et al.* 2016). Past research instructs us that large-

scale shifts in ecosystem productivity under alternative climate regimes can be propagated

through the system to influence fluctuations in fisheries catches (Chassot *et al.* 2010; Cheung,

Watson & Pauly 2013). Climate effects on marine species are often integrated over a series of

oceanic, ecological, and biological processes over time and space (Ottersen *et al.* 2001; Stenseth

et al. 2003). Fluctuations in population structure and size can, for example, emerge with some

time lag from age (or stage)-dependent demographic processes like birth and death (Jenouvrier *et*

al. 2022) modulated by seasonal and interannual variations in weather and habitat conditions

(Hutchings & Reynolds 2004; Lindegren *et al.* 2013). These transient fluctuations in the

population could sustain over generations through delayed feedbacks (Hastings 2004). In

contrast, fishing operations and management actions that regulate the operations can directly

modify short-term trends and variability in population structure and demographic rates, and thus

the transient dynamics of exploited species (Gamelon, Sandercock & Sæther 2019). Large-scale

commercial fishing often selectively removes larger, older members of the population, truncating

age (and size) structure, modifying life history traits, reducing the capacity to absorb the effects

of perturbations, and thus amplifying year-to-year variation in population size in variable

environments (Hsieh *et al.* 2006; Rouyer *et al.* 2012). Fishing-induced age truncations

(juvenescence) are widely reported for harvested marine populations (Hsieh *et al.* 2010; Rouyer

et al. 2012), many of which suffer from reduced reproductive capacity (Hixon, Johnson &

Sogard 2014). Climate-induced changes in the variability of ocean habitat conditions thus may

indirectly promote or impede the recovery of depleted populations with weakened demographic

resilience to perturbations (Lindegren *et al.* 2013; Harris *et al.* 2018).

In commercial fisheries, a harvested population is commonly managed through regulation of

fishing pressure based on its population status (Beddington, Agnew & Clark 2007). Because of

high socioeconomic values, exploitation of marine resource populations is regularly monitored

through scientific surveys and assessed based on population size estimates (Hilborn & Walters

1992). These population assessments form the scientific basis for the implementation of

management measures like catch limits. When population size declines below a predefined

threshold (a reference point set under the precautionary principle, ICES 2021; Goto *et al.* 2022a),

for example, fishing pressure may be adjusted to mitigate the risk of overharvesting. This

adaptive management cycle (human–natural system feedbacks through monitoring, assessment,

and management action) thus can be considered as natural experiment (Hilborn & Walters 1992;

Jensen, Branch & Hilborn 2012), providing opportunities to gain insights into the transient

dynamics of managed populations in variable environments. Understanding what drives these

transients would help develop robust strategies to sustainably manage human exploitation of

marine resource populations as we face increasingly uncertain conditions in the oceans.

This article reports a study that empirically explores how past fishing, management action, and

climate variability have influenced the transient dynamics of recovering fish populations in the

northeast Atlantic Ocean (FAO Major Fishing Area 27, Fig. 1a). These populations experienced

wide ranges of fishing pressure, climate, and management action in recent decades (Hidalgo *et

al.* 2011; Rouyer *et al.* 2011; Zimmermann & Werner 2019). Annual fishing pressures applied to

many northeast Atlantic stocks began to be eased in the late 1990s to rebuild biomass that

support sustainable harvesting (Sparholt, Bertelsen & Lassen 2007; Lassen, Kelly & Sissenwine

2014). As a result, these stocks experienced considerable changes in population size,

demographic structure, and productivity following the reductions in fishing pressure (Hidalgo *et

al.* 2012; Rouyer *et al.* 2012; Britten, Dowd & Worm 2016). During these periods the regional

climate also became increasingly warmer and more variable as greenhouse gas emissions rose,

which was reflected particularly in upper ocean heat content and transport (Delworth *et al.* 2016;

Zanna *et al.* 2019). Warming patterns, however, highly varied among geographic subregions

(ecoregions) owing to local air–sea feedbacks (Roemmich *et al.* 2015; Delworth *et al.* 2016),

triggering a series of oceanographic and ecological responses at variable rates (Beaugrand 2009;

Delworth *et al.* 2016). To glean insights into the recovery patterns of historically depleted fish

stocks, this study specifically asks how local and regional climate variability may contribute to

variations in the reproductive performance and transient growth of marine species exposed to

variable fishing pressures moderated by management actions.

Materials and Methods

Data

Time series (1946–2016) of annual estimates of age-specific information (mean body mass, abundance, maturity rate, and natural and fishing mortality rates) and management measures (reference points) of marine fishes in the northeast Atlantic Ocean were extracted from the 2017 stock assessment reports by the International Council of Exploration of the Sea (ICES) (Table 1). In the assessment of each stock (defined here as a population in a management unit area defined by the ICES) the demographic rates and life history traits were estimated using data from commercial catch and scientific monitoring surveys. Stock sizes and fishing mortality rates then were estimated annually using age-structured population models fitted to time series data of abundance indices (ICES 2017e; ICES 2017a; ICES 2017b; ICES 2017c; ICES 2017f; ICES 2017d). The reliability of the models and data had been evaluated by experts prior to assessments and extensively documented (e.g., ICES 2016). For analysis I excluded stocks with missing age-specific information, rejected assessments, or shorter time series (less than 15 years). The final dataset consists of 38 stocks comprising 11 species (cod *Gadus morhua*, haddock *Melanogrammus aeglefinus*, saithe *Pollachius virens*, herring *Clupea harengus*, plaice *Pleuronectes platessa*, turbot *Scophthalmus maximus*, sole *Solea solea*, whiting *Merlangius merlangus*, sprat *Sprattus sprattus*, megrim *Lepidorhombus spp.*, and four-spot megrim *Lepidorhombus boscii*) across seven ecoregions in the northeast Atlantic Ocean defined by the ICES: the Norwegian and Barents Seas (combined as a single ecoregion for analysis), the Greater North Sea, the Baltic Sea, Faroes, Icelandic waters, the Bay of Biscay and the Iberian Coast (hereafter the Bay of Biscay), and the Celtic Seas (Table 1 and Fig. 1a). These stocks

support some of the largest fisheries in the region, collectively yielding 1.9 to 2.6 million tonnes

of commercial landings annually in the recent decades (1997–2016).

Climate indices and sea surface temperatures (SST) were used as proxies for regional (North

Atlantic) and local (ecoregion) climate conditions (1946–2016). Monthly time series of three

regional climate indices were downloaded and averaged over winter (most active) months for

analysis (Fig. 1b): the North Atlantic Oscillation (NAO) and Atlantic Multidecadal Oscillation

(AMO) indices (both computed from the Hadley Centre Sea Ice and Sea Surface Temperature

data set [HadISST1, https://www.metoffice.gov.uk] and the extended reconstructed SST

[ERSSTv5, NOAA Physical Sciences Laboratory; https://psl.noaa.gov] and averaged over the

sources), and upper (0–700 m) ocean heat content (OHC) anomaly in the North Atlantic (NOAA

National Centers for Environmental Information; https://www.ncei.noaa.gov/). The NAO index

reflects the dominant mode of atmospheric variability between the Azores and Iceland, linked

with a range of large-scale oceanic features (such as surface temperature and heat content) in the

North Atlantic that regulate biological and ecosystem processes (Stenseth *et al.* 2003). The AMO

index, computed by averaging SSTs over the North Atlantic (25°–60°N at 7°–70°W) using the

method by van Oldenborgh *et al.* (2009), also reflects large-scale climate variability that may

influence the movement and migration of marine species (Faillettaz *et al.* 2019). OHC measures

the amount of excess heat (from increasing greenhouse gases) trapped in the ocean, integrating

basin-wide heat storage and transport (Zanna *et al.* 2019). Because of high signal-to-noise ratio,

OHC reflects more consistent decadal trends (less influenced by natural variability) than other

climate indices (Roemmich *et al.* 2015).

Reconstructed gridded monthly SST data were downloaded from and averaged over three

sources to account for systematic uncertainties in each source: HadISST1, COBE Sea Surface

Temperature SST (NOAA Physical Sciences Laboratory), and ERSSTv5. I first averaged the

SST data over ecoregion and for each ecoregion I computed SST anomalies by subtracting the

mean across years and dividing by the standard deviation for each season (Fig. 1c); winter

(December in the previous year–February, $wSST_{anom}$), spring (March–May, $spSST_{anom}$), summer

(June–July, $smSST_{anom}$), and autumn (September–November, $aSST_{anom}$).

Transient population growth and elasticity

I used transient population growth rate as an integrated measure that captures time-varying

demographic rates and age structures responding to non-stationary extrinsic factors (fishing

pressure, management action, and climate) and estimated the growth rates for each stock using

an age-structured matrix projection model (Caswell 2001). Leslie (state-transition) matrices (A)

were first constructed with annual estimates of age-specific survival and reproductive rates.

Survival rates ($S_{a,y}$) were estimated as $S_{a,y} = \exp(-Z_{a,y})$, where $Z_{a,y} = F_{a,y} + M_{a,y}$, and $F_{a,y}$ and $M_{a,y}$

are fishing and natural mortality (non-fishing like predation and starvation) rates for a-year olds

in year y. Egg production rates are often not estimated in the assessments of marine fish stocks,

and here I estimated recruitment success (integrating over reproduction and early life stage

survival) rate per spawner (reproductive fish) for a-year olds in year y ($RS_{a,y}$) (Rouyer *et al.*

2011) as a proxy for per capita fecundity rate as

178
$$RS_{a,y} = \frac{R_y m_{a,y-a_R}}{\sum_{a=a_{min}}^{a=a_{max}} m_{a,y-a_R} N_{a,y-a_R}} \tag{1}$$

where R_y is the recruit number in year y, $m_{a,y-a_R}$ and $N_{a,y-a_R}$ are the maturity rate (fraction of

juveniles becoming adults) and abundance of a-year olds in year y-a_R (spawning year), a_R is the

age of recruits, which accounts for variations in age at recruitment (0–3 years old) among stocks

reported in the assessments. a_{min} and a_{max} are the youngest and oldest ages in the stock. I then

numerically estimated deterministic transient growth rates (λ_{trans}) by projecting age-specific

abundances using the annual state-transition matrices and the annual estimates of age-specific

numbers taken from the assessments as

186
$$\lambda_{\text{trans,y}} = \frac{n_{y+T_{gen,y}}}{n_y}$$

187
$$\boldsymbol{n}_{y+Tgen} = \boldsymbol{n}_y \boldsymbol{A}_y \qquad (2)$$

188   where $\boldsymbol{n}_y$ and $\boldsymbol{A}_y$ are a vector of initial age-specific numbers and an $a \times a$ state-transition matrix

189   in year y, and T_{gen} is the generation time (years) in year y. To account for variations in life-

190   history strategy among stocks, the transient growth rates of each stock were estimated over its

191   generation time, which is defined as the mean age of parents of a cohort (Caswell 2001),

192
$$T_{\text{gen},y} = \frac{\sum_a a(\prod_{j=1}^{a-1} S_{j,y}) R_{a,y}}{\sum_a (\prod_{j=1}^{a-1} S_{j,y}) R_{a,y}} \qquad (3)$$

The transient growth rates here were thus defined in the timescales of 2.6 to 6.7 years (Fig. S1),

relevant for fisheries management.

To evaluate the contributions of demographic variations I performed transient life table

response experiments (LTRE) by numerically estimating transient elasticities (e_y) to survival and

fecundity rates through matrix projections (Koons $et\ al.$ 2016); demographic parameters

(fecundity or survival) of all ages were first perturbed by a small amount (1%), and stock

numbers were projected to re-compute $\lambda_{\text{trans},y}$ (Durant $et\ al.$ 2013). Elasticity (relative sensitivity),

defined as the proportional change in λ_{trans} relative to a proportional change in a demographic

rate (Caswell 2001), was computed as $e_y = \frac{\Delta \log \lambda_{trans,y}}{\Delta \log \alpha_y}$, where α_y is the survival or fecundity rates

in year y that are perturbed (Coulson $et\ al.$ 2004). Age-specific elasticities were then summed for

analysis; changes in elasticities to juvenile and adult survival rates were evaluated separately by

summing age-specific elasticities weighted by maturity rates. To quantify how the transient

dynamics deviate from expected long-term equilibriums (asymptotic dynamics assuming stable

environments and age structures) across years I also computed asymptotic (geometric) growth

rates and stable age distributions from the logarithms of the dominant (largest) eigenvalues and

the right eigenvectors (respectively) of the annual state-transition matrices (Caswell 2001).

To evaluate how variations in demographic structure have contributed to the recovery patterns

of fish stocks, I used mean age and age diversity of spawners as indices for demographic

structures. Mean age (SA_{mean}) is the weighted (by abundance) average of spawner age as

212
$$AS_{mean,y} = \frac{\sum_{a=a_{min}}^{a=a_{max}} a m_{a,y} N_{a,y}}{\sum_{a=a_{min}}^{a=a_{max}} m_{a,y} N_{a,y}} \tag{4}$$

The Shannon diversity index was used as an indicator of age diversity ($SA_{diversity}$) and computed

as $SA_{diversity,y} = -\sum_{a=a_{min}}^{n} p_{a,y} \ln(p_{a,y})$, where $p_{a,y}$ is the proportion of a-year-old spawner number

in year y (Marteinsdottir & Thorarinsson 1998).

Management actions (here adjustment of fishing pressures) are based on perceived current

stock status and harvest control rules set by the ICES (which are designed to achieve sustainable

harvesting without compromising the reproductive capacity of the stock). The current status is

assessed using scientific monitoring and catch data before the provision of annual catch advice

(total allowable catch). Under the control rules, when the estimated SB remains above a fixed

threshold (B_{pa}) at the start of the year following the assessment, the catch limit is computed based

on target exploitation rate (F_{msy}). These two parameters (B_{pa} and F_{msy}) are designed to prevent

overharvesting by accounting for uncertainty in population and harvest dynamics (ICES 2019).

When the SB falls below B_{pa}, exploitation rate is adjusted to F_{msy} scaled to the proportion of SB

relative to B_{pa}, thereby allowing the population to rebuild (adaptive management). For analysis

here fishing pressure and SB relative to stock-specific reference points ($\bar{F}/F_{msy}$ and SB/B_{pa}) were

used as proxies to evaluate the influence of the management measures on the transient dynamics

of fish stocks. Fishing pressure is defined as the instantaneous rate of fishing mortality ($\bar{F}$),

which is commonly estimated by averaging age-specific fishing mortality rates over dominant

ages in fishery catch (Table 1).

Data analysis

To explore how management-guided changes in fishing pressure and climate change have

influenced the transient dynamics and recovery patterns of the northeast Atlantic fish stocks, I

applied multilevel (hierarchical) modeling to derived demographic metrics to account for

nonindependence in data structure (Gelman & Hill 2006) with stock, ecoregion, and year as

grouping factors (random intercepts). Data exploration including identification of collinearity

(through visual inspection correlation plots and variance inflation factors) and outliners (>three

standard deviations) was performed before analysis (Zuur *et al.* 2009). The data exploration

revealed two groups of stocks with diverging (increasing and declining) patterns in stock status

based on relative spawner biomass (SB/B_{pa}). To explore mechanisms underlying the differences I

applied generalized additive mixed effects models (GAMMs, Wood 2017) to these groups

separately. In the following analyses the stocks were grouped based on stock status in the last 15

years of the time series (2002–2016); if spawner biomass is larger than 80% of a biological

threshold (B_{pa}) for at least 50% of the period (eight years) a stock is classified as 'recovering',

otherwise 'non-recovering'. This classification resulted in 27 recovering and 10 non-recovering

stocks (Table 1). Trend analyses were first performed with year as a covariate to capture shared

long-term historical patterns in relative spawner biomass, spawner demographic structure (mean

age and age diversity), relative fishing pressure ($\bar{F}/F_{\text{msy}}$), and transient growth. The response

variables were assumed to follow a Gamma distribution with continuous, positive expected

values μ and a logarithmic link function as

251                             $$Y_{jk} \sim \text{Gamma}(\mu_{ijk},\ \tau)$$

$$E(Y_{ijk}) = \mu_{ijk}$$

$$\text{var}(Y_{ijk}) = \mu_{ijk}{}^2/\tau$$

$$\log(\mu_{ijk}) = \beta_0 + f_1(\text{Year}_{ij}) + \text{stock}_i + \text{ecoregion}_j + \text{year}_{jk}$$

$$\text{stock}_i \sim N(0, \sigma^2_{stock}), \text{ecoregion}_j \sim N(0, \sigma^2_{ecoregion}), \text{year}_{jk} \sim N(0, \sigma^2_{year}) \qquad (5)$$

where Y_{jik} is the response of stock i in ecoregion j in year k, τ is a Gamma distributed parameter,

β_0 is the intercept, and f_1 is a smoother function (thin-plate cubic splines with ten maximum

degrees of freedom or knots) for year; stock_k, ecoregion_i, and year_{jk} are normally distributed

random intercepts with mean zero and standard distribution σ for stock, ecoregion, and year

nested within ecoregion. The structure of random effects was selected based on likelihood ratio

tests applied to ten structures fitted with a restricted maximum likelihood method (REML)

(Table S1). Further, I assessed direct effects of fishing on spawner biomass and age structure by

evaluating relationships between relative fishing pressure and relative biomass and age structure

of spawners as

$$Y_{jk} \sim \text{Gamma}(\mu_{ijk}, \tau)$$

$$E(Y_{ijk}) = \mu_{ijk}$$

$$\text{var}(Y_{ijk}) = \mu_{ijk}{}^2/\tau$$

$$\log(\mu_{ijk}) = \beta_0 + f_1(\text{Fishing}_{ij}) + \text{stock}_i + \text{ecoregion}_j + \text{year}_{jk}$$

$$\text{stock}_i \sim N(0, \sigma^2_{stock}), \text{ecoregion}_j \sim N(0, \sigma^2_{ecoregion}), \text{year}_{jk} \sim N(0, \sigma^2_{year}) \qquad (6)$$

where $Y_{jik}, \tau, \beta_0, \sigma_{stock}, \sigma_{ecoregion}$, and σ_{year} are the same as above. f_1 is a smoother function for

relative fishing pressure.

The contributions of fishing, management action, and climate to the transient dynamics of fish

stocks with variable age structures were evaluated in two steps. I first tested for time-varying

contributions of these extrinsic factors to variations in vital rates and then evaluated how these

factors influenced the elasticities of transient population growth:

Covariates of recruitment variability. Because natural mortality rates were assumed time-

invariant in most of the assessments, variations in survival rates are in effect driven by fishing.

Here, RS was thus selected as a demographic response variable (which integrates probabilities of

survival (and maturation) to adult and spawning) for analysis. Further, because RS can vary with

variations in spawner age structure and biomass (intrinsic factors), these intrinsic factors were

also tested. To test for the time-varying contributions of the extrinsic and intrinsic factors, the

following model was fitted to the data:

283
$$\mathrm{RS}_{\mathrm{total}\,ijk} \sim \mathrm{Gamma}(\mu_{ijk},\ \tau)$$

284
$$E(\mathrm{RS}_{\mathrm{total},ijk}) = \mu_{ijk}$$

285
$$\mathrm{var}(\mathrm{RS}_{\mathrm{total},ijk}) = \mu_{ijk}{}^2/\tau$$

286
$$\log(\mu_{ijk}) = \beta_0 + f_1(\mathrm{Year}_{ij}) + f_2(\mathrm{Year}_{ij}, \mathrm{Biomass}_{ijk}) + f_3(\mathrm{Year}_{ij}, \mathrm{Demography}_{ijk}) +$$

287
$$f_4(\mathrm{Year}_{ij}, \mathrm{Fishing}_{(i-1)jk}) + f_5(\mathrm{Year}_{ij}, \mathrm{Climate}_{ij}) + \mathrm{stock}_i + \mathrm{ecoregion}_j + \mathrm{year}_{jk}$$

288
$$\mathrm{stock}_i \sim N(0, \sigma^2_{stock}), \ \mathrm{ecoregion}_j \sim N(0, \sigma^2_{ecoregion}), \ \mathrm{year}_{jk} \sim N(0, \sigma^2_{year}) \qquad (7)$$

where $\mathrm{RS}_{\mathrm{total},y}$ is the total per capita recruitment success rate of stock i in ecoregion j in year k, β_0

is the intercept, f_1–f_5 are smoother functions for covariates: year, year × density, year ×

demographics (spawner mean age and age diversity), year × climate, and year × fishing. *Climate*,

Biomass, and *Demography* were lagged 0–3 years (depending on stock) to align with climate

conditions and spawner status (biomass and age structure) in spawning year, whereas *Fishing*

was lagged by an additional year (spawning year–1) to reflect indirect effects of fishing on

recruitment success that are not accounted for by variations in spawners. Regional climate

covariates were also lagged an additional year (*Climate*$_{(i-1)j}$) to reflect indirect effects through

regulation of local climate and ecosystem processes. Time (year)-varying covariate smoothers

were tested as tensor product construction. For interpretability and model convergence one

covariate of each type was tested at a time; one local or regional climate index (see above) for

climate, SB or SB/B_{pa} for spawner density, $\bar{F}$ or $\bar{F}/F_{msy}$ for fishing pressure, and SA$_{mean}$ or

SA$_{diversity}$ for demographics.

Covariates of elasticity to recruitment variability. The transient dynamics of depleted fish

stocks with age truncation often become recruitment-dependent and thus sensitive to variations

in climate and habitat conditions. As the stocks recover (increases in abundance and age

diversity), the transient dynamics are expected to become less sensitive to recruitment variability.

To test for time-varying contributions of the extrinsic and intrinsic factors to variations in

elasticity to RS (e_{rec}), the following model was fitted to the data, assuming e_{rec} follows a Beta

distribution with expected values μ ($0 < \mu < 1$) and a logistic link function as

$$e_{rec,ijk} \sim \text{Beta}(\mu_{ijk}, \phi)$$

$$E(e_{rec,ijk}) = \mu_{ijk}$$

$$\text{var}(e_{rec,ijk}) = \mu_i (1 - \mu_i)/(\phi + 1)$$

$$\text{logit}(\mu_{ijk}) = \beta_0 + f_1(\text{Year}_{ij}) + f_2(\text{Year}_{ij}, \text{Biomass}_{ijk}) + f_3(\text{Year}_{ij}, \text{Demography}_{ijk}) +$$

$$f_4(\text{Year}_{ij}, \text{Fishing}_{(i-1)jk}) + f_5(\text{Year}_{ij}, \text{Climate}_{ij}) + \text{stock}_i + \text{ecoregion}_j + \text{year}_{jk}$$

$$\text{stock}_i \sim N(0, \sigma_{stock}^2), \text{ecoregion}_j \sim N(0, \sigma_{ecoregion}^2), \text{year}_{jk} \sim N(0, \sigma_{year}^2) \qquad (8)$$

where ϕ is a parameter (> 0), and $\beta_0, f_1-f_5, \sigma_{stock}, \sigma_{ecoregion}$, and σ_{year}, are the same as above.

All numeric covariates were standardized by subtracting the mean and dividing by the

standard deviation (z-scores). In model selection, intrinsic covariates were first added, followed

by indirect fishing, and then climate. With the random effect structure selected (above) the

models were tested sequentially for covariates by fitting with a maximum likelihood method,

followed by parameter (re)estimation with REML (Zuur *et al.* 2009) for the model with the

lowest Akaike information criterion (AIC) score and highest model weight (*w*). AIC is computed

as AIC $= -2\ln(L) + 2k$, where L is the likelihood of the model and k is the number of parameters,

and the model weight of model i (w_i) is computed as

$$w_i = \frac{\exp(-\Delta AIC_i / 2)}{\sum_j \exp(-\Delta AIC_i / 2)}$$ (9)

As diagnostics, model residuals were checked for homogeneity and regressed against

covariates that were both included and not included in the selected models (Zuur *et al.* 2009). All

analyses were performed in R (version 4.2.1, R Development Core Team 2022) with the R

packages of *gamm4* (v.0.2-6, Wood & Scheipl 2020) and *mgcv* (v.1.8-40, Wood 2022) for

model-fitting, *mgcv* and *gratia* (v.0.7.3, Simpson & Singmann 2022) for model diagnostics, and

popbio (v.2.7, Stubben, Milligan & Nantel 2020) for computations of demographic parameters

and projections of population dynamics.

Because fish populations in different ecoregions experience a wide range of climate

conditions and often have adapted to local environments, fish–covariate patterns may deviate

from shared patterns captured by the regional models. To test this hypothesis, ecoregion-scale

analyses were also performed using the same model structures for RS_{total} and e_{rec} and the model

selection steps as above except for random effects. Because random effect estimates become less

stable when the number of groups is fewer than eight (Bolker 2015), the random effect of the

ecoregion-scale models consisted of year only, with stock being a fixed effect, except for Celtic

Seas; the random effects of the Celtic Seas model consisted of year and stock.

Results

Regionwide analyses showed that relative fishing pressures on the recovering stocks on

average peaked in the early 1990s ($\bar{F}/F_{msy} = \sim 2.2$) and near-linearly declined since ($\sim 2.1\%$ per

year; ΔAIC = 88.4), followed by increasing mean age (ΔAIC = 35.7), relative spawner biomass

(ΔAIC = 17.2), and age diversity (ΔAIC = 21.1) (Fig. 2a). In contrast, relative fishing pressures

on the non-recovering stocks kept rinsing until the late 1990s (to $\bar{F}/F_{msy}$ = ~3.7 on average)

before declining (~3.0% per year, ΔAIC = 63.3), but their relative spawner biomass continued to

decline (ΔAIC = 92.7 and 31.5; Fig. 2a). Although the mean spawner age of the non-recovering

stocks began rising with declining relative fishing pressures, no change in age diversity was

detected (Fig. 2a). Direct effects of selective fishing consistently emerged as negative

correlations between relative fishing pressure and spawner biomass and age structure in both

stock groups (ΔAIC = 15.1–165.0, Fig. 2b). Contrasting temporal patterns between relative

fishing pressure and spawner features were more pronounced in some ecoregions captured in

year-specific (nested within ecoregion) random effect (variance) estimates, especially the

Norwegian–Barents Seas, which showed abrupt shifts in the early 1990s (Fig. S2).

The time series of transient population growth rates reflected the contrasting patterns between

the recovering and the non-recovering stocks, and variations among ecoregions (Fig. 3 and S3).

The regional average growth rates of the recovering stocks linearly increased over the study

period (ΔAIC = 15.0, Fig. 3). Although the relative spawner biomass of the non-recovering

stocks continued to decline, their average growth rates began rising in the late 1990s (ΔAIC =

2.8, Fig. 3) when the relative fishing pressures began declining (Fig. 2a). These transient

dynamics differed from asymptotic dynamics; the transient growth rates were lower than

asymptotic growth rates in most stocks, and the deviations varied over years, with ~50% (19) of

the stocks showing increasingly less differences in the last decade (Fig. S4). The observed age

distributions of many of these stocks also differed from theoretical stable age distributions and

skewed toward younger ages (Fig. S5) owing to disproportionately higher annual mortality rates

of older fish by fishing (Fig. S6).

Regionwide analyses detected declining trends in per capita recruitment success rate in both

recovering and non-recovering stocks (ΔAIC = 2.1 and 2.7, Fig. 4a), with high among-stock

variances (σ^2_{stock} = 0.31 and 0.48). Ecoregion-specific random intercepts also revealed generally

higher year-to-year fluctuations in the non-recovering than in the recovering stocks (Fig. S3).

After accounting for variations among stocks, ecoregions, and years, the recruitment rates of

both stock groups declined with increasing relative spawner biomass (Fig. 4c and Table 2),

which was more supported by data than absolute biomass (Table S2). But the temporal patterns

of density dependency differed between the groups; the density dependent effect in the

recovering stocks increased as the biomass rose, whereas that in the non-recovering stocks

emerged in the last two decades but declined over time (Fig. 4c). Likewise, the contribution of

spawner age structure to recruitment variability differed between the stock groups; the

relationship between recruitment success and mean age in the recovering stocks shifted from

positive to negative in the early 1990s, whereas that in the non-recovering stocks remained

largely positive (Fig. 4b). An indirect positive fishing effect on recruitment rate was detected in

the recovering stocks until the late 1980s but gradually diminished since (Fig. 4d). The models

for both stock groups showed weak support for climate effects (Table S2). Of the climate indices

tested for the non-recovering stocks (whose model had slightly more support than the recovering

stocks), the model with ocean heat content anomaly in the year of spawning showed shifting

(positive to negative) climate effects in the recent decades (Fig. 4e).

Ecoregion-scale analyses on recruitment success revealed patterns largely consistent with the

regional analyses with some system-specific deviations (Fig. 5). In ecoregions with non-

recovering stocks like the Baltic, Celtic, and North Seas the density-dependent effect on

recruitment on average became weakened over time; in contrast, the effect in the Norwegian–

Barents Seas (with only recovering stocks) became strengthened (Fig. 5c). Correlations with

spawner age structure metrics were detected in all but the Bay of Biscay and Faroes (Fig. 5b); in

the Baltic Sea a positive correlation with age diversity became detectable in the early 1990s

when fishing pressures began declining (Fig. 2). The time-varying effect of spawner age was

especially pronounced in the Norwegian–Barents Seas stocks, showing a shift from positive to

negative in the 1990s (Fig. 5b). An indirect fishing effect was (weakly) supported only in the

North Sea stocks (Table S3 and Fig. 5d). Climate anomalies (in the year of spawning)

contributed to variability in recruitment success in all ecoregions except the North Sea but the

indices with the most statistical support varied among ecoregions (Fig. 5e and Table S3) despite

high correlations among climate indices (Fig. S7). The ways climate anomalies contributed to

recruitment variability also differed (Fig. 5e). In the Baltic Sea and Faroes correlations with

climate anomalies remained largely positive during the past few decades. In contrast, the positive

effect in the Norwegian–Barents Seas began declining in the recent decades. In the Bay of

Biscay and Icelandic waters the climate effect shifted from positive to negative.

Trend analyses revealed consistent temporal patterns in elasticities to vital rates for the

recovering and non-recovering stocks. Both showed declining elasticities to (post-recruitment)

juvenile survival rate (ΔAIC = 15.6 and 18.9) and increasing elasticities to adult survival rate

over time (ΔAIC = 17.0 and 6.7) (Fig. S8). But the regionwide analyses detected a declining

trend in elasticities to recruitment success rate in the recovering stocks only (ΔAIC = 3.6, Fig.

6a). Ecoregion-specific variances in all the models fluctuated over years, but on average

elasticities to adult survival showed higher deviations from the regional trends than the other

vital rates (Fig. S8).

Regionwide analyses on elasticities to recruitment success largely reflected the time-varying

responses in recruitment to intrinsic and extrinsic drivers. For the recovering stocks the model

indicated that more diverse ages increasingly contributed to elasticities in more recent years,

whereas the contributions of mean ages gradually increased in the non-recovering stocks (Fig.

6b). Elasticities to recruitment declined with higher spawner biomass in both stock groups, but

the density-dependent effect in the non-recovering stocks diminished over time (Fig 6c). The

model with an indirect fishing effect in both stock groups showed a declining contribution to

variations in the elasticities after the reductions in fishing pressure (Fig. 6d). In contrast, the

models for both stock groups indicated an increasingly more positive climate effect (with the

AMO index and winter sea temperature anomaly) over time (Fig. 6e).

Ecoregion-scale analyses on elasticities to recruitment success detected declining trends in all

but the Bay of Biscay and the Norwegian–Barents Seas (Fig 7a), which showed little change in

trends. The analyses also indicated spatial heterogeneity in the ways time-varying intrinsic and

extrinsic drivers contributed to variations in the elasticities. For example, shifting contributions

of spawner age structure were detected in most ecoregions, showing both diminishing (e.g.,

Faroes and North Sea) and increasing (e.g., Baltic Sea and Icelandic waters) trends (Fig. 7b).

Negative correlations with spawner biomass were consistently detected except in the Baltic Sea

(Fig. 7c). This maternal effect increased over time except in the Norwegian–Barents Seas, which

showed the opposite trend (Fig. 7c). The models with an indirect fishing effect had statistical

support for Faroes, and the Celtic, North, and Norwegian–Barents Seas; the fishing effect

declined and/shifted over time except the Norwegian–Brents Sea, which showed an increasing

contribution of fishing since the 1990s (Fig. 7d). In the Baltic Sea and the Bay of Biscay a

positive climate effect emerged in the 2000s (Fig. 7e), reflecting the temporal patterns of shifting

local climate effects on recruitment success (Fig. 5e). In Faroes and Icelandic waters, where

climate anomalies (the NAO index and spring sea temperature) had a positive effect on

recruitment, the elasticities were inversely correlated with the indices, but this effect shifted to

the opposite in the 2000s (Fig. 7e).

Discussion

Analyses on the transient dynamics of northeast Atlantic fish stocks reveal that management-

aided reductions in fishing pressure and shifting climate conditions nonlinearly shaped

rebuilding patterns of these historically depleted stocks through restorations of demographic

structure (reversing age truncation) and reproductive capacity in recent decades. As the survival

rates and demographic structures of reproductive fish improved with declining fishing pressures,

transient population growth rates became less sensitive to variability in recruitment and juvenile

survival and more to that in adult survival, allowing the stocks to rebuild biomass despite

declining recruitment success. As the biomass of reproductive fish rebuilt, recruitment success

rates also became increasingly regulated by density-dependent intrinsic processes involving

increasingly higher numbers of older reproductive fish. Although warmer than average climates

promoted recruitment success through the early 2000s in some ecoregions, warming trends over

the past decades began adversely affecting reproductive performance. Variations in regulation of

fishing pressure among stocks also led to diverging trends in recoveries of the stocks exposed to

a range of local climate conditions. Insufficient or delayed reductions in fishing pressure can

further deplete a stock and erode reproductive fish demographic structure, thereby diminishing

fecundity and thus recovery potential, while also remaining sensitive to climate variability.

Rebuilding biomass by accounting for demographic structure through management are thus not

only key to sustainable use of marine resource populations but also the foundation for effective

applications of adaptive measures resilient to future environmental changes.

Transient dynamics of recovering populations from intense fishing pressures

Fishing-induced changes in the demography and reproductive capacity of an exploited

population can be reversible when management measures are effectively applied to regulate

fishing pressures. This pattern is displayed particularly in the Norwegian–Barents Seas, where

fishing pressures were reduced by more than 60% (compared to the regional average of ~50%)

since the early 1990s, followed by sharp rises in the average age and biomass of reproductive

fish. But when a population experiences long-term intensive harvesting, recovery time can

become longer (Hutchings 2015) and more uncertain (Neubauer *et al.* 2013), with the population

remaining in long transient states (Hastings *et al.* 2018). Although fishing pressures on many

northeast Atlantic fish stocks precipitously declined through stricter management measures

applied since the late 1990s, demographic restorations and stock recoveries on average

progressed gradually. Reducing fishing pressures can allow more adults to survive and contribute

to the production of new recruits but the restoration of the demographic structure takes place

over generations and may take longer (a decade or more) for severely depleted stocks. Further,

although the declining trends in reproductive fish biomass halted in the early 2000s in most

ecoregions, reduction levels in fishing pressure varied among ecoregions and stocks and were

reflected in recovery patterns. When reductions in fishing pressure were not sufficient or

delayed, the declining trends in reproductive capacity and deterioration of demographic structure

in some stocks continued. Slow recoveries of severely depleted fish stocks are reportedly

widespread (Hutchings 2000). Regionwide analyses on the non-recovering stocks in the

northeast Atlantic reveal that sustaining high fishing pressures for even a few more years can

result in vastly different recovery patterns, with further age truncations and reductions in

reproductive capacity, which could further diminish fishery yields (Hilborn *et al.* 2020).

Removal of some biomass from a population by harvesting can promote its productivity

(Neubauer *et al.* 2013) by attenuating density-dependent effects on early life stage processes

(competition for food and space for example, Anderson *et al.* 2008). Intense fishing pressures

before the implementation of stricter management measures–mediated through reproductive

fish–may have led to higher recruitment rates in depleted stocks in the northeast Atlantic Ocean.

Fishing not only truncates age structure but induces changes in life-history traits that may

influence recovery (Neubauer *et al.* 2013; Hutchings & Kuparinen 2020). The generation times

of these depleted northeast Atlantic stocks were shortened (by 14 to 50%), which is often

associated with high year-to-year variability in recruitment (Minto, Myers & Blanchard 2008)

and high average recruitment rates (Bjørkvoll *et al.* 2012). Higher growth rates of depleted (and

age-truncated) fish populations also likely benefited during the early stages of recovery as more

reproductive fish contributed to spawning events, thereby increasing the probability of

recruitment success. These indirect effects, however, diminished over time following the

reductions of fishing pressures. Despite the signs of recovery in reproductive fish, per capita

recruitment success rates on average declined in the northeast Atlantic stocks, as also reported in

other Atlantic ecoregions (Britten, Dowd & Worm 2016). But these apparent patterns were

shaped by different processes among stocks. In the recovering stocks density dependency

remained present in recruitment success throughout the past decades and its strength increased as

reproductive fish biomass rose, likely slowing recoveries as fishing pressures declined. In

contrast, in the non-recovering stocks density dependency in recruitment success emerged only

after the reductions in fishing pressure but its strength continued to decline with reproductive fish

biomass, suggesting contributions of other (extrinsic) drivers.

Density dependency in early life stages also perhaps explains nonlinear relationships (shifting

from positive to negative) between reproductive fish age and per capita recruitment success in

the recovering stocks (especially in the Norwegian–Barents Seas) following the reductions in

fishing pressure. Higher mean age and age diversity in reproductive fish are expected to promote

recruitment success (Rogers & Dougherty 2019). Age-dependent variations in spawning pattern

can influence egg viability and larval survival; older spawners tend to produce more viable

offspring (Hixon, Johnson & Sogard 2014). But the analyses on the northeast Atlantic fishes

show that only the non-recovering (age-truncated) stocks kept positive relationships between

reproductive fish age structure and recruitment success. In the recovering stocks, density

dependency may have played a larger role than maternal effects, revealing a shift in

compensatory processes. As adult survival improves and population density increases, pre-recruit

survival may decline (owing to competition for food and other resources including cannibalism

by adults), contributing to apparent increases in average ages of reproductive fish.

Short-term (year-to-year) fluctuations in recruitment can propagate through transient dynamics

of a population when its (demographic) buffering capacity is compromised by age truncation

(Anderson *et al.* 2008). Truncated age distributions in reproductive fish may produce volatile

population behaviors that reflect recruitment pulses (Anderson *et al.* 2008). As adult survival

improved and stock biomass rebuilt in many northeast Atlantic fish stocks, the transient growth

rates became less sensitive to recruitment variability and more sensitive to adult survival. The

regionwide analyses show that increases in transient growth rate and biomass during the past few

decades resulted largely from improved survival rates of older fish. Greater sensitivities to adult

survival also may make harvested populations more manageable. Nonlinear responses to age

structure in reproductive fish is however also evident in sensitivity to recruitment variability.

Although the buffering effect of increasingly age-diverse reproductive fish emerged in many

ecoregions especially in the early stages of recovery, this relationship also shifted as the

recoveries further progressed (e.g., the Norwegian–Barents Seas), likely responding to other

intrinsic dynamics.

Climate regulation of demographic variations

Fish populations experiencing persistent selective depletions of older adults often become

sensitive to variations in early life stage processes that reflect signals in environmental

conditions like climate variability (Ottersen, Hjermann & Stenseth 2006; Rouyer *et al.* 2011).

Warmer than average climate conditions in the northeast Atlantic Ocean increasingly contributed

to observed patterns in fish recruitment success, reflecting shifting climate effects on recruitment

processes. For example, stratification may become intensified as climate warms, limiting nutrient

movement to deeper waters and food accessibility for the early life stages of demersal and

benthic species (Moore *et al.* 2018). At the region scale, however, the models captured only

weak responses in recruitment by the fish stocks to regional climate signals likely because of

heterogeneous warming patterns among ecoregions, as also seen in other parts of the world

(Barange *et al.* 2018). Region-scale analyses may mask differential effects of local climate and

oceanographic features (climate-induced shifts in ocean heat content and primary productivity

for example) that marine species are adapted to and experience (Conover 1992).

Ecoregion-scale variability in recruitment success may better reflect how fish reproductive

behavior responds to regional climate mediated by local habitat conditions, as hypothesized in

past research (Myers 1998; Ottersen & Stenseth 2001). For stocks in high latitude systems like

the Norwegian–Barents Seas and Icelandic waters, for example, climate conditions in winter–

spring (December–May) that regulate phonologies like timings of spawning migration (Sundby

& Nakken 2008) contributed to recruitment variability. Warming-induced reductions in sea ice

extent in these ecoregions promoted ecosystem productivity in the late 1990s and early 2000s

and may have benefited upper trophic animals like fish (Stenevik & Sundby 2007), likely

through changes in growth rate and life stage duration (Houde 2016). When spawning times

coincide with the period of high prey abundance, pre-recruit fish can grow more and experience

less size-dependent mortalities (like predation), contributing to greater recruitment success

(Peck, Huebert & Llopiz 2012). In contrast, in semi-enclosed systems like the Baltic Sea

warming-induced changes in nearshore habitat conditions in summer (June–August) (lower

salinity caused by higher precipitation and freshwater inflows for example, MacKenzie *et al.*

2007) may have played a greater role in regulating recruitment success through biophysical

processes during early life-stages.

Demographic responses by fish populations to changing climate conditions also may vary

discontinuously (Brander 2007). Local and regional climate conditions (both favorable and

unfavorable) that early life stage fish experience can persist through time (Ottersen *et al.* 2001)

and may emerge as delayed effects on transient population dynamics. In some northeast Atlantic

ecoregions the ways that climate affects fish recruitment success shifted from positive to

negative and may have slowed their recoveries. These shifts suggest that even in the ecoregions

where warming climate promoted stock productivity in the early twentieth century, fish began to

experience adverse (direct and indirect) effects. Mobile species can adjust their movement

(deeper and poleward) to mitigate physiological stress from warming upper water columns

(Pinsky *et al.* 2013; Engelhard, Righton & Pinnegar 2014). But asynchronous shifts in spatial

distribution by spawning fish and seasonal events in lower trophic groups like plankton booms

(phenological mismatches) also could trigger a series of events unfavorable for early life fish

(Asch & Erisman 2018), shrinking the probability of recruitment success.

Climate variability that regulates local ocean conditions may propagate through vital rates to

shape transient population dynamics of marine species (Jenouvrier *et al.* 2018). Depleted

populations in particular may have heightened sensitivities to adverse environmental conditions

(Hidalgo *et al.* 2011; Rouyer *et al.* 2011), which are becoming increasingly more variable in

recent decades (Rathore *et al.* 2020). Although the sensitivities of transient growth to recruitment

variability in the northeast Atlantic fish stocks declined as they recovered in the recent decades,

the analyses reveal that increasingly warmer than average climate began influencing how

recruitment variability propagates through transient dynamics. For example, in high-latitude

systems like Faroes, where the sensitivity to recruitment variability negatively covaried with

climate, this stabilizing effect diminished (and shifted to the opposite) as the climate warmed. In

contrast, in the Baltic Sea, where the sensitivity to recruitment variability positively covaried

with climate, this amplifying effect became strengthened as the climate warmed. Climate effects

on transient population dynamics also may, however, depend on the status (including

demographic structure) of fish stocks. In the Norwegian–Barents Seas, where fish stocks

experienced higher reductions in fishing pressure early and thus higher biomass rebuilding rates

than other stocks, changes in climate conditions had little influence on the sensitivities of

transient growth.

Transients as an integrated measure of effects of fishing, management, and climate

Analysis of transient population dynamics can reveal how time-varying demographic structures

and vital rates contribute to recovery patterns of managed resource populations when also

exposed to low-frequency, nonstationary drivers like climate. Declined sensitivities to highly

variable demographic parameters like recruitment, as displayed by the northeast Atlantic fish

stocks, can provide a buffer against rising environmental variability. Increasing sensitivities to

adult survival further suggest that the stocks also are expected to become more responsive to

management actions through regulation of fishing pressure (enforcing positive feedback), and

future management plans would benefit from maintaining healthy (age-diverse) reproductive

populations of exploited species (Pinsky & Byler 2015). Combined with other management

measures (like use of marine protected areas and fish size limits, White *et al.* 2013; Hixon,

Johnson & Sogard 2014), restoring and maintaining the demographic structure of an exploited

species thus may enhance resilience against destabilizing effects of environmental variability

amplified by climate change.

Emerging non-negligible changes in demographic responses to rising climate variability

nevertheless may influence the efficacy of fisheries management. Although historical climate

conditions promoted the productivity of fish stocks in the northeast Atlantic Ocean, some stocks

in this region and elsewhere in the Northern Hemisphere already are experiencing adverse effects

of changing climate (Pershing *et al.* 2015; Free *et al.* 2019). As many biological responses to

climate-driven changes in surface temperature and other ocean properties are often nonlinear, we

would expect continued shifts in how climate influences marine resource populations and

management (Britten, Dowd & Worm 2016). Accounting for these climate-induced changes thus

may not only reduce uncertainties in assessment but also improve the performance of

management measures (Rouyer et al. 2011; Bahri et al. 2021).

Exploited marine fish species are an integral part of an ecosystem and changes in their

abundances and demographic structures may propagate through community and ecosystem

dynamics (Jennings & Kaiser 1998), triggering a series of changes in ecologically connected

species (Goto *et al.* 2020). As adult survival improves in historically overharvested populations,

ecological processes including resource competition and predation also may indirectly modify

vital rates and demographic structures of non-target species (van Gemert & Andersen 2018).

Improving our understanding of how warming and more variable climates continue to reshape

the transient dynamics (through changes in life history traits and demographics) would help

develop management plans to mitigate the risk of overexploitation and safeguard ecosystem

services against rising environmental variability.

Acknowledgements

This work was partially supported by the Institute of Marine Research's Reduced Uncertainty

in Stock Assessment (REDUS) project.

Data Availability Statement

The data that support the findings of this study are openly available in ICES Scientific

Reports at https://ices-library.figshare.com/, reference number ICES CM 2017/ACOM:06, ICES

CM 2017/ACOM:08, ICES CM 2017/ACOM:11, ICES CM 2017/ACOM:12, ICES CM

2017/ACOM:13, and ICES CM 2017/ACOM:21, and in the data archives of Met Office Hadley

Centre, and NOAA Physical Sciences Laboratory and National Centers for Environmental

Information at https://www.metoffice.gov.uk, https://psl.noaa.gov, and

https://www.ncei.noaa.gov.

References

Anderson, C.N., Hsieh, C.-h., Sandin, S.A., Hewitt, R., Hollowed, A., Beddington, J., May, R.M.

& Sugihara, G. (2008) Why fishing magnifies fluctuations in fish abundance. *Nature,* **452,**

835.

Asch, R.G. & Erisman, B. (2018) Spawning aggregations act as a bottleneck influencing climate

change impacts on a critically endangered reef fish. *Diversity and Distributions,* **24,** 1712-

1728.

Barange, M., Bahri, T., Beveridge, M.C., Cochrane, K.L., Funge-Smith, S. & Poulain, F. (2018)

Impacts of climate change on fisheries and aquaculture: synthesis of currrent knowledge,

adaptation and mitigation options. Rome, FAO. 628 pp.

Beaugrand, G. (2009) Decadal changes in climate and ecosystems in the North Atlantic Ocean

and adjacent seas. *Deep Sea Research Part II: Topical Studies in Oceanography,* **56,** 656-673.

Beddington, J.R., Agnew, D.J. & Clark, C.W. (2007) Current problems in the management of

marine fisheries. *Science,* **316,** 1713-1716.

Bjørkvoll, E., Grøtan, V., Aanes, S., Sæther, B.-E., Engen, S. & Aanes, R. (2012) Stochastic

population dynamics and life-history variation in marine fish species. *The American*

Naturalist, **180,** 372-387.

Bolker, B.M. (2015) Linear and generalized linear mixed models. *Ecological statistics:*

contemporary theory and application (eds G.A. Fox, S. Negrete-Yankelevich & V.J. Sosa),

pp. 309-333. Oxford University Press, Oxford, United Kingdom.

Brander, K.M. (2007) Global fish production and climate change. *Proceedings of the National*

Academy of Sciences, **104,** 19709-19714.

Britten, G.L., Dowd, M. & Worm, B. (2016) Changing recruitment capacity in global fish stocks.

Proceedings of the National Academy of Sciences, **113,** 134-139.

Caswell, H. (2001) *Matrix population models: construction, analysis, and interpretation,* 2 edn.

Sinauer Sunderland.

Chassot, E., Bonhommeau, S., Dulvy, N.K., Mélin, F., Watson, R., Gascuel, D. & Le Pape, O.

(2010) Global marine primary production constrains fisheries catches. *Ecology Letters,* **13,**

495-505.

Cheung, W.W., Watson, R. & Pauly, D. (2013) Signature of ocean warming in global fisheries

catch. *Nature,* **497,** 365-368.

Christensen, V., Guenette, S., Heymans, J.J., Walters, C.J., Watson, R., Zeller, D. & Pauly, D.

(2003) Hundred-year decline of North Atlantic predatory fishes. *Fish and fisheries,* **4,** 1-24.

Conover, D. (1992) Seasonality and the scheduling of life history at different latitudes. *Journal*

of Fish Biology, **41,** 161-178.

Coulson, T., Guinness, F., Pemberton, J. & Clutton-Brock, T. (2004) The demographic

consequences of releasing a population of red deer from culling. *Ecology,* **85,** 411-422.

Crowder, L.B., Hazen, E.L., Avissar, N., Bjorkland, R., Latanich, C. & Ogburn, M.B. (2008)

The impacts of fisheries on marine ecosystems and the transition to ecosystem-based

management. *Annual Review of Ecology, Evolution, and Systematics***,** 259-278.

Delworth, T.L., Zeng, F., Vecchi, G.A., Yang, X., Zhang, L. & Zhang, R. (2016) The North

Atlantic Oscillation as a driver of rapid climate change in the Northern Hemisphere. *Nature*

Geoscience, **9,** 509-512.

Durant, J.M., Hidalgo, M., Rouyer, T., Hjermann, D.Ø., Ciannelli, L., Eikeset, A.M., Yaragina,

N. & Stenseth, N.C. (2013) Population growth across heterogeneous environments: effects of

harvesting and age structure. *Marine Ecology Progress Series,* **480,** 277-287.

Engelhard, G.H., Righton, D.A. & Pinnegar, J.K. (2014) Climate change and fishing: a century

of shifting distribution in North Sea cod. *Global Change Biology,* **20,** 2473-2483.

Faillettaz, R., Beaugrand, G., Goberville, E. & Kirby, R.R. (2019) Atlantic Multidecadal

Oscillations drive the basin-scale distribution of Atlantic bluefin tuna. *Science Advances,* **5,**

eaar6993.

FAO (1995) Precautionary approach to fisheries. Part 1: Guidelines on the precautionary

approach to capture fisheries and species introductions. *FAO Fisheries Technical Paper*, pp.

52 p., Rome, FAO.

Free, C.M., Thorson, J.T., Pinsky, M.L., Oken, K.L., Wiedenmann, J. & Jensen, O.P. (2019)

Impacts of historical warming on marine fisheries production. *Science,* **363,** 979-983.

Gamelon, M., Sandercock, B.K. & Sæther, B.E. (2019) Does harvesting amplify environmentally

induced population fluctuations over time in marine and terrestrial species? *Journal of*

Applied Ecology, **56,** 2186-2194.

Gelman, A. & Hill, J. (2006) *Data analysis using regression and multilevel/hierarchical models*.

Cambridge university press.

Goto, D., Devine, J.A., Umar, I., Fischer, S.H., De Oliveira, J.A., Howell, D., Jardim, E.,

Mosqueira, I. & Ono, K. (2022a) Shaping sustainable harvest boundaries for marine

populations despite estimation bias. *Ecosphere,* **13,** e3923.

Goto, D., Dunlop, E.S., Young, J.D. & Jackson, D.A. (2020) Shifting trophic control of fishery–

ecosystem dynamics following biological invasions. *Ecological Applications,* **30,** e02190.

Goto, D., Filin, A.A., Howell, D., Bogstad, B., Kovalev, Y. & Gjøsæter, H. (2022b) Tradeoffs of

managing cod as a sustainable resource in fluctuating environments. *Ecological Applications,*

32, e2498.

Harris, R.M., Beaumont, L.J., Vance, T.R., Tozer, C.R., Remenyi, T.A., Perkins-Kirkpatrick,

S.E., Mitchell, P.J., Nicotra, A., McGregor, S. & Andrew, N. (2018) Biological responses to

the press and pulse of climate trends and extreme events. *Nature climate change,* **8,** 579-587.

Hastings, A. (2004) Transients: the key to long-term ecological understanding? *Trends in*

Ecology & Evolution, **19,** 39-45.

Hastings, A., Abbott, K.C., Cuddington, K., Francis, T., Gellner, G., Lai, Y.-C., Morozov, A.,

Petrovskii, S., Scranton, K. & Zeeman, M.L. (2018) Transient phenomena in ecology.

Science, **361,** eaat6412.

Hidalgo, M., Rouyer, T., Bartolino, V., Cervino, S., Ciannelli, L., Massuti, E., Jadaud, A.,

Saborido-Rey, F., Durant, J.M. & Santurtún, M. (2012) Context-dependent interplays between

truncated demographies and climate variation shape the population growth rate of a harvested

species. *Ecography,* **35,** 637-649.

Hidalgo, M., Rouyer, T., Molinero, J.C., Massutí, E., Moranta, J., Guijarro, B. & Stenseth, N.C.

(2011) Synergistic effects of fishing-induced demographic changes and climate variation on

fish population dynamics. *Marine Ecology Progress Series,* **426,** 1-12.

Hilborn, R., Amoroso, R.O., Anderson, C.M., Baum, J.K., Branch, T.A., Costello, C., de Moor,

C.L., Faraj, A., Hively, D. & Jensen, O.P. (2020) Effective fisheries management instrumental

in improving fish stock status. *Proceedings of the National Academy of Sciences*.

Hilborn, R., Maguire, J.-J., Parma, A.M. & Rosenberg, A.A. (2001) The precautionary approach

and risk management: can they increase the probability of successes in fishery management?

Canadian Journal of Fisheries and Aquatic Sciences, **58,** 99-107.

Hilborn, R. & Walters, C.J. (1992) *Quantitative fisheries stock assessment: choice, dynamics and*

uncertainty. Springer New York, NY.

Hixon, M.A., Johnson, D.W. & Sogard, S.M. (2014) BOFFFFs: on the importance of conserving

old-growth age structure in fishery populations. *ICES Journal of Marine Science,* **71,** 2171-

2185.

Houde, E.D. (2016) Recruitment variability. *Fish reproductive biology: implications for*

assessment and management (eds T. Jakobsen, M.J. Fogarty, B.A. Megrey & E. Moksness),

pp. 98-187. John Wiley & Sons Ltd, West Sussex, United Kingdom.

Hsieh, C.-h., Reiss, C.S., Hunter, J.R., Beddington, J.R., May, R.M. & Sugihara, G. (2006)

Fishing elevates variability in the abundance of exploited species. *Nature,* **443,** 859-862.

Hsieh, C.-h., Yamauchi, A., Nakazawa, T. & Wang, W.-F. (2010) Fishing effects on age and

spatial structures undermine population stability of fishes. *Aquatic Sciences,* **72,** 165-178.

Hutchings, J.A. (2000) Collapse and recovery of marine fishes. *Nature,* **406,** 882.

Hutchings, J.A. (2015) Thresholds for impaired species recovery. *Proceedings of the Royal*

Society B: Biological Sciences, **282,** 20150654.

Hutchings, J.A. & Kuparinen, A. (2020) Implications of fisheries-induced evolution for

population recovery: Refocusing the science and refining its communication. *Fish and*

Fisheries, **21,** 453-464.

Hutchings, J.A. & Reynolds, J.D. (2004) Marine fish population collapses: consequences for

recovery and extinction risk. *BioScience,* **54,** 297-309.

ICES (2016) Report of the Benchmark Workshop on North Sea Stocks (WKNSEA), 14–18

March 2016, Copenhagen, Denmark. ICES CM 2016/ACOM:37 704 pp.

ICES (2017a) Report of the Arctic Fisheries Working Group (AFWG), 19–25 April 2017,

Copenhagen, Denmark. ICES CM 2017/ACOM:06. 493 pp.

ICES (2017b) Report of the Baltic Fisheries Assessment Working Group (WGBFAS), 19- 26

April 2017, Copenhagen, Denmark. ICES CM 2017/ACOM:11. 810 pp.

ICES (2017c) Report of the North Western Working Group (NWWG), 27 April – 4 May 2017,

Copenhagen, Denmark. ICES CM 2017/ACOM:08. 642 pp.

ICES (2017d) Report of the Working Group for the Bay of Biscay and Iberian waters Ecoregion

(WGBIE), 4-11 May 2017, Cadiz, Spain. *ICES CM 2017/ACOM,* **12,** 552 pp.

ICES (2017e) Report of the Working Group on Assessment of Demersal Stocks in the North Sea

and Skagerrak (2017), 26 April–5 May 2017. ICES HQ. ICES CM 2017/ACOM:21. 1248 pp.

ICES (2017f) Report of the Working Group on Celtic Seas Ecoregion (WGCSE), 9–18 May

2017, Copenhagen, Denmark. ICES CM 2017/ACOM:13. 1464 pp.

ICES (2019). Workshop on Guidelines for Management Strategy Evaluations (WKGMSE2).

ICES Scientific Reports. 1:33. 162 pp.

ICES (2021) ICES fisheries management reference points for category 1 and 2 stocks. ICES

Technical Guidelines.

Jennings, S. & Kaiser, M.J. (1998) The effects of fishing on marine ecosystems. *Advances in*

Marine Biology, pp. 201-352. Elsevier.

Jenouvrier, S., Desprez, M., Fay, R., Barbraud, C., Weimerskirch, H., Delord, K. & Caswell, H.

(2018) Climate change and functional traits affect population dynamics of a long-lived

seabird. *Journal of Animal Ecology,* **87,** 906-920.

Jenouvrier, S., Long, M.C., Coste, C.F., Holland, M., Gamelon, M., Yoccoz, N.G. & Sæther,

B.E. (2022) Detecting climate signals in populations across life histories. *Global Change*

Biology, **28,** 2236-2258.

Jensen, O.P., Branch, T.A. & Hilborn, R. (2012) Marine fisheries as ecological experiments.

Theoretical Ecology, **5,** 3-22.

Juan-Jordá, M.J., Murua, H., Arrizabalaga, H., Merino, G., Pacoureau, N. & Dulvy, N.K. (2022)

Seventy years of tunas, billfishes, and sharks as sentinels of global ocean health. *Science,* **378,**

eabj0211.

Koons, D.N., Iles, D.T., Schaub, M. & Caswell, H. (2016) A life-history perspective on the

demographic drivers of structured population dynamics in changing environments. *Ecology*

Letters, **19,** 1023-1031.

Lassen, H., Kelly, C. & Sissenwine, M. (2014) ICES advisory framework 1977–2012: from F

max to precautionary approach and beyond. *ICES Journal of Marine Science,* **71,** 166-172.

Lindegren, M., Checkley, D.M., Rouyer, T., MacCall, A.D. & Stenseth, N.C. (2013) Climate,

fishing, and fluctuations of sardine and anchovy in the California Current. *Proceedings of the*

National Academy of Sciences, **110,** 13672-13677.

MacKenzie, B.R., Gislason, H., Möllmann, C. & Köster, F.W. (2007) Impact of 21st century

climate change on the Baltic Sea fish community and fisheries. *Global Change Biology,* **13,**

1348-1367.

Marteinsdottir, G. & Thorarinsson, K. (1998) Improving the stock-recruitment relationship in

Icelandic cod (*Gadus morhua*) by including age diversity of spawners. *Canadian Journal of*

Fisheries adn Aquatic Sciences, **55,** 1372-1377.

Melnychuk, M.C., Kurota, H., Mace, P.M., Pons, M., Minto, C., Osio, G.C., Jensen, O.P., de

Moor, C.L., Parma, A.M. & Richard Little, L. (2021) Identifying management actions that

promote sustainable fisheries. *Nature Sustainability,* **4,** 440-449.

Minto, C., Myers, R.A. & Blanchard, W. (2008) Survival variability and population density in

fish populations. *Nature,* **452,** 344.

Moore, J.K., Fu, W., Primeau, F., Britten, G.L., Lindsay, K., Long, M., Doney, S.C., Mahowald,

N., Hoffman, F. & Randerson, J.T. (2018) Sustained climate warming drives declining marine

biological productivity. *Science,* **359,** 1139-1143.

Myers, R.A. (1998) When do environment–recruitment correlations work? *Reviews in Fish*

Biology and Fisheries, **8,** 285-305.

Neubauer, P., Jensen, O.P., Hutchings, J.A. & Baum, J.K. (2013) Resilience and recovery of

overexploited marine populations. *Science,* **340,** 347-349.

Ottersen, G., Hjermann, D.Ø. & Stenseth, N.C. (2006) Changes in spawning stock structure

strengthen the link between climate and recruitment in a heavily fished cod (*Gadus morhua*)

stock. *Fisheries Oceanography,* **15,** 230-243.

Ottersen, G., Planque, B., Belgrano, A., Post, E., Reid, P.C. & Stenseth, N.C. (2001) Ecological

effects of the North Atlantic oscillation. *Oecologia,* **128,** 1-14.

Ottersen, G. & Stenseth, N.C. (2001) Atlantic climate governs oceanographic and ecological

variability in the Barents Sea. *Limnology and Oceanography,* **46,** 1774-1780.

Pauly, D., Watson, R. & Alder, J. (2005) Global trends in world fisheries: impacts on marine

ecosystems and food security. *Philos Trans R Soc Lond B Biol Sci,* **360,** 5-12.

Peck, M.A., Huebert, K.B. & Llopiz, J.K. (2012) Intrinsic and extrinsic factors driving match–

mismatch dynamics during the early life history of marine fishes. *Advances in Ecological*

Research, pp. 177-302. Elsevier.

Pershing, A.J., Alexander, M.A., Hernandez, C.M., Kerr, L.A., Le Bris, A., Mills, K.E., Nye,

J.A., Record, N.R., Scannell, H.A. & Scott, J.D. (2015) Slow adaptation in the face of rapid

warming leads to collapse of the Gulf of Maine cod fishery. *Science,* **350,** 809-812.

Pinsky, M.L. & Byler, D. (2015) Fishing, fast growth and climate variability increase the risk of

collapse. *Proceedings of the Royal Society B,* **282,** 20151053.

Pinsky, M.L., Worm, B., Fogarty, M.J., Sarmiento, J.L. & Levin, S.A. (2013) Marine taxa track

local climate velocities. *Science,* **341,** 1239-1242.

Punt, A.E. (2006) The FAO precautionary approach after almost 10 years: have we progressed

towards implementing simulation-tested feedback-control management systems for fisheries

management? *Natural Resource Modeling,* **19,** 441-464.

Rathore, S., Bindoff, N.L., Phillips, H.E. & Feng, M. (2020) Recent hemispheric asymmetry in

global ocean warming induced by climate change and internal variability. *Nature*

Communications, **11,** 1-8.

Roemmich, D., Church, J., Gilson, J., Monselesan, D., Sutton, P. & Wijffels, S. (2015) Unabated

planetary warming and its ocean structure since 2006. *Nature Climate Change,* **5,** 240-245.

Rogers, L.A. & Dougherty, A.B. (2019) Effects of climate and demography on reproductive

phenology of a harvested marine fish population. *Global Change Biology,* **25,** 708-720.

Rouyer, T., Ottersen, G., Durant, J.M., Hidalgo, M., Hjermann, D.Ø., Persson, J., Stige, L.C. &

Stenseth, N.C. (2011) Shifting dynamic forces in fish stock fluctuations triggered by age

truncation? *Global Change Biology,* **17,** 3046-3057.

Rouyer, T., Sadykov, A., Ohlberger, J. & Stenseth, N.C. (2012) Does increasing mortality

change the response of fish populations to environmental fluctuations? *Ecology Letters,* **15,**

658-665.

Simpson, G.L. & Singmann, H. (2022) gratia: Graceful 'ggplot'-Based Graphics and Other

Functions for GAMs Fitted Using 'mgcv'. *R package*.

Sparholt, H., Bertelsen, M. & Lassen, H. (2007) A meta-analysis of the status of ICES fish

stocks during the past half century. *ICES Journal of Marine Science,* **64,** 707-713.

Stenevik, E.K. & Sundby, S. (2007) Impacts of climate change on commercial fish stocks in

Norwegian waters. *Marine Policy,* **31,** 19-31.

Stenseth, N.C., Ottersen, G., Hurrell, J.W., Mysterud, A., Lima, M., Chan, K.S., Yoccoz, N.G. &

Ådlandsvik, B. (2003) Studying climate effects on ecology through the use of climate indices:

the North Atlantic Oscillation, El Nino Southern Oscillation and beyond. *Proceedings of the*

Royal Society of London. Series B: Biological Sciences, **270,** 2087-2096.

Stubben, C., Milligan, B. & Nantel, P. (2020) popbio: Construction and Analysis of Matrix

Population Models. *R package*.

Sundby, S. & Nakken, O. (2008) Spatial shifts in spawning habitats of Arcto-Norwegian cod

related to multidecadal climate oscillations and climate change. *ICES Journal of Marine*

Science, **65,** 953-962.

van Gemert, R. & Andersen, K.H. (2018) Challenges to fisheries advice and management due to

stock recovery. *ICES Journal of Marine Science,* **75,** 1864-1870.

van Oldenborgh, G.J., te Raa, L.A., Dijkstra, H.A. & Philip, S.Y. (2009) Frequency-or

amplitude-dependent effects of the Atlantic meridional overturning on the tropical Pacific

Ocean. *Ocean Science,* **5,** 293-301.

White, J.W., Botsford, L.W., Hastings, A., Baskett, M.L., Kaplan, D.M. & Barnett, L.A. (2013)

Transient responses of fished populations to marine reserve establishment. *Conservation*

Letters, **6,** 180-191.

Wood, S. (2022) mgcv: Mixed GAM Computation Vehicle with Automatic Smoothness

Estimation. *R package*.

Wood, S. & Scheipl, F. (2020) gamm4: generalized additive mixed models using mgcv and

lme4. *R package*

Wood, S.N. (2017) *Generalized additive models: an introduction with R,* 2nd edn. Chapman and

Hall/CRC.

Worm, B., Hilborn, R., Baum, J.K., Branch, T.A., Collie, J.S., Costello, C., Fogarty, M.J.,

Fulton, E.A., Hutchings, J.A. & Jennings, S. (2009) Rebuilding global fisheries. *Science,* **325,**

578-585.

Zanna, L., Khatiwala, S., Gregory, J.M., Ison, J. & Heimbach, P. (2019) Global reconstruction of

historical ocean heat storage and transport. *Proceedings of the National Academy of Sciences,*

116, 1126-1131.

Zimmermann, F. & Werner, K.M. (2019) Improved management is the main driver behind

recovery of Northeast Atlantic fish stocks. *Frontiers in Ecology and the Environment,* **17,** 93-

99.

Zuur, A.F., Ieno, E.N., Walker, N.J., Saveliev, A.A. & Smith, G.M. (2009) *Mixed effects models*

and extensions in ecology with R. Springer.

Tables

Table 1. Stock-specific information and management measures used in assessments for 38 fish stocks in the northeast Atlantic Ocean included in the study. Stock ID indicates a unique ID for each stock assigned by the International Council of Exploration of the Sea (ICES); period and years indicate the years and the total number of years that stock data are available; age indicates the age classes included in the assessments; F range indicates the age range used to compute mean fishing mortality rates for assessment; F_{msy} and B_{pa} indicate the management reference points set for each stock as target fishing mortality rates that produce maximum sustainable yields and biomasses that satisfy the precautionary approach; and status indicates recovery status assigned based on spawner biomass relative to B_{pa} for analysis (R: recovering and NR: non-recovering) in this study.

common name	scientific name	ecoregion	stock ID	period	years	age	F range	F_{msy}	B_{pa}	status	source¶
cod	*Gadus morhua*	Norwegian–Barents Seas	cod.27.12	1946-2016	71	3-15+	5 -10	0.40	460000	R	a
cod	*Gadus morhua*	North Sea	cod.27.47d20	1963-2016	54	1-6+	2-4	0.31	150000	NR	b
cod	*Gadus morhua*	Baltic Sea	cod.27.22-24	1994-2016	23	1-5+	3-5	0.26	38400	NR	c
cod	*Gadus morhua*	Icelandic Waters	cod-farp	1959-2016	58	1-10+	3-7	0.23	29226	NR	d
cod	*Gadus morhua*	Icelandic Waters	cod.27.5a	1955-2016	62	3-14+	5-10	0.20	160000	R	d
cod	*Gadus morhua*	Celtic Seas	cod.27.6a	1981-2016	36	1-7+	2-5	0.17	22000	NR	e
cod	*Gadus morhua*	Celtic Seas	cod.27.7a	1968-2016	49	0-6+	2-4	0.31	8161	NR	e
cod	*Gadus morhua*	Celtic Seas	cod.27.7e-k	1971-2016	46	1-7+	2-5	0.35	10300	NR	e
four-spot megrim	*Lepidorhombus boscii*	Bay of Biscay and Iberian Coast	ldb.27.8c9a	1986-2017	31	1-7+	2-5	0.19	4600	R	f
haddock	*Melanogrammus aeglefinus*	Norwegian–Barents Seas	had-arct	1950-2016	67	3-13+	4-7	0.35	80000	R	a
haddock	*Melanogrammus aeglefinus*	North Sea	had.27.46a20	1972-2016	45	0-8+	2-4	0.19	132000	R	b
haddock	*Melanogrammus aeglefinus*	Icelandic Waters	had-faro	1957-2016	60	1-10+	3-8	0.13	22843	R	d
haddock	*Melanogrammus aeglefinus*	Icelandic Waters	had.27.5a	1979-2016	38	2-11+	4-7	0.52	59000	R	d
haddock	*Melanogrammus aeglefinus*	Celtic Seas	had-rock	1991-2016	26	1-7+	2-5	0.20	10200	R	e
haddock	*Melanogrammus aeglefinus*	Celtic Seas	had.27.7b-k	1993-2016	24	0-8+	3-5	0.40	10000	R	e
herring	*Clupea harengus*	Baltic Sea	her.27.25-2932	1974-2016	43	1-8+	3-6	0.22	600000	R	c
herring	*Clupea harengus*	Baltic Sea	her.27.28	1977-2016	40	1-8+	3-7	0.32	57100	R	c
megrim	*Lepidorhombus whiffiagonis and L. boscii*	Bay of Biscay and Iberian Coast	meg.27.8c9a	1986-2016	31	1-7+	2-4	0.19	980	R	f
plaice	*Pleuronectes platessa*	North Sea	ple.27.7d	1980-2016	37	1-7+	3-6	0.25	25826	R	b
plaice	*Pleuronectes platessa*	North Sea	ple.27.420	1957-2016	60	1-10+	2-6	0.21	230000	R	b
plaice	*Pleuronectes platessa*	Baltic Sea	ple.27.7e	1999-2016	18	1-5+	3-5	0.37	5550	R	c
plaice	*Pleuronectes platessa*	Celtic Seas	ple.27.7a	1981-2016	36	1-8+	3-6	0.16	5825	R	e
plaice	*Pleuronectes platessa*	Celtic Seas	ple-echw	1980-2016	36	2-10+	3-6	0.24	2443	NR	e
saithe	*Pollachius virens*	North Sea	pok.27.3a46	1967-2016	50	3-9+	4-7	0.36	150000	R	b
saithe	*Pollachius virens*	Icelandic Waters	pok.27.5b	1961-2016	56	3-11+	4-8	0.30	41400	R	d
saithe	*Pollachius virens*	Icelandic Waters	pok.27.5a	1980-2016	37	3-14+	4-9	0.20	61000	R	d
saithe	*Pollachius virens*	Norwegian–Barents Seas	pok.27.1-2	1960-2016	57	3-8+	4-7	0.32†	220000	R	a
sole	*Solea solea*	Baltic Sea	sol.27.20-24	1984-2016	33	2-6+	3-5	0.23	2600	R	c
sole	*Solea solea*	Bay of Biscay and Iberian Coast	sol-bisc	1984-2016	33	2-8+	3-6	0.33	10600	R	f
sole	*Solea solea*	Celtic Seas	sol.27.7a	1970-2016	47	2-8+	4-7	0.20	3100	NR	e
sole	*Solea solea*	Celtic Seas	sol.27.7e	1969-2016	48	2-12+	3-9	0.29	2900	R	e
sole	*Solea solea*	Celtic Seas	sol.27.7fg	1971-2016	46	1-10+	4-8	0.27	2380	R	e
sprat	*Sprattus sprattus*	Baltic Sea	spr.27.22-32	1974-2016	43	1-8+	3-5	0.26	574000	R	c
summer-spawning herring	*Clupea harengus*	Icelandic Waters	her-vasu	1987-2016	30	3-13+	5-10	0.22	273000	R	e
turbot	*Scophthalmus maximus*	North Sea	tur.27.3a	1981-2016	36	1-8+	2-6	0.36*	4163*	R	b
whiting	*Merlangius merlangus*	North Sea	whg.27.47d	1990-2016	27	1-8+	2-7	0.15	166708	R	b

| whiting | *Merlangius merlangus* | Celtic Seas | whg.27.6a | 1981-2016 | 36 | 1-7+ | 2-4 | 0.23 | 44600 | NR | e |
| whiting | *Merlangius merlangus* | Celtic Seas | whg-iris | 1980-2016 | 37 | 0-6+ | 1-3 | 0.22 | 16300 | NR | e |

†F_{msy} was not defined for Barents Sea Saithe in 2017 and an alternative reference point applied (F_{mp}) were used for analysis in the study.

*The reference points for North Sea turbot were not defined until 2018 and the stock was excluded from multilevel modeling analysis.

¶Stock information is taken from ICES Scientific Reports: (a) ICES (2017a), (b) ICES (2017e), (c) ICES (2017b), (d) ICES (2017c) (e) ICES (2017f), (f) ICES (2017d).

Table 2. Parameter estimates and test statistics for the generalized additive mixed effects models selected for the northeast Atlantic fish stocks with per capita recruitment success (RS_{total}) as a response variable (Table S2). EDF, F, CI, t, and SD indicate the effective degrees of freedom, F-statistic, confidence intervals, t-value, and standard deviation.

model structure and parameter	smoother		fixed effect			random effect	
	EDF	F	estimate	95% CI	t	variance	SD
recovering							
$RS_{total} \sim f(Year) + f(Year, SA_{mean}) +$							
$f(Year, SB/B_{pa}) + f(Year, \bar{F}_{lag1})$							
$f(Year)$	1.00	7.75					
$f(Year, SA_{mean})$	4.49	3.12					
$f(Year, SB/B_{pa})$	5.43	19.58					
$f(Year, \bar{F}_{lag1})$	2.00	5.73					
β_0			2.26	(1.81, 2.7)	9.97		
β_{Year}			-0.15	(-0.26, -0.03)	-2.51		
$\beta_{Year \times SAmean1}$			-0.34	(-0.51, -0.17)	-3.99		
$\beta_{Year \times SAmean2}$			-0.04	(-0.14, 0.07)	-0.71		
$\beta_{Year \times SSB/Bpa1}$			0.43	(0.26, 0.61)	4.83		
$\beta_{Year \times SSB/Bpa2}$			-0.05	(-0.17, 0.07)	-0.81		
$\beta_{Year \times \bar{F}lag1,1}$			-0.06	(-0.18, 0.06)	-1.00		
$\beta_{Year \times \bar{F}lag1,2}$			0.16	(0.06, 0.26)	3.07		
stock						0.20	0.45
ecoregion/year						0.13	0.36
ecoregion						0.06	0.25
residual						0.34	0.58
non-recovering							
$RS_{total} \sim f(Year) + f(Year, SA_{mean}) +$							
$f(Year, SSB/B_{pa}) + f(Year, OHC)$							
$f(Year)$	1.00	7.95					
$f(Year, SA_{mean})$	4.47	0.58					
$f(Year, SB/B_{pa})$	3.51	17.8					
$f(Year, OHC)$	2.19	187.8					
β_0			2.40	(1.58, 3.22)	5.75		
β_{Year}			-0.26	(-0.48, -0.03)	-2.22		
$\beta_{Year \times SAmean1}$			-0.18	(-0.47, 0.11)	-1.20		
$\beta_{Year \times SAmean2}$			0.82	(0.6, 1.05)	7.15		
$\beta_{Year \times SSB/Bpa1}$			-0.40	(-0.63, -0.16)	-3.28		
$\beta_{Year \times SSB/Bpa2}$			0.05	(-0.15, 0.25)	0.46		
stock						0.42	0.64
ecoregion/year						0.17	0.41
ecoregion						1.81×10^{-5}	4.26×10^{-3}
residual						0.27	0.52

Table 3. Parameter estimates and test statistics for the generalized additive mixed effects models selected for the northeast Atlantic fish stocks with elasticity to recruitment success (e_{RS}) as a response variable (Table S2). EDF, F, 95% CI, t, and SD indicate the effective degrees of freedom, F-statistic, confidence intervals, t-value, and standard deviation.

model structure and parameter	smoother		fixed effect			random effect	
	EDF	F	estimate	95% CI	t	variance	SD
recovering							
$e_{RS} \sim f(\text{Year}) + f(\text{Year, SA}_{mean}) + f(\text{Year, SB}/B_{pa})$							
$+ f(\text{Year}, \bar{F}_{lag1}/F_{msy}) + f(\text{Year, AMO}_{lag1})$							
$f(\text{Year})$	1.00	0.60					
$f(\text{Year, SA}_{mean})$	3.51	1.61					
$f(\text{Year, SB}/B_{pa})$	7.61	39.16					
$f(\text{Year}, \bar{F}_{lag1}/F_{msy})$	5.06	3.71					
$f(\text{Year, AMO}_{lag1})$	7.15	3.09					
β_0			-5.16	(-5.25, -5.07)	-118.0		
stock						0.02	0.16
ecoregion/year						0.01	0.11
ecoregion						0.01	0.08
non-recovering							
$e_{RS} \sim f(\text{Year}) + f(\text{Year, SA}_{mean}) + f(\text{Year, SB}/B_{pa})$							
$+ f(\text{Year}, \bar{F}_{lag1}/F_{msy}) + f(\text{Year, wSST}_{anom})$							
$f(\text{Year})$	1.00	2.10					
$f(\text{Year, SA}_{mean})$	7.01	141.36					
$f(\text{Year, SB}/B_{pa})$	6.65	4.88					
$f(\text{Year}, \bar{F}_{lag1}/F_{msy})$	3.60	1.96					
$f(\text{Year, wSST}_{anom})$	3.02	1.52					
β_0			-5.16	(-5.36, -4.96)	-50.2		
stock						0.09	0.31
ecoregion/year						0.008	0.088
ecoregion						3.4×10^{-7}	5.8×10^{-4}

Figure legends

Figure 1. Study systems and climate conditions in the northeast Atlantic Ocean. (a) Geographic locations of the ecoregions, which are defined by the International Council of Exploration of the Sea (ICES): Norwegian and Barents Seas (62°0'N–82°0'N at 11°0'W–68°30'E), Greater North Sea (56°0'N at 2°55'E), Baltic Sea (53°35'N–65°54'N at 11°57'E–30°14'E), Faroes (60°0'N–63°0'N at 15°0'W–4°0'W), Icelandic waters (64°38'N at 18°55'W), the Bay of Biscay and the Iberian Coast (42°39'N at 8°53'W), and Celtic Seas (48°0'N–60°30'N at 18°0'W–1°21'W). (b) Regional climate indices (North Atlantic Oscillation (NAO) index, Atlantic Meridional Oscillation index, and upper (0–400m) ocean heat content (OHC) anomaly) during the study period (1946–2016). (c) Seasonal sea surface temperature anomalies in each ecoregion. The time series of temperature anomalies in each ecoregion are truncated to show only the years that the fish stock data and assessments included in the study are available.

Figure 2. Regional patterns in fishing pressure and reproductive metrics of the recovering and non-recovering northeast Atlantic fish stocks during the study period (1946–2016). (a) Temporal trends in relative fishing pressure, relative spawner biomass, mean spawner age, and spawner age diversity. Vertical dashed lines indicate the reference year (2002) used to define the recovery status of fish stocks based on relative spawner biomass in analysis. (b) Relationships between relative fishing pressure and relative spawner biomass and age structure metrics. Solid lines and ribbons indicate regional average trends in time (a) or relative fishing pressure (b) and 95% confidence intervals estimated by generalized additive mixed effects modeling with year (a) or relative fishing pressure (b) as a fixed effect, and ecoregion, year within ecoregion, and stock as random effects. Circles indicate partial residuals estimated from the models. Relative fishing pressure and relative spawner biomass are defined as estimated fishing mortality rate and

spawner biomass relative to management reference points (F_{msy} and B_{pa}) defined by the International Council of Exploration of the Sea (ICES).

Figure 3. Regional trends in transient population growths of the recovering and non-recovering northeast Atlantic fish stocks during the study period (1946–2016). Solid lines and ribbons indicate temporal trends and 95% confidence intervals estimated by generalized additive mixed effects modeling with year as a fixed effect, and ecoregion, year within ecoregion, and stock as random effects. Circles indicate partial residuals estimated from the models.

Figure 4. Regional trends in and covariates of per capita recruitment success (fecundity) of the recovering and non-recovering northeast Atlantic fish stocks during the study period (1946–2016). (a) Temporal trends. Solid lines and ribbons indicate temporal trends and 95% confidence intervals estimated by generalized additive mixed effects modeling with year as a fixed effect, and ecoregion, year within ecoregion, and stock as random effects. Red circles indicate partial residuals estimated by the models. (b–e) Time-varying effects of spawner age structure, spawner biomass, fishing pressure. and climate condition. Covariates selected for each type in the models are indicated in the y-axis labels. Color gradients and contour lines indicate the partial effects of covariates estimated (as tensor interaction terms) in the models (only the covariate types that are supported by data are shown). Warmer colors indicate higher values, and the ranges of the partial effects differ among the covariates and are indicated as values on the contour lines. Gray circles indicate back-transformed input data.

Figure 5. Ecoregion-scale trends and covariates of per capita recruitment success (fecundity) of the northeast Atlantic fish stocks during the study period (1946–2016). (a) Temporal trends. Solid lines and ribbons indicate temporal trends and 95% confidence intervals estimated by generalized additive mixed effects modeling with year as a fixed effect, and ecoregion, year

within ecoregion, and stock as random effects. Circles indicate partial residuals estimated by the models. (b–e) Time-varying effects of spawner age structure, spawner biomass, fishing pressure. and climate condition. Covariates selected in the models are indicated in the y-axis labels. Color gradients and contour lines indicate the partial effects of covariates estimated (as tensor interaction terms) in the models. Warmer colors indicate higher values, and the ranges of the partial effects differ among the covariates and are indicated as values on the contour lines. Gray circles indicate back-transformed input data.

Figure 6. Regional trends in and covariates of elasticity to per capita recruitment success of the recovering and non-recovering northeast Atlantic fish stocks during the study period (1946–2016). (a) Temporal trends. Solid lines and ribbons indicate temporal trends and 95% confidence intervals estimated by generalized additive mixed effects modeling with year as a fixed effect, and ecoregion, year within ecoregion, and stock as random effects. Circles indicate partial residuals estimated by the models. (b–e) Time-varying effects of spawner age structure, spawner biomass, fishing pressure, and climate condition. Covariates selected in the models are indicated in the y-axis labels. Color gradients and contour lines indicate the partial effects of covariates estimated (as tensor interaction terms) in the models. Warmer colors indicate higher values, and the ranges of the partial effects differ among the covariates and are indicated as values on the contour lines. Gray circles indicate back-transformed input data.

Figure 7. Ecoregion-scale trends and covariates of elasticity to per capita recruitment success of the northeast Atlantic fish stocks during the study period (1946–2016). (a) Temporal trends. Solid lines and ribbons indicate temporal trends and 95% confidence intervals estimated by generalized additive mixed effects modeling with year as a fixed effect, and ecoregion, year within ecoregion, and stock as random effects. Circles indicate partial residuals estimated by the

models. (b–e) Time-varying effects of spawner age structure, spawner biomass, fishing pressure. and climate condition. Covariates selected in the models are indicated in the y-axis labels. Color gradients and contour lines indicate the partial effects of covariates estimated (as tensor interaction terms) in the models. Warmer colors indicate higher values, and the ranges of the partial effects differ among the covariates and are indicated as values on the contour lines. Gray circles indicate back-transformed input data.