Transient demographic dynamics of recovering fish populations shaped by past climate

variability, harvest, and management

**Running Title**: Transient dynamics of recovering populations

Daisuke Goto[1,2,¶]

[1]Institute of Marine Research, P.O. Box 1870, Nordnes, 5817 Bergen, Norway

[2]Swedish University of Agricultural Sciences, Department of Aquatic Resources, Drottningholm,

Sweden.

[¶]Corresponding author: daisuke.goto2@gmail.com

**Abstract**

Large-scale commercial harvesting and climate-induced fluctuations in ocean properties shape

the dynamics of marine populations as interdependent drivers at varied timescales. Selective

removals of larger, older members of a population can distort its demographic structure, eroding

resilience to fluctuations in habitat conditions and thus amplifying volatility in transient

dynamics. Through the implementation of stricter management measures, many historically

depleted fish stocks began showing signs of recovery in recent decades. But these interventions

coincided with accelerated changes in the oceans triggered by increasingly warmer, more

variable climates. Applying multilevel models to annual estimates of demographic metrics of 38

stocks comprising 11 species across seven ecoregions in the northeast Atlantic Ocean, this study

explores how time-varying local and regional climates contributed to the transient dynamics of

recovering populations exposed to variable fishing pressures moderated by management actions.

Analyses reveal that progressive reductions in fishing pressure and shifting climate conditions

nonlinearly shaped rebuilding patterns of the stocks through restorations of maternal

demographic structure (reversing age truncation) and reproductive capacity. As the survival rate

and demographic structure of reproductive fish improved, transient growth became less sensitive

to variability in recruitment and juvenile survival and more to that in adult survival. As the

biomass of reproductive fish rose, recruitment success also became increasingly regulated by

density-dependent processes involving higher numbers of older fish. When reductions in fishing

pressure were insufficient or delayed, however, stocks became further depleted, with more

eroded demographic structures. Although warmer local climates in spawning seasons promoted

recruitment success in some ecoregions, changing climates in recent decades began adversely

affecting reproductive performances overall, amplifying sensitivities to recruitment variability.

These shared patterns underscore the value of demographic transients in developing robust

strategies for managing marine resources. Such strategies could form the foundation for effective

applications of adaptive measures resilient to future environmental change.

*Keywords*: age truncation, stock assessment, meta-analysis, demographic buffering, climate

change, ecosystem-based management, matrix population model, life history trait, life table

response experiment, hierarchical modeling

**Introduction**

Excessive human exploitation that persisted through the twentieth century depleted marine living resources worldwide, threatening biodiversity, food and nutrient security, and other ecosystem services that the oceans provide (Pauly, Watson & Alder 2005; Crowder *et al.* 2008; Worm *et al.* 2009). Long-lived, late-maturing vertebrate predators were disproportionately depleted by industrial fishing as target or incidental catch (Christensen *et al.* 2003; Juan-Jordá *et al.* 2022). As part of the efforts to halt and reverse the losses, national government and intergovernmental organizations began applying the precautionary principle to the management of marine fisheries (Hilborn *et al.* 2001; Punt 2006), following international agreements, including the Code of Conduct for Responsible Fisheries adopted by the Food and Agriculture Organization of the United Nations (FAO 1995; Melnychuk *et al.* 2021). Since the implementation of stricter management measures to regulate fishing operations in the early 1990s, an increasing number of historically overexploited fish stocks began showing signs of recovery (Hilborn *et al.* 2020; Juan-Jordá *et al.* 2022). These management interventions, however, coincided with accelerated changes in the oceans (Barange *et al.* 2018). Global climate is becoming increasingly warmer and more variable, modifying regional and local habitat conditions that support the biomass production of fish stocks, possibly shifting biologically sustainable levels of harvesting (Chassot *et al.* 2010; Free *et al.* 2019).

The dynamics of marine resource populations in many systems are shaped by drivers operating interactively at varied timescales, including high-frequency perturbations like fishing and low-frequency variability in ocean properties forced by climate (Rouyer *et al.* 2012; Goto *et al.* 2022b). When exposed to such persistent, nonstationary perturbations, near-term (transient) dynamics of an exploited population, which depend on both variable vital rates (survival, growth,

and fecundity) and demographic (age and stage) structures, may deviate from asymptotic

(equilibrium) dynamics (Hastings 2004; Koons *et al.* 2016). Past research instructs us that large-

scale shifts in ecosystem productivity under alternative climate regimes can be propagated

through the system to influence fluctuations in fisheries catches (Chassot *et al.* 2010; Cheung,

Watson & Pauly 2013). Climate effects on marine species are often integrated over a series of

oceanic, ecological, and biological processes over space and through time (Ottersen *et al.* 2001;

Stenseth *et al.* 2003). Fluctuations in population structure and size can, for example, emerge with

some time lag from age (or stage)-dependent demographic processes like birth and death

(Jenouvrier *et al.* 2022) modulated by seasonal and interannual variations in weather and habitat

conditions (Hutchings & Reynolds 2004; Lindegren *et al.* 2013). These transient fluctuations in

the population could sustain over generations through delayed feedbacks (Hastings 2004).

In contrast, fishing operations and management actions that regulate the operations can directly

modify short-term trends and variability in population structure and demographic rates, and thus

the transient dynamics of exploited species (Gamelon, Sandercock & Sæther 2019). Large-scale

commercial fishing often selectively removes larger, older members of the population, truncating

age (and size) structure, modifying life history traits, reducing the capacity to absorb the effects

of perturbations, and thus amplifying year-to-year variation in population size in variable

environments (Hsieh *et al.* 2006; Rouyer *et al.* 2012). Fishing-induced age truncations

(juvenescence) are widely reported for harvested marine populations (Hsieh *et al.* 2010; Rouyer

*et al.* 2012), many of which suffer from reduced reproductive capacity (Hixon, Johnson &

Sogard 2014). Climate-induced changes in the variability of ocean habitat conditions thus may

indirectly promote or impede the recovery of depleted populations with weakened demographic

resilience to perturbations (Lindegren *et al.* 2013; Harris *et al.* 2018).

In commercial fisheries a harvested population is commonly managed through regulation of fishing pressure based on its population status (Beddington, Agnew & Clark 2007). Because of high socioeconomic values, exploitation of marine resource populations is regularly monitored through scientific surveys and assessed based on population size estimates (Hilborn & Walters 1992). These population assessments form the scientific basis for the implementation of management measures like catch limits. When population size declines below a predefined threshold (a reference point set under the precautionary principle, ICES 2021; Goto *et al.* 2022a), for example, fishing pressure may be adjusted to mitigate the risk of overharvesting. This adaptive management cycle (human–natural system feedbacks through monitoring, assessment, and management action) thus can be considered as a natural experiment (Hilborn & Walters 1992; Jensen, Branch & Hilborn 2012), providing opportunities to gain insights into the transient dynamics of managed populations in variable environments. Understanding what drives these transients would help develop robust strategies to sustainably manage human exploitation of marine resource populations as we face increasingly uncertain conditions in the oceans.

This article reports a study that empirically explores how past fishing, management action, and climate variability influenced the transient dynamics of recovering fish populations in the northeast Atlantic Ocean (FAO Major Fishing Area 27, Fig. 1a). These populations experienced wide ranges of fishing pressure, climate, and management action in recent decades (Rouyer *et al.* 2011; Zimmermann & Werner 2019). Annual fishing pressures applied to many northeast Atlantic stocks began to be eased in the late 1990s to rebuild biomass that supports sustainable harvesting (Sparholt, Bertelsen & Lassen 2007; Lassen, Kelly & Sissenwine 2014). Following the reductions in fishing pressure, these intensively managed stocks experienced considerable changes in population size, demographic structure, and productivity (Hidalgo *et al.* 2012; Rouyer

*et al.* 2012; Britten, Dowd & Worm 2016). During this period the regional climate also became

increasingly warmer and more variable as greenhouse gas emissions rose, which was reflected

especially in upper ocean heat content and transport (Delworth *et al.* 2016; Zanna *et al.* 2019).

Warming patterns, however, highly varied among geographic subregions (ecoregions) owing to

local air–sea feedbacks (Roemmich *et al.* 2015; Delworth *et al.* 2016), triggering a series of

oceanic and ecological responses at variable rates (Beaugrand 2009; Delworth *et al.* 2016). To

glean insights into the recovery patterns of historically depleted fish stocks, this study

specifically asks how local and regional climate variability contributes to variations in the

reproductive performance and transient growth of marine species exposed to variable fishing

pressures moderated by management actions.

**Materials and Methods**

To investigate the transient dynamics of northeast Atlantic fish stocks at local and regional

scales, a series of multilevel statistical models were applied to the time series of population and

demographic metrics along with intrinsic and extrinsic covariates. In the following I describe (a)

development of matrix population models to quantify transient population growth and elasticities

to demographic rates through transient life table response experiments, (b) trend analyses to

characterize the dynamics of reproductive fish biomass and age structure, transient growth, and

fishing pressure, followed by analyses to test for direct fishing effects, and (c) evaluation of the

relative contributions of time-varying intrinsic (reproductive fish biomass and age structure) and

extrinsic (fishing pressure and climate) drivers to variations in recruitment success and in

elasticity to recruitment success.

*Data*

Time series (1946–2016) of annual estimates of age-specific information (mean body mass,

abundance, maturity rate, and natural and fishing mortality rates) and management reference

points (which define the management targets for fishing pressure and spawner biomass, as

described below in *Management action for fisheries*) of marine fishes in the northeast Atlantic

Ocean were extracted from the 2017 stock assessment reports by the International Council of

Exploration of the Sea (ICES) (Table 1). In the assessment of each stock (defined here as a

population in a management unit area defined by the ICES) the demographic rates and life

history traits were estimated using data from commercial catch reporting and scientific

monitoring surveys. Stock sizes and fishing mortality rates then were estimated annually using

age-structured population models fitted to time series data of abundance indices (ICES 2017e;

ICES 2017a; ICES 2017b; ICES 2017c; ICES 2017f; ICES 2017d). The reliability of the models

and data had been evaluated by experts prior to assessments and extensively documented (e.g.,

ICES 2016). For analysis I excluded stocks with missing age-specific information, rejected

assessments, or shorter time series (less than 15 years). The final dataset consists of 38 stocks

comprising 11 species (cod *Gadus morhua*, haddock *Melanogrammus aeglefinus*, saithe

*Pollachius virens*, herring *Clupea harengus*, plaice *Pleuronectes platessa*, turbot *Scophthalmus*

*maximus*, sole *Solea solea*, whiting *Merlangius merlangus*, sprat *Sprattus sprattus*, megrim

*Lepidorhombus spp*., and four-spot megrim *Lepidorhombus boscii*) across seven ecoregions in

the northeast Atlantic Ocean defined by the ICES: the Norwegian and Barents Seas (combined as

a single ecoregion for analysis), the Greater North Sea (hereafter the North Sea), the Baltic Sea,

Faroes, Icelandic waters, the Bay of Biscay and the Iberian Coast (hereafter the Bay of Biscay),

and the Celtic Seas (Table 1 and Fig. 1a). These stocks support some of the largest fisheries in

the region, collectively yielding 1.9 to 2.6 million tonnes of commercial landings annually in the

recent decades (1997–2016).

Climate indices and sea surface temperature (SST) anomalies were used as proxies that

represent variations in overall regional (North Atlantic) and local (ecoregion) climate conditions

(1946–2016), which are filtered through oceanic processes fish populations directly and

indirectly experience over interannual and decadal timescales (Ottersen & Stenseth 2001;

Stenseth *et al.* 2003). SST was chosen as a proxy for local habitat conditions because its

reconstructed data can provide adequate spatial and temporal coverage for this study. Further, the

climate data were used as covariates in multilevel models with early life-history processes as

response variables (as described below in *Covariates of recruitment success* and *Covariates of*

*elasticity to recruitment success*). Climate effects on these processes can, for example, be

mediated by lower trophic level (like zooplankton) production responding to regional climate-

driven SST anomalies (Beaugrand & Kirby 2018) and other ocean properties (like heat

transport), which may in turn manifest as delayed effects in young fish survival (Hjermann *et al.*

2007).

Monthly time series of three regional climate indices were downloaded and averaged over

winter months (when the atmosphere is most active in the region, Stenseth *et al.* 2003) for

analysis (Fig. 1b): the North Atlantic Oscillation (NAO) and Atlantic Multidecadal Oscillation

(AMO) indices (both computed from the Hadley Centre Sea Ice and Sea Surface Temperature

data set [HadISST1, https://www.metoffice.gov.uk] and the extended reconstructed SST

[ERSSTv5, NOAA Physical Sciences Laboratory; https://psl.noaa.gov] and averaged over the

sources), and upper (0–700 m) ocean heat content (OHC) anomaly in the North Atlantic (NOAA

National Centers for Environmental Information; https://www.ncei.noaa.gov). The NAO index

reflects the dominant mode of atmospheric variability between the Azores and Iceland, linked

with a range of large-scale oceanic features (such as surface temperature and heat content) in the

North Atlantic that regulate biological and ecosystem processes (Stenseth *et al.* 2003). The AMO

index, computed by averaging SSTs over the North Atlantic (25°–60°N at 7°–70°W) using the

method by van Oldenborgh *et al.* (2009), also reflects large-scale climate variability that may

influence the movement and migration of marine species (Faillettaz *et al.* 2019). OHC measures

the amount of excess heat (from increasing greenhouse gases) trapped in the ocean, integrating

basin-wide heat storage and transport (Zanna *et al.* 2019). Because of high signal-to-noise ratio,

OHC reflects more consistent decadal trends (less influenced by natural variability) than other

climate indices (Roemmich *et al.* 2015).

Reconstructed gridded monthly SST data were downloaded from and averaged over three

sources to account for systematic uncertainties in each source: HadISST1, COBE Sea Surface

Temperature SST (NOAA Physical Sciences Laboratory), and ERSSTv5. I first averaged the

SST data over ecoregion and for each ecoregion over season; winter (December in the previous

year–February, $wSST_{anom}$), spring (March–May, $spSST_{anom}$), summer (June–July, $smSST_{anom}$),

and autumn (September–November, $aSST_{anom}$). I then computed SST anomalies by subtracting

the mean across years (1946–2016) for each season and ecoregion (Fig. 1c).

*Transient population growth and elasticity*

I used transient population growth rate as an integrated measure that captures time-varying

demographic rates and age structures responding to non-stationary extrinsic factors (fishing

pressure, management action, and climate) and estimated the growth rates for each stock using

an age-structured matrix projection model (Caswell 2001). Leslie (state-transition) matrices (*A*)

were first constructed with annual estimates of age-specific survival and reproductive rates.

Survival rates ($S_{a,y}$) were estimated as $S_{a,y} = \exp(-Z_{a,y})$, where $Z_{a,y} = F_{a,y} + M_{a,y}$, and $F_{a,y}$ and $M_{a,y}$

are fishing and natural (non-fishing like predation and starvation) mortality rates for $a$-year olds

in year $y$. Egg production rates are often not estimated in the assessments of marine fish stocks,

and here I estimated recruitment success (integrating over reproduction and early life-history

stage survival) rate per spawner (reproductive fish) for $a$-year olds in year $y$ ($RS_{a,y}$) (Rouyer *et al.*

2011) as a proxy for per capita fecundity rate as

$$RS_{a,y} = \frac{R_y m_{a,y-a_R}}{\sum_{a=a_{min}}^{a=a_{max}} m_{a,y-a_R} N_{a,y-a_R}} \tag{1}$$

where $R_y$ is the recruit number in year $y$, $m_{a,y-a_R}$ and $N_{a,y-a_R}$ are the maturity rate (fraction of

juveniles becoming adults) and abundance of $a$-year olds in year $y-a_R$ (spawning year), $a_R$ is the

age of recruits, which accounts for variations in age at recruitment (0–3 years old) among stocks

reported in the assessments. $a_{min}$ and $a_{max}$ are the youngest and oldest ages in the stock. I then

numerically estimated deterministic transient growth rates ($\lambda_{trans}$) by projecting age-specific

abundances using the annual state-transition matrices and the annual estimates of age-specific

numbers taken from the assessments as

$$\lambda_{trans,y} = \frac{n_{y+T_{gen,y}}}{n_y}$$

$$n_{y+Tgen} = n_y A_y \tag{2}$$

213  where $n_y$ and $A_y$ are a vector of initial age-specific numbers and an $a \times a$ state-transition matrix

214  in year $y$, and $T_{gen}$ is the generation time (years) in year $y$. To account for variations in life-

215  history strategy among stocks, the transient growth rates of each stock were estimated over its

216  generation time, which is defined as the mean age of parents of a cohort (Caswell 2001),

$$T_{gen,y} = \frac{\sum_a a (\prod_{j=1}^{a-1} S_{j,y}) R_{a,y}}{\sum_a (\prod_{j=1}^{a-1} S_{j,y}) R_{a,y}} \tag{3}$$

The transient growth rates here were thus defined in timescales of 2.6 to 6.7 years (Fig. S1),

relevant for fisheries management.

To evaluate the contributions of demographic variations to transient growth I performed

transient life table response experiments (LTRE) by numerically estimating transient elasticities

($e_y$) to survival and fecundity rates through matrix projections (Koons *et al.* 2016); demographic

parameters (fecundity or survival) of all ages were first perturbed by a small amount (1%), and

stock numbers were projected to re-compute $\lambda_{\text{trans},y}$ (Durant *et al.* 2013). Elasticity (relative

sensitivity), defined as the proportional change in $\lambda_{\text{trans}}$ relative to a proportional change in a

demographic rate (Caswell 2001), was computed as $e_y = \frac{\Delta \log \lambda_{trans,y}}{\Delta \log \alpha_y}$, where $\alpha_y$ is the survival or

fecundity rates in year *y* that are perturbed (Coulson *et al.* 2004). Age-specific elasticities were

then summed for analysis; changes in elasticities to juvenile and adult survival rates were

evaluated separately by summing age-specific elasticities weighted by maturity rates. To

quantify how the transient dynamics deviate from expected long-term equilibriums (asymptotic

dynamics assuming stable environments and age structures) across years I also computed year-

specific asymptotic (geometric) growth rates and stable age distributions from the logarithms of

the dominant (largest) eigenvalues and the right eigenvectors (respectively) of the annual state-

transition matrices (Caswell 2001).

To evaluate how variations in demographic structure have contributed to the reproductive

performance and recovery patterns of fish stocks, I used mean age and age diversity of spawners

as indices for demographic structure. Mean age (SA$_{\text{mean}}$) is the weighted (by abundance) average

of spawner age as

239
$$\text{AS}_{\text{mean,y}} = \frac{\sum_{a=a_{min}}^{a=a_{max}} a m_{a,y} N_{a,y}}{\sum_{a=a_{min}}^{a=a_{max}} m_{a,y} N_{a,y}}$$
(4)

240 The Shannon diversity index was used as an indicator of age diversity ($SA_{diversity}$) and computed

241 as $SA_{diversity,y} = -\sum_{a=a_{min}}^{n} p_{a,y} \ln(p_{a,y})$, where $p_{a,y}$ is the proportion of $a$-year-old spawner number

242 in a given age group in year $y$ (Marteinsdottir & Thorarinsson 1998).

*Management action for fisheries*

Management action (here adjustment of fishing pressure) is based on perceived current stock

status and harvest control rule set by the ICES (which are designed to achieve sustainable

harvesting without compromising the reproductive capacity of the stock). The current status is

assessed using scientific monitoring and catch data before the provision of annual catch advice

(total allowable catch). Under the control rule, when the estimated SB remains above a fixed

threshold ($B_{pa}$) at the start of the year following the assessment, the catch limit is computed based

on target exploitation rate ($F_{msy}$). These two parameters ($B_{pa}$ and $F_{msy}$) are designed to prevent

overharvesting by accounting for uncertainty in population and harvest dynamics (ICES 2019a).

When the SB falls below $B_{pa}$, exploitation rate is adjusted to $F_{msy}$ scaled to the proportion of SB

relative to $B_{pa}$, thereby allowing the population to rebuild (adaptive management). For analysis

here fishing pressure and SB relative to stock-specific reference points ($\bar{F}/F_{msy}$ and $SB/B_{pa}$) were

used as proxies to evaluate the influence of the management measures on the demographic

metrics of fish stocks. Fishing pressure is defined as the instantaneous rate of fishing mortality

($\bar{F}$), which is commonly estimated by averaging age-specific fishing mortality rates over

dominant ages in fishery catch (Table 1).

*Data analysis*

To explore how management-guided changes in fishing pressure and climate change

influenced the vital rates, transient dynamics, and recovery patterns of the northeast Atlantic fish

stocks, I applied multilevel (hierarchical) modeling to derived demographic metrics to account

for nonindependence in data structure (Gelman & Hill 2006) with stock, ecoregion, and year as

grouping factors (random intercepts). Data exploration including identification of collinearity

(through visual inspection correlation plots and variance inflation factors) and outliers (>three

standard deviations) was performed before analysis (Zuur *et al.* 2009). The data exploration

revealed two groups of stocks with diverging (increasing and declining) patterns in stock status

based on relative spawner biomass (SB/$B_{pa}$). To explore correlations underlying the differences I

applied generalized additive mixed effects models (GAMMs, Wood 2017) to these groups

separately. In the following analyses the stocks were grouped based on stock status in the last 15

years of the time series (2002–2016); if spawner biomass is larger than 80% of a biological

threshold ($B_{pa}$, Costello *et al.* 2016) for at least 50% of the period (eight years) a stock is

classified as 'recovering', otherwise 'non-recovering' (sensitivity analyses showed that the

diverging patterns in SB/$B_{pa}$ are robust to variations in the criteria used, Fig. S2 and S3). This

classification resulted in 27 recovering and 10 non-recovering stocks (Table 1).

*Trends in populations, demographics, and fishing*. Trend analyses were first performed with

year as a covariate to capture shared long-term historical patterns in relative spawner biomass,

spawner demographic structure (mean age and age diversity), relative fishing pressure ($\bar{F}/F_{msy}$),

and transient growth. The response variables were assumed to follow a Gamma distribution with

continuous, positive expected values $\mu$ and a logarithmic link function as

$$Y_{jk} \sim \text{Gamma}(\mu_{ijk}, \ \tau)$$

$$E(Y_{ijk}) = \mu_{ijk}$$

$$\text{var}(Y_{ijk}) = \mu_{ijk}^2/\tau$$

$$\log(\mu_{ijk}) = \beta_0 + f_1(\text{Year}_{ij}) + \text{stock}_i + \text{ecoregion}_j + \text{year}_{jk}$$

$$\text{stock}_i \sim N(0, \sigma_{stock}^2), \ \text{ecoregion}_j \sim N(0, \sigma_{ecoregion}^2), \ \text{year}_{jk} \sim N(0, \sigma_{year}^2) \qquad (5)$$

where $Y_{jik}$ is the response of stock $i$ in ecoregion $j$ in year $k$, $\tau$ is a Gamma distributed parameter,

$\beta_0$ is the intercept, and $f_1$ is a smoother function (thin-plate cubic splines with five maximum

degrees of freedom or knots) for year; $stock_i$, $ecoregion_j$, and $year_{jk}$ are normally distributed

random intercepts with mean zero and standard deviation $\sigma$ for stock, ecoregion, and year nested

within ecoregion. The structure of random effects was selected based on likelihood ratio tests

applied to ten structures fitted with a restricted maximum likelihood method (REML).

Abundance and demographic parameters in stock assessments are estimated with several known

sources of uncertainty (including observation, process, and estimation errors, ICES 2019b),

which may bias post-hoc analyses when used as data (Brooks & Deroba 2015). To address this

issue, I performed sensitivity analyses by Monte Carlo simulations to illustrate how uncertainties

in the estimates can propagate through matrix population and multilevel modeling analyses.

Because uncertainty in age-specific parameters was not consistently reported in the assessments,

I first generated 1000 simulated datasets with Gaussian noise in recruit numbers and fishing

mortality rates using the stock- and year-specific coefficients of variation derived from the

assessments and propagated the estimation uncertainties through the computation of state-

transition and transient population growth rates (see Fig. S4 for more detail). This step was then

followed by multilevel model-fitting with transient population growth as a response variable and

year as a covariate (as described above) applied to each simulated dataset.

Next, I assessed direct effects of fishing on spawner biomass and age structure by evaluating

relationships between relative fishing pressure and relative biomass or age structure of spawners

as

307                             $$Y_{jk} \sim \mathrm{Gamma}(\mu_{ijk},\ \tau)$$

308                             $$E(Y_{ijk}) = \mu_{ijk}$$

309
$$\text{var}(Y_{ijk}) = \mu_{ijk}{}^2/\tau$$

310
$$\log(\mu_{ijk}) = \beta_0 + f_1(\text{Fishing}_{ij}) + \text{stock}_i + \text{ecoregion}_j + \text{year}_{jk}$$

311
$$\text{stock}_i \sim N(0, \sigma^2_{stock}), \text{ecoregion}_j \sim N(0, \sigma^2_{ecoregion}), \text{year}_{jk} \sim N(0, \sigma^2_{year}) \qquad (6)$$

where $Y_{jik}, \tau$, $\beta_0$, $\sigma_{stock}$, $\sigma_{ecoregion}$, and $\sigma_{year}$ are the same as above. $f_1$ is a smoother function for

relative fishing pressure.

The contributions of fishing, management action, and climate to the transient dynamics of fish

stocks with variable age structures were evaluated in two steps. I first tested for time-varying

contributions of these extrinsic factors to variations in vital rates and then evaluated how these

factors influenced the elasticities of transient population growth:

*Covariates of recruitment success*. Because natural mortality rates were assumed time-

invariant in most of the assessments, variations in survival rates are in effect driven by fishing.

Here, RS was thus selected as a demographic response variable (which integrates probabilities of

spawning and early life history stage survival) for analysis. Because RS can vary with variations

in spawner age structure and biomass (intrinsic factors), these intrinsic factors were also tested.

To test for the time-varying contributions of the extrinsic and intrinsic factors, the following

model was fitted to the data:

325
$$\text{RS}_{\text{total}\,ijk} \sim \text{Gamma}(\mu_{ijk}, \ \tau)$$

326
$$E(\text{RS}_{\text{total},ijk}) = \mu_{ijk}$$

327
$$\text{var}(\text{RS}_{\text{total},ijk}) = \mu_{ijk}{}^2/\tau$$

328
$$\log(\mu_{ijk}) = \beta_0 + f_1(\text{Year}_{ij}) + f_2(\text{Year}_{ij}, \text{Biomass}_{ijk}) + f_3(\text{Year}_{ij}, \text{Demography}_{ijk}) +$$

329
$$f_4(\text{Year}_{ij}, \text{Fishing}_{(i\text{-}1)jk}) + f_5(\text{Year}_{ij}, \text{Climate}_{ij}) + \text{stock}_i + \text{ecoregion}_j + \text{year}_{jk}$$

330
$$\text{stock}_i \sim N(0, \sigma^2_{stock}), \text{ecoregion}_j \sim N(0, \sigma^2_{ecoregion}), \text{year}_{jk} \sim N(0, \sigma^2_{year}) \qquad (7)$$

where $RS_{total,ijk}$ is the total per capita recruitment success rate of stock $i$ in ecoregion $j$ in year $k$,

$\beta_0$ is the intercept, $f_1$–$f_5$ are smoother functions for covariates: year, year × density, year ×

demographics (spawner mean age and age diversity), year × climate, and year × fishing. *Climate*,

*Biomass*, and *Demography* were lagged 0–3 years (depending on stock) to align with climate

conditions and spawner status (biomass and age structure) in spawning year, whereas *Fishing*

was lagged an additional year (spawning year–1) to reflect indirect effects of fishing on

recruitment success that are not accounted for by variations in spawners. Regional climate

covariates were also lagged an additional year (*Climate*$_{(i-1)j}$) to reflect indirect effects through

regulation of local climate and ecosystem processes. Time (year)-varying covariate smoothers

were tested as tensor product construction (with $5 \times 2$ maximum knots using the *t2* function in

the R package *mgcv*). For interpretability and model convergence one covariate of each type was

tested at a time; one local or regional climate index (see above) for climate, SB or SB/$B_{pa}$ for

spawner density, $\bar{F}$ or $\bar{F}/F_{msy}$ for fishing pressure, and SA$_{mean}$ or SA$_{diversity}$ for demographics.

*Covariates of elasticity to recruitment success*. The transient dynamics of depleted fish stocks

with age truncation often become recruitment-dependent and thus sensitive to variations in

climate and habitat conditions. As the stocks recover (increases in abundance and age diversity),

the transient dynamics are expected to become less sensitive to recruitment variability. To test

for the time-varying contributions of the extrinsic and intrinsic factors to variations in elasticity

to RS ($e_{rec}$), the following model was fitted to the data, assuming $e_{rec}$ follows a Beta distribution

with expected values $\mu$ ($0 < \mu < 1$) and a logit link function as

351
$$e_{rec,ijk} \sim \text{Beta}(\mu_{ijk}, \phi)$$

352
$$E(e_{rec,ijk}) = \mu_{ijk}$$

353
$$\text{var}(e_{rec,ijk}) = \mu_i (1 - \mu_i)/(\phi + 1)$$

$$\text{logit}(\mu_{ijk}) = \beta_0 + f_1(\text{Year}_{ij}) + f_2(\text{Year}_{ij}, \text{Biomass}_{ijk}) + f_3(\text{Year}_{ij}, \text{Demography}_{ijk}) +$$

$$f_4(\text{Year}_{ij}, \text{Fishing}_{(i\text{-}1)jk}) + f_5(\text{Year}_{ij}, \text{Climate}_{ij}) + \text{stock}_i + \text{ecoregion}_j + \text{year}_{jk}$$

$$\text{stock}_i \sim N(0, \sigma^2_{stock}), \text{ecoregion}_j \sim N(0, \sigma^2_{ecoregion}), \text{year}_{jk} \sim N(0, \sigma^2_{year}) \qquad (8)$$

where $\phi$ is a parameter ($> 0$), and $\beta_0, f_1\text{–}f_5, \sigma_{stock}, \sigma_{ecoregion}$, and $\sigma_{year}$, are the same as above

(here tensor product construction for time-varying covariate smoothers was modeled using the *te*

function in the R package *mgcv*).

All numeric covariates were standardized by subtracting the mean and dividing by the

standard deviation (*z*-scores). In model selection intrinsic covariates were first added, followed

by indirect fishing, and then climate. With the random effect structure selected (as described

above, Table S1) the models were tested sequentially for covariates by fitting with a maximum

likelihood method, followed by parameter (re)estimation with REML (Zuur *et al.* 2009) for the

model with the lowest Akaike information criterion (AIC) score and highest model weight (*w*).

AIC is computed as AIC $= -2\ln(L) + 2k$, where $L$ is the likelihood of the model and $k$ is the

number of parameters, and the model weight of model $i$ ($w_i$) is computed as

$$w_i = \frac{\exp\left(-\Delta AIC_i/2\right)}{\sum_j \exp\left(-\Delta AIC_i/2\right)} \qquad (9)$$

As diagnostics, model residuals were checked for homogeneity and regressed against

covariates that were both included and not included in the selected models (Zuur *et al.* 2009). All

analyses were performed in R (version 4.2.1, R Development Core Team 2022) with the R

packages *gamm4* (v.0.2-6, Wood & Scheipl 2020) and *mgcv* (v.1.8-40, Wood 2022) for model-

fitting, *mgcv* and *gratia* (v.0.7.3, Simpson & Singmann 2022) for model diagnostics, and *popbio*

(v.2.7, Stubben, Milligan & Nantel 2020) and *popReconstruct* (1.0-7, Wheldon 2019) for

computations of demographic parameters and projections of population dynamics.

Because fish populations in different ecoregions experience a wide range of climate

conditions and often have adapted to local environments, fish–covariate patterns may deviate

from shared patterns captured by the regional models. To test this hypothesis, ecoregion-scale

analyses were also performed using the same model structures for $RS_{total}$ and $e_{rec}$ and the model

selection steps as above except for random effects. Because random effect estimates become less

stable when the number of groups is fewer than eight (Bolker 2015), the random effect of the

ecoregion-scale models consisted of year only, with stock being a fixed effect, except for the

Celtic Seas; the random effects of the Celtic Seas model consisted of year and stock.

**Results**

Analyses were performed to address primarily (a) whether long-term changes in spawners

(biomass and age structure) shared by intensely managed fish stocks in the northeast Atlantic

reflect management-guided changes in fishing pressure, (b) how intrinsic (spawner biomass and

age structure) and extrinsic (fishing and climate) drivers contribute to variations in recruitment

success, and (c) how the intrinsic and extrinsic drivers influence the sensitivity of transient

dynamics to recruitment variability. Overall, the diverging patterns in spawner biomass between

the recovering and the non-recovering stocks also emerged in age structure and transient

population dynamics, reflecting differential reductions in fishing pressures the stocks

experienced. The differences in spawner biomass and age structure between the groups were

further reflected in variations in recruitment success and elasticity to recruitment success,

indicating divergence in the strength of density dependency and the contribution of older

spawners. Although climate effects varied among ecoregions, most showed shifting responses in

recruitment (from positive to negative) and in elasticity to recruitment (from negative to positive)

to increasingly warmer than average local and regional climates.

Regionwide analyses showed that relative fishing pressures on the recovering stocks on

average peaked in the early 1990s ($\bar{F}/F_{msy}$ = ~2.2) and near-linearly declined since (~2.1% per

year; $\Delta$AIC = 88.4), followed by increasing mean age ($\Delta$AIC = 35.7), relative spawner biomass

($\Delta$AIC = 17.2), and age diversity ($\Delta$AIC = 21.1) (Fig. 2a). In contrast, relative fishing pressures

on the non-recovering stocks kept rinsing until the late 1990s (to $\bar{F}/F_{msy}$ = ~3.7 on average)

before declining (~3.0% per year, $\Delta$AIC = 63.3), but their relative spawner biomass continued to

decline ($\Delta$AIC = 92.7 and 31.5; Fig. 2a). Although the mean spawner age of the non-recovering

stocks began rising with declining relative fishing pressures, no change in age diversity was

detected (Fig. 2a). Direct effects of selective fishing consistently emerged as negative

correlations between relative fishing pressure and spawner biomass and age structure in both

stock groups ($\Delta$AIC = 15.1–165.0, Fig. 2b). Contrasting temporal patterns between relative

fishing pressure and spawner features were more pronounced in some ecoregions captured in

year-specific (nested within ecoregion) random effect (variance) estimates, especially the

Norwegian–Barents Seas, which showed abrupt shifts in the early 1990s (Fig. S5).

The time series of transient population growth rates reflected the contrasting patterns between

the recovering and the non-recovering stocks, and variations among ecoregions (Fig. 3 and S6).

Sensitivity analyses showed that multilevel model parameter estimates in the trend analyses for

both stock groups were robust to uncertainties in select abundance and demographic rates used as

input (simulated parameter values < 2 standard deviations of the estimates, Fig. S7). The regional

average growth rates of the recovering stocks linearly increased over the study period ($\Delta$AIC =

15.0, Fig. 3). Although the relative spawner biomass of the non-recovering stocks continued to

decline, their average growth rates began rising in the late 1990s ($\Delta$AIC = 2.8, Fig. 3) when the

relative fishing pressures began declining (Fig. 2a). These transient dynamics differed from

asymptotic dynamics; the transient growth rates were lower than asymptotic growth rates in most

stocks, and the deviations varied over years, with ~50% (19) of the stocks showing increasingly

less differences in the last decade (Fig. S8). The observed age distributions of many of these

stocks also differed from theoretical stable age distributions and skewed toward younger ages

(Fig. S9) owing to disproportionately higher annual mortality rates of older fish by fishing (Fig.

S10).

Regionwide analyses detected declining trends in per capita recruitment success rate in both

recovering and non-recovering stocks ($\Delta$AIC = 2.1 and 2.7, Fig. 4a), with high among-stock

variances ($\sigma_{stock}^2$ = 0.31 and 0.48, Fig. 4f-h). Ecoregion-specific random intercepts also revealed

generally higher year-to-year fluctuations in the non-recovering than in the recovering stocks

(Fig. S6). After accounting for variations among stocks and ecoregions (and years within each

ecoregion), the recruitment rates of both stock groups declined with increasing relative spawner

biomass (Fig. 4c and Table 2), which was more supported by data than absolute biomass (Table

S2). But the temporal patterns of density dependency differed between the groups; the density

dependent effect in the recovering stocks increased as the biomass rose, whereas that in the non-

recovering stocks emerged in the last two decades but declined over time (Fig. 4c). Likewise, the

contribution of spawner age structure to recruitment variability differed between the groups; the

relationship between recruitment success and mean age in the recovering stocks shifted from

positive to negative in the early 1990s, whereas that in the non-recovering stocks remained

largely positive (Fig. 4b). An indirect positive fishing effect on recruitment rate was detected in

the recovering stocks until the late 1980s but gradually diminished since (Fig. 4d). The models

for both stock groups showed weak support for climate effects (Table S2). Of the climate indices

tested for the non-recovering stocks (whose model had slightly more support than the recovering

stocks), the model with ocean heat content anomaly in the year of spawning showed shifting

(positive to negative) climate effects in the recent decades (Fig. 4e).

Ecoregion-scale analyses on recruitment success revealed patterns largely consistent with the

regional analyses with some system-specific deviations (Fig. S11). In ecoregions with non-

recovering stocks like the Baltic, Celtic, and North Seas the density-dependent effect on

recruitment on average became weakened over time; in contrast, the effect in the Norwegian–

Barents Seas (with only recovering stocks) became strengthened (Fig. S11c). Correlations with

spawner age structure metrics were detected in all but the Bay of Biscay and Faroes (Fig. S11b);

in the Baltic Sea a positive correlation with age diversity became detectable in the early 1990s

when fishing pressures began declining (Fig. 2). The time-varying effect of spawner age was

especially pronounced in the Norwegian–Barents Seas stocks, showing a shift from positive to

negative in the 1990s (Fig. S11b). An indirect fishing effect was (weakly) supported in the North

Sea stocks only (Table S3 and Fig. S11d). Climate anomalies (in the year of spawning)

contributed to variability in recruitment success in all ecoregions except the North Sea but the

indices with the most statistical support varied among ecoregions (Fig. 5 and Table S3) despite

high correlations among climate indices (Fig. S12). The ways climate anomalies contributed to

recruitment variability also differed (Fig. 5). In the Baltic Sea and Faroes correlations with

climate anomalies remained largely positive during the past few decades (Fig. 5a,d). In contrast,

the positive effect in the Norwegian–Barents Seas began diminishing in the recent decades (Fig.

5f). In the Bay of Biscay and Icelandic waters the climate effect shifted from positive to negative

(Fig. 5b,e).

Trend analyses revealed consistent temporal patterns in elasticities to vital rates for the

recovering and non-recovering stocks. Both showed declining elasticities to (post-recruitment)

juvenile survival rate ($\Delta$AIC = 15.6 and 18.9) and increasing elasticities to adult survival rate

over time ($\Delta$AIC = 17.0 and 6.7) (Fig. S13). But the regionwide analyses detected a declining

trend in elasticities to recruitment success rate in the recovering stocks only ($\Delta$AIC = 3.6, Fig.

6a) with less among-stock variances than recruitment success (Fig. 6f-h). Ecoregion-specific

variances in all the models fluctuated over years, but on average elasticities to adult survival

showed higher deviations from the regional trends than the other vital rates (Fig. S13).

Regionwide analyses on elasticities to recruitment success largely reflected the time-varying

responses in recruitment to intrinsic and extrinsic drivers. For the recovering stocks the model

indicated that more diverse ages increasingly contributed to elasticities in more recent years,

whereas the contributions of mean ages gradually increased in the non-recovering stocks (Fig.

6b). Elasticities to recruitment declined with higher spawner biomass in both stock groups, but

the density-dependent effect in the non-recovering stocks diminished over time (Fig 6c). The

model with an indirect fishing effect in both stock groups showed a declining contribution to

variations in the elasticities after the reductions in fishing pressure (Fig. 6d). In contrast, the

models for both stock groups indicated an increasingly more positive response to climate

anomalies (the AMO index and winter sea temperature anomaly) over time (Fig. 6e).

Ecoregion-scale analyses on elasticities to recruitment success detected declining trends in all

but the Bay of Biscay and the Norwegian–Barents Seas (Fig S14a), which showed little change

in trends. The analyses also indicated spatial heterogeneity in the ways time-varying intrinsic and

extrinsic drivers contributed to variations in the elasticities. For example, shifting contributions

of spawner age structure were detected in most ecoregions, showing both diminishing (e.g.,

Faroes and the North Sea) and increasing (e.g., the Baltic Sea and Icelandic waters) trends (Fig.

S14b). Negative correlations with spawner biomass were consistently detected except in the

Baltic Sea (Fig. S14c). This maternal effect increased over time except in the Norwegian–

Barents Seas, which showed the opposite trend (Fig. S14c). The models with an indirect fishing

effect had statistical support for Faroes, and the Celtic, North, and Norwegian–Barents Seas; the

fishing effect declined and/or shifted over time except the Norwegian–Barents Sea, which

showed an increasing contribution of fishing since the 1990s (Fig. S14d). In the Baltic Sea and

the Bay of Biscay a positive climate effect emerged in the 2000s (Fig. 7a,b), reflecting the

temporal patterns of shifting local climate effects on recruitment success (Fig. 5a,b). In Faroes

and Icelandic waters, where climate anomalies (the NAO index and spring sea temperature) had

a positive effect on recruitment, the elasticities were inversely correlated with the indices, but

this effect shifted to the opposite in the 1990s–2000s (Fig. 7c,d).

**Discussion**

Analyses on the transient dynamics of northeast Atlantic fish stocks reveal that management-

aided reductions in fishing pressure and shifting climate conditions nonlinearly shaped

rebuilding patterns of these historically depleted stocks through restorations of demographic

structure (reversing age truncation) and reproductive capacity in recent decades. As the survival

rates and demographic structures of reproductive fish improved with declining fishing pressures,

transient population growth rates became less sensitive to variability in recruitment and juvenile

survival and more to that in adult survival, allowing the stocks to rebuild biomass despite

declining recruitment success. As the biomass of reproductive fish rebuilt, recruitment success

rates also became increasingly regulated by density-dependent processes involving increasingly

higher numbers of older reproductive fish. Although warmer than average climates promoted

recruitment success throughout the early 2000s in some ecoregions, warming trends over the past

decades began adversely affecting reproductive performance overall. Variations in regulation of

fishing pressure among stocks also led to diverging trends in recoveries of the stocks exposed to

a range of local climate conditions. Insufficient or delayed reductions in fishing pressure can

further deplete a stock and erode reproductive fish demographic structure, thereby diminishing

fecundity and thus recovery potential, while also remaining sensitive to climate variability.

Rebuilding biomass by accounting for demographic structure through management are thus not

only key to sustainable use of marine resource populations but also the foundation for effective

applications of adaptive measures resilient to future environmental change.

*Transient dynamics of recovering populations from intense fishing pressures*

Fishing-induced changes in the demography and reproductive capacity of an exploited

population can be reversible when management measures are effectively applied to regulate

fishing pressure. This pattern is displayed especially in the Norwegian–Barents Seas, where

fishing pressures were reduced by more than 60% (compared to the regional average of ~50%)

since the early 1990s, followed by sharp rises in the average age and biomass of reproductive

fish. But when a population experiences long-term intensive harvesting, recovery time can

become longer (Hutchings 2015) and more uncertain (Neubauer *et al.* 2013), with the population

remaining in long transient states (Hastings *et al.* 2018). Although fishing pressures on many

northeast Atlantic fish stocks precipitously declined through stricter management measures

applied since the late 1990s, demographic restorations and stock recoveries on average

progressed gradually. Reducing fishing pressures can allow more adults to survive and contribute

to the production of new recruits but the restoration of the demographic structure takes place

over generations and may take longer (a decade or more) for severely depleted stocks. Further,

although the declining trends in reproductive fish biomass halted in the early 2000s in most

ecoregions, reduction levels in fishing pressure varied among ecoregions and stocks and were

reflected in recovery patterns. When reductions in fishing pressure were insufficient or delayed,

the declining trends in reproductive capacity and deterioration of demographic structure in some

stocks continued. Slow recoveries of severely depleted fish stocks are reportedly widespread

(Hutchings 2000). Regionwide analyses on the non-recovering stocks in the northeast Atlantic

reveal that sustaining high fishing pressures for even a few more years can result in vastly

different recovery patterns, with further age truncations and reductions in reproductive capacity,

which could further diminish fishery yields (Hilborn *et al.* 2020).

Removal of some biomass from a population by harvesting can promote its productivity

(Neubauer *et al.* 2013) by attenuating density-dependent effects on early life stage processes

(competition for food and space for example, Anderson *et al.* 2008). Intense fishing pressures

before the implementation of stricter management measures–mediated through reproductive

fish–may have led to higher recruitment rates in depleted stocks in the northeast Atlantic Ocean.

Fishing not only truncates age structure but induces changes in life-history traits that may

influence recovery (Neubauer *et al.* 2013; Hutchings & Kuparinen 2020). The generation times

of these depleted northeast Atlantic stocks were shortened (by 14 to 50%), which is often

associated with high year-to-year variability in recruitment (Minto, Myers & Blanchard 2008)

and high average recruitment rates (Bjørkvoll *et al.* 2012). Higher growth rates of depleted (and

age-truncated) fish populations also likely benefited during the early stages of recovery as more

reproductive fish contributed to spawning events, thereby increasing the probability of

recruitment success. These indirect effects, however, diminished over time following the

reductions of fishing pressures. Despite the signs of recovery in reproductive fish, per capita

recruitment success rates on average declined in the northeast Atlantic stocks, as also reported in

other Atlantic ecoregions (Britten, Dowd & Worm 2016). But these apparent patterns were

shaped by different processes among stocks. In the recovering stocks density dependency

remained present in recruitment success throughout the past decades and its strength increased as

reproductive fish biomass rose, likely slowing recoveries as fishing pressures declined. In

contrast, in the non-recovering stocks density dependency in recruitment success emerged only

after the reductions in fishing pressure but its strength continued to decline with reproductive fish

biomass, suggesting contributions of other (extrinsic) drivers.

Density dependency in early life also perhaps explains discontinuous relationships (shifting

from positive to negative) between reproductive fish age and per capita recruitment success in

the recovering stocks (especially in the Norwegian–Barents Seas) following the reductions in

fishing pressure. Higher mean age and age diversity in reproductive fish are expected to promote

recruitment success (Rogers & Dougherty 2019). Age-dependent variations in spawning pattern

can influence egg viability and larval survival; older spawners tend to produce more viable

offspring (Hixon, Johnson & Sogard 2014). But the analyses on the northeast Atlantic fishes

show that only the non-recovering (age-truncated) stocks kept positive relationships between

reproductive fish age structure and recruitment success. In the recovering stocks, density

dependency may have played a larger role than maternal effects, revealing a shift in

compensatory processes. As adult survival improves and population density increases, pre-recruit

survival may decline (owing to competition for food and other resources including cannibalism

by adults), contributing to apparent increases in average ages of reproductive fish.

Short-term (year-to-year) fluctuations in recruitment can propagate through transient dynamics

of a population when its (demographic) buffering capacity is compromised by age truncation

(Anderson *et al.* 2008). Truncated age distributions in reproductive fish may produce volatile

population behaviors that reflect recruitment pulses (Anderson *et al.* 2008). As adult survival

improved and stock biomass rebuilt in many northeast Atlantic fish stocks, the transient growth

became less sensitive to recruitment variability and more sensitive to adult survival. The

regionwide analyses show that increases in transient growth rate and biomass during the past few

decades resulted largely from improved survival of older fish. Greater sensitivities to adult

survival also may make harvested populations more manageable. Nonlinear responses to age

structure in reproductive fish is however also evident in sensitivity to recruitment variability.

Although the buffering effect of increasingly age-diverse reproductive fish emerged in many

ecoregions especially in the early stages of recovery, this relationship also shifted as the

recoveries further progressed (as seen in the Norwegian–Barents Seas fishes), likely responding

to other intrinsic or extrinsic dynamics (such as catch limits being raised as stock biomass

rebuilds).

*Climate regulation of demographic variations*

Fish populations experiencing persistent selective depletions of older adults often become

sensitive to variations in early life-history stage processes that reflect signals in environmental

conditions like climate variability (Ottersen, Hjermann & Stenseth 2006; Rouyer *et al.* 2011).

Warmer than average climate conditions in the northeast Atlantic Ocean increasingly contributed

to observed patterns in fish recruitment success, reflecting shifting climate effects on recruitment

processes. For example, stratification becomes intensified as climate warms, limiting nutrient

movement to deeper waters and food accessibility for the early life of demersal and benthic

species (Moore *et al.* 2018). At the region scale, however, the models captured only weak

responses in recruitment to regional climate signals likely because of heterogeneous warming

patterns among ecoregions, as also seen in other parts of the world (Barange *et al.* 2018).

Region-scale analyses may mask differential effects of local climate and oceanographic features

(climate-induced shifts in ocean heat content and primary productivity for example) that marine

species are adapted to and experience (Conover 1992).

Ecoregion-scale variability in recruitment success may better reflect how fish reproductive

behavior responds to regional climate mediated by local habitat conditions, as hypothesized in

past research (Myers 1998; Ottersen & Stenseth 2001). For stocks in high latitude systems like

the Norwegian–Barents Seas and Icelandic waters, for example, climate conditions in winter–

spring (December–May) that regulate phonologies like timings of spawning migration (Sundby

& Nakken 2008) contributed to recruitment variability. Warming-induced reductions in sea ice

extent in these ecoregions promoted ecosystem productivity in the late 1990s and early 2000s

and may have benefited upper trophic animals like fish (Stenevik & Sundby 2007), likely

through changes in growth rate and life stage duration (Houde 2016). When spawning times

coincide with the period of high prey abundance, pre-recruit fish can grow more and experience

less size-dependent mortalities (like predation), contributing to greater recruitment success

(Peck, Huebert & Llopiz 2012). In contrast, in semi-enclosed systems like the Baltic Sea

warming-induced changes in nearshore habitat conditions in summer (June–August) (lower

salinity caused by higher precipitation and freshwater inflows for example, MacKenzie *et al.*

2007) may have played a greater role in regulating recruitment success through biophysical

processes during the early years of life.

Demographic responses by fish populations to changing climate conditions also may vary

discontinuously (Brander 2007). Local and regional climate conditions (both favorable and

unfavorable) that early life-history stage fish experience can persist through time (Ottersen *et al.*

2001) and may emerge as delayed effects on transient population dynamics. In some northeast

Atlantic ecoregions the ways that climate affects fish recruitment success shifted from positive to

negative and may have slowed their recoveries. These shifts suggest that even in the ecoregions

where warming climate promoted productivity in the early twentieth century, fish began to

experience adverse (direct and indirect) effects. Mobile species can adjust their movement

(deeper and poleward) to mitigate physiological stress from warming upper water columns

(Pinsky *et al.* 2013; Engelhard, Righton & Pinnegar 2014). But asynchronous shifts in spatial

distribution by spawning fish and seasonal events in lower trophic groups like spring plankton

blooms (phenological mismatches) also could trigger a series of events unfavorable for early life

fish (Asch & Erisman 2018), shrinking the probability of recruitment success.

Climate variability that regulates local ocean conditions may propagate through vital rates to

shape transient population dynamics of marine species (Jenouvrier *et al.* 2018). Depleted

populations in particular may have heightened sensitivities to adverse environmental conditions

(Hidalgo *et al.* 2011; Rouyer *et al.* 2011), which are becoming increasingly more variable in

recent decades (Rathore *et al.* 2020). Although the sensitivities of transient growth to recruitment

variability in the northeast Atlantic fish stocks declined as they recovered, the analyses reveal

that increasingly warmer than average climates began influencing how recruitment variability

propagates through transient dynamics. For example, in high-latitude systems like Faroes, where

fish stocks become less sensitive to recruitment variability under a warmer climate, this

stabilizing effect gradually diminished (and shifted to the opposite) as the climate further

warmed. In contrast, in the Baltic Sea, where fish stocks become more sensitive to recruitment

variability under a warmer climate, this amplifying effect became strengthened as the climate

further warmed. Climate effects on transient dynamics also may, however, depend on population

size and demographic structure. In the Norwegian–Barents Seas, where fish stocks experienced

higher reductions in fishing pressure early and thus higher biomass rebuilding rates than other

stocks, the stocks responded to recruitment variability independently of climate conditions.

*Transients as an integrated measure of effects of fishing, management, and climate*

Analysis of transient population dynamics can reveal how time-varying demographic structure

and vital rates contribute to recovery patterns of managed resource populations when also

exposed to low-frequency, nonstationary drivers like climate. Declined sensitivities to highly

variable demographic parameters like recruitment, as displayed by intensely managed northeast

Atlantic fish stocks in this study, can provide a buffer against rising environmental variability.

Increasing sensitivities to adult survival further suggest that the stocks also are expected to

become more responsive to management actions through regulation of fishing pressure

(enforcing positive feedback), and future management plans would benefit from maintaining

healthy (age-diverse) reproductive populations of exploited species (Pinsky & Byler 2015).

Combined with other management measures (like use of marine protected areas and fish size

limits, White *et al.* 2013; Hixon, Johnson & Sogard 2014), restoring and maintaining the

demographic structure of an exploited species thus may enhance resilience against destabilizing

effects of environmental variability amplified by climate change.

Emerging non-negligible changes in demographic responses to rising climate variability

nevertheless may influence the efficacy of fisheries management. Although historical climate

conditions promoted the productivity of fish stocks in the northeast Atlantic Ocean, some stocks

in this region and elsewhere in the Northern Hemisphere already are experiencing adverse effects

of changing climate (Pershing *et al.* 2015; Free *et al.* 2019). As many biological responses to

climate-driven changes in surface temperature and other ocean properties are often nonlinear, we

would expect continued shifts in how climate influences marine resource populations and

management (Britten, Dowd & Worm 2016). Accounting for these climate-induced changes thus

may not only reduce uncertainties in assessment but also improve the performance of

management measures (Rouyer et al. 2011; Bahri et al. 2021).

Exploited marine fish species are an integral part of an ecosystem and changes in their

abundances and demographic structures may propagate through community and ecosystem

dynamics (Jennings & Kaiser 1998), triggering a series of changes in ecologically connected

species (Goto *et al.* 2022; Pérez-Rodríguez *et al.* 2022). As adult survival improves in

historically overharvested populations, ecological processes including resource competition and

predation also may indirectly modify vital rates and demographic structures of non-target species

(van Gemert & Andersen 2018). Improving our understanding of how warming and more

variable climates continue to reshape the transient dynamics (through changes in life history

traits and demographics) would help develop management plans to mitigate the risk of

overexploitation and safeguard ecosystem services against rising environmental variability.

**Acknowledgements**

This work was partially supported by the Institute of Marine Research's Reduced Uncertainty

in Stock Assessment (REDUS) project.

**Data Availability Statement**

The data that support the findings of this study are openly available in ICES Scientific

Reports at https://ices-library.figshare.com/, reference number ICES CM 2017/ACOM:06, ICES

CM 2017/ACOM:08, ICES CM 2017/ACOM:11, ICES CM 2017/ACOM:12, ICES CM

2017/ACOM:13, and ICES CM 2017/ACOM:21, and in the data archives of Met Office Hadley

Centre, and NOAA Physical Sciences Laboratory and National Centers for Environmental

Information at https://www.metoffice.gov.uk, https://psl.noaa.gov, and

https://www.ncei.noaa.gov. Code and derived datasets (age-specific demographic parameter

estimates) will be available upon publication.

**References**

Anderson, C.N., Hsieh, C.-h., Sandin, S.A., Hewitt, R., Hollowed, A., Beddington, J., May, R.M.

& Sugihara, G. (2008) Why fishing magnifies fluctuations in fish abundance. *Nature,* **452,**

835.

Asch, R.G. & Erisman, B. (2018) Spawning aggregations act as a bottleneck influencing climate

change impacts on a critically endangered reef fish. *Diversity and Distributions,* **24,** 1712-

1728.

Barange, M., Bahri, T., Beveridge, M.C., Cochrane, K.L., Funge-Smith, S. & Poulain, F. (2018)

Impacts of climate change on fisheries and aquaculture: synthesis of currrent knowledge,

adaptation and mitigation options. Rome, FAO. 628 pp.

Beaugrand, G. (2009) Decadal changes in climate and ecosystems in the North Atlantic Ocean

and adjacent seas. *Deep Sea Research Part II: Topical Studies in Oceanography,* **56,** 656-673.

Beaugrand, G. & Kirby, R.R. (2018) How do marine pelagic species respond to climate change?

Theories and observations. *Annual Review of Marine Science,* **10,** 169-197.

Beddington, J.R., Agnew, D.J. & Clark, C.W. (2007) Current problems in the management of

marine fisheries. *Science,* **316,** 1713-1716.

Bjørkvoll, E., Grøtan, V., Aanes, S., Sæther, B.-E., Engen, S. & Aanes, R. (2012) Stochastic

population dynamics and life-history variation in marine fish species. *The American*

*Naturalist,* **180,** 372-387.

Bolker, B.M. (2015) Linear and generalized linear mixed models. *Ecological statistics:*

*contemporary theory and application* (eds G.A. Fox, S. Negrete-Yankelevich & V.J. Sosa),

pp. 309-333. Oxford University Press, Oxford, United Kingdom.

Brander, K.M. (2007) Global fish production and climate change. *Proceedings of the National*

*Academy of Sciences,* **104,** 19709-19714.

Britten, G.L., Dowd, M. & Worm, B. (2016) Changing recruitment capacity in global fish stocks.

*Proceedings of the National Academy of Sciences,* **113,** 134-139.

Brooks, E.N. & Deroba, J.J. (2015) When "data" are not data: the pitfalls of post hoc analyses

that use stock assessment model output. *Canadian Journal of Fisheries and Aquatic Sciences,*

**72,** 634-641.

Caswell, H. (2001) *Matrix population models: construction, analysis, and interpretation,* 2 edn.

Sinauer Sunderland.

Chassot, E., Bonhommeau, S., Dulvy, N.K., Mélin, F., Watson, R., Gascuel, D. & Le Pape, O.

(2010) Global marine primary production constrains fisheries catches. *Ecology Letters,* **13,**

495-505.

Cheung, W.W., Watson, R. & Pauly, D. (2013) Signature of ocean warming in global fisheries

catch. *Nature,* **497,** 365-368.

Christensen, V., Guenette, S., Heymans, J.J., Walters, C.J., Watson, R., Zeller, D. & Pauly, D.

(2003) Hundred-year decline of North Atlantic predatory fishes. *Fish and fisheries,* **4,** 1-24.

Conover, D. (1992) Seasonality and the scheduling of life history at different latitudes. *Journal*

*of Fish Biology,* **41,** 161-178.

Costello, C., Ovando, D., Clavelle, T., Strauss, C.K., Hilborn, R., Melnychuk, M.C., Branch,

T.A., Gaines, S.D., Szuwalski, C.S. & Cabral, R.B. (2016) Global fishery prospects under

contrasting management regimes. *Proceedings of the National Academy of Sciences,* **113,**

5125-5129.

Coulson, T., Guinness, F., Pemberton, J. & Clutton-Brock, T. (2004) The demographic

consequences of releasing a population of red deer from culling. *Ecology,* **85,** 411-422.

Crowder, L.B., Hazen, E.L., Avissar, N., Bjorkland, R., Latanich, C. & Ogburn, M.B. (2008)

The impacts of fisheries on marine ecosystems and the transition to ecosystem-based

management. *Annual Review of Ecology, Evolution, and Systematics***,** 259-278.

Delworth, T.L., Zeng, F., Vecchi, G.A., Yang, X., Zhang, L. & Zhang, R. (2016) The North

Atlantic Oscillation as a driver of rapid climate change in the Northern Hemisphere. *Nature*

*Geoscience,* **9,** 509-512.

Durant, J.M., Hidalgo, M., Rouyer, T., Hjermann, D.Ø., Ciannelli, L., Eikeset, A.M., Yaragina,

N. & Stenseth, N.C. (2013) Population growth across heterogeneous environments: effects of

harvesting and age structure. *Marine Ecology Progress Series,* **480,** 277-287.

Engelhard, G.H., Righton, D.A. & Pinnegar, J.K. (2014) Climate change and fishing: a century

of shifting distribution in North Sea cod. *Global Change Biology,* **20,** 2473-2483.

Faillettaz, R., Beaugrand, G., Goberville, E. & Kirby, R.R. (2019) Atlantic Multidecadal

Oscillations drive the basin-scale distribution of Atlantic bluefin tuna. *Science Advances,* **5,**

eaar6993.

FAO (1995) Precautionary approach to fisheries. Part 1: Guidelines on the precautionary

approach to capture fisheries and species introductions. *FAO Fisheries Technical Paper*, pp.

52 p., Rome, FAO.

Free, C.M., Thorson, J.T., Pinsky, M.L., Oken, K.L., Wiedenmann, J. & Jensen, O.P. (2019)

Impacts of historical warming on marine fisheries production. *Science,* **363,** 979-983.

Gamelon, M., Sandercock, B.K. & Sæther, B.E. (2019) Does harvesting amplify environmentally

induced population fluctuations over time in marine and terrestrial species? *Journal of*

*Applied Ecology,* **56,** 2186-2194.

Gelman, A. & Hill, J. (2006) *Data analysis using regression and multilevel/hierarchical models*.

Cambridge university press.

Goto, D., Devine, J.A., Umar, I., Fischer, S.H., De Oliveira, J.A., Howell, D., Jardim, E.,

Mosqueira, I. & Ono, K. (2022a) Shaping sustainable harvest boundaries for marine

populations despite estimation bias. *Ecosphere,* **13,** e3923.

Goto, D., Dunlop, E.S., Young, J.D. & Jackson, D.A. (2020) Shifting trophic control of fishery–

ecosystem dynamics following biological invasions. *Ecological Applications,* **30,** e02190.

Goto, D., Filin, A.A., Howell, D., Bogstad, B., Kovalev, Y. & Gjøsæter, H. (2022b) Tradeoffs of

managing cod as a sustainable resource in fluctuating environments. *Ecological Applications,*

**32,** e2498.

Harris, R.M., Beaumont, L.J., Vance, T.R., Tozer, C.R., Remenyi, T.A., Perkins-Kirkpatrick,

S.E., Mitchell, P.J., Nicotra, A., McGregor, S. & Andrew, N. (2018) Biological responses to

the press and pulse of climate trends and extreme events. *Nature climate change,* **8,** 579-587.

Hastings, A. (2004) Transients: the key to long-term ecological understanding? *Trends in*

*Ecology & Evolution,* **19,** 39-45.

Hastings, A., Abbott, K.C., Cuddington, K., Francis, T., Gellner, G., Lai, Y.-C., Morozov, A.,

Petrovskii, S., Scranton, K. & Zeeman, M.L. (2018) Transient phenomena in ecology.

*Science,* **361,** eaat6412.

Hidalgo, M., Rouyer, T., Bartolino, V., Cervino, S., Ciannelli, L., Massuti, E., Jadaud, A.,

Saborido-Rey, F., Durant, J.M. & Santurtún, M. (2012) Context-dependent interplays between

truncated demographies and climate variation shape the population growth rate of a harvested

species. *Ecography,* **35,** 637-649.

Hidalgo, M., Rouyer, T., Molinero, J.C., Massutí, E., Moranta, J., Guijarro, B. & Stenseth, N.C.

(2011) Synergistic effects of fishing-induced demographic changes and climate variation on

fish population dynamics. *Marine Ecology Progress Series,* **426,** 1-12.

Hilborn, R., Amoroso, R.O., Anderson, C.M., Baum, J.K., Branch, T.A., Costello, C., de Moor,

C.L., Faraj, A., Hively, D. & Jensen, O.P. (2020) Effective fisheries management instrumental

in improving fish stock status. *Proceedings of the National Academy of Sciences.*

Hilborn, R., Maguire, J.-J., Parma, A.M. & Rosenberg, A.A. (2001) The precautionary approach

and risk management: can they increase the probability of successes in fishery management?

*Canadian Journal of Fisheries and Aquatic Sciences,* **58,** 99-107.

Hilborn, R. & Walters, C.J. (1992) *Quantitative fisheries stock assessment: choice, dynamics and*

*uncertainty*. Springer New York, NY.

Hixon, M.A., Johnson, D.W. & Sogard, S.M. (2014) BOFFFFs: on the importance of conserving

old-growth age structure in fishery populations. *ICES Journal of Marine Science,* **71,** 2171-

2185.

Hjermann, D.Ø., Bogstad, B., Eikeset, A.M., Ottersen, G., Gjøsæter, H. & Stenseth, N.C. (2007)

Food web dynamics affect Northeast Arctic cod recruitment. *Proceedings of the Royal Society*

*of London B: Biological Sciences,* **274,** 661-669.

Houde, E.D. (2016) Recruitment variability. *Fish reproductive biology: implications for*

*assessment and management* (eds T. Jakobsen, M.J. Fogarty, B.A. Megrey & E. Moksness),

pp. 98-187. John Wiley & Sons Ltd, West Sussex, United Kingdom.

Hsieh, C.-h., Reiss, C.S., Hunter, J.R., Beddington, J.R., May, R.M. & Sugihara, G. (2006)

Fishing elevates variability in the abundance of exploited species. *Nature,* **443,** 859-862.

Hsieh, C.-h., Yamauchi, A., Nakazawa, T. & Wang, W.-F. (2010) Fishing effects on age and

spatial structures undermine population stability of fishes. *Aquatic Sciences,* **72,** 165-178.

Hutchings, J.A. (2000) Collapse and recovery of marine fishes. *Nature,* **406,** 882.

Hutchings, J.A. (2015) Thresholds for impaired species recovery. *Proceedings of the Royal*

*Society B: Biological Sciences,* **282,** 20150654.

Hutchings, J.A. & Kuparinen, A. (2020) Implications of fisheries-induced evolution for

population recovery: Refocusing the science and refining its communication. *Fish and*

*Fisheries,* **21,** 453-464.

Hutchings, J.A. & Reynolds, J.D. (2004) Marine fish population collapses: consequences for

recovery and extinction risk. *BioScience,* **54,** 297-309.

ICES (2016) Report of the Benchmark Workshop on North Sea Stocks (WKNSEA), 14–18

March 2016, Copenhagen, Denmark. ICES CM 2016/ACOM:37 704 pp.

ICES (2017a) Report of the Arctic Fisheries Working Group (AFWG), 19–25 April 2017,

Copenhagen, Denmark. ICES CM 2017/ACOM:06. 493 pp.

ICES (2017b) Report of the Baltic Fisheries Assessment Working Group (WGBFAS), 19- 26

April 2017, Copenhagen, Denmark. ICES CM 2017/ACOM:11. 810 pp.

ICES (2017c) Report of the North Western Working Group (NWWG), 27 April – 4 May 2017,

Copenhagen, Denmark. ICES CM 2017/ACOM:08. 642 pp.

ICES (2017d) Report of the Working Group for the Bay of Biscay and Iberian waters Ecoregion

(WGBIE), 4-11 May 2017, Cadiz, Spain. *ICES CM 2017/ACOM,* **12,** 552 pp.

ICES (2017e) Report of the Working Group on Assessment of Demersal Stocks in the North Sea

and Skagerrak (2017), 26 April–5 May 2017. ICES HQ. ICES CM 2017/ACOM:21. 1248 pp.

ICES (2017f) Report of the Working Group on Celtic Seas Ecoregion (WGCSE), 9–18 May

2017, Copenhagen, Denmark. ICES CM 2017/ACOM:13. 1464 pp.

ICES (2019a). Workshop on Guidelines for Management Strategy Evaluations (WKGMSE2).

ICES Scientific Reports. 1:33. 162 pp.

ICES (2019b) Workshop on North Sea stocks management strategy evaluation (WKNSMSE).

ICES Scientific Reports. 1:12. 378 pp.

ICES (2021) ICES fisheries management reference points for category 1 and 2 stocks. ICES

Technical Guidelines.

Jennings, S. & Kaiser, M.J. (1998) The effects of fishing on marine ecosystems. *Advances in*

*Marine Biology*, pp. 201-352. Elsevier.

Jenouvrier, S., Desprez, M., Fay, R., Barbraud, C., Weimerskirch, H., Delord, K. & Caswell, H.

(2018) Climate change and functional traits affect population dynamics of a long-lived

seabird. *Journal of Animal Ecology,* **87,** 906-920.

Jenouvrier, S., Long, M.C., Coste, C.F., Holland, M., Gamelon, M., Yoccoz, N.G. & Sæther,

B.E. (2022) Detecting climate signals in populations across life histories. *Global Change*

*Biology,* **28,** 2236-2258.

Jensen, O.P., Branch, T.A. & Hilborn, R. (2012) Marine fisheries as ecological experiments.

*Theoretical Ecology,* **5,** 3-22.

Juan-Jordá, M.J., Murua, H., Arrizabalaga, H., Merino, G., Pacoureau, N. & Dulvy, N.K. (2022)

Seventy years of tunas, billfishes, and sharks as sentinels of global ocean health. *Science,* **378,**

eabj0211.

Koons, D.N., Iles, D.T., Schaub, M. & Caswell, H. (2016) A life-history perspective on the

demographic drivers of structured population dynamics in changing environments. *Ecology*

*Letters,* **19,** 1023-1031.

Lassen, H., Kelly, C. & Sissenwine, M. (2014) ICES advisory framework 1977–2012: from F

max to precautionary approach and beyond. *ICES Journal of Marine Science,* **71,** 166-172.

Lindegren, M., Checkley, D.M., Rouyer, T., MacCall, A.D. & Stenseth, N.C. (2013) Climate,

fishing, and fluctuations of sardine and anchovy in the California Current. *Proceedings of the*

*National Academy of Sciences,* **110,** 13672-13677.

MacKenzie, B.R., Gislason, H., Möllmann, C. & Köster, F.W. (2007) Impact of 21st century

climate change on the Baltic Sea fish community and fisheries. *Global Change Biology,* **13,**

1348-1367.

Marteinsdottir, G. & Thorarinsson, K. (1998) Improving the stock-recruitment relationship in

Icelandic cod (*Gadus morhua*) by including age diversity of spawners. *Canadian Journal of*

*Fisheries adn Aquatic Sciences,* **55,** 1372-1377.

Melnychuk, M.C., Kurota, H., Mace, P.M., Pons, M., Minto, C., Osio, G.C., Jensen, O.P., de

Moor, C.L., Parma, A.M. & Richard Little, L. (2021) Identifying management actions that

promote sustainable fisheries. *Nature Sustainability,* **4,** 440-449.

Minto, C., Myers, R.A. & Blanchard, W. (2008) Survival variability and population density in

fish populations. *Nature,* **452,** 344.

Moore, J.K., Fu, W., Primeau, F., Britten, G.L., Lindsay, K., Long, M., Doney, S.C., Mahowald,

N., Hoffman, F. & Randerson, J.T. (2018) Sustained climate warming drives declining marine

biological productivity. *Science,* **359,** 1139-1143.

Myers, R.A. (1998) When do environment–recruitment correlations work? *Reviews in Fish*

*Biology and Fisheries,* **8,** 285-305.

Neubauer, P., Jensen, O.P., Hutchings, J.A. & Baum, J.K. (2013) Resilience and recovery of

overexploited marine populations. *Science,* **340,** 347-349.

Ottersen, G., Hjermann, D.Ø. & Stenseth, N.C. (2006) Changes in spawning stock structure

strengthen the link between climate and recruitment in a heavily fished cod (*Gadus morhua*)

stock. *Fisheries Oceanography,* **15,** 230-243.

Ottersen, G., Planque, B., Belgrano, A., Post, E., Reid, P.C. & Stenseth, N.C. (2001) Ecological

effects of the North Atlantic oscillation. *Oecologia,* **128,** 1-14.

Ottersen, G. & Stenseth, N.C. (2001) Atlantic climate governs oceanographic and ecological

variability in the Barents Sea. *Limnology and Oceanography,* **46,** 1774-1780.

Pauly, D., Watson, R. & Alder, J. (2005) Global trends in world fisheries: impacts on marine

ecosystems and food security. *Philos Trans R Soc Lond B Biol Sci,* **360,** 5-12.

Peck, M.A., Huebert, K.B. & Llopiz, J.K. (2012) Intrinsic and extrinsic factors driving match–

mismatch dynamics during the early life history of marine fishes. *Advances in Ecological*

*Research*, pp. 177-302. Elsevier.

Pérez-Rodríguez, A., Umar, I., Goto, D., Howell, D., Mosqueira, I. & González-Troncoso, D.

(2022) Evaluation of harvest control rules for a group of interacting commercial stocks using

a multispecies MSE framework. *Canadian Journal of Fisheries and Aquatic Sciences,* **79,**

1302-1320.

Pershing, A.J., Alexander, M.A., Hernandez, C.M., Kerr, L.A., Le Bris, A., Mills, K.E., Nye,

J.A., Record, N.R., Scannell, H.A. & Scott, J.D. (2015) Slow adaptation in the face of rapid

warming leads to collapse of the Gulf of Maine cod fishery. *Science,* **350,** 809-812.

Pinsky, M.L. & Byler, D. (2015) Fishing, fast growth and climate variability increase the risk of

collapse. *Proceedings of the Royal Society B,* **282,** 20151053.

Pinsky, M.L., Worm, B., Fogarty, M.J., Sarmiento, J.L. & Levin, S.A. (2013) Marine taxa track

local climate velocities. *Science,* **341,** 1239-1242.

Punt, A.E. (2006) The FAO precautionary approach after almost 10 years: have we progressed

towards implementing simulation-tested feedback-control management systems for fisheries

management? *Natural Resource Modeling,* **19,** 441-464.

Rathore, S., Bindoff, N.L., Phillips, H.E. & Feng, M. (2020) Recent hemispheric asymmetry in

global ocean warming induced by climate change and internal variability. *Nature*

*Communications,* **11,** 1-8.

Roemmich, D., Church, J., Gilson, J., Monselesan, D., Sutton, P. & Wijffels, S. (2015) Unabated

planetary warming and its ocean structure since 2006. *Nature Climate Change,* **5,** 240-245.

Rogers, L.A. & Dougherty, A.B. (2019) Effects of climate and demography on reproductive

phenology of a harvested marine fish population. *Global Change Biology,* **25,** 708-720.

Rouyer, T., Ottersen, G., Durant, J.M., Hidalgo, M., Hjermann, D.Ø., Persson, J., Stige, L.C. &

Stenseth, N.C. (2011) Shifting dynamic forces in fish stock fluctuations triggered by age

truncation? *Global Change Biology,* **17,** 3046-3057.

Rouyer, T., Sadykov, A., Ohlberger, J. & Stenseth, N.C. (2012) Does increasing mortality

change the response of fish populations to environmental fluctuations? *Ecology Letters,* **15,**

658-665.

Simpson, G.L. & Singmann, H. (2022) gratia: Graceful 'ggplot'-Based Graphics and Other

Functions for GAMs Fitted Using 'mgcv'. *R package*.

Sparholt, H., Bertelsen, M. & Lassen, H. (2007) A meta-analysis of the status of ICES fish

stocks during the past half century. *ICES Journal of Marine Science,* **64,** 707-713.

Stenevik, E.K. & Sundby, S. (2007) Impacts of climate change on commercial fish stocks in

Norwegian waters. *Marine Policy,* **31,** 19-31.

Stenseth, N.C., Ottersen, G., Hurrell, J.W., Mysterud, A., Lima, M., Chan, K.S., Yoccoz, N.G. &

Ådlandsvik, B. (2003) Studying climate effects on ecology through the use of climate indices:

the North Atlantic Oscillation, El Nino Southern Oscillation and beyond. *Proceedings of the*

*Royal Society of London. Series B: Biological Sciences,* **270,** 2087-2096.

Stubben, C., Milligan, B. & Nantel, P. (2020) popbio: Construction and Analysis of Matrix

Population Models. *R package.*

Sundby, S. & Nakken, O. (2008) Spatial shifts in spawning habitats of Arcto-Norwegian cod

related to multidecadal climate oscillations and climate change. *ICES Journal of Marine*

*Science,* **65,** 953-962.

van Gemert, R. & Andersen, K.H. (2018) Challenges to fisheries advice and management due to

stock recovery. *ICES Journal of Marine Science,* **75,** 1864-1870.

van Oldenborgh, G.J., te Raa, L.A., Dijkstra, H.A. & Philip, S.Y. (2009) Frequency-or

amplitude-dependent effects of the Atlantic meridional overturning on the tropical Pacific

Ocean. *Ocean Science,* **5,** 293-301.

Wheldon, M.C. (2019) popReconstruct: Reconstruct Human Populations of the Recent Past. *R*

*package*.

White, J.W., Botsford, L.W., Hastings, A., Baskett, M.L., Kaplan, D.M. & Barnett, L.A. (2013)

Transient responses of fished populations to marine reserve establishment. *Conservation*

*Letters,* **6,** 180-191.

Wood, S. (2022) mgcv: Mixed GAM Computation Vehicle with Automatic Smoothness

Estimation. *R package*.

Wood, S. & Scheipl, F. (2020) gamm4: generalized additive mixed models using mgcv and

lme4. *R package*

Wood, S.N. (2017) *Generalized additive models: an introduction with R,* 2nd edn. Chapman and

Hall/CRC.

Worm, B., Hilborn, R., Baum, J.K., Branch, T.A., Collie, J.S., Costello, C., Fogarty, M.J.,

Fulton, E.A., Hutchings, J.A. & Jennings, S. (2009) Rebuilding global fisheries. *Science,* **325,**

578-585.

Zanna, L., Khatiwala, S., Gregory, J.M., Ison, J. & Heimbach, P. (2019) Global reconstruction of

historical ocean heat storage and transport. *Proceedings of the National Academy of Sciences,*

**116,** 1126-1131.

Zimmermann, F. & Werner, K.M. (2019) Improved management is the main driver behind

recovery of Northeast Atlantic fish stocks. *Frontiers in Ecology and the Environment,* **17,** 93-

99.

Zuur, A.F., Ieno, E.N., Walker, N.J., Saveliev, A.A. & Smith, G.M. (2009) *Mixed effects models*

*and extensions in ecology with R*. Springer.

**Tables**

**Table 1.** Stock-specific information and management measures used in the assessments of 38 fish stocks in the northeast Atlantic Ocean included in the study. Stock ID indicates a unique ID for each stock assigned by the International Council of Exploration of the Sea (ICES); period and years indicate the years and the total number of years that stock data are available; age indicates the age classes included in the assessments; $F$ range indicates the age range used to compute mean fishing mortality rates for assessment; $F_{msy}$ and $B_{pa}$ indicate the management reference points set for each stock as target fishing mortality rates that produce maximum sustainable yields and biomasses that satisfy the precautionary approach; and status indicates recovery status assigned based on spawner biomass relative to $B_{pa}$ for analysis (R: recovering and NR: non-recovering) in this study.

| common name | scientific name | ecoregion | stock ID | period | years | age | $F$ range | $F_{msy}$ | $B_{pa}$ | status | source¶ |
|---|---|---|---|---|---|---|---|---|---|---|---|
| cod | *Gadus morhua* | Norwegian–Barents Seas | cod.27.12 | 1946-2016 | 71 | 3-15+ | 5 -10 | 0.40 | 460000 | R | a |
| cod | *Gadus morhua* | North Sea | cod.27.47d20 | 1963-2016 | 54 | 1-6+ | 2-4 | 0.31 | 150000 | NR | b |
| cod | *Gadus morhua* | Baltic Sea | cod.27.22-24 | 1994-2016 | 23 | 1-5+ | 3-5 | 0.26 | 38400 | NR | c |
| cod | *Gadus morhua* | Icelandic Waters | cod-farp | 1959-2016 | 58 | 1-10+ | 3-7 | 0.23 | 29226 | NR | d |
| cod | *Gadus morhua* | Icelandic Waters | cod.27.5a | 1955-2016 | 62 | 3-14+ | 5-10 | 0.20 | 160000 | R | d |
| cod | *Gadus morhua* | Celtic Seas | cod.27.6a | 1981-2016 | 36 | 1-7+ | 2-5 | 0.17 | 22000 | NR | e |
| cod | *Gadus morhua* | Celtic Seas | cod.27.7a | 1968-2016 | 49 | 0-6+ | 2-4 | 0.31 | 8161 | NR | e |
| cod | *Gadus morhua* | Celtic Seas | cod.27.7e-k | 1971-2016 | 46 | 1-7+ | 2-5 | 0.35 | 10300 | NR | e |
| four-spot megrim | *Lepidorhombus boscii* | Bay of Biscay and Iberian Coast | ldb.27.8c9a | 1986-2017 | 31 | 1-7+ | 2-5 | 0.19 | 4600 | R | f |
| haddock | *Melanogrammus aeglefinus* | Norwegian–Barents Seas | had-arct | 1950-2016 | 67 | 3-13+ | 4-7 | 0.35 | 80000 | R | a |
| haddock | *Melanogrammus aeglefinus* | North Sea | had.27.46a20 | 1972-2016 | 45 | 0-8+ | 2-4 | 0.19 | 132000 | R | b |
| haddock | *Melanogrammus aeglefinus* | Icelandic Waters | had-faro | 1957-2016 | 60 | 1-10+ | 3-8 | 0.13 | 22843 | R | d |
| haddock | *Melanogrammus aeglefinus* | Icelandic Waters | had.27.5a | 1979-2016 | 38 | 2-11+ | 4-7 | 0.52 | 59000 | R | d |
| haddock | *Melanogrammus aeglefinus* | Celtic Seas | had-rock | 1991-2016 | 26 | 1-7+ | 2-5 | 0.20 | 10200 | R | e |
| haddock | *Melanogrammus aeglefinus* | Celtic Seas | had.27.7b-k | 1993-2016 | 24 | 0-8+ | 3-5 | 0.40 | 10000 | R | e |
| herring | *Clupea harengus* | Baltic Sea | her.27.25-2932 | 1974-2016 | 43 | 1-8+ | 3-6 | 0.22 | 600000 | R | c |
| herring | *Clupea harengus* | Baltic Sea | her.27.28 | 1977-2016 | 40 | 1-8+ | 3-7 | 0.32 | 57100 | R | c |
| megrim | *Lepidorhombus whiffiagonis and L. boscii* | Bay of Biscay and Iberian Coast | meg.27.8c9a | 1986-2016 | 31 | 1-7+ | 2-4 | 0.19 | 980 | R | f |
| plaice | *Pleuronectes platessa* | North Sea | ple.27.7d | 1980-2016 | 37 | 1-7+ | 3-6 | 0.25 | 25826 | R | b |
| plaice | *Pleuronectes platessa* | North Sea | ple.27.420 | 1957-2016 | 60 | 1-10+ | 2-6 | 0.21 | 230000 | R | b |
| plaice | *Pleuronectes platessa* | Baltic Sea | ple.27.7e | 1999-2016 | 18 | 1-5+ | 3-5 | 0.37 | 5550 | R | c |
| plaice | *Pleuronectes platessa* | Celtic Seas | ple.27.7a | 1981-2016 | 36 | 1-8+ | 3-6 | 0.16 | 5825 | R | e |
| plaice | *Pleuronectes platessa* | Celtic Seas | ple-echw | 1980-2016 | 36 | 2-10+ | 3-6 | 0.24 | 2443 | NR | e |
| saithe | *Pollachius virens* | North Sea | pok.27.3a46 | 1967-2016 | 50 | 3-9+ | 4-7 | 0.36 | 150000 | R | b |
| saithe | *Pollachius virens* | Icelandic Waters | pok.27.5b | 1961-2016 | 56 | 3-11+ | 4-8 | 0.30 | 41400 | R | d |
| saithe | *Pollachius virens* | Icelandic Waters | pok.27.5a | 1980-2016 | 37 | 3-14+ | 4-9 | 0.20 | 61000 | R | d |
| saithe | *Pollachius virens* | Norwegian–Barents Seas | pok.27.1-2 | 1960-2016 | 57 | 3-8+ | 4-7 | 0.32† | 220000 | R | a |
| sole | *Solea solea* | Baltic Sea | sol.27.20-24 | 1984-2016 | 33 | 2-6+ | 3-5 | 0.23 | 2600 | R | c |
| sole | *Solea solea* | Bay of Biscay and Iberian Coast | sol-bisc | 1984-2016 | 33 | 2-8+ | 3-6 | 0.33 | 10600 | R | f |
| sole | *Solea solea* | Celtic Seas | sol.27.7a | 1970-2016 | 47 | 2-8+ | 4-7 | 0.20 | 3100 | NR | e |
| sole | *Solea solea* | Celtic Seas | sol.27.7e | 1969-2016 | 48 | 2-12+ | 3-9 | 0.29 | 2900 | R | e |
| sole | *Solea solea* | Celtic Seas | sol.27.7fg | 1971-2016 | 46 | 1-10+ | 4-8 | 0.27 | 2380 | R | e |
| sprat | *Sprattus sprattus* | Baltic Sea | spr.27.22-32 | 1974-2016 | 43 | 1-8+ | 3-5 | 0.26 | 574000 | R | c |
| summer-spawning herring | *Clupea harengus* | Icelandic Waters | her-vasu | 1987-2016 | 30 | 3-13+ | 5-10 | 0.22 | 273000 | R | e |
| turbot | *Scophthalmus maximus* | North Sea | tur.27.3a | 1981-2016 | 36 | 1-8+ | 2-6 | 0.36* | 4163* | R | b |
| whiting | *Merlangius merlangus* | North Sea | whg.27.47d | 1990-2016 | 27 | 1-8+ | 2-7 | 0.15 | 166708 | R | b |

| whiting | *Merlangius merlangus* | Celtic Seas | whg.27.6a | 1981-2016 | 36 | 1-7+ | 2-4 | 0.23 | 44600 | NR | e |
| whiting | *Merlangius merlangus* | Celtic Seas | whg-iris | 1980-2016 | 37 | 0-6+ | 1-3 | 0.22 | 16300 | NR | e |

†$F_{msy}$ was not defined for Barents Sea Saithe in 2017 and an alternative reference point applied ($F_{mp}$) were used for analysis in the study.

*The reference points for North Sea turbot were not defined until 2018 and the stock was excluded from multilevel modeling analysis.

¶Stock information is taken from ICES Scientific Reports: (a) ICES (2017a), (b) ICES (2017e), (c) ICES (2017b), (d) ICES (2017c) (e) ICES (2017f), (f) ICES (2017d).

**Table 2**. Parameter estimates and test statistics for the generalized additive mixed effects models selected for the northeast Atlantic fish stocks with per capita recruitment success (RS$_{total}$) as a response variable (Table S2). EDF, $F$, CI, $t$, and SD indicate the effective degrees of freedom, F-statistic, confidence intervals, t-value, and standard deviation. All time (year)-dependent smoother terms for covariates are modeled as tensor product construction (using the *t2* function in the R package *mgcv*).

| model structure and parameter | smoother | | fixed effect | | | random effect | |
|---|---|---|---|---|---|---|---|
| | EDF | $F$ | estimate | 95% CI | $t$ | variance | SD |
| recovering | | | | | | | |
| RS$_{total}$ ~ $f$(Year) + $f$(Year, SA$_{mean}$) + | | | | | | | |
| $f$(Year, SB/$B_{pa}$) + $f$(Year, $\bar{F}_{lag1}$) | | | | | | | |
| $f$(Year) | 1.00 | 7.75 | | | | | |
| $f$(Year, SA$_{mean}$) | 4.49 | 3.12 | | | | | |
| $f$(Year, SB/$B_{pa}$) | 5.43 | 19.58 | | | | | |
| $f$(Year, $\bar{F}_{lag1}$) | 2.00 | 5.73 | | | | | |
| $\beta_0$ | | | 2.26 | (1.81, 2.7) | 9.97 | | |
| $\beta_{Year}$ | | | -0.15 | (-0.26, -0.03) | -2.51 | | |
| $\beta_{Year \times SAmean1}$ | | | -0.34 | (-0.51, -0.17) | -3.99 | | |
| $\beta_{Year \times SAmean2}$ | | | -0.04 | (-0.14, 0.07) | -0.71 | | |
| $\beta_{Year \times SSB/Bpa1}$ | | | 0.43 | (0.26, 0.61) | 4.83 | | |
| $\beta_{Year \times SSB/Bpa2}$ | | | -0.05 | (-0.17, 0.07) | -0.81 | | |
| $\beta_{Year \times \bar{F}lag1,1}$ | | | -0.06 | (-0.18, 0.06) | -1.00 | | |
| $\beta_{Year \times \bar{F}lag1,2}$ | | | 0.16 | (0.06, 0.26) | 3.07 | | |
| stock | | | | | | 0.20 | 0.45 |
| ecoregion/year | | | | | | 0.13 | 0.36 |
| ecoregion | | | | | | 0.06 | 0.25 |
| residual | | | | | | 0.34 | 0.58 |
| non-recovering | | | | | | | |
| RS$_{total}$ ~ $f$(Year) + $f$(Year, SA$_{mean}$) + | | | | | | | |
| $f$(Year, SSB/$B_{pa}$) + $f$(Year, OHC) | | | | | | | |
| $f$(Year) | 1.00 | 7.95 | | | | | |
| $f$(Year, SA$_{mean}$) | 4.47 | 0.58 | | | | | |
| $f$(Year, SB/$B_{pa}$) | 3.51 | 17.8 | | | | | |
| $f$(Year, OHC) | 2.19 | 187.8 | | | | | |
| $\beta_0$ | | | 2.40 | (1.58, 3.22) | 5.75 | | |
| $\beta_{Year}$ | | | -0.26 | (-0.48, -0.03) | -2.22 | | |
| $\beta_{Year \times SAmean1}$ | | | -0.18 | (-0.47, 0.11) | -1.20 | | |
| $\beta_{Year \times SAmean2}$ | | | 0.82 | (0.6, 1.05) | 7.15 | | |
| $\beta_{Year \times SSB/Bpa1}$ | | | -0.40 | (-0.63, -0.16) | -3.28 | | |
| $\beta_{Year \times SSB/Bpa2}$ | | | 0.05 | (-0.15, 0.25) | 0.46 | | |
| stock | | | | | | 0.42 | 0.64 |
| ecoregion/year | | | | | | 0.17 | 0.41 |
| ecoregion | | | | | | $1.81 \times 10^{-5}$ | $4.26 \times 10^{-3}$ |
| residual | | | | | | 0.27 | 0.52 |

**Table 3**. Parameter estimates and test statistics for the generalized additive mixed effects models selected for the northeast Atlantic fish stocks with elasticity to recruitment success ($e_{RS}$) as a response variable (Table S2). EDF, $F$, 95% CI, $t$, and SD indicate the effective degrees of freedom, F-statistic, confidence intervals, t-value, and standard deviation. All time (year)-dependent smoother terms for covariates are modeled as tensor product construction (using the *te* function in the R package *mgcv*).

| model structure and parameter | smoother | | fixed effect | | | random effect | |
|---|---|---|---|---|---|---|---|
| | EDF | $F$ | estimate | 95% CI | $t$ | variance | SD |
| **recovering** | | | | | | | |
| $e_{RS} \sim f(\text{Year}) + f(\text{Year, SA}_{\text{mean}}) + f(\text{Year, SB}/B_{\text{pa}})$ | | | | | | | |
| $+ f(\text{Year, } \bar{F}_{\text{lag1}}/F_{\text{msy}}) + f(\text{Year, AMO}_{\text{lag1}})$ | | | | | | | |
| $f(\text{Year})$ | 1.00 | 0.60 | | | | | |
| $f(\text{Year, SA}_{\text{mean}})$ | 3.51 | 1.61 | | | | | |
| $f(\text{Year, SB}/B_{\text{pa}})$ | 7.61 | 39.16 | | | | | |
| $f(\text{Year, } \bar{F}_{\text{lag1}}/F_{\text{msy}})$ | 5.06 | 3.71 | | | | | |
| $f(\text{Year, AMO}_{\text{lag1}})$ | 7.15 | 3.09 | | | | | |
| $\beta_0$ | | | -5.16 | (-5.25, -5.07) | -118.0 | | |
| stock | | | | | | 0.02 | 0.16 |
| ecoregion/year | | | | | | 0.01 | 0.11 |
| ecoregion | | | | | | 0.01 | 0.08 |
| | | | | | | | |
| **non-recovering** | | | | | | | |
| $e_{RS} \sim f(\text{Year}) + f(\text{Year, SA}_{\text{mean}}) + f(\text{Year, SB}/B_{\text{pa}})$ | | | | | | | |
| $+ f(\text{Year, } \bar{F}_{\text{lag1}}/F_{\text{msy}}) + f(\text{Year, wSST}_{\text{anom}})$ | | | | | | | |
| $f(\text{Year})$ | 1.00 | 2.10 | | | | | |
| $f(\text{Year, SA}_{\text{mean}})$ | 7.01 | 141.36 | | | | | |
| $f(\text{Year, SB}/B_{\text{pa}})$ | 6.65 | 4.88 | | | | | |
| $f(\text{Year, } \bar{F}_{\text{lag1}}/F_{\text{msy}})$ | 3.60 | 1.96 | | | | | |
| $f(\text{Year, wSST}_{\text{anom}})$ | 3.02 | 1.52 | | | | | |
| $\beta_0$ | | | -5.16 | (-5.36, -4.96) | -50.2 | | |
| stock | | | | | | 0.09 | 0.31 |
| ecoregion/year | | | | | | 0.008 | 0.088 |
| ecoregion | | | | | | $3.4 \times 10^{-7}$ | $5.8 \times 10^{-4}$ |

**Figure legends**

Figure 1. Study systems and climate conditions in the northeast Atlantic Ocean. (a) Geographic locations of the ecoregions, which are defined by the International Council of Exploration of the Sea (ICES): Norwegian and Barents Seas (62°0'N–82°0'N at 11°0'W–68°30'E), Greater North Sea (56°0'N at 2°55'E), Baltic Sea (53°35'N–65°54'N at 11°57'E–30°14'E), Faroes (60°0'N–63°0'N at 15°0'W–4°0'W), Icelandic waters (64°38'N at 18°55'W), the Bay of Biscay and the Iberian Coast (42°39'N at 8°53'W), and Celtic Seas (48°0'N–60°30'N at 18°0'W–1°21'W). (b) Regional climate indices (North Atlantic Oscillation (NAO) index, Atlantic Meridional Oscillation index, and upper (0–400m) ocean heat content (OHC) anomaly) during the study period (1946–2016). (c) Seasonal sea surface temperature anomalies in each ecoregion. The time series of temperature anomalies in each ecoregion are truncated to show only the years that the fish stock data and assessments included in the study are available.

Figure 2. Regional patterns in fishing pressure and reproductive metrics of the recovering and non-recovering northeast Atlantic fish stocks during 1946–2016. (a) Temporal trends in relative fishing pressure, relative spawner biomass, mean spawner age, and spawner age diversity. Vertical dashed lines indicate the reference year (2002) used to define the recovery status of fish stocks based on relative spawner biomass in analysis. (b) Relationships between relative fishing pressure and relative spawner biomass and age structure metrics. Solid lines and ribbons indicate regional average trends in time (a) or relative fishing pressure (b) and 95% confidence intervals estimated by generalized additive mixed effects modeling with year (a) or relative fishing pressure (b) as a fixed effect, and ecoregion, year within ecoregion, and stock as random effects. Circles indicate partial residuals estimated from the models. Relative fishing pressure and relative spawner biomass are defined as estimated fishing mortality rate and spawner biomass

relative to management reference points ($F_{msy}$ and $B_{pa}$) defined by the International Council of

Exploration of the Sea (ICES).

Figure 3. Regional trends in transient population growths of the recovering and non-recovering

northeast Atlantic fish stocks during 1946–2016. Solid lines and ribbons indicate temporal trends

and 95% confidence intervals estimated by generalized additive mixed effects modeling with

year as a fixed effect, and ecoregion, year within ecoregion, and stock as random effects. Circles

indicate partial residuals estimated from the models. Vertical dashed lines indicate the reference

year (2002) used to define the recovery status of fish stocks based on relative spawner biomass in

analysis.

Figure 4. Regional trends in and covariates of per capita recruitment success (fecundity) of the

recovering and non-recovering northeast Atlantic fish stocks during 1946–2016. (a) Temporal

trends. Solid lines and ribbons indicate temporal trends and 95% confidence intervals estimated

by generalized additive mixed effects modeling with year as a fixed effect, and ecoregion, year

within ecoregion, and stock as random effects. Red circles indicate partial residuals estimated by

the models. (b–e) Time-varying effects of spawner age structure, spawner biomass, fishing

pressure. and climate condition. Covariates selected for each type in the models are indicated in

the y-axis labels. Color gradients and contour lines indicate the partial effects of covariates

estimated (as tensor interaction terms) in the models (only the covariate types that are supported

by data are shown). Warmer colors indicate higher values, and the ranges of the partial effects

differ among the covariates and are indicated as values on the contour lines. Gray circles indicate

back-transformed input data. (f–h) Random effects estimates (±95% confidence intervals) for

stock and ecoregion for the recovering and non-recovering stocks (ecoregion-specific estimates

for the non-recovering stocks are negligible; random effect estimates for year nested within ecoregion are given in Fig. S6b).

Figure 5. Ecoregion-scale climate effects on per capita recruitment success (fecundity) of northeast Atlantic fish stocks (the Baltic Sea, the Bay of Biscay, the Celtic Seas, Faroes, Icelandic waters, and the Norwegian–Barents Seas) during 1946–2016. Time-varying effects of climate condition are estimated by generalized additive mixed effects modeling with year (and stock for the Celtic Seas) as a random effect (see Fig. S11 for complete results of the models with the other covariates, spawner age structure, spawner biomass, and fishing pressure). Climate covariates selected in the models are indicated in the y-axis labels. Color gradients and contour lines indicate the partial effects of covariates estimated (as tensor interaction terms) in the models. Warmer colors indicate higher values, and the ranges of the partial effects differ among the covariates and are indicated as values on the contour lines. Gray circles indicate back-transformed input data.

Figure 6. Regional trends in and covariates of elasticity to per capita recruitment success of the recovering and non-recovering northeast Atlantic fish stocks during 1946–2016. (a) Temporal trends. Solid lines and ribbons indicate temporal trends and 95% confidence intervals estimated by generalized additive mixed effects modeling with year as a fixed effect, and ecoregion, year within ecoregion, and stock as random effects. Circles indicate partial residuals estimated by the models. (b–e) Time-varying effects of spawner age structure, spawner biomass, fishing pressure, and climate condition. Covariates selected in the models are indicated in the y-axis labels. Color gradients and contour lines indicate the partial effects of covariates estimated (as tensor interaction terms) in the models. Warmer colors indicate higher values, and the ranges of the partial effects differ among the covariates and are indicated as values on the contour lines. Gray

circles indicate back-transformed input data. (f–h) Random effects estimates (±95% confidence intervals) for stock and ecoregion for the recovering and non-recovering stocks (ecoregion-specific parameters for the non-recovering stocks are negligible; random effect estimates of year nested within ecoregion are given in Fig. S6c).

Figure 7. Ecoregion-scale climate effects on elasticity to per capita recruitment success of northeast Atlantic fish stocks (the Baltic Sea, the Bay of Biscay, Faroes, and Icelandic waters) during 1946–2016. Time-varying effects of climate condition are estimated by generalized additive mixed effects modeling with year (and stock for the Celtic Seas) as a random effect (see Fig. S14 for complete results of the models with the other covariates, spawner age structure, spawner biomass, and fishing pressure). Climate covariates selected in the models are indicated in the y-axis labels. Color gradients and contour lines indicate the partial effects of covariates estimated (as tensor interaction terms) in the models. Warmer colors indicate higher values, and the ranges of the partial effects differ among the covariates and are indicated as values on the contour lines. Gray circles indicate back-transformed input data.