1

Distances and their visualization in studies of spatial-temporal genetic variation using

single nucleotide polymorphisms (SNPs)

Arthur Georges[1*], Luis Mijangos[1], Andrzej Kilian[2], Hardip Patel[3], Mark Aitkens[4+] and Bernd Gruber[1]

[1] Institute for Applied Ecology, University of Canberra, ACT 2601, Australia

[2] Diversity Arrays Technology, University of Canberra, ACT 2601, Australia

[3] National Centre for Indigenous Genomics, Australian National University, ACT 2601, Australia

[4] RPS Australia Asia Pacific, Unit 2A, 45 Fitzroy Street, Carrington NSW 2294, Australia

*Corresponding Author: georges@aerg.canberra.edu.au

[+]Current address – ERM, Commercial Centre, Level 1/45 Watt St, Newcastle NSW 2300, Australia

Running Title: SNPs and genetic distances

Abstract

1. Distance measures are widely used for examining genetic structure in datasets that comprise many individuals scored for a very large number of attributes. Genotype datasets composed of single nucleotide polymorphisms (SNPs) typically contain bi-allelic scores for tens of thousands if not hundreds of thousands of loci.

2. We examine the application of distance measures to SNP genotypes and sequence tag presence-absences (SilicoDArT) and use real datasets and simulated data to illustrate pitfalls in the application of genetic distances and their visualization.

3. Euclidean Distance is the metric of choice in many distance studies. However, other measures may be preferable because of their underlying models of divergence, population demographic history and linkage disequilibrium, because it is desirable to down-weight joint absences, or because of other characteristics specific to the data or analyses. Distance measures for SNP genotype data that depend on the arbitrary choice of reference and alternate alleles (e.g. Bray-Curtis distance) should not be used. Careful consideration should be given to which state is scored zero when applying binary distance measures to sequence tag presence-absences (e.g. Jaccard distance).

4. Missing values that arise in the SNP discovery process can cause displacement of affected individuals from their natural groupings and artificial inflation of confidence envelopes, leading to potential misinterpretation. Filtering on missing values then imputing those that remain avoids distortion in visual representations. Failure of a distance measure to conform to metric and Euclidean properties is important but only likely to create unacceptable outcomes in extreme cases. Lack of randomness in the selection of individuals (e.g. inclusion of sibs) and lack of independence of both individuals and loci (e.g. polymorphic haploblocks), can lead to substantial and otherwise inexplicable distortions of the visual representations and again, potential misinterpretation.

Keywords: Principal Components Analysis, Principal Coordinates Analysis, Genetic distance, Metric distance, Euclidean distance, Structural variant

Introduction

Population genetics is the study of the interplay of genetic drift, gene flow, recombination and selection, and to a lesser extent mutation, as they come to influence the contemporary

genetic composition of populations. Many measures of genetic similarity and dissimilarity

have been developed to quantify inter-individual and inter-population variation arising from

these processes. The concept of genetic distance (Sanghvi, 1953) is now a fundamental tool

in genetics (Nei & Kumar, 2000).

Application of genetic distances in studies of spatio-temporal variation is common

because substructuring of populations can arise from interesting combinations of genetic,

demographic and biogeographic processes. One area that has received substantial attention is

isolation by geographic distance (Wright, 1943) which, in a homogenous and continuous

landscape, is perhaps the simplest instance of departure from panmixia in a widespread

population. Isolation by distance is typically examined by comparing a genetic distance

matrix with geographic distance between individuals, sub-populations or sampling localities.

Fst and Chord Distance are often chosen to examine any relationship with geographic

distance (Séré et al., 2017). Such a relationship is often used as the null hypothesis against

which to examine more complicated hypotheses to explain patterns of variation across the

landscape (Manel et al., 2003; Spear et al., 2015). In particular, isolation by distance can be

complicated by variation in resistance to gene flow in complex landscapes (McRae, 2006).

Introduction of the concept of landscape resistance delivers a continuous set of circumstances

between genetic structure arising from isolation by distance in otherwise freely breeding

populations and subdivision between populations of a species where gene flow is absent, rare

or an episodic event. More novel applications of genetic distances include stratified sampling

of a core collection of specimens from a larger pool of accessions to maximally capture

genetic diversity (Rogers D: Jansen & van Hintum, 2007) and tracking the relationships of

SARS-CoV-2 variants (Jaccard D: Yin, 2020). Another area of endeavour that has benefitted

from the advances afforded distance measures is phylogeography, both in the context of

human populations (Pugach & Stoneking, 2015; Tassi et al., 2015) and animal and plant

populations (Edwards et al., 2015). Here, genetic substructuring is mapped onto the landscape

and interpreted in the context of historical (Georges et al., 2018) and contemporary (Mijangos

et al., 2022) barriers to geneflow. SNPs and distances derived from them are also useful in

population assignment (Negrini et al., 2009) and for putatively identifying migrants within

source populations (Mamoozadeh et al., 2020). These analyses rely on genetic distances and

their visualization.

Genetic distances fall into several broad classes. Some are obtained by direct

measurement, such as immunological distance (Faith, 1985) or DNA-DNA hybridization (de

Ley et al., 1970; Hirayama et al., 1996; Kirsch et al., 1990). However, most genetic distances

are calculated from character states (genotypes) arranged as a matrix of individuals (entities)

by genetic loci (attributes) Genetic distances can be further classified by whether they will be

used to infer patterns of ancestry and descent among species (phylogenetics), the structure

and relationships among populations of a species at various scales of divergence with or

without gene flow (population genetics), or relationships among individuals (e.g. kinship).

In this paper, we deal with distances defined for both individuals and populations

and their visualization in the broader context of divergence among populations and

subpopulations. The array of available measures of distance and similarity/dissimilarity is

daunting (Deza & Deza, 2009). We specifically focus our attention to analyses of single

nucleotide polymorphisms (SNPs), markers that are commonly used in studies of spatial or

temporal variation among individuals and populations of a species or closely related species.

We also consider genetic distances as they apply to complementary dominant sequence tag

presence-absence markers (SilicoDArT markers, analogous to microarray DArTs first

described by Jaccoud et al. (2001) but extracted in silico to sequences obtained from genomic

representations e.g. Ali et al., 2020; Mahboubi et al., 2020; Elshibli and Korpelainen, 2021;

Nantongo et al., 2022). Although genetic distances are used in a wide range of contexts

(Jansen & van Hintum, 2007; Libiger et al., 2009; Yin, 2020), we also restrict our attention

(not exclusively) to the application of distances in studies of spatial and temporal genetic

structure among populations where allelic profiles are governed principally by recent or

contemporary processes of drift, selection and gene flow. We do not consider distance

measures used to reveal deeper historical patterns of ancestry and descent among lineages on

independent evolutionary trajectories where the pattern of mutational events dominates

(species-level phylogenetics). Nor do we consider distance measures used to quantify

distances between SNPs loci themselves as opposed distances between individuals or

populations (Müller et al., 2005).

Our treatment of suitable genetic distance measures is predicated on the criterion that

they should agree with our commonsense notion of a distance. They should satisfy, to an

adequate level of approximation, the properties of a metric distance (Legendre & Legendre,

2012). This can be particularly important when the genetic distances are to be represented

graphically in a space defined conventionally by Cartesian coordinates and, in particular,

Principal Coordinates Analysis (PCoA) (Gower, 1966). We also discuss Principal

Components Analysis (PCA) (Pearson, 1901; Hotelling, 1933; Jolliffe, 2002) which, although

it does not take a distance matrix as input, is a closely related technique that provides a

foundational underpinning to PCoA. The two analyses are collectively referred to as

ordination. These techniques have found application in many diverse fields such as ecology,

economics, psychology, meteorology, oceanography, human health and genetics as a

descriptive and an exploratory tool to generate hypotheses for further examination (Jolliffe &

Cadima, 2016) rather than a formal statistical analysis (but see Patterson et al., 2006).

We examine some practical aspects of applying distance analyses and their

visualization. We pay particular attention to underlying assumptions of the application of

distance analyses and visualization of spatio-temporal structure using ordination, many of

which are poorly appreciated. These assumptions include the properties of various distance

measures and the importance of metric and/or Euclidean properties for visualization, the

impact of missing values and the importance of randomness in sampling and independence of

individuals (relatedness), their genotypes and the loci selected for screening.

Finally, we appreciate that PCA has recently come under criticism (Elhaik, 2022), but

argue that many of these criticisms derive from the over-interpretation of the patterns of

genetic variation revealed by PCA. We argue strongly that PCA and PCoA are exploratory

tools that draw out patterns from multivariable data by aggregating correlated influences into

a relatively few dimensions. These techniques have little to contribute to understanding of the

underlying events and processes that led to those patterns. The patterns thus are suggestive

but not demonstrative of underlying causes. Interpretations of the causes of the patterns need

to be pursued by more definitive subsequent analysis, beyond the scope of this review, to

distinguish between competing explanations for the observed patterns.

Box 1. Glossary of terms

Population. A group of individuals of the same biological species that, by virtue of

occupying a well-defined geographic provenance, have opportunity to interbreed. Changes in

the allelic profiles of a subset of individuals will come to affect the whole.

Sub-population. A subset of individuals of a population, occupying a well-defined

geographic provenance and which, because of resistance to geneflow arising from partial or

episodic isolation, have a distinct allele frequency profile. For ease of communication, we

refer to populations on the implicit understanding that the point also applies also to

subpopulations.

| 152 | **SNP.** Single Nucleotide Polymorphism. A single base pair mutation that is polymorphic. |

SNP genotype data. SNPs scored across multiple loci for each individual. Codominant markers.

SilicoDArT data. Presence or Absence of the amplified sequence tags across multiple loci for each individual. Dominant markers scored as 0 for absence and 1 for presence. Absences are typically null alleles. Hemizygous state scored as present (1).

Entity. Something with distinct and independent existence. May refer to a sample, subject, individual or population from which the genotypes are scored.

Attribute. A character that is scored for each entity, in this case the loci.

State. Refers to the value of an attribute (locus) for a given entity (individual) in the data matrix. Typically scored as 0, 1, 2 for SNP genotype data; 0, 1 for SilicoDArT data.

Call Rate. Measures the rate at which a SNP or SilicoDArT markers are successfully called (scored). They may not be called because of a mutation at one or both of the restriction enzyme recognition site or they may be missed in the sequencing because of insufficient read depth. Either way, the score is missing.

Distance. Term used loosely to encompass distance and dissimilarity measures (Gower & Legendre, 1986).

Euclidean distance. Metric distance that further can be represented faithfully in a space defined by Cartesian coordinates.

Metric distance. Distance that satisfies the formal metric properties – positive definite, symmetric, and conforms to the triangle inequality.

Dissimilarity measure. A quantitative measure of how different two individuals or populations are. Does not necessarily conform to the properties of a metric.

PCA. Principal Components Analysis, a classical ordination technique drawing upon the covariance matrix (unstandardized) or the correlation matrix (standardized) for the entities by attributes matrix.

PCoA. Principal Coordinates Analysis, a generalization of PCA that admits a wider range of distance and dissimilarity measures.

Datasets and Methods

A dataset was constructed from a SNP matrix generated for the freshwater turtles in the genus

Emydura (Georges et al., 2018), a recent radiation of Chelidae in Australasia. The dataset

includes selected populations that vary in level of divergence to encompass variation within

species and variation between closely related species. Sampling localities with evidence of

admixture between species were removed. Monomorphic loci were removed, and the data

was filtered on call rate (>95%), repeatability (>99.5%) and read depth ($5x <$ read depth $<$

$50x$). Where there was more than one SNP per sequence tag, only one was retained at

random. The resultant dataset had 18,196 SNP loci scored for 381 individuals from 7

sampling localities or populations – *Emydura victoriae* [Ord River, NT, n=15], *E. tanybaraga*

[Holroyd River, Qld, n=10], *E. subglobosa worrelli* [Daly River, NT, n=25], *E. subglobosa*

subglobosa [Fly River, PNG, n=55], *E. macquarii macquarii* [Murray Darling Basin north,

NSW/Qld, n=152], *E. macquarii krefftii* [Fitzroy River, Qld, n=39] and *E. macquarii emmotti*

[Cooper Creek, Qld, n=85]. The missing data rate was 1.7%, subsequently imputed by

nearest neighbour (Beretta & Santaniello, 2016) to yield a fully populated data matrix. The

data are a subset of those published by Georges et al. (2018) for illustrative purposes only. A

companion SilicoDArT dataset for the same individuals was also generated, filtered on read

depth (> 12x), repeatability (>99.5%), and call rate (>95%).

The above manipulations were performed in R package dartR (Gruber et al., 2018;

Mijangos et al., 2022; refer supplementary text for the R Code). Principal Components

Analysis was undertaken using the glPCA function of the R adegenet package (as

implemented in dartR, Gruber et al., 2018; Jombart, 2008; Jombart & Ahmed, 2011).

Principal Coordinates Analysis was undertaken using the pcoa function in R package ape

(Paradis & Schliep, 2019) implemented in dartR (Mijangos et al., 2022).

To exemplify the effect of missing values on SNP visualisation using PCA (Figure 7),

we performed computer simulations in dartR (Mijangos et al., 2022). Briefly, we simulated

ten populations that reproduced over 200 non-overlapping generations. Simulated populations

were placed in a linear series with low dispersal between adjacent populations (one disperser

every ten generations). Each population had 100 individuals, of which 50 individuals were

sampled at random. Genotypes were generated for 1,000 neutral loci on one chromosome.

We then randomly selected 50% of genotypes and set them as missing data. Principal

Components Analysis was undertaken using the glPCA function of the R adegenet package

(as implemented in dartR, Gruber et al., 2018; Jombart, 2008; Jombart & Ahmed, 2011). R

script and input files to replicate this analysis are provided in the supplementary text.

The data for the Australian Blue Mountains skink *Eulamprus leuraensis* (Figure 8)

were generated for 372 individuals collected from 17 swamps isolated to varying degrees in

the Blue Mountains region of New South Wales. Tail snips were collected and stored in 95%

ethanol. The tissue samples were digested with proteinase K overnight and DNA was

extracted using a NucleoMag 96 Tissue Kit (MachereyNagel, Duren, Germany) coupled with

NucleoMag SEP (Ref. 744900) to allow automated separation of high-quality DNA on a

Freedom Evo robotic liquid handler (TECAN, Miinnedorf, Switzerland). SNP data were

generated by the commercial service of Diversity Arrays Technology Pty Ltd (Canberra,

Australia) using published protocols (Georges et al., 2018; Kilian et al., 2012; Sansaloni et

al., 2011). A total of 13,496 loci were scored which reduced to 7,935 after filtering out

secondary SNPs on the same sequence tag, filtering on reproducibility (threshold 0.99) and

call rate (threshold 0.95), and removal of monomorphic loci. The resultant data is used to

demonstrate the impact of a substantial inversion on the outcomes of a PCA.

To test the effect of having closely related individuals (parents and offspring) on the

PCoA pattern we ran a simulation using dartR (Gruber et al. 2018, Mijangos et al. 2022),

where we picked up two individuals to become the parents with 2-8 offspring. We ran a

PCoA for all of the simulated cases (Figure 9 shows the case of 0 and 8 offspring). R script

and input files to replicate this analysis are provided in the supplementary text.

Figure 1. Distance between two points A and B represented in two-dimensional space can be calculated from their Cartesian coordinates using Pythagoras' rule – the square of the hypotenuse of a right-angled triangle is equal to the sum of the squares of the two adjacent sides. Each axis can be considered to represent a locus (a), with the value taken by an individual (A or B) on that axis called as x_{Ai} and x_{Bi} = 0, 1 or 2 for SNP genotype. As such, each individual is represented by a point in a multidimensional space defined by the i = 1 to L loci. Populations A and B can be similarly depicted, as in (b), with the relative frequency of the alternate allele in the population (p_{Ai} and p_{Bi}) as the value taken on each SNP axis. Representation for SNP PA data can be similarly defined.

The Concept of Distance

Standard Euclidean distance is a common-sense notion derived as an abstraction of physical distance. Applying Standard Euclidean distance to SNP data is straightforward. The horizontal axis of Figure 1a represents SNP Locus 1 and the values x_{A1} and x_{B1} represent the scores for that locus (0 or 1 or 2 for SNPs; 0 or 1 for tag presence-absence) for individuals A

257 and B respectively; the vertical axis represents SNP Locus 2 with the values x_{A2} and x_{B2}

258 similarly defined.

$$d(A, B) = \sqrt{\sum_{i=1}^{L} (x_{Ai} - x_{Bi})^2}$$

259 Standard Euclidean Distance can be similarly defined for populations,

$$D(A, B) = \sqrt{\sum_{i=1}^{L} (p_{Ai} - p_{Bi})^2}$$

where p_{Ai} and p_{Bi} are the relative frequencies of the alternate allele at locus i in populations A

and B respectively (Figure 1b).

The equations for $d(A,B)$ and $D(A,B)$ can be scaled to fall in the range [0,1] on noting

that its maximum is achieved for SNP genotype data when all x_{Ai} are 0 and all x_{Bi} are 2; for

SilicoDArT data the maximum is achieved when all x_{Ai} are 0 and all x_{Bi} are 1. Note also that

the equations for $d(A,B)$ and $D(A,B)$ are symmetric with respect to choice of which allelic

state is assigned to reference (0 for homozygous reference) and which is assigned to alternate

(2 for homozygous alternate). Interchanging the reference and alternate alleles (that is

applying transformation $x' = 2 – x$ for SNP genotype data or $x' = 1 – x$ for SilicoDArT data)

has no impact on the value of the distance between A and B.

A Mantel Test (Guillot & Rousset, 2013; Mantel, 1967) can be used to compare two

genetic distance matrices (or a genetic distance and a geographic distance matrix) provided

they are constructed for the same entities and those entities are independent (i.e. not

autocorrelated).

Generalization of the concept of Distance

Standard Euclidean distance is just one of many distance measures. The concept of distance

more generally can be distilled down to three basic properties. For a metric distance we have:

d(A,B) = 0 if and only if A = B

d(A,B) = d(B,A)

d(A,B) ≤ d(A,C) + d(B,C)

The first condition asserts that indiscernible entities are one and the same. The second

condition asserts symmetry. The last condition is referred to as the triangle inequality which

enforces the notion that the distance between two points is the shortest path between them.

From these properties we can conclude that metric distances must be non-negative.

$d(A,B) \geq 0$

In essence, metric distances are well-behaved distances. Standard Euclidean Distance, as with

all Euclidean distances, is a metric distance.

Graphically, the metric properties make complete sense for a distance (Figure 2).

Given three points defined by the distances between them, the position of each of them is

defined (Figure 2a). This is necessary (though not sufficient) if we are to represent our

distances in Cartesian coordinate space a fundamental requirement of PCoA.

Figure 2. Visual representation of the triangle inequality as used to define a metric distance. **(a)** – if

three distances between points A, B and C satisfy the metric property $d(A,B) \leq d(A,C) + d(B,C)$,

then the position of each of A, B and C in space is well defined – when the lines represented by d(A,C)

and d(B,C) are swung through an arc (blue lines), they meet to define a unique point C **(b)** – if not, and

$d(A,B) > d(A,C) + d(B,C)$, then point C is undefined – when the lines represented by d(A,C) and

d(B,C) are swung through an arc, they fail to meet and so C is undefined by the information contained

in the distances.

While the metric properties of a distance are clearly important, many measures used in

genetics are non-metric. An example of a dissimilarity measure that fails to satisfy the

symmetry condition is one defined on private alleles. Private alleles are those uniquely

possessed by one population when compared to other populations. The number of private

alleles possessed by population A compared to population B will typically be different from

the number of private alleles possessed by population B compared with A. Thus, the resultant

distances will not satisfy the second metric criterion of symmetry. Other genetic distances in

use do not satisfy the triangle inequality. For example, calculating the percentage of fixed

allelic differences between two entities (the percentage of alleles that are homozygous for the

reference allele in one entity and homozygous for the alternate allele in the other entity) as a

measure of distance satisfies the first two conditions of a metric distance, but not the triangle

inequality and so is non-metric. Nor is Nei's D (Nei, 1972) a metric distance for the same

reason (Farris, 1981), but the common alternative of Rogers' D (Rogers, 1972) is metric

(essentially equivalent to Euclidean Distance). F_{ST} is non-metric (Abisser and Rosenberg,

2020). The Bray-Curtis dissimilarity measure is non-metric but is rank-order similar to the

Jaccard distance, which is metric. And so on (refer Legendre & Legendre, 2012-- tables 7.2

and 7.3).

Genetic Distances for Individuals

Binary Data

SilicoDArT (sequence tag presence-absence) data involves scoring loci as "called" [1] or "not

called" [0]. They are called because if the two restriction enzymes (used in the accompanying

DArTSeq or ddRAD) find their mark, the corresponding sequence tags are amplified and

sequenced, and the locus is scored as 1 for that locus. If, however, there is a mutation at one

or both of the restriction enzyme sites, then the restriction enzyme(s) does not find its mark,

the corresponding sequence tag in that individual is not amplified or sequenced (null allele),

or if it is amplified from a different start site, is no longer considered homologous during pre-

processing. The individual is scored as 0 for that locus. SilicoDArT markers are commonly

used in an agricultural setting, to generate datasets that are complementary to SNP datasets

(Mahboubi et al., 2020; Nantongo et al., 2022). They have less commonly been used in the

context of population genetics (but see Elshibli and Korpelainen, 2021).

Unlike the co-dominant SNP genotype markers, SilicoDArT markers are dominant

markers. Technical artefacts such as low DNA quality or quantity, or failure of experimental

protocol resulting in shallow sequencing depth, may occasionally lead to missing data which

will be erroneously interpreted as a null allele. Typically, this is overcome by quality control

processes. Specific preprocessing options are used to separate true null alleles (sequence tag

absences) from sequence tags present in the target genome but missed by the sequencing

because of low read depth. First, the sequence tags are filtered more stringently on read depth

than for SNPs, typically with a threshold of $8x$ or more. Second, the distributions of the

sequence tag counts are examined for bimodality (clustering on either 0 or 1) and those loci

showing a continuum of counts rather than two clusters are discarded. Hemizygous loci are

scored as missing (-). Finally, the scores are tested for reproducibility using technical

replicates, providing the analyst with the opportunity to filter loci with poor reproducibility in

the SilicoDArT scores. In this way, a reliable set of dominant markers are obtained, to

complement the SNP dataset generated from the same individuals. Note that although the

SilicoDArT and SNP genotypes are derived from the same samples and representation, the

SilicoDArT markers include sequence tags that do not have a SNP, that is, sequence tags that

are invariant across all individuals.

Many binary distance measures (Deza & Deza, 2009) have been co-opted for genetic

studies using these dominant markers (e.g. Jaccard Distance, Ali et al., 2020). The calculation

of genetic distance from SilicoDArT data involves four intermediate sums. Taking two

individuals A and B, we can count the different cases,

$$N_{00} = \sum_{i=1}^{L} \begin{array}{l} 1, where\ x_{Ai} = x_{Bi} = 0 \\ 0, where\ x_{Ai} \neq x_{Bi} \end{array}$$

350 ....(1)

351 where x_{Ai} and x_{Bi} are the SilicoDArT scores for individuals A and B respectively. N_{00} is the

352 sum of loci scored 0 (absent) for both individuals;

$$N_{10} = \sum_{i=1}^{L} \begin{array}{l} 1, where\ x_{Ai} = 1 \\ 0, where\ x_{Bi} = 0 \end{array}$$

353 ....(2)

354 that is, sum loci scored 1 (present) for Individual A and 0 (absent) for Individual B;

$$N_{01} = \sum_{i=1}^{L} \begin{array}{l} 1, where\ x_{Ai} = 0 \\ 0, where\ x_{Bi} = 1 \end{array}$$

355 ....(3)

356 that is, sum of loci scored 0 (absent) for Individual A and 1 (present) for Individual B;

$$N_{11} = \sum_{i=1}^{L} \begin{array}{l} 1, where\ x_{Ai} = x_{Bi} = 1 \\ 0, where\ x_{Ai} \neq x_{Bi} \end{array}$$

357 ....(4)

358 that is, sum of loci scored 1 (present) for both individuals. These summations do not include

359 loci for which data are missing (NA) for one or both individuals.

360 The number of loci L is given by

$$L = N_{00} + N_{01} + N_{10} + N_{11}$$

....(5)

Based on the above intermediates [(1) – (5)], there are several ways to calculate a binary

distance between two individuals (Choi et al., 2010), some of which are shown in Table 1.

The two most commonly used distance measures are Standard Euclidean Distance and

Simple Matching Distance (Sokal & Michener, 1958). Both are based on the number of

mismatches between two individuals ($N_{01} + N_{10}$), expressed as a proportion of the total

number of loci considered (L), and used when there is symmetry (equivalence) in the

information carried by 0 (absence) and 1 (presence). Simple Matching Distance is simply the

Standard Euclidean distance squared and, as such, will give greater weight to larger distances

than Euclidean distance. This might be of value when looking for clusters of entities in

subsequent analyses. Simple Matching Distance is a metric distance (Orlóci, 1978:62).

Table 1. Some genetic distances commonly applied to binary data and derived from sequence tag presence or

absence, such as to SilicoDArT data. Variables are described in the text. Formulae illustrate the adjustments

made to the denominator for Jaccard and Sørensen distances rather than being presented in their conventional

algebraic form. Software packages in R for calculating distances for binary data, including those listed above

using the same terminology, include vegan (Oksanen et al., 2014) and dartR (Gruber et al., 2018; Mijangos et

al., 2022).

Name	Formula	Alternate Names	Reference
Standard Euclidean Distance	$d_E = \sqrt{\dfrac{N_{01} + N_{10}}{L}}$		
Simple Matching Distance	$d_{SM} = \dfrac{N_{01} + N_{10}}{L}$	Scaled Hamming Distance	(Sokal & Michener, 1958)
Jaccard Distance	$d_J = \dfrac{(N_{01} + N_{10})}{L - N_{00}}$	Marczewski-Steinhaus D (Marczewski & Steinhaus, 1959); Ružička D; Soergel D	(Jaccard, 1912)

Sørensen Distance	$d_S = \dfrac{(N_{01} + N_{10})}{L - N_{00} + N_{11}}$	Dice D (Dice, 1945)	Sørensen D (Sørensen, 1948)

Jaccard Distance is a variation on the Simple Matching Distance that down-weights

joint absences and so is no longer symmetric with respect to 0 and 1 scores. Note that

multiple independent mutations can result in a 0 score for a sequence tag, and absences of a

sequence tag (again 0) may arise from a failure to detect; both can contribute artificially to

the similarity between two entities. One might therefore consider recoding the data ($x' = 1 -$

x) so that 1 represents presence of a mutation at one or both of the restriction enzyme sites

(i.e. absence of the amplified tag) and 0 represents absence of such a disruptive mutation (i.e.

success in amplifying the sequence tag). Having made this simple adjustment, the Jaccard

Distance will down-weight joint absence of a disruptive mutation. The Jaccard Distance is a

metric distance (Levandowsky & Winter, 1971).Sørensen Distance adjusts the denominator

to down-weight the joint absences (0,0) and up-weight joint presences (1,1) (Table 1) and so

is a stronger response to the uncertainty surrounding joint absences that in the Jaccard

Distance. As with the Jaccard Distance, you might consider reversing the scores for absence

(0) and presence (1) to 1 and 0 respectively when dealing with SilicoDArT data. Special

attention may be required to manage missing values when applying the Sørensen and Jaccard

Distances, because adjustment of the denominator in their formulae (Table 1) can lead to a

potential systematic bias (Orlóci, 1978:62). The Sørensen Distance is not a metric distance

(Orlóci, 1978:61).

SNP Genotype Data

SNP data take on three values at a locus: 0 for homozygous reference allele; 1for

heterozygous; and 2 for homozygous alternate allele, that is, the count of the alternate allele.

Software packages PLINK (Chang et al., 2015), SNPTEST (Marchini et al., 2007) and dartR

(Gruber et al., 2018; Mijangos et al., 2022) use this scoring scheme as the default. The

alternative scoring scheme of 0 for homozygous reference allele, 1 for homozygous alternate

allele and 2 for heterozygotes is commonly used in genome-wide association studies

(GWAS), and software such as GCTA (Genome-wide Complex Trait Analysis, Yang et al.,

2011) and by Diversity Arrays Technology Pty Ltd (Canberra, Australia).

SNPs have particular characteristics that influence the choice of a distance measure.

SNPs can have more than two alleles (i.e. are multiallelic) but in practice, sites with more

than two allelic states are filtered out during the selection of markers as part of pipelines to

eliminate non-homologous sequence tags. Fortunately, multiallelic SNP sites are rarely

observed. As a result of this filtering, SNP markers are bi-allelic and have characteristics that

substantially limit options for a distance measure. For example, the values of 0 and 2 carry

equal weight because the choice of reference allele and alternate allele is arbitrary. Any

distance measure that gives differential weight to joint zeros (e.g. Bray & Curtis, 1957

Distance) will yield values that depend on the arbitrary choice of which allele is reference

and which is alternate at each locus. Such distance measures can be eliminated from options

available for SNP genotype data. Standardization or normalization across attributes (loci) is

not required because the attributes (loci) are all measured as genotypes on the same scale (0,

1 or 2); this is not the case under some circumstances in multiallelic systems. That said, the

biallelic nature of SNP markers permits the easy calculation of the maximum distance

possible, which permits scaling distances to fall conveniently in the range of [0,1] while

maintaining comparability across studies. Finally, issues that arise in a multiallelic context do

not apply in a biallelic context. For example, Rogers D (Rogers, 1972), and therefore

Standard Euclidean Distance, can yield undesirable results when two populations are both

polymorphic at a site, but share no alleles (Nei & Kumar, 2000:246). This situation does not

arise in the biallelic case, and so Standard Euclidean Distance (or equivalently Rogers D) is

often the distance of choice in SNP studies for both individuals and populations.

Standard Euclidean Distance as applied to SNP genotype data is defined in the usual

way with loci as the axes in a coordinate space, and the value on each axis is 0, 1 or 2 as

defined above. The scaling factor of ½ arises because the maximum squared distance

between two individuals at a locus is $(2-0)^2 = 4$.

$$d_E(A, B) = \frac{1}{2} \sqrt{\sum_{i=1}^{L} \frac{(x_{Ai} - x_{Bi})^2}{L}}$$

431   where x_{Ai} and x_{Bi} are the counts of the alternate allele at locus i for individual A and B

432   respectively; L is the number of loci for which both x_{Ai} and x_{Bi} are non-missing.

433   The Simple Matching Distance uses the counts of shared alleles between two

434   individuals i and j at a locus is given by

$$c_{i,j} = 0, where\ no\ alleles\ are\ shared\ [0,2]\ or[2,0]$$
$$= 1, where\ one\ allele\ is\ shared\ [0,1]\ or\ [2,1]\ or\ [1,2]$$
$$= 2, where\ both\ alleles\ are\ shared\ [0,0]\ or\ [1,1]\ or\ [2,2]$$

435 and is calculated as

$$d_{SM}(A,B) = 1 - \frac{1}{2L}\sum_{i=1}^{L} c_{ij}$$

where *L* is the number of loci non-missing for both individuals *i* and *j*. It is metric and similar

to the Allele Sharing Distance (Gao & Starmer, 2007), differing from it by a factor of 2.

Simple Matching Distance lacks a square and so does not give greater weight to larger

distances as does Standard Euclidean Distance.

Czekanowski Distance (Czekanowski, 1913) is calculated by summing the scores

across the axes

$$d_{Cz}(A,B) = \frac{1}{2L}\sum_{i=1}^{L} |x_{Ai} - x_{Bi}|$$

where x_{Ai} and x_{Bi} are the counts of the alternate allele at locus *i* for individual A and B

respectively; *L* is again the number of loci for which both x_{Ai} and x_{Bi} are non-missing.

Referred to also as the Manhattan D or the City Block D, Czekanowski Distance is a metric

distance. Czekanowski Distance is often preferred over Standard Euclidean distance when

there are extreme outliers among the data.

Software packages in R for calculating distances for SNP genotypes, including those

listed above using the same terminology, include vegan (Oksanen et al., 2014), adegenet

(Jombart, 2008), poppr (Kamvar et al., 2014), pegas (Paradis, 2010) and dartR (Gruber et al.,

2018; Mijangos et al., 2022).

Genetic Distances for Populations

Standard Euclidean Distance as applied to allele frequencies is defined in the usual way with

loci (attributes) represented by the axes in a coordinate space, and the values (states) for each

population (entity) on each axis set to the frequency of the alternate allele for the respective

locus (Table 2). An appropriate scaling factor is applied to bring the value of the distance into

the range [0,1], which renders it identical to Roger's D (Rogers, 1972). Standard Euclidean

distance is a model-free metric distance, in that its formulation makes no assumptions

regarding evolutionary processes generating the genetic distances.

Nei's standard genetic distance (Nei, 1972) is favoured by some over Standard

Euclidean Distance because of its relationship to divergence time. When populations are in

mutation-drift balance throughout the evolutionary process and all mutations result in new

alleles in accordance with the infinite-allele model, Nei's D is expected to increase in

proportion to the time after divergence between two populations (Nei, 1972). Nei's D is non-

metric.

Reynolds genetic distance (Reynolds et al., 1983) is also approximately linearly

related to divergence time in theory, but unlike Nei's Standard Genetic Distance, it is based

solely on a drift model and does not incorporate mutation. Thus, it may be more appropriate

than Nei's distance for spatial population genetics (divergence on relatively short timescales)

and in particular, representation of genetic similarity in trees (phenograms) or networks

where branch lengths need to be interpretable. A better approximation (Reynolds et al., 1983)

of the linear relationship with time is given by

$$D'_{Reynolds}(A, B) = -ln[1 - D_{Reynolds}(A, B)]$$

472   Chord Distance (Edwards & Cavalli-Sforza, 1964) (Table 2) assumes divergence

473   between populations is via drift alone, and so again may be more appropriate than Nei's D for

474   spatial population genetics. Chord Distance is based on Geodesic or Angular Distance

475   (Bhattacharyya, 1946; Edwards, 1971; Edwards & Cavalli-Sforza, 1967),

$$cos\,\alpha = \sqrt{p_{Ai}p_{Bi}} + \sqrt{q_{Ai}q_{Bi}}$$

where α is the Angular Distance. For a geometric interpretation using biallelic data, see Nei

& Kumar (2000:267). Chord Distance approximates Angular Distance by replacing the arc

distance with the length of the corresponding straight-line segment (Edwards & Cavalli-

Sforza, 1964) (Table 2). It is a metric distance that can be transformed to be approximately

Euclidean (Edwards, 1971). As with Reynold's D, Chord Distance is proportional to shallow

divergence time (for relatively small values of D) under specified assumptions (Edwards,

1971). It may be preferred over the model-free Standard Euclidean Distance where an

underlying genetic model of divergence with time is preferred.

Bray-Curtis Distance applies to abundances and down-weights joint absences in a

manner analogous to the Jaccard Distance. There is some confusion arising from the

translation of Bray-Curtis Distance (Bray & Curtis, 1957) from an ecological perspective to a

genetics perspective. When applied to SNP data, the Bray-Curtis Distance uses the abundance

of the alternate allele which renders it asymmetric with respect to the arbitrary choice of

which is reference and which is alternate allele. Bray-Curtis Distance defined in this way

should thus not be used on SNP data. Some authors have suggested that the Bray-Curtis

equation be applied separately to the counts of each allele, averaged for the locus, for

computing genetic distance between individuals or populations (Allele Frequency Difference

or AFD, Berner, 2009; Bray-Curtis/Allele Frequency Difference or BCAFD, Sherwin, 2022).

However, when this formulation is applied to either individuals or populations, it becomes

algebraically equivalent to the Czekanowski Distance (=Manhattan D). It no longer has the

properties of Bray-Curtis Distance (having been rendered metric and symmetric). To avoid

confusion, members of this class of distances should not be referred to as Bray-Curtis, AFD

or BCAFD, but rather as Czekanowski Distance or the more familiar Manhattan D, given that

naming of a distance measure should not be context dependent. The same issue has arisen in

ecology (Yoshioka, 2008). Czekanowski Distance applied to SNP data and corrected for

maximum value dependency, referred to as ^{A}A Distance (Sherwin, 2022), restores the

asymmetry with respect to choice of reference and alternate allele.

Table 2. Some genetic distances commonly applied to populations. p_{Ai} is the proportion of the alternate allele

for Locus i in population A, p_{Bi} is the proportion of the alternate allele for Locus i in population B. q_{Ai} and q_{Bi}

are similarly defined for the reference alleles. L is the number of called loci. Software packages in R for

calculating distances for binary data, including those listed above using the same terminology, include vegan

(Oksanen et al., 2014), adegenet (Jombart, 2008), poppr (Kamvar et al., 2014), pegas (Paradis, 2010) and dartR

(Gruber et al., 2018; Mijangos et al., 2022). Package heirfstat (Goudet, 2005) is the software of choice for

calculating F statistics.

Name	Formula	Reference
Standard Euclidean Distance	***Binary P/A*** $$D_E(A,B) = \sqrt{\frac{1}{L}\sum_{i=1}^{L}(p_{Bi}-p_{Ai})^2}$$ ***SNP genotypes*** $$D_E(A,B) = \frac{1}{2}\sqrt{\frac{1}{L}\sum_{i=1}^{L}(p_{Bi}-p_{Ai})^2}$$	

Czekanowscki Distance (Manhattan D)	$D_{cz}(A,B) = \dfrac{1}{2L}\sum_{i=1}^{L}(|p_{Bi} - p_{Ai}|)$	(Berner, 2009, as AFD; Czekanowski, 1913)
Nei Standard Genetic Distance	$D_{Nei}(A,B)$ $= -ln\left(\dfrac{\sum_{i=1}^{L}([p_{Ai}p_{Bi} + q_{Ai}q_{Bi}]}{\sqrt{\sum_{i=1}^{L}[(p_{Ai}^2 + q_{Ai}^2]}\sqrt{\sum_{i=1}^{L}[(p_{Bi}^2 + q_{Bi}^2]}}\right)$	(Nei, 1972)
Reynolds Distance	$D_{Reynolds}(A,B)$ $= \sqrt{\dfrac{\sum_{i=1}^{L}[(p_{Ai} - p_{Bi})^2 + (q_{Ai} - q_{Bi})^2]}{2\sum_{i=1}^{L}(1 - p_{Ai}p_{Bi} - q_{Ai}q_{Bi})}}$	(Reynolds et al., 1983)
Chord Distance	$D_{Chord}(A,B)$ $= \sqrt{1 - \dfrac{1}{L}\sum_{i=1}^{L}\sqrt{p_{Ai}p_{Bi}} + \sqrt{q_{Ai}q_{Bi}}}$	(Edwards & Cavalli-Sforza, 1964)
Wright's paired F_{ST}	Paired F_{ST} requires estimation with approaches such as that used by R package *hierfstat* [genet.dist(gl, method="WC84")]	(Wright, 1951)

Wright's *F-statistics* (Wright, 1951) describe the distribution of genetic diversity within and between populations (Holsinger & Weir, 2009). *F-statistics* are defined in terms of the proportion of heterozygotes observed (H_{obs}) and the proportion of heterozygotes expected under Hardy-Weinberg equilibrium (H_{exp}), as follows:

$$F = 1 - \frac{H_{obs}}{H_{exp}}$$

A deficit in the observed proportion of heterozygotes compared with that expected under Hardy-Weinberg equilibrium will yield a Wright's *F* that deviates from zero. If, at a single locus, observed and expected heterozygosity are in agreement, then *F=0*. If, at the extreme, no heterozygotes are observed, then *F=1*. If there is an excess of heterozygotes, F will be less than zero. When *F* is averaged across a large number of independent loci for a population, *F* will typically fall between zero and one. If two populations that differ in allelic profiles are combined (pooled), then Hardy-Weinberg equilibrium is not a sensible null expectation. Divergence of the two allele frequency profiles will manifest as non-random mating and a

departure from Hardy-Weinberg equilibrium (the "Wahlund Effect", Wahlund, 1928).

Wright's *F* applied to the pooled populations is thus informative when examining population

subdivision, because the resultant deficit in heterozygotes is an indication of genetic

structure. *F* applied in this way incorporates two components -- one arising from departure

from Hardy-Weinberg expectation within each population and the second arising from

departure from Hardy-Weinberg expectation because of structure between the two

populations or sub-populations. For this reason, Wright's pairwise F_{ST} is most commonly

used as a measure of genetic distance between two populations or sub-populations (see Nei,

1977; Nei & Kumar, 2000:236). F_{ST} *is* a measure of the heterozygosity attributable to

differences in allelic frequency profiles between populations or sub-populations, having

partitioned out the contributions from departure from Hardy-Weinberg expectation within

populations or sub-populations. Although not a genetic distance in the strict sense, F_{ST} can be

viewed a non-metric distance that varies in value between 0 and 1, such as in studies of

isolation by distance. Several methods to estimate F_{ST} have been developed and some are

complex (Excoffier, 2001; Weir & Cockerham, 1984), but there are many software packages

available to estimate F_{ST} for SNP genotype data (e.g. R package hierfstat, Goudet, 2005).

Figure 3. A heatmap used to graphically represent Standard Euclidean Distances calculated pairwise for individual in the freshwater turtle SNP dataset, labelled by sampling locality. Individuals are ranked to highlight clusters. Names of individuals are omitted for clarity. The legend gives the colour scale used to depict the distances with blues for small distances (hence the blue diagonal) and red for the maximum distance of one. The heatmap is accompanied by a neighbour-joining tree. The existence of strong structure is immediately evident. The blocks of low distances aligned along the diagonal represent the putative taxa.

549   Finally, SNP genotype data at the population level can be converted to binary data by

550 counting the shared alleles between two populations *A* and *B* at a locus

$$c_{A,B} = 0, no\ alleles\ are\ shared\ by\ the\ two\ populations\ at\ that\ locus$$
$$= 1, one\ or\ both\ alleles\ are\ shared\ at\ that\ locus$$

whereby distance measures devised for binary data can be applied. These distances are, in

effect, considering only fixed allelic differences between the two populations.

Visualization of SNP data

There are several methods for visualizing a distance matrix. Perhaps the simplest is to

represent the distance matrix as a heat map where colour is used to signify the magnitude of

distance between pairs of individuals (Figure 3). Note the white diagonal which indicates

zero differences between individuals and themselves, and that the heatmap is redundant in the

sense that the lower and upper triangles are identical. A high degree of population structuring

is usually immediately evident. For example, individuals of *Emydura* from the north basin of

the Murray-Darling catchment are very similar to each other genetically (light blue blocks of

Figure 3), but very different from populations of *Emydura* from northern Australia and New

Guinea (dark blue blocks of Figure 3). Heatmaps are a very quick way of gaining a visual

indication of the level of structuring across the dataset.

The heatmap is accompanied by a neighbour-joining tree (Saitou & Nei, 1987) which

is another common way of graphically representing genetic similarity among entities (Figure

3), not to be confused with their phylogenetic application. An alternative clustering algorithm

is UGPMA (unweighted pair group method with arithmetic mean, Sokal & Michener, 1958).

Note that to be faithfully represented in tree form, that is, without distortion, the distance

measure must be both metric and satisfy the four-point condition (for any four individuals, a

branching structure linking them has an internal branch ≥ 0) (Buneman, 1974). This is rarely

the case, as real data typically contains conflicting signals as arising, for example, from

homoplasy.

By far the most common approach to visualizing genetic distances between entities is

Principal Components Analysis (Hotelling, 1933; Jolliffe, 2002; Pearson, 1901) or PCA.

PCA is not strictly a distance analysis, but understanding this technique is fundamental to

understanding the related Principal Coordinates Analysis (PCoA) which is introduced later.

There are at least four computational approaches to calculating a PCA, differing in

computational efficiency and tractability (Wu et al., 1997), but essentially PCA takes a data

matrix (SNP genotypes or SilicoDArT), represents genetic distance between the entities

(individuals or samples or specimens) in a space defined by the L loci, centres and realigns

that space by linear transformation (rotation) to yield new orthogonal axes ordered on the

contribution of variance (represented by their eigenvalues) in their direction (defined by their

eigenvector). This process maintains the relative positions of the entities. Because the

resultant axes are ordered on the amount of information they contain, the first few axes,

preferably the first two or three, tend to contain information on any structure in the data

(signal) and later axes tend to contain only noise (Gauch, 1982). This is a powerful visual

technique for examining structure in the SNP dataset. An example of a PCA is presented in

Figure 4. Important variations include the combination of PCA with discriminant analysis

(DAPC, Jombart et al., 2010), adjustment for confounding factors (AC-PCA, Lin et al., 2016)

and accommodating sample allelic dropout from low read depth sequence tags (PCAngsd,

Meisner & Albrechtsen, 2018). The potential for sNMF (Sparse Non-negative Matrix

Factorization), a PCA-related technique useful for disentangling signal from different dyes in

microscopy (Rabinovich et al., 2003), may have application in population genetics,

particularly for identifying cases of admixture (Frichot et al., 2013). For a broader discussion

on variations of PCA not restricted to SNP data, refer to Abegaz et al. (2019). Specific R

software for undertaking classical PCA on SNP data and displaying the results graphically

includes adegenet (glPCA, Jombart, 2008) and the implementation of adegenet::glPCA in

dartR (Gruber et al., 2018; Mijangos et al., 2022), though there is a wide range of packages

for undertaking and displaying this very popular analysis.

Figure 4. Principal Components Analysis (PCA) as applied to genotype datasets for the Australasian species of *Emydura* (Chelonia: Chelidae). Plot of individuals in a two-dimensional space (**a**) defined by Principal Component 1 (horizontal axis) and Principal Component 2 (vertical axis). Together they explain 61.7% of the total variance among individuals. A scree plot (**b**) shows the contributions to variance by the first nine principal components (those that exceed the Kaiser-Guttman threshold, Guttman, 1954) of which only two each explain more than 10% of variation. Proximity of *Emydura subglobosa/worrelli* to *E. victoriae/tanybaraga* in 2D obscures their distinction, evident when Principal Component 3 is considered (**c**). The variation explained by the first two principal components can be largely set aside by considering structure within the cluster *Emydura subglobosa/worrelli/victoriae/tanybaraga* only (**d**), effectively allowing consideration of deeper dimensions. In (**e**), a PCA using SilicoDArT markers extracts essentially the same structure as the SNP genotype data (albeit flipped horizontally).

Had the data been drawn from a panmictic population (arguably the null proposition), each of the original variables would, on average, be expected to capture the same quantity of variance, and the ordination would fail (the first two axes would each represent only a small and random percentage of the total variance). When there is structure, the percentage of variation explained by the top principal components will dominate the ordination. However, the value of the percentage variation explained cannot be compared across studies as a measure of the strength of the result as if it is some sort of absolute; the percentage variance

explained by a principal component needs to be taken in the context of the average percent

variation explained by all components (Guttman, 1954). Variances explained by the top 2 or

3 principal components in SNP analyses are typically quite low, but this does not necessarily

indicate a poor outcome, as these values need to be considered in the context of the number

of axes in the overall ordination.

Note also that a PCA plot can be misleading if one chooses, for convenience, only two or

three dimensions in which to visualize the solution. For example, separation in a 2-D plot can

be accepted as real but proximity cannot because further separation can occur in deeper

dimensions each coupled with a substantial explained variance (Figure 4c). A widely used

approach to determining the number of dimensions for the final solution is to examine a scree

plot (Cattell, 1966), where the eigenvalues (proportional to variance explained) associated

with each of the new ordered dimensions are plotted (Figure 4b). It is usual to apply the

Kaiser-Guttman criterion (Guttman, 1954) whereby only those dimensions with more than

the average eigenvalue are considered, or to apply a related but less conservative approach to

take into account sampling variability (Jolliffe, 1972). Even so, this may result in many

informative dimensions. One must decide how much information to discard (say, for

example, keeping only those components that explain more than 10% of total variation) or

adopt, as a threshold, a visually evident sudden drop in the percentage variation explained,

commonly referred to as an elbow. The Kaiser-Guttman criterion is used by dartR to

determine how many axes to report in the accompanying scree plot.

More formal techniques (Cangelosi & Goriely, 2007; Jackson, 1993; Peres-Neto et al.,

2005) include the broken-stick approach of Macarthur (1957), which provides a good

combination of simplicity of calculation and evaluation of suitable dimensionality (Jackson,

1993). The broken-stick model retains components that explain more variance than would be

expected by randomly dividing the variance into equal parts. Another related approach is to

observe that the eigenvalues of lower "noise" dimensions are likely to decline geometrically,

a trend that can be linearized by a log transformation. Informative dimensions are those for

which variance explained greatly exceeds that predicted by this linear trend line (more

strictly, exert disproportionate leverage on the regression). A more recent approach assesses

the statistical significance of the variation explained by each Principal Component (Patterson

et al., 2006). Under specified assumptions, the sampling distribution of the ordered

eigenvalues, under the null hypothesis of no structure in the data, follows Tracy-Widom

distribution (Tracy & Widom, 1993). Thus, it is possible to assign a probability to an

observed eigenvalue and retain for consideration only those principal components that are

statistically significant (Patterson et al., 2006). The technique is sufficiently robust to

violations of its underlying assumptions (e.g. normality) to be applicable to large genetic

biallelic data arrays. Application of these more formal techniques require custom R scripts to

be prepared by the analyst. Whatever method is used, delineating the informative axes from

those representing noise is a personal choice which guides the quantity of signal (as opposed

to noise) that is to be discarded in selecting a final visual representation.

Generalization of PCA

Principal Co-ordinates Analysis (PCoA, Gower, 1966) is an alternative visualization

technique that represents a distance matrix in a Euclidean space defined by an ordered set of

orthogonal axes, as does PCA. Important variations include adjustment for confounding

factors (AC-PCoA, Wang et al., 2022) and application of iterative procedures to best match

measured distances against distances in the visual representation for a specified number of

dimensions (Belbin, 1991; Kruskal, 1964; Shepard, 1962; Torgerson, 1952). R Software for

generating the input distance matrix, then undertaking and displaying graphically a PCoA is

provided by ape (function pcoa, Paradis & Schliep, 2019) and the implementation of

ape::pcoa in dartR (Gruber et al., 2018; Mijangos et al., 2022).

671   The primary difference between PCA and PCoA is that PCA works with the

covariance (or correlation) matrix derived from the original data whereas PCoA works with

any distance matrix. Because the mathematics of PCA moves forward from the covariance

(or correlation) matrix, the insight attributed to John Gower (1966) was to substitute any

distance matrix for the covariance (or correlation) matrix at this point in the analysis,

following a simple transformation. This yields an ordered representation of those distances,

metric or otherwise, in multivariable space akin to PCA, greatly expanding the range of

application of ordination. In PCA, the *N* individuals are represented in a space of *L*

dimensions defined by the loci whereas in PCoA, the individuals are represented in an *N-1*

space with coordinate axes based solely on their pairwise distances.

681   Choosing the number of dimensions to display in visualizing a PCoA is similar to

PCA. Missing values are less disruptive for PCoA than classical PCA because they are

accommodated in pairwise fashion rather than globally, but they nevertheless require

consideration. Missing values result in variation in the precision of estimates of allele

frequencies across loci and can break the Euclidean properties of a sample distance matrix

even when the chosen metric is Euclidean in theory. Other considerations arise in PCoA from

using distance matrices that are non-Euclidean. If a Euclidean distance matrix is used in

PCoA then, in the absence of missing values, the distances in the input matrix are represented

faithfully in the full ordinated space, that is, without distortion (Figure 5c). The same cannot

be said of metric distances in general, as the metric properties ensure that the individuals can

be represented in an ordinated space, but not in a rigid linear space (Figure 5a, after Gower,

1982) – some curvature of the space may be required to satisfy the triangle inequality (Figure

5b), which will potentially compromise ability to faithfully represent the distances between

entities in a space defined by Cartesian coordinates.

Figure 5. Metricity is not sufficient to represent distances in a rigid space defined by Cartesian coordinates; the distances must be Euclidean. **(a)** – distances between four individuals that satisfy the metric properties can nevertheless not be represented in a linear space defined by Cartesian coordinates because the position of individual D is not defined by the interindividual distances (after Gower, 1982). **(b)** – this distortion can be resolved by allowing non-linear links (geodesics) to represent distances between individuals as, in the case of four points shown, the edges of an irregular Reuleaux tetrahedron in three dimensions. **(c)** – in contrast, Euclidean distances between four individuals can always be represented by linear segments without distortion, as edges of an irregular conventional tetrahedron in three dimensions.

Distortion arising from using non-Euclidean distances manifests as displacement of

the points in the visualization, so that the distances among them no longer fundamentally

represent the input values; some eigenvalues will be negative (representing imaginary

eigenvectors) (Gower & Legendre, 1986). However, the level of distortion is likely to be of

concern only if the absolute magnitude of the largest negative eigenvalue is greater than that

of any of the dimensions chosen for the reduced representation (Cailliez & Pages, 1976). A

few small negative eigenvalues do not detract much from the visual representation if only a

few of the highest dimensions are retained in the final solution (Sibson, 1979).

Negative eigenvalues complicate calculation of the variance contributions. In

particular, one can no longer calculate the percentage variation explained by a PCoA axis by

expressing the value of its eigenvalue over the total sum of the eigenvalues. A correction is

necessary (Legendre & Legendre, 2012:506).

$$\% \; explained \; = \frac{e_i + k}{\sum_{i=1}^{N} e_i + (N-1)k}$$

where e_i is the eigenvalue for PCoA axis i, N is the number of entities, and k is the absolute

value of the largest negative eigenvalue.

If negative eigenvalues are considered problematic for the reduced space

representation, a transformation can render them all positive and the distance matrix

Euclidean. Common transformations put to this purpose are the square root (Legendre &

Legendre, 2012:501), Cailliez transformation (Cailliez, 1983; Gower & Legendre, 1986) and

the Lingoes transformation (Gower & Legendre, 1986; Lingoes, 1971).

Square root $D'(A, B) = \sqrt{D(A, B)}$

Cailliez $D'(A, B) = D(A, B) + c$ for all $A \neq B$

Lingoes $D'(A, B) = \sqrt{D(A, B) + c}$ for all $A \neq B$

The value of c is chosen to be the smallest value required to convert the most extreme

negative eigenvalue to zero.

Finally, the outcome of a PCoA with an input matrix comprised of Standard

Euclidean Distances calculated from individual genotypes is identical to the outcome of a

PCA (Cox & Cox, 2001:43-44). The interchangeability of the two, PCA and PCoA, in this

restricted case leads to confusion on the distinction between the two analyses.

Notwithstanding that the final arrangement of entities in the ordinated space is the same for

PCA and PCoA using Euclidean Distance on the same data, the percentage variation

explained by each axis will differ between the two approaches. This is because PCA

calculates the percentage variation explained by an axis as the percentage its eigenvalue

represents of the total of all eigenvalues. The eigenvalues are calculated from the covariances

generated from the input data matrix. PCoA in contrast will calculate the percentage variation

explained by an axis as the sums of squares of the distance between the entities in the

direction of the axis expressed as a percentage of the total sums of squares of the distances in

the input distance matrix. The two will differ even when PCA and PCoA using Euclidean

Distance are applied to the same data. Hence, it is not possible to simply compare the

performance of a PCA and a PCoA based on examination of the values of percentage

variation explained.

Missing values

SNP genotype datasets typically have substantial numbers of missing values. Missing data

can arise because the read depth is insufficient to detect SNP-containing sequence tags

consistently, or because mutations at one or both of the restriction enzyme recognition sites in

some individuals result in allelic drop-out (null alleles). Because those that arise from

mutation are inherited, they are subject to genetic drift (and potentially local selection)

differentially within each sampled population, and so are not randomly distributed across the

entire dataset. Indeed, in some populations the mutation(s) leading to missing values may

become fixed. Filtering on call rate with a threshold applied globally will potentially admit

high frequencies of missing data at particular loci for single populations, which has

consequences as outlined below. In SilicoDArT data, the null alleles are themselves the data,

scored as presence or absence of the amplified tag. In this context, the marker data extraction

pipeline has to apply adequate quality control of read depth, because hemizygous individuals

are scored as present (1).

Missing data are problematic. Techniques like classical PCA that access the raw data

matrix cannot accommodate missing data. PCoA, which accesses a distance matrix, is less

affected by missing values in the raw data (Rohlf, 1972) because they are eliminated in pair-

wise fashion during construction of the distance matrix, not globally. PCoA is nevertheless

affected by missing values in ways that are not immediately transparent – by potential

breakage of metric or Euclidean properties, and by disregarding variable precision of

estimates of population allele frequencies across loci. The trade-off is one of balancing data

loss with bringing distortion of the visual representation within acceptable levels.

Figure 6. Principal Components Analysis (PCA) applied to two
populations where five individuals in one population (*Emydura*
worrelli) have had percentages of SNP loci ranging from 10% to
100% set to missing. **(a)** – the impact of this on PCA where missing
data are filled with the average global allele frequencies (mean-
imputed missing data) is clear, and subject to misinterpretation as
hybridization or various levels of introgression. **(b)** – the issue can be
resolved by local imputation, in this case by nearest-neighbour
imputation.

Because classical PCA will not accept missing values, when a locus is not scored for a

particular individual, either the data for the entire locus must be deleted or the data for the

entire individual must be deleted. This is clearly very wasteful of information, and

unacceptable loss when working with SNP genotype data. The degree of data loss can be

controlled to an extent by pre-filtering on call rate by locus (say requiring a call rate > 95%)

or by individual (say requiring a call rate > 80%), noting that a missing value rate of <10%

causes only modest overall distortion using implicit mean-imputed missing data (Figure 6a).

Numerous ways for handling missing data in PCA have been suggested (Dray & Josse,

2015), but the most common method is to replace a missing value at a locus with the mean of

the allele frequencies for that locus calculated across the individuals for which it is non-

missing (mean-imputed missing data). Mean-imputation of missing data can lead to the

individuals with a high proportion of missing data being drawn out of their natural grouping

and toward the origin, leading to potential misinterpretation (Yi & Latch, 2021). For

example, if individuals in the PCA aggregate into natural clusters, perhaps representing

geographic isolates, and these clusters are on either side of the origin, then an individual with

a high frequency of missing values will be drawn out of its cluster when the missing values

are replaced by global average allele frequencies (Figure 6). Its location intermediate to the

two clusters might be falsely interpreted as a case of admixture. Individuals with missing

data corrected by mean-imputation will also distort confidence envelopes as applied to

clusters with consequences for interpretation, particularly in studies of population assignment

(Figure 7).

Figure 7. A PCA plot of simulated data showing aggregations and their associated 95% confidence

limits for (a) data with no missing values and (b) data with 50% missing values. Note the

inflationary effect missing values have on the spread of the points in each aggregation, manifested as

inflation of the 95% confidence limits, an artifact. This is of particular relevance to studies of

population assignment.

Several better ways than global mean-imputation for handling missing values in SNP are

appropriate:

1.　Replace missing values for an individual with the mean observed allele frequency for the

population from which the individual was drawn. In this way, the individual is displaced

toward the centroid of the population from which it was drawn, not the origin of the

PCA.

2. Replace missing values for an individual with the expected value for the population to which the individual belongs, based on the assumption of Hardy-Weinberg equilibrium. The individual will again be displaced toward the centroid of the population from which it was drawn, not the origin of the PCA.

3. Replace missing values for an individual by the value at the same locus from its nearest neighbour (the individual closest to the focal individual based on Standard Euclidean genetic distance) (Beretta & Santaniello, 2016) (Figure 6). The focal individual will be drawn toward its nearest neighbour, typically an individual within the same population. This method has the advantage of filling missing values even where all individuals collected from a single location are missing for a given locus.

4. Generate a Euclidean distance matrix with pairwise deletion of missing values (rather than global deletion required of PCA) and apply PCoA to generate the ordination. This relies on the broad equivalence of PCA and PCoA using Euclidean distance (Cox & Cox, 2001:43-44).

For SilicoDArT, because missing values are typically associated with hemizygous individuals, imputation can be achieved by recoding the missing values as 1.

Linkage

Distance analyses usually assume that each locus contributes independent information to the overall distance value (i.e. they segregate independently). With the large genotype arrays typical of SNP datasets, some SNP loci are likely to be linked (Waples et al., 2022). In extreme cases, linkage disequilibrium can seriously distort the genetic structure, confounding interpretation. Large blocks of sequence with limited haplotype variation (presumably a result of limited recombination) have been observed in humans (Daly et al., 2001; Patil et al., 2001) and plants (Battlay et al., 2022) and their impact modelled in simulations (Lotterhos, 2019). If many markers have been sequenced in such a haploblock, they will be tightly linked, and support for any population structure that they represent will be proportionately inflated. This will be evident in a PCA or PCoA as artifactual structure that will potentially defy explanation when working with organisms with little genomic information. An example of such artifactual structure in a PCA plot generated from linked SNP markers is provided by polymorphism in a large inversion in human chromosome 8 (Amos & Ma, 2012; Battlay et al., 2022). SNP markers associated with this inversion generate a coordinated signal which

manifests as a three-group pattern, one corresponding to inverted homozygotes, one

corresponding to heterozygotes and one corresponding to non-inverted homozygotes (refer

also to the simulations of Lotterhos, 2019). Such an explanation was invoked to explain the

disaggregation of individuals of the Australian dragon *Pogona vitticeps* into two mirrored

clusters that could not be explained by location of capture, sex effects or batch effects (Wild

et al., 2022). The signature three-group pattern characteristic of a large polymorphic

inversion was evident also in a SNP study of genetic variation across swamps occupied by

the Australian Blue Mountains Water Skink *Eulamprus leuraensis* (Figure 8).

It is wise as a data quality step to prune groups of SNPs with high linkage

disequilibrium (LD). In extreme cases of linkage arising from SNPs residing on a shared,

non-recombining block in the genome (Figure 8) can be resolved by identifying loci that are

strongly correlated with the Principal Component that separates out the artifactual groupings

and removing those loci from the analysis (Wild et al., 2022) or by SNP pruning (Lotterhos,

2019) albeit with substantial loss of data. The approach of Wild et al. (2022) can also be used

to remove batch effects (Lou and Therkildsen, 2022) when they dominate the PCA

visualization as an alternative to more intrusive approaches such as removing all private

alleles (Lou and Therkildsen, 2022). Batch effects are best managed appropriate quality

control at the time of generating the SNPs and SilicoDArT data. It is important also to ensure

that SNP and SilicoDArT scores are generated together and the analysis rerun for the

combined sequences rather than attempting to combine datasets posthoc.

Figure 8. An ordination of SNP genotypes for the Australian Blue

Mountains Water Skink *Eulamprus leuraensis* collected from 17 locations

with varying levels of isolation. The ordination shows the three-group

structure characteristic of a large autosomal inversion (Amos & Ma, 2012).

For sake of illustration, we have assumed that the inversion is the least

frequent of the two polymorphic states. This interpretation remains

speculative until sufficient genomic information is available for this

species to demonstrate the existence of the inversion and to associate the

SNP loci strongly correlated with PCA 2 to that inversion.

Relatedness among individuals

Representing distance between populations on the basis of samples typically assumes that the

individuals are sampled at random and their genotypes are independent. Systematic errors can

arise if some sampled individuals are more closely related than are individuals selected at

random from their population (Östergren et al., 2020); these individuals can be expected to

separate out from the main body of individuals in a PCA or PCoA (Figure 9a). This can occur

if, for example, individuals are selected from a single school of fish that may be more closely

related than individuals chosen at random. The effect is most pronounced if parents and their

offspring (siblings) are among the sampled individuals (Figure 9a). Care should also be

exercised when interpreting clinal structure if the unit of dispersal comprises groups of

related individuals (Fix, 1997). These closely related individuals should be identified and all

but one removed from the analysis, with a close eye to the consequences of overzealous

removal of related individuals   (Waples and Anderson, 2017).

Figure 9. A series of sub-populations of *Emydura macquarii* from the northern basin of the Murray-Darling drainage (a) in which one subpopulation (central) has two individuals identified to be manipulated artificially into a parent-offspring relationship and eight individuals into a parent-offspring relationship; (b) The ordination after the addition of the modified individuals. If such closely related individuals go undetected, and are retained, the spatio-temporal distance relationships between populations can be subject to misinterpretation.

Discussion

Species and populations usually do not constitute a single panmictic unit where individuals breed at random over their entire range. Population subdivisions typically exist, which may be hierarchically arranged if they reflect patterns of ancestry and descent, or not if they are each on independent random walks, perhaps periodically reset by episodic gene flow (Georges et al., 2018). It follows that patterns evident in distances between entities, be they individuals, aggregations of individuals, populations or taxa, will often be highly structured. The challenge for SNP analyses arises from dealing with an incredible volume of data, certainly in the thousands of markers but often in the hundreds of thousands or even millions of markers, each with its own contribution to overall variation. Summarising the data as a distance matrix and visualizing the important structure using ordination is a first step in examining structure and hypothesising on its potential causes.

Many distance measures are available for different types of genetic data (Kosman & Leonard, 2005; Libiger et al., 2009). Fortunately, characteristics of SNP data and considering contemporary rather than historical processes governing variation among populations, dramatically reduce the options from which to select an appropriate distance. SNP markers are biallelic in practice, so distinct distance measures in a multiallelic context can become the same in a biallelic context. Additionally, symmetry considerations in the arbitrary choice of reference and alternate allele eliminate use of distance measures that treat scores of homozygous reference (0) as fundamentally different from scores of homozygous alternate (2). Because genetic distances are likely to be interpreted in the context of everyday notions of a distance measure, even if subconsciously, distances that satisfy the basic metric properties are preferable over those that do not. The decision on which distance measure to use then comes down to one of context and whether or not underlying evolutionary models of mutation and drift are considered important.

Genetic structure within and among populations is commonly explored using ordination which is inherently reliant on the concept of genetic distance. Strictly, distances

with Euclidean properties are required to faithfully represent those distances in a space

defined by Cartesian coordinates using PCA and PCoA. Distances with metric properties

yield adequate representations in most cases (e.g. Jaccard D, Manhattan D, Chord D). Non-

metric distances should in principle be avoided when the goal is to undertake a distance

analysis and ordination, notwithstanding their value in a broader context (e.g. Fst). However,

methods for detecting distortion of a visual representation arising from gross departures in

Euclidean and non-metric properties (negative Eigenvalues) are available as are means for

redressing the deficiency. In practice, the distortion arising from selecting only the top two or

three axes for the visualization generates more distortion than does a relatively few negative

eigenvalues. One can thus be somewhat relaxed about departures from Euclidean or metric

properties of a distance if there are compelling reasons for choosing a non-metric distance

measure in the context of the study, one's philosophical position on the need for an

underlying evolutionary model and questions to be explored.

PCA has recently come under criticism (Elhaik, 2022). The deficiencies of PCA

identified by Elhaik are real, but we argue that they derive from over-interpretation of PCA to

address questions that are beyond its remit, and because of misunderstanding of what

ordination delivers as its ultimate solution. PCA is essentially an exploratory tool that draws

out patterns from multivariable data by aggregating correlated influences into a relatively few

dimensions. PCA is about extracting patterns. It has little to contribute to understanding of

the processes that delivered those patterns however tempting it might be to use the

visualizations to make inferences about underlying events and processes (McVean, 2009).

The patterns thus are suggestive but not demonstrative of underlying causes. Interpretations

of the causes of the patterns need to be pursued by more definitive subsequent analysis to

distinguish between competing explanations for the observed patterns (Novembre and

Stephens, 2008). For example, individuals that fall in an intermediate position between two

well defined groupings in a bivariate plot of PCA1 and 2 may be an indication of admixture.

Certainly, that is the position in the ordination plot that admixed individuals will fall, but the

reverse inference is not true. The intermediary position may not be sustained on examination

of deeper dimensions or may have arisen from other causes. Hybridization and introgression

are complex processes that cannot be simply inferred from an examination of the patterns

evident in a PCA (Gompert and Buerkle, 2016). Definitive demonstration of admixture

requires additional analyses (refer Figure 4 of Georges et al., 2018), such as the frequency of

heterozygotes at loci fixed and different for the two major groupings, or use of statistical

packages such as NewHybrids (Anderson & Thompson, 2002). Similarly, PCA may be

undertaken as a prelude to genome-wide association studies (GWAS) to identify genes under

selection (Xu et al., 2022). Serious difficulties arise in terms of defensible and reproducible

conclusions when interpretation is restricted to PCA without targeted and appropriate

analyses to follow the indications from the PCA. One cannot reliably use two or three-

dimensional representations in PCA alone to decide ancestry or demographic history, to

reliably identify and exclude outliers, to infer kinship, to identify ancestral clines in the data,

to detect genomic signatures of natural selection, or to identify cases of convergent evolution

(Elhaik, 2022 and references cited therein). Misleading artifacts can emerge if one attempts to

interpolate a surface between observed sample points in the ordinated space, artifacts such as

sinusoidal trends or 'waves' that can lead to misinterpretation of migration patterns

(Novembre and Stephens, 2008). PCA and PCoA can provide only suggestive indications

useful for formulating hypotheses for further rigorous examination. Therein lies the value of

ordination.

A second criticism of PCA is that the visual representation of distances between

individuals is sensitive to sample size (Elhaik, 2022). Small sample sizes will have a greater

associated error in their representation of the population from which they are drawn, and this

will be reflected in the outcome of the ordination. But the sample size dependency identified

by Elhaik arises because decreasing the sample size of one structural grouping (population,

aggregation of populations, taxon) will reduce its contribution to overall variance and reorient

the individuals in the space defined by the informative dimensions of the ordination. This

leads to potentially dramatic changes in their projection on to the top two dimensions. Thus

sample size effect can have major consequences for interpretation if the analyst's focus is

restricted to two or three-dimensional PCA plots.

The apparent lack of reproducibility identified by Elhaik (2022) arises from restricting

attention to only the top two (or three) of informative dimensions. Misinerpretation arising

from this effect can be overcome by appropriate examination of the ordination, including all

substantial informative dimensions.

Arguably, missing values, and how they are managed, is the issue with the most

significant consequences for genetic distances and their visualisation. Classical PCA requires

a complete input dataset, and adequate filtering missing values requires removal of loci or

individuals from the analysis to create the input matrix free of missing values. For SNP data,

which typically have high frequencies of missing scores, the consequent data loss is

unacceptable. To overcome unacceptable loss of data, the missing values are typically infilled

using the global average for the locus concerned. We have shown that this can result in

misleading displacement of individuals with high rates of missing data away from their

natural groupings and toward the global centroid, and inflation of confidence envelopes for

aggregations of individuals with the risk of misinterpretation, particularly in analyses such as

population assignment. We have reviewed the approaches to deal with missing values that

avoid these distortions, the most effective of which appears to be replacement of a missing

value with that of the nearest neighbour (Beretta & Santaniello, 2016). Even replacement

with a random value is preferable to replacement with the global average allele frequency

(the most common default approach) because random values simply add noise to the data

which is driven to lower dimensions in any ordination.

Other concerns are linkage, which is difficult to detect and filter when sample sizes

from collection localities are small and cannot be pooled because genetic structure among

collection localities contributes artificially to estimates of linkage disequilibrium. In gross

cases of linkage, such as those arising from a polymorphic chromosomal inversion (Figure 8),

SNPs associated with the inversion can be identified and removed by screening out loci that

load high with the PCA axis most affected by the inversion (Wild et al., 2022) or by SNP

thinning (Lotterhos, 2019) albeit with substantial loss of data.

In summary, we recommend selection of a distance measure for SNP genotype data

that does not give differing outcomes depending on the arbitrary choice of reference and

alternate alleles in the case of SNP data, and careful consideration of which state should be

considered as zero when applying binary distance measures to SilicoDArT data. As a broad

recommendation on choice of a distance measure, Standard Euclidean Distance is the default

choice of a genetic distance for SNP and SilocoDArT data. It is free of assumptions about the

processes that generate the variation and, in the biallelic case, does not suffer the

unacceptable distortions that arise in multiallelic data when two populations are polymorphic

at a site, but share no alleles (Nei & Kumar, 2000:246). Standard Euclidean Distance can be

faithfully represented in ordinated Cartesian space. Other metric distances may be chosen in

order to down weight joint absences in the case of SilicoDArT data (e.g. Jaccard Distance) or

because of the importance of a particular model of progressive divergence with time (e.g.

Nei's D, Reynold's D, Chord D) for SNP genotype data. For example, Chord Distance is

better correlated with time since divergence and is a metric distance, so may be a better

choice than Fst /(1- Fst) (a non-metric distance) when analysing isolation by distance (Séré et

al., 2017). Diagnostic examination of eigenvalues should be undertaken when a non-metric

distance has been selected, or if the analysis is to include substantial missing values which

breaks metricity. Action should be taken if the sum of negative eigenvalues is substantial to

avoid distortion in the final visual representation. We strongly recommend filtering heavily

on missing values, then imputing those that remain to create a full matrix prior to undertaking

a distance analysis. Failure to do so can substantially and artificially inflate confidence

envelopes for populations or aggregations of populations and lead to other distortions that can

lead to misinterpretation. Screening for closely related individuals (parent-offspring or sibling

relationships) is also important, and the impact of polymorphic haploblocks in the genomes

of target species or populations occasionally emerges as a challenge for studies of species

with limited genomic information.

Conflict of Interest

Author AK is the Managing Director of Diversity Arrays Technology Pty Ltd (Canberra) and

the originator of the DArTSeq pipelines and associated data formats (including SilicoDArT).

The other authors have no conflicts of interest to declare.

Data Availability

Real datasets and the scripts used to generate simulated datasets used in this paper are

available on Dryad (https://doi.org/10.5061/dryad.4b8gthtkn).

Author Contributions

All authors contributed to the development of ideas presented in this manuscript. AG led the

writing, LM and BG undertook the simulations, HP provided input on human studies, MA

provided access to the skink data and provided associated context.

Acknowledgements

We would like to thank Bill Sherwin, Fred Allendorf, Evsey Kosman and anonymous

reviewers for comments on an earlier draft of this manuscript, and for assisting with setting

the scope and direction of the manuscript. We would like also to thank Springvale Coal Pty

Ltd for allowing the publication of genetic data obtained for the Blue Mountains Water Skink

Research and Management Program coordinated by RPS Australia East Pty Ltd for the

Springvale Extension Project. This work was funded in part by contributions from The

Cooperative Research Network for Murray-Darling Basin Futures, Australian Department of Industry, Innovation, Science, Research and Tertiary Education (DIISRTE). Grant Code: MDB-CRN (2012-2014); ARC Linkage Program: Drivers of fine scale genetic spatial structuring in aquatic organisms, Australian Research Council LP140100521 (2014-2017); and the ACT Priority Investment Program: EcoKDDart Future Proofing Biodiversity and Biosecurity with a novel integrative Big Data Analysis Hub, ACT Department of Territories and Local Government, Grant 638238 (2020-2023); via their contributions to the development of dartR.

References

Abegaz, F., Chaichoompu, K., Génin, E., Fardo, D. W., König, I. R., Mahachie John, J. M., & Van Steen, K. (2019). Principals about principal components in statistical genetics. Briefings in Bioinformatics, 20, 2200–2216. https://doi.org/10.1093/bib/bby081

Ali, F., Nadeem, M.A., Barut, M., Habyarimana, E., Chaudhary, H.J., Khalil, I.H., Alsaleh, A., Hatipoğlu, R., Karaköy, T., Kurt, C., Aasim, M., Sameeullah, M., Ludidi, N., Yang, S.H., Chung, G. & Baloch, F.S. (2020). Genetic diversity, population structure and marker-trait association for 100-seed weight in international safflower panel using SilicoDArT marker information. *Plants* 9(5):652. https://doi.org/10.3390/plants9050652

Amos, J., & Ma, C. I. (2012). Investigation of inversion polymorphisms in the human genome using Principal Components Analysis. *PLoS One*, *7*, e40224. https://doi.org/10.1371/journal.pone.0040224

Anderson, E., & Thompson, E. (2002). A model-based method for identifying species hybrids using multilocus genetic data. *Genetics*, *160*, 1217–1229. https://doi.org/10.1093/genetics/160.3.1217

Arbisser, I.M., & Rosenberg, N.A. (2020). Fst and the triangle inequality for biallelic markers. *Theoretical Population Biology* 2020 133:117-129. http:/doi.org/10.1016/j.tpb.2019.05.003.

Battlay, P., Wilson, J., Bieker, V. C., Lee, C., Prapas, D., Petersen, B., Craig, S., van Boheemen, L., Scalone, R., de Silva, N. P., Sharma, A., Konstantinović, B., Nurkowski, K. A., Rieseberg, L., Connallon, T., Martin, M. D., & Hodgins, K. A. (2022). Large haploblocks underlie rapid adaptation in an invasive weed. *BioRxiv*, https://doi.org/10.1101/2022.03.02.482376.

Belbin, L. (1991). Semi-strong hybrid scaling, a new ordination algorithm. *Journal of*

Vegetation Science, *2*, 491–496. https://doi.org/10.2307/3236031

Beretta, L., & Santaniello, A. (2016). Nearest neighbor imputation algorithms: a critical

evaluation. *BMC Medical Informatics and Decision Making*, *16*(74), 198–208.

https://doi.org/10.1186/s12911-016-0318-z

Berner, D. (2009). Allele frequency difference AFD–an intuitive alternative to Fst for

quantifying genetic population differentiation. *Genes*, *10*, 308.

https://doi.org/10.3390/genes10040308

Bhattacharyya, A. (1946). On a measure of divergence between two multinomial populations.

Sankhya, *7*, 401–406.

Bray, J. R., & Curtis, J. T. (1957). An ordination of upland forest communities of southern

Wisconsin. *Ecological Monographs*, *27*, 325–349. https://doi.org/10.2307/1942268

Buneman, P. (1974). A note on the metric properties of trees. *Journal of Combinatorial*

Theory B, *17*, 48–50. https://doi.org/10.1016/0095-8956(74)90047-1

Cailliez, F. (1983). The analytical solution to the additive constant problem. *Psychometrika*,

48, 305–308. https://doi.org/10.1007/BF02294026

Cailliez, F., & Pages, J. P. (1976). *Introduction à l'analyse des données*. Society de

Mathematiques Appliquées et de Sciences.

Cangelosi, R., & Goriely, A. (2007). Component retention in principal component analysis

with application to cDNA microarray data. *Biology Direct*, *2(2)*,

https://doi.org/10.1186/1745-6150-2-2.

Cattell, R. B. (1966). The scree test for the number of factors. *Multivariate Behavioral*

Research, *1*, 245–276. https://doi.org/10.1207/s15327906mbr0102_10

Chang, C. C., Chow, C. C., Tellier, L. C., Vattikuti, S., Purcell, S. M., & Lee, J. J. (2015).

Second-generation PLINK: rising to the challenge of larger and richer datasets.

GigaScience, *4*, 1–16. https://doi.org/10.1186/s13742-015-0047-8

Choi, S. S., Cha, S. H., & Tappert, C. C. (2010). A survey of binary similarity and distance

measures. *Systematics, Cybernetics and Informatics*, *8*, 43–48.

Cox, T. F., & Cox, M. A. A. (2001). *Multidimensional scaling* (2nd ed.). Chapman & Hall.

Czekanowski, J. (1913). *Zarys Metod Statystycznych w Zastosowaniu do Antropologii*
[Outline of statistical methods: as applied to anthropology]. Prace Towarzystwa
Naukowego Warszawskiego [Works of the Warsaw Scientific Society].

Daly, M., Rioux, J., Schaffner, S., Hudson, T., & Lander, E. (2001). High-resolution
haplotype structure in the human genome. *Nature Genetics*, *29*, 229–232.
https://doi.org/10.1038/ng1001-229

de Ley, J., Cattair, H., & Reynaerts, A. (1970). The quantitative measurement of DNA
hybridization from renaturation rates. *European Journal of Biochemistry*, *12*, 133–142.
https://doi.org/10.1111/j.1432-1033.1970.tb00830.x

Deza, M. M., & Deza, E. (2009). *Encyclopedia of Distances*. Springer-Verlag.
https://doi.org/10.1007/978-3-642-00234-2

Dice, L. R. (1945). Measures of the amount of ecologic association between species.
Ecology, *26*, 297–302. https://doi.org/10.2307/1932409

Dray, S., & Josse, J. (2015). Principal component analysis with missing values: a
comparative survey of methods. *Plant Ecology*, *216*, 657–667.
https://doi.org/10.1007/s11258-014-0406-z

Edwards, A. W. F. (1971). Distances between populations on the basis of gene frequencies.
Biometrics, *27*, 873–881.

Edwards, A. W. F., & Cavalli-Sforza, L. L. (1964). Reconstruction of evolutionary trees. In
Phenetic and Phylogenetic Classification (pp. 67–76). Systematics Association.

Edwards, A. W. F., & Cavalli-Sforza, L. L. (1967). Analysis, phylogenetic procedures,
models and estimation. *American Journal of Human Genetics*, *19*, 233–257.

Edwards, S. V., Shultz, A. J., & Campbell-Staton, S. C. (2015). Next-generation sequencing
and the expanding domain of phylogeography. *Folia Zoologica*, *64*, 187–206.
https://doi.org/10.25225/fozo.v64.i3.a2.2015

Elhaik, E. (2022). Principal Component Analyses (PCA)-based findings in population genetic
studies are highly biased and must be reevaluated. *Scientific Reports*, *12*, 14683.
https://doi.org/10.1038/s41598-022-14395-4

Elshibli, S. & Korpelainen, H. (2021). Genetic Diversity and population structure of

Medemia argun (Mart.) Wurttenb. ex H.Wendl.based on genome-wide markers.

Frontiers in Ecology and Evolution 9:687188. https://doi.org/10.3389/fevo.2021.687188

Excoffier, L. (2001). Analysis of population subdivision. In D. J. Balding, M. Bishop, & C.

Cannings (Eds.), *Handbook of Statistical Genetics* (pp. 271–307). John Wiley & Sons

Ltd.

Faith, D. P. (1985). A model of immunological distances in systematics. *Journal of*

Theoretical Biology, *114*, 511–526. https://doi.org/10.1016/S0022-5193(85)80041-2

Farris, J.S. (1981). Distance data revisited. Cladistics 1(10, 67-86.

https://doi.org/10.1111/j.1096-0031.1985.tb00411.x

Fix, A. G. (1997). Gene frequency clines produced by kin-structured founder effects. *Human*

Biology, *69*, 663–673.

Frichot, E., Mathieu, F., Trouillon, T., Bouchard, G., & François, O. (2013). Fast inference of

admixture coefficients using sparse non-negative matrix factorization algorithms. *ArXiv*,

25, 1–31. https://doi.org/http://arxiv.org/abs/1309.6208

Gao, X., & Starmer, J. (2007). Human population structure detection via multilocus genotype

clustering. *BMC Genetics*, *8*, 34. https://doi.org/10.1186/1471-2156-8-34

Gauch, H. G. J. (1982). Noise reduction by eigenvector ordinations. *Ecology*, *63*, 1643–1649.

https://doi.org/10.2307/1940105

Georges, A., & Adams, M. (1992). A phylogeny for australian chelid turtles based on

allozyme electrophoresis. *Australian Journal of Zoology*, *40*(5).

https://doi.org/10.1071/ZO9920453

Georges, A., Gruber, B., Pauly, G. B., White, D., Adams, M., Young, M. J., Kilian, A.,

Zhang, X., Shaffer, H. B., & Unmack, P. J. (2018). Genomewide SNP markers breathe

new life into phylogeography and species delimitation for the problematic short-necked

turtles (Chelidae: *Emydura*) of eastern Australia. *Molecular Ecology*, *27*(24), 5195–

5213. https://doi.org/10.1111/mec.14925

Goudet, J. (2005). Hierfstat, a package for R to compute and test variance components and F-

statistics. *Molecular Ecology Notes*, *5*, 184–186. https://doi.org/10.1111/j.1471-

8286.2004.00828.x

Gower, J. C. (1966). Some distance properties of latent root and vector methods used in

multivariate analysis. *Biometrika*, *53*, 325–338. https://doi.org/10.1093/biomet/53.3-

4.325

Gower, J. C. (1982). Euclidean distance geometry. *Mathematical Scientist*, *7*, 1–14.

Gower, J. C., & Legendre, P. (1986). Metric and Euclidean properties of dissimilarity

coefficients. *Journal of Classification*, *3*, 5–48. https://doi.org/10.1007/BF01896809

Gruber, B., Unmack, P. J., Berry, O. F., & Georges, A. (2018). dartr: An r package to

facilitate analysis of SNP data generated from reduced representation genome

sequencing. *Molecular Ecology Resources*, *18*(3), 691–699.

https://doi.org/10.1111/1755-0998.12745

Guillot, G., & Rousset, F. (2013). Dismantling the Mantel tests. *Methods in Ecology and*

Evolution, 4, 336–344. https://doi.org/10.1111/2041-210x.12018

Guttman, L. (1954). Some necessary conditions for common factor analysis. *Psychometrika*,

19, 149–161. https://doi.org/10.1007/BF02289162

Hirayama, H., Tamaoka, J., & Horikoshi, K. (1996). Improved immobilization of DNA to

microwell plates for DNA–DNA hybridization. *Nucleic Acids Research*, *24*, 4098–4099.

https://doi.org/10.1093/nar/24.20.4098

Holsinger, K., & Weir, B. (2009). Genetics in geographically structured populations:

defining, estimating and interpreting FST. *Nature Reviews Genetics*, *10*, 639–650.

https://doi.org/10.1038/nrg2611

Hotelling, H. (1933). Analysis of a complex of statistical variables into principal components.

Journal of Educational Psychology, *24*, 417–441. https://doi.org/10.1037/h0071325

Jaccard, P. (1912). The distribution of the flora in the alpine zone. *New Phytologist*, *11*, 37–

50. https://doi.org/10.1111/j.1469-8137.1912.tb05611.x

Jaccoud, D,. Peng, K., Feinstein, D. & Kilian, A. (2001). Diversity arrays: a solid state

technology for sequence information independent genotyping. *Nucleic Acids Research,*

29:E25. https://doi.org/10.1093/nar/29.4.e25.

Jackson, D. A. (1993). Stopping rules in Principal Components Analysis: A comparison of

heuristical and statistical approaches. *Ecology*, *74*, 2204–2214.

https://doi.org/10.2307/1939574

Jansen, J., & van Hintum, H. (2007). Genetic distance sampling: a novel sampling method for
obtaining core collections using genetic distances with an application to cultivated
lettuce. *Theoretical and Applied Genetics*, *114*, 421–428.
https://doi.org/10.1007/s00122-006-0433-9

Jolliffe, I. T. (1972). Discarding variables in a principal component analysis. I. Artificial data.
Applied Statistics, *21*, 160–173. https://doi.org/10.2307/2346488

Jolliffe, I. T. (2002). *Principal Component Analysis* (Second). Springer International
Publishing.

Jolliffe, I. T., & Cadima, J. (2016). Principal component analysis: a review and recent
developments. *Philosophical Transactions of the Royal Society A*, *374*, 20150202.
https://doi.org/10.1098/rsta.2015.0202

Jombart, T. (2008). adegenet: a R package for the multivariate analysis of genetic markers.
Bioinformatics, *24*, 1403–1405. https://doi.org/10.1093/bioinformatics/btn129

Jombart, T., & Ahmed, I. (2011). adegenet 1.3-1: new tools for the analysis of genome-wide
SNP data. *Bioinformatics*, *27*, 3070–3071. https://doi.org/10.1093/bioinformatics/btr521

Jombart, T., Devillard, S., & Balloux, F. (2010). Discriminant analysis of principal
components: a new method for the analysis of genetically structured populations. *BMC*
Genetics, *11*(94), 1–15. https://doi.org/10.1186/1471-2156-11-94

Kamvar, Z. N., Tabima, J. F., & Grünwald, N. J. (2014). Poppr: an R package for genetic
analysis of populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ*,
2, e281. http://doi.org/10.7717/peerj.281

Kilian, A., Wenzl, P., Huttner, E., Carling, J., Xia, L., Blois, H., Caig, V., Heller-Uszynska,
K., Jaccoud, D., Hopper, C., Aschenbrenner-Kilian, M., Evers, M., Peng, K., Cayla, C.,
Hok, P., & Uszynski, G. (2012). Diversity Arrays Technology: a generic genome
profiling technology on open platforms. In F. Pompanon & A. Bonin (Eds.), *Data*
Production and Analysis in Population Genomics: Methods and Protocols (pp. 67–89).
Humana Press. https://doi.org/10.1007/978-1-61779-870-2_5

Kirsch, J. A., Springer, M. S., Krajewski, C., Archer, M., Aplin, K., & Dickerman, A. W.
(1990). DNA/DNA hybridization studies of the carnivorous marsupials. I: The

intergeneric relationships of bandicoots (Marsupialia: Perameloidea). *Journal of*

Molecular Evolution, *30*, 434–438. https://doi.org/10.1007/BF02101115

Kosman, E., & Leonard, K. J. (2005). Similarity coefficients for molecular markers in studies

of genetic relationships between individuals for haploid, diploid, and polyploid species.

Molecular Ecology, *14*(2), 415–424. https://doi.org/10.1111/j.1365-294X.2005.02416.x

Kruskal, J. B. (1964). Nonmetric multidimensional scaling: a numerical method.

Psychometrika, *29*, 115–129. https://doi.org/10.1007/BF02289694

Legendre, P., & Legendre, L. (2012). Numerical Ecology. *Developments in Environmental*

Modelling, *24*, 1–990. https://doi.org/10.1016/B978-0-444-53868-0.50001-0

Levandowsky, M., & Winter, D. (1971). Distance between sets [5]. *Nature*, *234*(5323), 34–

35. https://doi.org/10.1038/234034A0

Libiger, O., Nievergelt, C. M., & Schork, N. J. (2009). Comparison of genetic distance

measures using human SNP genotype data. *Human Biology*, *81*, 389–406.

https://doi.org/10.3378/027.081.0401

Lin, Z., Yang, C., Zhu, Y., Duchi, J., Fu, Y., Wang, Y., Jiang, B., Zamanighomi, M., Xu, X.,

Li, M., Sestan, N., Zhao, H., & Wong, W. H. (2016). Simultaneous dimension reduction

and adjustment for confounding variation. *Proceedings of the National Academy of*

Sciences, *113*, 14662–14667. https://doi.org/10.1073/pnas.1617317113

Lingoes, J. C. (1971). Some boundary conditions for a monotone analysis of symmetric

matrices. *Psychometrika*, *36*, 195–203. https://doi.org/10.1007/BF02291398

Lotterhos, K.E. (2019). The effect of neutral recombination variation on genome scans for

selection, *G3 Genes|Genomes|Genetics, 9(6),* 1851–1867.

https://doi.org/10.1534/g3.119.400088

Lou, R.N. & Therkildsen, N.O. (2022). Batch effects in population genomic studies with low-

coverage whole genome sequencing data: Causes, detection and mitigation. Molecular

Ecology Resources 22(5), 1678-1692. https://doi-org.virtual.anu.edu.au/10.1111/1755-

0998.13559

Macarthur, R. (1957). On the relative abundance of bird species. *Proceedings of the National*

Academy of Sciences USA, *43*, 293–295. https://doi.org/10.1073/pnas.43.3.293

Mahboubi M, Mehrabi R, Naji AM, Talebi R. (2020). Whole-genome diversity, population structure and linkage disequilibrium analysis of globally diverse wheat genotypes using genotyping-by-sequencing DArTseq platform. *3 Biotech 10, 48.* https://doi.org/10.1007/s13205-019-2014-z.

Mamoozadeh, N. R., Graves, J. E., & McDowell, J. R. (2020). Genome-wide SNPs resolve spatiotemporal patterns of connectivity within striped marlin (*Kajikia audax*), a broadly distributed and highly migratory pelagic species. *Evolutionary Applications*, *13*, 677–698. https://doi.org/10.1111/eva.12892

Manel, S., Schwartz, M. K., Luikart, G., & Taberlet, P. (2003). Landscape genetics: combining landscape ecology and population genetics. *Trends in Ecology and Evolution*, *18*, 189–197. https://doi.org/10.1016/S0169-5347(03)00008-9

Mantel, N. (1967). The detection of disease clustering and a generalized regression approach. *Cancer Research*, *27*, 209–220. https://doi.org/10.1038/ng2088

Marchini, J., Howie, B., Myers, S., McVean, G., & Donnelly, P. (2007). A new multipoint method for genome-wide association studies by imputation of genotypes. *Nature Genetics*, *39*, 906–913. https://doi.org/10.1038/ng2088

Marczewski, M., & Steinhaus, H. (1959). On the taxonomic distance of biotopes. *Zastosowania Mathematyki*, *4*, 195–203. https://doi.org/10.4064/am-4-3-195-203

McRae, B. H. (2006). Isolation by resistance. *Evolution*, *60*, 1551–1561. https://doi.org/10.1111/j.0014-3820.2006.tb00500.x

McVean, G. (2009) A genealogical interpretation of principal components analysis. *PLoS Genetics* 5(10): e1000686. https://doi.org/10.1371/journal.pgen.1000686

Meisner, J., & Albrechtsen, A. (2018). Inferring population structure and admixture proportions in low-depth NGS data. *Genetics*, *210*, 719–731. https://doi.org/10.1534/genetics.118.301336

Mijangos, J., Gruber, B., Berry, O., Pacioni, C., & Georges, A. (2022). dartR v2: an accessible genetic analysis platform for conservation, ecology, and agriculture. *Methods in Ecology and Evolution*, *3*, 2150–2158. https://doi.org/10.1111/2041-210X.13918

Mijangos, J. L., Bino, G., Hawke, T., Kolomyjec, S. H., Kingsford, R. T., Sidhu, H., Grant, T., Day, J., Dias, K. N., Gongora, J., & Sherwin, W. B. (2022). Fragmentation by major

dams and implications for the future viability of platypus populations. *Communications*
Biology, *5*, 1–9. https://doi.org/10.1038/s42003-022-04038-9

Müller, T., Selinski, S., & Ickstadt, K. (2005). Cluster analysis: a comparison of different
similarity measures for SNP data. In *Technical Report, Universität Dortmund,*
Sonderforschungsbereich 475 - Komplexitätsreduktion in Multivariaten
Datenstrukturen, Dortmund (Vol. 14). http://hdl.handle.net/10419/22605

Nantongo, J.S., Agaba, H., & Gwali, S. (2022). SilicoDArT and SNP markers for genetic
diversity and population structure analysis of *Trema orientalis*; a fodder species. PLoS
ONE 17(8): e0267464. https://doi.org/10.1371/journal.pone.0267464

Negrini, R., Nicoloso, L., Crepaldi, P., Milanesi, E., Colli, L., Chegdani, F., Pariset, L.,
Dunner, S., Leveziel, H., Williams, J. L., & Ajmone Marsan, P. (2009). Assessing SNP
markers for assigning individuals to cattle populations. *Animal Genetics*, *40*(1), 18–26.
https://doi.org/10.1111/j.1365-2052.2008.01800.x

Nei, M. (1972). Genetic distance between populations. *American Naturalist*, *106*, 283–292.
https://doi.org/10.1086/282771

Nei, M. (1977). F-statistics and analysis of gene diversity in subdivided populations. *Annals*
of Human Genetics, *41*, 225–233. https://doi.org/10.1111/j.1469-1809.1977.tb01918.x

Nei, M., & Kumar, S. (2000). *Molecular Evolution and Phylogenetics*. Oxford University
Press.

Novembre, J., & Stephens, M. (2008). Interpreting principal component analyses of spatial
population genetic variation. Nature Genetics 40, 646–649.
https://doi.org/10.1038/ng.139

Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R. B.,
Simpson, G. L., Solymos, P., Stevens, M. H. H., & Wagner, H. (2014). *Vegan:*
Community Ecology Package. R Package Version 2.2-0.

Orlóci, L. (1978). *Multivariate Analysis in Vegetation Research* (2nd ed.). Dr W Junk
Publishers.

Östergren, J., Palm, S., Gilbey, J., & Dannewitz, J. (2020). Close relatives in population
samples: Evaluation of the consequences for genetic stock identification. *Molecular*
Ecology Resources 20(2), 498-510. https://doi.org/10.1111/1755-0998.13131

Paradis, E. (2010). Pegas: an R package for population genetics with an integrated-modular approach. *Bioinformatics*, *26*, 419–420. https://doi.org/10.1093/bioinformatics/btp696

Paradis, E., & Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics*, *35*, 526–528.

Patil, N., Berno, A. J., Hinds, D. A., Barrett, W. A., Doshi, J. M., Hacker, C. R., Kautzer, C., Lee, D. H., Marjoribanks, C., McDonough, D. P., Nguyen, B. T. N., Norris, M. C., Sheehan, J. B., Shen, N., Stern, D., Stokowski, R., Thomas, D. J., Trulson, M., Vyas, K. R., … Cox, D. (2001). Blocks of limited haplotype diversity revealed by high-resolution scanning of human Chromosome 21. *Science*, *294*, 1719–1723. https://doi.org/10.1126/science.1065573

Patterson, N., Price, A. L., & Reich, D. (2006). Population structure and eigenanalysis. *PLoS Genetics*, *2*, e190. https://doi.org/10.1371/journal.pgen.0020190

Pearson, K. (1901). On lines and planes of closest fit to systems of points in space. *Philisophical Magazine*, *2*, 559–572. https://doi.org/10.1080/14786440109462720

Peres-Neto, P. R., Jackson, D. A., & Somers, K. M. (2005). How many principal components? Stopping rules for determining the number of non-trivial axes revisited. *Computational Statistics and Data Analysis*, *49*, 974–997. https://doi.org/10.1016/j.csda.2004.06.015

Pugach, I., & Stoneking, M. (2015). Genome-wide insights into the genetic history of human populations. *Investigative Genetics*, *6*, 1–20. https://doi.org/http://doi.org/10.1186/s13323-015-0024-0

Rabinovich, A., Agarwal, S., Laris, C. A., Price, J. H., & Belongie, S. (2003). Unsupervised color decomposition of histologically stained tissue samples. *Advances in Neural Information Processing Systems*, *2003*, 667–674.

Reynolds, J., Weir, B. S., & Cockerham, C. C. (1983). Estimation of the coancestry coefficient: Basis for a short-term genetic distance. *Genetics*, *105*, 767–779. https://doi.org/10.1093/genetics/105.3.767

Rogers, J. S. (1972). Measures of Genetic Similarity and Genetic Distance. *Studies in Genetics VII, University of Texas Publication, Austin, USA*, *7213*, 145–153.

Rohlf, F. J. (1972). An empirical comparison of three ordination techniques in numerical

taxonomy. *Systematic Biology*, *21*, 271–280.

https://doi.org/10.1093/sysbio/21.3.271%0A

Saitou, N., & Nei, M. (1987). The neighbor-joining method: a new method for reconstructing

phylogenetic trees. *Molecular Biology and Evolution*, *4*, 406–425.

Sanghvi, L. D. (1953). Comparison of genetical and morphological methods for a study of

biological differences. *American Journal of Physical Anthropology*, *11*, 385–404.

https://doi.org/10.1002/ajpa.1330110313

Sansaloni, C., Petroli, C., Jaccoud, D., Carling, J., Detering, F., Grattapaglia, D., & Kilian, A.

(2011). Diversity Arrays Technology (DArT) and next-generation sequencing

combined: genome-wide, high throughput, highly informative genotyping for molecular

breeding of Eucalyptus. *BMC Proceedings*, *5*, P54. https://doi.org/10.1186/1753-6561-

5-s7-p54

Séré, M., Thévenon, S., Belem, A. M. G., & de Meeûs, T. (2017). Comparison of different

genetic distances to test isolation by distance between populations. *Heredity*, *119*, 55–

63. https://doi.org/10.1038/hdy.2017.26

Shepard, R. N. (1962). The analysis of proximities: multidimensional scaling with an

unknown distance function. *Psychometrika*, *27*, 125–139.

https://doi.org/10.1007/BF02289630

Sherwin, W. B. (2022). Bray-Curtis (AFD) differentiation in molecular ecology: Forecasting,

an adjustment (AA), and comparative performance in selection detection. *Ecology and*

Evolution, *12*, e9176. https://doi.org/10.1002/ece3.9176

Sibson, R. (1979). Studies in the robustness of multidimensional scaling: perturbational

analysis of classical scaling. *Journal of the Royal Statistical Society*, *41B*, 217–229.

https://doi.org/10.1111/j.2517-6161.1979.tb01076.x

Sokal, R. R., & Michener, C. D. (1958). A statistical method for evaluating systematic

relationships. *University of Kansas Science Bulletin*, *28*, 409–1438.

Sørensen, T. (1948). A method of establishing groups of equal amplitude in plant sociology

based on similarity of species and its application to analyses of the vegetation on Danish

commons. *Kongelige Danske Videnskabernes Selskab*, *5*, 1–34.

Spear, S. F., Cushman, S. A., & McRae, B. H. (2015). Resistance Surface Modeling in

Landscape Genetics. In N. Balkenhol, S. A. Cushman, A. T. Storfer, & L. P. Waits (Eds.), *Landscape Genetics* (pp. 129–148). John Wiley & Sons Ltd.

Tassi, F., Ghirotto, S., Mezzavilla, M., Vilaça, S. T., De Santi, L., & Barbujani, G. (2015). Early modern human dispersal from Africa: Genomic evidence for multiple waves of migration. *Investigative Genetics*, *6*, 1–16. https://doi.org/http://10.1186/s13323-015-0030-2

Tian, C., Gregersen, P., & Seldin, M. (2008). Accounting for ancestry: population substructure and genome-wide association studies. *Human Molecular Genetics*, *17*, R143-150. https://doi.org/10.1093/hmg/ddn268

Torgerson, W. (1952). Multidimensional scaling: I. Theory and method. *Psychometrika*, *17*, 401–419. https://doi.org/10.1007/BF02288916

Tracy, C., & Widom, H. (1993). Level-spacing distributions and the Airy kernel. *Physics Letters B*, *305*, 115–118. https://doi.org/10.1016/0370-2693(93)91114-3

Uffelmann, E., Huang, Q. Q., Munung, N. S., de Vrie, J., Okada, Y., Martin, A. R., Martin, H. C., Lappalainen, T., & Posthuma, D. (2021). Genome-wide association studies. *Nature Review Methods: Primers*, *1*, 1–59. https://doi.org/10.1038/s43586-021-00056-9

Wahlund, S. (1928). Zusammensetzung von populationen und korrelationserscheinungen vom standpunkt der vererbungslehre aus betrachtet. *Hereditas*, *11*, 65–106. https://doi.org/10.1111/j.1601-5223.1928.tb02483.x

Wang, Y., Sun, F., Lin, W., & Zhang, S. (2022). AC-PCoA: Adjustment for confounding factors using principal coordinate analysis. *BMC Computational Biology*, *18*, e1010184. https://doi.org/10.1371/journal.pcbi.1010184

Waples, R.S., & Anderson, E.C. (2017). Purging putative siblings from population genetic data sets: a cautionary view. Molecular Ecology 26(5), 1211-1224. https://doi.org/10.1111/mec.14022

Waples, R. S., Waples, R. K., & Ward, E. J. (2022). Pseudoreplication in genomic-scale data sets. *Molecular Ecology Resources*, *22*, 503–518. https://doi.org/10.1111/1755-0998.13482

Weir, B. S., & Cockerham, C. C. (1984). Estimating F-statistics for the analysis of population-structure. *Evolution*, *38*, 358–1370. https://doi.org/10.1111/j.1558-

5646.1984.tb05657.x

Wild, K. H., Roe, J. H., Schwanz, L., Georges, A., & Sarre, S. D. (2022). Evolutionary stability inferred for a free ranging lizard with sex-reversal. *Molecular Ecology*, *31*, 2281–2292. https://doi.org/10.1111/mec.16404

Wright, S. (1943). Isolation by distance. *Genetics*, *8*, 114–138. https://doi.org/10.1093/genetics/28.2.114

Wright, S. (1951). The genetical structure of populations. *Annals of Eugenics*, *15*, 323–354. https://doi.org/10.1111/j.1469-1809.1949.tb02451.x

Wu, W., Massart, D., & de Jong, S. (1997). The kernel PCA algorithms for wide data. Part I: Theory and algorithms. *Chemometrics and Intelligent Laboratory Systems* 36, 165–172 http://dx.doi.org/10.1016/S0169-7439(97)00010-5

Xu, G., Zhang, X., Chen, W., Zhang, R., Li, Z., Wen, W., Warburton, M., Li, J., Li, H., & Yang, X. (2022). Population genomics of *Zea* species identifies selection signatures during maize domestication and adaptation. *BMC Plant Biology*, *22*, 72. https://doi.org/10.1186/s12870-022-03427-w

Yang, J., Lee, S. H., Goddard, M. E., & Visscher, P. M. (2011). GCTA: a tool for genome-wide complex trait analysis. *American Journal of Human Genetics*, *88*, 76–82. https://doi.org/10.1016/j.ajhg.2010.11.011

Yi, X., & Latch, E. K. (2021). Nonrandom missing data can bias Principal Component Analysis inference of population genetic structure. *Molecular Ecology Resources*, *22*, 602–611. https://doi.org/10.1111/1755-0998.13498

Yin, C. (2020). Genotyping coronavirus SARS-CoV-2: methods and implications. *Genomics*, *112*, 3588–3596. https://doi.org/10.1016/j.ygeno.2020.04.016

Yoshioka, P. M. (2008). Misidentification of the Bray-Curtis similarity index. *Marine Ecology Progress Series*, *368*, 309–310. https://doi.org/10.3354/meps07728