1

Precision of the Integrated Cognitive Assessment for the assessment of neurocognitive

performance in athletes

Daniel J. Glassbrook*[1,2¶], Paul L. Chazot[1,3¶], and Karen Hind[1¶]

[1]Wolfson Research Institute for Health and Wellbeing, Durham University, Durham, United

Kingdom.

[2]School of Science, Technology and Health, York St John University, York, United

Kingdom.

[3]Department of Biosciences, Durham University, Durham, United Kingdom.

*Corresponding author:

Email: d.glassbrook@yorksj.ac.uk (DG)

[¶]These authors contributed equally to this work.

Abstract

Choice reaction time tests are commonly used for the assessment of cognitive function, and may be useful to assess the effect of concussions. This study investigated the precision of the Integrated Cognitive Assessment (ICA; Cognetivity Neurosciences Ltd., Vancouver, Canada) test for the assessment of cognitive function in athletes. Thirty-one participants volunteered to take part in this study, from both contact ($n = 22$) and non-contact sports ($n = 9$). Participants performed the ICA test consecutively both before and after normal training session to simulate resting and post-sport conditions. Precision errors, relationships (Pearson's r), and internal consistency (Cronbach's Alpha) were calculated for three variables, ICA Index (overall information processing ability), ICA Speed (information processing speed) and ICA Accuracy (information processing accuracy). ICA precision errors [root mean squared-standard deviation, RMS-SD (coefficient of variation, %CV)] pre-sport were ICA Index: 5.18 (7.14%), ICA Speed: 3.98 (4.64%), and ICA Accuracy: 3.64 (5.00%); and post-sport were ICA Index: 3.96 (4.94%), ICA Speed: 2.14 (2.32%), and ICA Accuracy 3.40 (4.25%). The ICA test demonstrates high in-vivo precision with all variables except ICA Index (7.14%) demonstrating an acceptable precision error of ≤5% %CV. All variables demonstrated strong relationships between consecutive tests pre- and post-sport ($r \geq 0.8$) except for the ICA Index post-sport which demonstrated a moderate ($r \geq 0.5$) relationship. The ICA Index demonstrated good internal consistency ($\alpha \geq 0.8$) for both pre-and post-sport. The ICA Speed and ICA Accuracy variables demonstrated excellent internal consistency ($\alpha \geq 0.9$) for both pre-and post-sport. The ICA test is suitable for the assessment of cognitive function pre- and post-sport.

Key words: Information Processing, Reaction Time, Concussion, Rapid Categorisation Task, Contact Sport, Non-Contact Sport.

Introduction

The brain plays a vital role in sports participation, and is responsible for decision making, and good decision making is key to success in sport (1). Athletes are required to demonstrate high levels of technical and perceptual decision making ability, all while under pressure and fatigue (2). Assessing cognitive function, such as information processing and decision making ability is a valuable tool which may quantify psychological ability in athletes. Additionally, assessing cognitive function may give insight into the effect of sport participation on the brain, for example through the effect on reaction time and decision making ability. There are many positive effects of sport and exercise participation, including cognitive benefits such as neuroplasticity, the process of adaptive structural and functional changes to the brain (3). However, injury is an innate part of sport participation due to the physical nature of sport. In terms of cognitive function, a relevant injury with detrimental effects is concussion. A concussion can arise from rapid rotational or linear acceleration or deceleration of the head, resulting in a sudden movement of the brain within the cranium where the brain collides with the cranium wall (4). People who play contact sports may be at a higher risk of sustaining a concussion due to the nature of these sports and associated collisions (5). In fact, sports-related concussion is one of the highest reported injuries in rugby union players in the United Kingdom (6). A person with a concussion can experience symptoms such as loss of consciousness, nausea, headaches, vertigo and amnesia (7), and evidence has demonstrated links between concussion and persistent cognitive decline and long-term neurodegeneration (6, 8-10). However, concussion is currently difficult to objectively assess. Although, understanding cognitive function pre- and post-sport participation may be a starting place.

One proposed method to assess the effects of concussion and cognitive function clinically is via information processing and reaction time tests. Reduced information processing ability,

including memory, is linked with concussion (11-13). Indeed, altered mental state and confusion are also symptoms associated with concussion that can negatively affect memory and information processing ability (7). Several tests have been developed to diagnose if an athlete has sustained a concussion which assess information processing ability. In sport, an immediate on- or off-field method for an athlete suspected of concussion is to assess their orientation and ability to answer questions for example, about their whereabouts and details about the match, such as the Maddocks questions (14). In professional sport, additional off-field assessment tools such as the Sports Concussion Assessment Tool (SCAT) are used (15). There is also a SCAT which has been developed for children (16). The SCAT includes a symptom check, memory assessment and balance evaluation, and the test results are compared to a previously recorded baseline assessment, or to normative data. Additionally, information processing speed underpins several conditions of cognitive dysfunction, for example, multiple sclerosis (17, 18) and Alzheimer's disease (19).

Slower reaction time is a commonly used indicator of cognitive change following concussion (20, 21). Two common types of reaction test include measurement of simple reaction time (SRT) or choice reaction time (CRT) (22). SRT is recorded when there is only one possible stimulus (signal) and one possible response (action), for example tapping anywhere on a screen when any image appears. In CRT tasks there are two or more possible stimuli, each of which requires a quite different response, for example, tapping on the left of the screen when an image of an object appears on the screen, and tapping on the right when an image of an animal appears on the screen. Clinically measured reaction time has been shown to be a sensitive measure of concussion status. SRT tests such as a weighted object drop and catch (23), ruler drop test (24), and somatosensory assessment, where the participant held a device similar to a computer mouse and reacted to a vibration applied to one or both the index or middle finger, and responded by clicking with the finger (or both) that received the vibration (25), have all been

used to show slower reaction time in participants with concussion. Computerised tests have

also been shown to be sensitive and able to determine athletes who experienced concussion

versus those without, for example the CogSport choice reaction time test and Immediate Post-

Concussion Assessment and Cognitive Testing (imPACT) (26-29). The CogSport choice test

takes approximately 20 minutes, and consists of seven tasks including a SRT, CRT and

complex reaction time task, as well as tests of visual and working memory, all designed as card

games (30). The imPACT is performed with a computer screen and mouse and typically takes

20-25 minutes and comprises of six test modules assessing attention, memory, processing

speed and reaction time, through tests such as word memory and symbol matching (31).

Interestingly, reaction time has also been used to show the lasting effect of concussion on

reaction time, with reaction time scores only returning to baseline at 10 to 14 days post injury

(24), and up to 21 days post injury (32).

The Integrated Cognitive Assessment (ICA; Cognetivity Neurosciences Ltd., Vancouver,

Canada) (33, 34), is a newly developed method for the assessment of cognitive function, and

may be applicable to the assessment of concussion in athletic populations. The ICA is a short

computerised cognitive test of cognitive function via an assessment of information processing

(CRT) speed based on a rapid categorisation task (33, 35). The ICA test can be completed in a

short amount of time (~5 minutes), on a handheld device such as an iPhone or iPad, and is

therefore able to be administered easily pre- and post-sport. Performing an assessment of

cognitive function such as the ICA pre-sport in a rested state effectively provides a baseline

measure of cognitive function, such as done with the SCAT. Performing a test of cognitive

function post-sport then can be compared to the baseline measure of pre-sport and determine

the effect of sport participation on cognitive function. Additionally, the ICA has been shown

to accurately detect mild cognitive impairment and be moderately to highly correlated with

other popular pen-and-paper cognitive tests such as the Montreal Cognitive Assessment

(Pearsons r = 0.58) and Addenbrooke's Cognitive Examination (Pearsons r = 0.62) cognitive

tests (33).

The ICA has been shown to accurately measure cognitive impairment in patients in the early

stages of dementia (33). However, to date, no known study has investigated the intra-day

precision of the ICA test. Therefore, the purpose of this study was to determine the same-day,

in-vivo (i.e., within whole living organisms; within people) precision of the ICA test to assess

cognitive function. It is hypothesised that the ICA will demonstrate acceptable precision, and

suitability for the assessment of cognitive function pre- and post-sport.

Materials and Methods

Thirty-one participants volunteered to take part in this study. Participant characteristics are

presented in Table 1. Participants were recruited from March to May 2022, from the University

sporting facility via advertisement flyer, and from the wider sporting community through

existing relationships with sporting clubs. After initially expressing interest to take part in the

study, participants were provided with an information sheet and given at least 24 hours to

consider their participation in the study before a follow up was sent by the lead researcher.

After confirmation of eligibility, participants were then booked in at a time and date of their

convenience for data collection. Participants were eligible for participation if they were a

current contact sport or non-contact sport athlete (Table 2) aged 18-40 years, and healthy;

having no underlying medical issues that affect participation in sport. Participants were

excluded if they were injured, pregnant, or suffering from post-concussion syndrome. Post-

concussion syndrome is defined as the collective experience of symptoms after sustaining a

concussion, such as headaches, fatigue, depression, anxiety and irritability (36, 37). This study

was approved by the Durham University Sport and Exercise Sciences Ethics Committee

(reference: SPORT-2022-01-07T10_44_59-srhd22), and written informed consent was

provided by each participant prior to participation. Participant characteristics, demographics

and athletic history were collected via standardised questionnaire directly after consent to

participate was obtained. Ten participants indicated prior to testing a history of concussions

(experienced 1 or more); all participants presented symptom free at the time of testing and were

not experiencing post-concussion syndrome.

To simulate resting- and post-sport conditions, participants performed the ICA test (version

1.6.0 or 1.7.0) before and after a normal training session for their respective sports. Data

collection was performed in a quiet room to minimise distractions, with a maximum of two

participants (separated within the room) performing the test at a time. Prior to their sports

training, and within an hour of the training start time, participants completed two consecutive

ICA tests. The participants then completed a normal training session and then two consecutive

ICA tests again, within an hour of the training finishing.

Table 1: Participant Characteristics

	Age (Yr.)	Height (m)	Body Mass (kg)
Total ($n = 31$)	23.7 ± 5.7	1.78 ± 0.09	72.6 ± 8.3
Male ($n = 16$)	22.9 ± 4.7	1.82 ± 0.08	75.7 ± 7.4
Female ($n = 15$)	24.6 ± 6.6	1.71 ± 0.07	68.0 ± 9.9
Contact Sport ($n = 22$)	24.9 ± 6.3	1.80 ± 0.07	75.5 ± 6.0
Non-Contact Sport ($n = 9$)	20.6 ± 0.6	174.1 ± 8.3	77.2 ± 12.2

Data are presented as mean ± standard deviation. Yr., Years; m, Metres; kg, Kilograms; n, Number.

158   Table 2: Sports Breakdown

Contact Sports ($n = 22$)		Non-Contact Sports ($n = 9$)	
Rugby Union ($n = 7$)	Semi-professional, Amateur	Touch Rugby ($n = 5$)	Amateur
Boxing ($n = 6$)	Amateur	Athletics ($n = 4$)	Amateur
Muay Thai (Kickboxing) ($n = 5$)	Professional, Amateur		
Indoor Football ($n = 4$)	Amateur		

n, Number.

The ICA has been described in detail in published works (35, 38). In summary, the ICA is a

short computerised cognitive test of cognitive function via an assessment of information

processing (CRT) speed based on a rapid categorisation task, and in contrast to the CogSport

and imPACT tests is independent of language or colour. The ICA in this study was completed

on an iPad, with each test taking approximately five minutes, considerably shorter than the 20+

minute CogSport and imPACT tests; the imPACT test of which is also not portable (completed

on a computer with screen and mouse). Each ICA test utilises the brains powerful reaction to

animal stimulus (39) and comprises of 100 natural images, with 10 additional practice images

provided in a practice section prior to the test. The practice images are not included in the test

score. The images are a mixture of animals (e.g., birds, fish, mammals), or non-animals (e.g.,

objects, food, vehicles), and are presented in rapid succession. Images appear for 100ms,

followed by a 20ms inter-stimulus interval and a 250ms dynamic noisy mask. Participants react

to the image by tapping with their thumb on the screen; on the right-hand side to select 'animal'

and on the left-hand side of the screen to select 'non-animal'. Participants were instructed to

perform the test as quick as possible, and these instructions were given both verbally and

presented on the screen. There are no associated risks with completing the ICA, and the test

provides three variables: ICA Speed, the response reaction times in trials they responded

correctly; ICA Accuracy, choice accuracy, the percentage of correct test responses; and ICA

Index, overall information processing ability, a combination of ICA Speed and ICA Accuracy.

Each variable is calculated by the following equations (33, 38):

179
$$Accuracy = \frac{Number\ of\ correct\ categorisations}{Total\ number\ of\ images} \ x\ 100$$

180
$$Speed = \min\left[100, 100e^{-\frac{mean\ correct\ reaction\ time}{1025}+0.341}\right]$$

181
$$ICA\ Index = \left(\frac{Speed}{100}\ x\ \frac{Accuracy}{100}\right) x\ 100$$

There is a speed-accuracy trade-off in reaction test performance, and often scoring higher in

either speed or accuracy is achieved at the expense of the other capacity (40, 41). To combat

the potential negative reflection on overall information processing ability from a poor speed or

accuracy score, a common solution is the inverse efficiency score (42), whereby speed and

accuracy are combined into a single score. In the case of the ICA, this concept is applied and

manifests as the ICA Index variable. Test results are recorded via report immediately after test

completion.

ICA Accuracy, ICA Speed and ICA Index are represented by Arbitrary Units and results of

each equation are transformed and scored by a standardised scale ranging from 0 to 100. I.e., a

Speed score closer to 100 indicates faster reaction times, while a Speed score closer to 0

indicates slower reaction times; an Accuracy score closer to 100 indicates higher accuracy, and

an Accuracy score closer to 0 indicates lower accuracy; and an ICA Index score closer to 100

represents greater overall information processing ability, whereas an ICA Index score closer to

0 indicates poorer overall information processing ability. Additionally, the specific

components included in the Speed equation (i.e., division by 1025 and addition of 0.341)

further serve to standardise and scale the Speed score. These additional components are

determined empirically to ensure that the Speed scores fall within the desired range of 0 to 100

for the majority of participants. Natural Log transformation further aids in normalizing the

distribution of scores.

All data analysis was performed in Microsoft Excel (2016). Raw anonymised data for ICA

Index, ICA Speed, and ICA Accuracy were extracted and exported to Microsoft Excel for

analysis (43). Throughout the study the lead author had access to information that could

identify individual participants, however, all analysis was performed on anonymised data.

Precision has been previously used to define test suitability in similar cognitive tests to the

ICA, such as the CogSport choice reaction time test (44). Precision of ICA scores and least

significant change (LSC) were calculated at the 95% confidence level. Precision was

determined as root mean square standard deviation (RMS-SD), coefficient of variation (CV),

and percentage CV (%CV). RMS-SD represents the sample standard deviation of the

differences between predicted values and observed values, and is calculated via the following

formulae, where SD represents standard deviation and *n* represents the number of participants:

212
$$\sqrt{(\frac{\Sigma SD^2}{n})}$$

The %CV expresses test variation relative to the mean of two tests and is corrected for small

sample bias, and was defined as acceptable <5% (45). The LSC represents a true meaningful

change was calculated from the precision errors (LSC = RMS-SD * 2.77). Pearson's correlation

coefficient was calculated for each variable to determine reliability and the strength of the

relationship between consecutive tests, with 0.2, 0.5, and 0.8 representing weak, moderate, and

strong correlations, respectively (46). Cronbach's Alpha was calculated for each variable to

determine internal consistency, with 0.9, 0.8, 0.7, 0.6, 0.5 and < 0.5 representing excellent,

good, acceptable, questionable, poor and unacceptable internal consistency, respectively (47).

Results

Results of the precision analysis for each ICA variable pre- and post-sport are presented in

Table 3. All variables except for ICA Index pre-sport had a precision error of ≤5% %CV. LSC

results are presented in Table 4. Pearson correlation coefficient data is presented in *5*. All

variables demonstrated strong relationships between consecutive tests pre- and post-sport,

except for the ICA Index post-sport which demonstrated a moderate relationship. Cronbach's

Alpha data is presented in Table *5*. The ICA Index demonstrated good internal consistency for

both pre-and post-sport. The Speed and Accuracy variables demonstrated excellent internal

consistency for both pre-and post-sport.

Table 3: Precision Analysis Results

Variable	Precision (*n* = 31)		
	RMS-SD	CV	%CV
Pre			
ICA Index	5.18	0.07	7.14
ICA Speed	3.98	0.05	4.64
ICA Accuracy	3.64	0.05	5.00
Post			
ICA Index	3.96	0.05	4.94
ICA Speed	2.14	0.02	2.32
ICA Accuracy	3.40	0.04	4.25

n, Number; ICA, Integrated Cognitive Assessment; RMS-SD, Root Mean Square Standard Deviation; CV, Coefficient of Variation; %, Percentage.

231 Table 4: Least Significant Change Results

Variable	LSC (*n* = 31)		
	RMS-SD	CV	%CV
Pre			
ICA Index	14.36	0.20	19.78
ICA Speed	11.01	0.13	12.86
ICA Accuracy	10.09	0.14	13.9
Post			
ICA Index	10.96	0.14	13.7
ICA Speed	5.94	0.06	6.43
ICA Accuracy	9.43	0.12	11.78

LSC, Least Significant Change; n, Number; ICA, Integrated Cognitive Assessment; RMS-SD, Root Mean Square Standard Deviation; CV, Coefficient of Variation; %, Percentage.

Table 5: Relationships between consecutive tests

Variable	Test 1	Test 2	Correlation (r)	Cronbach's Alpha (α)
Pre				
ICA Index	76.6 ± 8.3	82.2 ± 6.1	0.82	0.88
ICA Speed	88.6 ± 8.1	92.3 ± 7.3	0.85	0.91
ICA Accuracy	86.9 ± 8.9	89.2 ± 6.2	0.87	0.90
Post				
ICA Index	81.5 ± 7.0	83.0 ± 6.7	0.68	0.81
ICA Speed	94.8 ± 5.2	95.6 ± 4.6	0.82	0.90
ICA Accuracy	86.1 ± 9.1	87.0 ± 8.3	0.85	0.92

n, Number; ICA, Integrated Cognitive Assessment; Data are presented as mean ± standard deviation.

Discussion

The purpose of this study was to determine the same-day, in-vivo precision of the ICA test to assess cognitive function. The results of this study support the ICA as a tool with acceptable precision to measure changes in cognitive ability pre- and post-sport, confirming the hypothesis. All ICA variables in this study, except for ICA Index pre-sport demonstrated a precision error of ≤5% %CV. All variables demonstrated moderate to strong relationships and good to excellent internal consistency between consecutive tests pre- and post-sport. The ICA may be clinically relevant for the assessment of cognitive ability, and the effect of sport on cognitive ability.

The higher ICA Index precision score (7.14 %CV) pre-sport than post-sport (4.94 %CV) in this study may be explained by the larger average difference between test one and test two pre-sport (76.6 ± 8.3 to 82.2 ± 6.1), compared to a smaller average difference in ICA Index scores between test one and test two post-sport (81.5 ± 7.0 to 83.0 ± 6.7). This is further exemplified by a larger ICA Index RMS-SD pre-sport than post-sport (5.18 and 3.96, respectively), which indicates more variance in observed data around the mean. This result may be due to an increased level of comfort with the test from the first pre-sport ICA test to the subsequent test, and possibly a learning effect. However, this is in contrast to previous work which showed no learning effect for the ICA test in healthy participants and those diagnosed with dementia (33).

Additionally, a strong relationship (r = 0.82) and good internal consistency (α = 0.88) between

ICA tests and the ICA Index variable was observed pre-sport (Test 1 and Test 2; Table *5*),

further suggesting a minimal learning effect, and supporting the test as reliable.

All variables showed greater precision post-sport compared to pre-sport. This may be due to

the many positive physiological benefits that exercise has, such as an increase in blood flow to

muscles and brain (48), structural and functional changes in the brain (3), and increases

information processing ability (49). Indeed, improvements in cognitive function after a bout of

exercise is supported by previous research (50, 51).

Previous research looking at precision in a similar cognitive test to the ICA, the CogSport

choice reaction time test, has shown lower %CV for mean choice reaction time (speed) (1.4

%CV), and higher %CV for choice reaction time accuracy (11.4 %CV) (44). These results are

interesting as the ICA is shown to be less precise in measuring reaction speed (2.32 - 4.64 %CV

vs 1.4 %CV), however, the ICA test is shown to be more precise in terms of accuracy (4.25 –

5.00 %CV vs 11.4 %CV). These results may indicate that the test you adopt needs to be specific

to the variable of interest (i.e., speed or accuracy), however, this should be negated in the case

of the ICA via the ICA Index variable as an inverse efficiency score (42), whereby speed and

accuracy are combined into a single score. The contrasting results between the present study

and that of Straume-Naesheim, Andersen (44) may be due to the populations used; the

CogSport test was used in elite football players only, whereas only a small percentage of the

participants in the present study are practicing professionally (Table 2). The present study

recruited participants from a variety of sports, each with their own decision making and

reaction characteristics, in comparison to only football. Additionally, a greater number of

participants (*n* = 289) performed the CogSport test in comparison to the ICA in the present

study (*n* = 31). However, the administration procedure of the CogSport test was similar to the

administration of the ICA test, and was also administered under controlled conditions, i.e., in a quiet room to minimise distractions and with more than one participant performing the test at a time.

A potential limitation to this study is that information about the specific duration an intensity of training sessions undergone by participants was not recorded. Only that it was a typical or normal training session. i.e., not a session of just running or conditioning, and included practice of the sport itself. Additionally, the optimal intensity and duration of exercise that may impact cognitive function was not assessed. A relatively small sample size was used, although the sample size exceeds the minimum of 30 participants required for precision analysis with two consecutive tests (52), a greater number of participants could strengthen the statistical analysis. Future studies should look to recruit larger numbers, and complete comparisons between contact and non-contact sports, and between amateur and professional level participants.

In conclusion, the ICA is a practical test which can be used to measure cognitive function before and after sport participation. The results of this study support the ICA as a precise (as determined by %CV values) measure of information processing speed and information processing accuracy, and overall information processing ability (ICA Index). The ICA can be used for the assessment of cognitive function, and may be useful as a method to assess the effects of sport on cognitive function.

References

1.   Hepler TJ. Decision-making in sport under mental and physical stress. International Journal of Kinesiology and Sports Science. 2015;3(4):79-83.

2.   Gabbett TJ, Carius J, Mulvey M. Does improved decision-making ability reduce the physiological demands of game-based activities in field sport athletes? The Journal of Strength & Conditioning Research. 2008;22(6):2027-35.

3.   Fernandes J, Arida RM, Gomez-Pinilla F. Physical exercise as an epigenetic

modulator of brain plasticity and cognition. Neuroscience & Biobehavioral Reviews.

2017;80:443-56.

4.   Jordan BD. The clinical spectrum of sport-related traumatic brain injury. Nature

Reviews Neurology. 2013;9(4):222-30.

5.   La Fountaine MF, Toda M, Testa AJ, Hill-Lombardi V. Autonomic nervous system

responses to concussion: Arterial pulse contour analysis. Frontiers in Neurology. 2016;7.

6.   Hume PA, Theadom A, Lewis GN, Quarrie KL, Brown SR, Hill R, et al. A

Comparison of cognitive function in former rugby union players compared with former non-

contact-sport players and the impact of concussion history. Sports Medicine.

2017;47(6):1209-20.

7.   Sharma A, Hind K, Hume P, Singh J, Neary JP. Neurovascular coupling by functional

near infra-red spectroscopy and sport-related concussion in retired rugby players: The UK

rugby health project. Frontiers in Human Neuroscience. 2020;14.

8.   Manley G, Gardner AJ, Schneider KJ, Guskiewicz KM, Bailes J, Cantu RC, et al. A

systematic review of potential long-term effects of sport-related concussion. British journal

of sports medicine. 2017;51(12):969-77.

9.   Brush CJ, Ehmann PJ, Olson RL, Bixby WR, Alderman BL. Do sport-related

concussions result in long-term cognitive impairment? A review of event-related potential

research. International journal of psychophysiology. 2018;132:124-34.

10.   Cunningham J, Broglio SP, O'Grady M, Wilson F. History of sport-related concussion

and long-term clinical cognitive health outcomes in retired athletes: a systematic review.

Journal of athletic training. 2020;55(2):132-58.

11.   Bernstein DM. Information processing difficulty long after self-reported concussion.

Journal of the International Neuropsychological Society. 2002;8(5):673-82.

12.   Gronwall D, Wrightson P. Memory and information processing capacity after closed

head injury. Journal of Neurology, Neurosurgery & Psychiatry. 1981;44(10):889-95.

13.   O'Jile JR, Ryan LM, Betz B, Parks-Levy J, Hilsabeck RC, Rhudy JL, et al.

Information processing following mild head injury. Archives of clinical neuropsychology.

2006;21(4):293-6.

14.   Maddocks DL, Dicker GD, Saling MM. The assessment of orientation following

concussion in athletes. Clinical journal of sport medicine: official journal of the Canadian

Academy of Sport Medicine. 1995;5(1):32-5.

15.   Echemendia RJ, Meeuwisse W, McCrory P, Davis GA, Putukian M, Leddy J, et al.

The sport concussion assessment tool 5th edition (SCAT5): background and rationale. British

journal of sports medicine. 2017;51(11):848-50.

16.   Davis GA, Purcell L, Schneider KJ, Yeates KO, Gioia GA, Anderson V, et al. The

child sport concussion assessment tool 5th edition (child SCAT5): background and rationale.

British journal of sports medicine. 2017;51(11):859-61.

17.   Costa SL, Genova HM, DeLuca J, Chiaravalloti ND. Information processing speed in

multiple sclerosis: Past, present, and future. Multiple Sclerosis Journal. 2016;23(6):772-89.

18.   DeLuca J, Chelune GJ, Tulsky DS, Lengenfelder J, Chiaravalloti ND. Is speed of

processing or working memory the primary information processing deficit in multiple

sclerosis? Journal of Clinical and Experimental Neuropsychology. 2004;26(4):550-62.

19.   Lu H, Chan SS, Lam LC. 'Two-level'measurements of processing speed as cognitive

markers in the differential diagnosis of DSM-5 mild neurocognitive disorders (NCD).

Scientific Reports. 2017;7(1):1-8.

20.   Collie A, Maruff P, Makdissi M, McCrory P, McStephen M, Darby D. CogSport:

reliability and correlation with conventional cognitive tests used in postconcussion medical

evaluations. Clinical Journal of Sport Medicine. 2003;13(1):28-32.

21.   Erlanger D, Saliba E, Barth J, Almquist J, Webright W, Freeman J. Monitoring

resolution of postconcussion symptoms in athletes: preliminary results of a web-based

neuropsychological test protocol. Journal of Athletic Training. 2001;36(3):280.

22.   Deary IJ, Liewald D, Nissan J. A free, easy-to-use, computer-based simple and four-

choice reaction time programme: The Deary-Liewald reaction time task. Behavior Research

Methods. 2011;43(1):258-68.

23.   Eckner JT, Kutcher JS, Richardson JK. Effect of concussion on clinically measured

reaction time in 9 NCAA division I collegiate athletes: a preliminary study. PM&R.

2011;3(3):212-8.

24.   Del Rossi G. Evaluating the recovery curve for clinically assessed reaction time after

concussion. Journal of Athletic Training. 2017;52(8):766-70.

25.   Tommerdahl M, Francisco E, Holden J, Lensch R, Tommerdahl A, Kirsch B, et al. An

accurate measure of reaction time can provide objective metrics of concussion. The Journal

of Science and Medicine. 2020;2(2).

26.   Kira J, Anna ES, Richard S, Alex K, John William O. Evaluation of CogSport for

acute concussion diagnosis in cricket. BMJ Open Sport & Exercise Medicine.

2021;7(2):e001061.

27.   Khurana VG, Kaye AH. An overview of concussion in sport. Journal of Clinical

Neuroscience. 2012;19(1):1-11.

28.   Makdissi M, Collie A, Maruff P, Darby D, Bush A, McCrory P, et al. Computerised

cognitive assessment of concussed Australian Rules footballers. British journal of sports

medicine. 2001;35(5):354-60.

29.   Schatz P, Pardini JE, Lovell MR, Collins MW, Podell K. Sensitivity and specificity of

the ImPACT Test Battery for concussion in athletes. Archives of clinical neuropsychology.

2006;21(1):91-9.

30.   Chen JK, Johnston KM, Collie A, McCrory P, Ptito A. A validation of the post

concussion symptom scale in the assessment of complex concussion using cognitive testing

and functional MRI. J Neurol Neurosurg Psychiatry. 2007;78(11):1231-8.

31.   Allen BJ, Gfeller JD. The Immediate Post-Concussion Assessment and Cognitive

Testing battery and traditional neuropsychological measures: a construct and concurrent

validity study. Brain Injury. 2011;25(2):179-91.

32.   Covassin T, Elbin RJ, Nakayama Y. Tracking Neurocognitive Performance following

Concussion in High School Athletes. The Physician and Sportsmedicine. 2010;38(4):87-93.

33.   Kalafatis C, Modarres MH, Apostolou P, Marefat H, Khanbagi M, Karimi H, et al.

Validity and cultural generalisability of a 5-minute ai-based, computerised cognitive

assessment in mild cognitive impairment and Alzheimer's dementia. Frontiers in Psychiatry.

2021;12.

34.   Khaligh-Razavi S, Habibi S. System for assessing mental health disorder. UK

Intellect Prop Off. 2013.

35.   Khaligh-Razavi S-M, Habibi S, Sadeghi M, Marefat H, Khanbagi M, Nabavi SM, et

al. Integrated cognitive assessment: Speed and accuracy of visual processing as a reliable

proxy to cognitive performance. Scientific Reports. 2019;9(1):1102.

36.   McInnes K, Friesen CL, MacKenzie DE, Westwood DA, Boe SG. Mild Traumatic

Brain Injury (mTBI) and chronic cognitive impairment: A scoping review. PLoS One.

2017;12(4):e0174847.

37.   Medicine ACoR. Definition of mild traumatic brain injury. Journal of Head Trauma

Rehabilitation. 1993;8(3):86-7.

38.   Khaligh-Razavi S-M, Sadeghi M, Khanbagi M, Kalafatis C, Nabavi SM. A self-

administered, artificial intelligence (AI) platform for cognitive assessment in multiple

sclerosis (MS). BMC neurology. 2020;20(1):1-13.

39.   Cichy RM, Pantazis D, Oliva A. Resolving human object recognition in space and

time. Nature neuroscience. 2014;17(3):455-62.

40.   Wickelgren WA. Speed-accuracy tradeoff and information processing dynamics. Acta

psychologica. 1977;41(1):67-85.

41.   Liesefeld HR, Janczyk M. Combining speed and accuracy to control for speed-

accuracy trade-offs. Behavior Research Methods. 2019;51(1):40-60.

42.   Townsend JT, Ashby FG. Stochastic Modeling of Elementary Psychological

Processes. New York, NY: Cambridge University Press; 1983.

43.   Glassbrook DJ, Chazot PL, Hind K. ICA Precision Raw Data.xlsx.: figshare; 2023.

44.   Straume-Naesheim TM, Andersen TE, Bahr R. Reproducibility of computer based

neuropsychological testing among Norwegian elite football players. British Journal of Sports

Medicine. 2005;39(suppl 1):i64.

45.   Machin D, Campbell MJ, Walters SJ. Medical Statistics a Textbook for the Health

Sciences: John Wiley & Sons, New York; 2007.

46.   Cohen J. Statistical power analysis. Current directions in psychological science.

1992;1(3):98-101.

47.   Cronbach LJ. Coefficient alpha and the internal structure of tests. Psychometrika.

1951;16(3):297-334.

48.   Poels MM, Ikram MA, Vernooij MW, Krestin GP, Hofman A, Messen WJ, et al.

Total cerebral blood flow in relation to cognitive function: The Rotterdam Scan Study.

Journal of Cerebral Blood Flow & Metabolism. 2008;28(10):1652-5.

49.   Davranche K, Audiffren M. Facilitating effects of exercise on information processing.

Journal of Sports Sciences. 2004;22(5):419-28.

50.   Lemmink KAPM, Visscher C. Effect of intermittent exercise on multiple-choice

reaction times of soccer players. Perceptual and Motor Skills. 2005;100(1):85-95.

51.   Niedermeier M, Weiss EM, Steidl-Müller L, Burtscher M, Kopp M. Acute effects of a

short bout of physical activity on cognitive function in sport students. International Journal of

Environmental Research and Public Health. 2020;17(10):3678.

52.   Baim S, Wilson CR, Lewiecki EM, Luckey MM, Downs Jr RW, Lentle BC. Precision

assessment and radiation safety for dual-energy X-ray absorptiometry: position paper of the

International Society for Clinical Densitometry. Journal of Clinical Densitometry.

2005;8(4):371-8.

432