Precision of the Integrated Cognitive Assessment for the assessment of neurocognitive

performance in athletes

Daniel J. Glassbrook*[a], Paul L. Chazot[a], and Karen Hind[a]

[a]Wolfson Research Institute for Health and Wellbeing, Durham University, Durham, United

Kingdom.

*__Corresponding author:__

Dr Daniel J. Glassbrook

Present Address:

School of Science, Technology and Health, York St John University

York St John University Sports Park, Haxby Rd, New Earswick, York YO31 8TA

Telephone: +61 4 7498980466 Email: d.glassbrook@yorksj.ac.uk

Funding: This study was funded by My Sports Wellbeing.

Conflicts of Interest/Competing Interests: The authors have no relevant financial or non-

financial interests to disclose.

Availability of data and materials: The datasets generated during and/or analysed during

the current study are available in the figshare repository,

https://figshare.com/articles/dataset/ICA_Precision_Raw_Data_xlsx/22211644

Abstract

Choice reaction time tests are commonly used for the assessment of cognitive function, and may be useful to assess the effect of sport participation. This study investigated the precision of the Integrated Cognitive Assessment (ICA; Cognetivity Neurosciences Ltd., Vancouver, Canada) test for the assessment of cognitive function in athletes. Thirty-one participants volunteered to take part in this study, from both contact ($n = 22$) and non-contact sports ($n = 9$). Participants performed the ICA test consecutively both before and after normal training session to simulate resting and post-sport conditions. Precision errors, relationships (Pearson's r), and internal consistency (Cronbach's Alpha) were calculated for three variables, ICA Index (overall information processing ability), ICA Speed (information processing speed) and ICA Accuracy (information processing accuracy). ICA precision errors [root mean squared-standard deviation, RMS-SD (coefficient of variation, %CV)] pre-sport were ICA Index: 5.18 (7.14%), ICA Speed: 3.98 (4.64%), and ICA Accuracy: 3.64 (5.00%); and post-sport were ICA Index: 3.96 (4.94%), ICA Speed: 2.14 (2.32%), and ICA Accuracy 3.40 (4.25%). The ICA test demonstrates high in-vivo precision with all variables except ICA Index (7.14%) demonstrating an acceptable precision error of ≤5% %CV. All variables demonstrated strong relationships between consecutive tests pre- and post-sport (r ≥ 0.8) except for the ICA Index post-sport which demonstrated a moderate (r ≥ 0.5) relationship. The ICA Index demonstrated good internal consistency (α ≥ 0.8) for both pre-and post-sport. The ICA Speed and ICA Accuracy variables demonstrated excellent internal consistency (α ≥ 0.9) for both pre-and post-sport. The ICA test is suitable for the assessment of cognitive function pre- and post-sport.

Key words: Information Processing, Reaction Time, Rapid Categorisation Task, Contact Sport, Non-Contact Sport.

1.0 Introduction

The brain plays a vital role in sports participation, and is responsible for decision making, and good decision making is key to success in sport [1]. Athletes are required to demonstrate high levels of technical and perceptual decision making ability, all while under pressure and fatigue [2]. Assessing cognitive function, such as information processing and decision making ability is a valuable tool which may quantify psychological ability in athletes. Additionally, assessing cognitive function may give insight into the effect of sport participation on the brain, for example through the effect on reaction time and decision making ability. There are many positive effects of sport and exercise participation, including cognitive benefits such as neuroplasticity, the process of adaptive structural and functional changes to the brain [3]. However, injury is an innate part of sport participation due to the physical nature of sport, and head impacts which may impair cognitive function are associated with participation in contact sports.

One proposed method to assess the effects of possible deleterious effects of head impacts sustained during normal sport participation, and cognitive function clinically is via information processing and reaction time tests. Reduced information processing ability, including memory, is linked with head impacts [4-6]. Indeed, altered mental state and confusion are also symptoms associated with head impacts that can negatively affect memory and information processing ability [7]. Additionally, information processing speed underpins several conditions of cognitive dysfunction, for example, multiple sclerosis [8, 9] and Alzheimer's disease [10].

Slower reaction time is a commonly used indicator of cognitive change following head impacts [11, 12]. Two common types of reaction test include measurement of simple reaction time (SRT) or choice reaction time (CRT) [13]. SRT is recorded when there is only one

possible stimulus (signal) and one possible response (action), for example tapping anywhere

on a screen when any image appears. In CRT tasks there are two or more possible stimuli,

each of which requires a quite different response, for example, tapping on the left of the

screen when an image of an object appears on the screen, and tapping on the right when an

image of an animal appears on the screen. SRT tests such as a weighted object drop and catch

[14], ruler drop test [15], and somatosensory assessment, where the participant held a device

similar to a computer mouse and reacted to a vibration applied to one or both the index or

middle finger, and responded by clicking with the finger (or both) that received the vibration

[16], have all been used to show slower reaction time in participants that have experienced

head impacts. Computerised tests have also been shown to be sensitive and able to determine

athletes who experienced head impacts versus those without, for example the CogSport

choice reaction time test and Immediate Post-Concussion Assessment and Cognitive Testing

(imPACT) [17-20]. The CogSport choice test takes approximately 20 minutes, and consists

of seven tasks including a SRT, CRT and complex reaction time task, as well as tests of

visual and working memory, all designed as card games [21]. The imPACT is performed

with a computer screen and mouse and typically takes 20-25 minutes and comprises of six

test modules assessing attention, memory, processing speed and reaction time, through tests

such as word memory and symbol matching [22]. Interestingly, reaction time has also been

used to show the lasting effect of head impacts on reaction time, with reaction time scores

only returning to baseline at 10 to 14 days post injury [15], and up to 21 days post injury [23].

The Integrated Cognitive Assessment (ICA; Cognetivity Neurosciences Ltd., Vancouver,

Canada) [24, 25], is a newly developed method for the assessment of cognitive function, and

may be applicable to the assessment of cognitive function in athletic populations. The ICA is

a short computerised cognitive test of cognitive function via an assessment of information

processing (CRT) speed based on a rapid categorisation task [24, 26]. The ICA test can be

completed in a short amount of time (~5 minutes), on a handheld device such as an iPhone or

iPad, and is therefore able to be administered easily pre- and post-sport. Performing an

assessment of cognitive function such as the ICA pre-sport in a rested state effectively

provides a baseline measure of cognitive function. Performing a test of cognitive function

post-sport then can be compared to the baseline measure of pre-sport and determine the effect

of sport participation on cognitive function. Additionally, the ICA has been shown to

accurately detect mild cognitive impairment and be moderately to highly correlated with

other popular pen-and-paper cognitive tests such as the Montreal Cognitive Assessment

(Pearsons r = 0.58) and Addenbrooke's Cognitive Examination (Pearsons r = 0.62) cognitive

tests [24].

The ICA has been shown to accurately measure cognitive impairment in patients in the early

stages of dementia [24]. However, to date, no known study has investigated the intra-day

precision of the ICA test. Therefore, the purpose of this study was to determine the same-day,

in-vivo (i.e., within whole living organisms; within people) precision of the ICA test to assess

cognitive function. It is hypothesised that the ICA will demonstrate acceptable precision, and

suitability for the assessment of cognitive function pre- and post-sport.

2.0 Materials and Methods

2.1 Participants

Thirty-one participants volunteered to take part in this study. The minimum number of

participants required for precision analysis with two consecutive tests is thirty [27].

Participant characteristics are presented in Table 1. Participants were recruited from March to

May 2022, from the University sporting facility via advertisement flyer, and from the wider

sporting community through existing relationships with sporting clubs. After initially

expressing interest to take part in the study, participants were provided with an information

sheet and given at least 24 hours to consider their participation in the study before a follow up

was sent by the lead researcher. After confirmation of eligibility, participants were then

booked in at a time and date of their convenience for data collection. Participants were

eligible for participation if they were a current contact sport or non-contact sport athlete

(Table 2) aged 18-40 years, and healthy; having no underlying medical issues that affect

participation in sport. Participants were excluded if they were injured, pregnant, or suffering

from post-concussion syndrome. Post-concussion syndrome is defined as the collective

experience of symptoms after sustaining a concussion, such as headaches, fatigue,

depression, anxiety and irritability [28, 29]. This study was approved by the Durham

University Sport and Exercise Sciences Ethics Committee (reference: SPORT-2022-01-

07T10_44_59-srhd22;), and written informed consent was provided by each participant prior

to participation. Participant characteristics, demographics and athletic history were collected

via standardised questionnaire directly after consent to participate was obtained.

2.2 Procedures

To simulate resting- and post-sport conditions, participants performed the ICA test (version

1.6.0 or 1.7.0) before and after a normal training session for their respective sports. Data

collection was performed in a quiet room to minimise distractions, with a maximum of two

participants (separated within the room) performing the test at a time. Prior to their sports

training, and within an hour of the training start time, participants completed two consecutive

ICA tests. The participants then completed a normal training session and then two

consecutive ICA tests again, within an hour of the training finishing.

2.3 The Integrated Cognitive Assessment (ICA)

The ICA has been described in detail in published works [26, 30]. In summary, the ICA is a

short computerised cognitive test of cognitive function via an assessment of information

processing (CRT) speed based on a rapid categorisation task, and in contrast to the CogSport

and imPACT tests is independent of language or colour. The ICA in this study was

completed on an iPad, with each test taking approximately five minutes, considerably shorter

than the 20+ minute CogSport and imPACT tests; the imPACT test of which is also not

portable (completed on a computer with screen and mouse). Each ICA test utilises the brains

powerful reaction to animal stimulus [31] and comprises of 100 natural images, with 10

additional practice images provided in a practice section prior to the test. The practice images

are not included in the test score. The images are a mixture of animals (e.g., birds, fish,

mammals), or non-animals (e.g., objects, food, vehicles), and are presented in rapid

succession. Images appear for 100ms, followed by a 20ms inter-stimulus interval and a

250ms dynamic noisy mask. Participants react to the image by tapping with their thumb on

the screen; on the right-hand side to select 'animal' and on the left-hand side of the screen to

select 'non-animal'. Participants were instructed to perform the test as quick as possible, and

these instructions were given both verbally and presented on the screen. There are no

associated risks with completing the ICA, and the test provides three variables: ICA Speed,

the response reaction times in trials they responded correctly; ICA Accuracy, choice

accuracy, the percentage of correct test responses; and ICA Index, overall information

processing ability, a combination of ICA Speed and ICA Accuracy. Each variable is

calculated by the following equations [24, 30]:

$$Accuracy = \frac{Number\ of\ correct\ categorisations}{Total\ number\ of\ images}\ x\ 100$$

$$Speed = \min\left[100, 100e^{-\frac{mean\ correct\ reaction\ time}{1025}+0.341}\right]$$

$$ICA\ Index = \left(\frac{Speed}{100}\ x\ \frac{Accuracy}{100}\right)\ x\ 100$$

There is a speed-accuracy trade-off in reaction test performance, and often scoring higher in

either speed or accuracy is achieved at the expense of the other capacity [32, 33]. To combat

the potential negative reflection on overall information processing ability from a poor speed

or accuracy score, a common solution is the inverse efficiency score [34], whereby speed and

accuracy are combined into a single score. In the case of the ICA, this concept is applied and

manifests as the ICA Index variable. Test results are recorded via report immediately after

test completion.

ICA Accuracy, ICA Speed and ICA Index are represented by Arbitrary Units and results of

each equation are transformed and scored by a standardised scale ranging from 0 to 100. I.e.,

a Speed score closer to 100 indicates faster reaction times, while a Speed score closer to 0

indicates slower reaction times; an Accuracy score closer to 100 indicates higher accuracy,

and an Accuracy score closer to 0 indicates lower accuracy; and an ICA Index score closer to

100 represents greater overall information processing ability, whereas an ICA Index score

closer to 0 indicates poorer overall information processing ability. Additionally, the specific

components included in the Speed equation (i.e., division by 1025 and addition of 0.341)

further serve to standardise and scale the Speed score. These additional components are

determined empirically to ensure that the Speed scores fall within the desired range of 0 to

100 for the majority of participants. Natural Log transformation further aids in normalizing

the distribution of scores.

2.4 Statistical Analysis

All data analysis was performed in Microsoft Excel (2016). Raw anonymised data for ICA

Index, ICA Speed, and ICA Accuracy were extracted and exported to Microsoft Excel for

analysis [35]. Throughout the study the lead author had access to information that could

identify individual participants, however, all analysis was performed on anonymised data.

Precision has been previously used to define test suitability in similar cognitive tests to the

ICA, such as the CogSport choice reaction time test [36]. Precision of ICA scores and least

significant change (LSC) were calculated at the 95% confidence level. Precision was

determined as root mean square standard deviation (RMS-SD), coefficient of variation (CV),

and percentage CV (%CV). RMS-SD represents the sample standard deviation of the

differences between predicted values and observed values, and is calculated via the following

formulae, where SD represents standard deviation and n represents the number of

participants:

$$\sqrt{(\frac{\Sigma SD^2}{n})}$$

The %CV expresses test variation relative to the mean of two tests and is corrected for small

sample bias, and was defined as acceptable <5% [37]. The LSC represents a true meaningful

change was calculated from the precision errors (LSC = RMS-SD * 2.77). Pearson's

correlation coefficient was calculated for each variable to determine reliability and the

strength of the relationship between consecutive tests, with 0.2, 0.5, and 0.8 representing

weak, moderate, and strong correlations, respectively [38]. Cronbach's Alpha was calculated

for each variable to determine internal consistency, with 0.9, 0.8, 0.7, 0.6, 0.5 and < 0.5

representing excellent, good, acceptable, questionable, poor and unacceptable internal

consistency, respectively [39].

3.0 Results

Results of the precision analysis for each ICA variable pre- and post-sport are presented in

Table 3. All variables except for ICA Index pre-sport had a precision error of ≤5% %CV.

LSC results are presented in Table 4. Pearson correlation coefficient data is presented in

Error! Reference source not found.. All variables demonstrated strong relationships

between consecutive tests pre- and post-sport, except for the ICA Index post-sport which

demonstrated a moderate relationship. Cronbach's Alpha data is presented in Table 5. The

ICA Index demonstrated good internal consistency for both pre-and post-sport. The Speed

and Accuracy variables demonstrated excellent internal consistency for both pre-and post-

sport.

217 4.0 Discussion

The purpose of this study was to determine the same-day, in-vivo precision of the ICA test to

assess cognitive function. The results of this study support the ICA as a tool with acceptable

precision to measure changes in cognitive ability pre- and post-sport, confirming the

hypothesis. All ICA variables in this study, except for ICA Index pre-sport demonstrated a

precision error of ≤5% %CV. All variables demonstrated moderate to strong relationships

and good to excellent internal consistency between consecutive tests pre- and post-sport. The

ICA may be clinically relevant for the assessment of cognitive ability, and the effect of sport

on cognitive ability.

The higher ICA Index precision score (7.14 %CV) pre-sport than post-sport (4.94 %CV) in

this study may be explained by the larger average difference between test one and test two

pre-sport (76.6 ± 8.3 to 82.2 ± 6.1), compared to a smaller average difference in ICA Index

scores between test one and test two post-sport (81.5 ± 7.0 to 83.0 ± 6.7). This is further

exemplified by a larger ICA Index RMS-SD pre-sport than post-sport (5.18 and 3.96,

respectively), which indicates more variance in observed data around the mean. This result

may be due to an increased level of comfort with the test from the first pre-sport ICA test to

the subsequent test, and possibly a learning effect. However, this is in contrast to previous

work which showed no learning effect for the ICA test in healthy participants and those

diagnosed with dementia [24]. Additionally, a strong relationship (r = 0.82) and good internal

consistency (α = 0.88) between ICA tests and the ICA Index variable was observed pre-sport

(Test 1 and Test 2; Table 5), further suggesting a minimal learning effect, and supporting the

test as reliable.

All variables showed greater precision post-sport compared to pre-sport. This may be due to

the many positive physiological benefits that exercise has, such as an increase in blood flow

to muscles and brain [40], structural and functional changes in the brain [3], and increases

information processing ability [41]. Indeed, improvements in cognitive function after a bout

of exercise is supported by previous research [42, 43].

Previous research looking at precision in a similar cognitive test to the ICA, the CogSport

choice reaction time test, has shown lower %CV for mean choice reaction time (speed) (1.4

%CV), and higher %CV for choice reaction time accuracy (11.4 %CV) [36]. These results

are interesting as the ICA is shown to be less precise in measuring reaction speed (2.32 - 4.64

%CV vs 1.4 %CV), however, the ICA test is shown to be more precise in terms of accuracy

(4.25 – 5.00 %CV vs 11.4 %CV). These results may indicate that the test you adopt needs to

be specific to the variable of interest (i.e., speed or accuracy), however, this should be

negated in the case of the ICA via the ICA Index variable as an inverse efficiency score [34],

whereby speed and accuracy are combined into a single score. The contrasting results

between the present study and that of Straume-Naesheim, Andersen and Bahr [36] may be

due to the populations used; the CogSport test was used in elite football players only,

whereas only a small percentage of the participants in the present study are practicing

professionally (table 2). The present study recruited participants from a variety of sports, each

with their own decision making and reaction characteristics, in comparison to only football.

Additionally, a greater number of participants ($n = 289$) performed the CogSport test in

comparison to the ICA in the present study ($n = 31$). However, the administration procedure

of the CogSport test was similar to the administration of the ICA test, and was also

administered under controlled conditions, i.e., in a quiet room to minimise distractions and

with more than one participant performing the test at a time.

A limitation to this study is that information about the specific duration an intensity of

training sessions undergone by participants was not recorded. Only that it was a typical or

normal training session. i.e., not a session of just running or conditioning, and included

practice of the sport itself. Additionally, the optimal intensity and duration of exercise that

may impact cognitive function was not assessed. A relatively small sample size was used,

although the sample size exceeds the minimum of 30 participants required for precision

analysis with two consecutive tests [27], a greater number of participants could strengthen the

statistical analysis. Future studies should look to recruit larger numbers, and complete

comparisons between contact and non-contact sports, and between amateur and professional

level participants.

In conclusion, the ICA is a practical test which can be used to measure cognitive function

before and after sport participation. The results of this study support the ICA as a precise (as

determined by %CV values) measure of information processing speed and information

processing accuracy, and overall information processing ability (ICA Index). The ICA can be

used for the assessment of cognitive function, and may be useful as a method to assess the

effects of sport on cognitive function.

5.0 References

[1] T.J. Hepler, Decision-making in sport under mental and physical stress, International

Journal of Kinesiology and Sports Science 3(4) (2015) 79-83.

[2] T.J. Gabbett, J. Carius, M. Mulvey, Does improved decision-making ability reduce the

physiological demands of game-based activities in field sport athletes?, The Journal of

Strength & Conditioning Research 22(6) (2008) 2027-2035.

285 [3] J. Fernandes, R.M. Arida, F. Gomez-Pinilla, Physical exercise as an epigenetic modulator

286 of brain plasticity and cognition, Neuroscience & Biobehavioral Reviews 80 (2017) 443-456.

287 [4] D.M. Bernstein, Information processing difficulty long after self-reported concussion,

288 Journal of the International Neuropsychological Society 8(5) (2002) 673-682.

289 [5] D. Gronwall, P. Wrightson, Memory and information processing capacity after closed

290 head injury, Journal of Neurology, Neurosurgery & Psychiatry 44(10) (1981) 889-895.

291 [6] J.R. O'Jile, L.M. Ryan, B. Betz, J. Parks-Levy, R.C. Hilsabeck, J.L. Rhudy, W.D.

292 Gouvier, Information processing following mild head injury, Archives of clinical

293 neuropsychology 21(4) (2006) 293-296.

294 [7] A. Sharma, K. Hind, P. Hume, J. Singh, J.P. Neary, Neurovascular coupling by functional

295 near infra-red spectroscopy and sport-related concussion in retired rugby players: The UK

296 rugby health project, Frontiers in Human Neuroscience 14 (2020).

297 [8] S.L. Costa, H.M. Genova, J. DeLuca, N.D. Chiaravalloti, Information processing speed in

298 multiple sclerosis: Past, present, and future, Multiple Sclerosis Journal 23(6) (2016) 772-789.

299 [9] J. DeLuca, G.J. Chelune, D.S. Tulsky, J. Lengenfelder, N.D. Chiaravalloti, Is speed of

300 processing or working memory the primary information processing deficit in multiple

301 sclerosis?, Journal of Clinical and Experimental Neuropsychology 26(4) (2004) 550-562.

302 [10] H. Lu, S.S. Chan, L.C. Lam, 'Two-level' measurements of processing speed as cognitive

303 markers in the differential diagnosis of DSM-5 mild neurocognitive disorders (NCD),

304 Scientific Reports 7(1) (2017) 1-8.

305 [11] A. Collie, P. Maruff, M. Makdissi, P. McCrory, M. McStephen, D. Darby, CogSport:

306 reliability and correlation with conventional cognitive tests used in postconcussion medical

307 evaluations, Clinical Journal of Sport Medicine 13(1) (2003) 28-32.

[12] D. Erlanger, E. Saliba, J. Barth, J. Almquist, W. Webright, J. Freeman, Monitoring

resolution of postconcussion symptoms in athletes: preliminary results of a web-based

neuropsychological test protocol, Journal of Athletic Training 36(3) (2001) 280.

[13] I.J. Deary, D. Liewald, J. Nissan, A free, easy-to-use, computer-based simple and four-

choice reaction time programme: The Deary-Liewald reaction time task, Behavior Research

Methods 43(1) (2011) 258-268.

[14] J.T. Eckner, J.S. Kutcher, J.K. Richardson, Effect of concussion on clinically measured

reaction time in 9 NCAA division I collegiate athletes: a preliminary study, PM&R 3(3)

(2011) 212-218.

[15] G. Del Rossi, Evaluating the recovery curve for clinically assessed reaction time after

concussion, Journal of Athletic Training 52(8) (2017) 766-770.

[16] M. Tommerdahl, E. Francisco, J. Holden, R. Lensch, A. Tommerdahl, B. Kirsch, R.

Dennis, O. Favorov, An accurate measure of reaction time can provide objective metrics of

concussion, The Journal of Science and Medicine 2(2) (2020).

[17] J. Kira, E.S. Anna, S. Richard, K. Alex, O. John William, Evaluation of CogSport for

acute concussion diagnosis in cricket, BMJ Open Sport & Exercise Medicine 7(2)

(2021) e001061.

[18] V.G. Khurana, A.H. Kaye, An overview of concussion in sport, Journal of Clinical

Neuroscience 19(1) (2012) 1-11.

[19] M. Makdissi, A. Collie, P. Maruff, D. Darby, A. Bush, P. McCrory, K. Bennell,

Computerised cognitive assessment of concussed Australian Rules footballers, British journal

of sports medicine 35(5) (2001) 354-360.

[20] P. Schatz, J.E. Pardini, M.R. Lovell, M.W. Collins, K. Podell, Sensitivity and specificity

of the ImPACT Test Battery for concussion in athletes, Archives of clinical neuropsychology

21(1) (2006) 91-99.

[21] J.K. Chen, K.M. Johnston, A. Collie, P. McCrory, A. Ptito, A validation of the post

concussion symptom scale in the assessment of complex concussion using cognitive testing

and functional MRI, J Neurol Neurosurg Psychiatry 78(11) (2007) 1231-8.

[22] B.J. Allen, J.D. Gfeller, The Immediate Post-Concussion Assessment and Cognitive

Testing battery and traditional neuropsychological measures: a construct and concurrent

validity study, Brain Injury 25(2) (2011) 179-191.

[23] T. Covassin, R.J. Elbin, Y. Nakayama, Tracking Neurocognitive Performance following

Concussion in High School Athletes, The Physician and Sportsmedicine 38(4) (2010) 87-93.

[24] C. Kalafatis, M.H. Modarres, P. Apostolou, H. Marefat, M. Khanbagi, H. Karimi, Z.

Vahabi, D. Aarsland, S.-M. Khaligh-Razavi, Validity and cultural generalisability of a 5-

minute ai-based, computerised cognitive assessment in mild cognitive impairment and

Alzheimer's dementia, Frontiers in Psychiatry 12 (2021).

[25] S. Khaligh-Razavi, S. Habibi, System for assessing mental health disorder, UK Intellect.

Prop. Off (2013).

[26] S.-M. Khaligh-Razavi, S. Habibi, M. Sadeghi, H. Marefat, M. Khanbagi, S.M. Nabavi,

E. Sadeghi, C. Kalafatis, Integrated cognitive assessment: Speed and accuracy of visual

processing as a reliable proxy to cognitive performance, Scientific Reports 9(1) (2019) 1102.

[27] S. Baim, C.R. Wilson, E.M. Lewiecki, M.M. Luckey, R.W. Downs Jr, B.C. Lentle,

Precision assessment and radiation safety for dual-energy X-ray absorptiometry: position

paper of the International Society for Clinical Densitometry, Journal of Clinical Densitometry

8(4) (2005) 371-378.

[28] K. McInnes, C.L. Friesen, D.E. MacKenzie, D.A. Westwood, S.G. Boe, Mild Traumatic

Brain Injury (mTBI) and chronic cognitive impairment: A scoping review, PLoS One 12(4)

(2017) e0174847.

[29] A.C.o.R. Medicine, Definition of mild traumatic brain injury, Journal of Head Trauma

Rehabilitation 8(3) (1993) 86-87.

[30] S.-M. Khaligh-Razavi, M. Sadeghi, M. Khanbagi, C. Kalafatis, S.M. Nabavi, A self-

administered, artificial intelligence (AI) platform for cognitive assessment in multiple

sclerosis (MS), BMC neurology 20(1) (2020) 1-13.

[31] R.M. Cichy, D. Pantazis, A. Oliva, Resolving human object recognition in space and

time, Nature neuroscience 17(3) (2014) 455-462.

[32] W.A. Wickelgren, Speed-accuracy tradeoff and information processing dynamics, Acta

psychologica 41(1) (1977) 67-85.

[33] H.R. Liesefeld, M. Janczyk, Combining speed and accuracy to control for speed-

accuracy trade-offs, Behavior Research Methods 51(1) (2019) 40-60.

[34] J.T. Townsend, F.G. Ashby, Stochastic Modeling of Elementary Psychological

Processes, Cambridge University Press, New York, NY, 1983.

[35] D.J. Glassbrook, P.L. Chazot, K. Hind, ICA Precision Raw Data.xlsx., figshare, 2023.

[36] T.M. Straume-Naesheim, T.E. Andersen, R. Bahr, Reproducibility of computer based

neuropsychological testing among Norwegian elite football players, British Journal of Sports

Medicine 39(suppl 1) (2005) i64.

[37] D. Machin, M.J. Campbell, S.J. Walters, Medical Statistics a Textbook for the Health

Sciences, John Wiley & Sons, New York2007.

[38] J. Cohen, Statistical power analysis, Current directions in psychological science 1(3)

(1992) 98-101.

[39] L.J. Cronbach, Coefficient alpha and the internal structure of tests, Psychometrika 16(3)

(1951) 297-334.

[40] M.M. Poels, M.A. Ikram, M.W. Vernooij, G.P. Krestin, A. Hofman, W.J. Messen, A.

Van der Lugt, M.M. Breteler, Total cerebral blood flow in relation to cognitive function: The

Rotterdam Scan Study, Journal of Cerebral Blood Flow & Metabolism 28(10) (2008) 1652-

1655.

[41] K. Davranche, M. Audiffren, Facilitating effects of exercise on information processing,

Journal of Sports Sciences 22(5) (2004) 419-428.

[42] K.A.P.M. Lemmink, C. Visscher, Effect of intermittent exercise on multiple-choice

reaction times of soccer players, Perceptual and Motor Skills 100(1) (2005) 85-95.

[43] M. Niedermeier, E.M. Weiss, L. Steidl-Müller, M. Burtscher, M. Kopp, Acute effects of

a short bout of physical activity on cognitive function in sport students, International Journal

of Environmental Research and Public Health 17(10) (2020) 3678.

Table 1: Participant Characteristics

	Age (Yr.)	Height (m)	Body Mass (kg)
Total ($n = 31$)	23.7 ± 5.7	1.78 ± 0.09	72.6 ± 8.3
Male ($n = 16$)	22.9 ± 4.7	1.82 ± 0.08	75.7 ± 7.4
Female ($n = 15$)	24.6 ± 6.6	1.71 ± 0.07	68.0 ± 9.9
Contact Sport ($n = 22$)	24.9 ± 6.3	1.80 ± 0.07	75.5 ± 6.0
Non-Contact Sport ($n = 9$)	20.6 ± 0.6	174.1 ± 8.3	77.2 ± 12.2

Data are presented as mean ± standard deviation. Yr., Years; m, Metres; kg, Kilograms; n, Number.

396   Table 2: Sports Breakdown

Contact Sports ($n = 22$)		Non-Contact Sports ($n = 9$)	
Rugby Union ($n = 7$)	Semi-professional, Amateur	Touch Rugby ($n = 5$)	Amateur
Boxing ($n = 6$)	Amateur	Athletics ($n = 4$)	Amateur
Muay Thai (Kickboxing) ($n = 5$)	Professional, Amateur		
Indoor Football ($n = 4$)	Amateur		

n, Number.

399   Table 3: Precision Analysis Results

Variable	Precision ($n = 31$)		
	RMS-SD	CV	%CV
Pre			
ICA Index	5.18	0.07	7.14
ICA Speed	3.98	0.05	4.64
ICA Accuracy	3.64	0.05	5.00
Post			
ICA Index	3.96	0.05	4.94
ICA Speed	2.14	0.02	2.32
ICA Accuracy	3.40	0.04	4.25

n, Number; ICA, Integrated Cognitive Assessment; RMS-SD, Root Mean Square Standard Deviation; CV, Coefficient of Variation; %, Percentage.

402 Table 4: Least Significant Change Results

Variable	LSC ($n = 31$)		
	RMS-SD	CV	%CV
Pre			
ICA Index	14.36	0.20	19.78
ICA Speed	11.01	0.13	12.86
ICA Accuracy	10.09	0.14	13.9
Post			
ICA Index	10.96	0.14	13.7
ICA Speed	5.94	0.06	6.43
ICA Accuracy	9.43	0.12	11.78

LSC, Least Significant Change; n, Number; ICA, Integrated Cognitive Assessment; RMS-SD, Root Mean Square Standard Deviation; CV, Coefficient of Variation; %, Percentage.

405 Table 5: Relationships between consecutive tests

Variable	Test 1	Test 2	Correlation (r)	Cronbach's Alpha (α)
Pre				
ICA Index	76.6 ± 8.3	82.2 ± 6.1	0.82	0.88
ICA Speed	88.6 ± 8.1	92.3 ± 7.3	0.85	0.91
ICA Accuracy	86.9 ± 8.9	89.2 ± 6.2	0.87	0.90
Post				
ICA Index	81.5 ± 7.0	83.0 ± 6.7	0.68	0.81
ICA Speed	94.8 ± 5.2	95.6 ± 4.6	0.82	0.90
ICA Accuracy	86.1 ± 9.1	87.0 ± 8.3	0.85	0.92

n, Number; ICA, Integrated Cognitive Assessment; Data are
presented as mean ± standard deviation.