Group A *Streptococcus* strains causing meningitis without distinct invasive phenotype

Laura Marquardt[a][*], Federica Andreoni[a][*], Mathilde Boumasmoud[a], Tiziano A. Schweizer[a],

Dorothea M. Heuberger[b], Elena Parietti[a], Jana Epprecht[a], Dario Mattle[a], Anna K. Raez[a],

Ewerton Marques-Maggio[c]; Reto A. Schuepbach[b], Barbara Hasse[a], Srikanth Mairpady-

Shambat[a], Silvio D. Brugger[a], Annelies S. Zinkernagel[a][**]

[a]Departement of Infectious Diseases and Hospital Epidemiology, University Hospital Zurich,

University of Zurich, Zurich, Switzerland

[b]Institute for Intensive Care Medicine, University Hospital Zurich, University of Zurich, Zurich,

Switzerland

[c]Division of Clinical Pathology, University Hospital Zurich, University of Zurich, Zurich,

Switzerland.

*These authors contributed equally to this manuscript

**Corresponding author

<u>**Keywords**</u>

Group A Streptococcus; meningitis; virulence determinants

<u>**Running title**</u>

Group A Streptococcus in meningitis

Corresponding author

Prof. Dr. Dr. Med. Annelies Zinkernagel

University Hospital Zurich

Raemistrasse 100

8091, Zurich

Tel. +41 44 255 12 59

annelies.zinkernagel@usz.ch

Alternative corresponding author

Dr. Federica Andreoni

University Hospital Zurich

Raemistrasse 100

8091, Zurich

Tel. +41 44 255 12 58

Abstract

Group A streptococcal (GAS) meningitis is a fulminant disease associated with high

morbidity and mortality. To elucidate the mechanisms underlying the invasiveness of GAS in

meningitis, we compared GAS isolates derived from five cases of meningitis, to otitis and

colonizing isolates. We did not observe differences in adherence to and invasion of human

brain microvascular endothelial cells, virulence factors activity or barrier disruption. Whole

genome sequencing did not reveal particular invasiveness traits. Most patients previously

suffered from otitis media suggesting that meningitis likely resulted from a continuous

spread of the infection rather than being attributable to changes in pathogen's virulence.

<u>Introduction</u>

Streptococcus pyogenes or group A *Streptococcus* (GAS) asymptomatically colonizes the

human throat and skin. In addition to asymptomatic colonization, GAS causes mild to life-

threatening infections, such as necrotizing fasciitis and meningitis. GAS meningitis cases are

rare and represent only 1-4% of all invasive GAS cases detected across Europe and the

United States [1-3]. GAS meningitis is a fulminant disease associated with high morbidity,

including neurologic sequelae, and mortality. It occurs mainly in patients with predisposing

factors such as otitis, sinusitis, recent head injury or neurosurgery, presence of a

neurosurgical device, cerebrospinal fluid leak, or alcoholism [4, 5]. This suggests extension

per continuitatem from a neighboring infection such as otitis media, resulting in otogenic

meningitis. However, in a third of cases there are no predisposing factors present.

Pathogen components such as streptococcal virulence factors Streptolysin O (SLO), DNase

Sda1, protease SpyCEP and the M-protein, play a key role in the invasiveness and ability of

the pathogen to survive within the host environment [6].

In this study, we investigated five patients suffering from GAS meningitis, analyzing both

the host's and pathogen's characteristics. On the host side, various clinical parameters were

analyzed. On the pathogen side, aiming to identify particular pathogenic traits, we compared

phenotypically and genetically five GAS isolates derived from meningitis cases with ten GAS

isolates derived either from otitis or asymptomatic colonization.

Material and methods

Patients and ethics

Five patients diagnosed with clinically confirmed community-acquired GAS meningitis between 2013 and 2017 in Switzerland were retrospectively included in this study (Tab.S1). No specific surveillance program for GAS meningitis exists in Switzerland and the cases were selected using a convenience sampling strategy. This study was approved by the regional committee for medical research ethics (BASEC-ID 2016-00145 and 2017-02225).

Bacterial strains and cell lines

All meningitis, otitis and colonizing GAS clinical isolates are listed in Tab.S2 and were cultured in THY (Todd-Hewitt broth (BD) plus 2% yeast extract (Oxoid)) at 37°C. Immortalized human brain microvascular endothelial cells (HBMECs) were cultured as previously described [7].

Adherence, invasion and viability assays

Adherence and invasion assays were performed as previously described [8], except lysis was carried out using 500 µl deionized water (dH$_2$O). Cell viability was assessed as previously described [9].

Virulence factors activity

Activity of the virulence factors SLO, Sda1 and SpyCEP was assessed as previously described [10].

Electric cell-substrate impedance sensing (ECIS®) assay

ECIS® was performed with the ECIS® Z-Theta instrument (Applied Biophysics) to monitor

the effects of the GAS strains on human brain microvascular endothelial cells (HBMECs) [7]

barrier integrity. Before seeding, the wells of the arrays (ECIS 8W10E+, IBIDI) were washed

twice with sterile, nuclease-free dH_2O, coated with 5 µg/ml collagen IV for 1h at room

temperature (RT), washed three times with dH_2O and incubated with 10 mM L-cysteine for

2h at RT. The wells were washed two times with dH_2O before 200 µl of culture medium were

added and the assay started (37ºC, 5% CO2). After 30 min, 200µl culture medium containing

10^5 HBMECs were seeded per well. After 22 h incubation 50 µl of medium were removed

and 2h later HBMECs were infected with log phase GAS strains resuspended in

DMEM+10%FCS at an MOI of 10. After 1.5h penicillin (10 µg/ml) and gentamycin (100 µg/ml)

were added. Barrier integrity was monitored at 4000 Hz from seeding to 70h post infection.

Resistance data were normalized to the time of infection.

Bacterial genomics

Whole genome sequencing and genomic data analysis were performed as previously

described [11]. Briefly, from 150 bp paired end reads, *de novo* assemblies were generated

with SPAdes v.3.10[11] and annotated with Prokka v.1.13 [12]. A pangenome was

constructed with Roary v.3.12 [13] and the resulting core-gene alignment was used to build

a maximum likelihood phylogenetic tree with Fasttree v.2.1.10 [14]. *Emm*-typing was

performed *in-silico* by querying assemblies against the curated CDC database[15] with blastn

v.2.9.0. Virulence genes were identified by querying the assemblies against the core

sequences of the VFDB [16], using ABRIcate v.0.5 with the parameter -minid 85. ENA

accession number: PRJEB57816.

Results

Cases description

Five GAS meningitis cases, occurring between 2013 and 2017 in Switzerland, were

included in this study and the corresponding GAS isolates were retrospectively retrieved for

analysis. The median age of the patients was 64 years (42 to 79 years) and 60% (n=3) were

female. The mortality rate was 20% (n=1), neurological sequelae included slight hearing loss

and vertigo. The median duration of the hospital stay was 17 days (3 to 29 days) and 4

patients needed intensive care treatment with endotracheal intubation. Relevant clinical and

laboratory findings are depicted in table 1. Mastoiditis was the underlying diagnosis in three

cases. During the course of the disease one patient had a septic thrombosis and one

developed a brain abscess (Fig.1A-1B). Surgical treatment such as antrotomy,

mastoidectomy, or burr-hole trepanation was required in three patients. In patient 4,

histological analysis from the surgically removed mastoid tissue showed necrotic tissue and

individual mucosal fragments covered by simple cuboid epithelium with severe acute and

chronic inflammation and showing extracellular accumulation of bacterial cells (Fig.1C-1D).

Bacterial virulence

A detailed analysis of streptococcal virulence factors, adherence to and invasion of

HBMECs as well as barrier integrity of HBMECs upon GAS challenge was carried out on

meningitis, otitis and colonizing clinical isolates. We found no significant difference between

adherence and invasion behavior to HBMECs upon infection with meningitis, otitis or

colonizing GAS isolates (Fig.2A-2B). Furthermore, no differences in virulence factors activity

or the isolates potential to disrupt the barrier formed by HBMECs among the three groups

were observed (Fig.2C-2F). Typing showed a high prevalence of *emm*-1/ST28 (n=4/5, Tab.S1).

This was also the most common genotype among otitis and colonizing isolates (n=5/5 and

4/5 respectively, Tab.S1). Only one meningitis clinical isolate belonged to the *emm*-28/ST52

and one colonizing isolate to the *emm*-28/ST458. The meningitis isolates did not form a

monophyletic group (Fig.S1A-S1B). The presence of genes for virulence factors determinants

was analyzed and this showed no specific pattern for meningitis isolates when compared to

otitis or colonizing ones (Fig. S1C).

Discussion

We found no particular pathogenic characteristics in the five GAS isolates causing meningitis. The only common denominator was preceding otitis and mastoiditis on the host's side. GAS meningitis is an extremely rare condition, representing only the 0.2%-0.9% of cases of bacterial meningitis in the United States and Brazil [17-19] and 1-4% of invasive GAS cases in Europe and the United States [1-3]. A report from Switzerland described nine cases of GAS meningitis registered between 1983 and 1999 across all major Swiss hospitals, again emphasizing the rarity of this disease [20].

Meningitis occurs either by hematogenous spread or by direct spread of bacteria from a neighboring infection. Hematogenous spread normally occurs when colonizing bacteria breach the nasopharyngeal mucosa, reach the blood [21] and subsequently cross the blood-brain barrier. Spread from the upper respiratory tract's mucosa may be due to an altered balance in the host microbe population, possibly favoring GAS number increase [5]. On the other hand, direct spread to the brain occurs *per continuitatem* from infected adjacent anatomic structures, such as the ear or the mastoid bone [21].

We searched for common traits in the clinical presentation of our patients and characteristics shared by the clinical isolates to find possible explanations for the spreading of GAS to the brain. In our patients' cohort, the presence of mastoiditis in four patients and thrombosis in one supports the idea of direct spread through bone erosion from coalescent oto-mastoiditis and osteo-thrombophlebitis. Moreover otitis, a condition often associated with GAS meningitis [5], was present in the majority of cases suggesting continuous spread as a likely cause of infection. In contrast, hematogenous spreading was less common and only observed in one patient (patient 3). The presence of a local focus of infection may explain the high surgery rate in our GAS meningitis cohort, which has been reported

previously. In a nationwide cohort study in the Netherlands, 46% of GAS meningitis patients

required surgical intervention, while this remains an absolute rarity only described in case

reports for the more common meningitis pathogens (pneumococcus, meningococcus) [4,

22].

On the pathogen's side, no common features explaining bacterial spread to the brain

were found among the meningitis isolates. This points to host rather than pathogen factors

playing a role in facilitating bacterial invasion, as previously described for *S. aureus* [11].

Adherence and invasion assays towards HBMECs confirmed the absence of a particular

tropism towards brain microvascular tissue for the meningitis isolates, excluding an

advantage in crossing or damaging the blood brain barrier for these isolates, as compared to

otitis and colonizing ones, in the tested settings. Moreover, virulence factors activity did not

significantly vary among the three groups of isolates. Isolates from the *emm*-1 and *emm*-28

lineage represented both in the meningitis and colonizing groups, behaved very similarly in

terms of virulence factors activity.

In accordance with previous studies stating an association of *emm*-1/ST28 with GAS

meningitis [23], 4/5 meningitis isolates belonged to the *emm*-1/ST28 lineage. The fifth

isolate belonged to the *emm*-28 lineage, which is another prevalent *emm*-type previously

described in GAS meningitis [23] and, together with *emm*-1, among the most representative

emm-types for invasive GAS infections across Europe and Nord America [24]. All otitis and

colonizing isolates belonged either to the *emm*-1 or the *emm*-28 lineage as well. The

presence or absence of GAS virulence determinants did not reveal a specific pattern for

meningitis isolates when compared to otitis and colonizing ones.

GAS meningitis occurs very rarely, leading to the availability of a small number of cases

that could be included in this study. Nevertheless, despite the small simple size, our data

suggest that the most common pathophysiological cause in this patients' cohort was local

spread from a site of bacterial infection in neighboring tissues, with hematogenous spread

found in only one case. The invasive GAS strains did not exhibit particular virulence traits

when compared to colonizing or otitis strains, suggesting that host factors such as breach of

the mucosa or local infections play the main role in transition of a bacterial strain from

colonization to invasion, without distinct virulence determinants.

Author contributions

LM: Experimental design, acquisition, analysis and interpretation of data, writing of the

manuscript.

FA: Experimental design, acquisition, analysis and interpretation of data, writing of the

manuscript.

MB: Acquisition, analysis and interpretation of data, writing of the manuscript

TAS: Experimental design, data acquisition, data analysis and critical reading of the

manuscript

DMH: Experimental design, data acquisition, data analysis

EP: Experimental design, data acquisition, data analysis

JE: Data acquisition, critical reading of the manuscript

DM: Data acquisition

AKR: Data acquisition

EMM: Histological analysis

RAS: Funding

BH: Funding

SMS: Experimental design, funding, critical reading of the manuscript

SDB: Conceptualization, funding, critical reading of the manuscript

ASZ: Conceptualization, funding, critical reading of the manuscript

Funding

This work was supported by the "Schweizerischer Nationalfonds (SNF)" [grants

31003A_169962 and 310030_204343 (to A.S.Z.)], by the "University of Zürich CRPP

Personalized medicine of persisting bacterial infections aiming to optimize treatment and

outcome" (to S.D.B, B.H. and A.S.Z.), by the "Promedica Foundation" [grant 1449/M] (to

S.D.B.), by the "Swedish Society for Medical Research (SSMF) foundation" [grant P17-0179

(to S.M.S)] and by the Vontobel stiftung (to R.A.S.).

227   Conflict of interest

The   authors   have   no   conflict   of   interest   to   declare.

<u>**References**</u>

1. Imöhl M, Reinert RR, Ocklenburg C, van der Linden M. Epidemiology of invasive

Streptococcus pyogenes disease in Germany during 2003-2007. FEMS immunology and

medical microbiology **2010**; 58:389-96.

2. Lamagni TL, Darenberg J, Luca-Harari B, et al. Epidemiology of severe Streptococcus

pyogenes disease in Europe. Journal of clinical microbiology **2008**; 46:2359-67.

3. O'Loughlin RE, Roberson A, Cieslak PR, et al. The epidemiology of invasive group A

streptococcal infection and potential vaccine implications: United States, 2000-2004. Clinical

infectious diseases : an official publication of the Infectious Diseases Society of America

2007; 45:853-62.

4. Lucas MJ, Brouwer MC, Bovenkerk S, Man WK, van der Ende A, van de Beek D. Group A

Streptococcal meningitis in adults. The Journal of infection **2015**; 71:37-42.

5. van de Beek D, de Gans J, Spanjaard L, Sela S, Vermeulen M, Dankert J. Group a

streptococcal meningitis in adults: report of 41 cases and a review of the literature. Clinical

infectious diseases : an official publication of the Infectious Diseases Society of America

2002; 34:e32-6.

6. Cole JN, Barnett TC, Nizet V, Walker MJ. Molecular insight into invasive group A

streptococcal disease. Nature reviews Microbiology **2011**; 9:724-36.

7. Nizet V, Kim KS, Stins M, et al. Invasion of brain microvascular endothelial cells by group B

streptococci. Infection and immunity **1997**; 65:5074-81.

8. Andreoni F, Ogawa T, Ogawa M, et al. The IL-8 protease SpyCEP is detrimental for Group A

Streptococcus host-cells interaction and biofilm formation. Frontiers in microbiology **2014**;

5:339.

9. Seidl K, Zinkernagel AS. The MTT assay is a rapid and reliable quantitative method to

assess Staphylococcus aureus induced endothelial cell damage. Journal of microbiological

methods **2013**; 92:307-9.

10. Andreoni F, Zürcher C, Tarnutzer A, et al. Clindamycin Affects Group A Streptococcus

Virulence Factors and Improves Clinical Outcome. J Infect Dis **2017**; 215:269-77.

11. Räz AK, Andreoni F, Boumasmoud M, et al. Limited adaptive evolution of

Staphylococcus aureus during transition from colonization to invasive

infection. bioRxiv **2021**:2021.10.25.465068.

12. Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics **2014**; 30:2068-

9.

13. Page AJ, Cummins CA, Hunt M, et al. Roary: rapid large-scale prokaryote pan genome

analysis. Bioinformatics **2015**; 31:3691-3.

14. Price MN, Dehal PS, Arkin AP. FastTree: Computing Large Minimum Evolution Trees with

Profiles instead of a Distance Matrix. Molecular Biology and Evolution **2009**; 26:1641-50.

15. Control CoD. Available at: ftp://ftp.cdc.gov/pub/infectious_diseases/biotech/tsemm/.

16. VFDB. Available at: http://www.mgc.ac.cn/VFs/download.htm.

17. Schlech WF, 3rd, Ward JI, Band JD, Hightower A, Fraser DW, Broome CV. Bacterial

meningitis in the United States, 1978 through 1981. The National Bacterial Meningitis

Surveillance Study. Jama **1985**; 253:1749-54.

18. Santos MS, Ribeiro GS, Oliveira TQ, et al. Burden of group A streptococcal meningitis in

Salvador, Brazil: report of 11 years of population-based surveillance. Int J Infect Dis **2009**;

13:456-61.

19. Randhawa E, Woytanowski J, Sibliss K, Sheffer I. Streptococcus pyogenes and invasive

central nervous system infection. SAGE Open Med Case Rep **2018**; 6:2050313x18775584.

20. Sommer R, Rohner P, Garbino J, et al. Group A beta-hemolytic streptococcus meningitis:

clinical and microbiological features of nine cases. Clinical infectious diseases : an official

publication of the Infectious Diseases Society of America **1999**; 29:929-31.

21. Hersi K GF, Kondamudi NP. Meningitis. **2021**.

22. Perin A, Nascimben E, Longatti P. Decompressive craniectomy in a case of intractable

intracranial hypertension due to pneumococcal meningitis. Acta Neurochir (Wien) **2008**;

150:837-42; discussion 42.

23. Plainvert C, Doloy A, Joubrel C, et al. Characterization of Streptococcus pyogenes isolates

responsible for adult meningitis in France from 2003 to 2013. Diagnostic microbiology and

infectious disease **2016**; 84:350-2.

24. Gherardi G, Vitali LA, Creti R. Prevalent emm Types among Invasive GAS in Europe and

North America since Year 2000. Front Public Health **2018**; 6:59-.

 Figure legend

Figure 1 - Patients scans and histology results. **A)** CT-Scan: septic thrombosis with

mastoiditis and otitis media on the right side. Arrow indicates thrombosis **B)** MRI-Scan: T2

sequence showing penetration of lower temporal lobe with abscess formation. Arrow

indicates abscess **C)** HE Staining of mastoid tissue: arrow indicates necrotic mastoid tissue,

mixed neutrocytic and lymphocytic infiltration, foam cells and cholesterol crystals (not

depicted) indicating chronic inflammation. **D)** Brown-Brenn staining of mastoid tissue: arrow

indicates accumulation of extracellular bacteria.

Figure 2 - Bacterial strains virulence determinants. **A)** Adherence and **B)** invasion to HBMEC

cells shows no difference between isolates groups. Bacterial virulence factors activity of **C)**

the nuclease Sda1, **D)** the IL-8 protease SpyCEP and **E)** SLO as well as **F)** HBMEC cells barrier

function disruption did not vary among isolates groups. For adherence, invasion as well as

Sda1 and SpyCEP activity, each point on the graph represents the average of at least three

biological replicates for one strain. Excluding barrier disruption experiments, which were

carried out in duplicate, all experiment were conducted at least in triplicate. Control

experiments were carried out using medium only.

A CT scan

B MRI

C H&E stain

D Brown-Brenn stain

Patient ID Sex Age(y) Isolate Nr.	Associated health conditions	Clinical presentation	Complications	Site of GAS identification	Antibiotic treatment regimen	Death
1 F 73 CI1224	• Perforated otitis media • Diabetes • Hypothyreosis • Depression	• Altered vigilance and mental state • Vertigo • Hypothermia • Hypotension • Acute abdomen	• Partial septic sinus transversus thrombosis • Coma • Irreversible hypoxic brain damage • ICU involvement	CSF Blood Ear swab	3d ceftriaxon 3d amoxicillin 3d clindamycin	yes
2 M 55 CI1271	• Otitis media • Depression	• Altered vigilance • Meningism • Slight one-sided arm paralysis • Otalgia • Diarrhea • Nausea	• Coma • Mastoiditis • Cerebral abscess • ICU involvement	Blood Mastoid tissue	2d ceftriaxon 1d amoxicillin Escalation to: 18d penicillin 5d metronidazole 5d clindamycin + Surgical intervention	no
3 F 64 CI1293	• Bicuspid aortic valve	• Acute reduction of general condition • Meningism • Right sided paresis of upper and lower limbs • Right sided tongue deviation • Anisocoria • Petechial bleeds	• Thrombocytopenia • Acute kidney failure • Aortitis • Nosocomial Influenza infection	CSF Blood	Initially ceftriaxon then change to penicillin due to elevated transaminases, total 18d 3d gentamycin 15d steroids 10d vancomycin 10d clindamycin	no
4 F 79 CI407	• Otitis media (perforated) • Coronary heart disease • Pre-diabetic metabolism	• High temperatures • Vertigo • Altered mental state and vigilance	• Tegmen tympani penetration • Mastoiditis • Labyrinthitis • Coma • Long-term one sided deafness • ICU involvement	Mastoid tissue	Initially 1d ceftriaxon, then escalation to meropeneme for 2d, then back to ceftriaxone for 20 days Initially 2d acyclovir 1d amoxicillin 4d mephamesone + Surgical intervention	no
5 M 42 CI543	• Otitis media • Bullous • Myringitis with insertion of tympanostomy tube	• One day after insertion of tympanostomy tube acute reduction of vigilance • Altered mental state and meningism	• Mastoiditis • Subdural empyema • Cerebritis • Ventrikulitis • Coma • ICU involvement	Blood CSF Mastoid tissue	9d ceftriaxon, then escalation to 18d ceftazidime and de-escalation back to ceftriaxone for another 35d 2d amoxicillin + clavulanic acid 2d acyclovir 2d metronidazol 2d, pause 3d, then another 2d 9d intrathecal gentamycin 9d intrathecal vancomycin + Surgical intervention	no

Table 1 – Patients description. Clinical findings, complications and treatment regimen of the patients included in this study. Abbreviations: CSF=cerebrospinal fluid; d=day; ICU=intensive care unit.