Evolution of a Functionally Intact but Antigenically Distinct DENV Fusion Loop

Rita M. Meganck[1], Deanna Zhu[2], Stephanie Dong[2], Lisa J. Snoderly-Foster[1], Yago R.

Dalben[1], Devina Thiono[3], Laura J. White[3], Aravinda M. DeSilva[3], Ralph S. Baric[2],

Longping V. Tse[1]*

[1] Department of Molecular Microbiology and Immunology, Saint Louis University

[2] Department of Epidemiology, University of North Carolina at Chapel Hill

[3] Department of Microbiology, University of North Carolina at Chapel Hill

*Corresponding author: victor.tse@health.slu.edu

ABSTRACT

A hallmark of Dengue virus (DENV) pathogenesis is the potential for antibody-dependent enhancement, which is associated with deadly DENV secondary infection, complicates the identification of correlates of protection, and negatively impacts the safety and efficacy of DENV vaccines. ADE is linked to antibodies targeting the fusion loop (FL) motif of the envelope protein, which is completely conserved in mosquito-borne flaviviruses and required for viral entry and fusion. In the current study, we utilized saturation mutagenesis and directed evolution to engineer a functional variant with a mutated FL (D2-FL) which is not neutralized by FL-targeting monoclonal antibodies. The FL mutations were combined with our previously evolved prM cleavage site to create a mature version of D2-FL (D2-FLM), which evades both prM- and FL-Abs but retains sensitivity to other type-specific and quaternary cross-reactive (CR) Abs. CR serum from heterotypic (DENV4) infected non-human primates (NHP) showed lower neutralization titers against D2-FL and D2-FLM than isogenic wildtype DENV2 while similar neutralization titers were observed in serum from homotypic (DENV2) infected NHP. We propose D2-FL and D2-FLM as valuable tools to delineate CR Ab subtypes in serum as well as an exciting platform for safer live attenuated DENV vaccines suitable for naïve individuals and children.

INTRODUCTION

Dengue virus (DENV) is a member of the *Flavivirus* genus and is a major global public health threat, with four major serotypes of DENV found worldwide. Dengue causes ~400 million infections each year, of which ~20% of cases present clinically, a subset of which may progress to severe Dengue Hemorrhagic Fever/Dengue Shock Syndrome (DHF/DSS).[1,2] DENV is transmitted through *Aedes* mosquito vectors, and globalization and global warming are increasing the endemic range of Dengue worldwide.[3,4] The pathogenesis of Dengue is complex, as first-time infections are rarely severe and lead to serotype-specific immunity. However, re-infection with a different serotype increases the risk of developing DHF/DSS.[5] This is thought to be due to the phenomenon of antibody-dependent enhancement (ADE), in which poorly neutralizing cross-reactive (CR) antibodies (Abs) lead to enhanced viral uptake and infection of unique cell populations in an Fcγ-receptor-mediated manner.[6]

ADE remains a major challenge for DENV vaccine development.[7] The leading DENV vaccine platforms in clinical testing are tetravalent live attenuated virus mixtures of all four serotypes. However, creating formulations that elicit a balanced response has proven challenging.[8] Additionally, lab-grown strains differ from patient-derived DENVs in both maturation status and antigenicity.[9] In particular, Abs targeting the fusion loop (FL) have been reported to neutralize lab and patient strains with differing strengths and have been observed to facilitate Fcγ-receptor uptake *in vitro* and therefore ADE.[10–12] Currently, there is a single FDA-approved DENV vaccine, Dengvaxia. However, it is only approved for use in individuals aged 9-16 with previous DENV infection living in endemic areas and is contraindicated for use in naïve individuals and younger children. In naïve children,

vaccination stimulated non-neutralizing CR Abs that increased the risk of severe disease

after DENV infection.[13,14] Other DENV vaccines have been tested or are currently

undergoing clinical trial, but thus far none have been approved for use in the United

States.[15] The vaccine Qdenga has been approved in the European Union, Indonesia, and

Brazil, although vaccine efficacy in adults, naïve individuals, and with all serotypes has

not yet been shown.

The DENV FL is located in Envelope (E) protein domain II (EDII) and is involved

in monomer-monomer contacts with EDIII.[16] During the DENV infection cycle, low pH

triggers a conformational change in the E protein.[17] The structure of the virion rearranges,

and individual monomers form a trimer with all three FLs in the same orientation, ready

to initiate membrane fusion.[16,17] The core FL motif (DRGWGNGCGLFGK, AA 98-110) is

highly conserved, with 100% amino acid conservation in all DENV serotypes and other

mosquito-borne flaviviruses, including Yellow fever virus (YFV), Zika virus (ZIKV), West

Nile virus (WNV), Kunjin virus (KUNV), Murray Valley encephalitis virus (MVEV),

Japanese encephalitis virus (JEV), Usutu virus (USUV), and Saint Louis encephalitis virus

(SLEV; Figure 1A). Although the extreme conservation and critical role in entry have led

to it being considered extremely difficult to change the FL, we successfully tested the

hypothesis that massively parallel directed-evolution could produce viable DENV FL

mutants that were still capable of fusion and entry, while altering the antigenic footprint.

The FL mutations, in combination with optimized prM cleavage site mutations, ablate

neutralization by the prM- and FL-Abs, retain sensitivity to other protective Abs, and

provide a novel vaccine strategy for DENV.

RESULTS

To engineer a virus with a novel antigenic footprint at the FL, we targeted the core conserved FL motif. We generated two different saturation mutagenesis libraries, each with 5 randomized amino acids: DRG**X**G**X**G**XXX**FGK (Library 1; AA 101, 103, 105-107) and DRG**XXXXX**GLFGK (Library 2 AA 101-105). Library 1 was designed to mutate known residues targeted by FL mAbs while Library 2 focused on a continuous linear peptide that is the epitope for FL-Abs to maximally alter antigenicity.[18] Saturation mutagenesis plasmid libraries were used to produce viral libraries in either C6/36 (*Aedes albopictus* mosquito) or Vero 81 (African green monkey) cells and passaged three times in their respective cell types. Following directed evolution, viral genomes were extracted and subjected to deep sequencing to identify surviving and enriched variants (Figure 1B). Due to the high level of conservation, it was not surprising that most mutational combinations failed to yield viable progeny. In fact, evolutions carried out on Library 2 only yielded wild-type sequences. For Library 1, wild-type sequences dominated in Vero 81 evolved libraries. However, a novel variant emerged in C6/36 cells with two amino acid changes: DRGWG**S**GC**L**LFGK. The major variant comprised ~95% of the population, while the next most populous variant (DRGWG**S**GC**W**LFGK) comprised only 0.25% (Figure 1C). Bulk Sanger sequencing revealed an additional Env-T171A mutation outside of the FL region. Residues W101, C105, and L107 were preserved in our final sequence, supporting the importance of these residues.[16] When modeled on the pre-fusion DENV2 structure, the N103S and G106L mutations are located at the interface with the neighboring monomer EDIII domain, protected from the aqueous environment. In the post-fusion form, the two residues are located between W101 and F108 and form the

bowl concavity above the chlorine ion in the post-fusion trimer (Figure 1D). We used

reverse genetics to re-derive the FL N103S/G106L/T171A mutant, which we term D2-FL.

As enhancing Abs also target prM,[19] we also created a mature version of D2-FL termed

D2-FLM, containing both the evolved FL motif and our previously published evolved prM

furin cleavage site, which results in a more mature virion like those found in infected

patients (Figure 1E).[9,20]

We performed growth curves comparing DENV2, D2-FL, and D2-FLM in both

C6/36 and Vero 81 cells. In C6/36 cells, the growth of all three viruses was comparable,

reaching high titers of 10^6-10^7 FFU/mL. However, in Vero 81 cells, both FL mutant viruses

were highly attenuated, with a 2-2.5 log reduction in titer (Figure 2A). The species-specific

phenotype in culture involved a change from insect to mammalian cells, as well as a

change in growth temperature. To investigate if the mutant viruses were more unstable

at higher temperatures, we performed a thermostability assay, comparing viruses

incubated at temperatures ranging from 4-55°C before infection. The three viruses had

comparable thermostabilities, indicating that this does not explain the attenuation of the

FL mutants (Figure 2B). Because the D2-FLM virus contains mutations that increase prM

cleavage frequency, we also assayed the maturation status of the three viruses by

western blot. D2-FL had a comparable prM:E ratio to the isogenic wildtype DENV2 (DV2-

WT), while, as expected, D2-FLM had a reduced prM:E ratio, indicating a higher degree

of maturation (Figure 2C).

Next, we characterized the ability of Abs targeting the FL to recognize DV2-WT,

D2-FL, and D2-FLM with a panel of monoclonal antibodies (mAbs). Importantly, D2-FL

and D2-FLM were resistant to mAbs targeting the FL. Neutralization by 1M7 is reduced

by ~2-logs in both variants, 1N5 neutralization is reduced by ~1-log for D2-FL and reduced

to background levels for D2-FL, and no neutralization was observed for 1L6 or 4G2 for

either variant (Figure 3A).[18] Focusing on the D2-FLM virus containing both evolved motifs,

we then characterized the antigenicity of the whole virion with a panel of mAbs. As

expected, D2-FLM was unable to be neutralized by the prM Abs 1E16 and 5M22; the Ab

2H2 does not neutralize either DV2-WT or D2-FLM (Figure 3B). For Abs targeting

epitopes in non-mutated regions, including the ED1 and EDE epitopes that target EDII

and EDIII, $FRNT_{50}$ values were generally comparable, although EDE1-C10 shows a

moderate but statistically significant reduction between DV2-WT and D2-FLM, indicating

that the overall virion structural integrity was intact (Figure 3B).

Next, we analyzed neutralization of the D2-FLM virus using serum derived from

convalescent humans and experimental infected non-human primates (NHPs). Overall,

we tested serum from 6 humans and 9 NHPs at different time points with a total of 27

samples. Serum from a homotypic infected NHP (n=3) did not display a difference in

neutralization between DV2-WT, D2-FL, and D2-FLM, confirming that prM and FL

epitopes are not significant contributors to the homotypic type-specific (TS) neutralizing

Ab response in primates (Figure 3C). In heterotypic vaccination and infection, most of the

serum (18/24) did not cross-neutralize ($FRNT_{50}$ < 1:40) DV2-WT, confirming the serotypic

difference of DENVs (Table 1). However, in two NHPs infected with DENV4, strong

neutralization potency ($FRNT_{50}$ between 1: 100 – 1:1,000) was demonstrated against

DV2-WT (Figure 3C). Heterologous cross-neutralization was significantly reduced to

background levels ($FRNT_{50}$ < 1:40) against the D2-FLM virus at 90 days post-infection

(dpi). Of note, one DENV4 animal (3Z6) showed low levels of neutralization against D2-

FLM at early time points (20- and 60-days post-infection), which was eventually lost at

later time points. Neutralization observed against D2-FL, in general, fell between DV2-

WT and D2-FLM. Interestingly, in animal 3Z6, at 20 dpi, neutralization against DENV-FL

was comparable to DV2-WT, while D2-FLM was greatly reduced, indicating that

antibodies in the sera targeting the immature virion formed a large portion of the CR

response. In contrast, animal 0Y0 displayed less difference in neutralization between D2-

FL and D2-FLM, suggesting that FL antibodies were more prominent in this animal. These

data suggest that after a single infection, much of the CR Ab responses target prM and

the FL and tend to wane over time (Figure 3C). The collection of FL, mature, and FL-

mature variants provides new opportunity to delineate antibody composition in complex

polyclonal serum from DENV natural infection and vaccination.

DISCUSSION

Mechanistic understanding of vaccine protection and identification of correlates of

protection are immensely important for DENV vaccine development. The dual protective

and enhancing properties of DENV Abs create major challenges for dissecting the role of

various Ab populations in disease protection. Cross-reactive weakly neutralizing prM- and

FL-Abs are often immunodominant after primary DENV infection,[10,21–24] and can lead to

overestimation of the levels of heterotypic protection in traditional neutralization assays.

Since these same antibodies are also associated with ADE,[19] inaccurate conclusions

could have dire consequences if protection *in vitro* translates to the enhancement of

disease in human vaccinees. Unfortunately, Ab profiling in polyclonal serum is mainly

performed by ELISA, and a neutralization assay that can discriminate Abs does not exist.

The D2-FLM variant is not neutralized by FL- and prM-mAbs and appears insensitive to

neutralization by these Abs in polyclonal serum. Of note, EDE1-C10 neutralization

potency was also reduced in our FL-M variant, further indicating our mutations are

affecting the tip of the EDII region which partially overlaps with the EDE1 epitope[25]. In

combination with other chimeric DENVs,[26–28] D2-FLM provides a reagent to distinguish

between TS, protective CR (e.g. E dimer epitope, EDE),[29,30] and ADE-prone CR (e.g. FL

and prM)[24] Ab subclasses in neutralization assays after infection and vaccination.

Due to the ADE properties of DENV Abs, studies to understand and eliminate ADE

phenotype are under active investigation. For example, mAbs can be engineered to

eliminate binding to the Fcγ receptor, abolishing ADE.[31] While this methodology holds

potential for Ab therapeutic development and passive immunization strategies, it is not

relevant for vaccination. As FL and prM targeting Abs are the major species demonstrated

to cause ADE *in vitro* and are thought to be responsible for ADE-driven negative

outcomes after primary infection and vaccination,[10–12,32] we propose that genetic ablation

of the FL and prM epitopes in vaccine strains will minimize the production of these

subclasses of Abs responsible for undesirable vaccine responses. Indeed, efforts have

been made to reduce the availability of the FL or reduce the ability of FL Abs to drive

ADE. Covalently locked E-dimers and E-dimers with FL mutations have been engineered

as subunit vaccines that reduce the availability of the FL, thereby reducing the production

of FL Abs.[33–36] DENV subunit vaccines are an area of active study;[37] however,

monomer/dimer subunits can also expose additional, interior-facing epitopes not normally

exposed to the cell. Furthermore, dimer subunits are not a complete representation of the

DENV virion which presents other structurally important interfaces such as the 3-fold and

5-fold symmetries. Concerns about balanced immunity to all four serotypes also apply to

subunit vaccine platforms. Given the complexity of the immune response to DENV, live

virus vaccine platforms have thus far been more successful. However, the fusion loop is

strongly mutationally intolerant. Previous reported mutations in the fusion loop of

mosquito-borne flaviviruses include a L107F mutation in WNV [38] and JEV associated with

attenuation, and mutations at position 106 in ZIKV(G106A) and DENV(G106V) which

were tolerated. Interestingly, we also recovered a mutation in position 106 (G106L) Using

directed-evolution, we successfully generated our D2-FLM variant that combines viability

with the desired Ab responses. Therefore, the D2-FLM variant is a novel candidate for a

vaccine strain which presents all the native structures and complex symmetries of DENV

necessary for T-cell mediated responses and which can elicit more optimal protective Ab

responses.[39–41]

Other considerations of high importance when designing a live DENV vaccine

include strain selection and serotype balance.[42] In the current study we used DENV2

S16803, a prototype for DENV2.[43] However, S16803 was isolated several decades ago,

and it may be beneficial to utilize more contemporaneous strains.[8,44] Work is currently

ongoing to demonstrate the portability of the evolved FL motif on additional DENV2

strains and other serotypes, which is essential for tetravalent vaccine production. D2-FLM

was highly attenuated in Vero cells, creating a challenge for vaccine production.

Therefore, further adaptation of this strain to grow efficiently in mammalian cells while

retaining its antigenic properties is needed. Taken together, the FLM variant holds

exciting new possibilities for a new generation of DENV vaccines, as well as a platform

to readily measure TS and CR ADE-type responses and thereby assess the true

protective potential of any DENV vaccine trials and safeguard approval of DENV vaccines

for human use.

ACKNOWLEDGEMENTS

We thank members of the Tse, Baric, and DeSilva laboratories for helpful discussions. This work was supported by NIAID R01AI107731 to A.D. and R.S.B., P01AI106695 to R.S.B., NIAID F30AI160898 to D.R.Z. L.J.W. is supported by NIAID P01 5112869.

AUTHOR CONTRIBUTION

R.M.M. and L.V.T. designed the study. R.M.M. performed high-throughput sequencing preparation and analysis. R.M.M., D.Z., S.D., L.J.S., Y.D., and L.V.T. performed experiments. D.T., L.J.W., A.M.D.S., and R.S.B. provided reagents. L.V.T. and R.S.B. provided oversight of the project and funding. R.M.M. wrote the manuscript. L.V.T. reviewed and revised the final version. All authors approved the final version of the manuscript.

CONFLICT DISCLOSURE

R.M.M., R.S.B., and L.V.T. are inventors on a patent application filed on the subject matter of the manuscript.

MATERIALS AND METHODS

<u>Cells and viruses</u>

C6/36 (ATCC CRL-1660) were grown in MEM (Gibco) with 5% FBS (HyClone), 1% penicillin/streptomycin (Gibco), 0.1mM nonessential amino acids (Gibco), 1% HEPES (Gibco), and 2mM GlutaMAX (Gibco), cultured at 32°C with 5% CO_2. Vero 81 cells (ATCC CCL-81) were grown in DMEM/F12 (Gibco) with 10% FBS, 1% penicillin/streptomycin (Gibco), 0.1mM nonessential amino acids (Gibco), and 1% HEPES (Gibco), cultured at 37°C with 5% CO_2. DENV viruses were grown in C6/36 or Vero 81 cells maintained in infection media. C6/36 infection media consists of Opti-MEM (Gibco) with 2% FBS (HyClone), 1% penicillin/streptomycin (Gibco), 0.1mM nonessential amino acids (Gibco), 1% HEPES (Gibco), and 2mM GlutaMAX (Gibco). Vero 81 infection media consists of DMEM/F12 (Gibco) with 2% FBS, 1% penicillin/streptomycin (Gibco), 0.1mM nonessential amino acids (Gibco), and 1% HEPES (Gibco). DENV2 strain S16803 was used in this study.[43] Sequences used for the alignments include DENV1 WestPac-74 (U88535.1), DENV2 S-16803 (GU289914.1), DENV3 3001 (JQ411814.1), DENV4 Sri Lanka-92 (KJ160504.1), YFV 17D (NC_002031.1), SLEV Kern217 (NC_007580.2), JEV (NC_001437.1), USUV Vienna-2001 (NC_006551.1), MVEV (NC_000943.1), WNV-1 NY99 (NC_009942.1), and ZIKV MR-766 (NC_012532.1).

<u>DENV Reverse Genetics</u>

DENV2 S16803 was used in this study. Recombinant viruses were created using a four-plasmid system as previously described,[45] consisting of the DENV genome split into four segments, each cloned into a separate plasmid. The DENV plasmids were digested and ligated to form a single template for *in vitro* transcription. The resulting RNA was

electroporated into either C6/36 or Vero cells. Virus-containing supernatant was

harvested at 4-5 days post electroporation and passaged. DENV variants were created

through site-directed mutagenesis of the DENV plasmids.

<u>Library Generation and Directed Evolution</u>

DENV fusion loop libraries were generated through saturation mutagenesis of the

indicated resides, based on a previously published protocol.[20,46] Degenerate NNK

oligonucleotides were used to amplify the region, generating a library of mutated DNA

fragments. Q5 DNA Polymerase was used with less than 18 cycles to maintain accuracy.

The resulting library was cloned into the DENV reverse genetics system. The ligated

plasmids were electroporated into DH10B ElectroMax cells (Invitrogen) and directly

plated on 5,245mm^2 dishes (Corning) to avoid bias from suspension culture. Colonies

were pooled and purified using a Maxiprep kit (Qiagen), and the plasmid library used for

DENV reverse genetics (above). Viral libraries were passaged three times in the

corresponding cell type.

<u>High-throughput Sequencing and Analysis</u>

Viral RNA was isolated with a QIAamp viral RNA kit (Qiagen), and cDNA produced using

the Superscript IV Reverse Transcriptase (Invitrogen). Amplicons were prepared for

sequencing using the Illumina TruSeq system with two rounds of PCR using Q5 Hot Start

DNA polymerase (NEB). For the first round of PCR, primers were specific to the DENV2

E sequence surrounding the fusion loop motif with overhangs for the Illumina adapters.

After purification, this product was used as the template for the second round of PCR

using Illumina P5 and P7 primers containing 8-nucleotide indexes. Purified PCR products

were analyzed on a Bioanalyzer (Agilent Technologies) and quantified on a Qubit 4

fluorometer (Invitrogen). Amplicon libraries were run on a MiSeq system with 2x150bp

reads. Plasmid and P0 libraries were sequenced at a depth of ~4.5 million reads; later

passages were sequenced at a depth of ~750,000 reads. Custom perl and R scripts were

used to analyze and plot the data as previously published.[20]

DENV Growth Kinetics

One day before infection, $5x10^5$ cells were seeded in every well of a 6-well plate. Cells

with infected with an MOI of 0.05-0.1, estimating $1x10^6$ cells on the day of infection.

Infection was carried out for one hour in the incubator, followed by 3x washes with PBS

and replenishment with fresh infection medium. 300 uL of viral supernatant was collected

at 0, 24, 48, 72, 96, and 120 hours and stored at -80°C. All experiments were performed

independently at least three times.

DENV Focus-Forming Assay

Titers of viral supernatant were determined using a standard DENV focus-forming assay.

In brief, cells were seeded at $2x10^4$ cells per well of a 96-well plate one day before

infection. The next day, 50 uL of 10-fold serial dilution of viral supernatant were added to

each well for 1 hour in the incubator. After, 125uL of overlay (Opti-MEM, 2% FBS, NEAA,

P/S, and methylcellulose) was added to each well. Infection was allowed to continue for

48 hours in the incubator. Overlay was removed, and each well rinsed 3x with PBS

followed by a 30 minute fixation with 10% formalin in PBS. Cells were blocked in

permeabilization buffer (eBioscience) with 5% nonfat dried milk. Primary Abs anti-prM

2H2 and anti-E 4G2 from nonpurified hybridoma supernatant were used at a 1:500

dilution in blocking buffer. Goat anti-mouse HRP secondary (SeraCare KPL) were used

at a 1:1000 dilution in blocking buffer. Followed washing, foci were developed using

TrueBlue HRP substrate (SeraCare) and counted using an automated Immunospot

analyzer (Cellular Technology).

<u>Thermal Stability Assay</u>

The indicated viruses were thawed and incubated at temperatures ranging from 4°C to

55°C for one hour. Following, viral titers were determined by focus-forming assay as

described above.

<u>Western Blotting</u>

Viral supernatants were combined with 4X Laemmli Sample Buffer (Bio-Rad) and boiled

at 95°C for 5 minutes. After SDS-PAGE electrophoresis, samples were transferred to

PVDF membrane and blocked in 3% nonfat milk in PBS-T. A polyclonal rabbit anti-prM

(1:1000; Invitrogen PA5-34966) and polyclonal rabbit anti-Env (Invitrogen PA5-32246) in

2% BSA in PBS-T were incubated on the blot for 1 hour at 37C. Goat anti-rabbit HRP

(1:10,000 Jackson ImmunoLab) in 3% milk in PBS-T was incubated on the blot for 1 hour

at room temperature. Blots were developed by SuperSignal West Pico Plus

chemiluminescent substrate (ThermoFisher). Blots were imaged on an iBright FL1500

imaging system (Invitrogen). The pixel intensity of individual bands was measured using

ImageJ, and the relative maturation was calculated by using the following equation:

(prMExp/EnvExp)/(prMWT/EnvWT). All experiments were performed independently a

minimum of three times.

<u>FRNT Assay</u>

Focus reduction neuralization titer (FRNT) assays were performed as described

previously with C6/36 cells.[20] $1x10^5$ cells were seeded in a 96-well plate the day prior to

infection. Abs or sera were serially diluted and mixed with virus (~100 FFU/well) at a 1:1

volume and incubated for 1 hour in the incubator. The mixture was added onto the plate

with cells and incubated for 1 hour in the incubator, then overlay was added (see Focus-

Forming Assay) and plates were incubated for 48 hours. Viral foci were stained and

counted as described above (Focus-Forming Assay). A variable slope sigmoidal dose-

response curve was fitted to the data, and values were calculated with top or bottom

restraints of 100 and 0 using GraphPad Prism version 9.0. All experiments were

performed independently at least two times, due to limited amounts of human serum.

<u>Statistical Analysis</u>

GraphPad Prism version 9.0 was used for statistical analysis. Titer and % infection of D2-

FL and D2-FLM were compared to the DV2 using two-way ANOVA. $FRNT_{50}$s were

compared using Student's t-test. Significant symbols are as follows: *, $P < 0.05$; **, $P <$

0.005; ***, $P < 0.0005$; ****, $P < 0.00005$. The data are graphed as means ± standard

deviations.

Figures:

Figure 1: Generation of DENV2 fusion loop mutants via directed evolution. A) Alignment of Top: Dengue virus fusion loops; Bottom: Mosquito-borne flavivirus fusion loops, including Yellow Fever virus (YFV), Zika virus (ZIKV), West Nile virus (WNV), Kunjin virus (KUNV), Murray Valley Encephalitis virus (MVEV), Japanese Encephalitis virus (JEV), Usutu virus (USUV), and Saint Louis Encephalitis virus (SLEV). Amino acids are colored by functional groups: negatively charged (red), positively charged (blue), nonpolar (yellow), polar (green), aromatic (pink), and sulfide (dark red). B) Schematic of directed evolution procedure. Saturation mutagenesis libraries were used to produce viral libraries, which were passaged three times in either C6/36 or Vero 81 cells. At the end of the selection, viral genomes were isolated and mutations were identified by high-throughput sequencing. C) Left: Bubble plot of the sequences identified from either the unselected or selected (passage 3) C6/36 DENV libraries. Right: Pie chart of the sequences from passage 3 C6/36 DENV libraries. D) Structure of the DENV envelope with the fusion loop mutations highlighted in red. E) Sequences of the fusion loop and furin cleavage site of DENV2, D2-FL, and D2-FLM.

Figure 2: Biological and physical properties of mature DENV2 fusion loop mutants. A) Multistep growth curves (MOI = 0.05-0.1) of DV2-WT, D2-FL, and D2-FLM on C6/36 cells (left) or Vero 81 cells (right). B) Thermostability assay on DV2-WT, D2-FL, and D2-FLM. C) Western blot of virions, blotted against Envelope and prM proteins. The prM:E ratio was determined and normalized to the DV2-WT ratio. Averages of 3 biological replicates are shown. Two-way ANOVA was used for statistical comparison of growth curves and thermostability: ns = not significant; * < 0.05; ** < 0.005; *** < 0.0005.

A

B

C

Figure 3: Fusion loop mutant is insensitive to fusion loop mAbs, the major target for cross-

reactive Abs in NHPs. A) Left: $FRNT_{50}$ values for neutralization of DV2-WT, D2-FL, and

D2-FLM with mAbs against the FL (1M7, 1N5, 1L6, 4G2). All Abs were tested in at least

n=3 independent experiments, except 1N5 due to limited Ab. Right: Average

neutralization curves for neutralization of DV2-WT, D2-FL, and D2-FLM with mAbs

against the DENV2 fusion loop. B) $FRNT_{50}$ values for neutralization of DV2-WT and D2-

FLM with mAbs against DENV2 prM (2H2, 1E16, 5M22), EDI (3F9), EDE (C10, B7), and

EDIII (2D22). All Abs were tested in at least n=3 independent experiments. C)

Neutralization of DV2-WT, D2-FL, and D2-FLM with sera from NHPs infected with either

DENV4 or DENV2. $FRNT_{50}$s were compared using Student's t-test. Significant symbols

are as follows: *, $P < 0.05$; **, $P < 0.005$; ***, $P < 0.0005$; ****, $P < 0.00005$. The data are

graphed as means ± standard deviations.

Human #	Infected Serotype	Collection Time	DV2	D2-FL	D2-FLM
DS1500	DENV1	Convalescent serum	1 : 75	ND	1 : 53
DS2499	DENV1		1 : 80	ND	1 : 52
DS1136	DENV3		1 : 133	ND	1 : 45
DS1160	DENV3		1 : 86	ND	1: 208
DS0275	DENV4		1 : 94	ND	1 : 94
DS2239	DENV4		1 : 156	ND	1 : 42

NHP #	Infected Serotype	Collection Time	DV2	D2-FL	D2-FLM
R628	DENV4	30dpi	< 1:40	< 1:40	< 1:40
		60dpi	< 1:40	< 1:40	< 1:40
		90dpi	1 : 42*	< 1:40	< 1:40
R737		30dpi	< 1:40	< 1:40	< 1:40
		60dpi	< 1:40	< 1:40	< 1:40
		90dpi	< 1:40	< 1:40	< 1:40
R1132		30dpi	< 1:40	< 1:40	< 1:40
		60dpi	< 1:40	< 1:40	< 1:40
		90dpi	< 1:40	< 1:40	< 1:40
R1160		30dpi	1 : 58*	< 1:40	< 1:40
		60dpi	1 : 57*	1 : 50*	< 1:40
		90dpi	< 1:40	1 : 57*	1 : 91*

Table 1: Summary of FRNT$_{50}$s of human convalescent serum and NHP infection serum against DV2, D2-FL and D2-FLM.

REFERENCES

1. Bhatt, S. *et al.* The global distribution and burden of dengue. *Nature* **496**, (2013).
2. Brady, O. J. *et al.* Refining the Global Spatial Limits of Dengue Virus Transmission by Evidence-Based Consensus. doi:10.1371/journal.pntd.0001760.
3. Ebi, K. L. & Nealon, J. Dengue in a changing climate. *Environ Res* **151**, 115–123 (2016).
4. Messina, J. P. *et al.* The current and future global distribution and population at risk of dengue. *Nat Microbiol* doi:10.1038/s41564-019-0476-8.
5. Halstead, S. B. Pathogenesis of Dengue: Challenges to Molecular Biology. *Science (1979)* **239**, 476–481 (1988).
6. Katzelnick, L. C. *et al.* Antibody-dependent enhancement of severe dengue disease in humans. *Science (1979)* **358**, 929–932 (2017).
7. Rey, F. A., Stiasny, K., Vaney, M., Dellarole, M. & Heinz, F. X. The bright and the dark side of human antibody responses to flaviviruses: lessons for vaccine design. *EMBO Rep* **19**, 206–224 (2018).
8. Rabaa, M. A. *et al.* Genetic epidemiology of dengue viruses in phase III trials of the CYD tetravalent dengue vaccine and implications for efficacy. *Elife* **6**, 1–22 (2017).
9. Raut, R. *et al.* Dengue type 1 viruses circulating in humans are highly infectious and poorly neutralized by human antibodies. *Proc Natl Acad Sci U S A* **116**, 227–232 (2019).
10. Lai, C. Y. *et al.* Analysis of Cross-Reactive Antibodies Recognizing the Fusion Loop of Envelope Protein and Correlation with Neutralizing Antibody Titers in Nicaraguan Dengue Cases. *PLoS Negl Trop Dis* **7**, 1–11 (2013).
11. Beltramello, M. *et al.* The human immune response to dengue virus is dominated by highly cross-reactive antibodies endowed with neutralizing and enhancing activity. *Cell Host Microbe* **8**, 271–283 (2010).
12. Costin, J. M. *et al.* Mechanistic Study of Broadly Neutralizing Human Monoclonal Antibodies against Dengue Virus That Target the Fusion Loop. *J Virol* **87**, 52–66 (2013).
13. Wilder-Smith, A. *et al.* Personal View Deliberations of the Strategic Advisory Group of Experts on Immunization on the use of CYD-TDV dengue vaccine The Strategic Advisory Group of Experts (SAGE) on Immunization advises WHO on global policies for vaccines. In. *Lancet Infect Dis* **19**, e31–e38 (2019).
14. Dengue vaccine: WHO position paper, September 2018 – Recommendations. *Vaccine* **37**, 4848–4849 (2019).
15. Izmirly, A. M., Alturki, S. O., Alturki, S. O., Connors, J. & Haddad, E. K. Challenges in Dengue Vaccines Development: Pre-existing Infections and Cross-Reactivity. *Front Immunol* **11**, 1055 (2020).
16. Modis, Y., Ogata, S., Clements, D. & Harrison, S. C. Structure of the dengue virus envelope protein after membrane fusion. *Nature* **427**, 313–319 (2004).
17. Klein, D. E., Choi, J. L. & Harrison, S. C. Structure of a Dengue Virus Envelope Protein Late-Stage Fusion Intermediate. *J Virol* **87**, 2287–2293 (2013).

424   18.   Smith, S. A. *et al.* The potent and broadly neutralizing human dengue virus-
425   specific monoclonal antibody 1C19 reveals a unique cross-reactive epitope on the
426   bc loop of domain II of the envelope protein. *mBio* **4**, (2013).
427   19.   Rodenhuis-Zybert, I. A. *et al.* Immature Dengue Virus: A Veiled Pathogen? *PLoS*
428   *Pathog* **6**, e1000718 (2010).
429   20.   Tse, L. v. *et al.* Generation of Mature DENVs via Genetic Modification and
430   Directed Evolution. *mBio* **13**, 1–18 (2022).
431   21.   Smith, S. A. *et al.* Persistence of Circulating Memory B Cell Clones with Potential
432   for Dengue Virus Disease Enhancement for Decades following Infection. *J Virol*
433   **86**, 2665–2675 (2012).
434   22.   Oliphant, T. *et al.* Antibody Recognition and Neutralization Determinants on
435   Domains I and II of West Nile Virus Envelope Protein. *J Virol* **80**, 12149–12159
436   (2006).
437   23.   Lai, C.-Y. *et al.* Antibodies to Envelope Glycoprotein of Dengue Virus during the
438   Natural Course of Infection Are Predominantly Cross-Reactive and Recognize
439   Epitopes Containing Highly Conserved Residues at the Fusion Loop of Domain II.
440   *J Virol* **82**, 6631–6643 (2008).
441   24.   Dejnirattisai, W. *et al.* Cross-Reacting Antibodies Enhance Dengue Virus Infection
442   in Humans. *Science (1979)* **328**, 745–748 (2010).
443   25.   Sharma, A. *et al.* The epitope arrangement on flavivirus particles contributes to
444   Mab C10's extraordinary neutralization breadth across Zika and dengue viruses.
445   *Cell* **184**, 6052-6066.e18 (2021).
446   26.   Andrade, D. v. *et al.* Tracking the polyclonal neutralizing antibody response to a
447   dengue virus serotype 1 type-specific epitope across two populations in Asia and
448   the Americas. *Scientific Reports 2019 9:1* **9**, 1–12 (2019).
449   27.   Young, E. *et al.* Identification of Dengue Virus Serotype 3 Specific Antigenic Sites
450   Targeted by Neutralizing Human Antibodies. (2020)
451   doi:10.1016/j.chom.2020.04.007.
452   28.   Gallichotte, E. N. *et al.* A new quaternary structure epitope on dengue virus
453   serotype 2 is the target of durable type-specific neutralizing antibodies. *mBio* **6**,
454   (2015).
455   29.   Barba-Spaeth, G. *et al.* Structural basis of potent Zika–dengue virus antibody
456   cross-neutralization. *Nature 2016 536:7614* **536**, 48–53 (2016).
457   30.   Dejnirattisai, W. *et al.* A new class of highly potent, broadly neutralizing antibodies
458   isolated from viremic patients infected with dengue virus. *Nature Immunology*
459   *2014 16:2* **16**, 170–177 (2014).
460   31.   Kotaki, T. *et al.* An affinity-matured human monoclonal antibody targeting fusion
461   loop epitope of dengue virus with in vivo therapeutic potency. *Sci Rep* **11**, 1–14
462   (2021).
463   32.   de Alwis, R. *et al.* Dengue Viruses Are Enhanced by Distinct Populations of
464   Serotype Cross-Reactive Antibodies in Human Immune Sera. *PLoS Pathog* **10**,
465   e1004386 (2014).
466   33.   Rouvinski, A. *et al.* Covalently linked dengue virus envelope glycoprotein dimers
467   reduce exposure of the immunodominant fusion loop epitope. *Nat Commun* **8**,
468   (2017).

34. Rockstroh, A. *et al.* Recombinant Envelope-Proteins with Mutations in the Conserved Fusion Loop Allow Specific Serological Diagnosis of Dengue-Infections. *PLoS Negl Trop Dis* **9**, 1–12 (2015).

35. Slon-Campos, J. L. *et al.* A protective Zika virus E-dimer-based subunit vaccine engineered to abrogate antibody-dependent enhancement of dengue infection. *Nature Immunology 2019 20:10* **20**, 1291–1298 (2019).

36. Metz, S. W. *et al.* In Vitro Assembly and Stabilization of Dengue and Zika Virus Envelope Protein Homo-Dimers. *Scientific Reports 2017 7:1* **7**, 1–8 (2017).

37. Kudlacek, S. T. *et al.* Designed, highly expressing, thermostable dengue virus 2 envelope protein dimers elicit quaternary epitope antibodies. *Sci Adv* **7**, (2021).

38. Zhang, S. *et al.* A mutation in the envelope protein fusion loop attenuates mouse neuroinvasiveness of the NY99 strain of West Nile virus. *Virology* **353**, 35–40 (2006).

39. Tian, Y., Grifoni, A., Sette, A. & Weiskopf, D. Human T Cell Response to Dengue Virus Infection. *Front Immunol* **10**, 2125 (2019).

40. Ngono, A. E. & Shresta, S. Cross-reactive T Cell immunity to dengue and zika viruses: New insights into vaccine development. *Front Immunol* **10**, 1316 (2019).

41. Waickman, A. T. *et al.* Dissecting the heterogeneity of DENV vaccine-elicited cellular immunity using single-cell RNA sequencing and metabolic profiling. *Nature Communications 2019 10:1* **10**, 1–16 (2019).

42. Gallichotte, E. N. *et al.* Genetic Variation between Dengue Virus Type 4 Strains Impacts Human Antibody Binding and Neutralization. *CellReports* **25**, 1214–1224 (2018).

43. Halstead, S. B. & Marchette, N. J. BIOLOGIC PROPERTIES OF DENGUE VIRUSES FOLLOWING SERIAL PASSAGE IN PRIMARY DOG KIDNEY CELLS: STUDIES AT THE UNIVERSITY OF HAWAII. *Am J Trop Med Hyg* **69**, 5–11 (2003).

44. Juraska, M. *et al.* Viral genetic diversity and protective efficacy of a tetravalent dengue vaccine in two phase 3 trials. *Proc Natl Acad Sci U S A* **115**, E8378–E8387 (2018).

45. Messer, W. B. *et al.* Development and characterization of a reverse genetic system for studying dengue virus serotype 3 strain variation and neutralization. *PLoS Negl Trop Dis* **6**, (2012).

46. Tse, L. V. *et al.* Structure-guided evolution of antigenically distinct adeno-associated virus variants for immune evasion. *Proc Natl Acad Sci U S A* **114**, E4812–E4821 (2017).