Can increased prenatal exposure to thyroid hormones alter physiology and behaviour in the long-term? Insights from an experimental study in Japanese quails.

Kalle Aho[1,2], Antoine Stier[1,3,4], Tom Sarraude[1,5], Bin-Yan Hsu[1,6], Suvi Ruuskanen[1,7]

[1] Department of Biology, University of Turku FI-20014 Turku, Finland
[2] School of Medicine, Faculty of Health Sciences, University of Eastern Finland, Kuopio, Finland
[3] Univ Lyon, Université Claude Bernard Lyon 1, CNRS, ENTPE, UMR 5023 LEHNA, F-69622, Villeurbanne, France
[4] Institut Pluridisciplinaire Hubert Curien, UMR7178, Université de Strasbourg, CNRS, Strasbourg, France
[5] Groningen Institute for Evolutionary Life Sciences, University of Groningen, the Netherlands
[6] Department of Life Science, Tunghai University, Taichung, Taiwan
[7] Department of Biological and Environmental Science, University of Jyväskylä, Finland

Corresponding Authors:
Kalle Aho[1]
Department of Biology, University of Turku FI-20014 Turku, Finland
Email address: ahokal9@gmail.com; Suvi Ruuskanen: suvi.k.ruuskanen@jyu.fi

Abstract

Maternal thyroid hormones (triiodothyronine T3 and thyroxine T4) are important regulators of embryonic development and gene expression. While maternal thyroid disorders are known to cause developmental issues in humans, variation in maternal thyroid hormones in healthy mothers have also been found to correlate with infant and child phenotypes. This suggests a potential impact of maternal thyroid hormones on offspring phenotype in an eco-evolutionary context. In chickens, prenatal thyroid hormone treatment has been shown to influence embryonic gene expression and postnatal treatment to influence imprinting and learning. However, the potential long-term effects of maternal thyroid hormones on physiology and behaviour are unclear. This study aims to investigate the long-term effects of maternal thyroid hormones on behaviour, plasma thyroid hormone levels and brain gene expression using the Japanese quail (*Coturnix japonica*) as a model. Egg hormone levels were elevated by injecting unincubated eggs with either saline (control), T3, T4 or a mixture of T3 and T4. Social motivation, boldness and fearfulness to predators were tested shortly after hatching and as adults. Plasma thyroid hormone levels and pallial expression of thyroid hormone receptor A, type 2 deiodinase, and nuclear receptor coactivator 1 were measured in adulthood. We found no evidence that elevated thyroid hormone levels in eggs affected behaviour, plasma hormone levels, or gene expression. This is the first study examining the potential long-term effects of elevated maternal thyroid hormones within the natural range. Although we found no evidence of long-term effects, other traits may still be affected and remain to be studied.

Introduction

Maternal effects refer to the influence from maternal phenotype on their
offspring's phenotype apart from genomic inheritance (Mousseau and Fox, 1998; Moore
et al., 2019; Yin et al., 2019). This influence can be mediated by hormones passed from
the mother to the offspring. As the levels of maternal hormones are influenced by the
maternal environment, they can transmit information about the maternal environment to
the offspring. This allows the development of the offspring phenotype to be shaped by
environmental factors apart from the inherited genome (Marshall and Uller, 2007;
Mousseau and Fox, 1998). Potentially, if offspring environment matches with maternal
environment, such hormone-mediated maternal effects can have a great adaptive
potential (sensu "anticipatory maternal effects", Marshall and Uller, 2007). Over the
past decades, effects of maternally-transferred hormones on offspring phenotype have
been observed in a wide range of traits across the whole animal kingdom, including
physiology and behaviour (e.g. Brown et al 2014; Dantzer et al. 2013; Groothuis et al.,
2019, 2005; Ruuskanen and Hsu, 2018), and can have long-lasting effects into
adulthood with potential fitness consequences (e.g. metabolic rate, predator-avoidance
behavior, survival, reproduction; Ruuskanen & Laaksonen 2010, 2012: Groothuis et al.,
2005; Schoech et al., 2011; von Engelhardt and Groothuis, 2011). Nevertheless, a recent
meta-analysis (Mentesana et al., 2025) found nearly no overall effect, in contrast with
previous studies (Podmokła et al., 2018; Moore et al. 2019), suggesting that more
needsto be done in order to better understand hormone-mediated maternal effects
(Groothuis, Hsu, Kumar 2020).

Compared to the more extensively studied androgens and glucocorticoids
(Podmokła et al., 2018), the effects of maternally transferred thyroid hormones (THs)
have been less explored (Darras, 2019; Ruuskanen and Hsu, 2018). THs
(triiodothyronine, T3, and thyroxine, T4) play important roles in metabolism and
development, where they regulate the expression of numerous genes (Bernal, 2017;
Darras, 2019; Moog et al., 2017). TH action is regulated by several components of the
thyroid system. TH receptors (THRA and THRB) and coactivators control the
transcription of TH-responsive genes by forming a complex with the active form of the
hormone, T3 (Brent, 2012). The intracellular availability of T3 is regulated by several
transmembrane TH transporters (e.g. MCT8, OATP1c1) that differentially transport
either T4 or T3 (Groeneweg et al., 2020). Inside the cell, T3 availability can be further
regulated by deiodinase enzymes that either convert T4 to T3 (deiodinase 2, D2) or
inactivate both T3 and T4 (deiodinase 3, D3) (Gereben et al., 2008). Interestingly, the
expression of these regulatory components (TH receptors, transporters and deiodinases)
begins as early as embryonic day 4 in chickens (Flamant and Samarut, 1998) and
embryonic day 1 in passerines (Ruuskanen et al. 2022), allowing the embryo to regulate
TH levels in an organ and cell-specific manner (Darras, 2019; Darras et al., 2011;
Flamant and Samarut, 1998; Ruuskanen et al., 2022; Van Herck et al., 2012) , while the
development of the thyroid gland occurs later in development, around mid-incubation
(Forhead and Fowden, 2014; McNichols and McNabb, 1988; Porazzi et al., 2009). This
suggests that these molecular components are in place to regulate maternally-transferred
hormones. Indeed, in almost all vertebrates studied so far, proper embryonic

development depends on maternally transferred THs (Brown et al., 2014; Darras, 2019;
Moog et al., 2017; Patel et al., 2011).
In mammals, maternal THs are transferred across the placenta, whereas in egg-
laying animals such as birds, maternal THs are deposited in the egg yolk. In birds, high
natural variation in egg TH levels has been observed, up to 5-fold in magnitude
(Ruuskanen and Hsu, 2018). This variation may be due to environmental factors such as
food availability or temperature (Hsu et al., 2016; Ruuskanen et al., 2016c), and has
been reported at both the inter- and intra-individual levels (Hsu et al., 2016, 2019b;
Ruuskanen et al., 2016a, 2016c). As THs are important in development (Darras et al.,
2009; Decuypere et al., 2005; McNabb, 2007; Wilson and McNabb, 1997), and as
increasing yolk TH levels has been found to increase embryonic TH levels (Van Herck
et al., 2012), high variation in egg TH levels could therefore be a source of offspring
phenotypic variation.
To date, most of what is known about phenotypic differences due to maternal
THs comes mainly from biomedical studies (Moog et al., 2017). However, studies using
pathological TH concentrations do not provide relevant information on the ecology or
evolution of maternal THs. Therefore, the effects of maternal THs within the natural
range need to be studied to understand whether variation has effects on traits affecting
offspring fitness. In addition, experiments manipulating maternal THs in mammals
would inevitably interfere with maternal physiology, making it difficult to attribute
results solely to maternal THs. Egg-laying animals, such as birds, have therefore
become common models for studying the functions of maternal hormones since
maternal THs deposited in eggs can be manipulated directly and independently. So far,
elevated TH levels in the egg within the natural range have been found to affect
hatching success, offspring growth, chick levels of T3 and telomere length in altricial
bird species (Hsu et al., 2023a, 2019a, 2017; Ruuskanen et al., 2016b; Sarraude et al.,
2020; Stier et al., 2020). Nevertheless, some of the reported effects appear to be
transient during the early post-hatching period (Hsu et al., 2019a). Whether the effects
of maternal THs can be long-lasting and potentially influence offspring fitness remains
unclear. In terms of precocial bird species, previous studies studying maternal THs were
scarce and typically applied supra-physiological doses (e.g., Wilson and McNabb, 1997;
van Herck et al., 2012).
To investigate the potential long-term effects of variation in maternal THs, we
experimentally elevated egg TH levels within the natural range in Japanese quail
(*Coturnix japonica*). With these experimental quails, we have previously reported a
higher hatching success caused by the elevated yolk TH levels, but no major effects on
morphology or physiology (Sarraude et al., 2020). Now we further studied changes in
chick and adult behaviour and adult plasma TH levels. In addition, we studied brain
gene expression at a later age. While increasing maternal TH levels beyond normal
physiological range (maternal hyperthyroidism) has been shown to affect behaviour, for
example both decreasing and increasing anxiety (Laureano-Melo et al., 2019; Zhang et
al., 2010, 2008), it is not known whether increasing maternal THs within the natural
range affects similar or linked behaviors. Also, as postnatal THs have been shown to
influence filial imprinting and learning behaviour (Yamaguchi et al., 2012), it is
interesting to study whether maternal THs, acting earlier in development, may also have

similar behavioural effects either in chicks or in adulthood. The underlying mechanistic
links between thyroid hormones and anxiety are expected to link TH-induced changes in
early brain development (in humans, for example altered hippocampal neurotransmitter
release or other neurochemical changes, even changes in neurogenesis, summarized in
Laureano-Melo et al. 2019 Horm Behav). We therefore studied traits related to anxiety
(Kumar et al. 2013) in our animal model, using existing validated protocols testing
boldness/fearfulness and social stress (Kumar et al. 2013, Forkman et al. 2007). Such
traits related to boldness and fearfulness are very relevant in any animal model as they
have clear links to for example foraging success and risk of predation. Potential
behavioural effects were investigated in both chicks and adults, allowing for a broader
timescale. Due to the lack of studies with natural levels of elevated maternal THs and
inconsistent results from studies on maternal hyperthyroidism and offspring anxiety
(Laureano-Melo et al., 2019; Zhang et al., 2010, 2008), the direction of potential
behavioural changes is difficult to predict. In addition, plasma TH levels were measured
in adults, and changes in these would indicate long lasting effects on TH function.
While plasma TH levels have been reported to be affected by elevated egg TH levels
(within the natural range) in a sex-dependent manner in nestling rock pigeons (Hsu et
al., 2017), some more recent studies in wild passerines found no clear effects of higher
natural levels of egg THs on nestling plasma THs (Hsu et al., 2023b, 2020). In terms of
gene expression, previous studies have examined the effect of the natural variation in
maternal THs on brain gene expression only in bird embryos or young chicks (Van
Herck et al., 2015, 2012; Yamaguchi et al., 2012) focusing on developmental effects.
However, the potential long-term effects on TH action in the adult brain remain
unknown. Here we studied several genes involved in TH function: THRA (thyroid
hormone receptor A), NCOA1 (nuclear receptor coactivator 1) and DIO2 (type 2
deiodinase) in the pallium of adults. Expression of these genes has been shown to be
influenced by THs in other species (Misiti et al., 1998; Walpita et al., 2007). As these
genes are part of the cellular pathway of TH action, changes in their expression could
indicate long-term changes in the regulation of TH action in the brain, and
consequently, brain function.

Materials and methods

Materials

This study was conducted in accordance with the Finnish legislation and approved by
the Finnish Animal Ethics Committee (Eläinkoelautakunta; ESAVI / 1018 / 04.10.07 /
2016). Japanese quail (*Coturnix japonica*), a commonly used laboratory model with a
reference genome, was used. This study was done with same individuals previously
used in Sarraude et al. (2020).

Parental generation and egg collection

The parental generation was composed of adult quails that were supplied by private
breeders in Finland and kept in two acclimatised rooms. Twenty-four breeding pairs
were formed by pairing birds from different breeders. Individuals were identified using
metal leg rings. The floor was covered with 3–5 cm of sawdust bedding, and a hiding

place, sand and calcium grit were provided. Each pair was housed in an indoor aviary
divided into pens with a floor area of 1 m². The ambient temperature was maintained at
20°C, and the photoperiod was set to 16 light hours (06:00 am-22:00pm, +3UCT) and 8
dark hours. The birds were fed ad libitum (Poultry Complete Feed, "Kanan Paras
Täysrehu", Hankkija, Finland), and their water was replaced daily. Pairs were
monitored every morning to collect eggs for 7 days (longer storage time is known to
decrease hatchability). The eggs were marked with a non-toxic marker, weighed, and
stored in a climate-controlled chamber at 15°C and 50% relative humidity. On the final
day of the collection period, a solution was injected into a total of 4–8 eggs per pair (the
variation was due to not all females producing an egg every day, average = 7, standard
deviation = 1)
204
Treatment of eggs, rearing of chicks and collection of samples
We manipulated egg TH levels by injecting hormones directly into the eggs prior to
embryonic development at a dose that mimicked increased maternal transfer (see details
in Sarraude et al., 2020). Eggs from each breeding pair were randomly assigned to four
treatment groups: T3, T4, T3 + T4 and control (0.9% saline), 24 clutches for all groups
except T3 that had 22 clutches. We used saline control following previous studies on
injecting THs into eggs (van Herck et al. 2015), and because THs are not very well
soluble in oil. Eggs laid by the same pair were always evenly distributed among all
treatment groups. The injected hormone solutions were prepared from crystallised
powder (T3: CAS number 6893-02-3, Sigma-Aldrich, St. Louis, MO, USA and T4:
CAS number 51-48-9, Sigma-Aldrich, St. Louis, MO, USA). The amount of hormone
injected (T4: 8.9 ng/egg, equivalent to 1.79 pg/mg yolk; T3: 4.7 ng/egg, equivalent to
1.24 pg/mg yolk) was based on previous measurements in the parental generation (mean
± SD: T4, 15.3 ± 4.43 ng / egg; T3, 7.58 ± 2.32 ng / egg) and the aim was to increase
hormone contents in the yolk by two standard deviations (as recommended by
Podmokla et al., 2018). We were unsure how much of T4 is converted to T3 and
therefore used 2SD of both hormones for the T3T4 group. The injection protocol has
been described in detail in Sarraude et al. (2020). Briefly, each egg was injected with 50
µl of either the respective hormone solution (T3, T4 or T3+T4, in 0.9% saline) or 0.9%
saline (control, CO), delivered by using 27G insulin syringes (BD Microlance^TM). After
sealing the hole with sterile plaster (OPSITE Flexigrid, Smith & Nephew), the eggs
were incubated artificially (37.8 °C and 55% relative humidity). A total of 158 eggs
were treated (39 T3, 39 T4, 40 T3+T4 and 40 CO), of which 66 chicks hatched (15 T3,
20 T4, 21 T3 + T4 and 10 CO), with sex distributions of chicks surviving to adulthood
(females / males): 7/4 for T3, 10/8 for T4, 9/12 for T3 + T4 and 3/4 for CO. Breeding
pairs/pairs of siblings per treatment: CO 7/0, T3 9/2, T3T4 15/4 and T4 13/2. Sex was
determined at maturity by examining the presence of the cloacal gland. The overall
hatching rate (~40%) was consistent with previous studies in domestic quail
(Okuliarová et al., 2007).
Twelve hours post-hatching, the chicks were marked with a unique combination
of coloured rings and nail coding, and transferred to two cages (each with a floor area of
1 m2 and a height of 30 cm). The cages contained approximately 30 chicks, sex and
treatments mixed. The chicks were provided with heating mats and lamps as

supplementary heat sources for the first two weeks. The chicks were fed with sieved
commercial poultry feed ("Punaheltta paras poikanen", Hankkija, Finland) and provided
with calcium and bathing sand. Two weeks after hatching, the chicks were separated
into four 1 m2 cages (ca. 30 cm high) of approximately 16 individuals. Approximately
three weeks after hatching, the coloured rings were replaced with unique metal rings. At
week 4 after hatching, the birds were transferred to eight pens of 1 m2 floor area
(average of 7.1 birds/pen, range = 4–9), under the same conditions as the parents. At
around the age of sexual maturity (ca. 6–8 weeks after hatching), the birds were
separated by sex in twelve 1 m2 pens (average of 4.8 birds/pen, range = 4–5).
Photoperiod was 16L:8D for the first 7 months and 8L:16D thereafter. At approximately
4.5 months of age, we collected blood samples (200 µl) from the brachial vein,
separated plasma, and stored it at -80° C. As Japanese quails have reached adulthood at
4.5 months, this time point would have already shown if there are long-lasting,
organizational effects of prenatal THs effects into adulthood. All birds were euthanised
by decapitation at 10 months of age. After decapitation, brain tissue was collected for
the gene expression analysis from the pallium (average 44.0 mg, std 2.8 mg, taken at the
surface of the region) in PBS buffer, snap frozen in liquid nitrogen and stored at -80° C.
While differences in photoperiod limit comparisons between gene expression results
and other results, our aim was to study whether there are persistent effects of prenatal
THs independent of any environmental factors that individuals may face later in life, for
which our experimental design is sufficient.
259

Behavioural experiments
To assess the potential effects of maternal THs on behaviour, we conducted a series of
behavioural tests (open arena, emergence and tonic immobility tests) at different life
stages. We followed previously published protocols that have been used extensively
with many different model organisms. The tests were conducted in separate rooms
where the birds had no visual or auditory contact with other individuals. The exception
was the open arena test, where the subject was able to hear the vocalisations of another
bird being tested at the same time in another room. All experiments were recorded, and
the videos were analysed blindly of the treatment. All videos of the same test at both
ages (chick/adult) were analysed by one person, except for the open arena test, where
one person analysed chick videos and another person analysed adult videos. This may
have confounded the age difference in behavior scores with the observer effect. Since
the videos from different treatments at each age were analysed by the same person, the
fact that chick and adult videos were analyzed by different people should not confound
the effect of prenatal TH exposure.
The open arena test measured social motivation/stress and boldnes/fearfulness and
it followed a setup developed by Forkman et al. (2007). The test was performed once on
both sexes at three days (F/M/undetermined: CO 3/3/1, T3 6/3/2, T3T4 8/12, T4 8/8)
and once approximately 4.5 months of age (F/M: CO 3/4, T3 8/3, T3T4 9/12, T4 9/8).
In the test, the bird was placed in a circular illuminated arena and the movement
duration and the number of escape attempts (attempt to jump/fly over the wall) were
measured from videos for 10 minutes (from the time the researcher left the room). Only
10 of the 56 adults (1 CO, 2 T3, 2 T4 and 5 T3T4) attempted to escape, so adult escape

attempts were not analysed. Subjects who moved less and made fewer escape attempts
were interpreted as less bold and less socially motivated.
The emergence test measured social motivation and boldness/fearfulness
(especially the timidity aspect, see e.g Davis et al. 2008). The test was performed on
both sexes, twice on 6-9d old chicks (as there is less data on repeatability, Forkman et
al. 2007; F/M/undetermined: CO 3/4/1, T3 7/4/1 T3T4 9/12, T4 10/8) and once on 4.5
months old adults (F/M: CO 3/4, T3 7/4, T3T4 8/11, T4 9/7) . The test followed a setup
developed by Forkman et al. (2007), in which a bird was placed in a dark chamber
connected to another, bright chamber. One minute after the subject was placed in the
dark chamber, a door connecting the chambers was removed. After removing the door,
the researcher left the observation room and the time it took for the subject to
completely exit the dark chamber was measured. Subjects who left the dark chamber
more quickly were interpreted as bolder and more socially motivated. The test lasted
five minutes. Whether the subject remained in the dark chamber (0) or left it (1) during
the test was also recorded.
The tonic immobility test (Mills and Faure, 1991) measured fearfulness to
predators (Forkman et al 2007). The test was performed on once both sexes at 12 days
(F/M/undetermined: CO 3/4, T3 7/4/1, T3T4 9/12, T4 10/8) and once approximately 4.5
months of age (F/M: CO 3/4, T3 7/4, T3T4 8/12, T4 9/7). To induce immobility, the
bird was placed on its back in the centre of a V-shaped test apparatus and restrained by
hand for 15 seconds, after which the hand was removed, and the time taken for the bird
to resume a standing posture was measured. If the subject resumed the standing posture
in less than 10 seconds, this was recorded as a failed induction and the induction
procedure was repeated. Four adult subjects who failed five times were excluded. If the
subject remained stationary for more than 10 seconds, then the time to recovery was
measured. The longer a subject took to recover, the more fearful it was interpreted. The
test lasted 10 minutes. Five subjects (1 chick and 4 adults) did not recover for 10
minutes and were excluded from the analysis (approximately 4% of all tests).
311
Plasma thyroid hormone levels
To test whether elevated maternal THs affect plasma TH levels later in life (4.5 months
of age), we extracted THs from plasma samples and measured them using a previously
validated LC-MS / MS protocol (Ruuskanen et al., 2018, where CV <10%). Due to
practical limitations (*i.e.* volume of plasma available and haemolysis in some samples),
the sample size was reduced to 48 individuals (F/M: T3 6/4, T4 8/8, T3T4 6/9 and 3/4 in
the control). Hormone concentrations are expressed in pmol/ml plasma. The
repeatability of the TH measurements, was validated in Ruuskanen et al. (2018) with
CV% less than 10% (3.5-9.7%, depending on the hormone and sample type)
321
Gene expression
To test for potential long-term changes in the expression of TH related genes (*DIO2*,
DIO3, *THRA*, and *NCOA1*), we measured gene expression from pallium samples (10
months old). RNA was extracted from the homogenised pallium samples using the
NucleoSpin RNA Plus kit (MACHEREY-NAGEL). Since the sample size of the control
group was the main factor limiting statistical power, we decided to analyse all control

individuals and only an equivalent number of individuals (randomly selected) in the
three treatment groups (N = 31, 3F/4M in control and 4F/4M per other treatment
groups). Extraction was done from 50 µl of the homogenised brain sample according to
the manufacturer's instructions, except that the gDNA removal column was centrifuged
longer (1 min, 11000 g) and the RNA was eluted once with 40 µl of Rnase-free water.
RNA concentration and purity were quantified by optical density, and all samples not
meeting our quality criteria (i.e. RNA concentration > 30 ng/µl and 260/280 > 1.80)
were excluded from further analysis. RNA integrity was checked using an E-Gel 2%
electrophoresis system (Invitrogen), and the ribosomal RNA 18S vs. 28S band intensity
and was deemed satisfactory (the visual inspection revealed the expected two bands of
rRNA with 28S band being > 2 times more intense than 18S band). Samples were stored
at -80° C for 2 weeks prior to cDNA synthesis. cDNA synthesis was performed using
the SensiFAST ™ cDNA Synthesis Kit (Bioline) according to the manufacturer's
instructions. The reaction was performed using 500 ng of RNA as the starting material.
A control (noRT) was prepared without reverse transcriptase enzyme. The cDNA was
diluted with water to a concentration of 1.2 ng/µl and stored at -20° C. The 1.2ng/uL
was calculated from RNA pre-RT, and slight variations in cDNA starting quantity (due
to differences in RT efficiency) are controlled for using the reference genes when
calculating relative expression.
GAPDH and *PGK* were used as control genes, using primers previously designed
and validated in Japanese quail (Vitorino Carvalho et al., 2019). Stability of expression
between experimental groups was further verified using geNorm software (geNorm M <
0.5, geNorm V < 0.15), and thus the geometric mean of these two genes was used as our
normalization factor. Primers for the target genes were designed using the Primer-
BLAST tool (NCBI). The primer pairs (Table 1) were designed to have at least one
intron between them. The suitability of the primers in qPCR was tested prior to the
actual analysis using different primer concentrations and different amounts of cDNA.
The tests confirmed the sufficient amplification efficiency of the primers (between 90
and 110% based on the standard curve) and the formation of only one product of the
expected size based on both the presence of a single narrow peak in the melting curve,
and the presence of a single band of the expected size on the agarose gel. *DIO3* had
extremely poor amplification, meaning that we could not accurately quantify it, and was
excluded from the analysis.
qPCR assays were performed on a Mic qPCR (Bio Molecular Systems) in a final
volume of 12 µL containing 6 ng of cDNA, 800 nM of forward and reverse primers and
SensiFAST SYBR Lo-ROX (Bioline). We ran gene-specific qPCR plates and each
sample was analysed in duplicate. Two plates were run for each gene and experimental
groups and sex were always balanced within each plate. We used a cDNA reference
sample (*i.e.* ratio = 1), which was a pool of 10 different individuals on each plate. We
used a two-step cycling with the following conditions: 2 min at 95°C; then 45 cycles of
5s at 95°C followed by 20s at 60°C (fluorescence reading) for all reactions, as
recommended by manufacturer. The expression of each gene was calculated as
$(1+Ef_{Target})^{\Delta Cq(Target)}$/ geometric mean $[(1+Ef_{GAPDH})^{\Delta CqGAPDH}; (1+Ef_{PGK1})^{\Delta CqPGK1}]$, where
Ef is the amplification's efficiency and ΔCq is the difference between the Cq-values of
the reference sample and the sample of interest. Amplification in the no-RT controls

never occurred until at least 8 cycles after the lower Cq sample, and thus contamination
by genomic DNA could not interfere with our results. Technical precision and
repeatability were generally good (Table 1), but a few samples with a CV% of
duplicates (based on the final ratio) greater than 15% were excluded from the analysis
(final sample size: $N_{DIO2} = 27$, $N_{NCOA1} = 28$, $N_{THRa} = 26$)
378

Statistical analysis
All statistical analyses were performed with R software (version 3.6.2), using the
packages lme4 version 1.1-23 (Bates et al., 2015) and emmeans version 1.6.0 (Lenth,
2021). Normality and homoscedasticity of the residual distributions were checked
visually (quantile-quantile and residuals against fitted values plots)with additionally
confirmation by Shapiro-Wilk test of normality. For the behavioural analysis, recovery
time (tonic immobility) and exit time (emergence test), were natural log-transformed to
ensure that the residuals were normally distributed. Possible outliers were tested with R
using the *simulateResiduals* function in DHARMa package (Hartig 2024).
388

Linear mixed models (LMM, *lmer* function in lme4) were used to analyse time to exit in
the emergence test, movement duration and number of escape attempts in the open
arena test, recovery time in the tonic immobility test, plasma TH concentrations and
qPCR results. Generalised linear mixed model (GLMM, g*lmer* function in lme4) with a
binomial error distribution was used to analyse the proportion of individuals leaving the
dark chamber in the emergence test. Fixed effects were treatment, sex, and test
number/age (in repeated tests). Interactions between sex and treatment, and between
treatment and age (in repeated tests), were also analysed. Random effects were
individual ID (in repeated tests) and parental ID (to account for relatedness between
siblings in all experiments). F- and p-values for LMMs were obtained using the
Kenward-Roger approximation on the denominator degree of freedom (ddf),
implemented in the package *lmerTest* (*anova* function) (Kuznetsova et al., 2017). For
binomial regression models, main effects and interactions were tested by comparing the
model with and without the factor. For all models, we removed non-significant (p >
0.05) interactions to ensure correct interpretation on the main effects. Post-hoc tests
were performed using Tukey tests (*emmeans* function from emmeans package, Lenth,
2021) where appropriate.
406

Results

Results
Effects of prenatal thyroid hormone supplementation on behaviour
In general, no statistically significant differences in chick or adult behavioural traits
were observed between the egg TH manipulation groups (T3, T4, T3 + T4, control).
Elevated THs had no significant effect on social motivation or boldness (Table 2, Fig. 1-
2), nor on fearfulness to predators (Table 2, Fig. 3). A statistically significant interaction
between treatment and age was found in the emergence test (F = 3.58, ddf = 67.2, p =
0.018, Fig. 2): In all treatments except the control group (Tukey t=0.59, p=0.56), chick
exit times were significantly longer than those of adults (all t > 3.21, p < 0.002) while
no differences between treatments were found within each age-category (all t < 2.48, all

p> 0.07). There was also a statistically significant interaction between treatment and sex
in the tonic immobility test (F = 2.92, ddf = 37.5, p = 0.047, Fig. 3): In males, the
recovery time was longer in the T4-group (marginal mean ± SE: 177.1 ± 47.8 s) than in
the T3-group (marginal mean ± SE: 56.1 ± 20.8 s, Tukey t = 2.71, p = 0.047), whereas
no differences between treatments were observed in females (all t < 2.29, all p >0.12).
In the T4-group, the recovery time of males (marginal mean ± SE: 177.1 ± 47.8 s) was
longer than that of females (marginal mean ± SE: 85.2 ± 19.9 s) (t=2.16, p= 0.037),
while no sex differences were observed within the other treatments (all t<1.80, all
p>0.079).
426
Effects of prenatal thyroid hormone supplementation on plasma thyroid
hormone levels
Increasing TH levels in eggs did not significantly affect plasma TH levels in 4.5-month-
old birds (T3: F = 0.13, ddf = 37.7, p = 0.94, T4: F = 2.1, ddf = 41.5, p = 0.12, Fig. 4).
Treatment also had no sex-dependent effect on plasma T3 or T4 (treatment-sex
interactions, p > 0.36).
433
Effects of prenatal thyroid hormone supplementation on pallial gene
expression
Increasing TH levels in eggs did not affect pallial gene expression of TH-related genes
in 10-month-old birds (*DIO2*: F = 2.49, ddf = 17.3, p = 0.09, *NCOA1*: F = 0.092, ddf =
21.1, p = 0.96, *THRA*: F = 0.47, ddf = 18.9, p = 0.71, Fig. 5). Treatment also did not
have sex-dependent effect on gene expression (treatment-sex interactions, p > 0.26).
Females had higher expression of *NCOA1* (F=7.68, p=0.012) and *THRA* (F=5.17,
p=0.034) than males.
442

Discussion

In this study, we tested for long-term changes in behaviour, plasma TH levels
and pallial gene expression following increased exposure to maternal THs. In terms of
behaviour, we observed no significant changes in boldness, social motivation or
fearfulness to predators in chicks (3-12 d) or adults (4.5 months) when the egg TH level
was increased. To our knowledge, there are no similar previous studies, and it is
therefore difficult to assess whether there may have been changes in other behavioural
traits. The detection of changes was also partly limited by the small sample size, which
meant that we may not have had enough statistical power to detect subtle changes in
behaviour. Nevertheless, we found some age-dependent and sex-dependent effects of
TH treatment on behaviour. The age-dependent effect in the emergence test suggests
that elevated maternal TH leads to a decrease in boldness and social motivation with age
(from chick to adult), whereas such a decrease is not observed in controls. However, the
lack of differences between control and treatment within a given age suggests that such
an effect (if not associated with type I statistical error) is rather mild. The sex-dependent
effect in the tonic immobility test also seems relatively unconvincing because no sex
differences were observed within treatments, nor did T3T4 females or T4 males differ

from the same-sex controls. Given the important role of THs in brain development, we
expected that elevated maternal THs would have led to behavioural changes. Yet, our
data suggest that an experimental increase in maternal THs within the natural range has
no clear and large effect on the behavioural traits measured. However, in contrast to
maternal THs, post-hatch TH treatment twenty hours after hatching has been shown to
affect imprinting and learning in four-day-old chicks (Yamaguchi et al., 2012). The
discrepancy between our results and Yamaguchi et al. (2012) could suggest that the
sensitive period during which THs can influence behaviour might occur at a later timing
of development, but this requires further investigation.
We found that elevated maternal THs did not significantly affect plasma TH
levels in adulthood (4.5 months of age), although transient effects prior to measurement,
or downregulation of THs,cannot be ruled out. This was the first study to test for long-
term changes, whereas previously changes in response to elevated prenatal THs have
been observed at younger ages: Experimental elevation of egg THs was found to
increase male and decrease female plasma TH levels in rock pigeons at 14 days of age
(Hsu et al., 2017), but such effects were not observed in two other wild passerine
species (Hsu et al., 2023b, 2020). In contrast to these studies in altricial species, not
only was this study conducted on a precocial species, but our measurements were also
made at a different life stage (adulthood). Overall, this suggests that an increase in
maternal THs may only lead to transient changes in offspring plasma TH levels and that
the effects may not be universal across species.
We also found no significant effect of elevated maternal THs on the expression
of *THRA*, *DIO2* or *NCOA1* in the pallium of adult (10 months old) individuals. Since
we only collected samples from 10-month-old birds, we cannot rule out possible
changes at younger ages. Changes in gene expression have been observed in other TH-
related genes, but, at much shorter time spans: In chickens, injection of THs into the
yolk decreased the expression of the TH-transporter *OATP1C1* and increased the
expression of the TH-transporter *MCT8* in 4-day-old embryos (Van Herck et al., 2012).
The expressions of our target genes, *THRA*, *DIO2* and *NCOA1,* have also been shown to
be at least transiently responsive to THs in other species: increase in *THRA* and decrease
in *DIO2* mRNA levels in zebrafish embryos (Walpita et al., 2007), and increase in
NCOA1 mRNA levels in the anterior pituitary of adult rats (Misiti et al., 1998). It is
therefore possible that the gene regulation by maternal THs is restricted to early
development when the embryo is not yet able to produce THs.
We have previously shown in these same experimental birds, that an increase in
prenatal THs influences hatching success (Sarraude et al., 2020). Therefore, the dose
used in this study can have some effects during embryo development, which suggests
that the lack of large changes in long-term behaviour, plasma TH levels and brain gene
expression may be linked to an attenuation of prenatal TH effects by embryonic
regulation. It is indeed possible that embryos could buffer the effect of maternal THs by
altering the expression of deiodinases which are expressed already at very early stages
(Ruuskanen et al., 2022; Too et al., 2017). Metabolisation of THs to inactive forms
could prevent the elevated egg THs from exerting any effects. However, given the
effects previously observed in these same birds on hatching success (Sarraude et al.,
2020), the lack of changes in this study may also be explained by the stage of ontogeny,

prenatal THs not leading to long-lasting effects. Nevertheless, the lack of prenatal
measurements means that this cannot be fully confirmed, and it remains possible that
elevated maternal THs have no effect on the measured traits in Japanese quails.
508

Conclusions

In conclusion, we found no evidence that elevated maternal THs cause substantial long-
term changes in behaviour, plasma TH levels or pallial gene expression. This is
consistent with our previous findings in the same experimental birds, where elevation of
prenatal THs (T3, T4, or in combination) did not lead to long-term effects on oxidative
stress, moult or reproduction (Sarraude et al., 2020). While behavioural changes have
not been previously studied, plasma TH concentrations and gene expression have been
shown to respond to elevated maternal THs in the short- term (Hsu et al., 2017; Van
Herck et al., 2012). However, based on this study, these changes would appear to be
limited to early development or shortly after hatching with the offspring's own TH
production later replacing the effects of elevated maternal THs. Maternal THs could
theoretically influence the offspring's own TH production at a later age by altering the
function of the HPT-axis, but no evidence has yet been found to support this hypothesis.
The long-term effects of variation in maternal THs remain unclear, but this study has
provided new information on some previously untested traits. Given the tight
association between THs and metabolism, it would be interesting to study behaviors
related to thermoregulation and metabolism in the future, Understanding long-term
effects in addition to short-term effects will help to better understand the evolutionary
and ecological importance of maternal THs.
528

Acknowledgements

530 We thank personnel at the animal care facilities, Marjo Myllyrinne, Simo Laine for their
531 help.
532

Funding

534 The study was funded by the Academy of Finland (project #286278) to SR, the Finnish
535 National Agency for Education (grant no. TM-15-9960 to TS). AS was supported by a
536 Turku Collegium for Science and Medicine fellowship and an European Commission
537 Marie Sklodowska-Curie Fellowship #894963.
538

Data availability statement

540 Data is available on Figshare (DOI: https://doi.org/10.6084/m9.figshare.22266211.v1).
541

Declaration of interests

543 The authors declare that they have no known competing financial interests or personal
544 relationships that could have appeared to influence the work reported in this paper.
545

References

Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects Models Using lme4. J Stat Softw 67. https://doi.org/10.18637/jss.v067.i01

Bernal, J., 2017. Thyroid hormone regulated genes in cerebral cortex development. Journal of Endocrinology 232, R83–R97. https://doi.org/10.1530/JOE-16-0424

Brent, G.A., 2012. Mechanisms of thyroid hormone action. Journal of Clinical Investigation 122, 3035–3043. https://doi.org/10.1172/JCI60047

Brown, C.L., Urbinati, E.C., Zhang, W., Brown, S.B., McComb-Kobza, M., 2014. Maternal Thyroid and Glucocorticoid Hormone Interactions in Larval Fish Development, and Their Applications in Aquaculture. Reviews in Fisheries Science & Aquaculture 22, 207–220. https://doi.org/10.1080/23308249.2014.918086

Darras, V.M., 2019. The Role of Maternal Thyroid Hormones in Avian Embryonic Development. Front Endocrinol (Lausanne) 10. https://doi.org/10.3389/fendo.2019.00066

Darras, V.M., Van Herck, S.L.J., Geysens, S., Reyns, G.E., 2009. Involvement of thyroid hormones in chicken embryonic brain development. Gen Comp Endocrinol 163, 58–62. https://doi.org/10.1016/j.ygcen.2008.11.014

Darras, V.M., Van Herck, S.L.J., Heijlen, M., De Groef, B., 2011. Thyroid Hormone Receptors in Two Model Species for Vertebrate Embryonic Development: Chicken and Zebrafish. J Thyroid Res 2011, 1–8. https://doi.org/10.4061/2011/402320

Davis KA, Schmidt JB, Doescher RM, Satterlee DG. Fear responses of offspring from divergent quail stress response line hens treated with corticosterone during egg formation. Poult Sci. 2008 Jul;87(7):1303-13. doi: 10.3382/ps.2008-00083.

Dantzer B, Newman AE, Boonstra R, Palme R, Boutin S, Humphries MM, McAdam AG. Density triggers maternal hormones that increase adaptive offspring growth in a wild mammal. Science. 2013 Jun 7;340(6137):1215-7. doi: 10.1126/science.1235765.

Decuypere, E., Van As, P., Van der Geyten, S., Darras, V.M., 2005. Thyroid hormone availability and activity in avian species: A review. Domest Anim Endocrinol 29, 63–77. https://doi.org/10.1016/j.domaniend.2005.02.028

Flamant, F., Samarut, J., 1998. Involvement of Thyroid Hormone and Its α Receptor in Avian Neurulation. Dev Biol 197, 1–11. https://doi.org/10.1006/dbio.1998.8872

Forhead, A.J., Fowden, A.L., 2014. Thyroid hormones in fetal growth and prepartum maturation. Journal of Endocrinology 221, R87–R103. https://doi.org/10.1530/JOE-14-0025

Forkman, B., Boissy, A., Meunier-Salaün, M.-C., Canali, E., Jones, R.B., 2007. A critical review of fear tests used on cattle, pigs, sheep, poultry

and horses. Physiol Behav 92, 340–374.
https://doi.org/10.1016/j.physbeh.2007.03.016

Gereben, B., Zeöld, A., Dentice, M., Salvatore, D., Bianco, A.C., 2008.
Activation and inactivation of thyroid hormone by deiodinases: Local
action with general consequences. Cellular and Molecular Life Sciences
65, 570–590. https://doi.org/10.1007/s00018-007-7396-0

Groeneweg, S., van Geest, F.S., Peeters, R.P., Heuer, H., Visser, W.E., 2020.
Thyroid Hormone Transporters. Endocr Rev 41, 146–201.
https://doi.org/10.1210/endrev/bnz008

Groothuis, T.G.G., Kumar, N., Hsu, B.-Y., 2020. Explaining discrepancies in
the study of maternal effects: the role of context and embryo. Current
Opinion in Behavioral Sciences 36, 185-192.
https://doi.org/10.1016/j.cobeha.2020.10.006.

Groothuis, T.G.G., Hsu, B.-Y., Kumar, N., Tschirren, B., 2019. Revisiting
mechanisms and functions of prenatal hormone-mediated maternal
effects using avian species as a model. Philosophical Transactions of the
Royal Society B: Biological Sciences 374, 20180115.
https://doi.org/10.1098/rstb.2018.0115

Groothuis, T.G.G., Müller, W., von Engelhardt, N., Carere, C., Eising, C.,
2005. Maternal hormones as a tool to adjust offspring phenotype in avian
species. Neurosci Biobehav Rev 29, 329–352.
https://doi.org/10.1016/j.neubiorev.2004.12.002

Hartig F (2024). _DHARMa: Residual Diagnostics for Hierarchical (Multi-
Level / Mixed) Regression Models_. R package version 0.4.7,
<https://CRAN.R-project.org/package=DHARMa>.

Hsu, B.-Y, Dijkstra, C., Darras, V.M., Vries, B., Groothuis, T.G.G., 2016.
Maternal adjustment or constraint: differential effects of food availability
on maternal deposition of macro⬜nutrients, steroids and thyroid
hormones in rock pigeon eggs. Ecol Evol 6, 397–411.
https://doi.org/10.1002/ece3.1845

Hsu, B.-Y., Cossin-Sevrin, N., Stier, A., Ruuskanen, S., 2023a. Prenatal
thyroid hormones accelerate postnatal growth and telomere shortening in
wild great tits. Journal of Experimental Biology.
https://doi.org/10.1242/jeb.243875

Hsu, B.-Y., Cossin-Sevrin, N., Stier, A., Ruuskanen, S., 2023b. Prenatal
thyroid hormones accelerate postnatal growth and telomere shortening in
wild great tits. Journal of Experimental Biology.
https://doi.org/10.1242/jeb.243875

Hsu, B.-Y., Dijkstra, C., Darras, V.M., de Vries, B., Groothuis, T.G.G., 2017.
Maternal thyroid hormones enhance hatching success but decrease
nestling body mass in the rock pigeon (Columba livia). Gen Comp
Endocrinol 240, 174–181. https://doi.org/10.1016/j.ygcen.2016.10.011

Hsu, B.-Y., Doligez, B., Gustafsson, L., Ruuskanen, S., 2019a. Transient
growth-enhancing effects of elevated maternal thyroid hormones at no

apparent oxidative cost during early postnatal period. J Avian Biol 50.

https://doi.org/10.1111/jav.01919

Hsu, B.-Y., Sarraude, T., Cossin-Sevrin, N., Crombecque, M., Stier, A.,

Ruuskanen, S., 2020. Testing for context-dependent effects of prenatal

thyroid hormones on offspring survival and physiology: an experimental

temperature manipulation. Sci Rep 10, 14563.

https://doi.org/10.1038/s41598-020-71511-y

Hsu, B.-Y., Verhagen, I., Gienapp, P., Darras, V.M., Visser, M.E.,

Ruuskanen, S., 2019b. Between- and Within-Individual Variation of

Maternal Thyroid Hormone Deposition in Wild Great Tits (Parus major

). Am Nat 194, E96–E108. https://doi.org/10.1086/704738

Kumar V, Bhat ZA, Kumar D. Animal models of anxiety: a comprehensive

review. J Pharmacol Toxicol Methods. 2013 Sep-Oct;68(2):175-183. doi:

10.1016/j.vascn.2013.05.003.

Kuznetsova, A., Brockhoff, P.B., Christensen, R.H.B., 2017. lmerTest

Package: Tests in Linear Mixed Effects Models. J Stat Softw 82.

https://doi.org/10.18637/jss.v082.i13

Laureano-Melo, R., Souza, J.S. de, da Conceição, R.R., Albuquerque, J.M.L.,

Rodrigues, N.C., Marinho, B.G., Olivares, E.L., Giannocco, G., Côrtes,

W. da S., 2019. Prenatal thyroxine treatment promotes anxiolysis in male

Swiss mice offspring. Horm Behav 108, 10–19.

https://doi.org/10.1016/j.yhbeh.2018.12.008

Marshall, D., Uller, T., 2007. When is a maternal effect adaptive? Oikos 116,

1957–1963. https://doi.org/10.1111/j.2007.0030-1299.16203.x

McNabb, F.M.A., 2007. The Hypothalamic-Pituitary-Thyroid (HPT) Axis in

Birds and Its Role in Bird Development and Reproduction. Crit Rev

Toxicol 37, 163–193. https://doi.org/10.1080/10408440601123552

McNichols, M.J., McNabb, F.M.A., 1988. Development of thyroid function

and its pituitary control in embryonic and hatchling precocial Japanese

quail and altricial Ring doves. Gen Comp Endocrinol 70, 109–118.

https://doi.org/10.1016/0016-6480(88)90099-8

Mentesana L, Hau M, D'Amelio PB, Adreani NM, Sánchez-Tójar A. Do Egg

Hormones Have Fitness Consequences in Wild Birds? A Systematic

Review and Meta-Analysis. Ecol Lett. 2025 Mar;28(3):e70100. doi:

10.1111/ele.70100.

Mills, A.D., Faure, J.-M., 1991. Divergent selection for duration of tonic

immobility and social reinstatement behavior in Japanese quail (Coturnix

coturnix japonica) chicks. J Comp Psychol 105, 25–38.

https://doi.org/10.1037/0735-7036.105.1.25

Misiti, S., Schomburg, L., M. Yen, P., Chin, W.W., 1998. Expression and

Hormonal Regulation of Coactivator and Corepressor Genes.

Endocrinology 139, 2493–2500. https://doi.org/10.1210/endo.139.5.5971

Moog, N.K., Entringer, S., Heim, C., Wadhwa, P.D., Kathmann, N., Buss, C.,

2017. Influence of maternal thyroid hormones during gestation on fetal

brain development. Neuroscience 342, 68–100.
https://doi.org/10.1016/j.neuroscience.2015.09.070

Moore MP, Whiteman HH, Martin RA. A mother's legacy: the strength of
maternal effects in animal populations. Ecol Lett. 2019 Oct;22(10):1620-
1628. doi: 10.1111/ele.13351.

Mousseau, T., Fox, C., 1998. The adaptive significance of maternal effects.
Trends Ecol Evol 13, 403–407. https://doi.org/10.1016/S0169-
5347(98)01472-4

Okuliarová, M., Škrobánek, P., Zeman, M., 2007. Effect of Increasing Yolk
Testosterone Levels on Early Behaviour in Japanese Quail Hatchlings.
Acta Veterinaria Brno 76, 325–331.
https://doi.org/10.2754/avb200776030325

Patel, J., Landers, K., Li, H., Mortimer, R.H., Richard, K., 2011. Thyroid
hormones and fetal neurological development. Journal of Endocrinology
209, 1–8. https://doi.org/10.1530/JOE-10-0444

Podmokła, E., Drobniak, S.M., Rutkowska, J., 2018. Chicken or egg?
Outcomes of experimental manipulations of maternally transmitted
hormones depend on administration method - a meta-analysis. Biological
Reviews 93, 1499–1517. https://doi.org/10.1111/brv.12406

Porazzi, P., Calebiro, D., Benato, F., Tiso, N., Persani, L., 2009. Thyroid
gland development and function in the zebrafish model. Mol Cell
Endocrinol 312, 14–23. https://doi.org/10.1016/j.mce.2009.05.011

Russell V. Lenth, 2021. emmeans: Estimated Marginal Means, aka Least-
Squares Means.

Ruuskanen, S., Doligez, B., Pitala, N., Gustafsson, L., Laaksonen, T. 2012.
Long-term fitness consequences of maternal hormones: sons pay the
costs. Funct Ecol 26, 884-894. https://doi.org/10.1111/j.1365-
2435.2012.01994.x. https://doi.org/10.1016/j.yhbeh.2009.09.017

Ruuskanen, S., Laaksonen, T. 2010. Yolk hormones have sex-specific long-
term effects on behavior in the pied flycatcher. Horm Behav 57, 119–127

Ruuskanen, S., Darras, V.M., de Vries, B., Visser, M.E., Groothuis, T.G.G.,
2016a. Experimental manipulation of food availability leads to short-
term intra-clutch adjustment in egg mass but not in yolk androgen or
thyroid hormones. J Avian Biol 47, 36–46.
https://doi.org/10.1111/jav.00728

Ruuskanen, S., Darras, V.M., Visser, M.E., Groothuis, T.G.G., 2016b. Effects
of experimentally manipulated yolk thyroid hormone levels on offspring
development in a wild bird species. Horm Behav 81, 38–44.
https://doi.org/10.1016/j.yhbeh.2016.03.006

Ruuskanen, S., Groothuis, T.G.G., Schaper, S. V., Darras, V.M., de Vries, B.,
Visser, M.E., 2016c. Temperature-induced variation in yolk androgen
and thyroid hormone levels in avian eggs. Gen Comp Endocrinol 235,
29–37. https://doi.org/10.1016/j.ygcen.2016.05.026

Ruuskanen, S., Hsu, B.-Y., 2018. Maternal Thyroid Hormones: An
Unexplored Mechanism Underlying Maternal Effects in an Ecological

Framework. Physiological and Biochemical Zoology 91, 904–916.
https://doi.org/10.1086/697380

Ruuskanen, S., Hsu, B.-Y., Heinonen, A., Vainio, M., Darras, V.M., Sarraude,
T., Rokka, A., 2018. A new method for measuring thyroid hormones
using nano-LC-MS/MS. Journal of Chromatography B 1093–1094, 24–
30. https://doi.org/10.1016/j.jchromb.2018.06.052

Ruuskanen, S., Hukkanen, M., Garcin, N., Cossin-Sevrin, N., Hsu, B.-Y.,
Stier, A., 2022. Altricial Bird Early-Stage Embryos Express the
Molecular "Machinery" to Respond to and Modulate Maternal Thyroid
Hormone Cues. Physiological and Biochemical Zoology 95, 544–550.
https://doi.org/10.1086/721556

Sarraude, T., Hsu, B.-Y., Groothuis, T., Ruuskanen, S., 2020. Testing the
short-and long-term effects of elevated prenatal exposure to different
forms of thyroid hormones. PeerJ 8, e10175.
https://doi.org/10.7717/peerj.10175

Schoech, S.J., Rensel, M.A., Heiss, R.S., 2011. Short- and long-term effects
of developmental corticosterone exposure on avian physiology,
behavioral phenotype, cognition, and fitness: A review. Curr Zool 57,
514–530. https://doi.org/10.1093/czoolo/57.4.514

Stier, A., Hsu, B.-Y., Marciau, C., Doligez, B., Gustafsson, L., Bize, P.,
Ruuskanen, S., 2020. Born to be young? Prenatal thyroid hormones
increase early-life telomere length in wild collared flycatchers. Biol Lett
16, 20200364. https://doi.org/10.1098/rsbl.2020.0364

Too, H.C., Shibata, M., Yayota, M., Darras, V.M., Iwasawa, A., 2017.
Expression of thyroid hormone regulator genes in the yolk sac membrane
of the developing chicken embryo. Journal of Reproduction and
Development 63, 463–472. https://doi.org/10.1262/jrd.2017-017

Van Herck, S.L.J., Delbaere, J., Bourgeois, N.M.A., McAllan, B.M.,
Richardson, S.J., Darras, V.M., 2015. Expression of thyroid hormone
transporters and deiodinases at the brain barriers in the embryonic
chicken: Insights into the regulation of thyroid hormone availability
during neurodevelopment. Gen Comp Endocrinol 214, 30–39.
https://doi.org/10.1016/j.ygcen.2015.02.021

Van Herck, S.L.J., Geysens, S., Delbaere, J., Tylzanowski, P., Darras, V.M.,
2012. Expression profile and thyroid hormone responsiveness of
transporters and deiodinases in early embryonic chicken brain
development. Mol Cell Endocrinol 349, 289–297.
https://doi.org/10.1016/j.mce.2011.11.012

Vitorino Carvalho, A., Couroussé, N., Crochet, S., Coustham, V., 2019.
Identification of Reference Genes for Quantitative Gene Expression
Studies in Three Tissues of Japanese Quail. Genes (Basel) 10, 197.
https://doi.org/10.3390/genes10030197

von Engelhardt, N., Groothuis, T.G.G., 2011. Maternal Hormones in Avian
Eggs, in: Hormones and Reproduction of Vertebrates. Elsevier, pp. 91–
127. https://doi.org/10.1016/B978-0-12-374929-1.10004-6

769   Walpita, C.N., Van der Geyten, S., Rurangwa, E., Darras, V.M., 2007. The
770   effect of 3,5,3'-triiodothyronine supplementation on zebrafish (Danio
771   rerio) embryonic development and expression of iodothyronine
772   deiodinases and thyroid hormone receptors. Gen Comp Endocrinol 152,
773   206–214. https://doi.org/10.1016/j.ygcen.2007.02.020
774   Wilson, C.M., McNabb, F.M.A., 1997. Maternal Thyroid Hormones in
775   Japanese Quail Eggs and Their Influence on Embryonic Development.
776   Gen Comp Endocrinol 107, 153–165.
777   https://doi.org/10.1006/gcen.1997.6906
778   Yamaguchi, S., Aoki, N., Kitajima, T., Iikubo, E., Katagiri, S., Matsushima,
779   T., Homma, K.J., 2012. Thyroid hormone determines the start of the
780   sensitive period of imprinting and primes later learning. Nat Commun 3,
781   1081. https://doi.org/10.1038/ncomms2088
782   Yin J, Zhou M, Lin Z, Li QQ, Zhang YY. Transgenerational effects benefit
783   offspring across diverse environments: a meta-analysis in plants and
784   animals. Ecol Lett. 2019 Nov;22(11):1976-1986. doi: 10.1111/ele.13373.
785   Zhang, L., Hernández, V.S., Medina-Pizarro, M., Valle-Leija, P., Vega-
786   González, A., Morales, T., 2008. Maternal hyperthyroidism in rats
787   impairs stress coping of adult offspring. J Neurosci Res 86, 1306–1315.
788   https://doi.org/10.1002/jnr.21580
789   Zhang, L., Medina, M.P., Hernández, V.S., Estrada, F.S., Vega-González, A.,
790   2010. Vasopressinergic network abnormalities potentiate conditioned
791   anxious state of rats subjected to maternal hyperthyroidism.
792   Neuroscience 168, 416–428.
793   https://doi.org/10.1016/j.neuroscience.2010.03.059
794

Figure 1. Movement duration for chicks (3 d) and adults (4.5 months) in the open arena test for different prenatal thyroid hormone treatment groups. CO = control, 50 µl 0.9% saline; T3: 4.7 ng/egg in 50 µl 0.9% saline, equivalent to 1.24 pg/mg yolk; T4: 8.9 ng/egg in 50 µl 0.9% saline, equivalent to 1.79 pg/mg yolk. The lines represent the medians of the treatments, and the bars represent the interquartile range of the original observations. Sample size (chicks/adults): CO 7/7, T3 11/11, T3T4 20/21 and T4 16/17.

Figure 2. Time to exit the dark chamber for chicks (6-9 d) and adults (4.5 months) in the emergence test for different prenatal thyroid hormone treatment groups. CO = control, 50 µl 0.9% saline; T3: 4.7 ng/egg in 50 µl 0.9% saline, equivalent to 1.24 pg/mg yolk; T4: 8.9 ng/egg in 50 µl 0.9% saline, equivalent to 1.79 pg/mg yolk. The test was performed twice for chicks and once for adults. Results for both chick studies are shown. The lines represent the medians of the treatments, and the bars represent the interquartile range of the original observations. Sample size (chicks/adults): CO 11/6, T3 15/9, T3T4 24/13 and T4 22/13.

Figure 3. Time to resume standing position for females and males, chicks (12 d) and adults (4.5 months) combined, in the tonic immobility test for different prenatal thyroid hormone treatment groups. CO = control, 50 µl 0.9% saline; T3: 4.7 ng/egg in 50 µl 0.9% saline, equivalent to 1.24 pg/mg yolk; T4: 8.9 ng/egg in 50 µl 0.9% saline, equivalent to 1.79 pg/mg yolk. The lines represent the medians of the treatments, and the bars represent the interquartile range of the original observations. Sample size (female/male: CO 6/8, T3 13/7, T3T4 14/23 and T4 18/14.

Figure 4. Plasma T3 and T4 concentrations in 4.5-month-old birds between different prenatal thyroid hormone treatment groups. CO = control, 50 µl 0.9% saline; T3: 4.7 ng/egg in 50 µl 0.9% saline, equivalent to 1.24 pg/mg yolk; T4: 8.9 ng/egg in 50 µl 0.9% saline, equivalent to 1.79 pg/mg yolk. The lines represent the medians of the treatments, and the bars represent the interquartile range of the original observations. Sample size (T3/T4 concentrations): CO 6/7, T3 10/10, T3T4 14/15 and T4 14/16.

Figure 5. Relative expressions of DIO2, THRA and NCOA1 in the pallium of 10-month-old birds between different prenatal thyroid hormone treatments. CO = control, 50 µl 0.9% saline; T3: 4.7 ng/egg in 50 µl 0.9% saline, equivalent to 1.24 pg/mg yolk; T4: 8.9 ng/egg in 50 µl 0.9% saline, equivalent to 1.79 pg/mg yolk. The lines represent the medians of the treatments, and the bars represent the interquartile range of the original observations. Sample sizes (CO/T3/T3T4/T4): DIO2 5/7/8/7, THRA 6/4/8/8 and NCOA1 6/7/8/7.

Gene	Primers	Cq ± SD	Efficiency	CV Cq	Repeatability expression	Product size
GAPDH	F: TCTCTGTTGTTGACCTGACCTG R: ATGGCTGTCACCATTGAAGTC	15.0±0.50	98.0%	0.59%		154bp
PGK1	F: CAAGCTCACCCTGGACAAGT R: GGACGGCTGCCTTGATTCTT	21.1±0.86	85.5%	0.48%		119bp
DIO2	F: CTGGAACAGCTTCCTCCTGG R: ATCCAGAAGACGGCACTTGG	23.6±0.83	92.9%	0.40%	0.91	157bp
NCOA1	F: CGACTCACGGAAGAGGAAGG R: GTGTCGATGTCCCCGATGTT	20.4±2.82	91.9%	0.38%	0.84	132bp
THRA	F: TGGTGAAGAGCAGCATGTCA R: AGCACCCATCGTACTTGCAG	21.4±0.82	96.0%	0.46%	0.82	189bp

Test: measurement	Age	Control	T3	T3T4	T4	F_{ddf}/χ^2 (df)	P
Tonic immobility: Recovery time (s)	12 d	92.0±25.5	84.6±21.1	68.4±13.1	123±24.7	$2.52_{36.2}$	0.073
	4.5 m	90.6±25.3	83.3±21.4	67.3±13.6	121±25.5		
Open arena: Movement duration (s)	3 d	181±31.6	204±28.1	177±19.3	167±21.3	$0.44_{46.6}$	0.72
	4.5 m	77.2±31.1	100±27.1	73.3±19.1	63.3±20.8		
Open arena: Escape attempts (number)	3 d	5.85±3.87	4.14±3.53	8.25±2.14	7.21±2.43	$0.35_{41.7}$	0.79
Emergence: Exit time (s)	6–9 d	34.8±14.7	42.7±16.4	74.7±20.9	121±36.1	$1.92_{39.6}$	0.14
	4.5 m	25.6±12.5	6.59±2.75	23.0±7.64	12.6±4.30		
Emergence: Likelihood of emergence (0-1)	6–9 d	0.80±0.11	0.70±0.12	0.62±0.10	0.58±0.10	2.71 (3)	0.44
	4.5 m	0.91±0.06	0.86±0.08	0.81±0.08	0.78±0.09		