Can increased prenatal exposure to thyroid hormones alter physiology

and behaviour in the long-term? Insights from an experimental study

in Japanese quails.

Kalle Aho[1,2]*, Antoine Stier[1,3,4], Tom Sarraude[1,6], Bin-Yan Hsu[1], Suvi Ruuskanen[1,5]*

[1] Department of Biology, University of Turku FI-20014 Turku, Finland

[2] School of Medicine, Faculty of Health Sciences, University of Eastern Finland, Kuopio,
Finland

[3] Univ Lyon, Université Claude Bernard Lyon 1, CNRS, ENTPE, UMR 5023 LEHNA, F-69622,
Villeurbanne, France

[4] Institut Pluridisciplinaire Hubert Curien, UMR7178, Université de Strasbourg, CNRS,
Strasbourg, France

[5] Department of Biological and Environmental Science, University of Jyväskylä, Finland

[6] Groningen Institute for Evolutionary Life Sciences, University of Groningen, the
Netherlands

*corresponding authors: email: ahokal9@gmail.com, suvi.ruuskanen@gmail.com

Abstract

Maternal thyroid hormones (triiodothyronine T3 and thyroxine T4) are important
regulators of embryonic development and gene expression. While maternal thyroid
disorders are known to cause developmental issues in humans, variation in maternal
thyroid hormones in healthy mothers have also been found to correlate with infant and
child phenotypes. This suggests a potential impact of maternal thyroid hormones on
offspring phenotype in an eco-evolutionary context. In chickens, prenatal thyroid
hormone treatment has been shown to influence embryonic gene expression and
postnatal treatment to influence imprinting and learning. However, the potential long-
term effects of maternal thyroid hormones on physiology and behaviour are unclear.
This study aims to investigate the long-term effects of maternal thyroid hormones on
behaviour, plasma thyroid hormone levels and brain gene expression using the Japanese
quail (*Coturnix japonica*) as a model. Egg hormone levels were elevated by injecting
unincubated eggs with either saline (control), T3, T4 or a mixture of T3 and T4. Social
motivation, boldness and fearfulness to predators were tested shortly after hatching
and as adults. Plasma thyroid hormone levels and pallial expression of thyroid hormone
receptor A, type 2 deiodinase, and nuclear receptor coactivator 1 were measured in
adulthood. We found no evidence that elevated thyroid hormone levels in eggs affected
behaviour, plasma hormone levels, or gene expression. This is the first study examining
the potential long-term effects of elevated maternal thyroid hormones within the
natural range. Although we found no evidence of long-term effects, other traits may still
be affected and remain to be studied.

Keywords: maternal hormones, maternal effects, endocrinology, phenotypic plasticity,
bird

Introduction

Introduction

Maternal effects refer to the effect of the maternal phenotype or maternal

environment on the phenotype of the offspring. The effect can be mediated by hormones

passed from the mother to the offspring, the maternal hormones. As the levels of maternal

hormones are influenced by the maternal environment, they can transmit information about

the maternal environment to the offspring. This allows the development of the offspring

phenotype to be controlled by environmental factors (Marshall and Uller, 2007; Mousseau

and Fox, 1998). Such hormone-mediated maternal effects can be observed in a wide range of

traits, including physiology and behaviour (e.g. Groothuis et al., 2019, 2005; Ruuskanen and

Hsu, 2018), and can have long-lasting effects into adulthood with potential fitness

consequences (e.g. metabolic rate, predator-avoidance behavior, survival, reproduction;

Ruuskanen et al., 2010, 2012: Groothuis et al., 2005; Schoech et al., 2011; von Engelhardt and

Groothuis, 2011).

Compared to the more extensively studied androgens and glucocorticoids (Podmokła

et al., 2018), the effects of maternally transferred thyroid hormones (THs) have been less

explored (Darras, 2019; Ruuskanen and Hsu, 2018). THs (triiodothyronine, T3, and thyroxine,

T4) play important roles in metabolism and development, where they regulate the expression

of numerous genes (Bernal, 2017; Darras, 2019; Moog et al., 2017). TH action is regulated by

several components of the thyroid system. TH receptors (THRA and THRB) and coactivators

control the transcription of TH-responsive genes by forming a complex with the active form

of the hormone, T3 (Brent, 2012). The intracellular availability of T3 is regulated by several

transmembrane TH transporters (e.g. MCT8, OATP1c1) that differentially transport either T4

or T3 (Groeneweg et al., 2020). Inside the cell, T3 availability can be further regulated by

deiodinase enzymes that either convert T4 to T3 (deiodinase 2, D2) or inactivates both T3 and

T4 (deiodinase 3, D3) (Gereben et al., 2008). The expression of these components (TH

receptors, transporters and deiodinases) begins as early as embryonic day 4 in chickens

(Flamant and Samarut, 1998) and embryonic day 1 in passerines (Ruuskanen et al. 2022),

allowing the embryo to regulate TH levels in an organ and cell-specific manner (Darras, 2019;

Darras et al., 2011; Flamant and Samarut, 1998; Ruuskanen et al., 2022; Van Herck et al.,

2012). However, the development of the thyroid gland occurs later in development, around

mid-incubation (Forhead and Fowden, 2014; McNichols and McNabb, 1988; Porazzi et al.,

2009). Thus, in almost all vertebrates studied so far, proper embryonic development depends

on maternally transferred THs (Brown et al., 2014; Darras, 2019; Moog et al., 2017; Patel et

al., 2011).

In mammals, maternal THs are transferred across the placenta, whereas in egg-laying

animals such as birds, maternal THs are deposited in the egg yolk. In birds, high natural

variation in egg TH levels has been observed, up to 5-fold in magnitude (Ruuskanen and Hsu,

2018). This variation may be due to environmental factors such as food availability or

temperature (Hsu et al., 2016; Ruuskanen et al., 2016c), and has been reported at both the

inter- and intra-individual levels (Hsu et al., 2016, 2019b; Ruuskanen et al., 2016a, 2016c). As

THs are important in development (Darras et al., 2009; Decuypere et al., 2005; McNabb, 2007;

Wilson and McNabb, 1997), and as increasing yolk TH levels has been found to increase

embryonic TH levels (Van Herck et al., 2012), high variation in egg TH levels could therefore

be a source of offspring phenotypic variation.

To date, most of what is known about phenotypic differences due to maternal THs

comes mainly from biomedical studies (Moog et al., 2017). However, studies using

pathological TH concentrations do not provide relevant information on the ecology or

evolution of maternal THs. Therefore, the effects of maternal THs within the natural range

need to be studied to understand whether variation has effects on traits affecting offspring

fitness. In addition, experiments manipulating maternal THs in mammals would inevitably

interfere with maternal physiology, making it difficult to attribute results solely to maternal

THs. Egg-laying animals, such as birds, have therefore become common models for studying

the functions of maternal hormones since maternal THs deposited in eggs can be manipulated

directly and independently. So far, elevated TH levels in the egg within the natural range have

been found to affect hatching success, offspring growth, chick levels of T3 and telomere

length in altricial bird species (Hsu et al., 2023a, 2019a, 2017; Ruuskanen et al., 2016b;

Sarraude et al., 2020; Stier et al., 2020). Nevertheless, some of the reported effects appear

to be transient during the early post-hatching period (Hsu et al., 2019a). Whether the effects

of maternal THs can be long-lasting and potentially influence offspring fitness remains

unclear. In terms of precocial bird species, previous studies studying maternal THs were

scarce and typically applied supra-physiological doses (e.g., Wilson and McNabb, 1997; van

Herck et al., 2012).

To investigate the potential long-term effects of variation in maternal THs, we

experimentally elevated egg TH levels within the natural range in Japanese quail (*Coturnix*

japonica). With these experimental quails, we have previously reported a higher hatching

success caused by the elevated yolk TH levels, but no major effects on morphology or

physiology (Sarraude et al., 2020). Now we further studied changes in chick and adult

behaviour and adult plasma TH levels. In addition, we studied brain gene expression at later

age. While increasing maternal TH levels beyond normal physiological range (maternal

hyperthyroidism) has been shown to affect behaviour, for example both decreasing and

increasing anxiety (Laureano-Melo et al., 2019; Zhang et al., 2010, 2008), it is not known

whether increasing maternal THs within the natural range affects behaviour. Also, as

postnatal THs have been shown to influence filial imprinting and learning behaviour

(Yamaguchi et al., 2012), it is interesting to study whether maternal THs, acting earlier in

development, may also have behavioural effects either in chicks or in adulthood. Potential

behavioural effects were investigated in both chicks and adults, allowing for a broader

timescale, by examining boldness, social motivation and fearfulness to predators. Due to the

lack of studies with natural levels of elevated maternal THs and inconsistent results from

studies on maternal hyperthyroidism and offspring anxiety (Laureano-Melo et al., 2019;

Zhang et al., 2010, 2008), the direction of potential behavioural changes is difficult to predict.

In addition, plasma TH levels were measured in adults, and changes in these would indicate

long lasting effects on TH function. While plasma TH levels have been reported to be affected

by elevated egg TH levels (within the natural range) in a sex-dependent manner in nestling

rock pigeons (Hsu et al., 2017), some more recent studies in wild passerines found no clear

effects of higher natural levels of egg THs on nestling plasma THs (Hsu et al., 2023b, 2020). In

terms of gene expression, previous studies have examined the effect of the natural variation

in maternal THs on brain gene expression only in bird embryos or young chicks (Van Herck et

al., 2015, 2012; Yamaguchi et al., 2012), while the potential long-term effects of prenatal TH

exposure in adulthood remain unknown. Here we studied several genes involved in TH

function: THRA (thyroid hormone receptor A), NCOA1 (nuclear receptor coactivator 1) and

DIO2 (type 2 deiodinase) in the pallium of adults. Expression of these genes has been shown

to be influenced by THs in other species (Misiti et al., 1998; Walpita et al., 2007).

Materials and methods

1.1 Materials

This study was conducted in accordance with the Finnish legislation and approved by

the Finnish Animal Ethics Committee (Eläinkoelautakunta; ESAVI / 1018 / 04.10.07 / 2016).

Japanese quail (*Coturnix japonica*), a commonly used laboratory model with a reference

genome, was used. Twenty-four breeding pairs were housed in cages with food and water

available ad libitum. The photoperiod was long (16L:8D) and the temperature stable at 20° C.

For seven days, eggs were collected from the cages in the morning and stored (15° C, 50%

relative humidity) until treatment.

1.2 Treatment of eggs, rearing of chicks and collection of samples

We manipulated egg TH levels by injecting hormones directly into the eggs prior to

embryonic development at a dose that mimicked increased maternal transfer (see details in

Sarraude et al., 2020). Eggs from each breeding pair were randomly assigned to four

treatment groups: T3, T4, T3 + T4 and control (0.9% saline). We used saline control following

previous studies on injecting THs into eggs (van Herck et al. 2015), and because THs are not

very well soluble in oil. Eggs laid by the same pair were always evenly distributed among all

treatment groups. The injected hormone solutions were prepared from crystallised powder

(T3: CAS number 6893-02-3, Sigma-Aldrich, St. Louis, MO, USA and T4: CAS number 51-48-9,

Sigma-Aldrich, St. Louis, MO, USA). The amount of hormone injected (T4: 8.9 ng/egg,

equivalent to 1.79 pg/mg yolk; T3: 4.7 ng/egg, equivalent to 1.24 pg/mg yolk) was based on

previous measurements in the parental generation (mean ± SD: T4, 15.3 ± 4.43 ng / egg; T3,

7.58 ± 2.32 ng / egg) and the aim was to increase hormone contents in the yolk by two

standard deviations (as recommended by Podmokla et al., 2018). We were unsure how much

of T4 is converted to T3 and therefore used 2SD of both hormones for the T3T4 group. The

injection protocol has been described in detail in Sarraude et al. (2020). Briefly, each egg was

injected with 50 µl of either the respective hormone solution (T3, T4 or T3+T4, in 0.9% saline)

or 0.9% saline (control, CO), delivered by using 27G insulin syringes (BD Microlance™). After

sealing the hole with sterile plaster (OPSITE Flexigrid, Smith & Nephew), the eggs were

incubated artificially (37.8 °C and 55% relative humidity). A total of 158 eggs were treated (39

T3, 39 T4, 40 T3+T4 and 40 CO), of which 66 chicks hatched (15 T3, 20 T4, 21 T3 + T4 and 10

CO), with sex distributions of chicks surviving to adulthood (females / males): 7/4 for T3, 10/8

for T4, 9/12 for T3 + T4 and 3/4 for CO. Sex was determined at maturity by examining the

presence of the cloacal gland. The overall hatching rate (~40%) was consistent with previous

studies in domestic quail (Okuliarová et al., 2007).

The rearing conditions of the chicks have been described in detail in a previous

publication (Sarraude et al., 2020). In short, chicks were reared in groups (4-16 individuals,

depending on age) with treatments and sexes mixed, until sexual maturity (ca. 6–8 weeks

after hatching in Japanese quails), after which the birds were separated by sex to smaller

groups (4-5 individuals). Photoperiod was 16L:8D for the first 7 months and 8L:16 thereafter.

At approximately 4.5 months of age, we collected blood samples (200 µl) from the brachial

vein, separated plasma, and stored it at -80° C. As Japanese quails have reached adulthood

at 4.5 months, this time point would have already shown if there are long-lasting,

organizational effects of prenatal THs effects into adulthood. Birds were euthanised by

decapitation at 10 months of age. After decapitation, brain tissue was collected for the gene

expression analysis from the pallium (average 44.0 mg, std 2.8 mg, taken at the surface of the

region) in PBS buffer, snap frozen in liquid nitrogen and stored at -80° C. While differences in

photoperiod limit comparisons between gene expression results and other results, our aim

was to study whether there are persistent effects of prenatal THs independent of any

environmental factors that individuals may face later in life, for which our experimental

design is sufficient.

1.3 Behavioural experiments

To assess the potential effects of maternal THs on behaviour, we conducted a series of

behavioural tests (open arena, emergence and tonic immobility tests) at different life stages.

We followed previously published protocols that have been used extensively with many

different model organisms. The tests were conducted in separate rooms where the birds had

no visual or auditory contact with other individuals. The exception was the open arena test,

where the subject was able to hear the vocalisations of another bird being tested at the same

time in another room. All experiments were recorded, and the videos were analysed blindly

of the treatment. All videos of the same test at both ages (chick/adult) were analysed by one

person, except for the open arena test, where one person analysed chick videos and another

person analysed adult videos. This may have confounded the age difference in behavior

scores with the observer effect. Since the videos from different treatments at each age

were analysed by the same person, the fact that chick and adult videos were analyzed by

different people should not confound the effect of prenatal TH exposure.

The open arena test measured social motivation and boldness and it followed a setup

developed by Forkman et al. (2007). The test was performed on both sexes at three days and

approximately 4.5 months of age. In the test, the bird was placed in a circular illuminated

arena and the movement duration and the number of escape attempts (attempt to jump/fly

over the wall) were measured from videos for 10 minutes (from the time the researcher left

the room). Only 10 of the 56 adults (1 CO, 2 T3, 2 T4 and 5 T3T4) attempted to escape, so

adult escape attempts were not analysed. Subjects who moved less and made fewer escape

attempts were interpreted as less bold and less socially motivated.

The emergence test measured social motivation and boldness. The test was

performed on both sexes, twice on chicks (6-9d old) and once on adults (4.5 months old). The

test followed a setup developed by Forkman et al. (2007), in which a bird was placed in a dark

chamber connected to another, bright chamber. One minute after the subject was placed in

the dark chamber, a door connecting the chambers was removed. After removing the door,

the researcher left the observation room and the time it took for the subject to completely

exit the dark chamber was measured. Subjects who left the dark chamber more quickly were

interpreted as bolder and more socially motivated. The test lasted five minutes. Whether the

subject remained in the dark chamber (0) or left it (1) during the test was also recorded.

The tonic immobility test (Mills and Faure, 1991) measured fearfulness to predators

(Forkman et al 2007). The test was performed on both sexes at 12 days and approximately

4.5 months of age. To induce immobility, the bird was placed on its back in the centre of a V-

shaped test apparatus and restrained by hand for 15 seconds, after which the hand was

removed, and the time taken for the bird to resume a standing posture was measured. If the

subject resumed the standing posture in less than 10 seconds, this was recorded as a failed

induction and the induction procedure was repeated. Four adult subjects who failed five times

were excluded. If the subject remained stationary for more than 10 seconds, then the time to

recovery was measured. The longer a subject took to recover, the more fearful it was

interpreted. The test lasted 10 minutes. Five subjects (1 chick and 4 adults) did not recover

for 10 minutes and were excluded from the analysis (approximately 4% of all tests).

 1.4 Plasma thyroid hormone levels

To test whether elevated maternal THs affect plasma TH levels later in life (4.5 months

of age), we extracted THs from plasma samples and measured them using a previously

validated LC-MS / MS protocol (Ruuskanen et al., 2018, where CV <10%). Due to practical

limitations (*i.e.* volume of plasma available and haemolysis in some samples), the sample size

was reduced to 48 individuals (T3: 10, T4: 16, T3T4: 15, and 7 in the control). Hormone

concentrations are expressed in pmol/ml plasma. The repeatability of the TH measurements,

was validated in Ruuskanen et al. (2018) with CV% less than 10% (3.5-9.7%, depending on the

hormone and sample type)

1.5 Gene expression

To test for potential long-term changes in the expression of TH related genes (*DIO2*,

DIO3, *THRA*, and *NCOA1*), we measured gene expression from pallium samples (10 months

old). RNA was extracted from the homogenised pallium samples using the NucleoSpin RNA

Plus kit (MACHEREY-NAGEL). Since the sample size of the control group was the main factor

limiting statistical power, we decided to analyse all control individuals and only an equivalent

number of individuals (randomly selected) in the three treatment groups (N = 31, 7 in control

and 8 per other treatment groups). Extraction was done from 50 µl of the homogenised brain

sample according to the manufacturer's instructions, except that the gDNA removal column

was centrifuged longer (1 min, 11000 g) and the RNA was eluted once with 40 µl of Rnase-

free water. RNA concentration and purity were quantified by optical density, and all samples

not meeting our quality criteria (i.e. RNA concentration > 30 ng/µl and 260/280 > 1.80) were

excluded from further analysis. RNA integrity was checked using an E-Gel 2% electrophoresis

system (Invitrogen), and the ribosomal RNA 18S *vs*. 28S band intensity and was deemed

satisfactory (the visual inspection revealed the expected two bands of rRNA with 28S band

being > 2 times more intense than 18S band). Samples were stored at -80° C for 2 weeks prior

to cDNA synthesis. cDNA synthesis was performed using the SensiFAST ™ cDNA Synthesis Kit

(Bioline) according to the manufacturer's instructions. The reaction was performed using 500

ng of RNA as the starting material. A control (noRT) was prepared without reverse

transcriptase enzyme. The cDNA was diluted with water to a concentration of 1.2 ng/μl and

stored at -20° C. The 1.2ng/uL was calculated from RNA pre-RT, and slight variations in cDNA

starting quantity (due to differences in RT efficiency) are controlled for by the use of the

reference genes when calculating relative expression.

GAPDH and *PGK* were used as control genes, using primers previously designed and

validated in Japanese quail (Vitorino Carvalho et al., 2019). Stability of expression between

experimental groups was further verified using geNorm software (geNorm M < 0.5, geNorm

V < 0.15), and thus the geometric mean of these two genes was used as our normalization

factor. Primers for the target genes were designed using the Primer-BLAST tool (NCBI). The

primer pairs (Table 1) were designed to have at least one intron between them. The suitability

of the primers in qPCR was tested prior to the actual analysis using different primer

concentrations and different amounts of cDNA. The tests confirmed the sufficient

amplification efficiency of the primers (between 90 and 110% based on the standard curve)

and the formation of only one product of the expected size based on both the presence of a

single narrow peak in the melting curve, and the presence of a single band of the expected

size on the agarose gel. *DIO3* had extremely poor amplification, meaning that we could not

accurately quantify it, and was excluded from the analysis.

Table 1: qPCR primer set information and performance for gene expression assays. Cq refers to
qPCR quantification cycle, efficiency was evaluated using the LinReg method based on individual well
efficiency. Technical precision estimates are given for Cq as coefficient of variation CV (%) and for
expression (ratio of target to reference genes) as technical repeatability I.

Gene	Primers	Cq ± SD	Efficiency	CV Cq	Repeatability expression	Product size
GAPDH	F: TCTCTGTTGTTGACCTGACCTG R: ATGGCTGTCACCATTGAAGTC	15.0±0.50	98.0%	0.59%		154bp
PGK1	F: CAAGCTCACCCTGGACAAGT R: GGACGGCTGCCTTGATTCTT	21.1±0.86	85.5%	0.48%		119bp
DIO2	F: CTGGAACAGCTTCCTCCTGG R: ATCCAGAAGACGGCACTTGG	23.6±0.83	92.9%	0.40%	0.91	157bp
NCOA1	F: CGACTCACGGAAGAGGAAGG R: GTGTCGATGTCCCCGATGTT	20.4±2.82	91.9%	0.38%	0.84	132bp
THRA	F: TGGTGAAGAGCAGCATGTCA R: AGCACCCATCGTACTTGCAG	21.4±0.82	96.0%	0.46%	0.82	189bp

qPCR assays were performed on a Mic qPCR (Bio Molecular Systems) in a final volume of 12

µL containing 6 ng of cDNA, 800 nM of forward and reverse primers and SensiFAST SYBR Lo-

ROX (Bioline). We ran gene-specific qPCR plates and each sample was analysed in duplicate.

Two plates were run for each gene and experimental groups and sex were always balanced

within each plate. We used a cDNA reference sample (*i.e.* ratio = 1), which was a pool of 10

different individuals on each plate. We used a two-step cycling with the following conditions:

2 min at 95°C; then 45 cycles of 5s at 95°C followed by 20s at 60°C (fluorescence reading) for

all reactions, as recommended by manufacturer. The expression of each gene was calculated

as $(1+Ef_{Target})^{\Delta Cq(Target)}$/ geometric mean $[(1+Ef_{GAPDH})^{\Delta CqGAPDH}; (1+Ef_{PGK1})^{\Delta CqPGK1}]$, where Ef is the

amplification's efficiency and ΔCq is the difference between the Cq-values of the reference

sample and the sample of interest. Amplification in the no-RT controls never occurred until at

least 8 cycles after the lower Cq sample, and thus contamination by genomic DNA could not

interfere with our results. Technical precision and repeatability were generally good (Table

1), but a few samples with a CV% of duplicates (based on the final ratio) greater than 15%

were excluded from the analysis (final sample size: N_{DIO2} = 27, N_{NCOA1} = 28, N_{THRa} = 26)

1.6 Statistical analysis

All statistical analyses were performed with R software (version 3.6.2), using the packages

lme4 version 1.1-23 (Bates et al., 2015) and emmeans version 1.6.0 (Lenth, 2021). Normality

and homoscedasticity of the residual distributions were checked visually (quantile-quantile

and residuals against fitted values plots). Normality was additionally confirmed with Shapiro-

Wilk test of normality. For the behavioural analysis, recovery time (tonic immobility) and exit

time (emergence test), were transformed (natural logarithm) to ensure that the residuals

were normally distributed. Linear mixed models (LMM) were used to analyse time to exit in

the emergence test, movement duration and number of escape attempts in the open arena

test, recovery time in the tonic immobility test, plasma TH concentrations and qPCR results.

Generalised linear mixed model (GLMM) with a binomial error distribution was used to

analyse the proportion of individuals leaving the dark chamber in the emergence test. Fixed

effects were treatment, sex, and test number/age (in repeated tests). Interactions between

sex and treatment, and between treatment and age (in repeated tests), were also analysed.

Random effects were individual ID (in repeated tests) and parental ID (to account for

relatedness between siblings in all experiments). F- and p-values for LMMs were obtained

using the Kenward-Roger approximation on the denominator degree of freedom,

implemented in the package *lmerTest* (Kuznetsova et al., 2017). For binomial regression models, main effects and interactions were tested by comparing the model with and without the factor. Non-significant (p > 0.05) interactions were excluded from the final models. Post-hoc tests were performed using Tukey tests (emmeans package, Lenth, 2021) where appropriate.

Results

Effects of prenatal thyroid hormone supplementation on behaviour

In general, no statistically significant differences in chick or adult behavioural traits were observed between the egg TH manipulation groups (T3, T4, T3 + T4, control). Elevated THs had no significant effect on social motivation or boldness (Table 2, Fig. 1-2), nor on fearfulness to predators (Table 2, Fig. 3). A statistically significant interaction between treatment and age was found in the emergence test (F = 3.58, ddf = 67.2, p = 0.018, Fig. 2): In all treatments except the control group (Tukey t=0.59, p=0.56), chick exit times were significantly longer than those of adults (all t > 3.21, p < 0.002) while no differences between treatments were found within each age-category (all t < 2.48, all p> 0.07). There was also a statistically significant interaction between treatment and sex in the tonic immobility test (F = 2.92, ddf = 37.5, p = 0.047, Fig. 3): In males, the recovery time was longer in the T4-group (marginal mean ± SE: 177.1 ± 47.8 s) than in the T3-group (marginal mean ± SE: 56.1 ± 20.8 s, Tukey t = 2.71, p = 0.047), whereas no differences between treatments were observed in females (all t < 2.29, all p >0.12). In the T4-group, the recovery time of males (marginal mean ± SE: 177.1 ± 47.8 s) was longer than that of females (marginal mean ± SE: 85.2 ± 19.9 s) (t=2.16, p= 0.037), while no sex differences were observed within the other treatments (all t<1.80, all p>0.079).

357 **Table 2.** Marginal means and standard errors of behavioural tests. Results for chicks (age: 3d, 6-
358 9d, 12d) and adults (age: 4.5 months) are shown separately for each variable. F- and p-values for
359 overall treatment, independent of age, from linear models (ANOVA, Kenward-Roger
360 approximation). Test statistics (χ^2) and p-values for binomial regressions (model with treatment
361 compared to model without treatment) were calculated with ANOVA.

Test: measurement	Age	Control	T3	T3T4	T4	F_{ddf}/χ^2 (df)	P
Tonic immobility:	12 d	92.0±25.5	84.6±21.1	68.4±13.1	123±24.7	$2.52_{36.2}$	0.073
Recovery time (s)	4.5 m	90.6±25.3	83.3±21.4	67.3±13.6	121±25.5		
Open arena:	3 d	77.2±31.1	100±27.1	73.3±19.1	63.3±20.8	$0.44_{46.6}$	0.72
Movement duration (s)	4.5 m	181±31.6	204±28.1	177±19.3	167±21.3		
Open arena:	3 d	5.85±3.87	4.14±3.53	8.25±2.14	7.21±2.43	$0.35_{41.7}$	0.79
Escape attempts (number)							
Emergence:	6–9 d	34.8±14.7	42.7±16.4	74.7±20.9	121±36.1	$1.92_{39.6}$	0.14
Exit time (s)	4.5 m	25.6±12.5	6.59±2.75	23.0±7.64	12.6±4.30		
Emergence:	6–9 d	0.80±0.11	0.70±0.12	0.62±0.10	0.58±0.10	2.71 (3)	0.44
Likelihood of emergence (0-1)	4.5 m	0.91±0.06	0.86±0.08	0.81±0.08	0.78±0.09		

Figure 1. Movement duration for chicks (3 d) and adults (4.5 months) in the open arena test for

different prenatal thyroid hormone treatment groups (CO = control, 50 µl 0.9% saline; T3: 4.7

ng/egg in 50 µl 0.9% saline, equivalent to 1.24 pg/mg yolk; T4: 8.9 ng/egg in 50 µl 0.9% saline,

equivalent to 1.79 pg/mg yolk). The lines represent the medians of the treatments, and the bars

represent the interquartile range of the original observations. Sample size (chicks/adults): CO 7/7,

T3 11/11, T3T4 20/21 and T4 16/17.

Figure 2. Time to exit the dark chamber for chicks (6-9 d) and adults (4.5 months) in the emergence test for different prenatal thyroid hormone treatment groups (CO = control, 50 μl 0.9% saline; T3: 4.7 ng/egg in 50 μl 0.9% saline, equivalent to 1.24 pg/mg yolk; T4: 8.9 ng/egg in 50 μl 0.9% saline, equivalent to 1.79 pg/mg yolk). The test was performed twice for chicks and once for adults. Results for both chick studies are shown. The lines represent the medians of the treatments, and the bars represent the interquartile range of the original observations. Sample size (chicks/adults): CO 11/6, T3 15/9, T3T4 24/13 and T4 22/13.

Figure 3. Time to resume standing position for females and males, chicks (12 d) and adults (4.5

months) combined, in the tonic immobility test for different prenatal thyroid hormone treatment

groups (CO = control, 50 µl 0.9% saline; T3: 4.7 ng/egg in 50 µl 0.9% saline, equivalent to 1.24

pg/mg yolk; T4: 8.9 ng/egg in 50 µl 0.9% saline, equivalent to 1.79 pg/mg yolk). The lines

represent the medians of the treatments, and the bars represent the interquartile range of the

original observations. Sample size (female/male: CO 6/8, T3 13/7, T3T4 14/23 and T4 18/14.

Effects of prenatal thyroid hormone supplementation on plasma thyroid

hormone levels

Increasing TH levels in eggs did not significantly affect plasma TH levels in 4.5-

month-old birds (T3: F = 0.13, ddf = 37.7, p = 0.94, T4: F = 2.1, ddf = 41.5, p = 0.12, Fig.

4). Treatment also had no sex-dependent effect on plasma T3 or T4 (treatment-sex

interactions, p > 0.36).

Figure 4. Plasma T3 and T4 concentrations in 4.5-month-old birds between different prenatal thyroid hormone treatment groups (CO = control, 50 µl 0.9% saline; T3: 4.7 ng/egg in 50 µl 0.9% saline, equivalent to 1.24 pg/mg yolk; T4: 8.9 ng/egg in 50 µl 0.9% saline, equivalent to 1.79 pg/mg yolk). The lines represent the medians of the treatments, and the bars represent the interquartile range of the original observations. Sample size (T3/T4 concentrations): CO 6/7, T3 10/10, T3T4 14/15 and T4 14/16.

Effects of prenatal thyroid hormone supplementation on pallial gene expression

Increasing TH levels in eggs did not affect pallial gene expression of TH-related genes in 10-month-old birds ($DIO2$: F = 2.49, ddf = 17.3, p = 0.09, $NCOA1$: F = 0.092, ddf = 21.1, p = 0.96, $THRA$: F = 0.47, ddf = 18.9, p = 0.71, Fig. 5). Treatment also did not have sex-dependent effect on gene expression (treatment-sex interactions, p > 0.26). Females had higher expression of $NCOA1$ (F=7.68, p=0.012) and $THRA$ (F=5.17, p=0.034) than males.

Figure 5. Relative expressions of *DIO2*, *THRA* and *NCOA1* in the pallium of 10-month-old birds
between different prenatal thyroid hormone treatments (CO = control, 50 µl 0.9% saline; T3: 4.7
ng/egg in 50 µl 0.9% saline, equivalent to 1.24 pg/mg yolk; T4: 8.9 ng/egg in 50 µl 0.9% saline,
equivalent to 1.79 pg/mg yolk). The lines represent the medians of the treatments, and the bars
represent the interquartile range of the original observations. Sample sizes (CO/T3/T3T4/T4):
DIO2 5/7/8/7, THRA 6/4/8/8 and NCOA1 6/7/8/7.

Discussion

In this study, we tested for long-term changes in behaviour, plasma TH levels and

pallial gene expression following increased exposure to maternal THs. In terms of

behaviour, we observed no significant changes in boldness, social motivation or

fearfulness to predators in chicks (3-12 d) or adults (4.5 months) when the egg TH level

was increased. To our knowledge, there are no similar previous studies, and it is

therefore difficult to assess whether there may have been changes in other behavioural

traits. The detection of changes was also partly limited by the small sample size, which

meant that we may not have had enough statistical power to detect subtle changes in

behaviour. Nevertheless, we found some age-dependent and sex-dependent effects of

TH treatment on behaviour. The age-dependent effect in the emergence test suggests

that elevated maternal TH leads to a decrease in boldness and social motivation with

age (from chick to adult), whereas such a decrease is not observed in controls. However,

the lack of differences between control and treatment within a given age suggests that

such an effect (if not associated with type I statistical error) is rather mild. The sex-

dependent effect in the tonic immobility test also seems relatively unconvincing because

no sex differences were observed within treatments, nor did T3T4 females or T4 males

differ from the same-sex controls. Given the important role of THs in brain development,

we expected that elevated maternal THs would have led to behavioural changes. Yet,

our data suggest that an experimental increase in maternal THs within the natural range

has no clear and large effect on the behavioural traits measured. However, in contrast

to maternal THs, post-hatch TH treatment twenty hours after hatching has been shown

to affect imprinting and learning in four-day-old chicks (Yamaguchi et al., 2012). The

discrepancy between our results and Yamaguchi et al. (2012) could suggest that the

sensitive period during which THs can influence behaviour might occur at a later timing

of development, but this requires further investigation.

We found that elevated maternal THs did not significantly affect plasma TH levels

in adulthood (4.5 months of age). This was the first study to test for long-term changes,

whereas previously changes in response to elevated prenatal THs have been observed

at younger ages: Experimental elevation of egg THs was found to increase male and

decrease female plasma TH levels in rock pigeons at 14 days of age (Hsu et al., 2017),

but such effects were not observed in two other wild passerine species (Hsu et al.,

2023b, 2020). In contrast to these studies in altricial species, not only was this study

conducted on a precocial species, but our measurements were also made at a different

life stage (adulthood). Overall, this suggests that an increase in maternal THs may only

lead to transient changes in offspring plasma TH levels and that the effects may not be

universal across species.

We also found no significant effect of elevated maternal THs on the expression

of *THRA*, *DIO2* or *NCOA1* in the pallium of adult (10 months old) individuals. Since we

only collected samples from 10-month-old birds, we cannot rule out possible changes at

younger ages. Changes in gene expression have been observed in other TH-related

genes, but, at much shorter time spans: In chickens, injection of THs into the yolk

decreased the expression of the TH-transporter *OATP1C1* and increased the expression

of the TH-transporter *MCT8* in 4-day-old embryos (Van Herck et al., 2012). The

expressions of our target genes, *THRA*, *DIO2* and *NCOA1,* have also been shown to be at

least transiently responsive to THs in other species: increase in *THRA* and decrease in

DIO2 mRNA levels in zebrafish embryos (Walpita et al., 2007), and increase in *NCOA1*

mRNA levels in the anterior pituitary of adult rats (Misiti et al., 1998). It is therefore

possible that the gene regulation by maternal THs is restricted to early development

when the embryo is not yet able to produce THs.

We have previously shown in these same experimental birds, that an increase in

prenatal THs influences hatching success (Sarraude et al., 2020). Therefore, the dose

used in this study can have some effects during embryo development, which suggests

that the lack of large changes in long-term behaviour, plasma TH levels and brain gene

expression may be linked to an attenuation of prenatal TH effects by embryonic

regulation. It is indeed possible that embryos could buffer the effect of maternal THs by

altering the expression of deiodinases which are expressed already at very early stages

(Ruuskanen et al., 2022; Too et al., 2017). Metabolisation of THs to inactive forms could

prevent the elevated egg THs from exerting any effects. However, given the effects

previously observed in these same birds on hatching success (Sarraude et al., 2020), the

lack of changes in this study may also be explained by the stage of ontogeny, prenatal

THs not leading to long-lasting effects. Nevertheless, the lack of prenatal measurements

means that this cannot be fully confirmed, and it remains possible that elevated

maternal THs have no effect on the measured traits in Japanese quails.

In conclusion, we found no evidence that elevated maternal THs cause

substantial long-term changes in behaviour, plasma TH levels or pallial gene expression.

This is consistent with our previous findings in the same experimental birds, where

elevation of prenatal THs (T3, T4, or in combination) did not lead to long-term effects on

oxidative stress, moult or reproduction (Sarraude et al., 2020). While behavioural

changes have not been previously studied, plasma TH concentrations and gene

expression have been shown to respond to elevated maternal THs in the short- term

(Hsu et al., 2017; Van Herck et al., 2012). However, based on this study, these changes

would appear to be limited to early development or shortly after hatching with the

offspring's own TH production later replacing the effects of elevated maternal THs.

Maternal THs could theoretically influence the offspring's own TH production at a later

age by altering the function of the HPT-axis, but no evidence has yet been found to

support this hypothesis. The long-term effects of variation in maternal THs remain

unclear, but this study has provided new information on some previously untested

traits. Understanding long-term effects in addition to short-term effects will help to

better understand the evolutionary and ecological importance of maternal THs.

Acknowledgements

We thank personnel at the animal care facilities, Marjo Myllyrinne, Simo Laine for their

help.

Funding

The study was funded by the Academy of Finland (project #286278) to SR, the Finnish

National Agency for Education (grant no. TM-15-9960 to TS). AS was supported by a

Turku Collegium for Science and Medicine fellowship and an European Commission Ma-

rie Sklodowska-Curie Fellowship #894963.

Data availability statement

Data will be available on Figshare (DOI: 10.6084/m9.figshare.22266211) at publication.

Declaration of interests

The authors declare that they have no known competing financial interests or personal

relationships that could have appeared to influence the work reported in this paper.

Author contributions

S.R., T.S., B-Y.H, A.S and K.A. designed the study. S.R. obtained funding. T.S., B-Y.H and

S.R conducted the experiment. K.A. and T.S. collected the behavioural data from videos.

K.A., A.S and S.R. collected the physiological data and conducted laboratory work. K.A.

analysed all data with support from all co-authors, and wrote the first draft of the

manuscript. A.S, B-Y.H, T.S. and S.R commented on the manuscript.

References

Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects Models Using lme4. J Stat Softw 67. https://doi.org/10.18637/jss.v067.i01

Bernal, J., 2017. Thyroid hormone regulated genes in cerebral cortex development. Journal of Endocrinology 232, R83–R97. https://doi.org/10.1530/JOE-16-0424

Brent, G.A., 2012. Mechanisms of thyroid hormone action. Journal of Clinical Investigation 122, 3035–3043. https://doi.org/10.1172/JCI60047

Brown, C.L., Urbinati, E.C., Zhang, W., Brown, S.B., McComb-Kobza, M., 2014. Maternal Thyroid and Glucocorticoid Hormone Interactions in Larval Fish Development, and Their Applications in Aquaculture. Reviews in Fisheries Science & Aquaculture 22, 207–220. https://doi.org/10.1080/23308249.2014.918086

Darras, V.M., 2019. The Role of Maternal Thyroid Hormones in Avian Embryonic Development. Front Endocrinol (Lausanne) 10. https://doi.org/10.3389/fendo.2019.00066

Darras, V.M., Van Herck, S.L.J., Geysens, S., Reyns, G.E., 2009. Involvement of thyroid hormones in chicken embryonic brain development. Gen Comp Endocrinol 163, 58–62. https://doi.org/10.1016/j.ygcen.2008.11.014

Darras, V.M., Van Herck, S.L.J., Heijlen, M., De Groef, B., 2011. Thyroid Hormone Receptors in Two Model Species for Vertebrate Embryonic Development: Chicken and Zebrafish. J Thyroid Res 2011, 1–8. https://doi.org/10.4061/2011/402320

Decuypere, E., Van As, P., Van der Geyten, S., Darras, V.M., 2005. Thyroid hormone availability and activity in avian species: A review. Domest Anim Endocrinol 29, 63–77. https://doi.org/10.1016/j.domaniend.2005.02.028

Flamant, F., Samarut, J., 1998. Involvement of Thyroid Hormone and Its α Receptor in Avian Neurulation. Dev Biol 197, 1–11. https://doi.org/10.1006/dbio.1998.8872

Forhead, A.J., Fowden, A.L., 2014. Thyroid hormones in fetal growth and
prepartum maturation. Journal of Endocrinology 221, R87–R103.
https://doi.org/10.1530/JOE-14-0025

Forkman, B., Boissy, A., Meunier-Salaün, M.-C., Canali, E., Jones, R.B., 2007. A
critical review of fear tests used on cattle, pigs, sheep, poultry and horses.
Physiol   Behav   92,   340–374.
https://doi.org/10.1016/j.physbeh.2007.03.016

Gereben, B., Zeöld, A., Dentice, M., Salvatore, D., Bianco, A.C., 2008. Activation
and inactivation of thyroid hormone by deiodinases: Local action with
general consequences. Cellular and Molecular Life Sciences 65, 570–590.
https://doi.org/10.1007/s00018-007-7396-0

Groeneweg, S., van Geest, F.S., Peeters, R.P., Heuer, H., Visser, W.E., 2020.
Thyroid   Hormone   Transporters.   Endocr   Rev   41,   146–201.
https://doi.org/10.1210/endrev/bnz008

Groothuis, T.G.G., Hsu, B.-Y., Kumar, N., Tschirren, B., 2019. Revisiting
mechanisms and functions of prenatal hormone-mediated maternal
effects using avian species as a model. Philosophical Transactions of the
Royal   Society   B: Biological   Sciences   374,   20180115.
https://doi.org/10.1098/rstb.2018.0115

Groothuis, T.G.G., Müller, W., von Engelhardt, N., Carere, C., Eising, C., 2005.
Maternal hormones as a tool to adjust offspring phenotype in avian
species.   Neurosci   Biobehav   Rev   29,   329–352.
https://doi.org/10.1016/j.neubiorev.2004.12.002

Hsu, B., Dijkstra, C., Darras, V.M., Vries, B., Groothuis, T.G.G., 2016. Maternal
adjustment or constraint: differential effects of food availability on
maternal deposition of macro-nutrients, steroids and thyroid hormones
in   rock   pigeon   eggs.   Ecol   Evol   6,   397–411.
https://doi.org/10.1002/ece3.1845

Hsu, B.-Y., Cossin-Sevrin, N., Stier, A., Ruuskanen, S., 2023a. Prenatal thyroid
hormones accelerate postnatal growth and telomere shortening in wild
great   tits.   Journal   of   Experimental   Biology.
https://doi.org/10.1242/jeb.243875

Hsu, B.-Y., Cossin-Sevrin, N., Stier, A., Ruuskanen, S., 2023b. Prenatal thyroid
hormones accelerate postnatal growth and telomere shortening in wild
great tits. Journal of Experimental Biology.
https://doi.org/10.1242/jeb.243875

Hsu, B.-Y., Dijkstra, C., Darras, V.M., de Vries, B., Groothuis, T.G.G., 2017.
Maternal thyroid hormones enhance hatching success but decrease
nestling body mass in the rock pigeon (Columba livia). Gen Comp
Endocrinol 240, 174–181. https://doi.org/10.1016/j.ygcen.2016.10.011

Hsu, B.-Y., Doligez, B., Gustafsson, L., Ruuskanen, S., 2019a. Transient growth-
enhancing effects of elevated maternal thyroid hormones at no apparent
oxidative cost during early postnatal period. J Avian Biol 50.
https://doi.org/10.1111/jav.01919

Hsu, B.-Y., Sarraude, T., Cossin-Sevrin, N., Crombecque, M., Stier, A.,
Ruuskanen, S., 2020. Testing for context-dependent effects of prenatal
thyroid hormones on offspring survival and physiology: an experimental
temperature manipulation. Sci Rep 10, 14563.
https://doi.org/10.1038/s41598-020-71511-y

Hsu, B.-Y., Verhagen, I., Gienapp, P., Darras, V.M., Visser, M.E., Ruuskanen, S.,
2019b. Between- and Within-Individual Variation of Maternal Thyroid
Hormone Deposition in Wild Great Tits (Parus major). Am Nat 194, E96–
E108. https://doi.org/10.1086/704738

Kuznetsova, A., Brockhoff, P.B., Christensen, R.H.B., 2017. lmerTest Package:
Tests in Linear Mixed Effects Models. J Stat Softw 82.
https://doi.org/10.18637/jss.v082.i13

Laureano-Melo, R., Souza, J.S. de, da Conceição, R.R., Albuquerque, J.M.L.,
Rodrigues, N.C., Marinho, B.G., Olivares, E.L., Giannocco, G., Côrtes, W.
da S., 2019. Prenatal thyroxine treatment promotes anxiolysis in male
Swiss mice offspring. Horm Behav 108, 10–19.
https://doi.org/10.1016/j.yhbeh.2018.12.008

Marshall, D., Uller, T., 2007. When is a maternal effect adaptive? Oikos 116,
1957–1963. https://doi.org/10.1111/j.2007.0030-1299.16203.x

McNabb, F.M.A., 2007. The Hypothalamic-Pituitary-Thyroid (HPT) Axis in Birds
and Its Role in Bird Development and Reproduction. Crit Rev Toxicol 37,
163–193. https://doi.org/10.1080/10408440601123552

McNichols, M.J., McNabb, F.M.A., 1988. Development of thyroid function and
its pituitary control in embryonic and hatchling precocial Japanese quail
and altricial Ring doves. Gen Comp Endocrinol 70, 109–118.
https://doi.org/10.1016/0016-6480(88)90099-8

Mills, A.D., Faure, J.-M., 1991. Divergent selection for duration of tonic
immobility and social reinstatement behavior in Japanese quail (Coturnix
coturnix japonica) chicks. J Comp Psychol 105, 25–38.
https://doi.org/10.1037/0735-7036.105.1.25

Misiti, S., Schomburg, L., M. Yen, P., Chin, W.W., 1998. Expression and
Hormonal Regulation of Coactivator and Corepressor Genes.
Endocrinology 139, 2493–2500.
https://doi.org/10.1210/endo.139.5.5971

Moog, N.K., Entringer, S., Heim, C., Wadhwa, P.D., Kathmann, N., Buss, C.,
2017. Influence of maternal thyroid hormones during gestation on fetal
brain development. Neuroscience 342, 68–100.
https://doi.org/10.1016/j.neuroscience.2015.09.070

Mousseau, T., Fox, C., 1998. The adaptive significance of maternal effects.
Trends Ecol Evol 13, 403–407. https://doi.org/10.1016/S0169-
5347(98)01472-4

Okuliarová, M., Škrobánek, P., Zeman, M., 2007. Effect of Increasing Yolk
Testosterone Levels on Early Behaviour in Japanese Quail Hatchlings.
Acta Veterinaria Brno 76, 325–331.
https://doi.org/10.2754/avb200776030325

Patel, J., Landers, K., Li, H., Mortimer, R.H., Richard, K., 2011. Thyroid
hormones and fetal neurological development. Journal of Endocrinology
209, 1–8. https://doi.org/10.1530/JOE-10-0444

Podmokła, E., Drobniak, S.M., Rutkowska, J., 2018. Chicken or egg? Outcomes
of experimental manipulations of maternally transmitted hormones
depend on administration method - a meta-analysis. Biological Reviews
93, 1499–1517. https://doi.org/10.1111/brv.12406

Porazzi, P., Calebiro, D., Benato, F., Tiso, N., Persani, L., 2009. Thyroid gland
development and function in the zebrafish model. Mol Cell Endocrinol
312, 14–23. https://doi.org/10.1016/j.mce.2009.05.011

Russell V. Lenth, 2021. emmeans: Estimated Marginal Means, aka Least-
Squares Means.

Ruuskanen, S., Doligez, B., Pitala, N., Gustafsson, L., Laaksonen, T. 2012. Long-
term fitness consequences of maternal hormones: sons pay the costs.
Funct   Ecol   26,   884-894.   https://doi.org/10.1111/j.1365-
2435.2012.01994.x. https://doi.org/10.1016/j.yhbeh.2009.09.017

Ruuskanen, S., Laaksonen, T. 2010. Yolk hormones have sex-specific long-term
effects on behavior in the pied flycatcher. Horm Behav 57, 119–127

Ruuskanen, S., Darras, V.M., de Vries, B., Visser, M.E., Groothuis, T.G.G.,
2016a. Experimental manipulation of food availability leads to short-term
intra-clutch adjustment in egg mass but not in yolk androgen or thyroid
hormones. J Avian Biol 47, 36–46. https://doi.org/10.1111/jav.00728

Ruuskanen, S., Darras, V.M., Visser, M.E., Groothuis, T.G.G., 2016b. Effects of
experimentally manipulated yolk thyroid hormone levels on offspring
development in a wild bird species. Horm Behav 81, 38–44.
https://doi.org/10.1016/j.yhbeh.2016.03.006

Ruuskanen, S., Groothuis, T.G.G., Schaper, S. V., Darras, V.M., de Vries, B.,
Visser, M.E., 2016c. Temperature-induced variation in yolk androgen and
thyroid hormone levels in avian eggs. Gen Comp Endocrinol 235, 29–37.
https://doi.org/10.1016/j.ygcen.2016.05.026

Ruuskanen, S., Hsu, B.-Y., 2018. Maternal Thyroid Hormones: An Unexplored
Mechanism Underlying Maternal Effects in an Ecological Framework.
Physiological   and   Biochemical   Zoology   91,   904–916.
https://doi.org/10.1086/697380

Ruuskanen, S., Hsu, B.-Y., Heinonen, A., Vainio, M., Darras, V.M., Sarraude, T.,
Rokka, A., 2018. A new method for measuring thyroid hormones using
nano-LC-MS/MS. Journal of Chromatography B 1093–1094, 24–30.
https://doi.org/10.1016/j.jchromb.2018.06.052

Ruuskanen, S., Hukkanen, M., Garcin, N., Cossin-Sevrin, N., Hsu, B.-Y., Stier, A.,
2022. Altricial Bird Early-Stage Embryos Express the Molecular

"Machinery" to Respond to and Modulate Maternal Thyroid Hormone
Cues. Physiological and Biochemical Zoology 95, 544–550.
https://doi.org/10.1086/721556

Sarraude, T., Hsu, B.-Y., Groothuis, T., Ruuskanen, S., 2020. Testing the short-
and long-term effects of elevated prenatal exposure to different forms of
thyroid hormones. PeerJ 8, e10175. https://doi.org/10.7717/peerj.10175

Schoech, S.J., Rensel, M.A., Heiss, R.S., 2011. Short- and long-term effects of
developmental corticosterone exposure on avian physiology, behavioral
phenotype, cognition, and fitness: A review. Curr Zool 57, 514–530.
https://doi.org/10.1093/czoolo/57.4.514

Stier, A., Hsu, B.-Y., Marciau, C., Doligez, B., Gustafsson, L., Bize, P., Ruuskanen,
S., 2020. Born to be young? Prenatal thyroid hormones increase early-life
telomere length in wild collared flycatchers. Biol Lett 16, 20200364.
https://doi.org/10.1098/rsbl.2020.0364

Too, H.C., Shibata, M., Yayota, M., Darras, V.M., Iwasawa, A., 2017. Expression
of thyroid hormone regulator genes in the yolk sac membrane of the
developing chicken embryo. Journal of Reproduction and Development
63, 463–472. https://doi.org/10.1262/jrd.2017-017

Van Herck, S.L.J., Delbaere, J., Bourgeois, N.M.A., McAllan, B.M., Richardson,
S.J., Darras, V.M., 2015. Expression of thyroid hormone transporters and
deiodinases at the brain barriers in the embryonic chicken: Insights into
the regulation of thyroid hormone availability during neurodevelopment.
Gen Comp Endocrinol 214, 30–39.
https://doi.org/10.1016/j.ygcen.2015.02.021

Van Herck, S.L.J., Geysens, S., Delbaere, J., Tylzanowski, P., Darras, V.M., 2012.
Expression profile and thyroid hormone responsiveness of transporters
and deiodinases in early embryonic chicken brain development. Mol Cell
Endocrinol 349, 289–297. https://doi.org/10.1016/j.mce.2011.11.012

Vitorino Carvalho, A., Couroussé, N., Crochet, S., Coustham, V., 2019.
Identification of Reference Genes for Quantitative Gene Expression
Studies in Three Tissues of Japanese Quail. Genes (Basel) 10, 197.
https://doi.org/10.3390/genes10030197

von Engelhardt, N., Groothuis, T.G.G., 2011. Maternal Hormones in Avian Eggs,

719   in: Hormones and Reproduction of Vertebrates. Elsevier, pp. 91–127.

720   https://doi.org/10.1016/B978-0-12-374929-1.10004-6

721   Walpita, C.N., Van der Geyten, S., Rurangwa, E., Darras, V.M., 2007. The effect

722   of 3,5,3'-triiodothyronine supplementation on zebrafish (Danio rerio)

723   embryonic development and expression of iodothyronine deiodinases

724   and thyroid hormone receptors. Gen Comp Endocrinol 152, 206–214.

725   https://doi.org/10.1016/j.ygcen.2007.02.020

726   Wilson, C.M., McNabb, F.M.A., 1997. Maternal Thyroid Hormones in Japanese

727   Quail Eggs and Their Influence on Embryonic Development. Gen Comp

728   Endocrinol 107, 153–165. https://doi.org/10.1006/gcen.1997.6906

729   Yamaguchi, S., Aoki, N., Kitajima, T., Iikubo, E., Katagiri, S., Matsushima, T.,

730   Homma, K.J., 2012. Thyroid hormone determines the start of the

731   sensitive period of imprinting and primes later learning. Nat Commun 3,

732   1081. https://doi.org/10.1038/ncomms2088

733   Zhang, L., Hernández, V.S., Medina-Pizarro, M., Valle-Leija, P., Vega-González,

734   A., Morales, T., 2008. Maternal hyperthyroidism in rats impairs stress

735   coping of adult offspring. J Neurosci Res 86, 1306–1315.

736   https://doi.org/10.1002/jnr.21580

737   Zhang, L., Medina, M.P., Hernández, V.S., Estrada, F.S., Vega-González, A.,

738   2010. Vasopressinergic network abnormalities potentiate conditioned

739   anxious state of rats subjected to maternal hyperthyroidism.

740   Neuroscience 168, 416–428.

741   https://doi.org/10.1016/j.neuroscience.2010.03.059