Can increased prenatal exposure to thyroid hormones alter physiology

and behaviour in the long-term? Insights from an experimental study

in Japanese quails.

Kalle Aho[1,2*], Antoine Stier[1,3,4], Tom Sarraude[1,6], Bin-Yan Hsu[1], Suvi Ruuskanen[1,5*]

[1] Department of Biology, University of Turku FI-20014 Turku, Finland

[2] School of Medicine, Faculty of Health Sciences, University of Eastern Finland, Kuopio,
Finland

[3] Univ Lyon, Université Claude Bernard Lyon 1, CNRS, ENTPE, UMR 5023 LEHNA, F-
69622, Villeurbanne, France

[4] Institut Pluridisciplinaire Hubert Curien, UMR7178, Université de Strasbourg, CNRS,
Strasbourg, France

[5] Department of Biological and Environmental Science, University of Jyväskylä, Finland

[6] Groningen Institute for Evolutionary Life Sciences, University of Groningen, the
Netherlands

*corresponding authors: email: ahokal9@gmail.com, suvi.ruuskanen@gmail.com

Abstract

Maternal thyroid hormones (triiodothyronine T3 and thyroxine T4) are important regulators of embryonic development and gene expression. While maternal thyroid disorders are known to cause developmental issues in humans, variation in maternal thyroid hormones in healthy mothers have also been found to correlate with infant and child phenotypes. This suggests a potential impact of maternal thyroid hormones on offspring phenotype in an eco-evolutionary context. In chickens, prenatal thyroid hormone treatment has been shown to influence embryonic gene expression and postnatal treatment to influence imprinting and learning. However, the potential long-term effects of maternal thyroid hormones on physiology and behaviour are unclear. This study aims to investigate the long-term effects of maternal thyroid hormones on behaviour, plasma thyroid hormone levels and brain gene expression using the Japanese quail (*Coturnix japonica*) as a model. Egg hormone levels were elevated by injecting unincubated eggs with either saline (control), T3, T4 or a mixture of T3 and T4. Social motivation, boldness and fearfulness to predators were tested shortly after hatching and as adults. Plasma thyroid hormone levels and pallial expression of thyroid hormone receptor A, type 2 deiodinase, and nuclear receptor coactivator 1 were measured in adulthood. We found no evidence that elevated thyroid hormone levels in eggs affected behaviour, plasma hormone levels, or gene expression. This is the first study examining the potential long-term effects of elevated maternal thyroid hormones within the natural range. Although we found no evidence of long-term effects, other traits may still be affected and remain to be studied.

Keywords: maternal hormones, maternal effects, endocrinology, phenotypic plasticity, bird

Introduction

Maternal effects refer to the effect of the maternal phenotype or environment on the phenotype of the offspring. The effect can be mediated by hormones passed from the mother to the offspring, the maternal hormones. As the levels of maternal hormones are influenced by the maternal environment, they can transmit information about the maternal environment to the offspring. This allows the development of the offspring phenotype to be controlled by environmental factors (Marshall and Uller, 2007; Mousseau and Fox, 1998). Such hormone-mediated maternal effects can be observed in a wide range of traits, including physiology and behaviour (Groothuis et al., 2019, 2005; Ruuskanen and Hsu, 2018), and can have long-lasting effects into adulthood with potential fitness consequences (Groothuis et al., 2005; Schoech et al., 2011; von Engelhardt and Groothuis, 2011).

Compared to the more extensively studied androgens and glucocorticoids (Podmokła et al., 2018), the effects of maternally transferred thyroid hormones (THs) have been less explored (Darras, 2019; Ruuskanen and Hsu, 2018). THs (triiodothyronine, T3, and thyroxine, T4) play important roles in metabolism and development, where they regulate the expression of numerous genes (Bernal, 2017; Darras, 2019; Moog et al., 2017). TH action is regulated by several components of the thyroid system. TH receptors (THRA and THRB) and coactivators control the transcription of TH-responsive genes by forming a complex with the active form, T3 (Brent, 2012). The intracellular availability of T3 is regulated by several transmembrane TH transporters (e.g. MCT8, OATP1c1) that differentially transport either T4 or T3 (Groeneweg et al., 2020). Inside the cell, T3 availability can be further regulated by deiodinase enzymes that either convert T4 to T3 (deiodinase 2, D2) or convert both to inactive forms (deiodinase 3, D3) (Gereben et al., 2008). The expression of these components (TH receptors, transporters and deiodinases) begins very early in embryonic

development, allowing the embryo to regulate TH levels in an organ and cell-specific

manner (Darras, 2019; Darras et al., 2011; Flamant and Samarut, 1998; Ruuskanen et al.,

2022; Van Herck et al., 2012). However, the development of the thyroid gland occurs much

later in development (Forhead and Fowden, 2014; McNichols and McNabb, 1988; Porazzi et

al., 2009). Thus, in almost all vertebrates studied so far, proper embryonic development

depends on maternally transferred THs (Brown et al., 2014; Darras, 2019; Moog et al., 2017;

Patel et al., 2011).

In mammals, maternal THs are transferred across the placenta, whereas in egg-laying

animals such as birds, maternal THs are deposited in the egg yolk. In birds, high natural

variation in egg TH levels has been observed, up to 5-fold in magnitude (Ruuskanen and Hsu,

2018). This variation may be due to environmental factors such as food availability or

temperature (Hsu et al., 2016; Ruuskanen et al., 2016c), and has been reported at both the

inter- and intra-individual levels (Hsu et al., 2016, 2019b; Ruuskanen et al., 2016a, 2016c).

As THs are important in development (Darras et al., 2009; Decuypere et al., 2005; McNabb,

2007; Wilson and McNabb, 1997), and as increasing yolk TH levels has been found to

increase embryonic TH levels (Van Herck et al., 2012), high variation in egg TH levels could

therefore be a source of offspring phenotypic variation.

To date, most of what is known about phenotypic differences due to maternal THs

comes mainly from biomedical studies (Moog et al., 2017). However, studies using

pathological TH concentrations do not provide relevant information on the ecology or

evolution of maternal THs. Therefore, the effects of maternal THs within the natural range

need to be studied to understand whether variation has effects on traits affecting offspring

fitness. In addition, experiments manipulating maternal THs in mammals would inevitably

interfere with maternal physiology, making it difficult to attribute results solely to maternal

THs. Egg-laying animals, such as birds, have therefore become common models for studying

the functions of maternal hormones since maternal THs deposited in eggs can be

manipulated directly and independently. So far, elevated TH levels in the egg within the

natural range have been found to affect hatching success, offspring growth, plasma T3 level

and telomere length (Hsu et al., 2023a, 2019a, 2017; Ruuskanen et al., 2016b; Sarraude et

al., 2020; Stier et al., 2020).   Nevertheless, some of the reported effects appear to be

transient during the early post-hatching period (Hsu et al., 2019a). Whether the effects of

maternal THs can be long-lasting and potentially influence offspring fitness remains unclear.

To investigate the potential long-term effects of variation in maternal THs, we

experimentally elevated egg TH levels within the natural range in Japanese quail (*Coturnix*

japonica). With these experimental quails, we have previously reported a higher hatching

success caused by the elevated yolk TH levels, but no major effects on morphology or

physiology (Sarraude et al., 2020). Now we further studied changes in behaviour, plasma TH

levels and brain gene expression. While increasing maternal TH levels beyond normal

physiological range (maternal hyperthyroidism) has been shown to affect behaviour, for

example anxiety (Laureano-Melo et al., 2019; Zhang et al., 2010, 2008), it is not known

whether increasing maternal THs within the natural range affects behaviour. Also, as

postnatal THs have been shown to influence filial imprinting and learning behaviour

(Yamaguchi et al., 2012), it is interesting to study whether maternal THs, acting earlier in

development, may also have behavioural effects. Potential behavioural effects were

investigated in both chicks and adults, allowing for a broader timescale, by examining

boldness, social motivation and fearfulness to predators. Due to the lack of studies with

natural levels of elevated maternal THs and inconsistent results from studies on maternal

hyperthyroidism and offspring anxiety (Laureano-Melo et al., 2019; Zhang et al., 2010,

2008), the direction of potential behavioural changes is difficult to predict. In addition,

plasma TH levels were measured in adults, and changes in these would indicate long lasting

effects on TH function. While plasma TH levels have been reported to be affected by

elevated egg TH levels (within the natural range) in a sex-dependent manner in the rock

pigeons (Hsu et al., 2017), some more recent studies in wild passerines found no clear

effects of higher natural levels of egg THs on nestling plasma THs (Hsu et al., 2023b, 2020).

In terms of gene expression, previous studies have examined the effect of the natural

variation in maternal THs on brain gene expression only in bird embryos or young chicks

(Van Herck et al., 2015, 2012; Yamaguchi et al., 2012), while the potential long-term effects

of prenatal TH exposure remain unknown. Here we studied several genes involved in TH

function: *THRA* (thyroid hormone receptor A), *NCOA1* (nuclear receptor coactivator 1) and

DIO2 (type 2 deiodinase) in the pallium of adults. Expression of these genes has been shown

to be influenced by THs in other species (Misiti et al., 1998; Walpita et al., 2007).

Materials and methods

1.1 Materials

This study was conducted in accordance with the Finnish legislation and approved by

the Finnish Animal Ethics Committee (Eläinkoelautakunta; ESAVI / 1018 / 04.10.07 / 2016).

Japanese quail (*Coturnix japonica*), a commonly used laboratory model with a reference

genome, was used. Twenty-four breeding pairs were housed in cages with food and water

available ad libitum. The photoperiod was long (16L:8D) and the temperature stable at 20°

C. Eggs were collected from the cages in the morning and stored (15° C, 50% relative

humidity) for seven days before treatment.

1.2 Treatment of eggs, rearing of chicks and collection of samples

We manipulated egg TH levels by injecting hormones directly into the eggs prior to

embryonic development at a dose that mimicked increased maternal transfer (see details in

Sarraude et al., 2020). Eggs from each breeding pair were randomly assigned to four

treatment groups: T3, T4, T3 + T4 and control (0.9% saline). Eggs laid by the same pair were

always evenly distributed among all treatment groups. The injected hormone solutions were

prepared from crystallised powder (T3: CAS number 6893-02-3, Sigma-Aldrich, St. Louis,

MO, USA and T4: CAS number 51-48-9, Sigma-Aldrich, St. Louis, MO, USA). The amount of

hormone injected was based on previous measurements in the parental generation (mean ±

SD: T4, 15.3 ± 4.43 ng / egg; T3, 7.58 ± 2.32 ng / egg) and the aim was to increase hormone

contents in the yolk by two standard deviations (as recommended by Podmokla et al.,

2018). The injection protocol has been described in detail in Sarraude et al. (2020). Briefly,

each egg was injected with 50 µl of either the respective hormone solution (T3, T4 or T3+T4,

in 0.9% saline) or 0.9% saline (control, CO), delivered by using 27G insulin syringes (BD

Microlance^{TM}). After sealing the hole with sterile plaster (OPSITE Flexigrid, Smith &

Nephew), the eggs were incubated artificially (37.8 °C and 55% relative humidity). A total of

158 eggs were treated (39 T3, 39 T4, 40 T3+T4 and 40 CO), of which 66 chicks hatched (15

T3, 20 T4, 21 T3 + T4 and 10 CO), with sex distributions of chicks surviving to adulthood

(females / males): 7/4 for T3, 10/8 for T4, 9/12 for T3 + T4 and 3/4 for CO. Sex was

determined at maturity by examining the presence of the cloacal gland. The overall hatching

rate (~40%) was consistent with previous studies in domestic quail (Okuliarová et al., 2007).

The rearing conditions of the chicks have been described in detail in a previous

publication (Sarraude et al., 2020). In short, chicks were reared in groups (4-16 individuals,

depending on age) with treatments and sexes mixed, until sexual maturity (ca. 6–8 weeks

after hatching in Japanese quails), after which the birds were separated by sex to smaller

groups (4-5 individuals). Photoperiod was 16L:8D for the first 7 months and 8L:16

thereafter. At approximately 4.5 months of age, we collected blood samples from the

brachial vein, separated plasma, and stored it at -80° C. Birds were euthanised by

decapitation at 10 months of age. After decapitation, 40 mg of tissue was collected from the

pallium in PBS buffer, snap frozen in liquid nitrogen and stored at -80° C.

1.3 Behavioural experiments

To assess the potential effects of maternal THs on behaviour, we conducted a series of

behavioural tests (open arena, emergence and tonic immobility tests) at different life

stages. We followed previously published protocols that have been used extensively with

many different model organisms. The tests were conducted in separate rooms where the

birds had no visual or auditory contact with other individuals. The exception was the open

arena test, where the subject was able to hear the vocalisations of another bird being tested

at the same time in another room. All experiments were recorded, and the videos were

analysed blindly of the treatment. All videos of the same test at both ages (chick/adult)

were analysed by one person, except for the open arena test, where one person analysed

chick videos and another person analysed adult videos.

The open arena test measured social motivation and boldness and it followed a setup

developed by Forkman et al. (2007). The test was performed on both sexes at three days

and approximately 4.5 months of age. In the test, the bird was placed in a circular

illuminated arena and the movement duration and the number of escape attempts (attempt

to jump/fly over the wall) were measured from videos for 10 minutes (from the time the

researcher left the room). Only 10 of the 56 adults attempted to escape, so adult escape

attempts were not analysed. Subjects who moved less and made fewer escape attempts

were interpreted as less bold and less socially motivated.

The emergence test measured social motivation and boldness. The test was

performed on both sexes, twice on chicks (6-9d old) and once on adults (4.5 months old).

The test followed a setup developed by Forkman et al. (2007), in which a bird was placed in

a dark chamber connected to another, bright chamber. One minute after the subject was

placed in the dark chamber, a door connecting the chambers was removed. After removing

the door, the researcher left the observation room and the time it took for the subject to

completely exit the dark chamber was measured. Subjects who left the dark chamber more

quickly were interpreted as bolder and more socially motivated. The test lasted five

minutes. Whether the subject remained in the dark chamber (0) or left it (1) during the test

was also recorded.

The tonic immobility test (Mills and Faure, 1991) measured fearfulness to predators.

The test was performed on both sexes at 12 days and approximately 4.5 months of age. To

induce immobility, the bird was placed on its back in the centre of a V-shaped test apparatus

and restrained by hand for 15 seconds, after which the hand was removed, and the time

taken for the bird to resume a standing posture was measured. If the subject resumed the

standing posture in less than 10 seconds, this was recorded as a failed induction and the

induction procedure was repeated. Four adult subjects who failed five times were excluded.

If the subject remained stationary for more than 10 seconds, then the time to recovery was

measured. The longer a subject took to recover, the more fearful it was interpreted. The

test lasted 10 minutes. Five subjects (1 chick and 4 adults) did not recover for 10 minutes

and were excluded from the analysis (approximately 4% of all tests).

1.4 Plasma thyroid hormone levels

To test whether elevated maternal THs affect plasma TH levels later in life (4.5 months

of age), we extracted THs from plasma samples and measured them using a previously

validated LC-MS / MS protocol (Ruuskanen et al., 2018). Due to practical limitations (*i.e.*

volume of plasma available and haemolysis in some samples), the sample size was reduced

to 48 individuals (T3: 10, T4: 16, T3T4: 15, and 7 in the control). Hormone concentrations are

expressed in pmol/ml plasma. The repeatability of the TH measurements, was validated in

Ruuskanen et al. (2018) with CV% less than 10% (3.5-9.7%, depending on the hormone and

sample type)

1.5 Gene expression

To test for potential long-term changes in the expression of TH related genes (*DIO2*,

THRA, and *NCOA1*), we measured gene expression from pallium samples (10 months old).

RNA was extracted from the homogenised pallium samples using the NucleoSpin RNA Plus

kit (MACHEREY-NAGEL). Since the sample size of the control group was the main factor

limiting statistical power, we decided to analyse all control individuals and only an

equivalent number of individuals in the three treatment groups (N = 31, 7 in control and 8

per treatment group). Extraction was done from 50 µl of the homogenised brain sample

according to the manufacturer's instructions, except that the gDNA removal column was

centrifuged longer (1 min, 11000 g) and the RNA was eluted once with 40 µl of RNase-free

water. RNA concentration and purity were quantified by optical density, and all samples not

meeting our quality criteria (i.e. RNA concentration > 30 ng/µl and 260/280 > 1.80) were

excluded from further analysis. RNA integrity was checked using an E-Gel 2% electrophoresis

system (Invitrogen), and the ribosomal RNA 18S vs. 28S band intensity and was deemed

satisfactory. Samples were stored at -80° C for 2 weeks prior to cDNA synthesis. cDNA

synthesis was performed using the SensiFAST ™ cDNA Synthesis Kit (Bioline) according to

the manufacturer's instructions. The reaction was performed using 500 ng of RNA as the

starting material. A control (noRT) was prepared without reverse transcriptase enzyme. The

cDNA was diluted with water to a concentration of 1.2 ng/µl and stored at -20° C.

GAPDH and *PGK* were used as control genes, using primers previously designed and

validated in Japanese quail (Vitorino Carvalho et al., 2019). Stability of expression between

experimental groups was further verified using geNorm software (geNorm M < 0.5, geNorm

V < 0.15), and thus the geometric mean of these two genes was used as our reference gene.

Primers for the target genes were designed using the Primer-BLAST tool (NCBI). The primer

pairs (Table 1) were designed to have at least one intron between them. The suitability of

the primers in qPCR was tested prior to the actual analysis using different primer

concentrations and different amounts of cDNA. The tests confirmed the sufficient

amplification efficiency of the primers (between 90 and 110% based on the standard curve)

and the formation of only one product of the expected size based on both the presence of a

single narrow peak in the melting curve, and the presence of a single band of the expected

size on the agarose gel.

Table 1: qPCR primer set information and performance for gene expression assays. Cq refers to qPCR quantification cycle, efficiency was evaluated using the LinReg method based on individual well efficiency. Technical precision estimates are given for Cq as coefficient of variation CV (%) and for expression (ratio of target to reference genes) as technical repeatability (R).

Gene	Primers	Cq ± SD	Efficiency	CV Cq	Repeatability expression
GAPDH	F: TCTCTGTTGTTGACCTGACCTG R: ATGGCTGTCACCATTGAAGTC	15.0±0.50	98.0%	0.59%	
PGK1	F: CAAGCTCACCCTGGACAAGT R: GGACGGCTGCCTTGATTCTT	21.1±0.86	85.5%	0.48%	
DIO2	F: CTGGAACAGCTTCCTCCTGG R: ATCCAGAAGACGGCACTTGG	23.6±0.83	92.9%	0.40%	0.91
NCOA1	F: CGACTCACGGAAGAGGAAGG R: GTGTCGATGTCCCCGATGTT	20.4±2.82	91.9%	0.38%	0.84
THRA	F: TGGTGAAGAGCAGCATGTCA R: AGCACCCATCGTACTTGCAG	21.4±0.82	96.0%	0.46%	0.82

qPCR assays were performed on a Mic qPCR (Bio Molecular Systems) in a final volume of 12

µL containing 6 ng of cDNA, 800 nM of forward and reverse primers and SensiFAST SYBR Lo-

ROX (Bioline). We ran gene-specific qPCR plates and each sample was analysed in duplicate.

Two plates were run for each gene and experimental groups and sex were always balanced

within each plate. We used a cDNA reference sample (*i.e.* ratio = 1), which was a pool of 10

different individuals on each plate. We used a two-step cycling with the following

conditions: 2 min at 95°C; then 45 cycles of 5s at 95°C followed by 20s at 60°C (fluorescence

reading) for all reactions. The expression of each gene was calculated as $(1+Ef_{Target})^{\Delta Cq^{(Target)}}/$

geometric mean $[(1+Ef_{GAPDH})^{\Delta Cq^{GAPDH}}; (1+Ef_{PGK1})^{\Delta Cq^{PGK1}}]$, where Ef is the amplification's

efficiency and ΔCq is the difference between the Cq-values of the reference sample and the

sample of interest. Amplification in the no-RT controls never occurred until at least 8 cycles

after the lower Cq sample, and thus contamination by genomic DNA could not interfere with

our results. Technical precision and repeatability were generally good (Table 1), but a few

samples with a CV% of duplicates (based on the final ratio) greater than 15% were excluded

from the analysis (final sample size: N_{DIO2} = 27, N_{NCOA1} = 28, N_{THRa} = 26)

1.6 Statistical analysis

All statistical analyses were performed with R software (version 3.6.2), using the packages

lme4 version 1.1-23 (Bates et al., 2015) and emmeans version 1.6.0 (Lenth, 2021). Normality

and homoscedasticity of the residual distributions were checked visually. For the

behavioural analysis, recovery time (tonic immobility) and exit time (emergence test), were

transformed (natural logarithm) to ensure that the residuals were normally distributed.

Linear mixed models (LMM) were used to analyse time to exit in the emergence test,

movement duration and number of escape attempts in the open arena test, recovery time

in the tonic immobility test, plasma TH concentrations and qPCR results. Generalised linear

mixed model (GLMM) with a binomial error distribution was used to analyse the proportion

of individuals leaving the dark chamber in the emergence test. Fixed effects were treatment,

sex, and test number/age (in repeated tests). Interactions between sex and treatment, and

between treatment and age (in repeated tests), were also analysed. Random effects were

individual ID (in repeated tests) and parental ID (to account for relatedness between siblings

in all experiments). F- and p-values for LMMs were obtained using the Kenward-Roger

approximation on the denominator degree of freedom, implemented in the package

lmerTest (Kuznetsova et al., 2017). For binomial regression models, main effects and

interactions were tested by comparing the model with and without the factor. Non-

significant (p > 0.05) interactions were excluded from the final models. Post-hoc tests were

performed using Tukey tests where appropriate.

Results

Effects of prenatal thyroid hormone supplementation on behaviour

In general, no statistically significant differences in chick or adult behavioural traits

were observed between the egg TH manipulation groups (T3, T4, T3 + T4, control). Elevated

THs had no significant effect on social motivation or boldness (Table 2, Fig. 1-2), nor on

fearfulness to predators (Table 2, Fig. 3). A statistically significant interaction between

treatment and age was found in the emergence test (F = 3.58, ddf = 67.2, p = 0.018, Fig. 2):

In all treatments except the control group (Tukey t=0.59, p=0.56), chick exit times were

significantly longer than those of adults (all t > 3.21, p < 0.002) while no differences between

treatments were found within each age-category (all t < 2.48, all p> 0.07). There was also a

statistically significant interaction between treatment and sex in the tonic immobility test (F

= 2.92, ddf = 37.5, p = 0.047, Fig. 3): In males, the recovery time was longer in the T4-group

(marginal mean ± SE: 177.1 ± 47.8 s) than in the T3-group (marginal mean ± SE: 56.1 ± 20.8

s, Tukey t = 2.71, p = 0.047), whereas no differences between treatments were observed in

females (all t < 2.29, all p >0.12). In the T4-group, the recovery time of males (marginal

mean ± SE: 177.1 ± 47.8 s) was longer than that of females (marginal mean ± SE: 85.2 ± 19.9

s) (t=2.16, p= 0.037), while no sex differences were observed within the other treatments

(all t<1.80, all p>0.079).

 Table 2. Marginal means and standard errors of behavioural tests. Results for chicks (age: 3d,
 6-9d, 12d) and adults (age: 4.5 months) are shown separately for each variable. F- and p-values
 for treatment from linear models (ANOVA, Kenward-Roger approximation). Test statistics (χ^2)
 and p-values for binomial regressions (model with treatment compared to model without
 treatment) were calculated with ANOVA.

Test: measurement	Age	Control	T3	T3T4	T4	F_{ddf}/χ^2 (df)	P
Tonic immobility:	12 d	92.0±25.5	84.6±21.1	68.4±13.1	123±24.7	$2.52_{36.2}$	0.073
Recovery time (s)	4.5 m	90.6±25.3	83.3±21.4	67.3±13.6	121±25.5		
Open arena:	3 d	77.2±31.1	100±27.1	73.3±19.1	63.3±20.8	$0.44_{46.6}$	0.72
Movement duration (s)	4.5 m	181±31.6	204±28.1	177±19.3	167±21.3		
Open arena:	3 d	5.85±3.87	4.14±3.53	8.25±2.14	7.21±2.43	$0.35_{41.7}$	0.79
Escape attempts (number)							
Emergence:	6–9 d	34.8±14.7	42.7±16.4	74.7±20.9	121±36.1	$1.92_{39.6}$	0.14
Exit time (s)	4.5 m	25.6±12.5	6.59±2.75	23.0±7.64	12.6±4.30		
Emergence:	6–9 d	0.80±0.11	0.70±0.12	0.62±0.10	0.58±0.10	2.71 (3)	0.44
Likelihood of emergence (0-1)	4.5 m	0.91±0.06	0.86±0.08	0.81±0.08	0.78±0.09		

Figure 1. Movement duration for chicks (3 d) and adults (4.5 months) in the open arena test

for different prenatal thyroid hormone treatment groups. The lines represent the medians of

the treatments, and the bars represent the interquartile range of the original observations.

Sample size (chicks/adults): CO 7/7, T3 11/11, T3T4 20/21 and T4 16/17.

Figure 2. Time to exit the dark chamber for chicks (6-9 d) and adults (4.5 months) in the

emergence test for different prenatal thyroid hormone treatment groups. The test was

performed twice for chicks and once for adults. The lines represent the medians of the

treatments, and the bars represent the interquartile range of the original observations. Sample

size (chicks/adults): CO 11/6, T3 15/9, T3T4 24/13 and T4 22/13.

Figure 3. Time to resume standing position for females and males, chicks (12 d) and adults (4.5 months) combined, in the tonic immobility test for different prenatal thyroid hormone treatment groups. The lines represent the medians of the treatments, and the bars represent the interquartile range of the original observations. Sample size (female/male: CO 6/8, T3 13/7, T3T4 14/23 and T4 18/14.

Effects of prenatal thyroid hormone supplementation on plasma thyroid hormone levels

Increasing TH levels in eggs did not significantly affect plasma TH levels in 4.5-month-old birds (T3: F = 0.13, ddf = 37.7, p = 0.94, T4: F = 2.1, ddf = 41.5, p = 0.12, Fig. 4). Treatment also had no sex-dependent effect on plasma T3 or T4 (treatment-sex interactions, p > 0.36).

Figure 4. Plasma T3 and T4 concentrations in 4.5-month-old birds between different prenatal

thyroid hormone treatment groups. The lines represent the medians of the treatments, and

the bars represent the interquartile range of the original observations. Sample size (T3/T4

concentrations): CO 6/7, T3 10/10, T3T4 14/15 and T4 14/16.

Effects of prenatal thyroid hormone supplementation on pallial gene

expression

Increasing TH levels in eggs did not affect pallial gene expression of TH-related

genes in 10-month-old birds (*DIO2*: F = 2.49, ddf = 17.3, p = 0.09, *NCOA1*: F = 0.092, ddf

= 21.1, p = 0.96, *THRA*: F = 0.47, ddf = 18.9, p = 0.71, Fig. 5). Treatment also did not

have sex-dependent effect on gene expression (treatment-sex interactions, p > 0.26).

Females had higher expression of *NCOA1* (F=7.68, p=0.012) and *THRA* (F=5.17,

p=0.034) than males.

Figure 5. Relative expressions of *DIO2*, *THRA* and *NCOA1* in the pallium of 10-month-old birds

between different prenatal thyroid hormone treatments. The lines represent the medians of

the treatments, and the bars represent the interquartile range of the original observations.

Sample sizes (CO/T3/T3T4/T4): DIO2 5/7/8/7, THRA 6/4/8/8 and NCOA1 6/7/8/7.

Discussion

In this study, we tested for long-term changes in behaviour, plasma TH levels

and pallial gene expression following increased exposure to maternal THs. In terms of

behaviour, we observed no significant changes in boldness, social motivation or

fearfulness to predators in chicks (3-12 d) or adults (4.5 months) when the egg TH level

was increased. To our knowledge, there are no similar previous studies, and it is

therefore difficult to assess whether there may have been changes in other

behavioural traits. The detection of changes was also partly limited by the small

sample size, which meant that we may not have had enough statistical power to detect

subtle changes in behaviour. Nevertheless, we found some age-dependent and sex-

dependent effects of TH treatment on behaviour. The age-dependent effect in the

emergence test suggests that elevated maternal TH leads to a decrease in boldness

and social motivation with age (from chick to adult), whereas such a decrease is not

observed in controls. However, the lack of differences between control and treatment

within a given age suggests that such an effect (if not associated with type I statistical

error) is rather mild. The sex-dependent effect in the tonic immobility test also seems

relatively unconvincing because no sex differences were observed within treatments,

nor did T3T4 females or T4 males differ from the same-sex controls. Given the

important role of THs in brain development, we expected that elevated maternal THs

would have led to behavioural changes. Yet, our data suggest that an experimental

increase in maternal THs within the natural range has no clear effect on the

behavioural traits measured. However, in contrast to maternal THs, post-hatch TH

treatment has been shown to affect imprinting and learning (Yamaguchi et al., 2012).

The discrepancy between our results and Yamaguchi et al. (2012) could suggest that

the sensitive period during which THs can influence behaviour might occur at a later

timing of development, but this requires further investigation.

We found that elevated maternal THs did not significantly affect plasma TH

levels in adulthood (4.5 months of age). This was the first study to test for long-term

changes, whereas previously changes in response to elevated prenatal THs have been

observed at younger ages: Experimental elevation of egg THs was found to increase

male and decrease female plasma TH levels in rock pigeons at 14 days of age (Hsu et

al., 2017), but such effects were not observed in two other wild passerine species (Hsu

et al., 2023b, 2020). In contrast to these studies in altricial species, not only was this

study conducted on a precocial species, but our measurements were also made at a

different life stage (adulthood). Overall, this suggests that an increase in maternal THs

may only lead to transient changes in offspring plasma TH levels and that the effects

may not be universal across species.

We also found no significant effect of elevated maternal THs on the expression

of *THRA*, *DIO2* or *NCOA1* in the pallium of adult (10 months old) individuals. Changes in

gene expression have been observed in other TH-related genes, but again, at much

shorter time spans: In chickens, injection of THs into the yolk decreased the expression

of the TH-transporter *OATP1C1* and increased the expression of the TH-transporter

MCT8 in 4-day-old embryos (Van Herck et al., 2012). The expressions of our target

genes, *THRA*, *DIO2* and *NCOA1,* have also been shown to be at least transiently

responsive to THs in other species: *THRA* and *DIO2* in zebrafish embryos (Walpita et al.,

2007), and *NCOA1* in adult rats (Misiti et al., 1998). It is therefore possible that the

gene regulation by maternal THs is restricted to early development when the embryo

is not yet able to produce THs.

We have previously shown in the same experimental birds, that an increase in

prenatal THs influences hatching success (Sarraude et al., 2020). Therefore, the dose

used in this study can have some effects during embryo development, which suggests

that the lack of changes in long-term behaviour, plasma TH levels and brain gene

expression may be linked to an attenuation of prenatal TH effects by embryonic

regulation. It is indeed possible that embryos could buffer the effect of maternal THs

by altering the expression of deiodinases which are expressed already at very early

stages (Ruuskanen et al., 2022; Too et al., 2017). Metabolisation of THs to inactive

forms could prevent the elevated egg THs from exerting any effects. However, given

the effects previously observed in the same experimental birds on hatching success

(Sarraude et al., 2020), the lack of changes in this study may also be explained by the

stage of ontogeny, prenatal THs not leading to long-lasting effects. Nevertheless, the

lack of prenatal measurements means that this cannot be fully confirmed, and it

remains possible that elevated maternal THs have no effect on the measured traits in

Japanese quails.

In conclusion, we found no evidence that elevated maternal THs cause long-

term changes in behaviour, plasma TH levels or pallial gene expression. This is

consistent with our previous findings in the same experimental birds, where elevation

of prenatal THs (T3, T4, or in combination) did not lead to long-term effects on

oxidative stress, moult or reproduction (Sarraude et al., 2020). While behavioural

changes have not been previously studied, plasma TH concentrations and gene

expression have been shown to respond to elevated maternal THs in the short-term

(Hsu et al., 2017; Van Herck et al., 2012). However, based on this study, these changes

would appear to be limited to early development or shortly after hatching with the

offspring's own TH production later replacing the effects of elevated maternal THs.

Maternal THs could theoretically influence the offspring's own TH production at a later

age by altering the function of the HPT-axis, but no evidence has yet been found to

support this hypothesis. The long-term effects of variation in maternal THs remain

unclear, but this study has provided new information on some previously untested

traits. Understanding long-term effects in addition to short-term effects will help to

better understand the evolutionary and ecological importance of maternal THs.

Acknowledgements

We thank personnel at the animal care facilities, Marjo Myllyrinne, Simo Laine for their

help.

Funding

The study was funded by the Academy of Finland (project #286278) to SR, the Finnish

National Agency for Education (grant no. TM-15-9960 to TS). AS was supported by a

Turku Collegium for Science and Medicine fellowship and an European Commission

Marie Sklodowska-Curie Fellowship #894963.

Data availability statement

Data will be available on Figshare (DOI: 10.6084/m9.figshare.22266211) at publication.

Author contributions

S.R., T.S., B-Y.H, A.S and K.A. designed the study. S.R. obtained funding. T.S., B-Y.H and

S.R conducted the experiment. K.A. and T.S. collected the behavioural data from

videos. K.A., A.S and S.R. collected the physiological data and conducted laboratory

work. K.A. analysed all data with support from all co-authors, and wrote the first draft

of the manuscript. A.S, B-Y.H, T.S. and S.R commented on the manuscript.

References

Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects
Models Using lme4. J Stat Softw 67. https://doi.org/10.18637/jss.v067.i01
Bernal, J., 2017. Thyroid hormone regulated genes in cerebral cortex development.
Journal of Endocrinology 232, R83–R97. https://doi.org/10.1530/JOE-16-0424

Brent, G.A., 2012. Mechanisms of thyroid hormone action. Journal of Clinical
Investigation 122, 3035–3043. https://doi.org/10.1172/JCI60047

Brown, C.L., Urbinati, E.C., Zhang, W., Brown, S.B., McComb-Kobza, M., 2014.
Maternal Thyroid and Glucocorticoid Hormone Interactions in Larval Fish
Development, and Their Applications in Aquaculture. Reviews in Fisheries
Science & Aquaculture 22, 207–220.
https://doi.org/10.1080/23308249.2014.918086

Darras, V.M., 2019. The Role of Maternal Thyroid Hormones in Avian Embryonic
Development. Front Endocrinol (Lausanne) 10.
https://doi.org/10.3389/fendo.2019.00066

Darras, V.M., Van Herck, S.L.J., Geysens, S., Reyns, G.E., 2009. Involvement of
thyroid hormones in chicken embryonic brain development. Gen Comp Endocrinol
163, 58–62. https://doi.org/10.1016/j.ygcen.2008.11.014

Darras, V.M., Van Herck, S.L.J., Heijlen, M., De Groef, B., 2011. Thyroid Hormone
Receptors in Two Model Species for Vertebrate Embryonic Development: Chicken
and Zebrafish. J Thyroid Res 2011, 1–8. https://doi.org/10.4061/2011/402320

Decuypere, E., Van As, P., Van der Geyten, S., Darras, V.M., 2005. Thyroid hormone
availability and activity in avian species: A review. Domest Anim Endocrinol 29,
63–77. https://doi.org/10.1016/j.domaniend.2005.02.028

Flamant, F., Samarut, J., 1998. Involvement of Thyroid Hormone and Its α Receptor in
Avian Neurulation. Dev Biol 197, 1–11. https://doi.org/10.1006/dbio.1998.8872

Forhead, A.J., Fowden, A.L., 2014. Thyroid hormones in fetal growth and prepartum
maturation. Journal of Endocrinology 221, R87–R103.
https://doi.org/10.1530/JOE-14-0025

Forkman, B., Boissy, A., Meunier-Salaün, M.-C., Canali, E., Jones, R.B., 2007. A
critical review of fear tests used on cattle, pigs, sheep, poultry and horses. Physiol
Behav 92, 340–374. https://doi.org/10.1016/j.physbeh.2007.03.016

Gereben, B., Zeöld, A., Dentice, M., Salvatore, D., Bianco, A.C., 2008. Activation and
inactivation of thyroid hormone by deiodinases: Local action with general
consequences. Cellular and Molecular Life Sciences 65, 570–590.
https://doi.org/10.1007/s00018-007-7396-0

Groeneweg, S., van Geest, F.S., Peeters, R.P., Heuer, H., Visser, W.E., 2020. Thyroid
Hormone Transporters. Endocr Rev 41, 146–201.
https://doi.org/10.1210/endrev/bnz008

Groothuis, T.G.G., Hsu, B.-Y., Kumar, N., Tschirren, B., 2019. Revisiting mechanisms
and functions of prenatal hormone-mediated maternal effects using avian species
as a model. Philosophical Transactions of the Royal Society B: Biological Sciences
374, 20180115. https://doi.org/10.1098/rstb.2018.0115

Groothuis, T.G.G., Müller, W., von Engelhardt, N., Carere, C., Eising, C., 2005.
Maternal hormones as a tool to adjust offspring phenotype in avian species.
Neurosci Biobehav Rev 29, 329–352.
https://doi.org/10.1016/j.neubiorev.2004.12.002

Hsu, B., Dijkstra, C., Darras, V.M., Vries, B., Groothuis, T.G.G., 2016. Maternal
adjustment or constraint: differential effects of food availability on maternal
deposition of macro nutrients, steroids and thyroid hormones in rock pigeon eggs.
Ecol Evol 6, 397–411. https://doi.org/10.1002/ece3.1845

Hsu, B.-Y., Cossin-Sevrin, N., Stier, A., Ruuskanen, S., 2023a. Prenatal thyroid
hormones accelerate postnatal growth and telomere shortening in wild great tits.
Journal of Experimental Biology. https://doi.org/10.1242/jeb.243875

Hsu, B.-Y., Cossin-Sevrin, N., Stier, A., Ruuskanen, S., 2023b. Prenatal thyroid
hormones accelerate postnatal growth and telomere shortening in wild great tits.
Journal of Experimental Biology. https://doi.org/10.1242/jeb.243875

Hsu, B.-Y., Dijkstra, C., Darras, V.M., de Vries, B., Groothuis, T.G.G., 2017. Maternal
thyroid hormones enhance hatching success but decrease nestling body mass in the
rock pigeon (Columba livia). Gen Comp Endocrinol 240, 174–181.
https://doi.org/10.1016/j.ygcen.2016.10.011

Hsu, B.-Y., Doligez, B., Gustafsson, L., Ruuskanen, S., 2019a. Transient growth-
enhancing effects of elevated maternal thyroid hormones at no apparent oxidative
cost during early postnatal period. J Avian Biol 50.
https://doi.org/10.1111/jav.01919

Hsu, B.-Y., Sarraude, T., Cossin-Sevrin, N., Crombecque, M., Stier, A., Ruuskanen, S.,
2020. Testing for context-dependent effects of prenatal thyroid hormones on
offspring survival and physiology: an experimental temperature manipulation. Sci
Rep 10, 14563. https://doi.org/10.1038/s41598-020-71511-y

Hsu, B.-Y., Verhagen, I., Gienapp, P., Darras, V.M., Visser, M.E., Ruuskanen, S.,
2019b. Between- and Within-Individual Variation of Maternal Thyroid Hormone
Deposition in Wild Great Tits (*Parus major*). Am Nat 194, E96–E108.
https://doi.org/10.1086/704738

Kuznetsova, A., Brockhoff, P.B., Christensen, R.H.B., 2017. **lmerTest** Package: Tests

in Linear Mixed Effects Models. J Stat Softw 82.

https://doi.org/10.18637/jss.v082.i13

Laureano-Melo, R., Souza, J.S. de, da Conceição, R.R., Albuquerque, J.M.L.,

Rodrigues, N.C., Marinho, B.G., Olivares, E.L., Giannocco, G., Côrtes, W. da S.,

2019. Prenatal thyroxine treatment promotes anxiolysis in male Swiss mice

offspring. Horm Behav 108, 10–19. https://doi.org/10.1016/j.yhbeh.2018.12.008

Marshall, D., Uller, T., 2007. When is a maternal effect adaptive? Oikos 116, 1957–

1963. https://doi.org/10.1111/j.2007.0030-1299.16203.x

McNabb, F.M.A., 2007. The Hypothalamic-Pituitary-Thyroid (HPT) Axis in Birds and

Its Role in Bird Development and Reproduction. Crit Rev Toxicol 37, 163–193.

https://doi.org/10.1080/10408440601123552

McNichols, M.J., McNabb, F.M.A., 1988. Development of thyroid function and its

pituitary control in embryonic and hatchling precocial Japanese quail and altricial

Ring doves. Gen Comp Endocrinol 70, 109–118. https://doi.org/10.1016/0016-

6480(88)90099-8

Mills, A.D., Faure, J.-M., 1991. Divergent selection for duration of tonic immobility

and social reinstatement behavior in Japanese quail (Coturnix coturnix japonica)

chicks. J Comp Psychol 105, 25–38. https://doi.org/10.1037/0735-7036.105.1.25

Misiti, S., Schomburg, L., M. Yen, P., Chin, W.W., 1998. Expression and Hormonal

Regulation of Coactivator and Corepressor Genes. Endocrinology 139, 2493–2500.

https://doi.org/10.1210/endo.139.5.5971

Moog, N.K., Entringer, S., Heim, C., Wadhwa, P.D., Kathmann, N., Buss, C., 2017.

Influence of maternal thyroid hormones during gestation on fetal brain

development. Neuroscience 342, 68–100.

https://doi.org/10.1016/j.neuroscience.2015.09.070

Mousseau, T., Fox, C., 1998. The adaptive significance of maternal effects. Trends Ecol

Evol 13, 403–407. https://doi.org/10.1016/S0169-5347(98)01472-4

Okuliarová, M., Škrobánek, P., Zeman, M., 2007. Effect of Increasing Yolk

Testosterone Levels on Early Behaviour in Japanese Quail Hatchlings. Acta

Veterinaria Brno 76, 325–331. https://doi.org/10.2754/avb200776030325

Patel, J., Landers, K., Li, H., Mortimer, R.H., Richard, K., 2011. Thyroid hormones and

fetal neurological development. Journal of Endocrinology 209, 1–8.

https://doi.org/10.1530/JOE-10-0444

Podmokła, E., Drobniak, S.M., Rutkowska, J., 2018. Chicken or egg? Outcomes of
experimental manipulations of maternally transmitted hormones depend on
administration method - a meta-analysis. Biological Reviews 93, 1499–1517.
https://doi.org/10.1111/brv.12406

Porazzi, P., Calebiro, D., Benato, F., Tiso, N., Persani, L., 2009. Thyroid gland
development and function in the zebrafish model. Mol Cell Endocrinol 312, 14–23.
https://doi.org/10.1016/j.mce.2009.05.011

Russell V. Lenth, 2021. emmeans: Estimated Marginal Means, aka Least-Squares
Means.

Ruuskanen, S., Darras, V.M., de Vries, B., Visser, M.E., Groothuis, T.G.G., 2016a.
Experimental manipulation of food availability leads to short-term intra-clutch
adjustment in egg mass but not in yolk androgen or thyroid hormones. J Avian Biol
47, 36–46. https://doi.org/10.1111/jav.00728

Ruuskanen, S., Darras, V.M., Visser, M.E., Groothuis, T.G.G., 2016b. Effects of
experimentally manipulated yolk thyroid hormone levels on offspring development
in a wild bird species. Horm Behav 81, 38–44.
https://doi.org/10.1016/j.yhbeh.2016.03.006

Ruuskanen, S., Groothuis, T.G.G., Schaper, S. V., Darras, V.M., de Vries, B., Visser,
M.E., 2016c. Temperature-induced variation in yolk androgen and thyroid
hormone levels in avian eggs. Gen Comp Endocrinol 235, 29–37.
https://doi.org/10.1016/j.ygcen.2016.05.026

Ruuskanen, S., Hsu, B.-Y., 2018. Maternal Thyroid Hormones: An Unexplored
Mechanism Underlying Maternal Effects in an Ecological Framework.
Physiological and Biochemical Zoology 91, 904–916.
https://doi.org/10.1086/697380

Ruuskanen, S., Hsu, B.-Y., Heinonen, A., Vainio, M., Darras, V.M., Sarraude, T.,
Rokka, A., 2018. A new method for measuring thyroid hormones using nano-LC-
MS/MS. Journal of Chromatography B 1093–1094, 24–30.
https://doi.org/10.1016/j.jchromb.2018.06.052

Ruuskanen, S., Hukkanen, M., Garcin, N., Cossin-Sevrin, N., Hsu, B.-Y., Stier, A.,
2022. Altricial Bird Early-Stage Embryos Express the Molecular "Machinery" to
Respond to and Modulate Maternal Thyroid Hormone Cues. Physiological and
Biochemical Zoology 95, 544–550. https://doi.org/10.1086/721556

Sarraude, T., Hsu, B.-Y., Groothuis, T., Ruuskanen, S., 2020. Testing the short-and long-term effects of elevated prenatal exposure to different forms of thyroid hormones. PeerJ 8, e10175. https://doi.org/10.7717/peerj.10175

Schoech, S.J., Rensel, M.A., Heiss, R.S., 2011. Short- and long-term effects of developmental corticosterone exposure on avian physiology, behavioral phenotype, cognition, and fitness: A review. Curr Zool 57, 514–530. https://doi.org/10.1093/czoolo/57.4.514

Stier, A., Hsu, B.-Y., Marciau, C., Doligez, B., Gustafsson, L., Bize, P., Ruuskanen, S., 2020. Born to be young? Prenatal thyroid hormones increase early-life telomere length in wild collared flycatchers. Biol Lett 16, 20200364. https://doi.org/10.1098/rsbl.2020.0364

Too, H.C., Shibata, M., Yayota, M., Darras, V.M., Iwasawa, A., 2017. Expression of thyroid hormone regulator genes in the yolk sac membrane of the developing chicken embryo. Journal of Reproduction and Development 63, 463–472. https://doi.org/10.1262/jrd.2017-017

Van Herck, S.L.J., Delbaere, J., Bourgeois, N.M.A., McAllan, B.M., Richardson, S.J., Darras, V.M., 2015. Expression of thyroid hormone transporters and deiodinases at the brain barriers in the embryonic chicken: Insights into the regulation of thyroid hormone availability during neurodevelopment. Gen Comp Endocrinol 214, 30–39. https://doi.org/10.1016/j.ygcen.2015.02.021

Van Herck, S.L.J., Geysens, S., Delbaere, J., Tylzanowski, P., Darras, V.M., 2012. Expression profile and thyroid hormone responsiveness of transporters and deiodinases in early embryonic chicken brain development. Mol Cell Endocrinol 349, 289–297. https://doi.org/10.1016/j.mce.2011.11.012

Vitorino Carvalho, A., Couroussé, N., Crochet, S., Coustham, V., 2019. Identification of Reference Genes for Quantitative Gene Expression Studies in Three Tissues of Japanese Quail. Genes (Basel) 10, 197. https://doi.org/10.3390/genes10030197

von Engelhardt, N., Groothuis, T.G.G., 2011. Maternal Hormones in Avian Eggs, in: Hormones and Reproduction of Vertebrates. Elsevier, pp. 91–127. https://doi.org/10.1016/B978-0-12-374929-1.10004-6

Walpita, C.N., Van der Geyten, S., Rurangwa, E., Darras, V.M., 2007. The effect of 3,5,3'-triiodothyronine supplementation on zebrafish (Danio rerio) embryonic development and expression of iodothyronine deiodinases and thyroid hormone receptors. Gen Comp Endocrinol 152, 206–214. https://doi.org/10.1016/j.ygcen.2007.02.020

Wilson, C.M., McNabb, F.M.A., 1997. Maternal Thyroid Hormones in Japanese Quail
Eggs and Their Influence on Embryonic Development. Gen Comp Endocrinol 107,
153–165. https://doi.org/10.1006/gcen.1997.6906

Yamaguchi, S., Aoki, N., Kitajima, T., Iikubo, E., Katagiri, S., Matsushima, T.,
Homma, K.J., 2012. Thyroid hormone determines the start of the sensitive period
of imprinting and primes later learning. Nat Commun 3, 1081.
https://doi.org/10.1038/ncomms2088

Zhang, L., Hernández, V.S., Medina-Pizarro, M., Valle-Leija, P., Vega-González, A.,
Morales, T., 2008. Maternal hyperthyroidism in rats impairs stress coping of adult
offspring. J Neurosci Res 86, 1306–1315. https://doi.org/10.1002/jnr.21580

Zhang, L., Medina, M.P., Hernández, V.S., Estrada, F.S., Vega-González, A., 2010.
Vasopressinergic network abnormalities potentiate conditioned anxious state of
rats subjected to maternal hyperthyroidism. Neuroscience 168, 416–428.
https://doi.org/10.1016/j.neuroscience.2010.03.059