Genetic and dietary modulators of the inflammatory response in the

gastro-intestinal tract of the BXD mouse genetic reference population

Xiaoxu Li[1], Jean-David Morel[1], Giorgia Benegiamo[1], Johanne Poisson[1], Alexis Bachmann[1],

Alexis Rapin[1], Evan Williams[3], Alessia Perino[2], Kristina Schoonjans[2], Maroun Bou

Sleiman[1,*], Johan Auwerx[1,*]

[1]Laboratory of Integrative Systems Physiology, Institute of Bioengineering, École

Polytechnique Fédérale de Lausanne, 1015 Lausanne, Switzerland.

[2]Laboratory of Metabolic Signaling, Institute of Bioengineering, École Polytechnique

Fédérale de Lausanne, 1015 Lausanne, Switzerland.

[3]Luxembourg Centre for Systems Biomedicine, University of Luxembourg, Luxembourg,

Luxembourg.

*e-mail for correspondence: maroun.bousleiman@epfl.ch (MBS) or admin.auwerx@epfl.ch

(JA)

Abstract

Inflammatory gut disorders, including inflammatory bowel disease (IBD), can be impacted by

dietary, environmental and genetic factors. While the incidence of IBD is increasing worldwide,

we still lack a complete understanding of the gene-by-environment interactions underlying

inflammation and IBD. Here, we profiled the colon transcriptome of 52 BXD mouse strains

fed with a chow or high-fat diet (HFD) and identified a subset of BXD strains that exhibit an

IBD-like transcriptome signature on HFD, indicating that an interplay of genetics and diet can

significantly affect intestinal inflammation. Using gene co-expression analyses, we identified

modules that are enriched for IBD-dysregulated genes and found that these IBD-related

modules share *cis*-regulatory elements that are responsive to the STAT2, SMAD3, and REL

transcription factors. We used module quantitative trait locus (ModQTL) analyses to identify

genetic loci associated with the expression of these modules. Through a prioritization scheme

involving systems genetics in the mouse and integration with external human datasets, we

identified *Muc4* and *Epha6* as the top candidates mediating differences in HFD-driven

intestinal inflammation. This work provides insights into the contribution of genetics and diet

to IBD risk and identifies two candidate genes, *MUC4* and *EPHA6*, that may mediate IBD

susceptibility in humans.

Keywords: BXD genetic reference population (GRP), System genetics, Inflammatory bowel

disease (IBD), High-fat diet (HFD), Chow diet (CD), Weighted gene correlation network

analysis (WGCNA), Transcription factors (TFs), Module quantitative trait locus (ModQTL)

mapping

Introduction

A long-term lipid-rich diet is associated with multiple metabolic disorders, such as obesity (Hasegawa *et al.*, 2020), cardiovascular disease (Lutsey, Steffen and Stevens, 2008; Maurya *et al.*, 2023), and systemic low-grade inflammation (Duan *et al.*, 2018; Christ, Lauterbach and Latz, 2019). The gastro-intestinal tract is the primary site of adaptation to dietary challenge, due to its roles in nutrient absorption, immunity and metabolism (Enriquez *et al.*, 2022). Dietary challenges or other environmental or genetic factors can lead to prolonged inflammation and eventually damage the gastro-intestinal tract (Huang *et al.*, 2017; Enriquez *et al.*, 2022). IBD encompasses several chronic inflammatory gut disorders, including ulcerative colitis (UC) and Crohn's disease (Chang, 2020; Adolph *et al.*, 2022). Patients with IBD, have a higher risk of developing colorectal cancer (CRC), one of the most lethal cancers (Kim and Chang, 2014; Shah and Itzkowitz, 2022). The incidence of IBD has increased worldwide (Alatab *et al.*, 2020; Freeman *et al.*, 2021) during the last decade, in part due to increased consumption of lipid-rich diets (Maconi *et al.*, 2010; Hou, Abraham and El-Serag, 2011). Furthermore, mouse studies show that HFD leads to more inflammation in the dextran sulfate sodium (DSS)-induced UC models compared to normal diet (Zhao *et al.*, 2020). However, the response to HFD is variable across individuals (Zeevi *et al.*, 2015) and the association between the lipid-rich diet and the risk of IBD in clinical studies is inconclusive (Kreuter *et al.*, 2019), possibly due to genetic factors underlying inter-individual variability in gut inflammation and dysbiosis (Baumgart and Sandborn, 2012). More than 200 risk genes associated with IBD were identified through human genome-wide association studies (GWAS) (Huang *et al.*, 2017), which have implicated epithelial function, microbe sensing and restriction, and adaptive immune response as drivers (Graham and Xavier, 2020; Kong *et al.*, 2023). However, there is still no effective treatment for IBD. Current therapies, such as anti-tumor necrosis factor alpha (TNF-α) antibodies (Rutgeerts *et al.*, 2005) and integrin α4β7 antibodies,

blocking leukocyte migration (Feagan *et al.*, 2013), can temporarily alleviate inflammation in

a subset of patients (Rutgeerts *et al.*, 2005; Feagan *et al.*, 2013) but cause adverse effects

(Harbord *et al.*, 2017) and fail to prevent relapses (Doherty *et al.*, 2018). Therefore, it is

important to understand the gene-by-environment (GxE) interactions underpinning pre-clinical

gut inflammation that eventually evolves into IBD, to aid in designing novel preventive and

therapeutic strategies for intestinal inflammatory disorders.

Heterogeneity in clinical presentations as well as diversity in diet and lifestyle among human

IBD patients render human genetic studies challenging (Molodecky *et al.*, 2011). Experiments

in laboratory mice allow to control several environmental factors, such as temperature and diet,

when exploring the genetic modulators of IBD and also enable the collection of several relevant

tissues to help elucidate tissue-specific mechanisms (Nadeau and Auwerx, 2019; Li and

Auwerx, 2020). In addition, to mirror the heterogeneity of human populations, genetically

diverse populations, such as mouse genetic reference populations (GRPs), can be used in a

systems genetics paradigm (Nadeau and Auwerx, 2019; Li and Auwerx, 2020). This not only

allows the mapping of clinically relevant traits in controlled environments but also the

characterization of intermediate molecular phenotypes from tissues that cannot easily be

obtained in humans (Williams *et al.*, 2016; Li and Auwerx, 2020). For example, studies on the

molecular basis of non-alcoholic fatty liver disease in the Collaborative Cross founder strains

illustrated the importance of the genetic background in determining susceptibility to steatosis,

hepatic inflammation and fibrosis (Benegiamo *et al.*, 2023). Moreover, the BXD GRP was used

to identify genetic variants associated with metabolic phenotype variation, such as bile acid

homeostasis (Li *et al.*, 2022), lipid metabolism in plasma (Jha, McDevitt, Halilbasic, *et al.*,

2018) and liver (Jha, McDevitt, Gupta, *et al.*, 2018) as well as mitochondrial dysregulation

(Williams *et al.*, 2016), using GWAS or quantitative trait locus (QTL) mapping (Wu *et al.*,

2014; Williams *et al.*, 2016). Thus, large mouse GRPs are useful tools for identifying the tissue-

specific mechanisms of complex diseases.

In order to decipher the genetic and environmental contributions to the development of

intestinal inflammation, we measured the colon transcriptome of 52 BXD strains fed with CD

or HFD (Williams *et al.*, 2016). HFD feeding from 8 to 29 weeks of age induced an IBD-like

transcriptomic signature in colons of some, but not all, BXD strains, uncovering a subset of

BXD strains that could be susceptible to HFD-induced IBD-like state. Gene co-expression

analysis revealed two IBD-related modules in the colons of HFD-fed mice, one of which is

likely under the control of a ModQTL. Through a systems genetics prioritization of genes under

this ModQTL, we identified candidate IBD-related genes that we validated using GWAS in the

UK Biobank (UKBB) for human IBD.

Results

HFD feeding leads to highly variable transcriptomic adaptations in the colon of BXD strains

For this study, we used an extensively characterized BXD mouse panel of 52 BXD strains fed with a chow diet (CD) or high-fat diet (HFD) from 8 to 29 weeks of age (Williams *et al.*, 2016; Jha, McDevitt, Gupta, *et al.*, 2018; Jha, McDevitt, Halilbasic, *et al.*, 2018), in which we mapped genetic determinants of metabolic traits in the liver (Williams *et al.*, 2016; Jha, McDevitt, Gupta, *et al.*, 2018) and plasma (Jha, McDevitt, Halilbasic, *et al.*, 2018). These mice underwent metabolic phenotyping, with many metabolic traits being altered by HFD (**Figure 1A**), and multiple organs were harvested and flash-frozen for future use (Williams *et al.*, 2016). Here, we focused on proximal colon samples from this population and performed microarray-based transcriptome analysis of this tissue (**Figure 1A**).

Principal component analysis (PCA) of all transcriptomes (**Figure 1—figure supplement 1A**) showed that the first principal component (PC1) separated mice by diet, indicative of a global diet effect in the population. Nevertheless, transcriptomes of several strains (such as BXD12, BXD84 and BXD81) on HFD had very similar PC1 values to their CD counterparts (**Figure 1—figure supplement 1B**), suggesting that they were resistant to dietary changes. Similarly, BXD strains did not cluster completely by diet based on hierarchical clustering analysis indicating that the genetic differences can override the impact of diet on the transcriptome in the colon (**Figure 1—figure supplement 1C**). To obtain a global, strain-independent, view of the HFD effect, we performed a differential expression analysis and identified 115 up- and 295 down-regulated differentially expressed genes (DEGs, absolute Log$_2$(Fold Change) > 0.5 and Benjamini-Hochberg (BH)-adjusted P value < 0.05, **Figure 1B**). Of note, *Cldn4*, one of claudins implicated in intestinal permeability (Ahmad *et al.*, 2017), was significantly down-regulated and serum amyloid A (*Saa1* and *Saa3*), which have been involved in the inflammatory response (Ye and Sun, 2015; Tannock *et al.*, 2018), were up-regulated upon HFD

(**Figure 1B**). Furthermore, gene set enrichment analysis (GSEA) showed an upregulation of

inflammation, cell proliferation and translation, mitochondrial respiration, and stress response-

related pathways upon HFD, while genes involved in the intermediate filament - that contribute

to maintaining intestinal barriers (Misiorek *et al.*, 2016; Mun, Hur and Ku, 2022) - were down-

regulated (**Figure 1C**). All in all, the transcriptome data are consistent with an HFD-induced

downregulation of components of the intestinal barrier, enhanced permeability, induction of

the unfolded protein response (UPR) and increased inflammation in BXD colons, much like

HFD does in humans (Bischoff *et al.*, 2014). However, as in humans, not every strain exhibited

the same response to dietary challenges. GSEA analyses applied individually to the diet effect

in each strain showed a high degree of diversity in the inflammatory response (**Figure 1—**

figure supplement 1D). For example, BXD44, 45 and 55, highlighted in red, were the 3 most

susceptible strains to gut inflammation upon HFD, whereas BXD1, 67, and 85, colored in green,

showed no significant enrichment in gut inflammation. This diversity in responses provided

the basis for a systems genetics investigation of HFD-driven gut inflammation determinants in

the BXD.

The transcriptomic response to HFD of a subset of BXD strains resembles DSS-induced

ulcerative colitis (UC)

IBD is characterized by increasing inflammation in the gastro-intestinal tract (Adolph *et al.*,

2022). To investigate the disease relevance of the chronic inflammation seen in BXD colons

upon HFD, we extracted the transcriptomic signatures from DSS-induced mouse UC models

(Czarnewski *et al.*, 2019) and two IBD human studies (GSE16879 (Arijs *et al.*, 2009) and

GSE83687 (Peters *et al.*, 2017), **Materials and Methods**) and used these signatures as custom

gene sets in GSEA on the global HFD effect. DSS is widely used to induce UC in mouse

models and disease severity increases over time (Czarnewski *et al.*, 2019). GSEA analyses

showed that DSS-induced genes from days 4 (early inflammatory phase), 6 and 7 (acute

inflammatory phase) were significantly enriched in genes upregulated by HFD, especially the

dysregulated genes in the later stage of DSS-induced UC (**Figure 1D, bottom panel**). Similarly,

genes involved in human IBD (UC and Crohn's disease (CDs)) were also enriched in those

same genes (**Figure 1D, top panel**). The same trend was observed for downregulated genes in

mouse and human IBD, which were negatively enriched (**Figure 1D**), illustrating that HFD

induced an IBD-like transcriptomic signature in BXD colons.

While the average response across all BXDs shared features of mouse and human IBD, we

assessed the strain-specificity of this response by measuring each strain's response to IBD

using GSEA (**Figure 2A**). Hierarchical clustering of the normalized enrichment scores (NES)

in mouse IBD datasets classified the BXDs into three groups: susceptible strains highlighted

in red (19 strains), intermediate strains represented in blue (11 strains), and resistant strains

colored in green (17 strains) (**Figure 2A, top panel**). Of note, in line with colon histological

lesions comparison of DSS-induced colitis mouse models in the literature (Mähler *et al.*, 1998),

the C57BL/6J strain, one of the parental strains of the BXDs, was classified as one of the

susceptible strains while the other parental strain DBA/2J belonged to the resistant group

(**Figure 2A, top panel**), suggesting that genetic determinants inherited from the parental strains

may determine the susceptibility of BXD strains to HFD-induced IBD-like inflammation in the

colons.

To establish the functional relevance of this transcriptome-based classification on systemic

inflammation, we compared plasma cytokine levels of these three groups under HFD (Williams

et al., 2016). Interestingly, the susceptible group have significantly lower levels of the anti-

inflammatory cytokine - Interleukin (IL)-10 (**Figure 2B,** two-tailed t-test $p < 0.01$) and

increased the proinflammatory cytokine - IL-15 (**Figure 2C,** two-tailed t-test $p < 0.0001$)

compared to the resistant strains. *IL10* itself has been identified as an IBD-related candidate

gene using GWAS in humans (Franke *et al.*, 2008) and IL-10-deficient mice are also well-

known mouse model for IBD research (Keubler *et al.*, 2015). IL-15 is another important

cytokine involved in intestinal inflammation and is elevated in the human guts with IBD (Liu

et al., 2000). IL-15 knock-out mice are also reported to have less severe symptoms, such as

weight loss and histological scores, following DSS administration (Yoshihara *et al.*, 2006). In

summary, susceptibility to HFD-induced IBD-like inflammation in the colon, as assessed by

changes in levels of genes associated with IBD, correlates with markers of the general

inflammatory status of mice.

Identifying IBD-related gene modules in BXD colons

Since different BXD strains seem to exhibit different susceptibility to IBD, we set out to

explore gene expression signatures underlying these differences. For that, we used Weighted

Gene Co-expression Analysis (WGCNA) to construct CD- and HFD-specific gene co-

expression networks to identify modules of co-expressed genes (**Figure 3A, Appendix 1 -**

Table 1). Disease-associated modules were then defined as modules under HFD are

significantly enriched in mouse DSS-induced UC signatures by an over-representation analysis

(ORA, BH-adjusted P value < 0.05 and number of enriched genes > 5, **Figure 3A**). The HFD

co-expression network consisted of 39 modules ranging in size from 34 to 1,853 genes and

containing a total of 14,723 genes (**Appendix 1 - Table 1**). We visualized this network using

Uniform Manifold Approximation and Projection (UMAP) (**Figure 3B**), reflecting that the

majority modules were closely connected in the co-expression network.

Enrichment analyses indicated that modules HFD_M9, HFD_M16, and HFD_M28 were

enriched with genes that are upregulated by DSS-induced colitis, while HFD_M15, HFD_M24,

and HFD_M26 were significantly enriched with downregulated genes (**Figure 3C).** Of note,

more than 20% of genes involved in HFD_M9 and HFD_M28 were part of the dysregulated

genes of the acute phase of mouse UC (day6 and day7) (**Figure 3C**). Interestingly, genes

perturbed during IBD pathogenesis in humans were also enriched in HFD_M9 and HFD_M28

(**Figure 3C**).

While IBD-related genes were predominantly found in HFD modules, we also found that two

modules, CD_M28 and CD_M32, in CD-fed mouse colons were associated with IBD (**Figure**

3—figure supplement 1A). These two-modules significantly overlapped with the IBD-related

HFD_M9 and HFD_M28 modules, respectively (BH-adjusted P value < 0.05) (**Figure 3—**

figure supplement 1B). Moreover, the molecular signatures underlying human UC and

Crohn's disease were also clustered in these two modules (CD_M28 and CD_M32) under CD

(**Figure 3—figure supplement 1C**). Collectively, the co-expression and enrichment analyses

identify HFD_M9 and HFD_M28 as IBD-related modules on which we focus our subsequent

investigation.

Identifying biological roles and transcriptional regulation of the IBD-related modules

To identify the biological function of the IBD-related modules, we performed enrichment

analyses using the Hallmark database and the cell-type gene signatures (Kong *et al.*, 2023)

(**Materials and Methods**). Genes in HFD_M9 were enriched in KRAS signaling and

inflammation-related pathways, while HFD_M28 was enriched in IFN-α/γ responses (BH-

adjusted P value < 0.05) (**Figure 4A**). Both modules were enriched in IFN-γ response genes

(**Figure 4A**). IFN-γ is an essential cytokine for innate and adaptive intestinal immune responses

(Brasseit *et al.*, 2018). It has been reported to play a key role in mouse (Ito *et al.*, 2006) and

human (Tilg *et al.*, 2002) IBD pathogenesis, and was identified as a potential therapeutic target

to alleviate inflammatory response in IBD (Li *et al.*, 2021). In addition, genes that are

dysregulated in immune cells of Crohn's disease patients (Macrophages, B cell and immune

cycling cells) were enriched in HFD_M9 (**Figure 4B**). In contrast, genes of HFD_M28 were

not only enriched for genes that are dysregulated in immune cells, but also in intestinal

epithelial cells of diseased individuals, such as Goblet and stem cells (**Figure 4B**). Overall,

HFD_M9 and HFD_M28 are both involved in inflammatory response, while genes involved

in HFD_M28 also potentially influence intestinal epithelial barrier.

To identify transcriptional drivers of the two IBD-related modules, we performed a

transcription factor (TF) enrichment analysis (**Materials and Methods**) and found that ZIC2,

SMAD3, REL, FOSL1, and BATF are the top enriched transcription factors for the genes in

HFD_M9 (**Figure 4C**), while the expression of genes in module HFD_M28 may be regulated

by Interferon regulatory factors (IRFs, IRF1, IRF2, IRF7, and IRF9) and the signal transducer

and activator of transcription families (STAT, STAT2) (**Figure 4D**). In fact, most of these TFs

have been reported to be involved in gut inflammation. For example, *Smad3* mutant mice were

more susceptible to intestinal inflammation (Yang *et al.*, 1999). Moreover, the IFN-STAT axis

is essential to initiate the type-I IFN induction that is critical for human immune defense, such

as IBD diseases (Stolzer *et al.*, 2021) and primary immunodeficiency diseases (Mogensen,

2019) as well as for disease tolerance (Mottis *et al.*, 2022). Collectively, we have identified

TFs that likely control the expression of the two IBD-related modules to play an essential role

in gut inflammation regulation.

Identifying ModQTLs for IBD-related modules and filtering of candidate genes

To analyze how the genotype impacts the IBD-like inflammatory response associated to HFD,

we performed module QTL mapping analysis (ModQTL) for both IBD-related modules

(HFD_M9 and HFD_28) (**Figure 5A**). We found a suggestive QTL for HFD_M28 (P value <

0.1), on chromosome 16 containing 552 protein-coding genes (**Figure 5A, Appendix 1 - Table

2**). We annotated these candidate genes based on three criteria (**Figure 5B**): (1) presence of

high-impact genetic variants (such as missense and frameshift variants) in BXDs, (2)

association with inflammation based on literature mining (**Materials and methods**), (3)

presence of *cis*-expression QTLs (eQTLs), that is, whether the expression of the gene is

controlled by the QTL. The 27 genes satisfying at least two of the above criteria were

considered as candidate genes driving the expression of module HFD_M28 (**Figure 5C**).

To further prioritize candidate genes regulating module HFD_M28, we applied GWAS to

detect Crohn's disease- and UC-associated genetic variants using whole genome sequence

(WGS) dataset in UKBB (**Figure 5C, Materials and Methods**). Interestingly, the genetic

variants of two genes under the QTL peak, i.e, *ephrin type A receptor 6* (*EPHA6,* P value =

2.3E-06) (**Figure 5C**, **Figure 5—figure supplement 1A**) and *Mucin 4* (*MUC4,* P value = 1.2E-

06) (**Figure 5C**, **Figure 5—figure supplement 1B)** were also associated with UC in humans.

EPHA6 belongs to Eph/Ephrin Signaling and this pathway has been associated with gut

inflammation (Coulthard *et al.*, 2012) and proposed as a potential target to alleviate the

inflammatory response in IBD (Grandi *et al.*, 2019), but the association between *EPHA6* and

IBD is not explored yet. The Gene-Module Association Determination (G-MAD) (Li *et al.*,

2019) (https://systems-genetics.org/gmad) also revealed that expression of *Epha6* in mouse

gastro-intestinal tract correlates with genes involved in inflammation-related pathways, such

as IL-6 production and regulation of inflammatory response (**Figure 5D, Appendix 1 - Table**

3). *MUC4* is a transmembrane mucin (Gao *et al.*, 2021) and highly expressed in gastro-

intestinal   tract   according   to   the   human   protein   atlas   (Uhlén   *et   al.*,   2015)

(https://www.proteinatlas.org/humanproteome/tissue/intestine) (**Figure 5C**). The expression

of *MUC4* in the human gastro-intestinal tract correlates with genes that are enriched for CRC

and O-linked glycosylation based on G-MAD (Li *et al.*, 2019) (**Figure 5E, Appendix 1 - Table**

3). O-linked glycans are expressed by the intestinal epithelium to maintain barrier function,

especially mucin type O-glycans, and gut disorders can be affected by dysfunction of O-linked

glycosylation (Brazil and Parkos, 2022). Moreover, *MUC4* is upregulated in enterocytes and

Goblet cells in colons of Crohn's disease patients (**Figure 5F)**. *MUC4* hence is a strong

candidate because of its role in maintaining the intestinal epithelium and controlling the gut

inflammatory response (McGuckin *et al.*, 2011) and *EPHA6* might be a novel candidate gene

to impact gut inflammation. Based on the results of our QTL mapping, human GWAS in

UKBB, and existing literature, we hypothesize that *MUC4* and *EPHA6* impact on colon

integrity and inflammation and may be important players in gut inflammation or IBD triggered

by an unhealthy, lipid-rich diet.

Discussion

Dietary, environmental and genetic factors have all been reported to influence intestinal
inflammation (Adolph *et al.*, 2022). Indeed, HFD can impair the intestinal epithelial barrier
and trigger pre-clinical inflammation in the gastro-intestinal tract, eventually leading to
inflammatory disorders of the gut (Enriquez *et al.*, 2022). In addition, genetic factors identified
by GWAS can also predispose to IBD. For example, the interleukin-1 and -7 receptors (*IL-1R2*
and *IL-7R*) were identified as candidate genes that regulate the immune response in IBD (Khor,
Gardet and Xavier, 2011). However, the heterogeneity of diet and other environmental factors
in human studies limits our ability to identify GxE interactions and pinpoint the genes and
pathways involved in diet-induced gut inflammation. Studies in model organisms such as the
mouse, where the environment can be carefully controlled, provide a valuable complement to
human genetics studies that by nature are mainly observational (Nadeau and Auwerx, 2019; Li
and Auwerx, 2020). Unfortunately, most mouse studies only evaluate mice from a single
genetic background, limiting their generalizability and translatability to humans (Nadeau and
Auwerx, 2019; Li and Auwerx, 2020). Conversely, GRPs such as the BXDs can mimic at least
in part the heterogeneity of human populations and allow us to estimate the effect of GxE
interactions on complex diseases (Jha, McDevitt, Gupta, *et al.*, 2018; Li and Auwerx, 2020).

Here, we utilized a panel of 52 BXD genetically diverse mouse strains fed with either HFD or
CD to explore the genetic and dietary modulators of inflammation seen in the colon
transcriptomes using systems genetics approaches. The colon transcriptomic response to HFD
in this mouse population recapitulated several of the general features observed in DSS-induced
UC mouse models and human IBD patients. In particular, we identified the upregulation of
inflammation-related genes and the UPR as well as the downregulation of intercellular
adhesion-related genesets as common signatures induced by HFD (Kreuter *et al.*, 2019).
Moreover, our dataset not only was informative about the transcript changes of IBD at the

population level, but also unveiled extensive strain-specific effects that allowed us to classify strains based on their propensity to develop IBD-like signatures. The fact that these susceptibility groups also differed in anti- and pro-inflammatory plasma cytokine levels (IL-10 and IL-15, respectively) suggests a relation between these tissue-specific transcriptional signatures and systemic low-grade inflammation. Since gene interactions determine cellular processes and the molecular functions of correlated genes are often similar (Nayak *et al.*, 2009), we attempted to elucidate the mechanism underlying the diversity of IBD-like signatures and chronic inflammation in BXD colons using gene co-expression analyses. This led us to identify two IBD-related gene modules (HFD_M9 and HFD_M28).

As most differentially expressed genes are likely to be driven by and not be a cause of disease (Porcu *et al.*, 2021), we attempted to understand whether the signatures in the colon are causes or consequences of chronic inflammation. A first step was to characterize possible transcriptional and genetic regulators of IBD-related modules. Enrichment analyses showed that both IBD-associated modules largely consisted of immune response-related genes. Specifically, genes involved in HFD_M9 and HFD_M28 are both differentially expressed in immune cells in inflamed tissues of Crohn's disease patients (Kong *et al.*, 2023). Moreover, the HFD_M28 module was enriched for TF motifs of STAT2 and IRF family, and HFD_M9 for SMAD3 and REL, which were illustrated to control the expression of these gut inflammation-related genes, and influence the inflammatory response triggered by HFD in the colon.

While we found IBD-related gene modules and the TFs driving their expression, the genetic drivers of the diversity of gut inflammatory responses observed across the BXDs remained elusive. To find candidate genes causing gut inflammation upon HFD, we then performed Module QTL (ModQTL) analysis and allocated a suggestive ModQTL that may be controlling one of IBD-related module (HFD_M28) under HFD. Importantly, through our prioritization

scheme for the genes under the ModQTL, we identify two plausible candidates, *Epha6* and *Muc4*, that have high-impact variants in the BXDs, are related to inflammation, and harbor variants in humans that are associated with IBD based on UKBB GWAS result. In fact, *Muc4* knock-out mice have been shown to be more resistant to DSS-induced UC through upregulating the expression of *Muc2* (mucin secretion) and *Muc3* (transmembrane mucin) (Das *et al.*, 2016) . A GWAS study also indicated that mutations in *EPHA6* increase risk for CRC (Guda *et al.*, 2015), but its potential association with IBD is a new finding. Therefore, these results point to important potential roles of *Muc4* and *Epha6* in gut chronic inflammation leading to inflammatory gut disorders.

Although studies in the BXD cohort are limited to variants present in the parental strains, C57BL/6J and DBA/2J, our analysis nevertheless shows how genetic diversity in this population allows us to detect the genetic modulators of chronic intestinal inflammation, that are more difficult to identify in widely used IBD mouse models on a single genetic background. In support of the generalizability of our data, the identified candidate genes in our mouse models were also associated to human UC, demonstrating that chronic inflammation induced upon HFD feeding may indeed be a prelude to human UC.

In conclusion, our systems genetics investigation of the colon in a controlled GRP, complemented with human GWAS studies, enabled the prioritization of modulators of IBD susceptibility that were generalizable to the human situation and may have clinical value.

Materials and methods

Population handling

Mice were studied as previously described (Williams *et al.*, 2016) and multiple organs were harvested for further analysis. Briefly, in groups of 3-5 animals from the same strain and diet, in isolator cages with individual air filtration (500 cm2, GM500, Tecniplast) and provided water ad libitum. Mice were fed CD ad libitum until 8 weeks of age. From 8 weeks to 29 weeks, half of the cohort was fed ad libitum HFD and the rest continued to be fed a CD (**Figure 1A**). CD composition: 18% kCal fat, 24% kCal protein and 58% kCal of carbohydrates (Teklad Global 18% Protein Rodent Diet 2018 chow diet, Envigo, Indianapolis, USA). HFD composition: 60.3% kCal fat, 18.4% kCal protein and 27.3% kCal of carbohydrates (Teklad Custom Diet TD.06414, Envigo, Indianapolis, USA). All mice were fasted overnight (from 6pm to 9am) prior to euthanasia. All procedures were approved by the veterinary office of canton Vaud under animal experimentation license number VD2257. In this work, proximal colons were extracted from the bio-banked samples and we did not use any new animals.

Transcriptome of the proximal colon in BXDs

A ~1 cm portion of the proximal half part of the colon was excised following euthanasia, washed in PBS and immediately stored in liquid nitrogen. Approximately 5 animals of the same strain fed the same diet were pooled at equal mass concentration for further RNA extraction. Total RNA was extracted using Direct-zol (Zymo Research) including the DNase digestion step. 100ng of total RNA was amplified using the Ambion® WT Expression Kit from Life Technologies (part number 4411974) and 5,500ng of cDNA was fragmented and labeled using the Affymetrix WT terminal labeling kit (part number 900671) all following manufacturers protocols. Labeled cDNA was hybridized on an Affymetrix Clariom S Assay microarray platform (GPL23038) in ~16 hours of incubation, then washed and stained using

an Affymetrix 450 Fluidics Station according to Affymetrix protocols. Finally, arrays were

scanned on Affymetrix GSC3000 7G Scanner. Microarray data preprocessing was performed

using apt-probeset-summarize from the Array Power Tool (APT) suite (v2.11.3) with the gc-

sst-rma-sketch standard method and resulting expression values were log-transformed.

Microarray probes targeting polymorphic regions in the BXD population were ignored in the

process. For probesets targeting a same transcript, only the probeset with the highest value was

considered.

Differential gene expression analysis

General differences in mRNA expression profiles between diets was assessed using Principal

Component Analysis (PCA). Differential expression of individual transcripts between diets

was assessed using the limma R Bioconductor package (version 3.48.3) (Ritchie *et al.*, 2015).

Briefly, statistical significance was assessed using an empirical Bayes method (eBayes function)

with an additive linear model accounting for diet and strain effect and adjusted P values were

calculated by the Benjamini-Hochberg (BH) approach. Transcripts showing BH-adjusted P

value below 0.05 and absolute Log_2 (Fold Change) above 0.5 were considered significantly

associated with the effect of the diet.

Gene set enrichment analysis (GSEA) and Over-representation analysis (ORA)

Gene sets used in GSEA and ORA consisted of two parts: (1) the gene sets from the GO, KEGG,

Hallmark, and Reactome databases were retrieved through the msigdbr R package (version

7.2.1) (Liberzon *et al.*, 2011). (2) the gene signatures of mouse and human IBD were used as

custom gene sets (**Table 1**).

GSEA was performed using clusterProfiler R package (version 3.10.1) (Yu *et al.*, 2012) based

on the log_2(Fold Change) ranking using parameters (*nPerm* = 100000, *minGSSize* = 30,

maxGSSize = 5000, *pvalueCutoff* = 1). The gene sets with absolute NES higher than 1 and BH-

adjusted P value lower than 0.05 were identified as the significantly enriched gene sets.

ORA analysis was also performed using clusterProfiler R package (version 3.10.1) (Yu *et al.*,

2012) using parameters (minGSSize = 30, maxGSSize = 800). The gene sets with adjusted P

value calculated by BH lower than 0.05 were identified as the significantly enriched gene sets.

Weighted gene correlation network analysis (WGCNA)

We used *WGCNA* R package (v1.51) (Langfelder and Horvath, 2008) to construct co-

expression networks under CD and HFD, respectively. Firstly, the correlations between all

pairs of gene across all BXDs fed with CD or HFD were calculated by Pearson correlation.

Then, a best soft-thresholding power of 4 and 3 was chosen using pickSoftThreshold function

with parameters (*networkType* = "signed hybrid", *blockSize* = 25000, *corFnc* = "bicor") for CD

and HFD datasets in BXD colons separately. According to the calculated correlation

coefficients, a network was constructed using parameters (*networkType* = "signed hybrid",

minModuleSize = 30, *reassignThreshold* = 1e-6, *mergeCutHeight* = 0.15, *maxBlockSize* =

25000). The constructed co-expression gene modules were assigned color names and the

module eigengenes were also identified for further analyses. To detect the preserved CD-

modules in the co-expression modules under HFD, we defined gene modules under CD as

custom genesets and performed ORA on each HFD-modules.

Transcription factor (TF) enrichment analysis

We first constructed a lognormal background distribution using the sequences of + 5kb region

around the transcription starting site (TSS) of all genes and then downloaded the mouse

HOCOMOCO-v10 (Kulakovskiy *et al.*, 2018) motifs from R package motifDB to perform TF

enrichment analyses using R package *PWMenrich*. The significantly enriched motifs (P value

< 0.001) were selected and then ranked based on the percentage of enriched promoters.

Module Quantitative Trait Locus (ModQTL) mapping in the BXDs

We first downloaded genotype information of each BXD mice from GeneNetwork

(https://gn1.genenetwork.org/webqtl/main.py?FormID=sharinginfo&GN_AccessionId=600)

and generated the kinship matrix of BXD mice using the leave-one-chromosome-out (LOCO)

method. We then used the eigengenes of each module as phenotype input to perform Module

QTL (ModQTL) with the R package *qtl2* (version 0.28) (Broman *et al.*, 2019) and the threshold

of each QTL mapping analysis was obtained from a permutation test with 10,000 repeats. The

peaks of QTL were calculated by *find_peaks* function with parameter: *prob=0.95.*

The same methods were also applied to gene expression QTL mapping (eQTL) and the

significance threshold of each gene was obtained from a permutation test with 1,000 repeats.

The significant peaks overlapped with the location of their corresponding gene were identified

as *cis*-eQTL.

Literature mining

To explore the inflammation related genes, we first used candidate gene names and keywords

("IBD", "inflammatory bowel disease", "Ulcerative colitis", "Inflammation", "Inflammatory",

"Crohn's disease") to search the title or abstract of associated literature using R package

easyPubMed (version 2.13). Then, the genes involved in inflammation were confirmed by

manual curation.

Genome-wide association study (GWAS) in UKBB

The phenotype data of inflamed Ulcerative colitis (Data-Field 131629, n = 6,459) and Crohn's

disease (Data-Field 131627, n = 3,358) were firstly downloaded from UKBB (Bycroft *et al.*,

2018). 200,030 individuals with whole genome sequencing (WGS) (Halldorsson *et al.*, 2022)

in UK Biobank were selected and then the population of European descent (including with

1,173 patients with Crohn's disease and 2,295 patients with UC) was extracted for further

GWAS analyses. Control individuals (n = 143,194) were included based on the following

criteria: (1) Individuals without non- inflamed colitis (Data-Field 131631), Crohn's disease,

and UC. (2) Individuals not taking any IBD-related medicine (**Appendix 1 - Table 4**).

WGS data provided by UK Biobank and used for GWAS were processed starting from pVCF

files. We used REGENIE step1 to estimate population structure and then REGENIE step2 were

applied to test associations between phenotypes and genetic variants and also included the

following covariates in our model: the first 10 genetic principal components, age, sex, age:sex

interaction, Body Mass Index (BMI), and smoking status. All data preparation and GWAS

steps were run on DNAnexus.

Acknowledgments

We thank the Schoonjans' and Auwerx's lab members for technical assistance and discussions and Giacomo von Alvensleben for providing the GWAS analysis pipeline in human UKBB. The work in the JA laboratory was supported by grants from the Ecole Polytechnique Fédérale de Lausanne (EPFL), the European Research Council (ERC-AdG-787702), the Swiss National Science Foundation (SNSF 31003A_179435) and the Global Research Laboratory (GRL) National Research Foundation of Korea (NRF 2017K1A1A2013124). XL was supported by the China Scholarship Council (201906050019).

Author contributions

The study was conceived by XL, MBS and JA. EW, MBS and AB performed laboratory experiments. Data analyses were carried out by XL, AB, AR and JP. XL and JA wrote the original manuscript. XL, MBS, JDM, GB, AP, KS and JA reviewed and edited the manuscript with contributions from all co-authors.

Competing interests

Authors declare no conflict of interest related to the work reported.

Data availability

The data that support the findings are available upon request to the corresponding authors (MBS and JA). The microarray data are available under the GEO numbers GSE225791. To review this dataset, please use this link: https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE225791 and the review token is

exwtuucwpdqlzsr. Methods, materials and external resources are included in the **Materials**

and Methods.

References

Adolph, T.E. *et al.* (2022) 'The metabolic nature of inflammatory bowel diseases', *Nature*
Reviews Gastroenterology & Hepatology, 19(12), pp. 753–767. Available at:
https://doi.org/10.1038/s41575-022-00658-y.

Ahmad, R. *et al.* (2017) 'Obesity-induces Organ and Tissue Specific Tight Junction
Restructuring and Barrier Deregulation by Claudin Switching', *Scientific Reports*, 7(1), p. 5125.
Available at: https://doi.org/10.1038/s41598-017-04989-8.

Alatab, S. *et al.* (2020) 'The global, regional, and national burden of inflammatory bowel
disease in 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden
of Disease Study 2017', *The Lancet Gastroenterology & Hepatology*, 5(1), pp. 17–30.
Available at: https://doi.org/10.1016/S2468-1253(19)30333-4.

Arijs, I. *et al.* (2009) 'Mucosal Gene Expression of Antimicrobial Peptides in Inflammatory
Bowel Disease Before and After First Infliximab Treatment', *PLOS ONE*, 4(11), p. e7984.
Available at: https://doi.org/10.1371/journal.pone.0007984.

Baumgart, D.C. and Sandborn, W.J. (2012) 'Crohn's disease', *The Lancet*, 380(9853), pp.
1590–1605. Available at: https://doi.org/10.1016/S0140-6736(12)60026-9.

Benegiamo, G. *et al.* (2023) 'The genetic background shapes the susceptibility to mitochondrial
dysfunction and NASH progression', *Journal of Experimental Medicine*, 220(4), p. e20221738.
Available at: https://doi.org/10.1084/jem.20221738.

Bischoff, S.C. *et al.* (2014) 'Intestinal permeability – a new target for disease prevention and
therapy', *BMC Gastroenterology*, 14(1), p. 189. Available at: https://doi.org/10.1186/s12876-
014-0189-7.

Brasseit, J. *et al.* (2018) 'Divergent Roles of Interferon-γ and Innate Lymphoid Cells in Innate
and Adaptive Immune Cell-Mediated Intestinal Inflammation', *Frontiers in Immunology*, 9.
Available at: https://www.frontiersin.org/articles/10.3389/fimmu.2018.00023 (Accessed: 24
February 2023).

Brazil, J.C. and Parkos, C.A. (2022) 'Finding the sweet spot: glycosylation mediated regulation
of intestinal inflammation', *Mucosal Immunology*, 15(2), pp. 211–222. Available at:
https://doi.org/10.1038/s41385-021-00466-8.

Broman, K.W. *et al.* (2019) 'R/qtl2: Software for Mapping Quantitative Trait Loci with High-
Dimensional Data and Multiparent Populations', *Genetics*, 211(2), pp. 495–502. Available at:
https://doi.org/10.1534/genetics.118.301595.

Bycroft, C. *et al.* (2018) 'The UK Biobank resource with deep phenotyping and genomic data',
Nature, 562(7726), pp. 203–209. Available at: https://doi.org/10.1038/s41586-018-0579-z.

Chang, J.T. (2020) 'Pathophysiology of Inflammatory Bowel Diseases', *New England Journal*
of Medicine, 383(27), pp. 2652–2664. Available at: https://doi.org/10.1056/NEJMra2002697.

Christ, A., Lauterbach, M. and Latz, E. (2019) 'Western Diet and the Immune System: An
Inflammatory Connection', *Immunity*, 51(5), pp. 794–811. Available at:
https://doi.org/10.1016/j.immuni.2019.09.020.

Coulthard, M.G. *et al.* (2012) 'Eph/Ephrin Signaling in Injury and Inflammation', *The*
American Journal of Pathology, 181(5), pp. 1493–1503. Available at:
https://doi.org/10.1016/j.ajpath.2012.06.043.

Czarnewski, P. *et al.* (2019) 'Conserved transcriptomic profile between mouse and human
colitis allows unsupervised patient stratification', *Nature Communications*, 10(1), p. 2892.
Available at: https://doi.org/10.1038/s41467-019-10769-x.

Das, S. *et al.* (2016) 'Mice deficient in Muc4 are resistant to experimental colitis and colitis-
associated colorectal cancer', *Oncogene*, 35(20), pp. 2645–2654. Available at:
https://doi.org/10.1038/onc.2015.327.

Doherty, G. *et al.* (2018) 'European Crohn's and Colitis Organisation Topical Review on
Treatment Withdrawal ["Exit Strategies"] in Inflammatory Bowel Disease', *Journal of*
Crohn's and Colitis, 12(1), pp. 17–31. Available at: https://doi.org/10.1093/ecco-jcc/jjx101.

Duan, Y. *et al.* (2018) 'Inflammatory Links Between High Fat Diets and Diseases', *Frontiers*
in Immunology, 9. Available at:
https://www.frontiersin.org/articles/10.3389/fimmu.2018.02649 (Accessed: 19 January 2023).

Enriquez, J.R. *et al.* (2022) 'A dietary change to a high-fat diet initiates a rapid adaptation of
the intestine', *Cell Reports*, 41(7). Available at: https://doi.org/10.1016/j.celrep.2022.111641.

Feagan, B.G. *et al.* (2013) 'Vedolizumab as Induction and Maintenance Therapy for Ulcerative
Colitis', *New England Journal of Medicine*, 369(8), pp. 699–710. Available at:
https://doi.org/10.1056/NEJMoa1215734.

Franke, A. *et al.* (2008) 'Sequence variants in IL10, ARPC2 and multiple other loci contribute
to ulcerative colitis susceptibility', *Nature Genetics*, 40(11), pp. 1319–1323. Available at:
https://doi.org/10.1038/ng.221.

Freeman, K. *et al.* (2021) 'The incidence and prevalence of inflammatory bowel disease in UK
primary care: a retrospective cohort study of the IQVIA Medical Research Database', *BMC*
Gastroenterology, 21(1), p. 139. Available at: https://doi.org/10.1186/s12876-021-01716-6.

Fujiyama, Y. *et al.* (2007) 'Butter feeding enhances TNF-α production from macrophages and
lymphocyte adherence in murine small intestinal microvessels', *Journal of Gastroenterology*
and Hepatology, 22(11), pp. 1838–1845. Available at: https://doi.org/10.1111/j.1440-
1746.2007.04905.x.

Gao, X.-P. *et al.* (2021) 'Integrative Analysis of MUC4 to Prognosis and Immune Infiltration
in Pan-Cancer: Friend or Foe?', *Frontiers in Cell and Developmental Biology*, 9. Available at:
https://www.frontiersin.org/articles/10.3389/fcell.2021.695544 (Accessed: 10 February 2023).

Graham, D.B. and Xavier, R.J. (2020) 'Pathway paradigms revealed from the genetics of
inflammatory bowel disease', *Nature*, 578(7796), pp. 527–539. Available at:
https://doi.org/10.1038/s41586-020-2025-2.

Grandi, A. *et al.* (2019) 'Targeting the Eph/Ephrin System as Anti-Inflammatory Strategy in
IBD', *Frontiers in Pharmacology*, 10, p. 691. Available at:
https://doi.org/10.3389/fphar.2019.00691.

Guda, K. *et al.* (2015) 'Novel recurrently mutated genes in African American colon cancers',
Proceedings of the National Academy of Sciences, 112(4), pp. 1149–1154. Available at:
https://doi.org/10.1073/pnas.1417064112.

Halldorsson, B.V. *et al.* (2022) 'The sequences of 150,119 genomes in the UK Biobank',
Nature, 607(7920), pp. 732–740. Available at: https://doi.org/10.1038/s41586-022-04965-x.

Harbord, M. *et al.* (2017) 'Third European Evidence-based Consensus on Diagnosis and
Management of Ulcerative Colitis. Part 2: Current Management', *Journal of Crohn's and*
Colitis, 11(7), pp. 769–784. Available at: https://doi.org/10.1093/ecco-jcc/jjx009.

Hasegawa, Y. *et al.* (2020) 'Long-term effects of western diet consumption in male and female
mice', *Scientific Reports*, 10(1), p. 14686. Available at: https://doi.org/10.1038/s41598-020-
71592-9.

Hou, J.K., Abraham, B. and El-Serag, H. (2011) 'Dietary Intake and Risk of Developing
Inflammatory Bowel Disease: A Systematic Review of the Literature', *Official journal of the*
American College of Gastroenterology | ACG, 106(4), p. 563. Available at:
https://doi.org/10.1038/ajg.2011.44.

Huang, H. *et al.* (2017) 'Fine-mapping inflammatory bowel disease loci to single-variant
resolution', *Nature*, 547(7662), pp. 173–178. Available at:
https://doi.org/10.1038/nature22969.

Ito, R. *et al.* (2006) 'Interferon-gamma is causatively involved in experimental inflammatory
bowel disease in mice', *Clinical and Experimental Immunology*, 146(2), pp. 330–338.
Available at: https://doi.org/10.1111/j.1365-2249.2006.03214.x.

Jha, P., McDevitt, M.T., Halilbasic, E., *et al.* (2018) 'Genetic Regulation of Plasma Lipid
Species and Their Association with Metabolic Phenotypes', *Cell Systems*, 6(6), pp. 709-721.e6.
Available at: https://doi.org/10.1016/j.cels.2018.05.009.

Jha, P., McDevitt, M.T., Gupta, R., *et al.* (2018) 'Systems Analyses Reveal Physiological Roles
and Genetic Regulators of Liver Lipid Species', *Cell Systems*, 6(6), pp. 722-733.e6. Available
at: https://doi.org/10.1016/j.cels.2018.05.016.

Keubler, L.M. *et al.* (2015) 'A Multihit Model: Colitis Lessons from the Interleukin-10–
deficient Mouse', *Inflammatory Bowel Diseases*, 21(8), pp. 1967–1975. Available at:
https://doi.org/10.1097/MIB.0000000000000468.

Khor, B., Gardet, A. and Xavier, R.J. (2011) 'Genetics and pathogenesis of inflammatory
bowel disease', *Nature*, 474(7351), pp. 307–317. Available at:
https://doi.org/10.1038/nature10209.

Kim, E.R. and Chang, D.K. (2014) 'Colorectal cancer in inflammatory bowel disease: The risk, pathogenesis, prevention and diagnosis', *World Journal of Gastroenterology*, 20(29), pp. 9872–9881. Available at: https://doi.org/10.3748/wjg.v20.i29.9872.

Kong, L. *et al.* (2023) 'The landscape of immune dysregulation in Crohn's disease revealed through single-cell transcriptomic profiling in the ileum and colon', *Immunity*, 0(0). Available at: https://doi.org/10.1016/j.immuni.2023.01.002.

Kreuter, R. *et al.* (2019) 'The role of obesity in inflammatory bowel disease', *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease*, 1865(1), pp. 63–72. Available at: https://doi.org/10.1016/j.bbadis.2018.10.020.

Kulakovskiy, I.V. *et al.* (2018) 'HOCOMOCO: towards a complete collection of transcription factor binding models for human and mouse via large-scale ChIP-Seq analysis', *Nucleic Acids Research*, 46(Database issue), pp. D252–D259. Available at: https://doi.org/10.1093/nar/gkx1106.

Langfelder, P. and Horvath, S. (2008) 'WGCNA: an R package for weighted correlation network analysis', *BMC Bioinformatics*, 9(1), p. 559. Available at: https://doi.org/10.1186/1471-2105-9-559.

Li, H. *et al.* (2019) 'Identifying gene function and module connections by the integration of multispecies expression compendia', *Genome Research* [Preprint]. Available at: https://doi.org/10.1101/gr.251983.119.

Li, H. *et al.* (2022) 'Integrative systems analysis identifies genetic and dietary modulators of bile acid homeostasis', *Cell Metabolism*, 34(10), pp. 1594-1610.e4. Available at: https://doi.org/10.1016/j.cmet.2022.08.015.

Li, H. and Auwerx, J. (2020) 'Mouse Systems Genetics as a Prelude to Precision Medicine', *Trends in genetics: TIG*, 36(4), pp. 259–272. Available at: https://doi.org/10.1016/j.tig.2020.01.004.

Li, J.-Y. *et al.* (2021) 'IRF/Type I IFN signaling serves as a valuable therapeutic target in the pathogenesis of inflammatory bowel disease', *International Immunopharmacology*, 92, p. 107350. Available at: https://doi.org/10.1016/j.intimp.2020.107350.

Liberzon, A. *et al.* (2011) 'Molecular signatures database (MSigDB) 3.0', *Bioinformatics*,
27(12), pp. 1739–1740. Available at: https://doi.org/10.1093/bioinformatics/btr260.

Liu, Z. *et al.* (2000) 'IL-15 is highly expressed in inflammatory bowel disease and regulates
local T cell-dependent cytokine production', *Journal of Immunology (Baltimore, Md.: 1950)*,
164(7), pp. 3608–3615. Available at: https://doi.org/10.4049/jimmunol.164.7.3608.

Lutsey, P.L., Steffen, L.M. and Stevens, J. (2008) 'Dietary Intake and the Development of the
Metabolic   Syndrome',   *Circulation*,   117(6),   pp.   754–761.   Available   at:
https://doi.org/10.1161/CIRCULATIONAHA.107.716159.

Maconi, G. *et al.* (2010) 'Pre-illness changes in dietary habits and diet as a risk factor for
inflammatory bowel disease: A case-control study', *World Journal of Gastroenterology*,
16(34), pp. 4297–4304. Available at: https://doi.org/10.3748/wjg.v16.i34.4297.

Mähler, M. *et al.* (1998) 'Differential susceptibility of inbred mouse strains to dextran sulfate
sodium-induced colitis', *American Journal of Physiology-Gastrointestinal and Liver*
Physiology,   274(3),   pp.   G544–G551.   Available   at:
https://doi.org/10.1152/ajpgi.1998.274.3.G544.

Maurya, S.K. *et al.* (2023) 'Western Diet Causes Heart Failure With Reduced Ejection Fraction
and Metabolic Shifts After Diastolic Dysfunction and Novel Cardiac Lipid Derangements',
JACC: Basic   to   Translational   Science,   0(0).   Available   at:
https://doi.org/10.1016/j.jacbts.2022.10.009.

McGuckin, M.A. *et al.* (2011) 'Mucin dynamics and enteric pathogens', *Nature Reviews*
Microbiology, 9(4), pp. 265–278. Available at: https://doi.org/10.1038/nrmicro2538.

Misiorek, J.O. *et al.* (2016) 'Keratin 8-deletion induced colitis predisposes to murine colorectal
cancer enforced by the inflammasome and IL-22 pathway', *Carcinogenesis*, 37(8), pp. 777–
786. Available at: https://doi.org/10.1093/carcin/bgw063.

Mogensen, T.H. (2019) 'IRF and STAT Transcription Factors - From Basic Biology to Roles
in Infection, Protective Immunity, and Primary Immunodeficiencies', *Frontiers in Immunology*,
9. Available at: https://www.frontiersin.org/articles/10.3389/fimmu.2018.03047 (Accessed: 19
January 2023).

Molodecky, N.A. *et al.* (2011) 'Challenges associated with identifying the environmental
determinants of the inflammatory bowel diseases', *Inflammatory Bowel Diseases*, 17(8), pp.
1792–1799. Available at: https://doi.org/10.1002/ibd.21511.

Mottis, A. *et al.* (2022) 'Tetracycline-induced mitohormesis mediates disease tolerance against
influenza', *The Journal of Clinical Investigation*, 132(17). Available at:
https://doi.org/10.1172/JCI151540.

Mun, J., Hur, W. and Ku, N.-O. (2022) 'Roles of Keratins in Intestine', *International Journal*
of Molecular Sciences, 23(14), p. 8051. Available at: https://doi.org/10.3390/ijms23148051.

Nadeau, J.H. and Auwerx, J. (2019) 'The virtuous cycle of human genetics and mouse models
in drug discovery', *Nature Reviews. Drug Discovery*, 18(4), pp. 255–272. Available at:
https://doi.org/10.1038/s41573-018-0009-9.

Nayak, R.R. *et al.* (2009) 'Coexpression network based on natural variation in human gene
expression reveals gene interactions and functions', *Genome Research*, 19(11), pp. 1953–1962.
Available at: https://doi.org/10.1101/gr.097600.109.

Peters, L.A. *et al.* (2017) 'A functional genomics predictive network model identifies
regulators of inflammatory bowel disease', *Nature Genetics*, 49(10), pp. 1437–1449. Available
at: https://doi.org/10.1038/ng.3947.

Porcu, E. *et al.* (2021) 'Differentially expressed genes reflect disease-induced rather than
disease-causing changes in the transcriptome', *Nature Communications*, 12(1), p. 5647.
Available at: https://doi.org/10.1038/s41467-021-25805-y.

Ritchie, M.E. *et al.* (2015) 'limma powers differential expression analyses for RNA-
sequencing and microarray studies', *Nucleic Acids Research*, 43(7), p. e47. Available at:
https://doi.org/10.1093/nar/gkv007.

Rutgeerts, P. *et al.* (2005) 'Infliximab for Induction and Maintenance Therapy for Ulcerative
Colitis', *New England Journal of Medicine*, 353(23), pp. 2462–2476. Available at:
https://doi.org/10.1056/NEJMoa050516.

Serino, M. *et al.* (2012) 'Metabolic adaptation to a high-fat diet is associated with a change in
the gut microbiota', *Gut*, 61(4), pp. 543–553. Available at: https://doi.org/10.1136/gutjnl-
2011-301012.

Shah, S.C. and Itzkowitz, S.H. (2022) 'Colorectal Cancer in Inflammatory Bowel Disease:
Mechanisms and Management', *Gastroenterology*, 162(3), pp. 715-730.e3. Available at:
https://doi.org/10.1053/j.gastro.2021.10.035.

Stolzer, I. *et al.* (2021) 'An IFN-STAT Axis Augments Tissue Damage and Inflammation in a
Mouse Model of Crohn's Disease', *Frontiers in Medicine*, 8. Available at:
https://www.frontiersin.org/articles/10.3389/fmed.2021.644244 (Accessed: 12 January 2023).

Tannock, L.R. *et al.* (2018) 'Serum amyloid A3 is a high density lipoprotein-associated acute-
phase protein', *Journal of Lipid Research*, 59(2), pp. 339–347. Available at:
https://doi.org/10.1194/jlr.M080887.

Tilg, H. *et al.* (2002) 'Treatment of Crohn's disease with recombinant human interleukin 10
induces the proinflammatory cytokine interferon γ', *Gut*, 50(2), pp. 191–195. Available at:
https://doi.org/10.1136/gut.50.2.191.

Uhlén, M. *et al.* (2015) 'Tissue-based map of the human proteome', *Science*, 347(6220), p.
1260419. Available at: https://doi.org/10.1126/science.1260419.

Williams, E.G. *et al.* (2016) 'Systems proteomics of liver mitochondria function', *Science*,
352(6291). Available at: https://doi.org/10.1126/science.aad0189.

Wu, Y. *et al.* (2014) 'Multilayered genetic and omics dissection of mitochondrial activity in a
mouse reference population', *Cell*, 158(6), pp. 1415–1430. Available at:
https://doi.org/10.1016/j.cell.2014.07.039.

Yang, X. *et al.* (1999) 'Targeted disruption of SMAD3 results in impaired mucosal immunity
and diminished T cell responsiveness to TGF-β', *The EMBO Journal*, 18(5), pp. 1280–1291.
Available at: https://doi.org/10.1093/emboj/18.5.1280.

Ye, R.D. and Sun, L. (2015) 'Emerging functions of serum amyloid A in inflammation',
Journal of Leukocyte Biology, 98(6), pp. 923–929. Available at:
https://doi.org/10.1189/jlb.3VMR0315-080R.

Yoshihara, K. *et al.* (2006) 'Role of interleukin 15 in colitis induced by dextran sulphate

sodium   in   mice', *Gut*,   55(3),   pp.   334–341.   Available   at:

https://doi.org/10.1136/gut.2005.076000.

Yu, G. *et al.* (2012) 'clusterProfiler: an R Package for Comparing Biological Themes Among

Gene Clusters', *OMICS: A Journal of Integrative Biology*, 16(5), pp. 284–287. Available at:

https://doi.org/10.1089/omi.2011.0118.

Zeevi, D. *et al.* (2015) 'Personalized Nutrition by Prediction of Glycemic Responses', *Cell*,

163(5), pp. 1079–1094. Available at: https://doi.org/10.1016/j.cell.2015.11.001.

Zhao, D. *et al.* (2020) 'High-Fat Diet Promotes DSS-Induced Ulcerative Colitis by

Downregulated FXR Expression through the TGFB Pathway', *BioMed Research International*,

2020, p. e3516128. Available at: https://doi.org/10.1155/2020/3516128.

715
716

Figure Legends

Figure 1. The effect of long-term HFD on gene expression in BXD colons. (**A**) Graphical representation of a pipeline of a previously described BXD mouse study (Williams *et al.*, 2016). Mice were fed HFD or CD starting from 8 weeks of age and metabolic phenotyping was performed as indicated. Mice were sacrificed at 29 weeks of age and multiple organs were collected and frozen for further analyses. BXD colon transcriptomes were analyzed in this study. CD: chow diet indicated in blue, HFD: high-fat diet indicated in red. P values were calculated by two-tailed Student's t-test and indicated as follows: * P <0.05; ** P <0.01; *** P <0.001; **** P <0.0001. (**B**) Volcano plot showing the HFD effect on BXD colon transcriptomes compared to CD and the up- and down-regulated differentially expressed genes (DEGs, absolute Log_2(Fold Change) > 0.5 and BH-adjusted P value (adj. P) < 0.05) were highlighted in red and blue, respectively. (**C**) Gene set enrichment analysis showing the effect of HFD on gene expression in BXD colons. Gene sets were grouped into five categories: Translation, Inflammation, Mitochondria, Stress, and Intermediate filament. Normalized enrichment scores (NES) were represented by color and $-\text{Log}_{10}$ (BH-adj. P) were represented by dot size and indicated as follows: * Adjusted P value <0.05; ** Adjusted P value <0.01; *** Adjusted P value <0.001. (**D**) Enrichment analysis of molecular signatures of mouse and human IBD on the transcriptome of BXD colons. UC: Ulcerative colitis, CDs: Crohn's disease

The following figure supplement is available for figure 1:

Figure supplement 1. Transcriptome profile in BXD colons.

Figure 2. Identifying susceptible strains to HFD-induced IBD-like inflammation and their effect on plasma cytokines. (**A**) Enrichment analysis of the molecular signatures in human (**bottom panel**) and mouse IBD (**top panel**) models on the gene expression of individual BXD colons. Dendrogram showing the propensity to develop a mouse IBD-like signature among BXD strains upon HFD. BXD strains were divided into three clusters: susceptible (in red), intermediate (in blue), and resistant strains (in green). Normalized enrichment scores (NES) were represented by color and -Log$_{10}$(BH-adjusted P values) were represented by dot size and indicated as follows: * Adjusted P value <0.05; ** Adjusted P value <0.01; *** Adjusted P value <0.001. UC: Ulcerative colitis, CDs: Crohn's disease (**B, C**) Boxplots showing the effect of susceptible strains on plasma IL-10 (**B**) and IL-15 (**C**) level compared to resistant strains. P values were calculated by two-tailed Student's t-test and indicated as follows: * P <0.05; ** P <0.01; *** P <0.001; **** P <0.0001.

Figure 3. Identifying IBD-related gene modules. (**A**) Pipeline for exploring the IBD-associated gene modules. A co-expression gene network was constructed based on the transcriptome of BXD colons under HFD. IBD-associated modules were then defined as gene modules under HFD that are significantly clustered in mouse DSS-induced UC signatures. (**B**) UMAP representation of the co-expression gene network under HFD. 39 co-expression modules are represented in the corresponding color and the correlated modules (Spearman correlation coefficient between the eigengene of modules > 0.7) were linked by a grey line. (**C**) Heatmap showing the enrichment of co-expression modules in mouse and human IBD gene signatures and the number and percentage of enriched genes were labeled. The number of enriched genes divided by the number of genes involved in the respective module was defined as the percentage of enriched genes. Signed –Log$_{10}$(adj. P) indicated BH-adjusted P value and the enriched gene set. Enriched modules of up- and down-regulated genes upon disease were highlighted in red and blue, respectively. UC: Ulcerative colitis, CDs: Crohn's disease.

The following figure supplement is available for figure 3:

Figure supplement 1. Exploring IBD-associated co-expression modules under CD.

Figure 4. Biological interrogation of identified IBD-related modules. (**A**, **B**) Dot plots showing the enrichment of IBD-related modules in hallmark genesets (**A**) and cell-type gene signatures of inflamed colon in Crohn's disease patients (**B**). Gene ratios higher than 0.1 are shown and represented by dot size. Dots are colored by -log$_{10}$(BH-adjusted P values). (**C**, **D**) The enriched motifs for promoters of the genes involved in module HFD_M9 (**C**) and HFD_M28 (**D**). The significantly enriched motifs (P value < 0.001) were ranked based on the percentage of enriched promoters (In top motifs) and then the top five TFs were selected. TF: Transcription factor. PWM: Positional weight matrix.

Figure 5. ModQTL mapping for two IBD-related modules and the prioritization of candidate genes. (**A**) Manhattan plot showing the ModQTL mapping result for disease-related modules HFD_M9 and HFD_M28. ModQTL maps of HFD_M9 and HFD_M28 are indicated in orange and purple, respectively. The threshold calculated by permutation test (P < 0.1) for HFD_M28 is represented by a purple dashed line. (**B**) The filtering criteria for selecting candidate genes under the ModQTL peak for HFD_M28. Genes with 2 of the described criteria are considered as candidate genes. (**C**) The most significant associations between 27 candidate genes under the ModQTL peak and Crohn's disease or UC identified through GWAS according to whole genome sequence in the human UKBB are shown in the scatter plot (**top panel**). Crohn's disease and UC are indicated by pink circle and green triangle, respectively. The threshold (-Log$_{10}$P value = 5) is represented by a red dashed line. Heatmap showing the identified 27 candidate genes of module HFD_M28 (**bottom panel**). Variants colored in blue indicate genes with high-impact genetic variants in BXD mice (including missense, frameshift, initiator codon, splice donor, splice acceptor, in-frame deletion, in-frame insertion, stop lost, stop gained). Inflammation is indicated in red and represents genes associated with inflammation based on literature mining. *Cis*-eQTL colored in purple indicates genes with *Cis*-

eQTLs. Tissue specificity colored in orange means genes that are highly expressed in human

intestine (data were downloaded from human protein atlas,

https://www.proteinatlas.org/humanproteome/tissue/intestine (Uhlén *et al.*, 2015)). GWAS

result of UC in UKBB colored in green indicates that genes are significantly associated with

human UC. (**D**, **E**) Manhattan plots showing the associated gene expression modules of *Epha6*

in mouse gastro-intestinal tract (**D**) and that of *MUC4* in human gastro-intestinal tract (**E**) (data

from https://systems-genetics.org/gmad (Li *et al.*, 2019)). The threshold is represented by the

red dashed line (absolute Gene-Module Association Score (GMAS) >= 0.268). Terms above

the threshold are identified as the significant associated terms. GO terms or gene modules are

ranked by similarity. Known associated terms are shown as red dots and new significant

associated terms are colored in black. (**F**) Dot plot showing that the expression of *MUC4* was

higher in four cell types of human inflamed colon with Crohn's disease.

The following figure supplement is available for figure 5:

Figure supplement 1. Genome-wide association studies (GWAS) for Ulcerative colitis (UC)
and Crohn's disease (CDs) in humans.

Figure 1—figure supplement 1. Transcriptome profile in BXD colons. (**A**) Principal-component analysis (PCA) of the microarray profiles of BXD colons under high-fat (HFD, indicated in red) or chow diet (CD, represented in blue). (**B**) Bar plot showing the primary principle (PC1) calculated by PCA of the colon transcriptomes in each BXD strain fed with CD (blue) or HFD (red). BXD strains highlighted in bold mean they are more resistant to dietary challenges. (**C**) Heatmap showing unsupervised hierarchical clustering of colon transcriptome in both the CD and HFD fed BXDs. (**D**) Enrichment analysis of inflammation-related genesets showing the effect of HFD on gene expression in individual BXD colon. Normalized enrichment scores (NES) were represented by color and -Log$_{10}$(BH-adjusted P values) were represented by dot size and indicated as follows: * Adjusted P value < 0.05; ** Adjusted P value < 0.01; *** Adjusted P value < 0.001. BXD strains highlighted in red represent the 3 most susceptible strains to gut inflammation upon HFD. BXD strains colored in green show no significant enrichment in gut inflammation.

Figure 3—figure supplement 1. Exploring IBD-associated co-expression modules under CD. (**A**) Heatmap showing modules enriched in mouse IBD gene signatures and the number and percentage of enriched genes were labeled. The number of enriched genes divided by the number of genes involved in the respective module was defined as the percentage of enriched genes. Signed −Log$_{10}$(adj. P) indicated BH-adjusted P value and the enriched gene set. Enriched modules of up- and down-regulated genes upon disease were highlighted in red and blue, respectively. (**B**) Heatmap showing the similarity of co-expression modules identified under CD or HFD. The number and percentage of overlapped genes were labeled. The number of overlapped genes divided by the number of genes involved in the respective module in HFD was defined as the percentage of overlapped genes. Adjusted P values calculated by BH were indicated by color. (**C**) Heatmap showing the modules under CD enriched in human IBD gene signatures and the number and percentage of enriched genes were labeled. UC: Ulcerative colitis, CDs: Crohn's disease.

Figure 5—figure supplement 1. Genome-wide association studies (GWAS) for Ulcerative

colitis (UC) and Crohn's disease (CDs) in humans. (**A**, **B**) Lollipop plots showing UC- or

CDs-associated genetic variants of *EPHA6* (**A**) and *MUC4* (**B**) based on UKBB whole genome

sequence data. Variants effect were predicted and their classification are represented by color.

VEP: Variant Effect Prediction.

Tables

Table 1: Gene signatures of mouse and human IBDs

Disease	Source	DEGs	Thresholds	Species	Tissues
DSS-induced UC	GSE131032 (Czarnewski *et al.*, 2019)	Author-provided DEGs	absolute log2FC >1 & BH-adjusted P value < 0.05	Mouse	Colon
Crohn's disease & UC	GSE16879 (Arijs *et al.*, 2009; Li *et al.*, 2019)	Author-provided DEGs	absolute FC >1 & BH-adjusted P value <0.01	Human	Colon
Crohn's disease & UC	GSE83687 (Peters *et al.*, 2017)	DEGs computed by limma package (Ritchie et al., 2015)	absolute log2FC >1 & BH-adjusted P value <0.001	Human	Colon
Crohn's disease	SCP1884 (Kong *et al.*, 2023)	Author-provided DEGs	DE coefficient >1 & BH-adjusted P value <0.05	Human	Colon

Table Supplements and Legends

Appendix 1 - Table 1. Co-expression networks under CD or HFD. Related to Figure 3

Appendix 1 - Table 2. Genes under QTL peak of module HFD_M28. Related to Figure 5

Appendix 1 - Table 3. G-MAD result for *Epha6* in mice and *MUC4* in humans. Related to

Figure 5.

Appendix 1 - Table 4. Medicine for human IBD.