Self-formation of concentric zones of

telencephalic and ocular tissues and

directional retinal ganglion cell axons

Wei Liu [1,2,3¶]*, Rupendra Shrestha* [1,2,3]*, Albert Lowe* [1,2]*, Xusheng Zhang* [2]*, Ludovic Spaeth* [4]

[1]Department of Ophthalmology and Visual Sciences

[2]Department of Genetics

[3]The Ruth L. and David S. Gottesman Institute for Stem Cell Biology and Regenerative

Medicine

[4]Dominick P. Purpura Department of Neuroscience

Albert Einstein College of Medicine, Bronx, NY 10461

[¶]*Lead Contact.* Email: Wei Liu, wei.liu@einsteinmed.edu

Key words: human pluripotent stem cell, organoid, telencephalon, optic stalk, optic disc,

neuroretina, tissue patterning, self-organization, retinal ganglion cell, axon growth and

pathfinding, cell surface protein, cell isolation, glaucoma

Summary

The telencephalon and eye in mammals are originated from adjacent fields at the anterior

neural plate. Morphogenesis of these fields generates telencephalon, optic-stalk, optic-disc, and

neuroretina along a spatial axis. How these telencephalic and ocular tissues are specified

coordinately to ensure directional retinal ganglion cell (RGC) axon growth is unclear. Here, we

report the self-formation of human telencephalon-eye organoids comprising concentric zones of

telencephalic, optic-stalk, optic-disc, and neuroretinal tissues along the center-periphery axis.

Initially-differentiated RGCs grew axons towards and then along a path defined by adjacent

PAX2+ optic-disc cells. Single-cell RNA sequencing of CONCEPT organoids not only confirmed

telencephalic and ocular identities but also identified expression signatures of early optic-disc,

optic-stalk, and RGCs. These signatures were similar to those in human fetal retinas. Optic-disc

cells in CONCEPT organoids differentially expressed *FGF8* and *FGF9*; FGFR inhibitions

drastically decreased RGC differentiation and directional axon growth. Through the identified

RGC-specific cell-surface marker CNTN2, electrophysiologically-excitable RGCs were isolated

under a native condition. Our findings provide insight into the coordinated specification of early

telencephalic and ocular tissues in humans and establish resources for studying RGC-related

diseases such as glaucoma.

(184 words)

Impact statement

A human telencephalon-eye organoid model that exhibited axon growth and pathfinding from

retinal ganglion cell (RGC) axons is reported; via cell surface marker CNTN2 identified using

scRNA-seq, early RGCs were isolated under a native condition.

Introduction

Our understanding of eye and brain development in humans is mostly deduced from animal studies. In mice, fate mapping of the anterior neural plate reveals that the eye field is in rostral regions surrounded anteriorly and laterally by the telencephalic field and caudally and medially by the hypothalamic field, indicating the proximity of their embryonic origins (Inoue et al., 2000). Subsequent evagination of the eye field generates bilateral optic vesicles and optic stalks, and the optic stalk connects the optic vesicle to the forebrain. The optic vesicle then invaginates ventrally, resulting in the formation of double-layered optic cups in which the inner and outer layers develop into the neuroretina and retinal pigment epithelium (RPE), respectively. The posterior pole of the optic-cup forms the optic disc (also known as optic nerve head). In the central neuroretina close to the nascent optic disc, retinal ganglion cells (RGCs) start to appear as the optic fissure nearly closes. Early RGC axons find their path toward the optic disc and then enter the optic stalk to reach the brain. Concentrically organized growth-promoting and growth-inhibitory guidance cues around the optic disc regulate RGC axon growth and pathfinding through multiple mechanisms (Erskine and Herrera, 2007, 2014; Oster et al., 2004). Therefore, early eye morphogenesis leads to coordinated specification of the telencephalic, optic stalk, optic disc, and retinal tissues along a spatial axis.

Early telencephalic and eye development is marked and regulated by a group of tissue-specific transcription factors and signal transduction molecules (Wilson and Rubenstein, 2000). In mice, Foxg1 is specifically expressed in the presumptive telencephalon and is essential for the development of the cerebral hemispheres (Xuan et al., 1995). Pax6 is specifically expressed in the eye field and is essential for the development of multiple ocular tissues, such as the neuroretina, lens, and RPE (Ashery-Padan et al., 2000; Baumer et al., 2003; Marquardt et al., 2001; Zuber et al., 2003). Vsx2 and Mitf are specifically expressed in the neuroretina and RPE, respectively, and are essential for retinal development (Bharti et al., 2012; Horsford et al., 2005;

Liu et al., 2010; Rowan et al., 2004). Pax2 is expressed in the optic stalk, optic vesicles, central

neuroretina, and optic disc; Pax2 is essential for optic stalk and nerve development (Macdonald

et al., 1997; Martinez-Morales et al., 2001; Schwarz et al., 2000; Torres et al., 1996). Fgf8 is

specifically expressed at the rostral forebrain at early stages, induces Foxg1 expression

(Shimamura and Rubenstein, 1997), and regulates telencephalic patterning in a dose-

dependent manner (Storm et al., 2006). Fgf8 also maintains Pax2 expression in the optic stalk

(Soukkarieh et al., 2007), and Fgf8 and Fgf3 coordinate the initiation of retinal differentiation in

chicks (Martinez-Morales et al., 2005; Vogel-Hopker et al., 2000). Despite these findings in

vertebrates, it is unclear how telencephalic and ocular tissues are specified in humans.

Human organoids are transformative since they enable us to study the mechanisms of cell

specification and differentiation directly in human tissues (Lancaster and Knoblich, 2014). Self-

organized three-dimensional (3-D) retinal organoids are originally reported by Sasai's group and

further improved in follow-up studies (Cowan et al., 2020; Eiraku et al., 2011; Fligor et al., 2021;

Kim et al., 2019; Lowe et al., 2016; Meyer et al., 2011; Nakano et al., 2012; Reichman et al.,

2014; Zhong et al., 2014). We and others have demonstrated that three-dimensional (3-D)

retinal organoids derived from human embryonic stem cells (hESCs) display a stratified

structure containing all major types of retinal cells. Although RGCs are differentiated in 3-D

retinal organoids, there is no proper RGC axon outgrowth toward the optic disc as seen *in vivo*

since the optic disc-like tissue is missing in 3-D retinal organoids. When these organoids are

dissociated into single cells or cut into pieces for adherent culture, RGCs generate neurites

(Fligor et al., 2018; Langer et al., 2018; Teotia et al., 2017). A variety of brain organoids have

been described (Kadoshima et al., 2013; Lancaster et al., 2013; Mariani et al., 2012; Qian et al.,

2017; Velasco et al., 2019). Although rudimentary ocular tissues are occasionally found in some

brain organoids (Gabriel et al., 2021), optic disc- and stalk-like tissues are absent. Collectively,

RGC axon outgrowth and pathfinding directed by optic disc- and stalk-like tissues in organoids

have not been reported.

Tissue patterning and coordinated specification are fundamental for body plan formation *in vivo.*

Remarkably, concentric zones of trophectoderm, endoderm, mesoderm, and ectoderm (Etoc et

al., 2016; Minn et al., 2020), neural plate and neural plate border (Xue et al., 2018), and

ectodermal cells (Hayashi et al., 2016) are self-organized from single pluripotent stem cells,

indicating patterning and coordinated specification of these tissues *in vitro.* Nevertheless, optic

disc and stalk tissues are not reported in any of these structures.

We hypothesize that coordinated specification from the anterior ectodermal epithelium via

morphogen gradients leads to self-organization of telencephalic and ocular tissues, including

optic disc- and stalk-like tissues that provide guidance cues for RGC axon growth and

pathfinding. In support of this hypothesis, we generated self-formed human telencephalon-eye

organoids that comprise <u>con</u>centric zones of anterior <u>e</u>ctodermal <u>p</u>rogeni<u>t</u>ors (CONCEPT),

including FOXG1+ telencephalic, PAX2+ optic stalk, PAX2+ optic disc, and VSX2+ neuroretinal

cells along the center-periphery axis. We call this structure as an organoid since it displays a

spatially-organized structure with cell identities mimicking those tissues *in vivo.* In CONCEPT

organoids, early differentiated RGCs grew their axons toward and then along a path defined by

adjacent PAX2+ optic-disc cells. Single-cell RNA sequencing of CONCEPT organoids not only

confirmed telencephalic and ocular identities but also discovered expression signatures of cell

clusters. Additionally, CONCEPT organoids and human fetal retinas had similar expression

signatures. Furthermore, PAX2+ optic-disc cells differentially expressed *FGF8* and *FGF9*;

inhibition of FGF signaling with an FGFR inhibitor during early RGC differentiation drastically

decreased the number of RGCs somas and nearly ablated directional axon growth. Using RGC-

specific cell-surface marker CNTN2 identified in our single-cell RNA-sequencing, we developed

a one-step method for isolating electrophysiologically-excitable RGCs under a native condition.

Our studies provide deeper insight into the coordinated specification of telencephalic and ocular

tissues in humans and establish resources for studying neurodegenerative diseases such as

glaucoma.

Results

Generation of telencephalon-eye organoids that are composed of <u>conc</u>entric zones of <u>a</u>nterior <u>e</u>ctodermal <u>p</u>rogeni<u>t</u>ors (CONCEPT)

The telencephalon and eye in mammals are originated from adjacent fields at the anterior neuroepithelium (Inoue et al., 2000). Morphogenesis of these embryonic fields leads to the formation of telencephalon, optic stalk, optic disc, and neuroretina along a spatial axis. Early differentiated retinal ganglion cells (RGCs) in the neuroretina grow axons toward the optic disc and then along the optic stalk to reach the brain. How these telencephalic and ocular tissues are specified coordinately to ensure directional RGC axon growth is unclear in humans.

Cysts are hollow spheres of a columnar epithelium that is induced from human pluripotent stem cells via embedding hESC sheets in Matrigel; they are used to generate retinal cells (Kim et al., 2019; Lowe et al., 2016; Zhu et al., 2013). Nevertheless, developmental potentials of cysts have not been characterized.

Using the cysts, here we generate telencephalon-eye organoids that are composed of <u>conc</u>entric zones of anterior <u>e</u>ctodermal <u>p</u>rogeni<u>t</u>ors (CONCEPT) (Fig. 1A, B). Cysts were generated as previously described (Kim et al., 2019; Lowe et al., 2016). The epithelial structure of cysts was demonstrated by apical localization of the TJP1::GFP reporter at the lumen (Fig. 1C), consistent with previous findings (Lowe et al., 2016; Zhu et al., 2013). To assess developmental potentials of cysts, individual cysts were manually picked and then seeded onto the Matrigel-coated surface at low densities (Fig. 1A). After attaching to the culture surface, cysts grew as dome-shaped individual colonies. Subsequent culture of the colonies in a KSR medium (see Materials and Methods) led to the self-formation of concentric zones of anterior ectodermal progenitors: an elevated central zone surrounded by multiple zones. This

morphology was identifiable under a stereomicroscope (Fig. 1D) or an inverted microscope. If

multiple cysts were fused together or cysts for seeding were too big or small, CONCEPT

structures were affected or incomplete. Overall, the concentric structure was abundant.

CONCEPT structures expressed gene markers for telencephalon and eye in concentric patterns.

In mice, Foxg1 is specifically expressed in the E10 telencephalon at high levels; it was also

expressed in the optic stalk, invaginating optic cup, and lens at lower levels, with the rostral

optic stalk connecting the telencephalic vesicle to the invaginating optic cup (Fig. 1E) (Xuan et

al., 1995); Vsx2 is specifically expressed in the E10 neuroretina (Fig. 1F); Pax6 is specifically

expressed in the E10.5 neuroretina, RPE, lens vesicles, and surface ectoderm (Fig. 1G) (Liu et

al., 2010). In CONCEPT structures, the expression of FOXG1, VSX2, and PAX6 generally

exhibited concentric patterns spanning from the center to the periphery at days 15-17 (Fig. 1H-

1K, supp. Fig. S1; a scheme in the left panel in B). At this stage, PAX6 was expressed in

multiple concentric zones at distinct levels, with a higher level at a circular zone that also

expressed VSX2 (Fig. 1J, supp. Fig. S1). Concentric zones of FOXG1, VSX2, and PAX6

expression were also found at later stages. At day 26, VSX2 expression expanded peripherally

compared to its expression at day 17 (supp. Figs. S1C, S2C) and largely overlapped with PAX6

expression (supp. Fig. S2A-S2C). Additionally, VSX2 expression was higher in a zone close to

the center and relatively lower in more peripheral zones (supp. Fig. S2C). Therefore, the

CONCEPT structures comprise spatially organized telencephalic and ocular tissues and thus

are named as telencephalon-eye organoids.

Morphogens FGFs and BMPs play crucial roles in patterning the forebrain and eye *in vivo*. In

mice, *Fgf8* is specifically expressed at the rostral forebrain at early stages, induces *Foxg1*

expression (Shimamura and Rubenstein, 1997), and regulates telencephalic patterning in a

dose-dependent manner (Storm et al., 2006). *Bmp4* and *Bmp7* are expressed in the

dorsomedial telencephalon, optic vesicles, and presumptive lens placodes (Danesh et al., 2009;

Dudley and Robertson, 1997; Furuta and Hogan, 1998; Solloway et al., 1998). *Bmp4* is required

for lens induction (Furuta and Hogan, 1998), and *Bmp7* is required for proper patterning of the

optic fissure (Morcillo et al., 2006). In CONCEPT telencephalon-eye organoids, *FGF8, BMP4,*

and *BMP7* were expressed in attached cell colonies starting from early stages (Fig. 1P, S, T)

and subsequently exhibited circular patterns. At day 10, the expression of *BMP4* and *BMP7*

delineated multiple rings, with a big ring mostly at the center surrounded by numerous small

rings (Fig. 1P, S). These observations suggest that the attachment of a single-lumen cyst to the

culture surface caused differences in cell behaviors: some cells separated from the original cyst

and migrated peripherally; some of the separated cells formed small ring-like structures; the

cells that remained at the center formed a big ring-like structure. At day 17, circular expression

patterns of *BMP4* and *FGF8* emerged (Fig. 1Q, U). At day 25, *BMP4, FGF8,* and *BMP7*

expression clearly marked circular zones (Fig. 1R, V, W). Expression profiles of *FGFs* and

BMPs were highly reproducible (supp. Fig. S3A-S3D). Hence, the expression of *FGFs* and

BMPs spontaneously form circular gradients, which would dictate coordinated cell specification

in CONCEPT telencephalon-eye organoids.

Early differentiated RGCs grow directional long axons toward and then along a path

defined by PAX2+ cell populations in CONCEPT telencephalon-eye organoids

To assess cell differentiation in CONCEPT telencephalon-eye organoids, we examined marker

expression for RGCs, the first type of cells that differentiate in the neuroretina. In mice,

transcription factor Pou4f2 is expressed in the early differentiated RGCs and required for the

development of a large set of RGCs (Gan et al., 1996). Tubb3 is expressed in the somas and

axons of differentiating RGCs.

In CONCEPT organoids, POU4F2 and TUBB3 were detectable as early as day 17 and

increased to higher levels at day 22. RGC somas were marked by POU4F2 and TUBB3 co-

expression; RGC axons were marked by TUBB3 expression. Interestingly, RGCs grew

directional long axons that followed a circular path (Fig. 2A, B). The circular path of RGC axon

outgrowth became more evident in organoids at day 26 (Fig. 2C, D). In mice, the initially

differentiated RGCs are adjacent to the optic disc; their axons grow towards the optic disc to exit

the eye and then navigate within the optic stalk to reach their targets in the brain. The optic disc

and stalk specifically expressed Pax2 at distinct levels (Fig. 2E-G), consistent with previous

findings (Bassett et al., 2010; Martinez-Morales et al., 2001; Mui et al., 2005). Additionally,

presumptive optic disc cells expressed VSX2 whereas optic stalk cells do not (Figs. 1F; 2E) (Liu

et al., 2010). In CONCEPT organoids, there were two PAX2+ cell populations that formed two

adjacent rings (Fig. 2H-L). POU4F2+ RGCs grew TUBB3+ axons toward and then navigated

along the adjacent PAX2+ VSX2+ cell population (Fig. 2H-L), mimicking axon growth from the

nascent RGCs toward the PAX2+ optic disc *in vivo*. Meanwhile, the PAX2+ VSX2- cell

population at the inner zone set up an inner boundary of the path for RGC axon growth,

mimicking the PAX2+ optic stalk that spatially confines RGC axon growth *in vivo*. Based on

these findings, we designate PAX2+ VSX2+ cells and PAX2+ VSX2- cells in CONCEPT

organoids as optic disc and optic stalk cells, respectively. Taken together, our findings

demonstrate that RGCs grow directional axons toward and then along a path defined by PAX2+

cell populations in CONCEPT telencephalon-eye organoids.

CONCEPT telencephalon-eye organoids also contain lens cells that undergo terminal

differentiation

Besides anterior neuroectodermal cells, we wondered whether CONCEPT organoids also

contain non-neural ectodermal cells. CONCEPT telencephalon-eye organoids at around day 25

contained transparent structures reminiscent of the ocular lens. To determine their cell identity,

we performed immunostaining. Starting at day 22, lens markers CRYAA and beta crystallin

(CRY B) were found (Fig. 3A, B) and continuously expressed in the transparent cell structures

(Fig. 3C, D, K). Interestingly, these transparent structures were not stained by DAPI (Fig. 3C-F),

indicating denucleation in these lens cells. When CONCEPT organoids were detached using

Dispase at around day 28 and continuously grown as suspension cultures, crystal-like clusters

with fused transparent spheres -- lentoid bodies-- were found, and they continuously survived

for months (Fig. 3I, J). Lentoid bodies highly expressed gamma crystallin (CRY G) (Fig. 3L),

confirming their lens identity. In mice, terminally-differentiated lens fiber cells are free of

organelles and featured by specialized interlocking cell membrane domains shown as ball-and-

sockets and protrusions (Biswas et al., 2010). Using transmission electron microscopy, we

found that our lentoid bodies were free of organelles and exhibited ball-and-socket structures

(Fig. 3M, N). Taken together, our findings demonstrate that CONCEPT telencephalon-eye

organoids also contain lens cells that undergo terminal differentiation; FGFs in CONCEPT

organoids likely promote terminal lens differentiation as seen in other settings (Lovicu and

McAvoy, 2001).

Single-cell RNA sequencing analysis identifies telencephalic and ocular cell populations

in CONCEPT telencephalon-eye organoids

To fully characterize cell populations in CONCEPT organoids and the mechanisms underlying

RGC axon pathfinding, we performed single-cell RNA sequencing (10x Genomics) of the

organoids at day 24, around the stage when RGCs grew long axons toward and along the

PAX2+ VSX2+ cell population. Cell Ranger mapping showed that 11158 single cells were

sequenced at a depth of 27,842 reads and 2,967 genes per cell, and the dataset were further

analyzed using Seurat (v3.2.0) (Stuart et al., 2019). Cell filtration (nFeature_RNA > 200 &

nFeature_RNA < 6000 & percent.mt < 20) resulted in 10218 cells. Cell clustering grouped cells

into 14 clusters (Fig. 4A; Tab. 1), and cell cycle phases were identified based on cell cycle

scores using an established method (Tirosh et al., 2016) (Fig. 4B). Several cell clusters, *e.g.,*

clusters 4, 8, 9, were separated along cell cycle phases (Fig. 4A, B).

We next assessed cell identities using established markers. Mesoderm, endoderm, and neural

crest are identified by a group of gene markers (Poh et al., 2014; Simoes-Costa and Bronner,

2015; Varga et al., 2021; Yao et al., 2017). In CONCEPT organoids, gene markers for

mesoderm (*TBXT, GATA2, HAND1*), endoderm (*GATA1, GATA4, SOX17*), and neural crest

(*SNAI1, SOX10, FOXD3*) were not expressed (supp. Fig. S4). In contrast, gene markers for the

anterior neuroectoderm were widely expressed. CONCEPT organoids at day 24 were mostly

composed of FOXG1+ telencephalic cells (clusters 10, 3, 12, 1, 4, 8, 9, 13) and PAX6+ and/or

VSX2+ retinal cells (clusters 2, 7, 5, 6, 11) (Fig. 4A-4E). Cluster 0, which comprised both

telencephalic and retinal cells (Fig. 4C-4E), was marked by negative gene markers (supp. Fig.

S5A-S5D). Cluster 0, together with cluster 10, had much lower counts of captured RNA and

genes compared to other clusters (supp. Fig. S5E-S5G) and was therefore not dissected further.

In contrast to telencephalon and retinal markers, diencephalon markers (*GBX2, WNT3, SOX14*)

(Chatterjee and Li, 2012; Martinez-Ferre and Martinez, 2012) and midbrain/hindbrain markers

(*EN2, PAX7, TFAP2B*) (Yao et al., 2017) were rarely expressed (supp. Fig. S6). Lens markers

CRYAA and *FOXE3* were expressed in a very small cell population. Those denucleated lens

cells were not captured in the scRNA-seq since the absence of nucleus prevented active

transcription. Therefore, lens cells did not form a separate cluster probably due to their very low

abundance. Taken together, these findings indicate that CONCEPT telencephalon-eye

organoids at day 24 are mostly composed of FOXG1+ telencephalic cells and PAX6+ and/or

VSX2+ ocular (mostly retinal) cells.

We next characterized FOXG1+ telencephalic cells via assessing differentially expressed genes

(DEGs) of clusters. Top DEGs in clusters 4, 8, 9, 1 included *SOX3, FGFR2, PRRX1,* and

EDNRB (supp. Fig. S7A-S7D)*,* which mouse orthologs are specifically expressed in the dorsal

telencephalon (supp. Fig. S8B-S8E). Top DEGs in clusters 12, 3, 10 included *DLX2, DLX6-AS1,*

DLX1, and *RGS16* (supp. Fig. S7E-S7H), which mouse orthologs are specifically expressed in

the ventral telencephalon (supp. Figs. S8F-S8I). Consistently, *Foxg1* is expressed in both dorsal

and ventral telencephalon in E14.5 mouse embryos (supp. Fig. S8A). Therefore, telencephalic

cells in CONCEPT organoids at day 24 comprise both dorsal and ventral telencephalic cells.

Cluster 13 also expressed *FOXG1*, but it was separated from other FOXG1+ cells. Top DEGs in

cluster 13 included *TRIB3, DWORF, SLC7A11, GDF15,* and *UNC5B;* cell identities of cluster 13

were undetermined.

PAX6+ and/or VSX2+ retinal cells were grouped into several clusters. *VSX2* was expressed in

clusters 2, 7, 5, and 0. Clusters 2 was at G1 phase, cluster 7 was at G1 and S phases, and

cluster 5 was at S and G2M phases (Fig. 4A, B, E). Cluster 6 did not express *VSX2* but

expressed *PAX6* and was at the G1 phase (Fig. 4A, B, D, E). Cluster 6 differentially expressed

RPE markers, *e.g., PMEL, HSD17B2, DCT,* and *MITF* (supp. Fig. S9), indicating they were

mostly differentiating RPE cells.

Taken together, our single-cell RNA sequencing analysis of CONCEPT telencephalon-eye

organoids at day 24 confirms their telencephalic and ocular identities, establishing a valuable

transcriptomic dataset for mechanistic studies.

Identification of two PAX2+ cell populations that mimic to the optic disc and optic stalk,

respectively, in the scRNA-seq dataset of CONCEPT telencephalon-eye organoids

To identify PAX2+ cell populations that defined the path for RGC axon outgrowth, we examined

PAX2 expression in the dataset. *PAX2* was mostly expressed in cluster 2 and subsets of

clusters 7, 5, 0, 4, 8, and 9 (Fig. 4A, F). Since clusters 2, 7, 5, and 0 expressed *VSX2* (Fig. 4E,

F), we deduced that PAX2+ VSX2+ cells in cluster 2 and subsets of clusters 7, 5, and 0

corresponded to those PAX2+ VSX2+ cells that defined the path for RGC axon growth (Figs.

2H-L) and therefore were assigned as optic disc (OD) cells (Fig. 4F). PAX2+ VSX2+ cells also

expressed optic disc and optic stalk marker *SEMA5A* (Oster et al., 2003) (supp. Figs. S10, S12).

We particularly focused on cluster 2 since PAX2+ VSX2+ cells were mostly found in this cluster.

Interestingly, *COL9A3* and *COL13A1* were differentially expressed in cluster 2 (Fig. 4G, 4H;

supp. Fig. S10). Previous *in situ* hybridization results indicate that *COL13A1* is prominently

expressed in the optic disc of human fetal retinas (Sandberg-Lall et al., 2000). These findings

support that PAX2+ VSX2+ cells in CONCEPT telencephalon-eye organoids corresponded to

optic disc cells in human fetal retinas. Cluster 2 also differentially expressed glaucoma gene

CYP1B1, morphogen-encoding genes *LEFTY2, FGF9,* and *FGF8* (Fig. 4I-L; supp. Fig. S10).

Restricted expression of *FGF8, PAX2, SEMA5A, CYP1B1,* and *LEFTY2* in CONCEPT

organoids was validated using in *situ* hybridization (Figs. 1T-V; supp. Fig. S10).

In contrast to assigned optic disc cells, PAX2+ VSX2- cells were at edges of telencephalic

clusters 4, 8, 9 (Fig. 4A, E, F) and expressed the optic stalk marker *VAX1* (Fig. 4F, M). These

cells corresponded to those PAX2+ VSX2- cells that set up the inner boundary of the path for

RGC axon growth in CONCEPT organoids (Fig. 2H-L). Therefore, PAX2+ VSX2- cells in

subsets of clusters 4, 8, 9 were assigned as optic stalk cells (OS) (Fig. 4F). PAX2+ VSX2- optic

stalk cells also differentially expressed *CNTNAP2, ALDHA3,* and *LAMP5* (Fig. 4N-P), which

mouse orthologs are expressed in the optic stalk/nerve (supp. Fig. S11). Compared to assigned

optic-disc cells (PAX2+ VSX2+), assigned optic-stalk cells (PAX2+ VSX2-) were closer to

telencephalic cells in both the UMAP graph (Fig. 4M-P; supp. Fig. S7A-S7D) and their

positioning in CONCEPT organoids (Fig. 2H-L), mimicking the sequence in spatial positions of

forebrain, optic stalk, and optic disc in E13.5 mouse embryos.

We also assessed the expression of known axon guidance genes in CONCEPT organoids.

SEMA5a and *EFNB1* were expressed in both assigned optic disc and stalk cells, *EFNB2* was

highly expressed in assigned optic disc cells, and *NTN1* was mostly expressed in assigned optic

cells (supp. Fig. S12).

Collectively, we identify two PAX2+ cell populations that mimic the optic disc and optic stalk,

respectively, in the scRNA-seq dataset of CONCEPT telencephalon-eye organoids.

Identification of differentiating retinal neurons and RGC-specific cell-surface marker

CNTN2

We next assessed cell identities of cluster 11. Interestingly, cluster 11 differentially expressed

neurogenic retinal progenitor marker *ATOH7* (Fig. 4Q), indicating that these cells underwent

retinal differentiation. A large portion of cluster 11 differentially expressed *POU4F2* (Fig. 4R),

indicating their RGC identity. Notably, *CNTN2,* which encodes a glycosylphosphatidylinositol

(GPI)-anchored cell membrane protein, was largely co-expressed with *POU4F2* (Fig. 4R, 4S).

Cluster 11 also differentially expressed RGC marker *SNCG* (Fig. 4T). Two small portions of

cluster 11 differentially expressed photoreceptor markers *OTX2* and *CRX* (Fig. 4U, 4V) and

amacrine/horizontal cell markers *TFAP2C* and *PTF1A* (Fig. 4W-4X)*,* respectively. Therefore,

cluster 11 comprises differentiating retinal neurons, predominantly early RGCs; cell surface

protein gene *CNTN2* is differentially expressed in early RGCs.

Transcriptomic comparisons between CONCEPT organoids, brain/optic organoids, and

human fetal retinas

We next performed transcriptomic comparisons between CONCEPT organoids, brain/optic

organoids, and human fetal retinas. "Optic vesicle-containing brain organoids" with "axon-like

projections" were recently reported (Gabriel et al., 2021). We were curious about the similarities

and differences between CONCEPT organoids and Gabriel et al.'s organoids. We loaded

Gabriel et al.'s scRNA-seq dataset

(https://github.com/Gpasquini/Brain_organoids_with_OpticCups) and were able to reproduce

the cell clustering described in their paper (supp. Fig. S13A, S13C). Then, we assessed *PAX2*

expression in Gabriel et al.'s organoids. Interestingly, PAX2+ cells were extremely rare; there

were only eight and five PAX2+ cells scattering across Gabriel et al.'s datasets at day 30 and

day 60, respectively (supp. Fig. S13B, S13D), indicating that Gabriel et al.'s organoids do not

have PAX2+ cell clusters that were found in CONCEPT organoids. In Gabriel et al.'s organoids

on day 30, FOXG1+ telencephalic progenitors (supp. Fig. S13E), SIX3+ neuroretinal progenitors

(supp. Fig. S13F), and ATOH7+ neurogenic progenitors (supp. Fig. S13G) were extremely rare.

VSX2 was not even found in the dataset of Gabriel et al.'s organoids on day 30; *VSX2* was

filtered out, probably due to its extremely low expression. Based on these findings, we conclude

that PAX2+ optic disc, PAX2+ optic stalk, FOXG1+ telencephalic, and VSX2+ neuroretinal cell

clusters that were found in CONCEPT organoids did not exist in Gabriel et al.'s organoids,

indicating striking differences between Gabriel et al.'s organoids and our CONCEPT

telencephalon-eye organoids.

Transcriptomic comparisons between CONCEPT organoids at day 24 and human fetal retinas

at HWG9 (Lu et al., 2020) indicate that CONCEPT organoids and human fetal retinas had

similar expression signatures. Cell clustering of HGW9 retinas identified 21 clusters (Fig. 5A).

Notably, cluster 18 differentially expressed *PAX2, COL9A3, CYP1B1, SEMA5A,* and *FGF9* (Fig.

5B-5F), which were top DEGs of cluster 2 in CONCEPT organoids (Fig. 4F, 4G, 4I, 4K; supp.

Fig. S12A). Overall, 64/113 DEGs of cluster 18 in HGW9 were also DEGs of cluster 2 in

CONCEPT organoids. In both HGW9 and CONCEPT organoids, expression of *OLIG2, CD44,*

and *GFAP* was undetectable (supp. Fig. S14), indicating that astrocytes had not been

generated yet at these stages. In the HGW9 dataset, clusters 0-3, 9, 10, 12, 13, 17, 19 highly

expressed *VSX2*, indicating that these cells were retinal progenitor cells (Fig. 5G). Additionally,

cluster 18 was a protrusion from VSX2+ cell clusters in the UMAP graph (Fig. 5A-5G), which

pattern was similar to the position of cluster 2 relative to VSX2+ clusters in the UMAP graph of

CONCEPT organoids (Fig. 4A, 4E, 4F). In the HGW9 dataset, *ATOH7* expression marked

neurogenic retinal progenitors (Fig. 5H), *OTX2* expression marked early photoreceptors (Fig. 5I),

and *TFAP2C* and *PTF1A* expression marked amacrine and/or horizontal cells (Fig. 5J, 5K).

Notably, *CNTN2* was also largely co-expressed with *POU4F2* in human fetal RGCs (Fig. 5L, M).

Relative positions of ATOH7+, POU4F2+, CNTN2+, OTX2+ , TFAP2C+, and PTF1A+ cells in

HGW9 and CONCEPT retinas were similar (Fig. 5H-5M, Fig. 4Q-4S, 4U, 4W, 4X). When cells in

cluster 18 and retinal progenitor clusters from the HGW9 dataset were combined with cells in

clusters 2, 4, 5, 7 from the CONCEPT dataset for Seurat anchor-based clustering, cells in

cluster 18 from HGW9 (H18) were grouped with cluster 2 from CONCEPT organoids (C2,

assigned optic disc; N), and these cells expressed both *PAX2* and *VSX2* (arrowheads in Fig.

5N-5R). A small portion of H18 cells were grouped with cluster 4 from CONCEPT organoids (C4,

assigned optic stalk; N), and these cells expressed *PAX2* but not *VSX2* (arrows in Fig. 5N-5R).

These findings indicate that most of PAX2+ cells from the HGW9 dataset are optic disc cells

whereas a small portion are optic stalk cells, which is consistent with the expectation that optic

disc cells are preserved whereas optic stalk cells are mostly lost in whole retina preparations.

We would like to point out that human fetal retinas HGW9 were at a more advanced stage than

CONCEPT organoids, as indicated by the counts of days and the proportion of RGCs (Fig. 5L,

5M; Fig. 4R, 4S). It is conceivable that more transcriptomic similarities will be identified when

stage-matched samples are used for comparisons.

We then compared functional annotations of DEGs (top 200 genes) of cluster 2 in CONCEPT

organoids and DEGs (113 genes) of cluster 18 in human fetal retinas HGW9. Top GO terms in

GO:MF, GO:CC, and GO:BP are shown (Fig. 6). For DEGs of cluster 2 in CONCEPT organoids,

top enriched GO terms in GO:MF, GO:CC, and GO:BP were extracellular matrix structural

constituent, collagen-containing extracellular matrix, and system development, respectively.

Additional interesting GO:BP terms included axon development, astrocyte development, eye

development, response to growth factor, cell adhesion, cell motility, neuron projection

development, glial cell differentiation, and signal transduction. For DEGs of cluster 18 in human

fetal retinas HGW9, top enriched GO terms in GO:MF, GO:CC, and GO:BP were cell adhesion

molecule binding, extracellular space, and developmental process, respectively. A large number

of GO terms were enriched in both samples, further indicating transcriptomic similarities in

PAX2+ optic-disc cells between CONCEPT organoids and human fetal retinas.

Collectively, transcriptomic comparisons indicate that CONCEPT telencephalon-eye organoids

are innovative and share similar expression signatures with human fetal retinas, such as

specific *CNTN2* expression in early RGCs and co-expression of *PAX2* and *VSX2* in assigned

optic-disc cells.

396 One-step isolation of developing human RGCs via native marker CNTN2

Differential *CNTN2* expression in early RGCs of both CONCEPT organoids and human fetal

retinas caught our attention because CNTN2 is a cell-surface protein that may be used as a

native marker for RGC isolation. In literature, Cntn2 is specifically expressed in developing

RGCs in mice and chicks (Chatzopoulou et al., 2008; Yamagata and Sanes, 2012).

In CONCEPT at day 25, CNTN2 exhibited an expression pattern very similar to that of TUBB3

(Figs. 2, 7). CNTN2 was found in the cell membrane of RGCs that expressed POU4F2 in the

cell nucleus (Fig. 7A, D). PAX2+ cells formed inner and outer concentric zones, which were

designated as optic-stalk and optic-disc cells, respectively. POU4F2+ RGCs formed a dense

circular zone adjacent to PAX2+ optic-disc cells; POU4F2+ RGCs were sparse in more

peripheral areas (Fig. 7B). These findings indicate that early differentiated RGCs were adjacent

to PAX2+ optic-disc cells. Interestingly, RGCs in the dense POU4F2+ zone grew CNTN2+

axons toward and then along the path defined by adjacent PAX2+ optic-disc cells (Fig. 7A-E).

In areas where there was a gap in PAX2+ optic-disc cells, CNTN2+ axons exited the circular

path (diamond arrow in Fig. 7A; a gap between two arrowheads in Fig. 7B, F). In regions where

POU4F2+ RGCs were a few hundred micrometers away from PAX2+ optic-disc cells, RGCs

grew axons in centrifugal directions (arrow in Fig. 7C). Very similar findings were found using

hiPSCs (supp. Fig. S15), indicating the reproducibility of CONCEPT organoids in multiple cell

lines. PAX2+ optic-disc cells did not express ALDH1A3; instead, the path for RGC axon growth

was bordered by two cell populations that highly expressed ALDH1A3 (Fig. 7G-H). In mice,

Aldh1a3 expression was low in differentiating RGCs in the central retina but high in peripheral

retinal progenitors (Fig. 7K) and in the optic stalk (Li et al., 2000), consistent with ALDH1A3

expression in CONCEPT organoids (Fig. 7G, H). In mice, Vax1 and Vax2 are expressed in the

optic stalk and ventral retina and are jointly required for the optic stalk development (Mui et al.,

2005; Take-uchi et al., 2003). In CONCEPT organoids, cells that set up the inner boundary for

RGC axon growth also expressed optic-stalk marker VAX1/2 (Fig. 7I, J; the antibody recognizes

both VAX1 and VAX2). Taken together, immunostaining of CONCEPT organoids confirms that

CNTN2 is a specific cell-surface marker for differentiating human RGCs; RGCs grow axons

toward and then along a path that is defined by adjacent optic disc cells (PAX2+ VSX2+

ALDH1A3-).

Since cell surface protein CNTN2 was specifically expressed in developing human RGCs, we

sought to test whether CNTN2 can be used as a biomarker for isolating human RGCs under a

native condition. To that goal, retinal organoids in suspension culture on days 41 - 70 were

dissociated using Accutase to generate a single cell suspension, which was then subject to

magnetic-activated cell sorting (MACS) with an antibody against CNTN2. From 100 retinal

organoids on day 41 to 48, around 385,000 RGCs were isolated. Isolated RGCs were plated

onto Matrigel-coated chamber slides for 10-day adherent culture. These cells exhibited neuronal

morphology and widely expressed RGC markers TUBB3 and POU4F2 (Fig. 7L). Isolated RGCs

tended to form clusters in culture. In contrast to directional axon growth found in CONCEPT

organoids, isolated RGCs grew neurites in random directions (Fig. 7L), indicating the

differences in axon pathfinding cues between CONCEPT organoids and purified RGC cultures.

RGC neurites were also marked by CNTN2 expression (Fig. 7M). A portion of isolated RGCs

expressed POU4F2 but not CNTN2 (Fig. 7M), indicating that CNTN2 was downregulated in

dissociated RGC cultures. Isolated RGCs also expressed RGC markers ISL1, RBPMS, and

SNCG (Fig. 7N-P). Collectively, we have developed a one-step method for the isolation of

differentiating human RGCs in a native condition.

Isolated RGCs exhibit electrophysiological signature of excitable cells

In order to determine the functional properties of isolated RGCs in culture, we examined their

electrophysiological properties using whole-cell patch clamp recordings (Fig. 8, *see Methods*).

Using current-clamp configuration, we found that RGCs displayed a hyperpolarized resting

membrane potential (*mean ±SD: -20.1 ±6.4mV*, Fig. 8A). When cells were held at -70 mV by

current injection, in most cases (6/9), depolarizing currents steps could trigger an action

potential, which was followed by a depolarization plateau if the current was injected for more

than 10ms (Fig. 8B). In voltage clamp recordings we examined the nature of the voltage-gated

conductances (*see Methods*). From a holding membrane potential of -80 mV, both inward and

outward currents (I_m) were observed in response to depolarizing voltage steps (V_m, Fig. 8C).

Outward currents were primarily mediated by voltage-gated potassium channels, as application

of 20mM TEA significantly reduced their amplitude (*mean I_K ±SD at 60mV; 964±537 pA in*

control; 279±201 pA in TEA, U=24.0; p=0.019, two-sided MannWhitneyU test, Fig. 8D).

Conversely, 1µM TTX abolished all inward currents, demonstrating that they were mediated by

activation of voltage-gated sodium channels (*mean I_{Na} ±SD at -10mV; -338.2±121 in control; -*

18.3±13 pA in TTX; U=0.0; p=0.035, two-sided MannWhitneyU test, Fig. 8E). Taken together,

these results indicate that isolated RGCs exhibit functional features traditionally found in

excitable cells, such as in neurons.

FGF signaling mediated by FGFRs is required for early RGC differentiation and

directional axon growth in human cells

Directional RGC axon growth toward and then along PAX2+ optic-disc cells in CONCEPT

organoids could be mediated by signaling molecules secreted from PAX2+ optic-disc cells.

scRNA-seq analysis identified expression signatures of cluster 2, the major component of

PAX2+ optic disc cells in CONCEPT organoids (Fig. 6; supp. Fig. S10). Interestingly, *FGF8* and

FGF9 were differentially expressed in cluster 2 (Figs. 4K, 4L; 9A; supp. Fig. S10). To assess the

roles of FGF8 and FGF9 in CONCEPT organoids, we validated that *FGF8* and *FGF9* were

differentially expressed in PAX2+ optic-disc cells of CONCEPT organoids (Fig. 9B-E). TUBB3+

RGC axons grew towards and then along the regions with high-level *FGF8* and *FGF9*

expression (Fig. 9D, E), suggesting that FGF8 and FGF9 may attract RGC axon growth.

Additionally, *FGFR1, FGFR2, FGFR3, MAP2K1,* and *MAP2K2* were expressed in multiple types

of cells in CONCEPT organoids. In RGCs, *FGFR1* and *MAP2K2* were clearly expressed (supp.

Fig. S16), indicating that RGCs expressed the components that transduce the signals from FGF

ligands. Since FGF8 and FGF9 are probably redundant and it is challenging to inactivate both

FGF8 and FGF9 in CONCEPT organoids, we decided to inactivate FGF signaling with

FGFR1/2/3 inhibitor PD 161570 during days 17-24, a stage at early RGC differentiation. After

the FGFR inhibition, FGF8 expression was grossly unaffected. Despite persistent FGF8, the

number of RGC somas were drastically reduced (Fig. 9F-9K). Notably, remaining RGCs nearly

did not grow directional axons (arrowheads in Fig. 9K), and a few remaining axons wandered

around (arrow in Fig. 9K). These findings indicate that a) PAX2+ optic-disc cells differentially

expressed FGF8 and FGF9; b) FGF signaling mediated by FGFRs is required for early RGC

differentiation and directional axon growth in human cells.

Discussion

In this study, we report the self-formation of concentric zones of telencephalic and ocular tissues in CONCEPT telencephalon-eye organoids from human pluripotent stem cells, establishing a model for studying the early development of telencephalic and ocular tissues in humans. RGCs grew axons toward and then along a path defined by PAX2+ cell populations, setting up a model for studying RGC axon growth and pathfinding. We identified expression signatures of cell clusters in CONCEPT organoids using single-cell RNA sequencing and revealed mechanisms of early RGC differentiation and axon growth. We demonstrated that CONCEPT organoids and human fetal retinas had similar expression signatures. Lastly, we established a one-step method for the isolation of human electrophysiologically-excitable RGCs via CNTN2 under a native condition. Our study not only provides deeper insight into coordinated specification of telencephalic and ocular tissues for directional RGC axon growth in humans, but also generates resources for therapeutic studies of RGC-related diseases such as glaucoma.

Cysts are hollow spheres of a columnar epithelium mimicking the anterior ectoderm

Cysts are hollow spheres of a columnar epithelium with the apical surface at the lumen. They are induced from pluripotent stem cells via embedding small sheets of hESCs into Matrigel and subsequent growth either in a solid thin film (Zhu et al., 2013) or suspensions (Kim et al., 2019; Lowe et al., 2016). Cyst growth in suspensions is cost-effective and scalable. The formation of cysts induced by Matrigel mimics the epithelization of the epiblast by the extracellular matrix (ECM) in the blastocyst (Bedzhov and Zernicka-Goetz, 2014; Coucouvanis and Martin, 1995).

In this study, individual cysts efficiently generate telencephalic and ocular tissues in the absence of any extrinsic factors, indicating their default cell fates of the anterior ectoderm. In literature, anterior neural tissues are generated from re-aggregated single pluripotent stem cells through

dual inhibition of Smad signaling (Chambers et al., 2009) or inhibition of Wnt/ß-catenin signaling

(Nakano et al., 2012). Besides that, undirected cultures without extrinsic factors are used for the

generation of retinal cultures (Fernando et al., 2022; Reichman et al., 2014; Zhong et al., 2014).

In our procedure, we did not add any extrinsic factors; the epithelial structure of cysts and

subsequent adherent growth at a low density facilitate neural induction and differentiation. Our

findings are consistent with the reports that epiblast cells are epigenetically primed for

ectodermal fates (Argelaguet et al., 2019) and neuroectodermal tissues are the default

differentiation from pluripotent stem cells (Munoz-Sanjuan and Brivanlou, 2002; Tropepe et al.,

2001). Collectively, cysts mimic the anterior ectoderm in the epithelial structure and cell fates.

Pattern formation and coordinated cell differentiation in CONCEPT telencephalon-eye

organoids

Tissue patterning is fundamental for the formation of a body plan, which is defined by the

anteroposterior, dorsoventral, and left-right axes. Self-formation of CONCEPT telencephalon-

eye organoids establishes an efficient way to pattern early telencephalic and ocular tissues

along a spatial axis. In CONCEPT organoids on days 22-26, FOXG1+ telencephalon, PAX2+

optic stalk, PAX2+ optic disc, and VSX2+ neuroretinal tissues are positioned along the center-

periphery axis, mimicking the sequence of their spatial positions in E13.5 mouse embryos. Our

findings indicate that the formation of telencephalic and retinal tissues are highly coordinated,

consistent with the prosomere model (Martinez-Morales et al., 2004; Redies and Puelles, 2001).

Collectively, we establish CONCEPT telencephalon-eye organoids for studying the early

formation of telencephalic and ocular tissues in humans.

The tissue patterning in CONCEPT telencephalon-eye organoids is originated from the

attachment of a radially-symmetric epithelium to the culture surface for growth as colonies since

individual cysts kept in suspensions do not form any apparent tissue patterning. When a floating

cyst initially contacted the culture surface, the contact resulted in ECM-cell adhesions. The initial

ECM-cell adhesions caused additional ECM-cell contacts in neighboring cells, resulting in the

flattening and spreading of a cyst onto the culture surface. Since the cyst is a radially-symmetric

epithelium, ECM-cell adhesions between the cyst and the culture surface formed sequentially in

a concentric manner. Remodeling of cell-cell and ECM-cell adhesions and self-organization

underlay the tissue patterning. Molecular events in concentric patterns during the attachment of

a cyst onto the culture surface were eventually translated to concentric gradients of morphogens

that specify cell fates. The radially-symmetric epithelial structure of cysts is important for the

formation of a concentric pattern since such pattern would not be generated when amorphous

embryoid bodies are attached to the culture surface for growth as colonies. In literature, timed

BMP4 treatment is shown to promote neuroretinal differentiation from pluripotent stem cells

(Kuwahara et al., 2015), and FGF8 promotes telencephalic and eye development (Martinez-

Morales et al., 2005; Wilson and Rubenstein, 2000). In our system, BMP4 and FGF8, along with

BMP7 and other FGFs, were highly expressed starting at early stages and gradually formed

concentric morphogen gradients, which would dictate tissue patterning, resulting in coordinated

specification of telencephalic, optic stalk, optic disc, and neuroretinal tissues along the center-

periphery axis in CONCEPT telencephalon-eye organoids. Co-culture of these tissues in the

concentric zones provides mutual cues for coordinated tissue specification and growth.

Nevertheless, anchorage culture of cell colonies prevents three-dimensional morphogenesis as

seen in embryos, impeding proper tissue interactions at more advanced stages. When these

cell colonies are detached using dispase for suspension culture, retinal progenitor cells self-

organize to form protruding, translucent spheres called retinal organoids whereas other types of

cells form structures with dark appearance; concentric zones no longer exist (Lowe et al., 2016).

It is conceivable that the simple suspension culture does not have proper physical and chemical

supports provided by periocular tissues *in vivo* and therefore optic-stalk cells do not form a tube-

like structure. It is a challenge ahead to reconstruct this three-dimensional environment to mimic

the morphogenesis *in vivo.*

Concentric patterns of stem cell-derived cultures are also reported in a few other experiments.

When dissociated single pluripotent stem cells are grown in micropatterned culture surface at

certain cell densities in a medium supplemented with BMP4, concentric zones of progenitors

expressing markers for trophectoderm, endoderm, mesoderm, and ectoderm are found (Etoc et

al., 2016; Minn et al., 2020). Cell density and colony geometry dictate cell fate specification from

pluripotent stem cells. A concentric gradient of BMP4 activity regulates the patterning (Etoc et

al., 2016). When dissociated single pluripotent stem cells are grown in a pre-patterned

geometrically-confined culture surface in a medium supplemented with dual inhibitors for TGF-β

and BMP4, concentric zones of progenitors expressing the markers for neural plate and neural

plate border are observed. Morphogenetic cues – cell shape and cytoskeletal contractile force –

dictate the patterning of the neural plate and neural plate border via BMP-SMAD signaling (Xue

et al., 2018). When dissociated single pluripotent stem cells are grown as individual colonies in

a medium supplemented with knockout serum replacement, multiple zones of ectodermal cells

autonomously form (Hayashi et al., 2016). In all three experiments, dissociated single cells are

used to generate cell colonies through either cell re-aggregation or proliferation. In our case,

adherent culture of a radially-symmetric epithelium – the cyst – was used to generate

CONCEPT organoids. Among the three previous and our experiments, starting cells differ in

their developmental potentials, tissue structures, and treatments. Despite the differences, all

these organoids comprise concentric zones of distinct cell populations. Formation of CONCEPT

telencephalon-eye organoids mimics the coordinated specification of early telencephalic and

ocular tissues in humans.

RGC axon pathfinding cues in CONCEPT telencephalon-eye organoids and humans

In the mouse retina, multiple RGC axon guidance cues are concentrically organized around the

optic disc, regulating RGC axon growth and exit from the eye through the optic stalk (Oster et al.,

2004). Early differentiated RGCs are in a short distance from the nascent optic disc, and axons

of later differentiated RGCs in more peripheral retinal regions follow the path of the initial axons.

It is accepted that the optic-disc regions provide growth-promoting guidance cues whereas

peripheral retinal regions provide inhibitory guidance cues (Oster et al., 2004). Pax2 is

specifically expressed in the ventral optic stalk, optic vesicles, central neuroretina, optic disc,

and optic stalk; Pax2 is essential for optic stalk and nerve development in mice (Macdonald et

al., 1997; Martinez-Morales et al., 2001; Torres et al., 1996).

In CONCEPT telencephalon-eye organoids, coordinated specification of telencephalic and

ocular tissues generates PAX2+ optic-disc and optic-stalk cells that define the path for RGC

axon growth. To the best of our knowledge, directional RGC axon growth guided by optic-disc

and optic-stalk tissues has not been reported.

Single-cell RNA sequencing of CONCEPT organoids identified DEGs of PAX2+ optic-disc and

optic-stalk cells. Interestingly, *FGF8* and *FGF9* were differentially expressed in PAX2+ optic-disc

cells; inhibition of FGF signaling with an FGFR inhibitor during early RGC differentiation

drastically decreased the number of RGC somas and nearly ablated directional axon growth,

indicating that early RGC differentiation and directional axon growth require FGFR-mediated

FGF signaling in human cells. In chicks, FGF8 and FGF3 coordinate the initiation of retinal

differentiation (Martinez-Morales et al., 2005; Vogel-Hopker et al., 2000); FGF8 maintains PAX2

expression in optic nerve explants (Soukkarieh et al., 2007). In mice, FGF8 and FGF9 trigger

axon outgrowth in motor neuron column explants (Shirasaki et al., 2006). Therefore, the roles of

FGF signaling in the regulation of early RGC differentiation and directional axon growth in

CONCEPT organoids are consistent with the literature. Dissecting DEGs of PAX2+ optic-disc

cells in CONCEPT telencephalon-eye organoids elucidates the mechanisms of RGC axon

growth and pathfinding in humans.

CONCEPT organoids and human fetal retinas shared expression signatures. For example,

assigned optic disc cells from both CONCEPT organoids and human fetal retinas differentially

expressed *COL9A3, SEMA5A,* and *FGF9,* which may be axon guidance cues for RGC axon

growth in humans. Additionally, DEGs of assigned optic disc cells from both CONCEPT

organoids and human fetal retinas were enriched for GO terms astrocyte development, neuron

projection development, and glial cells differentiation. Furthermore, RGC axon growth and

pathfinding guided by optic disc cells were active before astrocytes were generated in both

CONCEPT organoids and human fetal retinas. It is conceivable that CONCEPT organoids are a

testable model for studying the functions of axon guidance molecules that are identified using

scRNA-seq datasets.

One-step isolation of early human RGCs under a native condition for both organoids and

human fetal retinas

RGCs are degenerated in glaucoma, a major cause of vision impairment in developed countries.

Disease modeling and drug discovery to suppress RGC death will have a huge impact on

saving vision. The use of human RGC models is critical for therapeutic studies since humans

and rodents differ significantly in RGCs. Additionally, cell replacement therapies for glaucoma

are extensively evaluated. Therefore, efficient isolation of human RGCs in a native condition will

have a substantial impact on therapeutic studies of RGC-related diseases such as glaucoma. In

literature, cell surface marker Thy1 is used for the isolation of adult mouse RGCs, but it is

unsuccessful in the isolation of RGCs from 3D retinal organoids (Sluch et al., 2017). A study

reports the purification of human RGCs via Thy1 from adherent cultures (Rabesandratana et al.,

2020). Tagging human RGCs with an engineered marker Thy1.2 leads to efficient isolation of

human RGCs (Sluch et al., 2017), but may not be unsuitable for clinical uses. Using single-cell

RNA sequencing, we identified cell-surface protein CNTN2 as a specific marker for early RGCs

in both human organoids and fetal retinas. We isolated electrophysiologically-excitable RGCs

from retinal organoids using MACS via CNTN2, and it is conceivable that this method is

applicable for human fetal retinas. Therefore, we establish a one-step method for isolating early

human RGCs under a native condition, facilitating therapeutic studies for RGC-related retinal

diseases such as glaucoma.

Disclosure – A patent application is pending.

Author Contributions

WL conceived and led the project, performed experiments, analyzed data, drew conclusions,

and wrote the paper. RD performed experiments, analyzed data, and edited the paper. AL

performed experiments. XZ analyzed the single-cell RNA sequencing dataset. LS designed,

performed, and analyzed patch clamp experiments and wrote the section. All authors discussed

the results and contributed to the final manuscript.

Acknowledgements

We are grateful to Dr. Roy Chuck for support, M.R.G. Stem Cell Institute (supported by

NYSTEM C029154) for service, Leslie Cummins and Frank P Macaluso at the Analytical

Imaging Facility of AECOM for electron microscopy (supported by P30CA013330), David

Reynolds at the Genomics Core of AECOM for the preparation of single-cell libraries, Dr.

Kamran Khodakhah for technical support and critical readings, and grants from NEI, NIH

(R01EY022645 and R21EY029806 to W.L.). Albert Lowe was supported by training grants

T32GM007491 and C30292GG. Ludovic Spaeth was supported by a grant from NIH

(5R01NS105470-06 to Dr. Khodakhah).

 Materials and Methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
cell line (*Homo-sapiens*)	H1 hESCs	WiCell	WA01	
	hiPSCs	Corriell Institute	AICS 0023	
antibody	anti-FOXG1 (Rabbit polyclonal)	Abcam	Cat# ab18259	IF (1:500)
	anti-TUBB3 (mouse monoclonal)	Covance	Cat# MMS-435P	IF (1:1000)
	anti-FGF8 (mouse monoclonal)	R&D	Cat# MAB323	ICC (1:500)
	anti-RBPMS (rabbit polyclonal)	PhosphoSolution	Cat# 1830-RBPMS	IF (1:200)
	anti-ISL1 (mouse monoclonal)	DSHB	40.2D6	IF (1:500)

	anti-SNCG (rabbit polyclonal)	Abcam	Cat# ab55424	IF (1:200)
	anti-PAX2 (rabbit polyclonal)	Invitrogen	Cat# 716000	IF (1:200)
	anti- alpha A crystallin (rabbit polyclonal)	Santa Cruz sc	Cat# sc-22743	IF (1:500)
	Anti- beta crystallin (rabbit polyclonal)	Santa Cruz	Cat# sc-22745	IF (1:100)
	Anti-gamma crystallin (rabbit polyclonal)	Santa Cruz	Cat# sc-22746	Western (1:1000)
	Anti-CNTN2 (mouse monoclonal)	DSHB	Cat# 4D7	IF (1:100)
	Anti-ALDH1A3 (rabbit polyclonal)	Invitrogen	Cat# PA529188	IF (1:500)
	Anti-VAX1/2 (rabbit polyclonal)	Santa Cruz	Cat# sc-98613	IF (1:200)
	Anti-PAX6 (rabbit polyclonal)	Covance	Cat# PRB-278P	IF (1:500)
	Anti-POU4F2 (goat polyclonal)	Santa Cruz	Cat# SC-6026	IF (1:200)

	Anti-VSX2 (sheep polyclonal)	Millipore	Cat# AB9016	IF (1:500)
commercial assay or kit	DIG RNA Labeling Mix	Millipore Sigma	Cat# 11277073910	
	MagnaBind goat anti-mouse IgG beads	ThermoScientific	Cat# 21354	
sequence-based reagent	BMP4, forward	This paper	PCR primers	CGGAAGCT AGGTGAGT GTGG
	BMP4, reverse	This paper	PCR primers	GAGtaatacg actcactatagg gGGAAGCC CCTTTCCC AATCA
	BMP7, forward	This paper	PCR primers	gaggtccctctc cattccct
	BMP7, reverse	This paper	PCR primers	GAGtaatacg actcactatagg gtgcacccatca gacctccta
	FGF8, forward	This paper	PCR primers	GTTGCACT TGCTGGTC CTCT

	FGF8, reverse	This paper	PCR primers	GAGtaatacg actcactatagg gTTGAGTTT TGGGTGCC CTAC
	PAX2, forward	This paper	PCR primers	gctgtctgtgctgt gagagt
	PAX2, reverse	This paper	PCR primers	GAGtaatacg actcactatagg gccggggacatt tagcaggtt
	SEMA5A, forward	This paper	PCR primers	CAGAGGCT CAGGCACA ATGA
	SEMA5A, reverse	This paper	PCR primers	GAGtaatacg actcactatagg gTCCGTGT CTACCCAG GACTT
	CYP1B1, forward	This paper	PCR primers	cccagcggttctt catgagt
	CYP1B1, reverse	This paper	PCR primers	GAGtaatacg actcactatagg ggcacacttggtt gcgttagt
	LEFTY2, forward	This paper	PCR primers	agccctctaact gaacgtgtg

	LEFTY2, reverse	This paper	PCR primers	GAGtaatacg actcactatagg gtcttctgagtatc tacattcaattgct
	EMX2, forward	This paper	PCR primers	ACCGAGAA AGGGAGAG GGAA
	EMX2, reverse	This paper	PCR primers	GAGtaatacg actcactatagg gTCGGCCA ATTTCTCCA ACCA
	FGF9, forward	This paper	PCR primers	GTCCGCTA TGAACCTG TGGT
	FGF9, reverse	This paper	PCR primers	GAGtaatacg actcactatagg gATAGTCTC GCTTGCCC AAGG
chemical compound, drug	PD 161570	Tocris	Cat# 3724	1 µM
software, algorithm	Seurat	Stuart et al., 2019.	Seurat v3.2.0	
	CellRanger	10x Genomics	CellRanger (3.1.0)	

Maintenance of hESCs

ESCRO and IRB committees at AECOM approved the use of hESCs in this project.

Undifferentiated H1 hESCs (WiCell WA01) or hiPSCs (Corriell Institute AICS 0023) were grown

on Matrigel-coated 6-well plates in mTeSR1 medium and passaged using ReLeSR (STEMCELL

technologies) following manufacturer instructions.

Retinal cell differentiation

CONCEPT telencephalon-eye organoids were generated as follows. A humidified incubator at

37°C with 5% CO2 was used for cell culture. H1 hESCs or iPSCs that were passaged using

ReLeSR two or three days before experiments were detached using Dispase (GIBCO 17105041)

and then harvested using centrifugation. After that, the cell pellets were suspended in ice-cold

Matrigel. After gelling at 37 °C for 15-20 minutes, the hESC/Matrigel clump was gently

dispersed in a N2B27 Medium (DMEM/F12+GlutaMAX (GIBCO):Neurobasal medium (GIBCO)

= 1:1, 0.5 x B27 supplement (GIBCO), 0.5 x N2 supplement (GIBCO), 0.1 mM β-

mercaptoethanol, and 0.2 mM L-GlutaMax) for floating culture. With the starting day of cell

differentiation designated as day 0, cysts with a single lumen formed on day 1. Cysts with a

single lumen constituted over 85% of cultures. On day 4 or 5, individual cysts with a diameter at

around 150-200 μm were manually picked using a curved Pasteur pipet under an inverted

microscope and then seeded onto Matrigel-coated 24-well plates at a density of 2-6 cysts per

well. Cysts spontaneously attached to the culture surface and grew. From a time during days

13-16, attached cell colonies were grown in a KSR medium (GMEM medium supplemented with

10% knockout serum replacement, 1 mM sodium pyruvate, 0.1 mM non-essential amino acids,

2 mM l-glutamine, and 55 μM 2-mercaptoethanol; all the culture reagents were from Life

Technologies). Culture mediums were changed every two or three days. Overall, the procedure

is efficient and robust. Images in the manuscript represent 80-90% of cultures.

Inhibition of FGF signaling in CONCEPT telencephalon-eye organoids with FGFR

inhibitor PD 161570

To inactivate FGF signaling in CONCEPT organoids, FGFR1/2/3 inhibitor PD 161570 (1 µM;

Tocris) was supplemented to the culture medium starting on day 17, with vehicle DMSO as a

control. Treated CONCEPT organoids were harvested on day 24 for assays.

Magnetic-activated cell sorting (MACS) of developing human RGCs

Retinal organoids in suspension culture were generated using the established method (Kim et

al., 2019; Lowe et al., 2016). For each MACS experiment, 84 – 140 retinal organoids at stages

of day 41 – 70 were dissociated into single cells using Accutase (GIBCO A1110501). Non-

retinal cells were trimmed if there were any. Dissociated single cells were harvested using

centrifugation and then incubated for 25 minutes at room temperature with MagnaBind goat

anti-mouse IgG (ThermoScientific 21354) beads that were previously coupled with a CNTN2

antibody (DSHB 4D7) following manufacturer instructions. Cells bound to the beads were

isolated using a magnetic rack and then washed one time with the KSR medium supplemented

with antibiotic:Antimycotic (GEMINI 400101) while the tube was still against the magnetic rack.

After the wash, the cells were released from the beads via Accutase digestion for 30 minutes

and then harvested using centrifugation. The isolated cells were plated onto a chamber slide

(ibidi 80826, ibidiTreat µ-Slide 8 Well, coated with poly-ornithine and Matrigel, 30,000-50,000

cells/200 µl/well) in BrainPhys neuronal medium (Stem Cell Technology 05790) supplemented

with N2 and B27 (GIBCO 17502001, A3582801). From 100 retinal organoids on day 41 to 48,

around 385,000 RGCs were isolated. After 10 days of culture in chamber slides, RGCs were

fixed in 4% paraformaldehyde (PFA) for 10 minutes and then processed for immunostaining.

Immunostaining, antibodies, and light microscopy

CONCEPT telencephalon-eye organoids were fixed in 4% PFA for 15-30 minutes at room

temperature and processed for immunostaining. See Supplemental Information for details.

In situ **hybridization**

DIG-labeled anti-sense RNA probes for in situ hybridization were generated via in vitro

transcription using a DIG RNA labeling kit (Millipore Sigma-Aldrich 11175025910). See

Supplemental Information for details. To assess the *in vivo* expression of DEGs identified by

single-cell RNA sequencing, in situ hybridization images of their mouse orthologs in E14.5

mouse brains were downloaded from a public database (Visel et al., 2004) (https://gp3.mpg.de/)

with permission and then assembled in supplemental Figures.

Electron microscopy (EM)

EM was performed by Analytical Imaging Facility at Albert Einstein College of Medicine with a

standard method. Lens cell clusters were fixed in 0.1M Cacodylate buffer containing 2%

paraformaldehyde and 2.5% glutaraldehyde for 60 minutes at room temperature and then

processed for EM.

Single-cell RNA sequencing

CONCEPT telencephalon-eye organoids at day 24 (referred to as CR24) from one culture well

were dissociated into single cells using activated Papain (Worthington Biochemical Corporation

LS003126) following manufacturer's instructions. Then, 10,000 dissociated cells were captured

using Chromium Controller (10x Genomics), followed by library preparation using Single Cell 3'

version 3.1 kit (10x Genomics) per manufacturer's instructions. The library was sequenced in

one lane of Illumina HiSeq (2x150 bp) by GeneWIZ company.

Fastq sequences were mapped to the human genome (GRCh38-3.0.0) using CellRanger (3.1.0)

to generate a count matrix, which was further analyzed using the Seurat Package (v3.2.0)

(Stuart et al., 2019). Sequenced cells were filtered (nFeature_RNA > 200 & nFeature_RNA <

6000 & percent.mt < 20), resulting in 10218 cells. Dimension reduction and clustering were

performed using the following functions: *NormalizeData, FindVariableFeatures*

(selection.method = "vst", nfeatures = 2000), ScaleData, RunPCA, ElbowPlot, FindNeighbors

(dims = 1:17), FindClusters (resolution = 0.5), and *RunUMAP(dims = 1:17).* Differentially

expressed genes were identified using the function *FindAllMarkers (only.pos = FALSE, min.pct*

= 0.25, logfc.threshold = 0.25). Cells in human fetal retinas dataset HGW9 (GSE138002) were

also clustered using the Seurat package, with filtration of nFeature_RNA > 200 &

nFeature_RNA < 5000 & percent.mt < 5. For integration of the datasets of CONCEPT organoids

and human fetal retinas HGW9 (GSE138002), cells in cluster 18 and retinal progenitor clusters

from the HGW9 dataset were combined with cells in clusters 2, 4, 5, 7 from the CONCEPT

dataset for Seurat anchor-based clustering using functions *FindIntegrationAnchors* and

IntegrateData. The integrated dataset was then used for cell clustering. For enriched GO term

analysis, DEGs (top 200 genes) of cluster 2 in CONCEPT organoids and DEGs (113 genes) of

cluster 18 in human fetal retinas HGW9 were used for as inputs for running the function *gost* in

the gprofiler2 package (Kolberg et al., 2020).

Whole-cell patch clamp recordings of isolated RGCs in culture.

Standard methods were used. See Supplemental Information for details.

Statistical analysis

Statistical analysis in DEG identification was performed using the Seurat Package (v3.2.0)

(Stuart et al., 2019). Adjusted p-values were shown.

Data availability

Single-cell RNA sequencing data of CONCEPT telencephalon-eye organoids at day 24 is

available in Gene Expression Omnibus (http://www.ncbi.nlm.nih.gov/geo) using the accession

number GSE191017.

Reference:

Argelaguet, R., Clark, S.J., Mohammed, H., Stapel, L.C., Krueger, C., Kapourani, C.A., Imaz-
Rosshandler, I., Lohoff, T., Xiang, Y., Hanna, C.W.*, et al.* (2019). Multi-omics profiling of
mouse gastrulation at single-cell resolution. Nature *576*, 487-491.
Ashery-Padan, R., Marquardt, T., Zhou, X., and Gruss, P. (2000). Pax6 activity in the lens
primordium is required for lens formation and for correct placement of a single retina in the eye.
Genes Dev *14*, 2701-2711.
Bassett, E.A., Williams, T., Zacharias, A.L., Gage, P.J., Fuhrmann, S., and West-Mays, J.A.
(2010). AP-2alpha knockout mice exhibit optic cup patterning defects and failure of optic stalk
morphogenesis. Hum Mol Genet *19*, 1791-1804.
Baumer, N., Marquardt, T., Stoykova, A., Spieler, D., Treichel, D., Ashery-Padan, R., and Gruss,
P. (2003). Retinal pigmented epithelium determination requires the redundant activities of Pax2
and Pax6. Development *130*, 2903-2915.
Bedzhov, I., and Zernicka-Goetz, M. (2014). Self-organizing properties of mouse pluripotent
cells initiate morphogenesis upon implantation. Cell *156*, 1032-1044.
Bharti, K., Gasper, M., Ou, J., Brucato, M., Clore-Gronenborn, K., Pickel, J., and Arnheiter, H.
(2012). A regulatory loop involving PAX6, MITF, and WNT signaling controls retinal pigment
epithelium development. PLoS Genet *8*, e1002757.
Biswas, S.K., Lee, J.E., Brako, L., Jiang, J.X., and Lo, W.K. (2010). Gap junctions are
selectively associated with interlocking ball-and-sockets but not protrusions in the lens. Mol Vis
16, 2328-2341.
Chambers, S.M., Fasano, C.A., Papapetrou, E.P., Tomishima, M., Sadelain, M., and Studer, L.
(2009). Highly efficient neural conversion of human ES and iPS cells by dual inhibition of
SMAD signaling. Nat Biotechnol *27*, 275-280.
Chatterjee, M., and Li, J.Y. (2012). Patterning and compartment formation in the diencephalon.
Front Neurosci *6*, 66.
Chatzopoulou, E., Miguez, A., Savvaki, M., Levasseur, G., Muzerelle, A., Muriel, M.P.,
Goureau, O., Watanabe, K., Goutebroze, L., Gaspar, P.*, et al.* (2008). Structural requirement of
TAG-1 for retinal ganglion cell axons and myelin in the mouse optic nerve. J Neurosci *28*, 7624-
7636.
Coucouvanis, E., and Martin, G.R. (1995). Signals for death and survival: a two-step mechanism
for cavitation in the vertebrate embryo. Cell *83*, 279-287.
Cowan, C.S., Renner, M., De Gennaro, M., Gross-Scherf, B., Goldblum, D., Hou, Y., Munz, M.,
Rodrigues, T.M., Krol, J., Szikra, T.*, et al.* (2020). Cell Types of the Human Retina and Its
Organoids at Single-Cell Resolution. Cell *182*, 1623-1640 e1634.
Danesh, S.M., Villasenor, A., Chong, D., Soukup, C., and Cleaver, O. (2009). BMP and BMP
receptor expression during murine organogenesis. Gene Expr Patterns *9*, 255-265.
Dudley, A.T., and Robertson, E.J. (1997). Overlapping expression domains of bone
morphogenetic protein family members potentially account for limited tissue defects in BMP7
deficient embryos. Dev Dyn *208*, 349-362.
Eiraku, M., Takata, N., Ishibashi, H., Kawada, M., Sakakura, E., Okuda, S., Sekiguchi, K.,
Adachi, T., and Sasai, Y. (2011). Self-organizing optic-cup morphogenesis in three-dimensional
culture. Nature *472*, 51-56.
Erskine, L., and Herrera, E. (2007). The retinal ganglion cell axon's journey: insights into
molecular mechanisms of axon guidance. Dev Biol *308*, 1-14.

Erskine, L., and Herrera, E. (2014). Connecting the retina to the brain. ASN Neuro *6*.

Etoc, F., Metzger, J., Ruzo, A., Kirst, C., Yoney, A., Ozair, M.Z., Brivanlou, A.H., and Siggia,

E.D. (2016). A Balance between Secreted Inhibitors and Edge Sensing Controls Gastruloid Self-

Organization. Developmental cell *39*, 302-315.

Fernando, M., Lee, S., Wark, J.R., Xiao, D., Lim, B.Y., O'Hara-Wright, M., Kim, H.J., Smith,

G.C., Wong, T., Teber, E.T.*, et al.* (2022). Differentiation of brain and retinal organoids from

confluent cultures of pluripotent stem cells connected by nerve-like axonal projections of optic

origin. Stem Cell Reports *17*, 1476-1492.

Fligor, C.M., Langer, K.B., Sridhar, A., Ren, Y., Shields, P.K., Edler, M.C., Ohlemacher, S.K.,

Sluch, V.M., Zack, D.J., Zhang, C.*, et al.* (2018). Three-Dimensional Retinal Organoids

Facilitate the Investigation of Retinal Ganglion Cell Development, Organization and Neurite

Outgrowth from Human Pluripotent Stem Cells. Sci Rep *8*, 14520.

Fligor, C.M., Lavekar, S.S., Harkin, J., Shields, P.K., VanderWall, K.B., Huang, K.C., Gomes,

C., and Meyer, J.S. (2021). Extension of retinofugal projections in an assembled model of human

pluripotent stem cell-derived organoids. Stem Cell Reports.

Furuta, Y., and Hogan, B.L. (1998). BMP4 is essential for lens induction in the mouse embryo.

Genes Dev *12*, 3764-3775.

Gabriel, E., Albanna, W., Pasquini, G., Ramani, A., Josipovic, N., Mariappan, A., Schinzel, F.,

Karch, C.M., Bao, G., Gottardo, M.*, et al.* (2021). Human brain organoids assemble functionally

integrated bilateral optic vesicles. Cell Stem Cell.

Gan, L., Xiang, M., Zhou, L., Wagner, D.S., Klein, W.H., and Nathans, J. (1996). POU domain

factor Brn-3b is required for the development of a large set of retinal ganglion cells. Proc Natl

Acad Sci U S A *93*, 3920-3925.

Hayashi, R., Ishikawa, Y., Sasamoto, Y., Katori, R., Nomura, N., Ichikawa, T., Araki, S., Soma,

T., Kawasaki, S., Sekiguchi, K.*, et al.* (2016). Co-ordinated ocular development from human iPS

cells and recovery of corneal function. Nature *531*, 376-380.

Horsford, D.J., Nguyen, M.T., Sellar, G.C., Kothary, R., Arnheiter, H., and McInnes, R.R.

(2005). Chx10 repression of Mitf is required for the maintenance of mammalian neuroretinal

identity. Development *132*, 177-187.

Inoue, T., Nakamura, S., and Osumi, N. (2000). Fate mapping of the mouse prosencephalic

neural plate. Dev Biol *219*, 373-383.

Kadoshima, T., Sakaguchi, H., Nakano, T., Soen, M., Ando, S., Eiraku, M., and Sasai, Y. (2013).

Self-organization of axial polarity, inside-out layer pattern, and species-specific progenitor

dynamics in human ES cell-derived neocortex. Proc Natl Acad Sci U S A *110*, 20284-20289.

Kim, S., Lowe, A., Dharmat, R., Lee, S., Owen, L.A., Wang, J., Shakoor, A., Li, Y., Morgan,

D.J., Hejazi, A.A.*, et al.* (2019). Generation, transcriptome profiling, and functional validation of

cone-rich human retinal organoids. Proc Natl Acad Sci U S A *116*, 10824-10833.

Kolberg, L., Raudvere, U., Kuzmin, I., Vilo, J., and Peterson, H. (2020). gprofiler2 -- an R

package for gene list functional enrichment analysis and namespace conversion toolset g:Profiler.

F1000Res *9*.

Kuwahara, A., Ozone, C., Nakano, T., Saito, K., Eiraku, M., and Sasai, Y. (2015). Generation of

a ciliary margin-like stem cell niche from self-organizing human retinal tissue. Nature

communications *6*, 6286.

Lancaster, M.A., and Knoblich, J.A. (2014). Organogenesis in a dish: modeling development and

disease using organoid technologies. Science *345*, 1247125.

Lancaster, M.A., Renner, M., Martin, C.A., Wenzel, D., Bicknell, L.S., Hurles, M.E., Homfray,
T., Penninger, J.M., Jackson, A.P., and Knoblich, J.A. (2013). Cerebral organoids model human
brain development and microcephaly. Nature *501*, 373-379.
Langer, K.B., Ohlemacher, S.K., Phillips, M.J., Fligor, C.M., Jiang, P., Gamm, D.M., and Meyer,
J.S. (2018). Retinal Ganglion Cell Diversity and Subtype Specification from Human Pluripotent
Stem Cells. Stem Cell Reports *10*, 1282-1293.
Li, H., Wagner, E., McCaffery, P., Smith, D., Andreadis, A., and Drager, U.C. (2000). A retinoic
acid synthesizing enzyme in ventral retina and telencephalon of the embryonic mouse. Mech Dev
95, 283-289.
Liu, W., Lagutin, O., Swindell, E., Jamrich, M., and Oliver, G. (2010). Neuroretina specification
in mouse embryos requires Six3-mediated suppression of Wnt8b in the anterior neural plate. J
Clin Invest *120*, 3568-3577.
Lovicu, F.J., and McAvoy, J.W. (2001). FGF-induced lens cell proliferation and differentiation is
dependent on MAPK (ERK1/2) signalling. Development *128*, 5075-5084.
Lowe, A., Harris, R., Bhansali, P., Cvekl, A., and Liu, W. (2016). Intercellular Adhesion-
Dependent Cell Survival and ROCK-Regulated Actomyosin-Driven Forces Mediate Self-
Formation of a Retinal Organoid. Stem Cell Reports *6*, 743-756.
Lu, Y., Shiau, F., Yi, W., Lu, S., Wu, Q., Pearson, J.D., Kallman, A., Zhong, S., Hoang, T., Zuo,
Z.*, et al.* (2020). Single-Cell Analysis of Human Retina Identifies Evolutionarily Conserved and
Species-Specific Mechanisms Controlling Development. Developmental cell.
Macdonald, R., Scholes, J., Strahle, U., Brennan, C., Holder, N., Brand, M., and Wilson, S.W.
(1997). The Pax protein Noi is required for commissural axon pathway formation in the rostral
forebrain. Development *124*, 2397-2408.
Mariani, J., Simonini, M.V., Palejev, D., Tomasini, L., Coppola, G., Szekely, A.M., Horvath,
T.L., and Vaccarino, F.M. (2012). Modeling human cortical development in vitro using induced
pluripotent stem cells. Proc Natl Acad Sci U S A *109*, 12770-12775.
Marquardt, T., Ashery-Padan, R., Andrejewski, N., Scardigli, R., Guillemot, F., and Gruss, P.
(2001). Pax6 is required for the multipotent state of retinal progenitor cells. Cell *105*, 43-55.
Martinez-Ferre, A., and Martinez, S. (2012). Molecular regionalization of the diencephalon.
Front Neurosci *6*, 73.
Martinez-Morales, J.R., Del Bene, F., Nica, G., Hammerschmidt, M., Bovolenta, P., and
Wittbrodt, J. (2005). Differentiation of the vertebrate retina is coordinated by an FGF signaling
center. Developmental cell *8*, 565-574.
Martinez-Morales, J.R., Rodrigo, I., and Bovolenta, P. (2004). Eye development: a view from the
retina pigmented epithelium. Bioessays *26*, 766-777.
Martinez-Morales, J.R., Signore, M., Acampora, D., Simeone, A., and Bovolenta, P. (2001). Otx
genes are required for tissue specification in the developing eye. Development *128*, 2019-2030.
Meyer, J.S., Howden, S.E., Wallace, K.A., Verhoeven, A.D., Wright, L.S., Capowski, E.E.,
Pinilla, I., Martin, J.M., Tian, S., Stewart, R.*, et al.* (2011). Optic vesicle-like structures derived
from human pluripotent stem cells facilitate a customized approach to retinal disease treatment.
Stem Cells *29*, 1206-1218.
Minn, K.T., Fu, Y.C., He, S., Dietmann, S., George, S.C., Anastasio, M.A., Morris, S.A., and
Solnica-Krezel, L. (2020). High-resolution transcriptional and morphogenetic profiling of cells
from micropatterned human ESC gastruloid cultures. Elife *9*.

Morcillo, J., Martinez-Morales, J.R., Trousse, F., Fermin, Y., Sowden, J.C., and Bovolenta, P.
(2006). Proper patterning of the optic fissure requires the sequential activity of BMP7 and SHH.
Development *133*, 3179-3190.
Mui, S.H., Kim, J.W., Lemke, G., and Bertuzzi, S. (2005). Vax genes ventralize the embryonic
eye. Genes Dev *19*, 1249-1259.
Munoz-Sanjuan, I., and Brivanlou, A.H. (2002). Neural induction, the default model and
embryonic stem cells. Nat Rev Neurosci *3*, 271-280.
Nakano, T., Ando, S., Takata, N., Kawada, M., Muguruma, K., Sekiguchi, K., Saito, K.,
Yonemura, S., Eiraku, M., and Sasai, Y. (2012). Self-formation of optic cups and storable
stratified neural retina from human ESCs. Cell Stem Cell *10*, 771-785.
Oster, S.F., Bodeker, M.O., He, F., and Sretavan, D.W. (2003). Invariant Sema5A inhibition
serves an ensheathing function during optic nerve development. Development *130*, 775-784.
Oster, S.F., Deiner, M., Birgbauer, E., and Sretavan, D.W. (2004). Ganglion cell axon
pathfinding in the retina and optic nerve. Semin Cell Dev Biol *15*, 125-136.
Poh, Y.C., Chen, J., Hong, Y., Yi, H., Zhang, S., Chen, J., Wu, D.C., Wang, L., Jia, Q., Singh, R.*,
et al.* (2014). Generation of organized germ layers from a single mouse embryonic stem cell. Nat
Commun *5*, 4000.
Qian, X., Nguyen, H.N., Jacob, F., Song, H., and Ming, G.L. (2017). Using brain organoids to
understand Zika virus-induced microcephaly. Development *144*, 952-957.
Rabesandratana, O., Chaffiol, A., Mialot, A., Slembrouck-Brec, A., Joffrois, C., Nanteau, C.,
Rodrigues, A., Gagliardi, G., Reichman, S., Sahel, J.A.*, et al.* (2020). Generation of a
Transplantable Population of Human iPSC-Derived Retinal Ganglion Cells. Front Cell Dev Biol
8, 585675.
Redies, C., and Puelles, L. (2001). Modularity in vertebrate brain development and evolution.
Bioessays *23*, 1100-1111.
Reichman, S., Terray, A., Slembrouck, A., Nanteau, C., Orieux, G., Habeler, W., Nandrot, E.F.,
Sahel, J.A., Monville, C., and Goureau, O. (2014). From confluent human iPS cells to self-
forming neural retina and retinal pigmented epithelium. Proc Natl Acad Sci U S A *111*, 8518-
8523.
Rowan, S., Chen, C.M., Young, T.L., Fisher, D.E., and Cepko, C.L. (2004). Transdifferentiation
of the retina into pigmented cells in ocular retardation mice defines a new function of the
homeodomain gene Chx10. Development *131*, 5139-5152.
Sandberg-Lall, M., Hagg, P.O., Wahlstrom, I., and Pihlajaniemi, T. (2000). Type XIII collagen is
widely expressed in the adult and developing human eye and accentuated in the ciliary muscle,
the optic nerve and the neural retina. Exp Eye Res *70*, 401-410.
Schwarz, M., Cecconi, F., Bernier, G., Andrejewski, N., Kammandel, B., Wagner, M., and Gruss,
P. (2000). Spatial specification of mammalian eye territories by reciprocal transcriptional
repression of Pax2 and Pax6. Development *127*, 4325-4334.
Shimamura, K., and Rubenstein, J.L. (1997). Inductive interactions direct early regionalization of
the mouse forebrain. Development *124*, 2709-2718.
Shirasaki, R., Lewcock, J.W., Lettieri, K., and Pfaff, S.L. (2006). FGF as a target-derived
chemoattractant for developing motor axons genetically programmed by the LIM code. Neuron
50, 841-853.
Simoes-Costa, M., and Bronner, M.E. (2015). Establishing neural crest identity: a gene
regulatory recipe. Development *142*, 242-257.

Sluch, V.M., Chamling, X., Liu, M.M., Berlinicke, C.A., Cheng, J., Mitchell, K.L., Welsbie,
D.S., and Zack, D.J. (2017). Enhanced Stem Cell Differentiation and Immunopurification of
Genome Engineered Human Retinal Ganglion Cells. Stem Cells Transl Med *6*, 1972-1986.
Solloway, M.J., Dudley, A.T., Bikoff, E.K., Lyons, K.M., Hogan, B.L., and Robertson, E.J.
(1998). Mice lacking Bmp6 function. Dev Genet *22*, 321-339.
Soukkarieh, C., Agius, E., Soula, C., and Cochard, P. (2007). Pax2 regulates neuronal-glial cell
fate choice in the embryonic optic nerve. Dev Biol *303*, 800-813.
Storm, E.E., Garel, S., Borello, U., Hebert, J.M., Martinez, S., McConnell, S.K., Martin, G.R.,
and Rubenstein, J.L. (2006). Dose-dependent functions of Fgf8 in regulating telencephalic
patterning centers. Development *133*, 1831-1844.
Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., 3rd, Hao, Y.,
Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive Integration of Single-Cell Data.
Cell *177*, 1888-1902 e1821.
Take-uchi, M., Clarke, J.D., and Wilson, S.W. (2003). Hedgehog signalling maintains the optic
stalk-retinal interface through the regulation of Vax gene activity. Development *130*, 955-968.
Teotia, P., Chopra, D.A., Dravid, S.M., Van Hook, M.J., Qiu, F., Morrison, J., Rizzino, A., and
Ahmad, I. (2017). Generation of Functional Human Retinal Ganglion Cells with Target
Specificity from Pluripotent Stem Cells by Chemically Defined Recapitulation of Developmental
Mechanism. Stem Cells *35*, 572-585.
Tirosh, I., Izar, B., Prakadan, S.M., Wadsworth, M.H., 2nd, Treacy, D., Trombetta, J.J., Rotem,
A., Rodman, C., Lian, C., Murphy, G.*, et al.* (2016). Dissecting the multicellular ecosystem of
metastatic melanoma by single-cell RNA-seq. Science *352*, 189-196.
Torres, M., Gomez-Pardo, E., and Gruss, P. (1996). Pax2 contributes to inner ear patterning and
optic nerve trajectory. Development *122*, 3381-3391.
Tropepe, V., Hitoshi, S., Sirard, C., Mak, T.W., Rossant, J., and van der Kooy, D. (2001). Direct
neural fate specification from embryonic stem cells: a primitive mammalian neural stem cell
stage acquired through a default mechanism. Neuron *30*, 65-78.
Varga, B.V., Faiz, M., Yang, H., Pivonkova, H., Gao, S., Khelifi, G., Linderoth, E., Zhen, M.,
Hussein, S.M., and Nagy, A. (2021). Signal requirement for cortical potential of transplantable
human neuroepithelial stem cells. biorxiv *https://doi.org/10.1101/2021.03.27.437311*.
Velasco, S., Kedaigle, A.J., Simmons, S.K., Nash, A., Rocha, M., Quadrato, G., Paulsen, B.,
Nguyen, L., Adiconis, X., Regev, A.*, et al.* (2019). Individual brain organoids reproducibly form
cell diversity of the human cerebral cortex. Nature *570*, 523-527.
Visel, A., Thaller, C., and Eichele, G. (2004). GenePaint.org: an atlas of gene expression patterns
in the mouse embryo. Nucleic Acids Res *32*, D552-556.
Vogel-Hopker, A., Momose, T., Rohrer, H., Yasuda, K., Ishihara, L., and Rapaport, D.H. (2000).
Multiple functions of fibroblast growth factor-8 (FGF-8) in chick eye development. Mech Dev
94, 25-36.
Wilson, S.W., and Rubenstein, J.L. (2000). Induction and dorsoventral patterning of the
telencephalon. Neuron *28*, 641-651.
Xuan, S., Baptista, C.A., Balas, G., Tao, W., Soares, V.C., and Lai, E. (1995). Winged helix
transcription factor BF-1 is essential for the development of the cerebral hemispheres. Neuron *14*,
1141-1152.
Xue, X., Sun, Y., Resto-Irizarry, A.M., Yuan, Y., Aw Yong, K.M., Zheng, Y., Weng, S., Shao,
Y., Chai, Y., Studer, L.*, et al.* (2018). Mechanics-guided embryonic patterning of neuroectoderm
tissue from human pluripotent stem cells. Nat Mater *17*, 633-641.

Yamagata, M., and Sanes, J.R. (2012). Expanding the Ig superfamily code for laminar specificity
in retina: expression and role of contactins. J Neurosci *32*, 14402-14414.
Yao, Z., Mich, J.K., Ku, S., Menon, V., Krostag, A.R., Martinez, R.A., Furchtgott, L.,
Mulholland, H., Bort, S., Fuqua, M.A.*, et al.* (2017). A Single-Cell Roadmap of Lineage
Bifurcation in Human ESC Models of Embryonic Brain Development. Cell Stem Cell *20*, 120-
134.
Zhong, X., Gutierrez, C., Xue, T., Hampton, C., Vergara, M.N., Cao, L.H., Peters, A., Park, T.S.,
Zambidis, E.T., Meyer, J.S.*, et al.* (2014). Generation of three-dimensional retinal tissue with
functional photoreceptors from human iPSCs. Nature communications *5*, 4047.
Zhu, Y., Carido, M., Meinhardt, A., Kurth, T., Karl, M.O., Ader, M., and Tanaka, E.M. (2013).
Three-dimensional neuroepithelial culture from human embryonic stem cells and its use for
quantitative conversion to retinal pigment epithelium. PLoS One *8*, e54552.
Zuber, M.E., Gestri, G., Viczian, A.S., Barsacchi, G., and Harris, W.A. (2003). Specification of
the vertebrate eye by a network of eye field transcription factors. Development *130*, 5155-5167.
991

 Figure legends

Figure 1. Generation of telencephalon-eye organoids comprising <u>con</u>centric zones of

anterior <u>e</u>ctodermal <u>p</u>rogeni<u>t</u>ors (CONCEPT). (A) A scheme of the procedure. **(B)** Diagrams

of developing CONCEPT organoids showing concentric zones of the anterior ectodermal

progenitors. A summary of Figs.1, 2, 7, S1-S3, S15. **(C)** Morphology of cysts at day 2 showing

the epithelial structure indicated by apical localization of the reporter TJ::GFP at the lumen. **(D)**

Morphology of CONCEPT organoids at day 26. **(E-G)** Expression of telencephalon (Tel) marker

Foxg1, neuroretinal (NR) markers Vsx2 and Pax6 in mouse eyes at E10-10.5. Rostral optic stalk

(OS) connected the telencephalic vesicle to the optic cup. **(H-O)** FOXG1+ telencephalic

progenitors, VSX2+ and/or PAX6+ retinal progenitors formed concentric zones in CONCEPT

organoids. N > 5 experiments. **(P-W)** In CONCEPT organoids, morphogens FGF8, BMP4, and

BMP7 mRNA expression started at early stages and subsequently formed circular gradients. N >

5 experiments. Scale bars, 100 μm (C, E, M, O, P, S, T), 200 μm (I, K), 500 μm (Q, U), 1 mm (D,

H, J, L, N, R, V, W).

Fig. 2. Retinal ganglion cells (RGCs) grow axons toward and then along a path defined by

PAX2+ cell populations in CONCEPT telencephalon-eye organoids. N > 5 experiments. **(A-**

D) POU4F2+ RGCs grew TUBB3+ axons toward and then along a path with a circular or a

portion of circular shape. **(E, F)** In mice, Pax2 was expressed in central regions of the retina and

optic stalk at E10.5 (E) and in the optic disc and optic stalk at E13.5 (F). Tubb3+ axons from the

initial RGCs grew toward the optic disc, exited the eye, and navigated within the optic stalk (G).

(H-L) In CONCEPT organoids at day 26, TUBB3+ RGC axons grew toward and then along a

path defined by an adjacent PAX2+ VSX2+ cell population (arrowhead in H, brackets in I, J); the

PAX2+ VSX2- cell population sets up an inner boundary of RGC axon growth. **(L)** A diagram

summarizing RGC axon growth, PAX2+ optic disc (OD), and PAX2+ optic stalk (OS) in

CONCEPT organoids. The area labeled by the asterisk may appear as false signals in a low-

resolution printout but it is clearly a background in digital display. **(M)** A count of CONCEPT

organoids showing directional retinal ganglion cell axons. Scale bars, 50 µm (E, F, G), 100 µm

(B), 200 µm (A, H, I).

Fig. 3. CONCEPT telencephalon-eye organoids contain lens cells that undergo terminal

differentiation. N > 5 experiments. **(A-L)** In CONCEPT organoids, lens markers CRYAA and

beta crystallin (CRY B) were expressed at day 22 (A, B) and day 39 (C, D, K; a count in H).

Lens cells were not stained by DAPI (E, F); they exhibited a crystal-like shape (G). When

CONCEPT organoids were detached using Dispase at around day 28 and grown in suspension,

crystal-like clusters, named as lentoid bodies, were found (I) and survived for months (J). These

lentoid bodies highly expressed gamma crystallin (CRY G), as revealed by Western blot (L). **(M-**

N) These lentoid bodies were free of organelles and exhibited ball-and-socket structures (K, L),

as revealed by electron microscopy. Scale bars, 100 µm (A, B, C, D, I), 200 µm (J), 5 µm (K),

200 nm (L).

Fig. 4. scRNA-seq of CONCEPT organoids identifies telencephalic and ocular cells,

including PAX2+ VSX2+ optic disc cells, PAX2+ VSX2- optic stalk cells, and CNTN2+

RGCs. CONCEPT organoids at day 24 were used for profiling. **(A)** Identification of 14 cell

clusters. **(B)** Cell cycle phases revealed by cell cycle scores. **(C)** *FOXG1* expression marked

telencephalic cells. **(D, E)** The expression of *PAX6* and/or *VSX2* marked retinal cells. **(F)** PAX2+

cells were found in two major cell populations: PAX2+ VSX2+ cells were assigned as the optic

disc (OD), whereas PAX2+ VSX2- FOXG1+ cells were assigned as the optic stalk (OS). **(G-L)**

The expression of major DEGs in cluster 2, the major cell population that mimics the optic disc.

(M-P) The expression of major gene markers for PAX2+ VSX2- optic stalk cells. **(Q-T)**

Identification of CNTN2 as a specific marker for early human RGCs. A large portion of cluster 11

differentially expressed neurogenic retinal progenitor marker *ATOH7* and RGC markers

POU4F2 and *SNCG.* The expression of *CNTN2* and *POU4F2* largely overlapped. **(U-X)** Two

small portions of cluster 11 differentially expressed early photoreceptor cell markers (*OTX2* and

CRX, U, V) and amacrine/horizontal cell markers (*TFAP2C* and *PTF1A,* W, X), respectively.

Fig. 5. Expression signatures of human fetal retinas HGW9 are similar to those of

CONCEPT organoids. (A) Cell clustering of human fetal retinas HGW9 (GSE138002) identified

21 clusters. **(B-F)** Cluster 18 differentially expressed *PAX2, COL9A3, CYP1B1, SEMA5A,* and

FGF9, which were top DEGs of cluster 2 in CONCEPT organoids. **(G)** *VSX2* expression marked

retinal progenitor cells. **(H-M)** Identification of neurogenic retinal progenitor cells (H), early

photoreceptor cells (I), amacrine/horizontal cells (J, K), and early RGCs (L, M). *POU4F2* and

CNTN2 were largely co-expressed in early RGCs, consistent with their expression profiles in

CONCEPT organoids. **(N-R)** When cells in cluster 18 and retinal progenitors from HGW9 were

combined with cells in clusters 2, 4, 5, 7 from CONCEPT organoids (CR24) for Seurat anchor-

based clustering, cells in cluster 18 from HGW9 (H18) were grouped with cluster 2 from

CONCEPT organoids (C2, assigned optic disc; N), and these cells expressed both *PAX2* and

VSX2 (arrowheads in N-R). A small portion of H18 cells were grouped with C4 cells (assigned

optic stalk; N), and these cells expressed *PAX2* but not *VSX2* (arrows in N-R).

Fig. 6. Comparisons of enriched GO terms in DEGs (top 200 genes) of cluster 2 in

CONCEPT organoids and DEGs (113 genes) of cluster 18 in human fetal retinas HGW9. A

large number of GO terms were enriched in both samples.

Fig. 7. RGCs grow CNTN2+ axons toward and then along a defined path in CONCEPT

telencephalon-eye organoids and can be isolated in one step via CNTN2 in a native

condition. N > 5 experiments. **(A-F)** PAX2+ cells formed two concentric zones mimicking the

optic stalk (OS) and optic disc (OD), respectively (A-C; high magnifications in D, E). POU4F2+

RGCs grew CNTN2+ axons toward and then along a path defined by adjacent PAX2+ optic-disc

cells (A-E). RGCs at a few hundreds of micrometers away from PAX2+ optic-disc cells grew

axons centrifugally (arrow in C). At regions where there was a gap in PAX2+ optic-disc cells,

CNTN2+ RGC axons exited the circular path and grew centrifugally (diamond arrowhead in A,

double arrowheads in B and F). PAX2+ optic-stalk cells set up an inner boundary for RGC axon

growth. **(G, H)** PAX2+ optic-disc cells did not express ALDH1A3; the cells that set up the

boundaries of the path highly expressed ALDH1A3. **(I, J)** Cells that set up the inner boundary

for RGC axon growth expressed VAX1/VAX2 (the antibody recognizes both VAX1 and VAX2).

(K) In E13.5 mouse eye, Aldh1a3 expression was high in the peripheral retina and was low or

nearly absent in the central retina. **(L-P)** One-step isolation of RGCs. RGCs from floating retinal

organoids at day 41 (L, M) and day 70 (N-O) were dissociated into single cells using Accutase

and then isolated using MACS via CNTN2 for 10-day growth. Isolated RGCs expressed

POU4F2 and grew TUBB3+ neurites in random directions (L). RGCs also expressed CNTN2

(M), ISL1 (N), RBPMS (N), and SNCG (O); positive cells were counted (P). Scale bars, 200 µm

(A,G,I), 100 µm (D-F,H,J,K), 50 µm (L).

Fig. 8. Electrophysiological features of RGCs. RGCs from retinal organoids on day 48 were

isolated using MACS via a CNTN2 antibody and then grown on polymer coverslips in a chamber

slide for 20-25 days before whole-cell patch clamp recordings. **(A)** Resting membrane potential

of RGCs. *Black triangle: average; black line: median; box: interquartile range, n=9.* **(B)** RGCs

can fire action potentials. Cells were patched in current-clamp mode. A steady current (I_m) was

injected to maintain the membrane potential at -70mV and depolarizing current steps of 10ms

(*left*), 100ms (*middle*) or 1s (*right*) were injected to elicit action potentials (V_m). **(C)** RGCs show

functional voltage-gated currents. Cells were recorded in voltage-clamp (holding=-80mV) and

depolarizing voltage steps (200ms, +10mV steps up to 80mV, V_m) were applied to record inward

and outward voltage-gated currents (I_m). *Inset*: zoom on inward currents. **(D)** Outward currents

are primarily due to voltage-gated potassium channels. *Left*, representative example of a

current-voltage experiment performed in presence of 20mM Tetraethylammonium (TEA), a

blocker of voltage-gated potassium channels. *Inset*: zoom on inward currents. *Right*, amplitude

of potassium current as a function of membrane potential (*mean±SEM; $n_{control}$=5, n_{TEA}=5*). **(E)**

Inward currents result from activity of voltage-gated sodium channels. *Left*, representative

example of a current-voltage experiment performed in presence of 1µM Tetrodotoxin (TTX), a

blocker of voltage-gated sodium channels. *Inset*: zoom on inward currents. *Right*, amplitude of

sodium current as a function of membrane potential (*mean±SEM; $n_{control}$=5, n_{TTX}=3*).

Fig. 9. FGF signaling mediated by FGFRs is required for early RGC differentiation and

directional axon growth. (A) *PAX2, FGF9,* and *FGF8* are differentially expressed in cluster 2,

the major component of PAX2+ optic-disc cells. **(B)** *PAX2* mRNA expression in CONCEPT

organoids on day 25. Two PAX2+ concentric zones corresponding to the optic stalk (OS) and

optic disc (OD) are labeled. **(C)** Dual-color immunocytochemistry indicates the co-localization of

FGF8 and PAX2 in the optic-disc zone of CONCEPT organoids on day 25. **(D-E)** TUBB3+

axons grew towards and then along the cells that expressed high levels of *FGF8* (D) and *FGF9*

mRNA (E) in CONCEPT organoids on day 25. Immunocytochemistry of TUBB3 was performed

after *in situ* hybridization. **(F-K)** After the inhibition of FGF signaling with FGFR inhibitor PD

161570 during days 17-24, FGF8 expression still largely remained, but the number of RGC

somas drastically reduced (J, K). Notably, remaining RGCs nearly did not grow directional

axons (arrowheads in K), and a few remaining axons wandered around (arrow in K). CNTN2

immunocytochemistry before (F, H) and after FGF8 immunocytochemistry (G, I, J, K) are shown.

N = 3/3 experiments. Scale bar, 250 µm (B), 100 µm (C-E), 1 mm (F), 200 µm (K).

1112

1113   **Tab. 1.** Cell counts for clusters in the scRNA-seq dataset of CONCEPT organoids at day 24.

Clusters	0	1	2	3	4	5	6	7	8	9	10	11	12	13
Idents	Mixed (lower counts)	dTel	OD	vTel	dTel/ OS	NR	RPE	NR	dTel/ OS	dTel/ OS	vTel	RGC/ PR/ AC/ HC	vTel	UD
# Cells	1295	1158	964	931	921	871	747	743	729	603	450	393	333	80
percent	0.127	0.113	0.094	0.091	0.09	0.085	0.073	0.073	0.071	0.059	0.044	0.038	0.033	0.008

1114

1115   Abbreviations: Idents, assigned cell identities; # Cells, cell number; Tel, telencephalon; NR,

1116   neural retina; OD, optic disc; OS, optic stalk; RPE, retinal pigment epithelial cells; RGC, retinal

1117   ganglion cells; PR, photoreceptor cells; AC, amacrine cells; HC, horizontal cells; UD,

1118   undetermined.

1119

1120

A N2B27 — KSR

Day 0 → Day 1 → Day 4, 5 → Day 6 → Day 13 to 16 onward

Embed hESC sheets in Matrigel → Disperse gently → Clumps of Matrigel/hESCs → Grow in suspension → Cysts → Passage 2-6 cysts per well → Attached colonies → Concentric zones of anterior ectodermal progenitors (CONCEPT)

B
FOXG1+ / VSX2+ PAX6+ — Day 15-17
RGC / OS: PAX2+ VSX2- / OD: PAX2+ VSX2+ / VSX2+ PAX6+ / FOXG1+ — Day 22-26

C Cysts Day 2, TJP1::GFP, 100 u

D CONCEPT Day 26, 1mm

E E10 Foxg1, NR, OS, Tel

F E10 Vsx2

G E10.5 Pax6

H Day 15 FOXG1 G VSX2 R, 1 mm

I 200 u

J Day 17 PAX6 G VSX2 R, 1 mm

K 200 u

L Day 26 FOXG1 G VSX2 R, 1 mm

M 100 u

N Day 26 PAX6 G VSX2 R, 1mm

O 100 u

P BMP4 Day 10, 100 u

Q BMP4 Day 17, 500 u

R BMP4 Day 25, 1 mm

S BMP7 Day 10, 100 u

T FGF8 Day 10, 100 u

U FGF8 Day 17, 500 u

V FGF8 Day 25, 1 mm

W BMP7 Day 25, 1 mm

A Day 22 TUBB3 G POU4F2 R — 200 u

B — 100 u

C Day 26 TUBB3 G POU4F2 R

D

E Pax2 E10.5 — 50 u

F Pax2 E13.5

G E13.5 Pax2 R Tubb3 G — 50 u

H Day 26 PAX2 R TUBB3 G — 200 u

I PAX2

J PAX2 R VSX2 G

K VSX2

L RGC FOXG1+ OS: PAX2+ VSX2− OD: PAX2+ VSX2+

M

group

With Directional RGC Axons

Without

A Day 22 CRYAA
B Day 22 CRY B
C Day 39 CRYAA
D Day 39 CRY B
H 25 With Lens Cells, Without 2
I Day 75
J Day 117
K CRYAA R TUBB3 G Day 39
L kD 50 37 25 20 CRY G ACTB Lentoid bodies 1: Day 81 2: Day 113
M Day 117 Cell 1 Cell 2 Cell 3 Cell 4 5 u
N Cell 2 Cell 3 200 nm

id	source	term_id	term_name	term_size		p_value C2		p_value H18
1	GO:MF	GO:0005201	extracellular matrix structural constituent	174		4.2e−05		1.6e−01
2	GO:MF	GO:0050839	cell adhesion molecule binding	562		2.5e−03		1.3e−05
3	GO:CC	GO:0062023	collagen−containing extracellular matrix	430		6.2e−10		4.3e−05
4	GO:CC	GO:0005615	extracellular space	3384		2.9e−05		3.9e−09
5	GO:BP	GO:0061564	axon development	509		2.5e−04		4.2e−01
6	GO:BP	GO:0048708	astrocyte differentiation	85		2.0e−02		4.1e−04
7	GO:BP	GO:0001654	eye development	404		4.4e−16		2.4e−05
8	GO:BP	GO:0070848	response to growth factor	710		1.7e−09		3.3e−06
9	GO:BP	GO:0007155	cell adhesion	1530		1.2e−08		1.6e−06
10	GO:BP	GO:0048870	cell motility	1705		1.8e−09		1.7e−05
11	GO:BP	GO:0031175	neuron projection development	990		3.0e−07		1.3e−04
12	GO:BP	GO:0010001	glial cell differentiation	235		5.4e−06		1.3e−09
13	GO:BP	GO:0007165	signal transduction	6034		3.4e−03		3.2e−04
14	GO:BP	GO:0048731	system development	3987		5.8e−25		5.7e−13
15	GO:BP	GO:0032502	developmental process	6489		1.0e−21		2.6e−13

g:Profiler (biit.cs.ut.ee/gprofiler)

A
Day 25
CNTN2 G
POU4F2 R
PAX2 M
200 u

B
OS
OD

C

D
100 u

E
100 u

F
100 u

G
Day 25
CNTN2 G
POU4F2 R
ALDH1A3 M
200 u

H
100 u

I
Day 25
CNTN2 G
POU4F2 R
VAX1/2 M
200 u

J
100 u

K
E13.5
Tubb3 G
Aldh1a3 R
100 u

L
TUBB3 G
POU4F2 R
50 u

M
CNTN2 G
POU4F2 R

N
ISL1 R
RBPMS G

O
POU4F2 R
SNCG G

P

Cell Counts			
	Pos.	DAPI	%
SNCG	66	66	100
TUBB3	53	55	96.4
POU4F2	37	55	67.3
ISL1	15	35	42.9
RBPMS	7	35	20

gene	p_val	avg_logFC	pct.1	pct.2	p_val_adj	cluster
PAX2	0	0.930533006	0.769	0.169	0	2
FGF9	0	0.784668891	0.69	0.161	0	2
FGF8	1.77E-213	0.465810314	0.381	0.07	5.94E-209	2