Self-formation of concentric zones of

telencephalic and ocular tissues and

directional retinal ganglion cell axons

Wei Liu [1,2,3¶], Rupendra Shrestha [1,2,3], Albert Lowe [1,2], Xusheng Zhang [2], Ludovic Spaeth [4]

[1]Department of Ophthalmology and Visual Sciences

[2]Department of Genetics

[3]The Ruth L. and David S. Gottesman Institute for Stem Cell Biology and Regenerative

Medicine

[4]Dominick P. Purpura Department of Neuroscience

Albert Einstein College of Medicine, Bronx, NY 10461

[¶]*Lead Contact.* Email: Wei Liu, wei.liu@einsteinmed.edu

Key words: human pluripotent stem cell, organoid, telencephalon, optic stalk, optic disc,

neuroretina, tissue patterning, self-organization, retinal ganglion cell, axon growth and

pathfinding, cell surface protein, cell isolation, glaucoma

Summary

The telencephalon and eye in mammals are originated from adjacent fields at the anterior

neural plate. Morphogenesis of these fields generates telencephalon, optic-stalk, optic-disc, and

neuroretina along an axis. How these telencephalic and ocular tissues are specified

coordinately to ensure directional retinal ganglion cell (RGC) axon growth is unclear. Here, we

report the self-formation of human telencephalon-eye organoids comprising concentric zones of

telencephalic, optic-stalk, optic-disc, and neuroretinal tissues along the center-periphery axis.

Initially-differentiated RGCs grew axons towards and then along a path defined by adjacent

PAX2+ optic-disc cells. Single-cell RNA sequencing identified expression signatures of two

PAX2+ cell populations that mimic optic-disc and optic-stalk, respectively, mechanisms of early

RGC differentiation and axon growth, and RGC-specific cell-surface protein CNTN2, leading to

one-step purification of electrophysiologically-excitable RGCs. Our findings provide insight into

the coordinated specification of early telencephalic and ocular tissues in humans and establish

resources for studying RGC-related diseases such as glaucoma.

(150 words)

Introduction

Our understanding of eye and brain development in humans is mostly deduced from animal studies. In mice, fate mapping of the anterior neural plate reveals that the eye field is in rostral regions surrounded anteriorly and laterally by the telencephalic field and caudally and medially by the hypothalamic field, indicating the proximity of their embryonic origins [1]. Subsequent evagination of the eye field generates bilateral optic vesicles and optic stalks; the optic stalks connect the optic vesicles to the forebrain, forming a midline-periphery axis. The optic vesicle then invaginates ventrally, resulting in the formation of double-layered optic cups in which the inner and outer layers develop into the neuroretina and retina pigment epithelium (RPE), respectively. The posterior pole of the optic-cup forms the optic disc (also known as optic nerve head). In the central neuroretina close to the nascent optic disc, retinal ganglion cells (RGCs) start to appear as the optic fissure nearly closes. Early RGC axons find their path toward the optic disc and then enter the optic stalk to reach the brain. Concentrically organized growth-promoting and growth-inhibitory guidance molecules around the optic disc regulate RGC axon growth and pathfinding through multiple mechanisms [2-4]. Coinciding with the optic-cup formation, the surface ectoderm facing the optic vesicle develops into lens. Collectively, the early development of telencephalic and ocular tissues is highly coordinated in mammals.

Early telencephalic and eye development is marked and regulated by a group of tissue-specific transcription factors and signal transduction molecules [5]. In mice, Foxg1 is specifically expressed in the presumptive telencephalon and is essential for the development of the cerebral hemispheres [6]. Pax6 is specifically expressed in the eye field and is essential for the development of multiple ocular lineages, such as the neuroretina, lens, and RPE [7-10]. Vsx2 and Mitf are specifically expressed in the neuroretina and RPE, respectively, and are essential for retinal development [11-14]. Pax2 is expressed in the optic stalk, optic vesicles, central neuroretina, and optic disc; Pax2 is essential for optic stalk and nerve development [15-18]. Fgf8 is specifically

expressed at the rostral forebrain at early stages, induces Foxg1 expression [19], and regulates

telencephalic patterning in a dose-dependent manner [20]. Fgf8 also maintains Pax2 expression

in the optic stalk [21], and Fgf8 and Fgf3 coordinate the initiation of retinal differentiation in chicks

[22,23]. Despite these findings in vertebrates, it is unclear how telencephalic and ocular tissues are

specified in humans.

Human organoids are transformative since they enable us to study the mechanisms of cell

specification and differentiation directly in humans [24]. Self-organized three-dimensional (3-D)

retinal organoids are originally reported by Sasai's group and further improved in follow-up

studies [25-33]. We and others have demonstrated that retinal organoids derived from human

embryonic stem cells (hESCs) display a stratified structure containing all major types of retinal

cells. Although RGCs are differentiated in 3-D retinal organoids, there is no proper RGC axon

outgrowth toward the optic disc as seen in vivo since the optic disc-like tissue is missing in

retinal organoids. When these organoids are dissociated into single cells or cut into pieces for

adherent culture, RGCs generate neurites [34-36]. A variety of brain organoids have been

described [37-41]. Although rudimentary ocular tissues are occasionally found in some brain

organoids [42], ocular and brain tissues are not patterned along any defined axis. Assembloids, in

which retinal, thalamic, and cortical organoids are assembled [32], do not contain any optic disc

and stalk tissues. Collectively, RGC axon outgrowth and pathfinding directed by optic disc and

stalk tissues in organoids have not been reported. Since RGCs degenerate in glaucoma and

cell replacement therapies are actively evaluated, it is imperative to establish models for

studying RGC axon growth and pathfinding in humans.

Tissue patterning and coordinated specification are fundamental for body plan formation in vivo.

Remarkably, concentric zones of trophectoderm, endoderm, mesoderm, and ectoderm [43,44],

neural plate and neural plate border [45], and ectodermal cells [46] are self-organized from single

pluripotent stem cells, indicating patterning and coordinated specification of these tissues in

vitro. Nevertheless, optic disc and stalk tissues are not reported in any of these structures.

We hypothesize that coordinated specification from the anterior ectodermal epithelium via

morphogen gradients leads to self-organization of telencephalic and ocular tissues, including

optic disc- and stalk-like tissues that provide guidance cues for RGC axon growth and

pathfinding. In support of this hypothesis, we generated self-formed human telencephalon-eye

organoids that comprise concentric zones of anterior ectodermal progenitors (CONCEPT),

including tissues mimicking FOXG1+ telencephalon, PAX2+ optic stalk, PAX2+ optic disc,

VSX2+ neuroretina, and PAX6+ multi-lineage tissues along the center-periphery axis. We refer

these tissues as telencephalon, optic stalk, optic disc, and neuroretina for the sake of gene

marker expression, tissue functions, and the ease of description; we do not claim these tissues

represent in vivo tissues in all aspects. We call this structure as an organoid since it displays a

well-organized structure with cell identities mimicking those tissues in vivo. In CONCEPT

organoids, early differentiated RGCs grew their axons toward and then along a path defined by

adjacent PAX2+ optic-disc cells. Single-cell RNA sequencing of CONCEPT organoids not only

confirmed telencephalic and ocular identities but also discovered expression signatures of cell

clusters. We found that PAX2+ optic-disc cells differentially expressed FGF8 and FGF9;

inhibition of FGF signaling with an FGFR inhibitor during early RGC differentiation drastically

decreased RGCs somas and nearly ablated directional axon growth. Using cell-surface marker

CNTN2 identified in our single-cell RNA-sequencing, we developed a one-step method for

isolating human electrophysiologically-excitable RGCs under a native condition. Our studies

provide deeper insight into the coordinated specification of telencephalic and ocular tissues to

ensure guided RGC axon growth in humans and establish resources for studying

neurodegenerative diseases such as glaucoma.

Results

Generation of a telencephalon-eye organoid that is composed of <u>con</u>centric zones of

anterior <u>e</u>ctodermal <u>p</u>rogeni<u>t</u>ors (CONCEPT)

The telencephalon and eye in mammals are originated from adjacent fields at the anterior

neuroepithelium [1]. Morphogenesis of these embryonic fields leads to the formation of

telencephalon, optic stalk, optic disc, and neuroretina along an axis. Early differentiated retinal

ganglion cells (RGCs) in the neuroretina grow axons toward the optic disc and then along the

optic stalk to reach the brain. How these telencephalic and ocular tissues are specified

coordinately to ensure directional RGC axon growth is unclear in humans.

Cysts are hollow spheres of an columnar epithelium induced from human pluripotent stem cells

through embedding hESC sheets in Matrigel; they are used to generate retinal cells [28,33,47].

Nevertheless, developmental potentials of cysts have not been characterized.

Using the cysts, here we generate a telencephalon-eye organoid that is composed of <u>con</u>centric

zones of anterior <u>e</u>ctodermal <u>p</u>rogeni<u>t</u>ors (CONCEPT) from cysts (Fig. 1A, B). Cysts were

generated as previously described [28,33]. The epithelial structure of cysts was demonstrated by

apically localized TJP1::GFP reporter at the lumen (Fig. 1C), consistent with previous findings

[28,47]. To assess developmental potentials of cysts, individual cysts were manually picked and

then seeded onto the Matrigel-coated surface at low densities (Fig. 1A). After attaching to the

culture surface, cysts initially grew as dome-shaped individual colonies. Subsequent culture of

the colonies in a KSR medium (see Materials and Methods) led to the self-formation of

concentric zones of anterior ectodermal progenitors: an elevated central zone surrounded by

multiple zones; this morphology was distinguishable under a steoromicroscope (Fig. 1D) or an

inverted microscope. CONCEPT structures were highly reproducible, which is demonstrated by

consistent gene expression profiles of multiple CONCEPT organoids in whole culture wells (Fig.

S1). If multiple cysts were fused together or cysts for seeding were too big or small, CONCEPT

structures were affected or incomplete. CONCEPT structures expressed gene markers for

telencephalon and eye in concentric patterns. In mice, Foxg1 is specifically expressed in the

E10 telencephalon at high levels; it was also expressed in the optic stalk, invaginating optic cup,

and lens at lower levels, with the rostral optic stalk connecting the telencephalic vesicle to the

invaginating optic cup (Fig. 1E) [6]; Vsx2 is specifically expressed in the E10.5 neuroretina (Fig.

1F); Pax6 is specifically expressed in the E10.5 neuroretina, RPE, lens vesicles, and surface

ectoderm (Fig. 1G) [13]. In CONCEPT structures, the expression of FOXG1, VSX2, and PAX6

generally exhibited concentric patterns spanning from the center to the periphery (Fig. 1H-O).

Similar to FOXG1, telencephalon marker EMX2 [48] was also expressed at the central region (Fig.

S1F). PAX6 was expressed in multiple concentric zones at distinct levels, mimicking its

expression patterns in multiple ocular tissues in E10.5 mouse eyes (Fig. 1G, J, K, N, O).

Therefore, the CONCEPT structure is a telencephalon-eye organoid mimicking telencephalic

and ocular tissues in the aspects of cell types and patterning at early stages.

Morphogens FGFs and BMPs play crucial roles in patterning the forebrain and eye in vivo. In

mice, Fgf8 is specifically expressed at the rostral forebrain at early stages, induces Foxg1

expression [19], and regulates telencephalic patterning in a dose-dependent manner [20]. Bmp4 and

Bmp7 are expressed in the dorsomedial telencephalon, optic vesicles, and presumptive lens

placodes [49-52]. Bmp4 is required for lens induction [50], and Bmp7 is required for proper patterning

of the optic fissure [53]. In CONCEPT telencephalon-eye organoids, FGF8, BMP4, and BMP7

were expressed in attached cell colonies starting from early stages (Fig. 1P, S, T) and

subsequently exhibited circular patterns (Fig. 1R, V, W). At day 10, the expression of BMP4 and

BMP7 delineated multiple rings, with a big ring mostly at the center surrounded by numerous

small rings (Fig. 1P, S). These observations suggest that the attachment of a single-lumen cyst

to the culture surface caused differences in cell behaviors: some cells separated from the

original cyst and migrated peripherally; some of the separated cells formed small ring-like

structures; the cells that remained at the center formed a big ring-like structure. At day 17,

circular patterns of BMP4 and FGF8 emerged (Fig. 1Q, U). At day 25, BMP4, FGF8, and BMP7

clearly exhibited circular patterns (Fig. 1R, V, W). Hence, FGFs and BMPs spontaneously form

circular gradients, which would dictate coordinated cell specification in CONCEPT

telencephalon-eye organoids.

Early differentiated RGCs grow directional long axons toward and then along a path

defined by PAX2+ cell populations in CONCEPT telencephalon-eye organoids

To assess cell differentiation in CONCEPT telencephalon-eye organoids, we examined marker

expression for RGCs, the first type of cells that differentiate in the neuroretina. In mice,

transcription factor Pou4f2 is expressed in the early differentiated RGCs and required for the

development of a large set of RGCs [54]. Tubb3 is expressed in the somas and axons of

differentiating RGCs.

In CONCEPT organoids, POU4F2 and TUBB3 were detectable as early as day 17 and

increased to higher levels at day 22. RGC somas were marked by POU4F2 and TUBB3 co-

expression; RGC axons were marked by TUBB3 expression. Interestingly, RGCs grew

directional long axons that followed a circular path (Fig. 2A, B). The circular path of RGC axon

outgrowth became more evident in organoids at day 26 (Fig. 2C, D). In mice, the initially

differentiated RGCs are adjacent to the optic disc; their axons grow towards the optic disc to exit

the eye and then navigate within the optic stalk to reach their targets in the brain; the optic disc

and stalk specifically express Pax2 at distinct levels (Fig. 2E-G) [17,55,56]; presumptive optic disc

cells expressed VSX2 whereas optic stalk cells do not [13] (Fig. 1F; 2E). In CONCEPT organoids,

there were two PAX2+ cell populations that formed two adjacent rings (Fig. 2H-L). POU4F2+

RGCs grew TUBB3+ axons toward and then navigated along the adjacent PAX2+ VSX2+ cell

population (Fig. 2H-L), mimicking axon growth from the initial RGCs toward the PAX2+ optic

disc in vivo. Meanwhile, PAX2+ VSX2- cell population at the inner zone set up an inner

boundary of the path for RGC axon growth, mimicking PAX2+ optic stalk that spatially confines

RGC axon growth in vivo. Based on these findings, we designate PAX2+ VSX2+ cells and

PAX2+ VSX2- cells as the optic disc and optic stalk, respectively. Taken together, our findings

demonstrate that RGCs grow directional axons toward and then along a path defined by PAX2+

cell populations in CONCEPT telencephalon-eye organoids.

CONCEPT telencephalon-eye organoids contain lens cells that undergo terminal

differentiation

CONCEPT telencephalon-eye organoids at stages around day 25 contained transparent

structures reminiscent of the ocular lens. To determine their cell identity, we performed

immunostaining. Starting at day 22, lens markers CRYAA and CRY B were found (Fig. 3A, B)

and continuously expressed in the transparent cell structures (Fig. 3C, D). Interestingly, these

transparent structures were not stained by DAPI (Fig. 3C-F), indicating denucleation in these

lens cells. When CONCEPT organoids were detached using Dispase at around day 28 and

continuously grown as suspension cultures, crystal-like clusters with fused transparent spheres

were found, and they continuously survived for months (Fig. 3I, J). In mice, terminally-

differentiated lens fiber cells are free of organelles and featured by specialized interlocking cell

membrane domains shown as ball-and-sockets and protrusions [57]. Using transmission electron

microscopy, we found that the crystal-like lens clusters were free of organelles and exhibited

ball-and-socket structures (Fig. 3K, L). Taken together, our findings demonstrate that

CONCEPT telencephalon-eye organoids contain lens cells that undergo terminal differentiation;

FGFs in CONCEPT organoids likely promote terminal lens differentiation as seen in other

settings [58].

Single-cell RNA sequencing analysis identifies telencephalic and ocular cell populations in CONCEPT telencephalon-eye organoids

To fully characterize cell populations in CONCEPT organoids and the mechanisms underlying RGC axon pathfinding, we performed 10x single-cell RNA sequencing of the organoids at day 24, around the stage when RGCs grew long axons toward and along the PAX2+ VSX2+ cell population. Cell Ranger mapping showed that 11158 single cells were sequenced at a depth of 27,842 reads and 2,967 genes per cell, and the dataset were then analyzed using Seurat (v3.2.0) [59]. Cell filtration (nFeature_RNA > 200 & nFeature_RNA < 6000 & percent.mt < 20) resulted in 10218 cells with high quality. Cell clustering grouped cells into 14 clusters (Fig. 4A; Tab. 1). In addition, cell cycle scores were plotted using an established method (Fig. 4B) [60]. Several cell clusters, *e.g.,* clusters 4, 8, 9, were separated along cell cycle scores (Fig. 4A, B), indicating the differences in cell cycle phases across these cell clusters.

We next assessed cell identities using known markers. Mesoderm, endoderm, and neural crest are identified by a group of gene markers [61-64]. In CONCEPT organoids, gene markers for mesoderm (TBXT, GATA2, HAND1), endoderm (GATA1, GATA4, SOX17), and neural crest (SNAI1, SOX10, FOXD3) were not expressed (Fig. S2). In contrast, gene markers for the anterior neuroectoderm were widely expressed. Telencephalic progenitor marker FOXG1 was expressed in clusters 10, 3, 12, 1, 4, 8, 9, 13; retinal progenitor markers PAX6 and VSX2 were expressed in clusters 2, 7, 5, 6, 11 (Fig. 4C-E). Cluster 0 was defined by a group of negative markers (Fig. S3); they expressed both FOXG1+ telencephalic cells and PAX6+ and/or VSX2+ retinal cells. In contrast to telencephalon and retinal markers, diencephalon markers (GBX2, WNT3, SOX14) [65,66] and midbrain/hindbrain markers (EN2, PAX7, TFAP2B) [61] were rarely expressed (Fig. S4). Lens markers CRYAA and FOXE3 were expressed in a very small cell population; those denucleated lens cells were not captured in the scRNA-seq since the absence of nucleus prevented active mRNA transcription. Therefore, lens cells did not form a separate

cluster probably due to their very low abundance. Taken together, these findings indicate that

CONCEPT telencephalon-eye organoids at day 24 are mostly composed of FOXG1+

telencephalic cells and PAX6+ and/or VSX2+ retinal cells.

We next characterized the FOXG1+ telencephalic cells via assessing differentially expressed

genes (DEGs). DEGs in clusters 4, 8, 9, 1 included POU3F3, RGMA, EDNRB, and SOX3,

which orthologs in mice are expressed in the pallium. DEGs in clusters 12, 3, 10 included DLX2,

RGS16, DLX1, DLL1, DLX6-AS1, NEFL, DCX, and RTN1, which orthologs in mice are

expressed in the subpallium (Figs. S5, S6). Therefore, FOXG1+ telencephalic cells in

CONCEPT organoids at day 24 comprise cell clusters in both the pallium and subpallium.

PAX6+ and/or VSX2+ retinal cells were also grouped into several clusters. VSX2 was

expressed in clusters 2, 7, 5, and 0, with cluster 2 at G1 phase, cluster 7 in G1 and S phases,

and cluster 5 in S and G2M phases (Fig. 4A, B, E). Cluster 6 did not VSX2 but expressed PAX6

and was at the G1 phase, suggesting cells in cluster 6 exited the retinal progenitor cell state and

underwent cell differentiation (Fig. 4A, B, D, E). Consistently, cluster 6 differentially expressed

RPE markers, e.g., PMEL, HSD17B2, DCT, and MITF (Fig. S7), indicating they were mostly

differentiating RPE. Cluster 11 did not express VSX2; instead, it expressed PAX6 and was at

the G1 phase (Fig. 4A, B, D, E), suggesting that cells in cluster 11 exited the retinal progenitor

cell state and underwent cell differentiation.

Taken together, our single-cell RNA sequencing analysis of CONCEPT telencephalon-eye

organoids at day 24 confirms their telencephalic and ocular identities, establishing a valuable

transcriptomic dataset for mechanistic studies.

Identification of two PAX2+ cell populations that mimic to the optic disc and optic stalk,

respectively, in the scRNA-seq dataset of CONCEPT telencephalon-eye organoids

To identify PAX2+ cell populations that defined the path for RGC axon outgrowth, we examined

PAX2 expression in the dataset. PAX2 was mostly expressed in cluster 2 and subsets of

clusters 7, 5, 0, 4, 8, and 9 (Fig. 4A, F). Since clusters 2, 7, 5, and 0 expressed VSX2 (Fig. 4E,

F), we deduced that PAX2+ VSX2+ cells in cluster 2 and subsets of clusters 7, 5, and 0

corresponded to those PAX2+ VSX2+ cells that defined the path for RGC axon growth (Figs.

2H-L). Therefore, these PAX2+ VSX2+ cells in the dataset of single-cell RNA sequencing were

assigned as the optic disc (OD) (Fig. 4F). PAX2+ VSX2+ cells also expressed optic disc and

optic stalk marker SEMA5A (Fig. 4F, G) [67]. We particularly focused on cluster 2 since PAX2+

VSX2+ cells were mostly found in this cluster. Interestingly, COL13A1 was differentially

expressed in cluster 2 (Fig. 4H; Fig. S10); in situ hybridization indicates that COL13A1 is

specifically expressed in the optic disc of human fetal retinas [68]. These findings firmly support

that PAX2+ VSX2+ cells in CONCEPT telencephalon-eye organoids corresponded to optic disc

cells in human fetal retinas. Cluster 2 also differentially expressed glaucoma gene CYP1B1,

signaling molecules LEFTY2, FGF9, and FGF8 (Fig. 4I-L; Fig. S10). The expression of PAX2,

SEMA5A, CYP1B1, LEFTY2, FGF9, and FGF8 in CONCEPT organoids was validated using in

situ hybridization (Figs. 1T-V; 5B-E; S10).

In contrast, PAX2+ VSX2- cells in subsets of clusters 4, 8, 9 (Fig. 4A, E, F) expressed the optic

stalk marker VAX1 (Fig. 4F, M); these cells corresponded to those PAX2+ VSX2- cells that set

up the inner boundary of the path for RGC axon growth in CONCEPT organoids (Fig. 2H-L).

Therefore, PAX2+ VSX2- cells in subsets of clusters 4, 8, 9 were assigned as the optic stalk

(OS) (Fig. 4F). PAX2+ VSX2- optic stalk cells also differentially expressed CNTNAP2, ALDHA3,

and LAMP5 (Fig. 4N-P), which orthologs in mice are expressed in the optic stalk/nerve (Fig. S8).

Compared to PAX2+ VSX2+ optic disc cells, PAX2+ VSX2- optic stalk cells were closer to

telencephalic cells in both the UMAP plot (Fig. 4M-P; Fig. S5A-D) and their positioning in

CONCEPT organoids (Fig. 2H-L), mimicking relative positions of telencephalon, optic stalk, and

optic disc along the midline-periphery axis in E10.5-E13.5 mouse embryos (Figs. 1E, 2E-G).

Consistently, several markers for PAX2+ VSX2- optic stalk cells were also expressed in the

telencephalon (Fig. S8). Collectively, we identify two PAX2+ cell populations that mimic the

optic disc and optic stalk, respectively, in the scRNA-seq dataset of CONCEPT telencephalon-

eye organoids.

Identification of glycosylphosphatidylinositol (GPI)-anchored cell membrane protein

CNTN2 as a specific marker for developing human RGCs

To identify RGCs in the dataset, we checked the expression of RGC markers. We found that

RGC markers ATOH7, POU4F2, SNCG, and CNTN2 were differentially expressed in cluster 11,

indicating its RGC identity (Fig. 4A, 4Q-T). CNTN2 caught our attention because it is a GPI-

anchored cell membrane protein, which may be used as a native marker for RGC isolation. In

literature, Cntn2 is specifically expressed in developing RGCs in mice and chicks [69,70].

In CONCEPT at day 25, CNTN2 exhibited an expression pattern very similar to that of TUBB3

(Figs. 2, 5). CNTN2 was found in the cell membrane of RGCs that expressed POU4F2 in the

cell nucleus (Fig. 5A, D). PAX2+ cells formed inner and outer concentric zones, mimicking the

optic stalk and optic disc, respectively. POU4F2+ RGCs formed a dense circular zone adjacent

to PAX2+ optic disc cells; POU4F2+ RGCs were sparse in more peripheral areas (Fig. 5B).

These findings indicate that early differentiated RGCs were adjacent to PAX2+ optic disc cells.

Interestingly, RGCs in the dense POU4F2+ zone grew CNTN2+ axons toward and then along

the path defined by adjacent PAX2+ optic disc cells (Fig. 5A-E). In areas where there were a

gap in PAX2+ optic disc cells, CNTN2+ axons exited the circular path (diamond arrow in Fig. 5A;

a gap between two arrowheads in Fig. 5B, F). In regions where POU4F2+ RGCs were a few

hundreds of micrometers away from PAX2+ optic disc cells, RGCs grew axons in centrifugal

directions (arrow in Fig. 5C). Very similar findings were found using hiPSCs (Fig. S9), indicating

the reproducibility of CONCEPT organoids in multiple cell lines. PAX2+ optic disc cells did not

express ALDH1A3; instead, the path for RGC axon growth was bordered by two cell populations

that highly expressed ALDH1A3 (Fig. 5G-H). In mice, Aldh1a3 expression was low in

differentiating RGCs in the central retina but high in peripheral retinal progenitors and optic stalk

(Fig. 5K) [71], consistent with ALDH1A3 expression in CONCEPT organoids (Fig. 5G, H). In mice,

Vax1 and Vax2 are expressed in the optic stalk and ventral retina and are jointly required for the

optic stalk development [55,72]. In CONCEPT organoids, cells that set up the inner boundary for

RGC axon growth also expressed optic stalk marker VAX1/2 (Fig. 5I, J; the antibody recognizes

both VAX1 and VAX2). Taken together, our data demonstrate that CNTN2 is a specific cell-

surface marker for differentiating human RGCs; RGCs grow axons toward and then along a

path that is defined by adjacent PAX2+ VSX2+ ALDH1A3- optic disc cells.

One-step isolation of developing human RGCs via native marker CNTN2

Since cell surface protein CNTN2 was specifically expressed in developing human RGCs, we

sought to test whether CNTN2 can be used as a biomarker for isolating human RGCs under a

native condition. To that goal, retinal organoids in suspension culture on days 41 - 70 were

dissociated using Accutase to generate a single cell suspension, which was then subject to

magnetic-activated cell sorting (MACS) with an antibody against CNTN2. From 100 retinal

organoids on day 41 to 48, around 385,000 RGCs were isolated. Isolated RGCs were plated

onto Matrigel-coated chamber slides for 10-day adherent culture. These cells exhibited neuronal

morphology and widely expressed RGC markers TUBB3 and POU4F2 (Fig. 5L). Isolated RGCs

tended to form clusters in culture. In contrast to directional axon growth found in CONCEPT

organoids, isolated RGCs grew neurites in random directions (Fig. 5L), indicating the

differences in axon pathfinding cues between CONCEPT organoids and isolated RGC cultures.

RGC neurites were also marked by CNTN2 expression (Fig. 5M). A portion of isolated RGCs

expressed POU4F2 but not CNTN2 (Fig. 5M), indicating that CNTN2 was downregulated in

dissociated RGC cultures. Isolated RGCs also expressed RGC markers ISL1, RBPMS, and

SNCG (Fig. 5N-P). Collectively, we have developed a one-step method for the isolation of

differentiating human RGCs in a native condition.

Isolated RGCs exhibit electrophysiological signature of excitable cells

In order to determine the functional properties of isolated RGCs in culture, we examined their

electrophysiological properties using whole-cell patch clamp recordings (Fig. 6, *see Methods*).

Using current-clamp configuration, we found that RGCs displayed had a hyperpolarized resting

membrane potential (*mean ±SD: -20.1 ±6.4mV*, Fig. 6A). When cells were held at -70 mV by

current injection, in most cases (6/9), depolarizing currents steps could trigger an action

potential, which were often followed by a depolarization plateau if the current was injected for

more than 10ms (Fig. 6B). In voltage clamp recordings we examined the nature of the voltage-

gated conductances (*see Methods*). From a holding membrane potential of -80 mV, both inward

and outward currents (I_m) were observed in response to depolarizing voltage steps (V_m, Fig. 6C).

Outward currents were primarily mediated by voltage-gated potassium channels, as application

of 20mM TEA significantly reduced their amplitude (*mean I_K ±SD at 60mV; 964±537 pA in*

control; 279±201 pA in TEA, U=24.0; p=0.019, two-sided MannWhitneyU test, Fig. 6D).

Conversely, 1µM TTX abolished all inward currents, demonstrating that they were mediated by

activation of voltage-gated sodium channels (*mean I_{Na} ±SD at -10mV; -338.2±121 in control; -*

18.3±13 pA in TTX; U=0.0; p=0.035, two-sided MannWhitneyU test, Fig. 6E). Taken together,

these results indicate that isolated RGCs exhibit functional features traditionally found in

excitable cells, such as in neurons.

FGF signaling mediated by FGFRs is required for early RGC differentiation and

directional axon growth in humans

Directional RGC axon growth toward and then along PAX2+ optic disc cells in CONCEPT

organoids could be mediated by signaling molecules secreted from PAX2+ optic disc cells.

scRNA-seq analysis identified expression signatures of cluster 2, the major component of

PAX2+ optic disc cells (Fig. S10). Interestingly, FGF8 and FGF9 were differentially expressed in

cluster 2 (Figs. 4K, 4L; 7A; S10). To assess the roles of FGF8 and FGF9 in CONCEPT

organoids, we validated that FGF8 and FGF9 were expressed in PAX2+ optic disc cells of

CONCEPT organoids (Fig. 7B-E). TUBB3+ RGC axons grew towards and then along the

regions with high-level FGF8 and FGF9 (Fig. 7D, E), suggesting that FGF8 and FGF9 may

attract RGC axon growth. FGFR1, FGFR2, FGFR3, MAP2K1, and MAP2K2 were expressed in

multiple types of cells in CONCEPT organoids. In RGCs, FGFR1 and MAP2K2 were clearly

expressed (Fig. S11), indicating that RGCs expressed the components that transduce FGF

signaling. Since FGF8 and FGF9 are probably redundant and it is challenging to inactivate both

FGF8 and FGF9 in CONCEPT organoids, we decided to inactivate FGF signaling with

FGFR1/2/3 inhibitor PD 161570 during days 17-24, a stage at early RGC differentiation. After

the FGFR inhibition, FGF8 expression was grossly unaffected. On the other hand, RGC somas

were drastically reduced, and directional axon growth was nearly absent (Fig. 7F-7K). These

findings indicate that a) PAX2+ optic disc cells differentially expressed FGF8 and FGF9; b) FGF

signaling mediated by FGFRs is required for early RGC differentiation and directional axon

growth in humans.

Discussion

In this study, we report the self-formation of concentric zones of telencephalic and ocular tissues

in CONCEPT telencephalon-eye organoids from human pluripotent stem cells, establishing a

model for studying the early development of telencephalic and ocular tissues in humans. RGCs

grew axons toward and then along a path defined by PAX2+ cell populations, setting up a

model for studying RGC axon pathfinding. We identified expression signatures of cell clusters in

CONCEPT organoids using single-cell RNA sequencing and revealed mechanisms of

mechanisms of early RGC differentiation and axon growth. Lastly, we established a one-step

method for the isolation of human electrophysiologically-excitable RGCs via CNTN2 under a

native condition. Our studies not only provide deeper insight into coordinated specification of

telencephalic and ocular tissues for directional RGC axon growth in humans, but also generate

resources for therapeutic studies of RGC-related diseases such as glaucoma.

The cyst is a radially symmetric epithelium mimicking the anterior ectoderm

Cysts are hollow spheres composed of homogeneous columnar epithelial cells with the apical

surface at the lumen. They are induced from pluripotent stem cells via embedding small sheets

of hESCs into Matrigel and subsequent growth either in a solid thin film [47] or suspensions [28,33].

Cyst growth in suspensions is cost-effective and scalable. The formation of cysts induced by

Matrigel mimics the epithelization of the epiblast by the extracellular matrix (ECM) in the

blastocyst [73,74].

In this study, individual cysts efficiently generate telencephalic and ocular tissues in the absence

of any extrinsic factors, indicating their default cell fates of the anterior ectoderm. In literature,

anterior neural tissues are generated from re-aggregated single pluripotent stem cells through

dual inhibition of Smad signaling [75] or inhibition of Wnt/ß-catenin signaling [27]. Besides that,

undirected cultures without extrinsic factors are used for the generation of retinal cultures [29,30,76].

In our procedure, we did not add any extrinsic factors; the epithelial structure of cysts and

subsequent adherent growth at a low density facilitate neural induction and differentiation. Our

findings are consistent with the reports that epiblast cells are epigenetically primed for

ectodermal fates [77] and neuroectodermal tissues are the default differentiation from pluripotent

stem cells [78,79]. Collectively, cysts mimic the anterior ectoderm in the epithelial structure and cell

fates.

Pattern formation and coordinated cell differentiation in CONCEPT telencephalon-eye

organoids

Tissue patterning is fundamental for the formation of a body plan, which is defined by the

anteroposterior, dorsoventral, and left-right axes. Self-formation of CONCEPT telencephalon-

eye organoids establishes an efficient way to pattern early telencephalic and ocular tissues

along an axis. In CONCEPT organoids on days 22-26, FOXG1+ telencephalon, PAX2+ optic

stalk, PAX2+ optic disc, and VSX2+ neuroretinas are positioned along the center-periphery axis,

mimicking the relative positions of those tissues in E10.5-13.5 mouse embryos. Our findings

indicate that the formation of telencephalic and retinal tissues are highly coordinated, consistent

with the prosomere model [80,81]. Collectively, we establish CONCEPT telencephalon-eye

organoids for studying the early formation of telencephalic and ocular tissues in humans.

The tissue patterning in CONCEPT telencephalon-eye organoids is originated from the

attachment of a radially symmetric epithelium to the culture surface for growth as colonies since

individual cysts kept in suspensions do not form any apparent tissue patterning. When a floating

cyst initially contacted the culture surface, the contact resulted in ECM-cell adhesions. The initial

ECM-cell adhesions caused additional ECM-cell contacts in neighboring cells, resulting in the

flattening and spreading of a cyst onto the culture surface. Since the cyst is a radially symmetric

epithelium, ECM-cell adhesions between the cyst and the culture surface formed sequentially in

a concentric manner. Remodeling of cell-cell and ECM-cell adhesions and self-organization

underlay the tissue patterning. Molecular events in concentric patterns during the attachment of

a cyst onto the culture surface were eventually translated to concentric gradients of morphogens

that specify cell fates. The radially symmetric epithelial structure of cysts is important for the

formation of a concentric pattern since such pattern would not be generated when amorphous

embryoid bodies are attached to the culture surface for growth as colonies. In literature, timed

BMP4 treatment is shown to promote neuroretinal differentiation from pluripotent stem cells [82],

and FGF8 promotes telencephalic and eye development [5,22]. In our system, BMP4 and FGF8,

along with BMP7 and other FGFs, were highly expressed starting at early stages and gradually

formed concentric morphogen gradients, which would dictate tissue patterning, resulting in

coordinated specification of telencephalic, optic stalk, optic disc, and neuroretinal tissues along

the center-periphery axis in CONCEPT telencephalon-eye organoids. Co-culture of these

tissues in the concentric zones provides mutual cues for coordinated tissue specification and

growth. Nevertheless, anchorage culture of cell colonies prevents three-dimensional

morphogenesis as seen in embryos, impeding proper tissue interactions at more advanced

stages. When these cell colonies are detached using dispase for suspension culture, retinal

progenitor cells self-organize to form protruding, translucent spheres called retinal organoids

whereas other types of cells form structures with dark appearance; concentric zones no longer

exist [28]. It is conceivable that the simple suspension culture does not have proper physical and

chemical supports provided by periocular tissues and other surrounding tissues in vivo and

therefore optic stalk cells do not form a tube-like structure. It is a challenge ahead to reconstruct

this three-dimensional environment to mimic the morphogenesis in vivo.

Concentric patterns of stem cell-derived cultures are also reported in a few other experiments.

When dissociated single pluripotent stem cells are grown in micropatterned culture surface at

442 certain cell densities in a medium supplemented with BMP4, concentric zones of progenitors

443 expressing markers for trophectoderm, endoderm, mesoderm, and ectoderm are found [43,44].

444 Cell density and colony geometry dictate cell fate specification from pluripotent stem cells. A

445 concentric gradient of BMP4 activity regulates the patterning [43]. When dissociated single

446 pluripotent stem cells are grown in pre-patterned geometrically confined culture surface in a

447 medium supplemented with dual inhibitors for TGF-β and BMP4, concentric zones of

448 progenitors expressing the markers for neural plate and neural plate border are observed.

449 Morphogenetic cues—cell shape and cytoskeletal contractile force—dictate the patterning of the

450 neural plate and neural plate border via BMP-SMAD signaling [45]. When dissociated single

451 pluripotent stem cells are grown as individual colonies in a medium supplemented with knockout

452 serum replacement, multiple zones of ectodermal cells autonomously form [46]. In all three

453 experiments, dissociated single cells are used to generate cell colonies through either cell re-

454 aggregation or proliferation. In our case, adherent culture of a radially-symmetric epithelium –

455 the cyst – was used to generate CONCEPT organoids. Among the three previous and our

456 experiments, starting cells differ in their developmental potentials, tissue structures, and

457 treatments. Despite the differences, all these organoids comprise concentric zones of distinct

458 cell populations. Formation of CONCEPT telencephalon-eye organoids mimics the coordinated

459 specification of early telencephalic and ocular tissues in humans.

460 **RGC axon pathfinding cues in CONCEPT telencephalon-eye organoids**

461 In the mouse retina, multiple RGC axon guidance cues are concentrically organized around the

462 optic disc, regulating RGC axon growth and exit from the eye through the optic stalk [2]. Early

463 differentiated RGCs are in a short distance from the nascent optic disc, and axons of later

464 differentiated RGCs in more peripheral retinal regions follow the path of the initial axons. It is

465 accepted that the optic-disc regions provide growth-promoting guidance cues whereas

466 peripheral retinal regions provide inhibitory guidance cues [2]. Pax2 is specifically expressed in

the ventral optic stalk, optic vesicles, central neuroretina, optic disc, and optic stalk; Pax2 is

essential for optic stalk and nerve development in mice [15-17].

In CONCEPT telencephalon-eye organoids, coordinated specification of telencephalic and

ocular tissues generates PAX2+ optic-disc and optic-stalk cells that define the path for RGC

axon growth. To the best of our knowledge, directional RGC axon growth guided by optic-disc

and optic-stalk tissues has not been reported.

Single-cell RNA sequencing of CONCEPT organoids identified DEGs of PAX2+ optic-disc and

optic-stalk cells. Interestingly, FGF8 and FGF9 were differentially expressed in PAX2+ optic-disc

cells; inhibition of FGF signaling with an FGFR inhibitor during early RGC differentiation

drastically decreased the number of RGC somas and nearly ablated directional axon growth,

indicating that early RGC differentiation and directional axon growth require FGFR-mediated

FGF signaling in humans. In chicks, Fgf8 and Fgf3 coordinate the initiation of retinal

differentiation [22,23]; FGF8 maintains Pax2 expression in optic nerve explants [21]. In mice, FGF8

and FGF9 trigger axon outgrowth in motor neuron column explants [83]. Therefore, the roles of

FGF signaling in the regulation of early RGC differentiation and directional axon growth in

CONCEPT organoids are consistent with literature. Dissecting DEGs of PAX2+ optic-disc cells

in CONCEPT telencephalon-eye organoids elucidates the mechanisms of RGC axon growth

and pathfinding in humans.

RGCs described in this study are still at their early stages in development. One prominent

feature of these RGCs are the growth of their axons towards and long the path defined by optic

disc cells, which is quite clear in CONCEPT organoids. Optic stalk cells in CONCEPT organoids

confine RGC axon growth at inner regions, preventing RGC axons to enter telencephalic areas.

We did not observe any clear evidence of synaptic or dendritic connections between RGCs and

other cells at tested stages. Engineering assembloids may be needed to establish connections

between RGCs and their visual targets.

One-step isolation of developing human RGCs under a native condition

RGCs are degenerated in glaucoma, a major cause of vision impairment in developed countries.

Disease modeling and drug discovery to suppress RGC death will have a huge impact on

saving vision. The use of human RGC models is critical for therapeutic studies since humans

and rodents differ significantly in RGCs. Additionally, cell replacement therapies for glaucoma

are extensively evaluated. Therefore, efficient isolation of human RGCs in a native condition will

have a substantial impact on therapeutic studies of RGC-related diseases such as glaucoma. In

literature, cell surface marker Thy1 is used for the isolation of adult mouse RGCs, but it is

unsuccessful in the isolation of RGCs from 3D retinal organoids [84]. A study reports the

purification of human RGCs via Thy1 from adherent cultures [85]. Tagging human RGCs with an

engineered marker Thy1.2 leads to efficient isolation of human RGCs [84], but may not be

unsuitable for clinical uses. Using single-cell RNA sequencing, we identified cell surface protein

CNTN2 as a specific marker for developing human RGCs. We isolated human

electrophysiologically-excitable RGCs using MACS via CNTN2. Therefore, we establish a one-

step method for isolating human RGCs under a native condition, facilitating therapeutic studies

for RGC-related retinal diseases such as glaucoma.

Disclosure – A patent application is pending.

Author Contributions

WL conceived and led the project, performed experiments, analyzed data, drew conclusions,

and wrote the paper. RD performed experiments, analyzed data, and edited the paper. AL

performed experiments. XZ analyzed the single-cell RNA sequencing dataset. LS designed,

performed, and analyzed patch clamp experiments and wrote the section. All authors discussed

the results and contributed to the final manuscript.

Acknowledgements

We are grateful to Dr. Roy Chuck for support, M.R.G. Stem Cell Institute (supported by

NYSTEM C029154) for service, Leslie Cummins and Frank P Macaluso at the Analytical

Imaging Facility of AECOM for electron microscopy (supported by P30CA013330), David

Reynolds at the Genomics Core of AECOM for the preparation of single-cell libraries, Dr.

Kamran Khodakhah for technical support and critical readings, and grants from NEI, NIH

(R01EY022645 and R21EY029806 to W.L.). Albert Lowe was supported by training grants

T32GM007491 and C30292GG. Ludovic Spaeth was supported by a grant from NIH

(5R01NS105470-06 to Dr. Khodakhah).

Materials and Methods

Maintenance of hESCs

ESCRO and IRB committees at AECOM approved the use of hESCs in this project.

Undifferentiated H1 hESCs (WiCell WA01) or hiPSCs (Corriell Institute AICS 0023) were grown

on Matrigel-coated 6-well plates in mTeSR1 medium and passaged using ReLeSR (STEMCELL

technologies) following manufacturer instructions.

Retinal cell differentiation

CONCEPT telencephalon-eye organoids were generated as follows. A humidified incubator at

37°C with 5% CO2 was used for cell culture. H1 hESCs or iPSCs that were passaged using

ReLeSR two or three days before experiments were detached using Dispase (GIBCO 17105041)

and then harvested by centrifugation. After that, the cell pellets were suspended in ice-cold

Matrigel. After gelling at 37 °C for 15-20 minutes, the hESC/Matrigel clump was gently

dispersed in a N2B27 Medium (DMEM/F12+GlutaMAX (GIBCO):Neurobasal medium (GIBCO)

= 1:1, 0.5 x B27 supplement (GIBCO), 0.5 x N2 supplement (GIBCO), 0.1 mM β-

mercaptoethanol, and 0.2 mM L-GlutaMax) for floating culture. With the starting day of cell

differentiation designated as day 0, cysts with a single lumen formed on day 1. Cysts with a

single lumen constituted over 85% of cultures. On day 4 or 5, individual cysts with a diameter at

around 150-200 μm were manually picked using a curved Pasteur pipets under an inverted

microscope and then seeded onto Matrigel-coated 24-well plates at a density of 2-6 cysts per

well. Cysts spontaneously attached to the culture surface and grew. From a time during days

13-16, attached cell colonies were grown in a KSR medium (GMEM medium supplemented with

10% knockout serum replacement, 1 mM sodium pyruvate, 0.1 mM non-essential amino acids,

2 mM l-glutamine, and 55 μM 2-mercaptoethanol; all the culture reagents were from Life

Technologies). Culture mediums were changed every two or three days. Overall, the procedure

is efficient and robust. Images in the manuscript represent 80-90% of cultures.

Inhibition of FGF signaling in CONCEPT telencephalon-eye organoids with FGFR

inhibitor PD 161570

To inactivate FGF signaling in CONCEPT organoids, FGFR1/2/3 inhibitor PD 161570 (1 µM;

Tocris) was supplemented to the culture medium staring on day 17, with vehicle DMSO as a

control. Treated CONCEPT organoids were harvested on day 24 for assays.

Magnetic-activated cell sorting (MACS) of developing human RGCs

Retinal organoids in suspension culture were generated using the established method [28,33]. For

each MACS experiment, 84 – 140 retinal organoids at stages of day 41 – 70 were dissociated

into single cells using Accutase (GIBCO A1110501). Non-retinal cells were trimmed if there

were any. Dissociated single cells were harvested using centrifugation and then incubated for

25 minutes at room temperature with MagnaBind goat anti-mouse IgG (ThermoScientific 21354)

beads that were previously coupled with a CNTN2 antibody (DSHB 4D7) following manufacturer

instructions. Cells bound to the beads were isolated using a magnetic rack and then washed

one time with the KSR medium supplemented with antibiotic:Antimycotic (GEMINI 400101)

while the tube was still against the magnetic rack. After the wash, the cells were released from

the beads via Accutase digestion for 30 minutes and then harvested using centrifugation. The

isolated cells were plated onto a chamber slide (ibidi 80826, ibidiTreat µ-Slide 8 Well, coated

with poly-ornithine and Matrigel, 30,000-50,000 cells/200 µl/well) in BrainPhys neuronal medium

(Stem Cell Technology 05790) supplemented with N2 and B27 (GIBCO 17502001, A3582801).

From 100 retinal organoids on day 41 to 48, around 385,000 RGCs were isolated. After 10 days

of culture in chamber slides, RGCs were fixed in 4% paraformaldehyde (PFA) for 10 minutes

and then processed for immunostaining.

Immunostaining, antibodies, and light microscopy

CONCEPT telencephalon-eye organoids were fixed in 4% PFA for 15-30 minutes at room

temperature and processed for immunostaining. See Supplemental Information for details.

In situ hybridization

DIG-labeled anti-sense RNA probes for in situ hybridization were generated via in vitro

transcription using a DIG RNA labeling kit (Millipore Sigma-Aldrich 11175025910). See

Supplemental Information for details. To assess the in vivo expression of DEGs identified by

single-cell RNA sequencing, in situ hybridization images of their mouse orthologs in E14.5

mouse brains were downloaded from a public database [86] (https://gp3.mpg.de/) with permission

and then assembled in supplemental Figs.

Electron microscopy (EM)

EM was performed by Analytical Imaging Facility in Albert Einstein College of Medicine with a

standard method. Lens cell clusters were fixed in 0.1M Cacodylate buffer containing 2%

paraformaldehyde and 2.5% glutaraldehyde for 60 minutes at room temperature and then

processed for EM.

Single-cell RNA sequencing

CONCEPT telencephalon-eye organoids at day 24 from one culture well were dissociated into

single cells using activated Papain (Worthington Biochemical Corporation LS003126) following

manufacturer's instructions. Then, 10,000 dissociated cells were captured using Chromium

Controller (10x Genomics), followed by library preparation using Single Cell 3' version 3.1 kit

(10x Genomics) following manufacturer's instructions. The library was sequenced in one lane of

Illumina HiSeq (2x150 bp) in GeneWIZ company.

Fastq sequences were mapped to the human genome (GRCh38-3.0.0) using CellRanger (3.1.0)

to generate a count matrix, which was then analyzed using the Seurat Package (v3.2.0) [59].

Sequenced cells were filtered (nFeature_RNA > 200 & nFeature_RNA < 6000 & percent.mt <

20), resulting in 10218 cells with high quality. Dimension reduction and clustering were

performed using the following functions: *NormalizeData, FindVariableFeatures, ScaleData,*

RunPCA, ElbowPlot, FindNeighbors (dims = 1:17), and *FindClusters (resolution = 0.5).*

Differentially expressed genes were identified using the function *FindAllMarkers (only.pos =*

FALSE, min.pct = 0.25, logfc.threshold = 0.25).

Whole-cell patch clamp recordings of isolated RGCs in culture.

Standdard methods were used. See Supplemental Information for details.

Statistical analysis

Statistical analysis in DEG identification was performed using the Seurat Package (v3.2.0) [59].

Adjusted p-values were shown.

Data availability

Single-cell RNA sequencing data of CONCEPT telencephalon-eye organoids at day 24 is

available in Gene Expression Omnibus (http://www.ncbi.nlm.nih.gov/geo) using the accession

number GSExxx.

Reference:

615   1.   Inoue, T., Nakamura, S., and Osumi, N. (2000). Fate mapping of the mouse
616   prosencephalic neural plate. Dev Biol *219*, 373-383. 10.1006/dbio.2000.9616.
617   2.   Oster, S.F., Deiner, M., Birgbauer, E., and Sretavan, D.W. (2004). Ganglion cell axon
618   pathfinding in the retina and optic nerve. Semin Cell Dev Biol *15*, 125-136.
619   10.1016/j.semcdb.2003.09.006.
620   3.   Erskine, L., and Herrera, E. (2007). The retinal ganglion cell axon's journey: insights into
621   molecular mechanisms of axon guidance. Dev Biol *308*, 1-14.
622   10.1016/j.ydbio.2007.05.013.
623   4.   Erskine, L., and Herrera, E. (2014). Connecting the retina to the brain. ASN Neuro *6*.
624   10.1177/1759091414562107.
625   5.   Wilson, S.W., and Rubenstein, J.L. (2000). Induction and dorsoventral patterning of the
626   telencephalon. Neuron *28*, 641-651. S0896-6273(00)00171-9 [pii].
627   6.   Xuan, S., Baptista, C.A., Balas, G., Tao, W., Soares, V.C., and Lai, E. (1995). Winged
628   helix transcription factor BF-1 is essential for the development of the cerebral
629   hemispheres. Neuron *14*, 1141-1152. 10.1016/0896-6273(95)90262-7.
630   7.   Zuber, M.E., Gestri, G., Viczian, A.S., Barsacchi, G., and Harris, W.A. (2003).
631   Specification of the vertebrate eye by a network of eye field transcription factors.
632   Development *130*, 5155-5167. 10.1242/dev.00723.
633   8.   Marquardt, T., Ashery-Padan, R., Andrejewski, N., Scardigli, R., Guillemot, F., and
634   Gruss, P. (2001). Pax6 is required for the multipotent state of retinal progenitor cells. Cell
635   *105*, 43-55.
636   9.   Ashery-Padan, R., Marquardt, T., Zhou, X., and Gruss, P. (2000). Pax6 activity in the
637   lens primordium is required for lens formation and for correct placement of a single
638   retina in the eye. Genes Dev *14*, 2701-2711. 10.1101/gad.184000.
639   10.   Baumer, N., Marquardt, T., Stoykova, A., Spieler, D., Treichel, D., Ashery-Padan, R.,
640   and Gruss, P. (2003). Retinal pigmented epithelium determination requires the redundant
641   activities of Pax2 and Pax6. Development *130*, 2903-2915.
642   11.   Horsford, D.J., Nguyen, M.T., Sellar, G.C., Kothary, R., Arnheiter, H., and McInnes, R.R.
643   (2005). Chx10 repression of Mitf is required for the maintenance of mammalian
644   neuroretinal identity. Development *132*, 177-187. dev.01571 [pii]
645   10.1242/dev.01571.
646   12.   Rowan, S., Chen, C.M., Young, T.L., Fisher, D.E., and Cepko, C.L. (2004).
647   Transdifferentiation of the retina into pigmented cells in ocular retardation mice defines a
648   new function of the homeodomain gene Chx10. Development *131*, 5139-5152.
649   10.1242/dev.01300
650   131/20/5139 [pii].
651   13.   Liu, W., Lagutin, O., Swindell, E., Jamrich, M., and Oliver, G. (2010). Neuroretina
652   specification in mouse embryos requires Six3-mediated suppression of Wnt8b in the
653   anterior neural plate. J Clin Invest *120*, 3568-3577. 10.1172/JCI43219.
654   14.   Bharti, K., Gasper, M., Ou, J., Brucato, M., Clore-Gronenborn, K., Pickel, J., and
655   Arnheiter, H. (2012). A regulatory loop involving PAX6, MITF, and WNT signaling
656   controls retinal pigment epithelium development. PLoS Genet *8*, e1002757.
657   10.1371/journal.pgen.1002757.

658   15.   Torres, M., Gomez-Pardo, E., and Gruss, P. (1996). Pax2 contributes to inner ear
659   patterning and optic nerve trajectory. Development *122*, 3381-3391.
660   16.   Macdonald, R., Scholes, J., Strahle, U., Brennan, C., Holder, N., Brand, M., and Wilson,
661   S.W. (1997). The Pax protein Noi is required for commissural axon pathway formation in
662   the rostral forebrain. Development *124*, 2397-2408.
663   17.   Martinez-Morales, J.R., Signore, M., Acampora, D., Simeone, A., and Bovolenta, P.
664   (2001). Otx genes are required for tissue specification in the developing eye.
665   Development *128*, 2019-2030.
666   18.   Schwarz, M., Cecconi, F., Bernier, G., Andrejewski, N., Kammandel, B., Wagner, M.,
667   and Gruss, P. (2000). Spatial specification of mammalian eye territories by reciprocal
668   transcriptional repression of Pax2 and Pax6. Development *127*, 4325-4334.
669   19.   Shimamura, K., and Rubenstein, J.L. (1997). Inductive interactions direct early
670   regionalization of the mouse forebrain. Development *124*, 2709-2718.
671   20.   Storm, E.E., Garel, S., Borello, U., Hebert, J.M., Martinez, S., McConnell, S.K., Martin,
672   G.R., and Rubenstein, J.L. (2006). Dose-dependent functions of Fgf8 in regulating
673   telencephalic patterning centers. Development *133*, 1831-1844. 133/9/1831 [pii]
674   10.1242/dev.02324.
675   21.   Soukkarieh, C., Agius, E., Soula, C., and Cochard, P. (2007). Pax2 regulates neuronal-
676   glial cell fate choice in the embryonic optic nerve. Dev Biol *303*, 800-813.
677   10.1016/j.ydbio.2006.11.016.
678   22.   Martinez-Morales, J.R., Del Bene, F., Nica, G., Hammerschmidt, M., Bovolenta, P., and
679   Wittbrodt, J. (2005). Differentiation of the vertebrate retina is coordinated by an FGF
680   signaling center. Developmental cell *8*, 565-574. 10.1016/j.devcel.2005.01.022.
681   23.   Vogel-Hopker, A., Momose, T., Rohrer, H., Yasuda, K., Ishihara, L., and Rapaport, D.H.
682   (2000). Multiple functions of fibroblast growth factor-8 (FGF-8) in chick eye
683   development. Mech Dev *94*, 25-36. 10.1016/s0925-4773(00)00320-8.
684   24.   Lancaster, M.A., and Knoblich, J.A. (2014). Organogenesis in a dish: modeling
685   development and disease using organoid technologies. Science *345*, 1247125.
686   10.1126/science.1247125.
687   25.   Eiraku, M., Takata, N., Ishibashi, H., Kawada, M., Sakakura, E., Okuda, S., Sekiguchi,
688   K., Adachi, T., and Sasai, Y. (2011). Self-organizing optic-cup morphogenesis in three-
689   dimensional culture. Nature *472*, 51-56. 10.1038/nature09941.
690   26.   Meyer, J.S., Howden, S.E., Wallace, K.A., Verhoeven, A.D., Wright, L.S., Capowski,
691   E.E., Pinilla, I., Martin, J.M., Tian, S., Stewart, R., et al. (2011). Optic vesicle-like
692   structures derived from human pluripotent stem cells facilitate a customized approach to
693   retinal disease treatment. Stem Cells *29*, 1206-1218. 10.1002/stem.674.
694   27.   Nakano, T., Ando, S., Takata, N., Kawada, M., Muguruma, K., Sekiguchi, K., Saito, K.,
695   Yonemura, S., Eiraku, M., and Sasai, Y. (2012). Self-formation of optic cups and storable
696   stratified neural retina from human ESCs. Cell Stem Cell *10*, 771-785.
697   10.1016/j.stem.2012.05.009.
698   28.   Lowe, A., Harris, R., Bhansali, P., Cvekl, A., and Liu, W. (2016). Intercellular Adhesion-
699   Dependent Cell Survival and ROCK-Regulated Actomyosin-Driven Forces Mediate Self-
700   Formation of a Retinal Organoid. Stem Cell Reports *6*, 743-756.
701   10.1016/j.stemcr.2016.03.011.
702   29.   Zhong, X., Gutierrez, C., Xue, T., Hampton, C., Vergara, M.N., Cao, L.H., Peters, A.,
703   Park, T.S., Zambidis, E.T., Meyer, J.S., et al. (2014). Generation of three-dimensional

704   retinal tissue with functional photoreceptors from human iPSCs. Nature communications
705   *5*, 4047. 10.1038/ncomms5047.

706 30.   Reichman, S., Terray, A., Slembrouck, A., Nanteau, C., Orieux, G., Habeler, W., Nandrot,
707   E.F., Sahel, J.A., Monville, C., and Goureau, O. (2014). From confluent human iPS cells
708   to self-forming neural retina and retinal pigmented epithelium. Proc Natl Acad Sci U S A
709   *111*, 8518-8523. 10.1073/pnas.1324212111.

710 31.   Cowan, C.S., Renner, M., De Gennaro, M., Gross-Scherf, B., Goldblum, D., Hou, Y.,
711   Munz, M., Rodrigues, T.M., Krol, J., Szikra, T., et al. (2020). Cell Types of the Human
712   Retina and Its Organoids at Single-Cell Resolution. Cell *182*, 1623-1640 e1634.
713   10.1016/j.cell.2020.08.013.

714 32.   Fligor, C.M., Lavekar, S.S., Harkin, J., Shields, P.K., VanderWall, K.B., Huang, K.C.,
715   Gomes, C., and Meyer, J.S. (2021). Extension of retinofugal projections in an assembled
716   model of human pluripotent stem cell-derived organoids. Stem Cell Reports.
717   10.1016/j.stemcr.2021.05.009.

718 33.   Kim, S., Lowe, A., Dharmat, R., Lee, S., Owen, L.A., Wang, J., Shakoor, A., Li, Y.,
719   Morgan, D.J., Hejazi, A.A., et al. (2019). Generation, transcriptome profiling, and
720   functional validation of cone-rich human retinal organoids. Proc Natl Acad Sci U S A
721   *116*, 10824-10833. 10.1073/pnas.1901572116.

722 34.   Langer, K.B., Ohlemacher, S.K., Phillips, M.J., Fligor, C.M., Jiang, P., Gamm, D.M., and
723   Meyer, J.S. (2018). Retinal Ganglion Cell Diversity and Subtype Specification from
724   Human Pluripotent Stem Cells. Stem Cell Reports *10*, 1282-1293.
725   10.1016/j.stemcr.2018.02.010.

726 35.   Fligor, C.M., Langer, K.B., Sridhar, A., Ren, Y., Shields, P.K., Edler, M.C., Ohlemacher,
727   S.K., Sluch, V.M., Zack, D.J., Zhang, C., et al. (2018). Three-Dimensional Retinal
728   Organoids Facilitate the Investigation of Retinal Ganglion Cell Development,
729   Organization and Neurite Outgrowth from Human Pluripotent Stem Cells. Sci Rep *8*,
730   14520. 10.1038/s41598-018-32871-8.

731 36.   Teotia, P., Chopra, D.A., Dravid, S.M., Van Hook, M.J., Qiu, F., Morrison, J., Rizzino,
732   A., and Ahmad, I. (2017). Generation of Functional Human Retinal Ganglion Cells with
733   Target Specificity from Pluripotent Stem Cells by Chemically Defined Recapitulation of
734   Developmental Mechanism. Stem Cells *35*, 572-585. 10.1002/stem.2513.

735 37.   Lancaster, M.A., Renner, M., Martin, C.A., Wenzel, D., Bicknell, L.S., Hurles, M.E.,
736   Homfray, T., Penninger, J.M., Jackson, A.P., and Knoblich, J.A. (2013). Cerebral
737   organoids model human brain development and microcephaly. Nature *501*, 373-379.
738   10.1038/nature12517.

739 38.   Kadoshima, T., Sakaguchi, H., Nakano, T., Soen, M., Ando, S., Eiraku, M., and Sasai, Y.
740   (2013). Self-organization of axial polarity, inside-out layer pattern, and species-specific
741   progenitor dynamics in human ES cell-derived neocortex. Proc Natl Acad Sci U S A *110*,
742   20284-20289. 10.1073/pnas.1315710110.

743 39.   Mariani, J., Simonini, M.V., Palejev, D., Tomasini, L., Coppola, G., Szekely, A.M.,
744   Horvath, T.L., and Vaccarino, F.M. (2012). Modeling human cortical development in
745   vitro using induced pluripotent stem cells. Proc Natl Acad Sci U S A *109*, 12770-12775.
746   10.1073/pnas.1202944109.

747 40.   Qian, X., Nguyen, H.N., Jacob, F., Song, H., and Ming, G.L. (2017). Using brain
748   organoids to understand Zika virus-induced microcephaly. Development *144*, 952-957.
749   10.1242/dev.140707.

41. Velasco, S., Kedaigle, A.J., Simmons, S.K., Nash, A., Rocha, M., Quadrato, G., Paulsen,
B., Nguyen, L., Adiconis, X., Regev, A., et al. (2019). Individual brain organoids
reproducibly form cell diversity of the human cerebral cortex. Nature *570*, 523-527.
10.1038/s41586-019-1289-x.
42. Gabriel, E., Albanna, W., Pasquini, G., Ramani, A., Josipovic, N., Mariappan, A.,
Schinzel, F., Karch, C.M., Bao, G., Gottardo, M., et al. (2021). Human brain organoids
assemble functionally integrated bilateral optic vesicles. Cell Stem Cell.
10.1016/j.stem.2021.07.010.
43. Etoc, F., Metzger, J., Ruzo, A., Kirst, C., Yoney, A., Ozair, M.Z., Brivanlou, A.H., and
Siggia, E.D. (2016). A Balance between Secreted Inhibitors and Edge Sensing Controls
Gastruloid Self-Organization. Developmental cell *39*, 302-315.
10.1016/j.devcel.2016.09.016.
44. Minn, K.T., Fu, Y.C., He, S., Dietmann, S., George, S.C., Anastasio, M.A., Morris, S.A.,
and Solnica-Krezel, L. (2020). High-resolution transcriptional and morphogenetic
profiling of cells from micropatterned human ESC gastruloid cultures. Elife *9*.
10.7554/eLife.59445.
45. Xue, X., Sun, Y., Resto-Irizarry, A.M., Yuan, Y., Aw Yong, K.M., Zheng, Y., Weng, S.,
Shao, Y., Chai, Y., Studer, L., and Fu, J. (2018). Mechanics-guided embryonic patterning
of neuroectoderm tissue from human pluripotent stem cells. Nat Mater *17*, 633-641.
10.1038/s41563-018-0082-9.
46. Hayashi, R., Ishikawa, Y., Sasamoto, Y., Katori, R., Nomura, N., Ichikawa, T., Araki, S.,
Soma, T., Kawasaki, S., Sekiguchi, K., et al. (2016). Co-ordinated ocular development
from human iPS cells and recovery of corneal function. Nature *531*, 376-380.
10.1038/nature17000.
47. Zhu, Y., Carido, M., Meinhardt, A., Kurth, T., Karl, M.O., Ader, M., and Tanaka, E.M.
(2013). Three-dimensional neuroepithelial culture from human embryonic stem cells and
its use for quantitative conversion to retinal pigment epithelium. PLoS One *8*, e54552.
10.1371/journal.pone.0054552.
48. Suda, Y., Kokura, K., Kimura, J., Kajikawa, E., Inoue, F., and Aizawa, S. (2010). The
same enhancer regulates the earliest Emx2 expression in caudal forebrain primordium,
subsequent expression in dorsal telencephalon and later expression in the cortical
ventricular zone. Development *137*, 2939-2949. 10.1242/dev.048843.
49. Danesh, S.M., Villasenor, A., Chong, D., Soukup, C., and Cleaver, O. (2009). BMP and
BMP receptor expression during murine organogenesis. Gene Expr Patterns *9*, 255-265.
10.1016/j.gep.2009.04.002.
50. Furuta, Y., and Hogan, B.L. (1998). BMP4 is essential for lens induction in the mouse
embryo. Genes Dev *12*, 3764-3775. 10.1101/gad.12.23.3764.
51. Dudley, A.T., and Robertson, E.J. (1997). Overlapping expression domains of bone
morphogenetic protein family members potentially account for limited tissue defects in
BMP7 deficient embryos. Dev Dyn *208*, 349-362. 10.1002/(SICI)1097-
0177(199703)208:3<349::AID-AJA6>3.0.CO;2-I.
52. Solloway, M.J., Dudley, A.T., Bikoff, E.K., Lyons, K.M., Hogan, B.L., and Robertson,
E.J. (1998). Mice lacking Bmp6 function. Dev Genet *22*, 321-339. 10.1002/(SICI)1520-
6408(1998)22:4<321::AID-DVG3>3.0.CO;2-8.

53.   Morcillo, J., Martinez-Morales, J.R., Trousse, F., Fermin, Y., Sowden, J.C., and
795   Bovolenta, P. (2006). Proper patterning of the optic fissure requires the sequential
796   activity of BMP7 and SHH. Development *133*, 3179-3190. 10.1242/dev.02493.
797   54.   Gan, L., Xiang, M., Zhou, L., Wagner, D.S., Klein, W.H., and Nathans, J. (1996). POU
798   domain factor Brn-3b is required for the development of a large set of retinal ganglion
799   cells. Proc Natl Acad Sci U S A *93*, 3920-3925.
800   55.   Mui, S.H., Kim, J.W., Lemke, G., and Bertuzzi, S. (2005). Vax genes ventralize the
801   embryonic eye. Genes Dev *19*, 1249-1259. 10.1101/gad.1276605.
802   56.   Bassett, E.A., Williams, T., Zacharias, A.L., Gage, P.J., Fuhrmann, S., and West-Mays,
803   J.A. (2010). AP-2alpha knockout mice exhibit optic cup patterning defects and failure of
804   optic stalk morphogenesis. Hum Mol Genet *19*, 1791-1804. 10.1093/hmg/ddq060.
805   57.   Biswas, S.K., Lee, J.E., Brako, L., Jiang, J.X., and Lo, W.K. (2010). Gap junctions are
806   selectively associated with interlocking ball-and-sockets but not protrusions in the lens.
807   Mol Vis *16*, 2328-2341.
808   58.   Lovicu, F.J., and McAvoy, J.W. (2001). FGF-induced lens cell proliferation and
809   differentiation is dependent on MAPK (ERK1/2) signalling. Development *128*, 5075-
810   5084.
811   59.   Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., 3rd, Hao,
812   Y., Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive Integration of
813   Single-Cell Data. Cell *177*, 1888-1902 e1821. 10.1016/j.cell.2019.05.031.
814   60.   Tirosh, I., Izar, B., Prakadan, S.M., Wadsworth, M.H., 2nd, Treacy, D., Trombetta, J.J.,
815   Rotem, A., Rodman, C., Lian, C., Murphy, G., et al. (2016). Dissecting the multicellular
816   ecosystem of metastatic melanoma by single-cell RNA-seq. Science *352*, 189-196.
817   10.1126/science.aad0501.
818   61.   Yao, Z., Mich, J.K., Ku, S., Menon, V., Krostag, A.R., Martinez, R.A., Furchtgott, L.,
819   Mulholland, H., Bort, S., Fuqua, M.A., et al. (2017). A Single-Cell Roadmap of Lineage
820   Bifurcation in Human ESC Models of Embryonic Brain Development. Cell Stem Cell *20*,
821   120-134. 10.1016/j.stem.2016.09.011.
822   62.   Poh, Y.C., Chen, J., Hong, Y., Yi, H., Zhang, S., Chen, J., Wu, D.C., Wang, L., Jia, Q.,
823   Singh, R., et al. (2014). Generation of organized germ layers from a single mouse
824   embryonic stem cell. Nat Commun *5*, 4000. 10.1038/ncomms5000.
825   63.   Varga, B.V., Faiz, M., Yang, H., Pivonkova, H., Gao, S., Khelifi, G., Linderoth, E., Zhen,
826   M., Hussein, S.M., and Nagy, A. (2021). Signal requirement for cortical potential of
827   transplantable human neuroepithelial stem cells. biorxiv
828   *https://doi.org/10.1101/2021.03.27.437311*. https://doi.org/10.1101/2021.03.27.437311.
829   64.   Simoes-Costa, M., and Bronner, M.E. (2015). Establishing neural crest identity: a gene
830   regulatory recipe. Development *142*, 242-257. 10.1242/dev.105445.
831   65.   Chatterjee, M., and Li, J.Y. (2012). Patterning and compartment formation in the
832   diencephalon. Front Neurosci *6*, 66. 10.3389/fnins.2012.00066.
833   66.   Martinez-Ferre, A., and Martinez, S. (2012). Molecular regionalization of the
834   diencephalon. Front Neurosci *6*, 73. 10.3389/fnins.2012.00073.
835   67.   Oster, S.F., Bodeker, M.O., He, F., and Sretavan, D.W. (2003). Invariant Sema5A
836   inhibition serves an ensheathing function during optic nerve development. Development
837   *130*, 775-784. 10.1242/dev.00299.
838   68.   Sandberg-Lall, M., Hagg, P.O., Wahlstrom, I., and Pihlajaniemi, T. (2000). Type XIII
839   collagen is widely expressed in the adult and developing human eye and accentuated in

the ciliary muscle, the optic nerve and the neural retina. Exp Eye Res *70*, 401-410.
10.1006/exer.1998.0826.

69. Chatzopoulou, E., Miguez, A., Savvaki, M., Levasseur, G., Muzerelle, A., Muriel, M.P.,
Goureau, O., Watanabe, K., Goutebroze, L., Gaspar, P., et al. (2008). Structural
requirement of TAG-1 for retinal ganglion cell axons and myelin in the mouse optic
nerve. J Neurosci *28*, 7624-7636. 10.1523/JNEUROSCI.1103-08.2008.

70. Yamagata, M., and Sanes, J.R. (2012). Expanding the Ig superfamily code for laminar
specificity in retina: expression and role of contactins. J Neurosci *32*, 14402-14414.
10.1523/JNEUROSCI.3193-12.2012.

71. Li, H., Wagner, E., McCaffery, P., Smith, D., Andreadis, A., and Drager, U.C. (2000). A
retinoic acid synthesizing enzyme in ventral retina and telencephalon of the embryonic
mouse. Mech Dev *95*, 283-289. 10.1016/s0925-4773(00)00352-x.

72. Take-uchi, M., Clarke, J.D., and Wilson, S.W. (2003). Hedgehog signalling maintains the
optic stalk-retinal interface through the regulation of Vax gene activity. Development *130*,
955-968. 10.1242/dev.00305.

73. Bedzhov, I., and Zernicka-Goetz, M. (2014). Self-organizing properties of mouse
pluripotent cells initiate morphogenesis upon implantation. Cell *156*, 1032-1044.
10.1016/j.cell.2014.01.023.

74. Coucouvanis, E., and Martin, G.R. (1995). Signals for death and survival: a two-step
mechanism for cavitation in the vertebrate embryo. Cell *83*, 279-287.

75. Chambers, S.M., Fasano, C.A., Papapetrou, E.P., Tomishima, M., Sadelain, M., and
Studer, L. (2009). Highly efficient neural conversion of human ES and iPS cells by dual
inhibition of SMAD signaling. Nat Biotechnol *27*, 275-280. 10.1038/nbt.1529.

76. Fernando, M., Lee, S., Wark, J.R., Xiao, D., Lim, B.Y., O'Hara-Wright, M., Kim, H.J.,
Smith, G.C., Wong, T., Teber, E.T., et al. (2022). Differentiation of brain and retinal
organoids from confluent cultures of pluripotent stem cells connected by nerve-like
axonal projections of optic origin. Stem Cell Reports *17*, 1476-1492.
10.1016/j.stemcr.2022.04.003.

77. Argelaguet, R., Clark, S.J., Mohammed, H., Stapel, L.C., Krueger, C., Kapourani, C.A.,
Imaz-Rosshandler, I., Lohoff, T., Xiang, Y., Hanna, C.W., et al. (2019). Multi-omics
profiling of mouse gastrulation at single-cell resolution. Nature *576*, 487-491.
10.1038/s41586-019-1825-8.

78. Tropepe, V., Hitoshi, S., Sirard, C., Mak, T.W., Rossant, J., and van der Kooy, D. (2001).
Direct neural fate specification from embryonic stem cells: a primitive mammalian neural
stem cell stage acquired through a default mechanism. Neuron *30*, 65-78. 10.1016/s0896-
6273(01)00263-x.

79. Munoz-Sanjuan, I., and Brivanlou, A.H. (2002). Neural induction, the default model and
embryonic stem cells. Nat Rev Neurosci *3*, 271-280. 10.1038/nrn786.

80. Redies, C., and Puelles, L. (2001). Modularity in vertebrate brain development and
evolution. Bioessays *23*, 1100-1111. 10.1002/bies.10014.

81. Martinez-Morales, J.R., Rodrigo, I., and Bovolenta, P. (2004). Eye development: a view
from the retina pigmented epithelium. Bioessays *26*, 766-777. 10.1002/bies.20064.

82. Kuwahara, A., Ozone, C., Nakano, T., Saito, K., Eiraku, M., and Sasai, Y. (2015).
Generation of a ciliary margin-like stem cell niche from self-organizing human retinal
tissue. Nature communications *6*, 6286. 10.1038/ncomms7286.

83.   Shirasaki, R., Lewcock, J.W., Lettieri, K., and Pfaff, S.L. (2006). FGF as a target-derived
chemoattractant for developing motor axons genetically programmed by the LIM code.
Neuron *50*, 841-853. 10.1016/j.neuron.2006.04.030.
84.   Sluch, V.M., Chamling, X., Liu, M.M., Berlinicke, C.A., Cheng, J., Mitchell, K.L.,
Welsbie, D.S., and Zack, D.J. (2017). Enhanced Stem Cell Differentiation and
Immunopurification of Genome Engineered Human Retinal Ganglion Cells. Stem Cells
Transl Med *6*, 1972-1986. 10.1002/sctm.17-0059.
85.   Rabesandratana, O., Chaffiol, A., Mialot, A., Slembrouck-Brec, A., Joffrois, C., Nanteau,
C., Rodrigues, A., Gagliardi, G., Reichman, S., Sahel, J.A., et al. (2020). Generation of a
Transplantable Population of Human iPSC-Derived Retinal Ganglion Cells. Front Cell
Dev Biol *8*, 585675. 10.3389/fcell.2020.585675.
86.   Visel, A., Thaller, C., and Eichele, G. (2004). GenePaint.org: an atlas of gene expression
patterns in the mouse embryo. Nucleic Acids Res *32*, D552-556. 10.1093/nar/gkh029.
898

     Figure legends

Figure 1. Generation of a telencephalon-eye organoid composed of <u>con</u>centric zones of

anterior <u>e</u>ctodermal <u>p</u>rogeni<u>t</u>ors (CONCEPT). (A) A scheme of the procedure. **(B)** A diagram

of CONCEPT organoids showing concentric zones of the anterior ectodermal progenitors. See

also Figs.1, 2, 3, 5, S1, S10. **(C)** Morphology of cysts at day 2 showing the epithelial structure

indicated by the apical reporter TJ::GFP at the lumen. **(D)** Morphology of CONCEPT organoids

at day 26. **(E-G)** Expression of telencephalon (Tel) marker Foxg1, neuroretinal (NR) markers

Vsx2 and Pax6 in mouse eyes at E10-10.5. Rostral optic stalk (OS) connected the telencephalic

vesicle to the optic cup. **(H-O)** FOXG1+ telencephalic progenitors, VSX2+ and/or PAX6+ retinal

progenitors formed concentric zones in CONCEPT organoids. N > 5 experiments. **(P-W)** In

CONCEPT organoids, morphogens FGF8, BMP4, and BMP7 mRNA expression started at early

stages and subsequently formed circular gradients. N > 5 experiments. Scale bars, 100 µm (C,

E, M, O, P, S, T), 200 µm (I, K), 500 µm (Q, U), 1 mm (D, H, J, L, N, R, V, W).

Fig. 2. Retinal ganglion cells (RGCs) grow axons toward and then along a path defined by

PAX2+ cell populations in CONCEPT telencephalon-eye organoids. N > 5 experiments. **(A-**

D) POU4F2+ RGCs grew TUBB3+ axons toward and then along a path with a circular or a

portion of circular shape. **(E, F)** In mice, Pax2 was expressed in central regions of the retina and

optic stalk at E10.5 (E) and in the optic disc and optic stalk at E13.5 (F). Tubb3+ axons from the

initial RGCs grew toward the optic disc, exited the eye, and navigated within the optic stalk (G).

(H-L) In CONCEPT organoids at day 26, TUBB3+ RGC axons grew toward and then along a

path defined by an adjacent PAX2+ VSX2+ cell population (arrowhead in H, brackets in I, J); the

PAX2+ VSX2- cell population sets up an inner boundary of RGC axon growth. **(L)** A diagram

summarizing RGC axon growth, PAX2+ optic disc (OD), and PAX2+ optic stalk (OS) in

CONCEPT organoids. The area labeled by the asterisk may appear as false signals in a low-

resolution printout but it is clearly a background in digital display. **(M)** A count of CONCEPT

925 organoids showing directional retinal ganglion cell axons. Scale bars, 50 μm (E, F, G), 100 μm

926 (B), 200 μm (A, H, I).

927 **Fig. 3. CONCEPT telencephalon-eye organoids contain lens cells that undergo terminal**

928 **differentiation.** N > 5 experiments. **(A-H)** In CONCEPT organoids, lens markers CRYAA and

929 CRY B were expressed at day 22 (A, B) and day 39 (C, D). Lens cells were not stained by DAPI

930 (E, F); they exhibited a crystal-like shape (G). A count of CONCEPT organoids with these lens

931 cells is shown (H). **(I, J)** When CONCEPT organoids were detached using Dispase at around

932 day 28 and grown in suspension, crystal-like clusters were found (I) and survived for months (J).

933 **(K-L)** Crystal-like lens clusters were free of organelles and exhibited ball-and-socket structures

934 (K, L), as revealed by electron microscopy. Scale bars, 100 μm (A, B, C, D, I), 200 μm (J), 5 μm

935 (K), 200 nm (L).

936 **Fig. 4. Cell clustering analysis identifies telencephalic cells, ocular cells, and two PAX2+**

937 **cell populations that mimic the optic disc and optic stalk, respectively.** See also suppl.

938 Figs. S2-S8. CONCEPT organoids at day 24 were used for single-cell RNA sequencing and

939 analyzed using Seurat (v3.2.0). **(A)** Identification of 14 cell clusters. **(B)** Cell cycle phases as

940 shown as cell cycle scores. **(C)** FOXG1 expression marked telencephalic cells. **(D, E)** The

941 expression of PAX6 and/or VSX2 marked retinal cells. **(F)** PAX2+ cells were found in two major

942 cell populations: PAX2+ VSX2+ cells were assigned as the optic disc (OD), whereas PAX2+

943 FOXG1+ VSX2- cells were assigned as the optic stalk (OS). **(G)** The expression of the optic

944 disc/stalk marker SEMA5A. **(H-L)** The expression of major DEGs in cluster 2, the major cell

945 population that mimic the optic disc. **(M-P)** The expression of major gene markers for PAX2+

946 VSX2- optic stalk cells. **(Q-T)** Identification of glycosylphosphatidylinositol (GPI)-anchored cell

947 membrane protein CNTN2 as a specific marker for developing human RGCs. Cluster 11

948 differentially expressed RGC markers, including CNTN2.

Fig. 5. RGCs grow CNTN2+ axons toward and then along a defined path in CONCEPT telencephalon-eye organoids and can be isolated in one step via CNTN2 in a native condition. N > 5 experiments. **(A-F)** PAX2+ cells formed two concentric zones mimicking the optic stalk (OS) and optic disc (OD), respectively (A-C; high magnifications in D, E). POU4F2+ RGCs grew CNTN2+ axons toward and then along a path defined by adjacent PAX2+ optic-disc cells (A-E). RGCs at a few hundreds of micrometers away from PAX2+ optic-disc cells grew axons centrifugally (arrow in C). At regions where there was a gap in PAX2+ optic-disc cells, CNTN2+ RGC axons exited the circular path and grew centrifugally (diamond arrowhead in A, double arrowheads in B and F). PAX2+ optic-stalk cells set up an inner boundary for RGC axon growth. **(G, H)** PAX2+ optic-disc cells did not express ALDH1A3; the cells that set up the boundaries of the path highly expressed ALDH1A3. **(I, J)** Cells that set up the inner boundary for RGC axon growth expressed VAX1/VAX2. **(K)** In E13.5 mouse eye, Aldh1a3 was highly expressed in the peripheral retina and a small region in the optic disc. Aldh1a3 expression was at low or nearly absent in the central retina. **(L-P)** One-step isolation of RGCs. RGCs from floating retinal organoids at day 41 (L, M) and day 70 (N-O) were dissociated into single cells using Accutase and then isolated using MACS via CNTN2 for 10-day growth. Isolated RGCs expressed POU4F2 and grew TUBB3+ neurites in random directions (L). RGCs also expressed CNTN2 (M), ISL1 (N), RBPMS (N), and SNCG (O); positive cells were counted (P). Scale bars, 200 μm (A,G,I), 100 μm (D-F,H,J,K), 50 μm (L).

Fig. 6. Electrophysiological features of RGCs. RGCs from retinal organoids on day 48 were isolated using MACS via a CNTN2 antibody and then grown on polymer coverslips in a chamber slide for 20-25 days before whole-cell patch clamp recordings. **(A)** Resting membrane potential of RGCs. *Black triangle: average; black line: median; box: interquartile range, n=9.* **(B)** RGCs can fire action potentials. Cells were patched in current-clamp mode. A steady current (I_m) was injected to maintain the membrane potential at -70mV and depolarizing current steps of 10ms

(*left*), 100ms (*middle*) or 1s (*right*) were injected to elicit action potentials (V_m). **(C)** RGCs show

functional voltage-gated currents. Cells were recorded in voltage-clamp (holding=-80mV) and

depolarizing voltage steps (200ms, +10mV steps up to 80mV, V_m) were applied to record inward

and outward voltage-gated currents (I_m). *Inset*: zoom on inward currents. **(D)** Outward currents

are primarily due to voltage-gated potassium channels. *Left*, representative example of a

current-voltage experiment performed in presence of 20mM Tetraethylammonium (TEA), a

blocker of voltage-gated potassium channels. *Inset*: zoom on inward currents. *Right*, amplitude

of potassium current as a function of membrane potential (*mean±SEM; $n_{control}$=5, n_{TEA}=5*). **(E)**

Inward currents result from activity of voltage-gated sodium channels. *Left*, representative

example of a current-voltage experiment performed in presence of 1µM Tetrodotoxin (TTX), a

blocker of voltage-gated sodium channels. *Inset*: zoom on inward currents. *Right*, amplitude of

sodium current as a function of membrane potential (*mean±SEM; $n_{control}$=5, n_{TTX}=3*).

Fig. 7. FGF signaling mediated by FGFRs is required for early RGC differentiation and

directional axon growth. (A) PAX2, FGF9, and FGF8 are differentially expressed in cluster 2,

the major component of PAX2+ optic-disc cells. **(B)** PAX2 mRNA expression in CONCEPT

organoids on day 25. Two PAX2+ concentric zones corresponding to the optic stalk (OS) and

optic disc (OD) are labeled. **(C)** Dual-color immunocytochemistry indicates the co-localization of

FGF8 and PAX2 in the optic-disc zone of CONCEPT organoids on day 25. **(D-E)** TUBB3+

axons grew towards and then along the cells that expressed high levels of FGF8 (D) and FGF9

mRNA (E) in CONCEPT organoids on day 25. Immunocytochemistry of TUBB3 was performed

after in situ hybridization. **(F-K)** After the inhibition of FGF signaling with FGFR inhibitor PD

161570 during days 17-24, the number of RGC somas drastically reduced, and directional axon

growth of RGCs was nearly absent, whereas FGF8 protein expression largely remained.

CNTN2 immunocytochemistry before (F, H) and after FGF8 immunocytochemistry (G, I, J, K)

are shown. N = 3/3 experiments. Scale bar, 250 µm (B), 100 µm (C-E), 1 mm (F), 200 µm (K).

Tab. 1. Cell counts of clusters in the scRNA-seq dataset.

Abbreviations: Idents, assigned cell identities; # Cells, cell number; Tel, telencephalon; R, retina;

OD, optic disc; OS, optic stalk; RPE, retina pigment epithelium; RGC, retinal ganglion cell; N,

early neurons; UD, undetermined.

 Supplemental Information

 Immunostaining, antibodies, and light microscopy

CONCEPT telencephalon-eye organoids were fixed in 4% PFA for 15-30 minutes at room

temperature and processed for immunostaining. These primary antibodies were used: FOXG1

(Abcam, ab18259, 1:500), TUBB3 (Covance MMS-435P, 1:1000), FGF8 (1:500, R&D MAB323),

RBPMS (1:200, PhosphoSolution 1830-RBPMS), ISL1 (1:500, DSHB 40.2D6), SNCG (1:200,

Abcam ab55424), PAX2 (Invitrogen, 716000, 1:200), alpha A crystallin (Santa Cruz sc 22743,

1:500, shown as CRYAA in figure panels), beta crystallin (Santa Cruz sc-22745, 1:100, shown

as CRY B in figure panels), CNTN2 (DSHB 4D7, 1:100), ALDH1A3 (Invitrogen, PA529188,

1:500), VAX1/2 (Santa Cruz sc-98613, 1:200), PAX6 (1:500, Covance PRB-278P), POU4F2

(Santa Cruz, SC-6026, 1:200), SIX3 (1:500, Rockland), and VSX2 (1:500, Millipore AB9016).

Primary antibodies were visualized using Alexa Fluor 488-, 568-, and 647-conjugated secondary

antibodies and imaged using a Zeiss AxioObserver Z1 microscope. When the sample did not fit

in one image, multiple images were stitched to obtain an overview. In dual-color

immunocytochemistry for PAX2 and FGF8, FGF8 was visualized using AP-conjugated anti

mouse secondary antibody (Invitrogen A16038), and PAX2 was detected by biotin-conjugated

anti rabbit secondary antibody (Invitrogen B2770) followed by HRP-conjugated streptavidin

(ThermoScientific 21130). For dual color immunocytochemistry of CNTN2 and FGF8 (both are

mouse antibodies; no working antibodies raised in different species are available),

immunocytochemistry of CNTN2 was performed first and visualized using HRP-conjugated

streptavidin. Then, the samples were fixed in 4% PFA for 15 minutes to avoid antibody crosstalk.

After that, FGF8 antibody was applied and then detected by an AP-conjugated secondary

antibody. Brown and blue colors are well separated in Fig. 5K, indicating that antibody

crosstalks did not exist.

PCR Primers for in situ hybridization probes

PCR primers are as follows: BMP4: forward, CGGAAGCTAGGTGAGTGTGG, reverse,

GAGtaatacgactcactatagggGGAAGCCCCTTTCCCAATCA; BMP7: forward, gaggtccctctccattccct,

reverse, GAGtaatacgactcactatagggtgcacccatcagacctccta; FGF8: forward,

GTTGCACTTGCTGGTCCTCT, reverse,

GAGtaatacgactcactatagggTTGAGTTTTGGGTGCCCTAC; PAX2: forward, gctgtctgtgctgtgagagt,

reverse, GAGtaatacgactcactatagggccggggacatttagcaggtt; SEMA5A: forward,

CAGAGGCTCAGGCACAATGA, reverse,

GAGtaatacgactcactatagggTCCGTGTCTACCCAGGACTT; CYP1B1: forward,

cccagcggttcttcatgagt, reverse, GAGtaatacgactcactataggggcacacttggttgcgttagt; LEFTY2: forward,

agccctctaactgaacgtgtg, reverse, GAGtaatacgactcactatagggtcttctgagtatctacattcaattgct; EMX2:

forward, ACCGAGAAAGGGAGAGGGAA, reverse,

GAGtaatacgactcactatagggTCGGCCAATTTCTCCAACCA; FGF9: forward,

GTCCGCTATGAACCTGTGGT, reverse,

GAGtaatacgactcactatagggATAGTCTCGCTTGCCCAAGG; FGF2: forward,

aacaccgaaatgctggaggt, reverse, GAGtaatacgactcactataggggagaccacatgtacacgcca.

Whole-cell patch clamp recordings of isolated RGCs in culture.

RGCs were isolated from retinal organoids on day 48 using MACS via a CNTN2 antibody and

then grown on polymer coverslips in chamber slides (ibidi 80826) for 20 – 27 days. At the time

of whole-cell patch clamping recordings, the polymer coverslips carrying RGCs were carefully

cut out with a scalpel blade and then placed in a recording chamber under an upright

microscope (Zeiss Examiner A1), containing artificial cerebrospinal fluid (aCSF) composed of (*in*

mM): NaCl (*140*), MgCl$_2$ (*1*), KCl (*5*), CaCl$_2$ (*2*), Hepes (*10*), Glucose (*10*). Osmolarity and pH

were adjusted to 300mOsm and 7.3 respectively. Whole-cell patch-clamp recordings in voltage

and current-clamp mode were obtained at room temperature using an Optopatch amplifier

(Cairn Research, UK) and acquired with WinWCP 5.2 freeware (John Dempster, SIPBS,

University of Strathclyde, UK). Patch pipettes (3-4MΩ when filled with corresponding solution)

were pulled from borosilicate capillaries using a horizontal puller (Sutter P97, USA) and coated

with dental wax to reduce the pipette capacitance. For current clamp recordings, patch pipettes

were filled with a solution containing (*in mM*): K-gluconate (*130*), Na-gluconate (*10*), NaCl (*4*),

Hepes (*10*), Phosphocreatin (*10*), MgATP (*4*), Na_2GTP (*0.3*). Osmolarity and pH were adjusted

at 295mOsm and 7.3 respectively. Resting membrane potential was obtained from averaging

membrane potential recorded for one minute in I=0 mode immediately after breaking the cell

membrane. In current clamp mode, with the cell hyperpolarized to -70 mV, current steps of 100

to 500pA were made to explore whether the cells were excitable. For voltage-clamp recordings,

cells were held at clamped potential of -80mV, and series resistance was monitored and

compensated (>80%). Membrane potentials were corrected for the liquid junction potential,

calculated at +5.4mV (https://swharden.com/LJPcalc). Patch pipettes were filled with a solution

containing (*in mM*): KCl (*140*), $MgCl_2$ (*5*), $CaCl_2$ (*2.5*), Hepes (*10*), MgATP (*4*), Na_2GTP (*0.3*).

Osmolarity and pH were adjusted at 295mOsm and 7.3 respectively. A -P/4 subtraction protocol

was used to isolate voltage-gated currents by removing linear leak current and capacitance

artifacts. In all cases, voltage and current recordings were low pass filtered at 3kHz and

digitized at 10-20kHz (Axon Digidata 1550b, Molecular Device, USA). Tetraethylammonium-

chloride (TEA, Sigma-Aldrich, USA) and Tetrodotoxin-citrate (TTX, Fisher Scientific) stocks

were diluted in aCSF to reach 20mM and 1µM respectively. All recordings were analyzed with

WinWcp 5.2 and custom scripts and routines written in Python 3.9. Statistical tests were

performed with corresponding function from *scipy.stats* package (v1.8).

1074

1075

Legends for supplemental Figures

Fig. S1. Reproducibility of CONCEPT telencephalon-eye organoids is demonstrated by consistent gene expression profiles of multiple organoids in whole culture wells. Related to Figs. 1, 7. **(A-C)** CONCEPT organoids on day 17 expressed FGF8 (A, n =5/5), BMP4 (B, n = 4/4; the broken colony was not counted since its morphology is not quite clear), and BMP7 (C, n = 4/4) in concentric zones. **(D-F)** CONCEPT organoids on day 25 expressed FGF8 (D, n = 5/8), RGC marker CNTN2 (E, n = 5/6), and telencephalon marker EMX2 (F, n = 2/2) in concentric zones. Scale bars, 1 mm (A-F).

Fig. S2. Mesoderm, endoderm, and neural crest markers are not expressed in CONCEPT telencephalon-eye organoids at 24. Related to Fig. 4. The dataset for Fig. 4 is used for plotting. Expression profiles of mesoderm markers TBXT, GATA2, and HAND1, endoderm markers GATA1, GATA4, and SOX17, neural crest markers SNAI1, SOX10, FOXD3 are shown.

Fig. S3. The expression of top DEGs in cluster 0. Related to Fig. 4. The dataset for Fig. 4 is used for plotting. Cluster 0 was marked by DEGs that were lowly expressed.

Fig. S4. Diencephalon markers and midbrain/hindbrain markers are rarely expressed in CONCEPT telencephalon-eye organoids at day 24. Related to Fig. 4. **(A-F)** Diencephalon markers GBX2, WNT3, and SOX14, and midbrain/hindbrain markers EN2, PAX7, and TFAP2B were rarely expressed. The dataset for Fig. 4 is used for plotting. **(G-L)** Expression of the diencephalon markers and midbrain/hindbrain markers in the E14.5 mouse brain. The images were downloaded from a public database (https://gp3.mpg.de/) with permission. Image IDs were shown following gene symbols.

Fig. S5. DEGs for telencephalic clusters in CONCEPT telencephalon-eye organoids include markers for both the pallium and subpallium. Related to Fig. 4 and Fig. S6. DEGs

for telencephalic clusters were used for plotting, and expression of their orthologs in E14.5

mouse brain was shown in Fig. S6. **(A-D)** Expression of the DEGs POU3F3, RGMA, EDNRB,

and SOX13, which orthologs are expressed in the pallium of E14.5 mouse brain (see also Fig.

S6B-E). **(E-L)** Expression of the DEGs DLX2, RGS16, DLX1, DLL1, DLX6-AS1, NEFL, DCX,

and RTN1, which orthologs are expressed in the subpallium of E14.5 mouse brain (see Fig.

S6F-M).

Fig. S6. Mouse orthologs of the telencephalic DEGs are expressed in both the pallium

and subpallium of the E14.5 mouse brain. Related to Fig. 4 and Fig. S5. The images were

downloaded from a public database (https://gp3.mpg.de/) with permission. Image IDs were

shown following gene symbols. **(A)** Expression of Foxg1. **(B-E)** Expression of the genes Pou3f3,

Rgma, Ednrb, and Sox3 in the pallium of E14.5 mouse brain. **(F-M)** Expression of the genes

Dlx2, Rgs16, Dlx1, Dll1, Dlx6os1, Nefl, Dcx, and Rtn1 in the subpallium of E14.5 mouse brain.

Fig. S7. DEGs in cluster 6 include RPE markers. Related to Fig. 4. Top DEGs in cluster 6

include RPE markers PMEL, HSD17B2, DCT, and MITF.

Fig. S8. Mouse orthologs of the markers for Pax2+ optic disc and Pax2+ optic stalk cell

clusters in CONCEPT organoids are indeed expressed in the optic disc and optic

stalk/nerve, respectively, in E14.5 mouse embryos. Related to Fig. 4. In situ hybridization

images of the E14.5 mouse brain were downloaded from a public database (https://gp3.mpg.de/)

with permission. Image IDs were shown following gene symbols. **(A-T)** Expression of the gene

markers in E14.5 brain is shown.

Fig. S9. CONCEPT telencephalon-eye organoids are generated using hiPSC line AICS

0023. Related to Fig. 7. hiPSCs (AICS 0023) were used for the generation of CONCEPT

telencephalon-eye organoids. **(A-D)** POU4F2+ RGCs grew CNTN2+ axons toward and then

along a circular path defined by PAX2+ cell populations. Scale bar, 500 µm (A), 100 µm (B).

Fig. S10. Expression signatures of cell cluster 2. Related to Figs. 4, 5. **(A)** A list of top 10

DEGs and additional DEG markers SEMA5A, PAX2, COL13A1, FGF9, and FGF8 in cluster 2

were identified using Seurat analysis. **(B-E)** Expression of PAX2, SEMA5A, CYP1B1, and

LEFTY2 in CONCEPT organoids at day 25 was revealed by in situ hybridization. The PAX2+

optic disc at the outer zone was marked (bracket in B).

Fig. S11. FGFR1, FGFR2, FGFR3, MAP2K1, and MAP2K2 are expressed in multiple types

of cells in CONCEPT organoids. Related to Fig. 4. In RGCs, FGFR1 and MAP2K2 were

expressed.

Fig. 1

A N2B27   KSR

Day 0   Day 1   Day 4, 5   Day 6   Day 13 to 16 onward

Embed hESC sheets in Matrigel → Disperse gently → Clumps of Matrigel/ hESCs → Grow in suspension → Cysts → Passage 2-6 cysts per well → Attached colonies → Concentric zones of anterior ectodermal progenitors (CONCEPT)

B RGC / FOXG1+ / PAX2+ / VSX2+

CONCEPT at day 22-26

80 CONCEPT / Non 9

Count

C Cysts Day 2 — TJP1:: GFP — 100 u

D CONCEPT Day 26 — 1mm

E NR / OS / Tel — E10 Foxg1

F E10 Vsx2

G E10.5 Pax6

H Day 15 FOXG1 G VSX2 R — 1 mm

I 200 u

J Day 17 PAX6 G VSX2 R — 1 mm

K 200 u

L Day 26 FOXG1 G VSX2 R — 1 mm

M 100 u

N Day 26 PAX6 G VSX2 R — 1mm

O 100 u

P BMP4 Day 10 — 100 u

Q BMP4 Day 17 — 500 u

R BMP4 Day 25 — 1 mm

S BMP7 Day 10 — 100 u

T FGF8 Day 10 — 100 u

U FGF8 Day 17 — 500 u

V FGF8 Day 25 — 1 mm

W BMP7 Day 25 — 1 mm

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Cell Counts			
	Pos.	DAPI	%
SNCG	66	66	100
TUBB3	53	55	96.4
POU4F2	37	55	67.3
ISL1	15	35	42.9
RBPMS	7	35	20

Fig. 6

Fig. 7

gene	p_val	avg_logFC	pct.1	pct.2	p_val_adj	cluster
PAX2	0	0.930533006	0.769	0.169	0	2
FGF9	0	0.784668891	0.69	0.161	0	2
FGF8	1.77E-213	0.465810314	0.381	0.07	5.94E-209	2

CNTN2 (Brown) FGF8 (Blue)

DMSO on day 17-24

FGFRi PD161570 on day 17-24

Tab.1. Cell counts of clusters in the scRNA-seq dataset

Cluster	0	1	2	3	4	5	6	7	8	9	10	11	12	13
Idents	Tel/R	Tel	OD	Tel	Tel/OS	NR	RPE	NR	Tel/OS	Tel/OS	Tel	RGC	Tel	UD
# Cells	1295	1158	964	931	921	871	747	743	729	603	450	393	333	80

Abbreviations: Idents, assigned cell identities; # Cells, cell number; Tel, telencephalon; R, retina; OD, optic disc; OS, optic stalk; RPE, retina pigment epithelium; RRG, retinal ganglion cell; UD, undetermined.