Title: Metabolic perturbations to an *E. coli*-based cell-free system reveal a trade-off between transcription and translation

Author list: Manisha Kapasiawala[1]*, Richard M. Murray[1]

Author address:

[1]Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, CA 91125, USA

*Corresponding author: mkapasia@caltech.edu

Abstract: Cell-free transcription-translation (TX-TL) systems have been used for diverse applications, but their performance and scope are limited by variability and poor predictability. To understand the drivers of this variability, we explored the effects of metabolic perturbations to an *E. coli* Rosetta2 TX-TL system. We targeted three classes of molecules: energy molecules, in the form of nucleotide triphosphates (NTPs); central carbon "fuel" molecules, which regenerate NTPs; and magnesium ions (Mg^{2+}). Using malachite green mRNA aptamer (MG aptamer) and destabilized enhanced Green Fluorescent Protein (deGFP) as transcriptional and translational readouts, respectively, we report the presence of a universal trade-off between optimizing total protein yield and optimizing total mRNA yield, as measured by integrating the area under the curve for mRNA time-course dynamics. We found that a system's position along the trade-off curve is strongly determined by Mg^{2+} concentration, cell lysate batch, and fuel type. Our results suggest that these previously identified sources of variability ultimately stem from the cell lysate, and that variability can be reduced by modulating these components. Our results further suggest the trade-off arises from limitations in translation regulation and inefficient energy regeneration. This work advances our understanding of the effects of fuel and energy metabolism on TX-TL in cell-free systems and lays a foundation for improving TX-TL performance, lifetime, standardization, and prediction.

Keywords: *energy metabolism, cell-free systems, trade-off, magnesium ion, transcription-translation (TX-TL), batch-to-batch variability*

Introduction:

Cell-free synthetic biology is a rapidly growing field that leverages synthetic biology tools and techniques to engineer biological systems outside the traditional context of living cells[1]. The cell-free approach offers many advantages over engineering cellular systems, including greater control over the reaction environment, the ability to produce toxic small molecules or proteins that are difficult to express in living cells, and scalability for biomanufacturing and other applications. In recent years, cell-free systems have been used for protein engineering[1,2], rapid and high-throughput prototyping of engineered biological components and systems[3–7], biosensing and diagnostics[8–10], and the development of synthetic life[11–13].

Most cell-free applications rely on cell-free protein synthesis, where crude cell lysates or purified proteins are supplemented with DNA and an energy buffer containing small molecules that enable *in vitro* transcription and translation (TX-TL)[14–16]. Although cell lysate-based TX-TL systems have many uses, their performance and scope are limited by issues such as batch-to-batch variability[17], lack of predictability[18], and limited lifetime[19]. These issues arise because cell lysate's proteomic composition largely reflects the cytosolic protein content of living cells. While some of these proteins are harnessed to regenerate energy molecules for TX-TL processes, most of them generate metabolites that are benign or even harmful for TX-TL.

To address these concerns, previous studies have sought to gain insight into TX-TL dynamics through a variety of experiment approaches. Most of these studies have focused on gaining detailed insight into how perturbations to TX-TL reaction conditions – including salts[20,21], fuel sources[22,23], crowding agents[20,24], and other components[15,19,25–28] – affect TX-TL performance. Some studies have taken a less targeted approach, with the authors opting instead to harness lab automation and machine learning along with experiments to rapidly optimize TX-TL systems[27,29]. While these studies have helped elucidate which reaction components and preparation steps affect TX-TL performance and predictability most significantly, more work is necessary to determine why these factors contribute to variability and limited TX-TL lifetime, as well as the broader implications they have for design principles for TX-TL systems.

Aiming to get a more holistic view of cell-free reaction dynamics, more recent studies have focused on measuring small molecules – including NTPs, amino acids, and central carbon metabolites – that participate in cell-free metabolism and studying or modeling their effects on TX-TL[30–33]. In one such study, the authors found that the majority of *E. coli* metabolic pathways are active during TX-TL[33], and they subsequently used these insights to form a phenomenological model of TX-TL coupled to cell-free metabolism, with the model containing "fuel," "energy," and "waste" species and their effects on TX-TL. By linking cell-free metabolism's effects on TX-TL dynamics to specific TX-TL reaction components, these studies have suggested that further insight into TX-TL predictability, variability, and lifetime could come from further investigation of the effects of metabolism of TX-TL.

Building off the experiments and insights from these previous studies, to understand the effect of cell-free metabolism on variability, we focused our efforts on the chemical composition of the

buffer that is used with cell lysate and DNA to form a TX-TL reaction. We targeted two classes of small molecules to modulate: "energy" molecules, specifically nucleotide triphosphates (NTPs), which power TX-TL processes; and "fuel" molecules, which regenerate NTPs via cell lysate metabolism. We also considered the effects of magnesium (Mg^{2+}), both because of its biological importance – in stabilizing nucleic acids and ribosomes, acting as an essential cofactor for many enzymes, and making ATP bioactive in the form of Mg-ATP – and because of several previous findings suggesting its role in modulating TX-TL dynamics[21,24,25,28].

Using mRNA aptamer Malachite Green (MG aptamer) and destabilized enhanced Green Fluorescent Protein (deGFP) as transcriptional and translational readouts, respectively, we report the presence of a trade-off between optimizing total protein yield and optimizing total mRNA yield, as measured by integrating the area under the curve for mRNA time-course dynamics. We found that the trade-off is universal across different fuel sources, that a system's position along the trade-off curve is determined strongly by Mg^{2+} concentration and cell lysate batch, and that the trade-off curve's location shifts and size becomes larger as DNA concentration is increased. In systems where transcription and translation were decoupled, we found that a distinct regime optimized for translation exists in the translation-only system, but that no individual regime was optimized for transcription in the transcription-only system, suggesting that the trade-off arises at the translational level. Finally, in systems where additional energy is supplied and where a fuel source is absent, the trade-off is absent. Overall, our results suggest the trade-off arises from limitations in translation regulation and inefficient energy regeneration. By improving our understanding of the effects of fuel and energy metabolism on TX-TL in cell-free systems, this work provides insight into design considerations for future studies aimed at improving TX-TL performance, lifetime, standardization, and prediction.

Results and Discussion:

E. coli Rosetta2 cell-free systems exhibit a trade-off between transcription and translation across fuel sources and concentrations

As most previous studies have utilized either a design-of-experiments approach (e.g., randomly surveying many reaction compositions) or have modulated components one at a time, we chose to simultaneously vary fuel and Mg^{2+} concentrations. We performed TX-TL reactions consisting of fuel versus Mg^{2+} panels, where the former was varied between 0 - 45 mM, the latter was simultaneously varied between 0-10mM, and all other reaction components were kept the same[14] (see **Materials and Methods** for more details) (**Figure 1A**). As the preparation of cell lysate varies significantly from lab to lab and is difficult to keep consistent[18], we performed experiments in two cell lysate batches to see if these chemical factors acted consistently.

For fuel molecules, we focused on four fuel metabolites with varying degrees of energy regeneration and waste generation capacity that spanned central carbon metabolism to gain insight into how different fuel molecules affect TX-TL through their effect on metabolism. We chose four central carbon metabolites: 3-phosphoglyceric acid (3PGA), maltose, pyruvate, and succinate (**Figure 1B**). We targeted 3PGA – whose conversion to pyruvate regenerates ATP

during glycolysis – because it is one of the standard molecules[14] used for energy regeneration in cell lysate-based TX-TL systems (the others being phosphoenolpyruvate (PEP) and pyruvate). We chose pyruvate because it is located downstream of 3PGA in the glycolysis pathway and can also participate in energy regeneration, albeit without contributing potential waste molecule inorganic phosphate (P_i) to the system[21]. We selected maltose, which consumes P_i and outputs a glucose molecule and glucose-6-phosphate (G6P) molecule, because it has been used previously to improve energy regeneration by recycling P_i [34]. However, as one maltose molecule can be used to generate two G6P molecules, and thus four 3PGA molecules, maltose produces more candidate waste molecules per molecule of fuel compared to the other three fuels. These candidate waste molecules include acetate, ethanol, lactate, and formate (and their conjugate organic acids), which may inhibit TX-TL either directly or indirectly, by reducing pH too low for optimal enzyme activity[35–37]. Finally, we considered succinate, whose conversion from succinyl-CoA regenerates GTP. As the succinyl-CoA to succinate reaction is reversible, we believed the addition of succinate could slow down GTP regeneration by favoring the reverse reaction, thereby reducing the overall rate of energy regeneration to a level where energy could be used more efficiently by TX-TL.

TX-TL reactions were supplied with one of two different batches of cell lysate along with plasmid DNA encoding the sequence for the expression of MG aptamer, which fluoresces upon binding to malachite green dye, and deGFP, under a P_{OR1OR2} promoter. For each reaction condition, the total integrated area under the MG aptamer fluorescence curve and maximum deGFP were calculated as measurements of transcription and translation, respectively. The results of these experiments are shown in **Figure 1**.

By plotting maximum deGFP versus integrated MG aptamer for all fuel sources across different fuel and Mg^{2+} concentrations, we observed a trade-off between optimizing maximum deGFP and optimizing integrated MG aptamer (**Figure 1C, D**). The same trade-off curve was observed across all fuel sources but was not observed when no fuel was added (**Figure 1C**). Furthermore, a system's position along the trade-off curves was also unaffected by the concentration of fuel (**Figure 1D**). Control experiments indicated that the fluorescence of MG aptamer and deGFP were robust to changes in Mg^{2+} and pH across the relevant reaction values[22,23] and across different 3PGA and Mg^{2+} values (**Figure S1**), which suggested that expression of MG aptamer and deGFP, rather than their fluorescence alone, were changing across the various fuel and Mg^{2+} concentrations. These results suggest the presence of a universal trade-off between transcription and translation for the fuels considered here.

The results of this initial experiment were surprising, because while a system's position along the trade-off curve was weakly dependent on fuel type (**Figure 1C**), all points fell along the same trade-off curve, regardless of fuel type or lysate batch. These results seem to suggest that while these TX-TL systems with different fuel types and concentrations exhibit variability in transcriptional and translational performance, their performance is constrained to a single trade-off curve that could be traversed for a given fuel type and desired performance. These results also suggest the TX-TL trade-off originates from a fundamental trade-off in fuel and energy metabolism, perhaps arising from limitations in TX-TL and metabolism machinery.

Figure 1: A universal TX-TL trade-off. **(A)** Overview of experimental workflow. **(B)** Simplified map of *E. coli* core metabolism with NTP consumption and regeneration. A simplified glycolysis, TCA cycle, and interacting branch of the Pentose Phosphate Pathway are shown. Consumption or regeneration of other cofactors – such as NADH, FADH, QH_2, H_2O, and CO_2 – is omitted for simplicity. **(C)** and **(D)** show maximum deGFP values versus integrated MG aptamer values, colored by either (C) fuel type or (D) fuel concentration. Each point represents one of three replicates of a particular set of fuel and Mg^{2+} concentrations. The data shown reflect experiments performed in two batches of cell lysate, Batch 1 and Batch 2 (see Methods and Materials for details), and all experiments were performed using 5 nM DNA.

Variability along the trade-off curve is controlled by Mg²⁺ concentration and lysate batch

To investigate which factors controlled a system's position along the trade-off curve, we colored the data in **Figure 1** by Mg^{2+} concentration and cell lysate batch, the results of which are shown in **Figure 2**. We found that as the concentration of added Mg^{2+} increases, a TX-TL system generally shifts from a translation-optimized regime to a transcription-optimized regime (**Figure 2A**). We also found that while data for the two batches of lysate fell along the same curve, variability along the curve could be further explained by lysate batch (**Figure 2B**).

Figure 2: Effects of varying Mg^{2+} concentration and lysate batch on the TX-TL trade-off. These panels show the same data from Figures 1C and 1D colored by **(A)** the concentration of added Mg^{2+} (in addition to the Mg^{2+} that is already in the cell lysate from its preparation in S30 buffer) or **(B)** lysate batch. Each point represents one of three replicates of a particular set of fuel and Mg^{2+} concentrations. The data shown reflect experiments performed in two batches of cell lysate, Batch 1 and Batch 2, in systems using 5 nM DNA.

These provide clear design principles for experimental efforts aimed at improving performance and predictability of, and reducing variability in, TX-TL systems. With regards to predictability and performance, as the Mg^{2+} trend is universal across a variety of fuel sources (**Figure S2**), our results suggest that Mg^{2+} concentration can be tuned for a given combination of cell lysate batch, fuel type, and fuel concentration to design a system optimized for a particular behavior, whether that be optimizing transcription, optimizing translation, or some combination of both. For efforts aimed at reducing batch-to-batch variability, our results suggest that TX-TL systems prepared from different batches of lysate can be adjusted to behave similarly by tuning their fuel and Mg^{2+} concentrations such that the systems fall along the same position on the trade-off curve. As Mg^{2+} is an essential cofactor for many proteins, these results also suggest the TX-TL trade-off originates in part from Mg^{2+}-dependent regulation of cell-free machinery.

Inspecting **Figure 2A** more closely revealed that the Mg^{2+} trend was not completely consistent; plotting the trade-off separately for individual fuel sources revealed that 3PGA-fueled systems had a weaker translation optimization to transcription optimization trend as Mg^{2+} concentration was increased, when compared to the other fuels (**Figure S2A**). Specifically, while TX-TL systems fueled by pyruvate, succinate, and maltose were able to optimize translation at 0mM added Mg^{2+} and optimize transcription at 10mM added Mg^{2+}, some 3PGA-fueled systems were still able to achieve high deGFP yields at high concentrations of added Mg^{2+}.

In 3PGA-fueled systems, increasing fuel pushes a system from a TX-optimized to a TL-optimized regime

To understand why 3PGA-fueled systems deviated from the Mg^{2+} trend, we plotted the data from **Figure 1D** separately for each fuel type, the results of which are shown in **Figure 3**.

Figure 3: Effects of varying fuel concentration on the TX-TL trade-off by fuel type. The same data are shown as in Figures 1C and 1D with points colored by fuel concentration. Each point is one of three replicates of a set of fuel and Mg^{2+} concentrations, where fuel is either **(A)** 3PGA, **(B)** maltose, **(C)** pyruvate, or **(D)** succinate. Experiments were performed in two batches of cell lysate, Batch 1 and Batch 2, in systems using 5 nM DNA.

While systems fueled by pyruvate, succinate, and maltose did not display any trend along the trade-off curve with increasing fuel concentration (**Figure 3B-D**), we found that in 3PGA-fueled systems, increasing fuel concentration generally shifted a TX-TL system from a transcription-optimized regime to a translation-optimized regime (**Figure 3A**). This phenomenon was also observed in 3PGA-fueled TX-TL systems where the volume fraction of cell lysate in a TX-TL reaction was increased or decreased by 25% (**Figure S3**), with the trend becoming stronger as the lysate fraction was increased (**Figure S3D**), extending the relevance of the trend to TX-TL systems using alternative cell lysate volume fractions[25].

We next sought to confirm whether the TX-TL trade-off and 3PGA trend extended to systems using alternative reporter plasmids. The results of these 3PGA versus Mg^{2+} panels are shown in **Figure 4**.

Figure 4: Effects of 3PGA concentration on the TX-TL trade-off across different reporter plasmids. Shown are trade-off curves corresponding to **(A)** P_{OR1OR2}-MG aptamer-deGFP, **(B)** P_{T7}-MG aptamer-deGFP, **(C)** P_{Tet}-MG aptamer-deGFP, and **(D)** P_{Tet}-F30 Pepper-mTurquoise2. Points are colored by 3PGA concentration. P_{OR1OR2}, a strong promoter and P_{Tet}, a medium strength promoter, are constitutive promoters, and P_{T7} is a strong non-*E. coli* promoter expressed in TX-TL reactions supplemented with 10µM T7 RNA polymerase (see Methods and Materials for sequences and additional information). Each point represents one of three replicates of a particular set of 3PGA and Mg^{2+} concentrations. The data shown in **(A)**, **(B)**, and **(C)** reflect experiments performed in cell lysate Batch 3, while the data in **(D)** reflect experiments performed in cell lysate Batch 2 (see Methods and Materials for details). All experiments were performed using 5 nM DNA.

We first switched the promoter from P_{OR1OR2} to P_{T7}, where we found that the TX-TL trade-off and 3PGA trend were present across P_{T7}-driven TX-TL systems (**Figure 4B**) just as they were in P_{OR1OR2}-driven systems (**Figure 4A**). When the promoter was changed to P_{Tet}, however, the TX-TL trade-off was no longer visible (**Figure 4C**). In P_{Tet}-driven TX-TL systems, it appeared that only some portion of the data, corresponding to about 15-30 mM 3PGA, fell along the common trade-off curve (**Figure S4**). The data corresponding to 0-5 mM 3PGA did not give the highest MG aptamer yields as in **Figure 4A** and **Figure 4B**, and at high 3PGA concentrations, the left edge of the curve dipped, reflecting decreased deGFP yields at high 3PGA concentrations. To determine if this was a behavior unique to this promoter, we also performed a 3PGA versus Mg^{2+} panel in systems expressing F30-Pepper aptamer and mTurquoise2 under a P_{Tet} promoter (**Figure 4D**). Here the trade-off curve and 3PGA trend were present as before, although as in systems using the P_{Tet}-MG aptamer-deGFP plasmid, there was a decrease in protein yields at higher 3PGA concentrations. While it is not clear what caused this change in the shape of the

trade-off curve in P_{Tet}-driven systems, this finding is consistent with the hypothesis that the TX-TL trade-off is indicative of a fundamental trade-off in fuel and energy metabolism. In such systems, weaker promoters would not be able to maximize rates of transcription and translation output as effectively, resulting in greater energy dissipation and translation-inhibiting waste accumulation compared to TX-TL under strong promoters, which could explain why deGFP yields tend to drop at high 3PGA concentrations in **Figure 4C** and **Figure 4D**, although this explanation is insufficient to explain the shape of the data in **Figure 4C**. While more promoters need to be tested to determine the generalizability of the TX-TL trade-off to systems using different genetic parts, the presence of the trade-off in P_{T7}-driven MG aptamer and deGFP expression was notable, considering that the T7 promoter is widely used in applications aimed at maximizing protein expression.

In 3PGA-fueled systems, the TX-TL trade-off and fuel trend are consistent across DNA concentrations

Having confirmed that the TX-TL trade-off was present in a wide variety of TX-TL systems, we next sought to confirm whether the trade-off curve scaled with DNA concentration, as previous experiments had all been performed with 5 nM DNA. For this set of experiments, we decided to focus on 3PGA-fueled systems, since 3PGA – and its downstream metabolic product phosphoenolpyruvate (PEP) – are the most widely used metabolites for energy regeneration in TX-TL systems, and gaining insight into the TX-TL trade-off in 3PGA-fueled systems would have greater relevance for studies aimed at characterizing and improving these systems. We performed 3PGA versus Mg^{2+} panels at DNA concentrations of 1, 2.5, 5, 7.5, and 10nM, the results of which are shown in **Figure 5**. We found that the TX-TL trade-off scaled as DNA concentration was increased, with a curve first becoming apparent at 2.5 nM DNA and then becoming more apparent as the DNA concentration was increased (**Figure 5A**).

Figure 5: Effects of varying DNA concentration on the TX-TL trade-off. **(A)** Each point represents one of three replicates of a particular set of 3PGA and Mg^{2+} concentrations, where points are colored by DNA concentrations ranging from 1-10 nM. **(B)** The same data are shown as in (A), albeit with the points colored by 3PGA concentration rather than DNA concentration. The data shown reflect experiments performed in cell lysate Batch 2 (see Methods and Materials for details).

As in the 5 nM DNA cases discussed in the previous section, the trend in shifting a system from a transcription-optimized regime to a translation-optimized regime at higher 3PGA

concentrations was also consistent in the trade-off curves at other DNA concentrations (**Figure 5B**). Preliminary analyses also revealed that maximum transcription along a given trade-off curve scaled linearly with increasing DNA concentration in systems with high Mg^{2+} concentration and 0mM 3PGA (**Figure S5**), although the fact that this occurs only in regimes where translation is poor suggests that behavior may exist only in TX-only systems. More experiments are needed to determine whether the linear scaling of maximum transcription with DNA is a general phenomenon in TX-TL systems.

The trade-off arises at the translational level in 3PGA-fueled systems

Having confirmed that there was a universal TX-TL trade-off that controlled a system's transcriptional and translational capacity as a function of Mg^{2+} concentration broadly, and additionally as a function of 3PGA concentration when that fuel source was used, we next sought to gain more insight into the cause of the trade-off in 3PGA-fueled systems. It was not clear from our previous data whether conditions that optimized transcription versus conditions that optimized translation were simply mutually exclusive, in that optimal Mg^{2+} and 3PGA concentrations were simply different for transcription versus translation. An alternative explanation was that either or both of the processes, transcription and translation, had an inhibitory effect on the other process; in the case where only one process inhibited the other, the first process could be inhibited by the reaction environment. A third possible explanation was that transcription and translation were processes competing for the same resources but that one or both processes had a preferred regime for optimization. Combinations of these explanations, or others, could also explain the observed trade-off.

To gain insight into the individual processes of transcription and translation, we decoupled the TX-TL system into a TX-only system and a TL-only system. In the TX-only system, tetracycline was added to the system to block translation via inhibition of translation initiation through its binding to the 30S subunit of the ribosome[38]. In the TL-only system, mRNA was added to the system instead of DNA. The results of these experiments are shown in **Figure 6**.

Using heatmaps to display integrated mRNA and maximum deGFP data, we found that in the TX-TL system (**Figure 6A**), the regime of Mg^{2+} and 3PGA concentrations that optimized transcription was distinct and different than the regime that optimized translation (**Figure 6D**). Specifically, optimal transcription preferred low 3PGA and high Mg^{2+} concentrations and optimal translation preferred high 3PGA and low Mg^{2+} concentrations; both findings are consistent with the trade-off curve. However, in the TX-only system (**Figure 6B**), MGaptamer expression was not optimized in any particular regime of Mg^{2+} or 3PGA concentrations (**Figure 6E**), although overall expression was higher, likely due to the abundance of energy in the absence of translation. Meanwhile, in the TL-only system (**Figure 6C**), higher deGFP yields were achieved, as before, at low Mg^{2+} and high 3PGA concentrations, despite reduced mRNA degradation in the regime of high Mg^{2+} and low 3PGA concentrations (**Figure 6F**).

Figure 6: Decoupling of transcription (TX) and translation (TL). **(A)**, **(B)**, and **(C)** show experimental schemes corresponding to the data shown in **(D)**, **(E)**, and **(F)**, respectively. **(A)** Simplified view of TX-TL: P_{OR1OR2}-MG aptamer-deGFP DNA is translated by *E. coli* RNA polymerase to mRNA, which fluoresces upon binding to Malachite Green (MG) dye. The mRNA is then translated by ribosomes into deGFP protein. **(B)** Simplified view of TX-only system: transcription proceeds as in (A), but translation is inhibited by tetracycline, which binds to ribosomes and inhibits translation initiation. **(C)** Simplified view of TL-only system: MG aptamer-deGFP mRNA, rather than DNA, is added to the system, and subsequently the mRNA is translated into deGFP protein. **(D)** Integrated mRNA measurements (top) and endpoint deGFP measurements (bottom) for the TX-TL system. **(E)** Integrated mRNA measurements (top) and endpoint deGFP measurements (bottom) for the TX-only system, to which 200 µg/mL tetracycline was added. **(F)** Integrated mRNA measurements (top) and endpoint deGFP measurements (bottom) for the TL-only system, where 0.3 µM mRNA was added at the beginning of the reaction, respectively. All reactions shown use Batch 2 lysate, and the TX-TL and TX-only systems use 5 nM DNA. For reference, standard TX-TL formulations for this lysate batch use 30 mM 3PGA and 8 mM Mg^{2+}. Each measurement is the average of three replicates for each experimental condition. Colorbars are scaled separately for each plot.

These data suggest that the TX-TL trade-off is caused by a combination of translation optimization at low Mg^{2+} and high 3PGA concentrations, increased MGaptamer stability at high Mg^{2+} and low 3PGA concentrations, and competition for resources in the TX-TL system. In other

words, optimal translation occurs where conditions are ideal for that process, and optimal transcription occurs where translation is sub-optimal and leaves more fuel and energy resources for transcription. While these experiments did not shed light on the mechanism governing translation optimization at lower Mg^{2+} and higher 3PGA concentrations, they suggest that the TX-TL trade-off arises at the translational level in 3PGA fueled systems.

The TX-TL trade-off is absent when TX-TL systems are supplied with no fuel or Mg^{2+} and additional energy

The TX-TL decoupling experiments had implied that the TX-TL trade-off arose, in part, due to competition for fuel and energy resources between transcription and translation. By this hypothesis, TX-TL systems with abundant energy should not fall along the trade-off curve, particularly in a regime of low Mg^{2+} concentration where both transcription and translation were favorable when not limited by a competing process (**Figures 6E, F**). To test this hypothesis, we next performed experiments where TX-TL systems were supplied with no central carbon fuel source and no added Mg^{2+}. To explore the space of different energy sources and concentrations, we performed a three-dimensional panel of ATP versus GTP versus NTPs, where each solution was added at a final concentration of 0, 5, 10, or 15mM (see Methods and Materials for more details regarding this experiment). The results of these experiments are shown in **Figure 7**.

Figure 7: Absent TX-TL trade-off in the absence of fuel or Mg^{2+}. **(A)** Overview of experimental workflow. **(B)** Maximum deGFP values versus integrated MG aptamer values (on a log scale for this axis). Each point represents one of three replicates of a particular set of NTP, added ATP, and added GTP concentrations, colored by the fuel type. Here, the "central carbon" fuel data is the same as in Figure 1, and those reactions all contain 4.8 mM NTPs. The "NTPs" fuel data reflect experiments where no central fuel or Mg^{2+} has been added. The data shown reflect experiments performed in two batches of cell lysate, Batch 1 and Batch 2 (see Methods and Materials for details), and all experiments were performed using 5 nM DNA.

As expected, TX-TL systems supplied with additional energy molecules and no fuel or no Mg^{2+} did not fall along the previously found trade-off curve corresponding to that DNA concentration (5 nM). These systems could achieve deGFP yields comparable to systems supplied with pyruvate or maltose and integrated MG aptamer yields 3-10X higher than in TX-TL systems supplied with those fuels and Mg^{2+}. While there was no clear trend in ATP, GTP, or total NTP

concentration for the data shown in **Figure 7B**, generally, as the concentration of NTPs increased, integrated MG aptamer values increased, and deGFP values decreased (**Figure S6**), which is consistent with a previous finding[28]. The decrease in deGFP yields despite the abundance of energy was likely due to NTP chelation of Mg^{2+}, which is necessary for ribosomes and other translation machinery. Although Mg^{2+} was not directly added to the reactions, it was present at a concentration of around 4mM due to the presence of S30B buffer (used in the preparation of cell lysate) in the reaction. Our results suggest that fuel-supplied TX-TL systems are indeed energy-limited, and that the TX-TL trade-off curve exists in part due to competition for a limited pool of energy for transcription and translation.

The data revealed two additional insights. First, systems with no fuel or no Mg^{2+} and high NTPs often exhibited unusual transcriptional and translational dynamics (**Figure S7**). Some systems exhibit long time delays in transcription (**Figure S7A**) or translation (**Figures S7E-F**). Other systems exhibited biphasic expression (**Figures S7B, D**) or steady state transcription rates as opposed to the typical transcription pulse (**Figures S7C**). Time delays in particular could be a desired property for TX-TL applications such as environmental event recording or genelet circuits with a large transcriptional circuit and few protein outputs.

Second, although these TX-TL systems did not achieve protein yields as high as those achieved in systems with 3PGA, supplying systems with an additional 5 NTPs can help achieve deGFP yields comparable to systems with about 5 mM NTPs and 10 mM 3PGA. This suggests that while energy regeneration is likely occurring in 3PGA-fueled systems, energy being regenerated is not being used efficiently. A large fraction of that regenerated energy, perhaps the majority, is likely fueling metabolic processes irrelevant to transcription or translation.

Discussion:

An improved understanding of cell-free TX-TL systems is essential for scaling up, expanding the scope of, and improving the performance of TX-TL applications in academic and industrial settings. By using metabolic perturbations as a tool to gain insight into the behavior of TX-TL systems in different metabolic regimes, this study has shed light on how Mg^{2+} and fuel contribute to TX-TL variability and how these two components and NTPs affect TX-TL lifetime. Our results suggest that variability ultimately stems from the cell lysate, specifically the varying abilities of different batches of cell lysate to regulate translation and perform energy regeneration, and that this variability can be reduced by modulating these components.

While it is not immediately clear why different batches of cell lysate have different translation regulation and energy regeneration capabilities, several possible hypotheses exist. It is possible that different methods of cell lysis (or even batch-to-batch differences in cell lysis) result in different proteomic compositions of cell lysate, perhaps because some proteins might be more robust to denaturation during sonication compared to other proteins. Another possibility is that, despite being grown to *approximately* the same optical density (OD), small differences in OD might result in cells being in vastly growth phases, which would also result in different proteomic compositions for each batch of lysate. As Mg^{2+} is an essential cofactor for many different

proteins in cell lysate, and as many different proteins in a cell lysate play a role in energy regeneration, it is hard to know which proteins are directly responsible for variability and the TX-TL trade-off, but it is likely dozens or even hundreds of different proteins.

For users of TX-TL systems, this study provides a new method of analyzing the effects of variability, via the analysis of plots of maximum deGFP versus integrated MG aptamer. The analysis is a novel approach to understanding batch-to-batch variability across cell lysates and perhaps one that may be useful in data calibration and standardization. In our study, perhaps the most successful application of this approach has been elucidating how Mg^{2+} affects TX-TL variability via controlling the extent to which a system is optimized for transcription versus translation. This approach was similarly valuable in gaining many new insights into TX-TL variability: the effects of different fuels, notably 3PGA; the performances of different lysates; and how TX-TL processes scale with DNA concentration.

Armed with these insights, other users of TX-TL systems aiming to reduce variability and standardize their systems may consider calibrating the concentrations of the relevant small molecules (fuel, Mg^{2+}, and/or DNA) for their application so that TX-TL curves for different systems have similar locations, shapes, and other characteristics. By characterizing different batches of cell lysate by the shape and location of their trade-off curves, for example, we can understand their respective transcriptional and translational limits. We can also calibrate transcription versus translation optimization in different batches of cell lysate by choosing the appropriate set of fuel types, fuel concentrations, and Mg^{2+} concentrations for a particular genetic program. For example, users of transcriptional circuits, where circuit computation occurs mostly or entirely at the transcriptional level, may be able to achieve better circuit performance in TX-TL systems by simply increasing the Mg^{2+} concentration by a few millimolar and optionally switching to a maltose-fueled system for better performance. Conversely, users desiring maximum protein yield, a common goal for TX-TL applications, may be able to achieve better performance by decreasing the amount of added Mg^{2+} or omitting it altogether, with further improvements coming from optimizing the fuel source and concentration.

Beyond reducing variability and improving standardization, this study also has relevance for efforts aimed at increasing the predictability and performance of TX-TL systems for desired uses. While the analyses performed in this study have focused mostly on integrated MG aptamer and maximum deGFP measurements, it is worth noting that many systems with similar values exhibited very different dynamics. As discussed in the Results section, systems provided with additional NTPs exhibited unique and potentially desirable dynamics, but even systems provided with central carbon fuels have revealed potentially relevant findings. For example, although TX-TL systems have been shown to be powered by a wide variety of fuel sources[22,23,39], 3PGA and PEP have been the fuels of choice due their energy regeneration capability. However, our data suggest that, at the right fuel and Mg^{2+} concentrations, succinate and pyruvate perform very similarly to 3PGA, particularly in achieving high deGFP yields. As pyruvate and succinate cost about 25X and 4000X cheaper than 3PGA, respectively, this finding may be relevant for further reducing the cost of TX-TL systems. The data included with this paper provide hundreds of unique reaction conditions that can serve as a starting point for

exploring conditions that are optimized for high protein yields, long reaction lifetime, or other desired behaviors.

In addition to providing insight into design considerations for TX-TL systems, the TX-TL trade-off curve, initially a surprising finding, has also provided a simple yet powerful way of gaining insight into fundamental trade-offs in fuel and energy metabolism. Taken together, our results suggest that the TX-TL trade-off is present in energy-limited systems and is indicative of a trade-off between energy regeneration and waste accumulation that results from fuel metabolism. In other words, while fuel molecules enable energy regeneration, their own metabolism also results in waste accumulation (through the generation of immediate or downstream products) that affects TX-TL processes. While the identities of the waste molecules remain unknown, our results suggest that in 3PGA-fueled systems, the waste is partially 3PGA-derived and affects translation. This insight, along with results from the TX-TL decoupling experiments, have helped us to create a new model of cell-free metabolism coupled to cell-free TX-TL (**Figure 8**). While this model has not been validated computationally, doing so may enable improved prediction in TX-TL systems, perhaps even in non-lysate-based TX-TL systems (e.g. PURE-based systems[16]), and this remains an active area of future work.

Figure 8: A new model for cell-free metabolism coupled to cell-free TX-TL. The conversion of fuel to waste regenerates energy, which can power transcription and translation, but the accumulation of waste inhibits translation. The fuel-energy-waste paradigm is derived from an existing model[33]. This model's key contribution is providing roles for Mg^{2+} via its inhibition of mRNA degradation and translation.

In revealing the fundamental metabolic trade-offs in TX-TL systems, this work has also shed light on why previous studies aimed at increasing protein yield have been largely unsuccessful. Our data suggest that low Mg^{2+} concentrations, strong promoters, and high DNA concentrations – and high 3PGA concentrations where 3PGA is used – are necessary for high deGFP yields. However, it is not possible to remove more Mg^{2+} from the system without compromising the function of proteins in the cell lysate, nor is adding more fuel possible, as the inability of TX-TL systems to take advantage of faster energy regeneration will make the faster waste

accumulation an issue at high fuel concentrations. Adding too much DNA may skew the systems towards transcription rather than translation. Previous studies aimed at improving deGFP yields have focused on other mechanisms of energy regeneration and waste mitigation, such as using combinations of fuels like 3PGA and maltose[34], and such approaches and others, including at the level of plasmid design, will be necessary for vastly improving protein yields in lysate-based TX-TL systems.

Overall, by elucidating the effects of cell-free metabolism on TX-TL systems, this work lays a foundation for improving TX-TL performance, lifetime, standardization, and prediction. Ultimately, it represents one of many such steps that will be necessary for further improvement of TX-TL systems, whether that be for applications in biomanufacturing, gaining insight into natural biological systems, or the development of synthetic life.

Materials and Methods:

Preparation of cell lysate

Cell lysate was prepared from *E. coli* BL21 Rosetta2 cells using the protocol established by Sun and coworkers[14] with some modifications. Briefly, cells were grown sequentially in 6 mL, 100 mL, and 1 L cultures, where smaller cultures were used to inoculate larger ones. Once 6L of culture had grown to roughly an OD600 of 2, cells were repeatedly centrifuged and washed with S30B buffer by re-suspension. Cells were then re-suspended in S30B buffer and subjected to cell lysis.

For the Batch 2 and Batch 3 lysates, cells were resuspended at 1.4 g cells per mL of S30B buffer and lysed by a French cell press at a pressure of 640 psi. For the Batch 1 lysate, cells were resuspended to 1 g cells per mL of S30B buffer and lysed by sonication at a probe frequency of 20 kHz and an amplitude of 50%. Cells were sonicated at 10-second ON/OFF intervals for a total of 60 seconds ON (6 ON/OFF cycles), resulting in approximately 300 Joules of energy delivered. After both methods of cell lysis, 3 µL of 1M DTT was added per mL of cell lysate, and the lysates were then centrifuged to remove cell debris. All batches of lysate were then subjected to a 1-hour run-off reaction at 37°C, after which they were subjected to centrifugation to remove additional debris.

The Batch 2 and Batch 3 lysates were additionally subjected to dialysis at 4°C via Slide-a-Lyzer G3 cassettes (10kDa, 15mL) placed in a beaker containing S30B buffer and a magnetic stir bar to facilitate diffusion. Excess air was removed using a syringe. The Batch 2 lysate was dialyzed for 4 hours (where cassettes were transferred to a beaker of fresh S30B at the 2-hour mark) at 140 rpm. The Batch 3 lysate was dialyzed for 2 hours (where cassettes were transferred to a beaker of fresh S30B at the 35-minute mark) at 160 rpm.

Finally, both batches of cell lysate were aliquoted, flash-frozen in liquid nitrogen, and stored at -80°C until further use. Bradford assays determined that the final protein concentrations for the cell lysates were 30-40 mg/mL.

Cloning plasmids used to express deGFP

The P_{OR1OR2}-MG aptamer-deGFP plasmid used to express deGFP was originally prepared by Dan Siegal and modified by Zoila Jurado via Site-Directed Mutagenesis to change the T7 promoter to a P_{OR1OR2} promoter and the T7 terminator to a T500 terminator. This plasmid is P_{OR1OR2}-MG aptamer-UTR1-deGFP-T500. The P_{Tet}-MG aptamer-UTR1-deGFP-T500 plasmid was subsequently constructed using Site-Directed Mutagenesis to change the P_{OR1OR2} promoter to a P_{Tet} promoter, to make expression results as comparable as possible. The P_{T7}-MG aptamer-UTR1-deGFP-T500 plasmid was originally prepared by Dan Siegal-Gaskins and modified using Site-Directed Mutagenesis to add a few extra bases to the T7 promoter. The P_{Tet}-F30Pepper-UTR1-mTurquoise2-ECK120029600 plasmid was constructed using Golden Gate Assembly of

CIDAR MoClo part plasmids. All plasmid sequences are in the Supplementary Information section.

Preparation of template DNA plasmids

In experiments where the P_{OR1OR2}-MG aptamer-UTR1-deGFP-T500 plasmid was used, template DNA for the cell-free reactions was prepared from *E. coli* KL740 cells (purchased from Arbor Biosciences) that had been transformed with the plasmid described above. Cells were grown overnight at 30°C, in Lysogeny Broth (LB) medium supplemented with 100µg/mL carbenicillin, and either Mini-prepped (using a Qiagen Mini-prep kit) or Midi-prepped (using a Macharey-Nagel NucleoBond Xtra Midi kit) the next day using the appropriate protocols. When Mini-prepping, DNA was eluted with nuclease-free water; when Midi-prepping, after the concentration step, DNA was eluted with IDTE Buffer (10 mM Tris-HCl, 0.1 mM EDTA).

The P_{Tet}-MGaptamer-UTR1-deGFP-T500, P_{T7}-MG aptamer-UTR1-deGFP-T500, and P_{Tet}-F30Pepper-UTR1-mTurquoise2-ECK120029600 plasmids were prepared from cells grown overnight at 37°C, in Lysogeny Broth (LB) medium supplemented with 100 µg/mL carbenicillin (for the deGFP plasmids) or 50 µg/mL kanamycin. The cells were Mini-prepped using a Qiagen Mini-prep kit, where DNA was eluted with nuclease-free water.

Preparation of cell-free reactions

Cell-free reactions were prepared as per the protocol by Sun and coworkers[14]. Unless otherwise noted, each reaction consisted of the following: 33% (by volume) of cell lysate, 1.5 mM of each amino acid (except for Leucine, which was added at 1.25 mM), 4.8 nM of NTP mix (containing 1.5 mM each of ATP and GTP, 0.9 mM each of CTP and UTP, pH adjusted to 7.5 using KOH), 50mM HEPES pH 8, 0.2 mg/mL tRNA, 0.26 mM Coenzyme A, 0.33 mM NAD$^+$, 0.75 mM cyclic AMP (cAMP), 0.068 mM Folinic acid, 1 mM Spermidine, 10 µM Malachite Green Dye, and 30 mM 3PGA. Depending on the batch of cell lysate, a standard reaction used either 8 mM magnesium glutamate and 80 mM potassium glutamate (for the Batch 2 lysate) or 6 mM magnesium glutamate and 140 mM potassium glutamate (for the Batch 1 lysate). For the Mg^{2+} versus Fuel panels, 3PGA and Mg^{2+} were omitted from the master mix, and the appropriate concentration of Mg^{2+} and 3PGA was added to reactions individually. Finally, depending on the experiments, a variable amount of DNA plasmid was added, and the remaining volume of the 10 µL reaction was filled with nuclease-free water.

As reactions were all performed in 384-well glass-bottom plates, water, DNA, and components were titrated into the TX-TL reaction using the Echo 525 liquid handler, while the remaining components were added to a bulk solution and electronically dispensed by a multi-dispense pipette. Unless otherwise specified, all reactions containing DNA were supplied with the P_{OR1OR2}-MG aptamer-deGFP plasmid. To prevent the reaction from starting before all the components had been added in, all reagents were kept on ice until they were added to the reaction and added sequentially to reduce evaporation and/or degradation. First, a master mix minus the cell lysate was created and kept on ice in an Eppendorf tube; then, components to be added by

Echo were added to the 384-well plate, where all stock concentrations were diluted to less than 500mM so that they could be effectively dispensed via Echo; next, the cell lysate was added to the master mix; finally, the master mix was pipetted to the wall of each well. Once the plate was covered with a plastic seal, it was centrifuged at 4000xg for 1 minute at 4°C. Finally, the bottom of the plate was quickly cleaned by Kimwipe to remove any debris and immediately transferred to a plate BioTek H1MF plate reader that had reached its setpoint of 29°C.

Preparation of no-fuel, no Mg²⁺ cell-free reactions

Reactions were prepared as described in the "Preparation of cell-free reactions" with minor changes. No fuel or additional Mg^{2+} were added to the reactions, apart from what was already there from the preparation of the cell lysate. For the addition of the NTP mix, we prepared a NTP stock solution at a molar ratio of 1.5:1.5:0.9:0.9 for ATP: GTP: CTP: UTP, pH adjusted to 7.5 using KOH. For the ATP and GTP solutions, we prepared stock solutions in nuclease-free water. Each solution was added at 0, 5, 10, or 15 mM.

Dynamic measurements of fluorescent molecules

Fluorescence measurements were performed by a BioTek H1MF plate reader at 29°C, and measurements were read from the bottom of the plate every 3 minutes (preceding by 5 seconds of linear shaking) at excitation/emission wavelengths suitable for MG aptamer (610/650 nm, Gain 150) and deGFP (485/515 nm, Gain 61) for 18 hours. For the experiment corresponding to Figure 4, excitation/emission wavelengths suitable for F30-Pepper (580/620 nm, Gain 150) and mTurquoise2 (434/474 nm, Gain 61) were used.

For MG aptamer and deGFP, arbitrary fluorescence units were converted to micromolar concentrations using a calibration curve prepared using purified deGFP protein and MG aptamer mRNA. To create a deGFP fluorescence calibration curve, serial dilutions of eGFP (purchased from Cell Biolabs) were performed in 1x PBS, after which 3 sample replicates of 10µL each were loaded onto a 384-well plate and read by a BioTek H1MF plate reader at 29°C and at a Ex/Em wavelengths of 485/515 and a Gain of 61. 3 technical replicates were read over 3 minutes at 1-minute intervals to generate 4 points per replicate, and the average of these 12 points was used as a single point at a given concentration through which the calibration line was fit with a zero intercept.

To create a MGaptamer fluorescence calibration curve, mRNA of the MGaptamer sequence only – specifically rArCrUrGrGrArUrCrCrCrGrArCrUrGrGrCrGrArGrArGrCrCrArGrGrUrArArCrG rArArUrGrGrArUrCrCrArArU – was purchased from IDT DNA. The lyophilized mRNA was re-suspended in nuclease-free water at its concentration was determined by a Nanodrop 2000c. Next, serial dilutions of MGaptamer mRNA were performed in 1X PBS containing 10 µM Malachite Green dye, after which 3-4 sample replicates of 10 µL each were loaded onto a 384-well plate and read by a BioTek H1MF plate reader at 29°C and at excitation/emission wavelengths of 610/650 and a Gain of 150. Four technical replicates were read over 10 minutes at 2.5-minute intervals to generate 5 points per replicate, and the average of these 20 points

was used as a single point at a given concentration through which the calibration line was fit with a zero intercept. The slope of the background- subtracted calibration curve was used to calibrate measurements.

mRNA purification for a translation-only system

To obtain an mRNA sequence containing MGaptamer-UTR1-deGFP-T500, a linear DNA template was first prepared, followed by DNA isolation, *in vitro* transcription, DNAse treatment, and mRNA isolation. First, for generation of the linear DNA template, primers of the following sequences were first used to amplify the relevant sequence off the P_{OR1OR2}-MG aptamer-deGFP plasmid by PCR: CCAGAAAACCGAATTTTGCTGG and ATGATAAAGAAGACAGTCATAAGTGCG. A 1-mL PCR reaction was performed using NEB Q5 2x Master Mix and run for 30 cycles with an elongation time of 45 seconds and an annealing temperature of 65°C.

After the PCR reaction was performed, the PCR was aliquoted in three 1.7-mL Eppendorf tubes (i.e., 333 µL per tube) and DNA was precipitated by adding 33 µL of 3 M sodium acetate and 1 mL 100% ethanol per tube. After chilling the tubes at -80°C for 20 minutes, the tubes were centrifuged at 16000x*g* for 30 minutes at 4°C. Finally, all supernatant was pipetted out, and residual ethanol was allowed to evaporate by leaving the tubes open and uncovered at room temperature. The DNA was resuspended in 50 µL nuclease-free water per tube.

In vitro transcription was performed using the HiScribe® T7 High Yield RNA Synthesis Kit, and instructions included in the kit's manual were used for DNase I treatment of the reaction.

Finally, the synthesized mRNA was isolated using the PureLink RNA Mini kit. 100% ethanol was first added to the reaction to a final concentration of 35%, after which the kit's "Protocol for RNA clean-up and purification from liquid samples" was used (omitting the Lysis step). mRNA concentration was determined by a Nanodrop 2000c.

Translation inhibition using tetracycline

The translation inhibition experiments were performed in Batch 2 lysate by adding 200 µg/mL tetracycline to TX-TL reactions. Stock tetracycline solutions were created at 80 mg/mL in DMSO, where they were stored at -80°C until further use. For addition to TX-TL reactions, the stock solution was diluted to 5 mg/mL in nuclease-free water and subsequently dispensed using the Echo 525 liquid handler. To determine the optimal concentration of tetracycline to use for translation-inhibition, titrations were performed (see **Figure S8**).

Data analysis

Fluorescence measurements of deGFP and MGaptamer were converted to micromolar concentrations via the calibration curves. Endpoint deGFP concentration was determined to be the maximum deGFP concentration achieved for a given experimental condition. Integrated

MGaptamer measurements were made by using the NumPy[40] **trapz** package for numerical integration using the trapezoidal method. The Jupyter notebooks that were used for creating tidy data and for subsequent data analysis and plot generation are included with this work, in addition to all the data that was collected for this study.

Acknowledgements:

The work described here would not be possible without the initial ideas that came out of extensive discussions with William Poole. His suggestions to probe the roles of Mg^{2+} and different fuel sources helped generate new questions and analyses that ultimately led to the experiments discussed here, and we thank him for his insights. Apart from Poole, additional members of the lab – specifically Ankita Roychoudhury, Zoila Jurado, David Alexander Johnson, and Miryong (Miki) Yun – also contributed to the work shown here. Zoila was present in and contributed to many of these early discussions with William, and she also prepared the P_{OR1OR2}-MG aptamer-deGFP plasmid used throughout this study. All four individuals helped prepare the cell lysate batches used in the experiments here, and we thank them for their help. The authors thank Paul Freemont for his helpful comments, particularly those pertaining to cell-free metabolism. Finally, ChatGPT was used to write part of the introduction section, which has been an interesting and helpful endeavor.

Research was sponsored by the Army Research Office and was accomplished under Cooperative Agreement Number W911NF-22-2-0210. The views and conclusions contained in this document are those of the authors and should not be interpreted as representing the official policies, either expressed or implied, of the Army Research Office or the U.S. Government. The U.S. Government is authorized to reproduce and distribute reprints for Government purposes notwithstanding any copyright notation herein.

Author Information:

Author Contributions

M.K. conceived and designed the study, performed experiments, acquired and analyzed the data, and wrote the manuscript. R.M.M. acquired funding and supervised this project.

Conflict of Interest

RMM has a financial stake in Tierra Biosciences, a private company that makes use of cell-free technologies such as those described in this article for protein expression and screening. The other authors have nothing to disclose.

References:

(1) Garenne, D.; Haines, M. C.; Romantseva, E. F.; Freemont, P.; Strychalski, E. A.; Noireaux, V. Cell-Free Gene Expression. *Nat. Rev. Methods Primer* **2021**, *1* (1), 49. https://doi.org/10.1038/s43586-021-00046-x.

(2) Brookwell, A.; Oza, J. P.; Caschera, F. Biotechnology Applications of Cell-Free Expression Systems. *Life* **2021**, *11* (12). https://doi.org/10.3390/life11121367.

(3) Karim, A. S.; Dudley, Q. M.; Juminaga, A.; Yuan, Y.; Crowe, S. A.; Heggestad, J. T.; Garg, S.; Abdalla, T.; Grubbe, W. S.; Rasor, B. J.; Coar, D. N.; Torculas, M.; Krein, M.; Liew, F. (Eric); Quattlebaum, A.; Jensen, R. O.; Stuart, J. A.; Simpson, S. D.; Köpke, M.; Jewett, M. C. In Vitro Prototyping and Rapid Optimization of Biosynthetic Enzymes for Cell Design. *Nat. Chem. Biol.* **2020**, *16* (8), 912–919. https://doi.org/10.1038/s41589-020-0559-0.

(4) Kelwick, R.; Webb, A. J.; MacDonald, J. T.; Freemont, P. S. Development of a Bacillus Subtilis Cell-Free Transcription-Translation System for Prototyping Regulatory Elements. *Metab. Eng.* **2016**, *38*, 370–381. https://doi.org/10.1016/j.ymben.2016.09.008.

(5) Takahashi, M. K.; Hayes, C. A.; Chappell, J.; Sun, Z. Z.; Murray, R. M.; Noireaux, V.; Lucks, J. B. Characterizing and Prototyping Genetic Networks with Cell-Free Transcription–Translation Reactions. *Bact. Archaeal Transcr.* **2015**, *86*, 60–72. https://doi.org/10.1016/j.ymeth.2015.05.020.

(6) Swank, Z.; Maerkl, S. J. CFPU: A Cell-Free Processing Unit for High-Throughput, Automated In Vitro Circuit Characterization in Steady-State Conditions. *BioDesign Res.* **2021**, *2021*, 2968181. https://doi.org/10.34133/2021/2968181.

(7) Vögeli, B.; Schulz, L.; Garg, S.; Tarasava, K.; Clomburg, J. M.; Lee, S. H.; Gonnot, A.; Moully, E. H.; Kimmel, B. R.; Tran, L.; Zeleznik, H.; Brown, S. D.; Simpson, S. D.; Mrksich, M.; Karim, A. S.; Gonzalez, R.; Köpke, M.; Jewett, M. C. Cell-Free Prototyping Enables Implementation of Optimized Reverse β-Oxidation Pathways in Heterotrophic and Autotrophic Bacteria. *Nat. Commun.* **2022**, *13* (1), 3058. https://doi.org/10.1038/s41467-022-30571-6.

(8) Sharpes, C. E.; McManus, J. B.; Blum, S. M.; Mgboji, G. E.; Lux, M. W. Assessment of Colorimetric Reporter Enzymes in the PURE System. *ACS Synth. Biol.* **2021**, *10* (11), 3205–3208. https://doi.org/10.1021/acssynbio.1c00360.

(9) Zhang, Y.; Steppe, P. L.; Kazman, M. W.; Styczynski, M. P. Point-of-Care Analyte Quantification and Digital Readout via Lysate-Based Cell-Free Biosensors Interfaced with Personal Glucose Monitors. *ACS Synth. Biol.* **2021**, *10* (11), 2862–2869. https://doi.org/10.1021/acssynbio.1c00282.

(10) Nguyen, P. Q.; Soenksen, L. R.; Donghia, N. M.; Angenent-Mari, N. M.; de Puig, H.; Huang, A.; Lee, R.; Slomovic, S.; Galbersanini, T.; Lansberry, G.; Sallum, H. M.; Zhao, E. M.; Niemi, J. B.; Collins, J. J. Wearable Materials with Embedded Synthetic Biology Sensors for Biomolecule Detection. *Nat. Biotechnol.* **2021**, *39* (11), 1366–1374. https://doi.org/10.1038/s41587-021-00950-3.

(11) Cho, E.; Lu, Y. Compartmentalizing Cell-Free Systems: Toward Creating Life-Like Artificial Cells and Beyond. *ACS Synth. Biol.* **2020**, *9* (11), 2881–2901. https://doi.org/10.1021/acssynbio.0c00433.

(12) Yewdall, N. A.; Mason, A. F.; van Hest, J. C. M. The Hallmarks of Living Systems: Towards Creating Artificial Cells. *Interface Focus* **2018**, *8* (5), 20180023. https://doi.org/10.1098/rsfs.2018.0023.

(13) Rothschild, L. J.; Averesch, N. J. H.; Strychalski, E. A.; Moser, F.; Glass, J. I.; Cruz Perez, R.; Yekinni, I. O.; Rothschild-Mancinelli, B.; Roberts Kingman, G. A.; Wu, F.; Waeterschoot, J.; Ioannou, I. A.; Jewett, M. C.; Liu, A. P.; Noireaux, V.; Sorenson, C.; Adamala, K. P. Building Synthetic Cells—From the Technology Infrastructure to Cellular Entities. *ACS*

Synth. Biol. **2024**. https://doi.org/10.1021/acssynbio.3c00724.

(14) Sun, Z. Z.; Hayes, C. A.; Shin, J.; Caschera, F.; Murray, R. M.; Noireaux, V. Protocols for Implementing an Escherichia Coli Based TX-TL Cell-Free Expression System for Synthetic Biology. *J. Vis. Exp.* **2013**, No. 79, e50762. https://doi.org/10.3791/50762.

(15) Gregorio, N. E.; Levine, M. Z.; Oza, J. P. A User's Guide to Cell-Free Protein Synthesis. *Methods Protoc.* **2019**, *2* (1). https://doi.org/10.3390/mps2010024.

(16) Shimizu, Y.; Inoue, A.; Tomari, Y.; Suzuki, T.; Yokogawa, T.; Nishikawa, K.; Ueda, T. Cell-Free Translation Reconstituted with Purified Components. *Nat. Biotechnol.* **2001**, *19* (8), 751–755. https://doi.org/10.1038/90802.

(17) Cole, S. D.; Beabout, K.; Turner, K. B.; Smith, Z. K.; Funk, V. L.; Harbaugh, S. V.; Liem, A. T.; Roth, P. A.; Geier, B. A.; Emanuel, P. A.; Walper, S. A.; Chávez, J. L.; Lux, M. W. Quantification of Interlaboratory Cell-Free Protein Synthesis Variability. *ACS Synth. Biol.* **2019**, *8* (9), 2080–2091. https://doi.org/10.1021/acssynbio.9b00178.

(18) Rhea, K. A.; McDonald, N. D.; Cole, S. D.; Noireaux, V.; Lux, M. W.; Buckley, P. E. Variability in Cell-Free Expression Reactions Can Impact Qualitative Genetic Circuit Characterization. *Synth. Biol.* **2022**, *7* (1), ysac011. https://doi.org/10.1093/synbio/ysac011.

(19) Roychoudhury, A. Understanding the Lifetime and Rate of Protein Production in Cell-Free Reactions While Maximizing Energy Use. Undergraduate Thesis, California Institute of Technology, 2021.

(20) Jiang, N.; Ding, X.; Lu, Y. Development of a Robust Escherichia Coli-Based Cell-Free Protein Synthesis Application Platform. *Biochem. Eng. J.* **2021**, *165*, 107830. https://doi.org/10.1016/j.bej.2020.107830.

(21) Kim, T.-W.; Kim, D.-M.; Choi, C.-Y. Rapid Production of Milligram Quantities of Proteins in a Batch Cell-Free Protein Synthesis System. *J. Biotechnol.* **2006**, *124* (2), 373–380. https://doi.org/10.1016/j.jbiotec.2005.12.030.

(22) Karim, A. S.; Heggestad, J. T.; Crowe, S. A.; Jewett, M. C. Controlling Cell-Free Metabolism through Physiochemical Perturbations. *Metab. Eng.* **2018**, *45*, 86–94. https://doi.org/10.1016/j.ymben.2017.11.005.

(23) Wang, Y.; Zhang, Y.-H. P. Cell-Free Protein Synthesis Energized by Slowly-Metabolized Maltodextrin. *BMC Biotechnol.* **2009**, *9* (1), 58. https://doi.org/10.1186/1472-6750-9-58.

(24) Vezeau, G. E.; Salis, H. M. Tuning Cell-Free Composition Controls the Time Delay, Dynamics, and Productivity of TX-TL Expression. *ACS Synth. Biol.* **2021**, *10* (10), 2508–2519. https://doi.org/10.1021/acssynbio.1c00136.

(25) Banks, A. M.; Whitfield, C. J.; Brown, S. R.; Fulton, D. A.; Goodchild, S. A.; Grant, C.; Love, J.; Lendrem, D. W.; Fieldsend, J. E.; Howard, T. P. Key Reaction Components Affect the Kinetics and Performance Robustness of Cell-Free Protein Synthesis Reactions. *Comput. Struct. Biotechnol. J.* **2022**, *20*, 218–229. https://doi.org/10.1016/j.csbj.2021.12.013.

(26) Batista, A. C.; Soudier, P.; Kushwaha, M.; Faulon, J.-L. Optimising Protein Synthesis in Cell-Free Systems, a Review. *Eng. Biol.* **2021**, *5* (1), 10–19. https://doi.org/10.1049/enb2.12004.

(27) Borkowski, O.; Koch, M.; Zettor, A.; Pandi, A.; Batista, A. C.; Soudier, P.; Faulon, J.-L. Large Scale Active-Learning-Guided Exploration for in Vitro Protein Production Optimization. *Nat. Commun.* **2020**, *11* (1), 1872. https://doi.org/10.1038/s41467-020-15798-5.

(28) Nagaraj, V. H.; Greene, J. M.; Sengupta, A. M.; Sontag, E. D. Translation Inhibition and Resource Balance in the TX-TL Cell-Free Gene Expression System. *Synth. Biol.* **2017**, *2* (1), ysx005. https://doi.org/10.1093/synbio/ysx005.

(29) Caschera, F.; Karim, A. S.; Gazzola, G.; d'Aquino, A. E.; Packard, N. H.; Jewett, M. C. High-Throughput Optimization Cycle of a Cell-Free Ribosome Assembly and Protein Synthesis System. *ACS Synth. Biol.* **2018**, *7* (12), 2841–2853. https://doi.org/10.1021/acssynbio.8b00276.

(30) Horvath, N.; Vilkhovoy, M.; Wayman, J. A.; Calhoun, K.; Swartz, J.; Varner, J. D. Toward a Genome Scale Sequence Specific Dynamic Model of Cell-Free Protein Synthesis in Escherichia Coli. *Metab. Eng. Commun.* **2020**, *10*, e00113. https://doi.org/10.1016/j.mec.2019.e00113.

(31) Vilkhovoy, M.; Horvath, N.; Shih, C.-H.; Wayman, J. A.; Calhoun, K.; Swartz, J.; Varner, J. D. Sequence Specific Modeling of E. Coli Cell-Free Protein Synthesis. *ACS Synth. Biol.* **2018**, *7* (8), 1844–1857. https://doi.org/10.1021/acssynbio.7b00465.

(32) Miguez, A. M.; Zhang, Y.; Piorino, F.; Styczynski, M. P. Metabolic Dynamics in Escherichia Coli-Based Cell-Free Systems. *ACS Synth. Biol.* **2021**, *10* (9), 2252–2265. https://doi.org/10.1021/acssynbio.1c00167.

(33) Poole, W. Compilation and Inference with Chemical Reaction Networks. Thesis, California Institute of Technology, 2021.

(34) Garamella, J.; Marshall, R.; Rustad, M.; Noireaux, V. The All E. Coli TX-TL Toolbox 2.0: A Platform for Cell-Free Synthetic Biology. *ACS Synth. Biol.* **2016**, *5* (4), 344–355. https://doi.org/10.1021/acssynbio.5b00296.

(35) Zhang, B.-Q.; Bu, H.-L.; You, D.; Ye, B.-C. Acetylation of Translation Machinery Affected Protein Translation in E. Coli. *Appl. Microbiol. Biotechnol.* **2020**, *104*, 1–13. https://doi.org/10.1007/s00253-020-10985-2.

(36) Haft, R. J. F.; Keating, D. H.; Schwaegler, T.; Schwalbach, M. S.; Vinokur, J.; Tremaine, M.; Peters, J. M.; Kotlajich, M. V.; Pohlmann, E. L.; Ong, I. M.; Grass, J. A.; Kiley, P. J.; Landick, R. Correcting Direct Effects of Ethanol on Translation and Transcription Machinery Confers Ethanol Tolerance in Bacteria. *Proc. Natl. Acad. Sci.* **2014**, *111* (25), E2576–E2585. https://doi.org/10.1073/pnas.1401853111.

(37) Warnecke, T.; Gill, R. T. Organic Acid Toxicity, Tolerance, and Production in Escherichia Coli Biorefining Applications. *Microb. Cell Factories* **2005**, *4* (1), 25. https://doi.org/10.1186/1475-2859-4-25.

(38) Barrenechea, V.; Vargas-Reyes, M.; Quiliano, M.; Milón, P. A Complementary Mechanism of Bacterial mRNA Translation Inhibition by Tetracyclines. *Front. Microbiol.* **2021**, *12*. https://doi.org/10.3389/fmicb.2021.682682.

(39) Dopp, B. J. L.; Tamiev, D. D.; Reuel, N. F. Cell-Free Supplement Mixtures: Elucidating the History and Biochemical Utility of Additives Used to Support in Vitro Protein Synthesis in E. Coli Extract. *Biotechnol. Adv.* **2019**, *37* (1), 246–258. https://doi.org/10.1016/j.biotechadv.2018.12.006.

(40) Harris, C. R.; Millman, K. J.; Walt, S. J. van der; Gommers, R.; Virtanen, P.; Cournapeau, D.; Wieser, E.; Taylor, J.; Berg, S.; Smith, N. J.; Kern, R.; Picus, M.; Hoyer, S.; Kerkwijk, M. H. van; Brett, M.; Haldane, A.; Río, J. F. del; Wiebe, M.; Peterson, P.; Gérard-Marchant, P.; Sheppard, K.; Reddy, T.; Weckesser, W.; Abbasi, H.; Gohlke, C.; Oliphant, T. E. Array Programming with NumPy. *Nature* **2020**, *585* (7825), 357–362. https://doi.org/10.1038/s41586-020-2649-2.

Table of contents:

Figure S1: Exploring the dependence of MGaptamer and deGFP fluorescence on pH, Mg^{2+}, and 3PGA concentration. In **(A)** and **(B)**, either **(A)** 0.2µM MGaptamer-UTR1-deGFP mRNA or **(B)** 6.02µM purified deGFP was added to 9uL of phosphate-buffered saline (PBS) of the appropriate pH, some amount of 100mM Mg-glutamate to the appropriate concentration, and nuclease-free water to 10µL. The 384-well plate was incubated for 18 hours in a plate-reader at 29°C, and the endpoint concentrations, as determined by the fluorescence calibration data, are indicated in the plots shown. **(C)** 10µM purified deGFP was added to TX-TL reactions and incubated at 29°C for 18 hours. The final deGFP concentrations, as reported by the fluorescence values that were calibrated to micromolar units, are shown in the figure.

Figure S2: Effects of Mg²⁺ concentration on the TX-TL trade-off by fuel type. Each point represents one of three replicates of a particular set of fuel and Mg²⁺ concentrations, where fuel is either **(A)** 3PGA, **(B)** maltose, **(C)** pyruvate, or **(D)** succinate. The data shown reflect experiments performed in two batches of cell lysate, Batch 1 and Batch 2 (see Methods and Materials for details), and all experiments were performed using 5 nM DNA.

Figure S3: Effects of 3PGA concentration on the TX-TL trade-off across different lysate volume fractions. Each point represents one of three replicates of a particular set of 3PGA and Mg^{2+} concentrations. **(A)** Trade-off curves from three different lysate volume fractions are overlaid, where the lysate volume fraction is the volume fraction of a TX-TL reaction that is made up of cell lysate. **(B) – (D)** Individual lysate fraction trade-off curves are shown and points are colored by 3PGA concentration, where the curves correspond to lysate fractions **(B)** F_L = 0.25, **(C)** F_L = 0.33, the nominal case; and **(D)** F_L = 0.41. The data shown reflect experiments performed in cell lysate Batch 2 (see Methods and Materials for details), and all experiments were performed using 5 nM DNA.

Figure S4: Effects of 3PGA concentration on the TX-TL trade-off in systems expressing MG aptamer and deGFP under different promoters. Each point represents one of three replicates of a particular set of 3PGA and Mg^{2+} concentrations. The data are the same shown in **Figure 4**, albeit with the data from panels **(A)**, **(B)**, and **(D)** overlaid and colored by promoter type. The data shown reflect experiments performed in Batch 3 of cell lysate (see Methods and Materials for details), and all experiments were performed using 5 nM DNA.

Figure S5: Scaling of transcription and TL with increasing DNA concentration. **(A)** Five lines were chosen to intersect the trade-off curves at roughly equi-angular intervals, and the point on each trade-off curve corresponding most closely to that line is highlighted. **(B)** Integrated MG aptamer versus DNA concentration for each of the intersecting lines for the points highlighted in (A). **(C)** Maximum deGFP versus DNA concentration for each of the intersecting lines for the points highlighted in (B). All experiments were performed in Batch 2 of cell lysate.

Figure S6: Exploring the effects of total ATP versus total GTP concentration in NTP-fueled systems. Here, the systems explored were TX-TL systems using 5 nM DNA to which no central carbon fuel nor Mg^{2+} had been added. Systems fueled by NTPs were performed using a three-way panel of ATP versus GTP versus NTP mix, where each solution was added at 0, 5, 10, or 15mM. As the NTP mix also contained ATP and GTP, the total concentrations of ATP and GTP were calculated for each system by summing the amount of ATP or GTP in the NTP mix with the added amount of ATP or GTP, respectively. Results were then averaged over three replicates and two batches of cell lysate (Batch 1 and Batch 2). **(A)** Integrated MG aptamer values for NTP-fueled systems. **(B)** Maximum deGFP values for the NTP-fueled systems.

Figure S7: Examples of unusual and potentially desirable transcription and translation dynamics in TX-TL systems with no fuel or no Mg²⁺ and with additional energy. These systems have the following (NTP, Added ATP, Added GTP) concentrations in mM, where a dashed line (-) indicates that a concentration is being varied in that plot: **(A)** (0, -, 15), **(B)** (-, 5, 5), **(C)** (-, 5, 10), **(D)** (0, 5, -), **(E)** (5, 5, -), **(F)** (-, 5, 5). **(A)** shows an example of a system (at 15mM ATP)

exhibiting time-delayed transcription albeit with a high MG aptamer yield that continues increasing even at 18 hours. **(B)** and **(C)** show examples of systems with biphasic and sustained transcription (both at 15mM NTPs), respectively. **(D)** shows examples of systems with varying translation dynamics, ranging from a near-linear increase in deGFP (0mM GTP), biphasic translation (10mM GTP), slow translation (5mM GTP), and extremely time-delayed translation (15mM GTP). **(E)** demonstrates that TX-TL systems that have time-delayed translation do not necessarily have lower deGFP yields, as indicated in the comparison between the 5mM GTP and 10mM GTP cases. **(F)** shows that under certain regimes (here at 5mM ATP and 5mM GTP), TX-TL systems can exhibit similar translation dynamics, albeit with time delays that are proportional to the amount of NTPs added to the system. Each point represents one of three replicates of a particular set of NTP, Added ATP, and Added GTP concentrations. All experiments were performed at 5 nM DNA in systems using the Batch 2 lysate. Note: axes have different scales for each plot.

Figure S8: Tetracycline titrations. Titrations were performed to determine the concentration of tetracycline necessary for translation inhibition albeit without adding enough DMSO to the reaction to additionally reduce transcription or translation. Controls were performed by adding the equivalent amount of DMSO to make sure the tetracycline alone was impacting translation. **(A)** and **(B)** show transcription and translation data, respectively, for TX-TL reactions performed with 200μg/mL tetracycline. **(C)** and **(D)** show transcription and translation data, respectively, for TX-TL reactions performed with the equivalent amount of DMSO for each of the conditions in **(A)** and **(B)**. TX-TL reactions were performed using the Batch 2 lysate with 5 nM DNA at 30mM 3PGA and 8mM Mg^{2+}.

Plasmid sequences

The following are the sequences of the four plasmids used in this study. Promoters are highlighted in red, mRNA aptamer in blue, untranslated region (5' UTR) in yellow, protein sequence in green, and terminator sequence in magenta.

P$_{OR1OR2}$-MGaptamer-UTR1-deGFP-T500 sequence (pZJQ.27):

TGAGCTAACACCGTGCGTGTTGACAATTTTACCTCTGGCGGTGATAATGGTTGCAGCTAGCGGGATCCCGACTG
GCGAGAGCCAGGTAACGAATGGATCCAATAATTTTGTTTAACTTTAAGAAGGAGATATACCATGGAGCTTTTCAC
TGGCGTTGTTCCCATCCTGGTCGAGCTGGACGGCGACGTAAACGGCCACAAGTTCAGCGTGTCCGGCGAGGGC
GAGGGCGATGCCACCTACGGCAAGCTGACCCTGAAGTTCATCTGCACCACCGGCAAGCTGCCCGTGCCCTGGC
CCACCCTCGTGACCACCCTGACCTACGGCGTGCAGTGCTTCAGCCGCTACCCCGACCACATGAAGCAGCACGA
CTTCTTCAAGTCCGCCATGCCCGAAGGCTACGTCCAGGAGCGCACCATCTTCTTCAAGGACGACGGCAACTACA
AGACCCGCGCCGAGGTGAAGTTCGAGGGCGACACCCTGGTGAACCGCATCGAGCTGAAGGGCATCGACTTCAA
GGAGGACGGCAACATCCTGGGGCACAAGCTGGAGTACAACTACAACAGCCACAACGTCTATATCATGGCCGAC
AAGCAGAAGAACGGCATCAAGGTGAACTTCAAGATCCGCCACAACATCGAGGACGGCAGCGTGCAGCTCGCCG
ACCACTACCAGCAGAACACCCCCATCGGCGACGGCCCCGTGCTGCTGCCCGACAACCACTACCTGAGCACCCA
GTCCGCCCTGAGCAAAGACCCCAACGAGAAGCGCGATCACATGGTCCTGCTGGAGTTCGTGACCGCCGCCGG
GATCTAACTCGAGCAAAGCCCGCCGAAAGGCGGGCTTTTCTGTGTCGACCGATGCCCTTGAGAGCCTTCAACCC
AGTCAGCTCCTTCCGGTGGGCGCGGGGCATGACTATCGTCGCCGCACTTATGACTGTCTTCTTTATCATGCAAC
TCGTAGGACAGGTGCCGGCAGCGCTCTTCCGCTTCCTCGCTCACTGACTCGCTGCGCTCGGTCGTTCGGCTGC
GGCGAGCGGTATCAGCTCACTCAAAGGCGGTAATACGGTTATCCACAGAATCAGGGGATAACGCAGGAAAGAA
CATGTGAGCAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAAAGGCCGCGTTGCTGGCGTTTTTCCATAGGCTC
CGCCCCCCTGACGAGCATCACAAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGATA
CCAGGCGTTTCCCCCTGGAAGCTCCCTCGTGCGCTCTCCTGTTCCGACCCTGCCGCTTACCGGATACCTGTCCG
CCTTTCTCCCTTCGGGAAGCGTGGCGCTTTCTCAATGCTCACGCTGTAGGTATCTCAGTTCGGTGTAGGTCGTTC
GCTCCAAGCTGGGCTGTGTGCACGAACCCCCCGTTCAGCCCGACCGCTGCGCCTTATCCGGTAACTATCGTCTT
GAGTCCAACCCGGTAAGACACGACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGT
ATGTAGGCGGTGCTACAGAGTTCTTGAAGTGGTGGCCTAACTACGGCTACACTAGAAGGACAGTATTTGGTATC
TGCGCTCTGCTGAAGCCAGTTACCTTCGGAAAAAGAGTTGGTAGCTCTTGATCCGGCAAACAAACCACCGCTGG
TAGCGGTGGTTTTTTTGTTTGCAAGCAGCAGATTACGCGCAGAAAAAAAGGATCTCAAGAAGATCCTTTGATCTT
TTCTACGGGGTCTGACGCTCAGTGGAACGAAAACTCACGTTAAGGGATTTTGGTCATGAGATTATCAAAAAGGAT
CTTCACCTAGATCCTTTTAAATTAAAAATGAAGTTTTAAATCAATCTAAAGTATATATGAGTAAACTTGGTCTGACA
GTTACCAATGCTTAATCAGTGAGGCACCTATCTCAGCGATCTGTCTATTTCGTTCATCCATAGTTGCCTGACTCCC
CGTCGTGTAGATAACTACGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGATACCGCGAGACCCA
CGCTCACCGGCTCCAGATTTATCAGCAATAAACCAGCCAGCCGGAAGGGCCGAGCGCAGAAGTGGTCCTGCAA
CTTTATCCGCCTCCATCCAGTCTATTAATTGTTGCCGGGAAGCTAGAGTAAGTAGTTCGCCAGTTAATAGTTTGC
GCAACGTTGTTGCCATTGCTACAGGCATCGTGGTGTCACGCTCGTCGTTTGGTATGGCTTCATTCAGCTCCGGTT
CCCAACGATCAAGGCGAGTTACATGATCCCCCATGTTGTGCAAAAAAGCGGTTAGCTCCTTCGGTCCTCCGATC
GTTGTCAGAAGTAAGTTGGCCGCAGTGTTATCACTCATGGTTATGGCAGCACTGCATAATTCTCTTACTGTCATG
CCATCCGTAAGATGCTTTTCTGTGACTGGTGAGTACTCAACCAAGTCATTCTGAGAATAGTGTATGCGGCGACCG
AGTTGCTCTTGCCCGGCGTCAATACGGGATAATACCGCGCCACATAGCAGAACTTTAAAAGTGCTCATCATTGGA
AAACGTTCTTCGGGGCGAAAACTCTCAAGGATCTTACCGCTGTTGAGATCCAGTTCGATGTAACCCACTCGTGCA
CCCAACTGATCTTCAGCATCTTTTACTTTCACCAGCGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAATGCCGCA
AAAAAGGGAATAAGGGCGACACGGAAATGTTGAATACTCATACTCTTCCTTTTTCAATATTATTGAAGCATTTATC
AGGGTTATTGTCTCATGAGCGGATACATATTTGAATGTATTTAGAAAAATAAACAAATAGGGGTTCCGCGCACATT
TCCCCGAAAAGTGCCACCTGACGTCTAAGAAACCATTATTATCATGACATTAACCTATAAAAATAGGCGTATCACG
AGGCCCTTTCGTCTTCAAGAATTCTGGCGAATCCTCTGACCAGCCAGAAAACGACCTTTCTGTGGTGAAACCGG
ATGCTGCAATTCAGAGCGGCAGCAAGTGGGGGACAGCAGAAGACCTGACCGCCGCAGAGTGGATGTTTGACAT
GGTGAAGACTATCGCACCATCAGCCAGAAAACCGAATTTTGCTGGGTGGGCTAACGATATCCGCGGATCCCGAC
TGATGCGTGAACGTGACGGACGTAACCACCGCGACATGTGTGTGCTGTTC

P$_{T7}$-MGaptamer-UTR1-deGFP-T500 sequence (pMK18):

TAATACGACTCACTATAGGGAGAGCTAGCGGGATCCCGACTGGCGAGAGCCAGGTAACGAATGGATCCAATAAT
TTTGTTTAACTTTAAGAAGGAGATATACCATGGAGCTTTTCACTGGCGTTGTTCCCATCCTGGTCGAGCTGGACG
GCGACGTAAACGGCCACAAGTTCAGCGTGTCCGGCGAGGGCGAGGGCGATGCCACCTACGGCAAGCTGACCC
TGAAGTTCATCTGCACCACCGGCAAGCTGCCCGTGCCCTGGCCCACCCTCGTGACCACCCTGACCTACGGCGT
GCAGTGCTTCAGCCGCTACCCCGACCACATGAAGCAGCACGACTTCTTCAAGTCCGCCATGCCCGAAGGCTAC
GTCCAGGAGCGCACCATCTTCTTCAAGGACGACGGCAACTACAAGACCCGCGCCGAGGTGAAGTTCGAGGGCG
ACACCCTGGTGAACCGCATCGAGCTGAAGGGCATCGACTTCAAGGAGGACGGCAACATCCTGGGGCACAAGCT
GGAGTACAACTACAACAGCCACAACGTCTATATCATGGCCGACAAGCAGAAGAACGGCATCAAGGTGAACTTCA
AGATCCGCCACAACATCGAGGACGGCAGCGTGCAGCTCGCCGACCACTACCAGCAGAACACCCCCATCGGCGA
CGGCCCCGTGCTGCTGCCCGACAACCACTACCTGAGCACCCAGTCCGCCCTGAGCAAAGACCCCAACGAGAAG
CGCGATCACATGGTCCTGCTGGAGTTCGTGACCGCCGCCGGGATCTAACTCGAGCTAGCATAACCCCTTGGGG
CCTCTAAACGGGTCTTGAGGGGTTTTTTGTCGACCGATGCCCTTGAGAGCCTTCAACCCAGTCAGCTCCTTCC
GGTGGGCGCGGGGCATGACTATCGTCGCCGCACTTATGACTGTCTTCTTTATCATGCAACTCGTAGGACAGGTG
CCGGCAGCGCTCTTCCGCTTCCTCGCTCACTGACTCGCTGCGCTCGGTCGTTCGGCTGCGGCGAGCGGTATCA
GCTCACTCAAAGGCGGTAATACGGTTATCCACAGAATCAGGGGATAACGCAGGAAAGAACATGTGAGCAAAAGG
CCAGCAAAAGGCCAGGAACCGTAAAAAGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCCCCTGACGAG
CATCACAAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGATACCAGGCGTTTCCCCC
TGGAAGCTCCCTCGTGCGCTCTCCTGTTCCGACCCTGCCGCTTACCGGATACCTGTCCGCCTTTCTCCCTTCGG
GAAGCGTGGCGCTTTCTCAATGCTCACGCTGTAGGTATCTCAGTTCGGTGTAGGTCGTTCGCTCCAAGCTGGGC
TGTGTGCACGAACCCCCCGTTCAGCCCGACCGCTGCGCCTTATCCGGTAACTATCGTCTTGAGTCCAACCCGGT
AAGACACGACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTATGTAGGCGGTGCT
ACAGAGTTCTTGAAGTGGTGGCCTAACTACGGCTACACTAGAAGGACAGTATTTGGTATCTGCGCTCTGCTGAA
GCCAGTTACCTTCGGAAAAAGAGTTGGTAGCTCTTGATCCGGCAAACAAACCACCGCTGGTAGCGGTGGTTTTT
TTGTTTGCAAGCAGCAGATTACGCGCAGAAAAAAAGGATCTCAAGAAGATCCTTTGATCTTTTCTACGGGGTCTG
ACGCTCAGTGGAACGAAAACTCACGTTAAGGGATTTTGGTCATGAGATTATCAAAAAGGATCTTCACCTAGATCC
TTTTAAATTAAAAATGAAGTTTTAAATCAATCTAAAGTATATATGAGTAAACTTGGTCTGACAGTTACCAATGCTTA
ATCAGTGAGGCACCTATCTCAGCGATCTGTCTATTTCGTTCATCCATAGTTGCCTGACTCCCCGTCGTGTAGATA
ACTACGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGATACCGCGAGACCCACGCTCACCGGCTC
CAGATTTATCAGCAATAAACCAGCCAGCCGGAAGGGCCGAGCGCAGAAGTGGTCCTGCAACTTTATCCGCCTCC
ATCCAGTCTATTAATTGTTGCCGGGAAGCTAGAGTAAGTAGTTCGCCAGTTAATAGTTTGCGCAACGTTGTTGCC
ATTGCTACAGGCATCGTGGTGTCACGCTCGTCGTTTGGTATGGCTTCATTCAGCTCCGGTTCCCAACGATCAAG
GCGAGTTACATGATCCCCCATGTTGTGCAAAAAAGCGGTTAGCTCCTTCGGTCCTCCGATCGTTGTCAGAAGTAA
GTTGGCCGCAGTGTTATCACTCATGGTTATGGCAGCACTGCATAATTCTCTTACTGTCATGCCATCCGTAAGATG
CTTTTCTGTGACTGGTGAGTACTCAACCAAGTCATTCTGAGAATAGTGTATGCGGCGACCGAGTTGCTCTTGCCC
GGCGTCAATACGGGATAATACCGCGCCACATAGCAGAACTTTAAAAGTGCTCATCATTGGAAAACGTTCTTCGG
GGCGAAAACTCTCAAGGATCTTACCGCTGTTGAGATCCAGTTCGATGTAACCCACTCGTGCACCCAACTGATCTT
CAGCATCTTTTACTTTCACCAGCGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAATGCCGCAAAAAAGGGAATAA
GGGCGACACGGAAATGTTGAATACTCATACTCTTCCTTTTTCAATATTATTGAAGCATTTATCAGGGTTATTGTCT
CATGAGCGGATACATATTTGAATGTATTTAGAAAAATAAACAAATAGGGGTTCCGCGCACATTTCCCCGAAAAGT
GCCACCTGACGTCTAAGAAACCATTATTATCATGACATTAACCTATAAAAATAGGCGTATCACGAGGCCCTTTCGT
CTTCAAGAATTCTGGCGAATCCTCTGACCAGCCAGAAAACGACCTTTCTGTGGTGAAACCGGATGCTGCAATTCA
GAGCGGCAGCAAGTGGGGGACAGCAGAAGACCTGACCGCCGCAGAGTGGATGTTTGACATGGTGAAGACTATC
GCACCATCAGCCAGAAAACCGAATTTTGCTGGGTGGGCTAACGATATCCGCGGATCCCGACTGATGCGTGAAC
GTGACGGACGTAACCACCGCGACATGTGTGTGCTGTTC

P_{Tet}-MGaptamer-UTR1-deGFP-T500 sequence (pMK19):

TCCCTATCAGTGATAGAGATTGACATCCCTATCAGTGATAGAGATACTGAGCACGCTAGCGGGATCCCGACTGG
CGAGAGCCAGGTAACGAATGGATCCAATAATTTTGTTTAACTTTAAGAAGGAGATATACCATGGAGCTTTTCACT
GGCGTTGTTCCCATCCTGGTCGAGCTGGACGGCGACGTAAACGGCCACAAGTTCAGCGTGTCCGGCGAGGGC
GAGGGCGATGCCACCTACGGCAAGCTGACCCTGAAGTTCATCTGCACCACCGGCAAGCTGCCCGTGCCCTGGC
CCACCCTCGTGACCACCCTGACCTACGGCGTGCAGTGCTTCAGCCGCTACCCCGACCACATGAAGCAGCACGA
CTTCTTCAAGTCCGCCATGCCCGAAGGCTACGTCCAGGAGCGCACCATCTTCTTCAAGGACGACGGCAACTACA
AGACCCGCGCCGAGGTGAAGTTCGAGGGCGACACCCTGGTGAACCGCATCGAGCTGAAGGGCATCGACTTCAA
GGAGGACGGCAACATCCTGGGGCACAAGCTGGAGTACAACTACAACAGCCACAACGTCTATATCATGGCCGAC
AAGCAGAAGAACGGCATCAAGGTGAACTTCAAGATCCGCCACAACATCGAGGACGGCAGCGTGCAGCTCGCCG
ACCACTACCAGCAGAACACCCCCATCGGCGACGGCCCCGTGCTGCTGCCCGACAACCACTACCTGAGCACCCA
GTCCGCCCTGAGCAAAGACCCCAACGAGAAGCGCGATCACATGGTCCTGCTGGAGTTCGTGACCGCCGCCGG
GATCTAACTCGAGCAAAGCCCGCCGAAGGCGGGCTTTTCTGTGTCGACCGATGCCCTTGAGAGCCTTCAACCC
AGTCAGCTCCTTCCGGTGGGCGCGGGGCATGACTATCGTCGCCGCACTTATGACTGTCTTCTTTATCATGCAAC
TCGTAGGACAGGTGCCGGCAGCGCTCTTCCGCTTCCTCGCTCACTGACTCGCTGCGCTCGGTCGTTCGGCTGC
GGCGAGCGGTATCAGCTCACTCAAAGGCGGTAATACGGTTATCCACAGAATCAGGGGATAACGCAGGAAAGAA
CATGTGAGCAAAAGGCCAGCAAAAGGCCAGGAACCGTAAAAAGGCCGCGTTGCTGGCGTTTTTCCATAGGCTC
CGCCCCCCTGACGAGCATCACAAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGATA
CCAGGCGTTTCCCCCTGGAAGCTCCCTCGTGCGCTCTCCTGTTCCGACCCTGCCGCTTACCGGATACCTGTCCG
CCTTTCTCCCTTCGGGAAGCGTGGCGCTTTCTCAATGCTCACGCTGTAGGTATCTCAGTTCGGTGTAGGTCGTTC
GCTCCAAGCTGGGCTGTGTGCACGAACCCCCCGTTCAGCCCGACCGCTGCGCCTTATCCGGTAACTATCGTCTT
GAGTCCAACCCGGTAAGACACGACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGT
ATGTAGGCGGTGCTACAGAGTTCTTGAAGTGGTGGCCTAACTACGGCTACACTAGAAGGACAGTATTTGGTATC
TGCGCTCTGCTGAAGCCAGTTACCTTCGGAAAAAGAGTTGGTAGCTCTTGATCCGGCAAACAAACCACCGCTGG
TAGCGGTGGTTTTTTTGTTTGCAAGCAGCAGATTACGCGCAGAAAAAAAGGATCTCAAGAAGATCCTTTGATCTT
TTCTACGGGGTCTGACGCTCAGTGGAACGAAAACTCACGTTAAGGGATTTTGGTCATGAGATTATCAAAAAGGAT
CTTCACCTAGATCCTTTTAAATTAAAAATGAAGTTTTAAATCAATCTAAAGTATATATGAGTAAACTTGGTCTGACA
GTTACCAATGCTTAATCAGTGAGGCACCTATCTCAGCGATCTGTCTATTTCGTTCATCCATAGTTGCCTGACTCCC
CGTCGTGTAGATAACTACGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGATACCGCGAGACCCA
CGCTCACCGGCTCCAGATTTATCAGCAATAAACCAGCCAGCCGGAAGGGCCGAGCGCAGAAGTGGTCCTGCAA
CTTTATCCGCCTCCATCCAGTCTATTAATTGTTGCCGGGAAGCTAGAGTAAGTAGTTCGCCAGTTAATAGTTTGC
GCAACGTTGTTGCCATTGCTACAGGCATCGTGGTGTCACGCTCGTCGTTTGGTATGGCTTCATTCAGCTCCGGTT
CCCAACGATCAAGGCGAGTTACATGATCCCCCATGTTGTGCAAAAAAGCGGTTAGCTCCTTCGGTCCTCCGATC
GTTGTCAGAAGTAAGTTGGCCGCAGTGTTATCACTCATGGTTATGGCAGCACTGCATAATTCTCTTACTGTCATG
CCATCCGTAAGATGCTTTTCTGTGACTGGTGAGTACTCAACCAAGTCATTCTGAGAATAGTGTATGCGGCGACCG
AGTTGCTCTTGCCCGGCGTCAATACGGGATAATACCGCGCCACATAGCAGAACTTTAAAAGTGCTCATCATTGGA
AAACGTTCTTCGGGGCGAAAACTCTCAAGGATCTTACCGCTGTTGAGATCCAGTTCGATGTAACCCACTCGTGCA
CCCAACTGATCTTCAGCATCTTTTACTTTCACCAGCGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAATGCCGCA
AAAAAGGGAATAAGGGCGACACGGAAATGTTGAATACTCATACTCTTCCTTTTTCAATATTATTGAAGCATTTATC
AGGGTTATTGTCTCATGAGCGGATACATATTTGAATGTATTTAGAAAAATAAACAAATAGGGGTTCCGCGCACATT
TCCCCGAAAAGTGCCACCTGACGTCTAAGAAACCATTATTATCATGACATTAACCTATAAAAATAGGCGTATCACG
AGGCCCTTTCGTCTTCAAGAATTCTGGCGAATCCTCTGACCAGCCAGAAAACGACCTTTCTGTGGTGAAACCGG
ATGCTGCAATTCAGAGCGGCAGCAAGTGGGGGACAGCAGAAGACCTGACCGCCGCAGAGTGGATGTTTGACAT
GGTGAAGACTATCGCACCATCAGCCAGAAACCGAATTTTGCTGGGTGGGCTAACGATATCCGCGGATCCCGAC
TGATGCGTGAACGTGACGGACGTAACCACCGCGACATGTGTGTGCTGTTC

pTet-F30Pepper-UTR1-mTurquoise2-ECK120029600 sequence (pMK10):

Here the F30 scaffold that flanks either side of the Pepper aptamer is highlighted in orange.

TCCCTATCAGTGATAGAGATTGACATCCCTATCAGTGATAGAGATACTGAGCACTACTTTGCCATGTGTATGTGG
GTTCGCCCACATACTCTGATGATCCCCAATCGTGGCGTGTCGGCCTGCTTCGGCAGGCACTGGCGCCGGGATC
ATTCATGGCAAAATAATTTTGTTTAACTTTAAGAAGGAGATATACCAATGGTAAGTAAGGGCGAAGAGTTGTTTAC
TGGGGTAGTCCCGATCCTGGTCGAGCTGGATGGGGACGTTAACGGCCATAAATTTAGCGTGTCGGGTGAAGGC
GAGGGGGACGCGACCTACGGGAAACTGACCTTGAAGTTTATCTGTACGACCGGGAAACTTCCGGTGCCATGGC
CTACACTCGTCACGACCTTGAGCTGGGGTGTTCAATGCTTTGCACGTTACCCAGACCATATGAAACAGCATGATT
TCTTCAAAAGTGCAATGCCAGAAGGGTATGTTCAAGAGCGCACCATCTTTTTTAAGGACGATGGCAACTATAAGA
CCCGTGCCGAAGTCAAGTTCGAAGGTGACACCCTTGTCAACCGTATCGAGTTGAAAGGTATCGACTTTAAGGAA
GATGGGAATATCCTGGGTCATAAACTTGAATATAATTACTTTTCCGACAATGTTTATATCACAGCCGACAAACAAA
AAAACGGTATCAAGGCCAATTTCAAGATTCGGCATAACATCGAAGATGGGGGGGTTCAACTCGCTGATCATTACC
AGCAGAATACACCTATCGGCGACGGCCCAGTCTTATTACCTGATAATCATTACCTTAGCACCCAATCCAAACTCA
GCAAGGATCCAAACGAAAGCGCGATCACATGGTGCTTTTGGAGTTTGTCACTGCTGCTGGTATTACATTGGGTA
TGGATGAATTGTACAAAGGGGGCGGCGGTAGCCATCATCATCATCACCATTAAAGGTTTCAGCCAAAAAACTTAA
GACCGCCGGTCTTGTCCACTACCTTGCAGTAATGCGGTGGACAGGATCGGCGGTTTTCTTTTCTCTTCTCAAGCT
TATGTCTTCTACTAGTAGCGGCCGCTGCAGTCCGGCAAAAAGGGCAAGGTGTCACCACCCTGCCCTTTTTCTTT
AAAACCGAAAGATTACTTCGCGTTATGCAGGCTTCCTCGCTCACTGACTCGCTGCGCTCGGTCGTTCGGCTGC
GGCGAGCGGTATCAGCTCACTCAAAGGCGGTAAGTCGCGTGTTATACGCCCGTTATCCATGGGTATGGACAGTT
TTCCCTTTGATATGTAACGCACGTTGTGTCTCAAAATCTCTGATGTTACATTGCACAAGATAAAAATATATCATCAT
GAACAATAAAACTGTCTGCTTACATAAACAGTAATACAAGGGGTGTTATGAGCCATATTCAACGGGAAACGTCTT
GCTCCCGTCCGCGCTTAAACTCCAACATGGACGCTGATTTATATGGGTATAAATGGGCTCGCGATAATGTCGGG
CAATCAGGTGCGACAATCTATCGCTTGTATGGGAAGCCCGATGCGCCAGAGTTGTTTCTGAAACATGGCAAAGG
TAGCGTTGCCAATGATGTTACAGATGAGATGGTCCGTCTCAACTGGCTGACGGAGTTTATGCCTCTCCCGACCAT
CAAGCATTTTATCCGTACTCCTGATGATGCGTGGTTACTCACCACCGCGATTCCTGGGAAAACAGCCTTCCAGGT
ATTAGAAGAATATCCTGATTCAGGTGAAAATATTGTTGATGCGCTGGCCGTGTTCCTGCGCCGGTTACATTCGAT
TCCTGTTTGTAATTGTCCTTTTAACAGCGATCGTGTATTTCGTCTTGCTCAGGCGCAATCACGCATGAATAACGGT
TTGGTTGATGCGAGTGATTTTGATGACGAGCGTAATGGCTGGCCTGTTGAACAAGTCTGGAAAGAAATGCACAA
GCTCTTGCCATTCTCACCGGATTCAGTCGTCACTCATGGTGATTTCTCACTTGATAACCTTATTTTTGACGAGGGG
AAATTAATAGGTTGTATTGATGTTGGACGGGTCGGAATCGCAGACCGTTACCAGGACCTTGCCATTCTTTGGAAC
TGCCTCGGTGAGTTTTCTCCTTCATTACAGAAACGGCTTTTTCAAAAATATGGTATTGATAATCCTGATATGAATAA
ATTGCAGTTTCATTTGATGCTCGATGAGTTTTTTCTAATAATACTAGCAGAAATCATCCTTAGCGAAAGCTAAGGAT
TTTTTTTATCTGATTACCGCCTTTGAGTGAGCGTCGACCTAGTCCTAGCCAGGAACCGTAAAAAGGCCGCGTTGC
TGGCGTTTTTCCATAGGCTCCGCCCCCCTGACGAGCATCACAAAAATCGACGCTCAAGTCAGAGGTGGCGAAAC
CCGACAGGACTATAAAGATACCAGGCGTTTCCCCCTGGAAGCTCCCTCGTGCGCTCTCCTGTTCCGACCCTGCC
GCTTACCGGATACCTGTCCGCCTTTCTCCCTTCGGGAAGCGTGGCGCTTTCTCATAGCTCACGCTGTAGGTATC
TCAGTTCGGTGTAGGTCGTTCGCTCCAAGCTGGGCTGTGTGCACGAACCCCCCGTTCAGCCCGACCGCTGCGC
CTTATCCGGTAACTATCGTCTTGAGTCCAACCCGGTAAGACACGACTTATCGCCACTGGCAGCAGCCACTGGTA
ACAGGATTAGCAGAGCGAGGTATGTAGGCGGTGCTACAGAGTTCTTGAAGTGGTGGCCTAACTACGGCTACACT
AGAAGAACAGTATTTGGTATCTGCGCTCTGCTGAAGCCAGTTACCTTCGGAAAAAGAGTTGGTAGCTCTTGATCC
GGCAAACAAACCACCGCTGGTAGCGGTGGTTTTTTTGTTTGCAAGCAGCAGATTACGCGCAGAAAAAAAGGATC
TCAAGAAGATCCTTTGATCTTTTCTACGGGGTCTGACGCGCATAAATAGGCGTATCACGAGGCAGAATTTCAGAT
AAAAAAAATCCTTAGCTTTCGCTAAGGATGATTTCTGGAATTCGCGGCCGCTTCTAGAGACTAGTGGAAGACATG
GAG