1

Enterococcus faecalis OG1RF Evolution at Low pH Selects Fusidate-sensitive Mutants in

Elongation Factor G and at High pH Selects Defects in Phosphate Transport

By

Bailey A. Fitzgerald,[1] Ayman Wadud,[1] Zachary Slimak, and Joan L. Slonczewski*

*Corresponding author: Joan L. Slonczewski, Kenyon College, Gambier, OH 43022

[1]These two authors contributed equally to this study.

ABSTRACT

Enterococcus bacteria inhabit human and soil environments that show a wide range of pH.

Strains include commensals as well as antibiotic-resistant pathogens. We investigated adaptation

to pH stress in *E. faecalis* OG1RF by conducting experimental evolution in acid (pH 4.8), neutral

pH (pH 7.0), and base (pH 9.0). Serial planktonic culture was performed for 500 generations, and

in high-pH biofilm culture for four serial bead transfers. Nearly all mutations led to

nonsynonomous codons, indicating adaptive selection. All acid-adapted clones from planktonic

culture showed a mutation in *fusA* (encoding elongation factor G). The acid-adapted *fusA*

mutants had a tradeoff of decreased resistance to fusidic acid (fusidate). All base-adapted clones

from planktonic cultures, and some from biofilm-adapted cultures, showed mutations affecting

the Pst phosphate ABC transporter (*pstA*, *pstB*, *pstB2*, *pstC*) and *pyrR* (pyrimidine biosynthesis

regulator/uracil phosphoribosyltransferase). Biofilm culture produced small-size colonies on

brain-heart infusion agar; these variants each contained a single mutation in *pstB2*, *pstC*, or *pyrR*.

The *pst* and *pyrR* mutants outgrew the ancestral strain at pH 9.2, with a tradeoff of lower growth

at pH 4.8. Additional genes that had a mutation in multiple clones evolved at high pH (but not at

low pH) include *oppBCDF* (oligopeptide ABC transporter), *ccpA* (catabolite control protein A),

and *ftsZ* (septation protein). Overall, experimental evolution of *E. faecalis* showed strong pH

dependence, favoring fusidate-sensitive elongation factor G modification at low pH and loss of

phosphate transport genes at high pH.

IMPORTANCE

E. faecalis bacteria are found in dental biofilms where they experience low pH as a result of

fermentative metabolism. Thus the effect of pH on antibiotic resistance has clinical importance.

In endodontal infections, enterococci can resist calcium hydroxide therapy that generates

extreme high pH. In other environments such as soil and plant rhizosphere, enterococci

experience acidification associated with climate change. Thus the pH modulation of natural

selection in enterococci is important for human health as well as for understanding soil

environments.

INTRODUCTION

Enterococcus faecalis are gram-positive fermenter bacteria found in a wide range of habitats, including dental biofilms and the gastrointestinal tract of humans (1–4) as well as soil and plants (5). Pathogenic strains cause a variety of infections throughout the body, which are of special concern in hospitals due to the acquired vancomycin-resistance in certain strains and an intrinsic antibiotic-resistance to numerous other antibiotics (6). *Enterococcus* species are known for their ability to tolerate a wide range of conditions such as salinity, temperature, and pH (6–9). Survival at low pH is of importance for enteric infections, and at high pH is relevant to dental biofilms that are treated with calcium hydroxide pastes (10, 11). Multidrug-resistant enterococci are isolated from environmental sources such as soil (12) and the digestive microbiomes of farm animals, insects and nematodes (13). At the same time, environmental enterococci with wide-ranging stress resistance offer potential benefits such as tolerance of toxic metals (14) and probiotics (15).

Yet surprisingly little is known about the genetics of enterococcal adaptation to acid or base. In *E. faecium*, acid shock leads to metabolomic changes such as elevated production of dipeptides (16). Acid stress in *Enterococcus* has been studied as part of general stress responses (17–19). At high pH, such as in the presence of the dental application of calcium hydroxide, *Enterococcus* survival requires maintenance of proton motive force (20). Cellular effects of high pH include changes in morphology (21) and in metabolism and protein synthesis (22, 23).

In Gram-negative bacteria, acid adaptation is associated with fitness tradeoffs such as the loss of antibiotic resistance. *Escherichia coli* K-12 experimental evolution at low pH (24, 25) or in the presence of benzoic acid (26, 27) leads to loss of genes encoding multidrug transporters

driven by the proton-motive force. It was therefore of interest to explore pH stress adaptation and

fitness tradeoffs in a Gram-positive model organism.

We investigated the genetics of pH stress in the model organism *Enterococcus faecalis*

OG1RF (2) by means of experimental evolution of planktonic and biofilm cultures. For

comparison, in *E. coli*, experimental evolution in acid or base yields mutations in a number of

stress response genes, such as amino-acid decarboxylases (24, 25, 28) and acid-resistance

regulators (29). In *E. faecalis* pH stress is also important for biofilm formation, including

endodontic infections (20, 30). A method to address stress effects in biofilms is experimental

evolution by serial bead transfer. Bead-transfer experiments yield informative genetic variants in

other environmental bacteria such as *Pseudomonas* and *Burkholderia* species (31, 32). In

Enterococcus species, experimental evolution has been applied to studies of virulence and

antimicrobial susceptibility (33, 34) but not to more fundamental stress factors such as pH.

We find that experimental evolution yields pH-adaptative mutations in a number of

genes, notably those encoding elongation factor G (*fusA*) at low pH and the phosphate ABC

transporter (*pstABC*) at high pH. Mutations in *fusA* may confer resistance to fusidic acid (35, 36);

but our *fusA* mutants show a tradeoff of fusidate sensitivity. The *pts* loci are associated with

virulence (37, 38). In our biofilm-evolved strains, base-selected mutations in *pstABC* and in *pyrR*

(pyrimidine biosynthesis regulator) led to small-colony growth, a class of phenotype associated

with virulence in clinical strains (39, 40).

RESULTS

Experimental evolution of acid- and base-adapted mutants. Experimental evolution of

E. faecalis OG1RF was conducted by microplate serial dilutions at 1:250, as described under

Methods. Eight evolving populations were maintained in brain-heart infusion (BHI) buffered at

each pH: pH 4.8, pH 7.0, and pH 9.0. After 500 generations, one isolate from each population

was resequenced for comparison with the OG1RF reference (2) using the *breseq* resequencing

pipeline (41, 42).

The resequenced clones had a range of 1-5 mutations per clone. The mutations predicted

were nearly all point mutations or small deletions (Table 1). Only one deletion encompassed

more than one gene (genes encoding catechol 2,3 dioxygenase and a multidrug ABC transporter,

in clone C06). Overall, the ratio of nonsynonymous to synonymous codon changes was

extremely high, with only one silent mutation out of 48. The high proportion of nonsynonymous

changes indicates the occurance of adaptive selection (43). Another sign of adaptive selection is

that two-thirds of the mutant genes were shared by at least two clones, although the mutations

were at different positions. The different pH classes showed a significant difference in mutation

frequency:

Acid-evolved *fusA* mutants are fusidate-sensitive at pH 7.0. In our acid-evolved

clones, the gene that showed the most mutations encodes elongation factor G (*fusA*), an essential

translation protein associated with virulence and biofilm formation (44–46). The ancestral strain

OG1RF is resistant to fusidic acid, associated with the *fusA* mutation C316A (2). We found an

added missense mutation in *fusA* in every acid-evolved clone and in five of our eight pH 7-

evolved clones. Six different missense mutations occurred, in different strains. No *fusA*

mutations were found in populations evolved at high pH. Two acid-evolved clones contained an

additional mutation affecting a gene of unknown function, RS05200, encoding a helix-turn-helix

protein. Another clone showed a missense mutation in *xdh* (xanthine dehydrogenase).

The mutant strains from pH 4.8-evolved populations were tested for growth advantage

during culture at low pH. (**Fig. 1**). At all three pH levels (pH 4.8, pH 7.0, pH 9.2) the growth

curves for mutants were indistinguishable from those of the ancestral strain. For all strains, the

steepest rise for log-phase growth occurred at the optimal condition of pH 7. No evidence for

directional selection at low pH was seen under the conditions tested.

Since all our acid-evolved clones and four of the eight clones evolved at pH 7 showed

additional point mutations in *fusA*, we tested all clones for fusidate resistance in comparison with

the ancestor OG1RF (**Fig. 2 and S1**). All clones containing a *fusA* mutation had lower fusidate

resistance than OG1RF. However, the clones lacking a *fusA* mutation (F08, F09, F10, C08)

showed fusidate resistance indistinguishable from that of the ancestor. Each of the six *fusA*

mutations incurred a single amino-acid substitution at sites different from the OG1RF mutation

(2). Thus it appears that various kinds of changes in FusA structure can reverse the OG1RF

resistance phenotype.

Base-evolved populations had mutations in phosphate transport (*pst*) and

metabolism (*pyr*). The base-evolved clones had the largest number of mutations overall (Table

1). All but one of the eight strains from pH 9.0 evolution showed a mutation in a gene involving

phosphate metabolism: *pstB*, *pstB2* (both encoding a phosphate ATP-binding protein of an ABC

transporter (47, 48)and *pyrR* (encoding a bifunctional pryimidine transcriptional regulator/ uracil

phosphribosyltransferase (49–51)). A majority of these mutations were frameshift or nonsense

codons that would knock out the gene. Each strain showed just one mutation in one of these three

genes. Several base-evolved clones had additional mutations. Four strains had deletions affecting

ccpA (catabolite control protein A), and four strains had deletions in various subunits of

oppBCDF (an oligopeptide ABC transporter). Two strains had missense mutations in an ArgR-

family transporter, and another two strains had mutations in septation protein FtsZ.

Several genes were mutated in populations evolved at pH 7.0, as well as in those at pH

9.0. These included *opp1BCDR*, *ccpA*, *walK* (cell wall metabolism sensor histidine kinase), and

rpoB, *rpoC* (RNA polymerase subunits beta and beta prime). The only gene mutated solely in

two pH 7-evolved clones was *rny* (ribonuclease Y).

Biofilm evolution by bead transfer yields small-colony clones. The evolution

experiment was modified to favor genetic variants that adhere to a bead forming a biofilm (31,

32). Cultures underwent four bead transfers over six days, generating approximately 25-50

doublings. The final populations were plated on BHI and inspected for variant phenotypes. No

phenotypes were observed after evolution in acid (pH 4.8) or at pH 7, but the pH 9-evolved

populations showed two colony size morphs, small and large. Three small-colony clones were

found that retained the phenotype upon reculturing: (strains M02, M06, and M08) (**Figure 3**).

The relative size of their colonies compared to those of the ancestor was approximately: M02,

60%; M06, 30% M08, 60%. The genomes of these clones were sequenced, and each revealed a

single mutation (Table 1). The mutations affected genes *pstB2* (M02), *pstC* (M06), and *pyrR*

(M08).

Base-evolved clones with *pst* and *pyr* mutations showed evidence of directional

selection. We sought evidence for directional selection by observing the batch growth profiles

for clones from populations evolved at pH 4.8, pH 7.0 and pH 9.0. Each of the clones was

cultured overnight in the three different pH-buffered media, for physiological adaptation to each

condition. The strongest evidence for directional selection was observed for pH 9-evolved strains

containing a mutation in one of the phosphate-related genes. **Figure 4** shows the growth curves

for the biofilm pH 9-evolved strains that each contain one mutation detected by *breseq* in *pstB2*

(M02), *pstC* (M06), and *pyrR* (M08). In each case, the log-phase growth (during approximately

0-7 h) was significantly lower at pH 4.8 for the mutant replicates than for replicate curves of the

ancestral strain OG1RF; and significantly higher than the ancestor at pH 9.2. (Significance was

tested at 6 h, $p < 0.05$). No consistent selection effect was observed at pH 7.0. These

observations are consistent with selection for a high pH-adaptive mutation following prolonged

serial culture at high pH, with the directional tradeoff of loss of growth rate at low pH.

Similar evidence of base-adaptive selection was demonstrated for seven of the eight

mutant clones obtained from high-pH planktonic evolution (**Fig. S2**). Only clone C10 showed

substantial decrease in growth. Of the eight base-evolved clones, all but clone C12 showed a

mutation in *pst* or *pyrR*, as well as mutations in other genes. All eight clones showed a small-

colony phenotype on BHI media.

The mutations in phosphate transport favored at high pH might be expected to lead to a

dependence on phosphate concentration. However, amendment of the culture media with 10 mM

or 50 mM phosphate did not significantly affect the growth curves of evolved strains or of the

ancestral strain, either at pH 4.8 or at pH 9.2.

DISCUSSION

The adaptability of *Enterococcus* bacteria to environmental stresses such as acid or base

is important for the human microbiome as well as for soil and plant-associated communities (9,

17–22, 30, 52). Few studies however employ the approach of experimental evolution to examine

stress factors under long-term adaptation. Most serial-culture studies of *Enterococcus* sp. focus

on virulence or antibiotic resistance rather than environmental factors (34, 53).

We show that experimental evolution of *E. faecalis* leads to distinct pH-dependent

patterns of mutations (Table 1). In most cases, genes or operons with variants in multiple clones

(that is, the most likely to result from adaptation) are affected only at one end of the pH range;

that is, in acid or base but not both. This finding is important as most studies of *Enterococcus*

adaptation do not account for pH.

Each of the eight acid-adapted clones had one non-synonymous codon replacement

affecting elongation factor G. Three clones from different populations had alanine replaced by

threonine; other clones had one of three different substitutions. Four out of eight populations

evolving at pH 7 also had a nonsynonymous substitution in *fusA*, but no *fusA* variants appeared

at high pH. A possible explanation could be that decrease of cell pH destabilizes the elongation

factor G protein, and that the mutations restore protein function. While directional selection for

acid adaptation was not shown under the conditions tested, the high frequency of *fusA* missense

mutations argues for the occurrence of seletion. Similarly, acid-dependent evolution in *E. coli*

leads to mutations in amino-acid decarboxylases known to help pH regulation, although

phenotypes are not detected (25).

The *fusA* mutations did show an important tradeoff, the loss of resistance to fusidic acid.

This finding implies that *E. faecalis* growth at low pH, such as in dental biofilms and in the

intestine (17, 52), could select against some forms of antibiotic resistance. Previously, acid (pH

5.5) increases the fusidate sensitivity of clinical isolates of methicillin-resistant *Staphylococcus*

aureus (54). Thus it might be of interest to investigate the role of pH in fusidate therapy.

High pH effects on *Enterococcus* species are important for dental microbiomes because

these bacteria can resist endodontic therapies involving application of high-pH pastes such as

calcium hydroxide (20, 30). The basis of high-pH resistance is poorly understood. A number of

genes are upregulated at high pH (21, 22). One of these, encoding the key septation protein FtsZ,

had an amino-acid substitution in two of our high-pH clones (Table 1).

We show evidence for directional selection of variants that knock out a component of the

PstABC phosphate uptake system, in most cases by frameshifting deletion or by nonsense

substitution (Table 1). Each *pst* mutation we found was associated with higher growth at high pH

but lower growth at low pH (Figure 2). The *pstC* mutant (strain M06) showed the smallest

colonies; this would make sense if *pstC* knocks out the entire phosphate uptake whereas

mutations in the *pstB* or *pstB2* redundent genes only partly cut phosphate uptake.

The growth rate change of *pst* and *pyrR* mutations led to visibly smaller colonies on BHI

media. While no change in antibiotic resistance was found under our conditions, in clinical

studies small-colony variants are associated with drug resistance and increased virulence (39, 40,

55). Mutations in *pst* genes are associated with clinical virulence (Lieberman) and altered

antibiotic resistance (48). In soil and plant root communities, phosphate release is maximized at

pH below 7 (56, 57) and has been proposed to contribute to bacterial pH regulation (58). In this

model, phosphate release is inhibited at low pH and activated at high pH. Blocking phosphate

uptake at high pH could enhance alkali-induced phosphate release.

The effect of *pyrR* loss and it relationship to phosphate is unclear. One possible

connection to phosphate transport is the uracil phosporibosyltransferase activity of this regulator

(50). This enzyme reaction releases pyrophosphate and has a high pH optimum. Thus it could be

that deletion of *pyrR* decreases phosphate release at high pH. In pathogenic staphylococci, *pyrR*

is associated with biofilm formation and virulence (51, 59).

Additional mutated genes in high-pH evolved population include two metabolic loci, the

oligotransport operon *oppBCD*, and the catabolite reculator *cppA*. Changes in these genes could

enhance metabolism under the lowered energy condition of high pH, where the transmembrane

pH difference is inverted and the proton potential consists solely of charge difference (60).

Overall, our study shows how *E. faecalis* exposure to acid or base can lead to evolutionary shifts

with potential importance for host adaptation.

MATERIALS AND METHODS

Bacterial strains and culture media. All strains of *E. faecalis* are derivatives of *E. faecalis*

OG1RF (2) kindly provided by Barbara Murray. For evolution serial culture, strains were

cultured in pH-buffered brain-heart infusion medium under BSL-2 conditions at 25°C. For

growth curves, incubation was at 37ºC. Buffers for experimental evolution included 100 mM

homopiperazine-1,4-bis(2-ethanesulfonic acid) (HOMOPIPES, pK_a=4.55), and 100 mM N-

cyclohexyl-3-aminopropanesulfonic acid (CAPS, pK_a=10.4). For some of the phenotype testing

at pH 7, the buffer included in growth medium was piperazine-N,N′-bis(2-ethanesulfonic acid)

(PIPES, pKa=6.8). pH was adjusted with KOH. Fusidic acid was from Millipore-Sigma. Some

colony isolations were performed on tryptic soy agar (TSA).

Experimental evolution of planktonic cells. Serial culture was performed by dilution into

deepwell microplates daily, or after two days (every sixth dilution). Eight serial cultures were

maintained at each pH: pH 4.8, pH 7.0, and pH 9.0. For each dilution, 2 μL of the previous

culture was transferred into 1000 μL (1:250) of fresh medium. All media consisted of brain heart

infusion media buffered with 0.1 M each of CAPS HOMOPIPES adjusted with KOH to the

appropriate pH. Two days growth was assumed to yield approximately 8 generations

(doublings). All evolving populations were propagated to at least 500 generations. From every

sixth dilution, 200 μl was combined with 100 μl of 50% glycerol and transferred to a deepwell

plate for storage at -80°C.

Once evolved populations reached ~500 generations, each well was streaked on individual

plates of brain-heart infusion agar (BHI) and cultured for 48 hours. Two clones were chosen

from each well population and used to create individual overnight cultures in BCH. 2 μL of the

first set of overnight solutions were used to inoculate 1 ml of fresh BCH overnights. A portion of

each clone was converted to freezer stock, and a portion was extracted for genomic DNA.

Experimental evolution of biofilms. Biofilm evolution was performed by serial bead transfer,

based on methods devised by Cooper and colleagues (31, 32). Four replicate populations of *E.*

faecalis OG1RF were grown at 25°C in the presence of 7-mm polystyrene beads suspended in 2

ml of BHI buffered with CAPS and HOMOPIPES, adjusted to the appropriate pH. The low pH

culture was grown at pH 4.8, while the neutral pH culture was grown at 7.0, and the high pH

culture was grown at pH 9.0. Populations were selected for reversible surface attachment through

transfer of a bead to a new test tube, in which cells must adhere to a new bead in order to persist,

after 48 hours. This process was repeated by alternating white and black beads for four serial

dilutions.

At the end of the dilution series, each bead was placed in 1 ml BHI medium to wash off

most planktonic cells; then placed again in 1 ml fresh BHI and vortexed for 1 min. The vortexed

suspension was then plated on BHI agar plates and incubated 48 h. Plates were then observed for

novel colony morphologies.

Growth Curves. From streaked TSA or BHI plates of ~500-generation evolution isolates, single

colonies were picked for overnight culture in BCH buffered at the same pH as the subsequent

microplate (pH 4.8, pH 7.0, or pH 9.2). 4 μL of the overnight culture of *E. faecalis* was

transferred into sterile 96-well microplates with 200 μL (1:100) of BCH or other media with 100

mM HOMOPIPES and 100 mM CAPS adjusted to the appropriate pH. Growth curves were

performed on a Spectramax microplate reader (Molecular Devices) at OD_{600} with OD reads

taken every 15 minutes for 22 hours. Comparison of batch growth was performed by T-test at

standard intervals post inoculation.

DNA sequence analysis. For the planktonic evolution cultures, DNA was extracted from one

selected ~500-generation isolate of each well (24 isolates) and from the OG1RF(B) ancestral

strain (2 isolates). The 24 sequenced clones were designated BF01-14 and C01-12. For the

biofilm evolution at pH 9, the final biofilm bead suspensions were streaked on BHI agar. After

48 hours, one large colony and one small colony variant (SCV) were isolated from each plate

and streaked on new BHI plates separately. The isolates were plated on BHI twice more to

ensure the persistence of the growth phenotype. Three large-colony isolates (M1, M5, M7) and

three small-colony isolates (M2, M6, M8), were selected for DNA preparation and sequencing.

Genomic DNA was extracted using the MasterPure Gram Positive DNA Purification Kit.

Purity and concentrations of DNA were determined using a NanoDrop 2000 spectrophotometer

(Thermo Fisher Scientific). DNA was sequenced by Admera Health (South Plainfield, NJ).

Library preparation was performed using NexteraXT library kit (Illumina, California, USA).

Sequencing was performed via Illumina NovaSeq (Illumina, California, USA) with a read length

of 150 PE for 40 M PE reads.

Mutations were called from the DNA sequences by alignment to the *E. faecalis* OG1RF

reference sequence with NCBI accession number NC_017316.1 using the *breseq* version

0.36.1 resquencing pipeline (41, 42). Resulting genomic data had approximately 300-fold

coverage. Our lab stock of OG1RF (the ancestor for all evolution experiments) showed two

differences from the reference that appeared in all our evolved clones: insertion of A at genome

position 260,670, causing frameshift in gene RS01405 (acyl-ACP thioesterase); and T deletion at

position 2,108,103 within an intergenic region.

Data availability

All sequence data are deposited at NCBI, SRA PRJNA933188. Reviewer link:

https://dataview.ncbi.nlm.nih.gov/object/PRJNA933188?reviewer=h51sokrkgscppqt0618086ihqj

AUTHOR CONTRIBUTIONS

BAF performed the planktonic evolution, and AW performed the biofilm evolution; they both

drafted a manuscript. BAF, AW and ZS conducted growth curves. JLS conceived the study,

mentored the students and completed the manuscript.

ACKNOWLEDGMENTS

This work was supported by National Science Foundation award MCB-1923077. B. F. and A.

W. designed the study, conducted experiments and drafted a manuscript. Z. S. conducted

experiments and analysis. J. S. conceived the study, mentored the students and completed the

manuscript for publication. We thank Daniel Barich for assistance with computing and statistics.

TABLES

Table 1. Mutations in clones from experimental evolution*

Table spanning strain columns (pH 4.8: F03–C04; pH 7.0: F07–C08; pH 9.0: F11–C12; Biofilm: M2–M8) with mutation annotations.

Position	F03	F04	F05	F06	C01	C02	C03	C04	F07	F08	F09	F10	C05	C06	C07	C08	F11	F12	F13	F14	C09	C10	C11	C12	M2	M6	M8	Mutation	Annotation	Gene(s)	Function
	___ pH 4.8 ___								___ pH 7.0 ___								___ pH 9.0 ___								Biofilm						
109,326																					X							C→T	R63H (C<u>G</u>T→C<u>A</u>T)	RS00485 ←	ArgR family transcriptional regulator
109,349																	X											A→T	N55K (AA<u>T</u>→AA<u>A</u>)	RS00485 ←	ArgR family transcriptional regulator
157,299									X				X															A→G	M16V (<u>A</u>TG→<u>G</u>TG)	fusA →	elongation factor G
157,452	X		X	X				X																				G→A	A67T (<u>G</u>CG→<u>A</u>CG)	fusA →	elongation factor G
157,461		X																										G→A	A70T (<u>G</u>CA→<u>A</u>CA)	fusA →	elongation factor G
157,623											X																	A→G	T124A (<u>A</u>CT→<u>G</u>CT)	fusA →	elongation factor G
157,638					X		X																					C→T	P129S (<u>C</u>CT→<u>T</u>CT)	fusA →	elongation factor G
158,374						X									X													C→T	A374V (G<u>C</u>T→G<u>T</u>T)	fusA →	elongation factor G
412,086											X																	C→A	P151Q (CC<u>A</u>→CA<u>A</u>)	RS02155 →	polysaccharide biosynthesis protein
669,400														X														C→T	M171I (AT<u>G</u>→AT<u>A</u>)	mvk ←	mevalonate kinase
674,675														X														G→T	G119W (G<u>G</u>G→T<u>G</u>G)	opp1B →	oligopeptide ABC permease Opp1B
674,889																							X					C→A	S190* (TC<u>G</u>→TA<u>G</u>)	opp1B →	oligopeptide ABC permease Opp1B
676,283											X																	G→C	R341P (C<u>G</u>T→C<u>C</u>T)	opp1C →	oligopeptide ABC permease Opp1C
676,885																				X	X							G→T	D188Y (<u>G</u>AT→<u>T</u>AT)	opp1D →	oligopeptide ABC ATP-binding protein Opp1D
677,878																	X											Δ9 bp	coding (497-505/957)	opp1F →	oligopeptide ABC ATP-binding protein Opp1F
756,900																	X											A→G	T253A (<u>A</u>CA→<u>G</u>CA)	ftsZ →	cell division protein FtsZ
756,942													X															G→A	G267S (<u>G</u>GT→<u>A</u>GT)	ftsZ →	cell division protein FtsZ
906,596					X																							C→T	Q120* (<u>C</u>AG→<u>T</u>AG)	ebpA →	endocarditis biofilm pilus tip protein
1,007,216																					X							C→T	S384F (T<u>C</u>T→T<u>T</u>T)	walK →	cell wall metabolism sensor His kinase
1,007,741														X														G→A	R559Q (C<u>G</u>A→C<u>A</u>A)	walK →	cell wall metabolism sensor His kinase
1,041,849		X																										C→T	Q117* (<u>C</u>AA→<u>T</u>AA)	RS05200 →	helix-turn-helix domain-containing protein
1,042,664					X																							C→A	C388* (TG<u>C</u>→TG<u>A</u>)	RS05200 →	helix-turn-helix domain-containing protein
1,137,647							X																					A→G	Q367Q (CA<u>A</u>→CA<u>G</u>)	RS05670 →	flotillin family protein
1,200,146															X													C→T	P32S (<u>C</u>CG→<u>T</u>CG)	dhaL ←	dihydroxyacetone kinase subunit L
1,213,195																						X						A→C	Q182P (CA<u>G</u>→CC<u>G</u>)	RS06020 →	CBS domain-containing protein
1,213,721																			X									Δ171 bp	coding(1071-1241/1320)	RS06020 →	CBS domain-containing protein
1,213,821																				X								(ATTTGC)₁→₂	coding (1171/1320)	RS06020 →	CBS domain-containing protein
1,301,608											X																	A→T	D272V (G<u>A</u>T→G<u>T</u>T)	rpoD →	RNA polymerase sigma factor
1,494,315																			X									Δ189 bp	coding (273-461/537)	pyrR ←	pyr transc. regulator/ U phosphoribosyltransferase
1,494,468													X															A→C	V103G (G<u>T</u>C→G<u>G</u>C)	pyrR ←	pyr transc. regulator/ U phosphoribosyltransferase
1,494,663																										X		+A	coding (113/537)	pyrR ←	pyr transc. regulator/ U phosphoribosyltransferase
1,514,647					X																							Δ210 bp	coding (511-720/1002)	ccpA ←	catabolite control protein A
1,514,742													X															Δ96 bp	coding (530-625/1002)	ccpA ←	catabolite control protein A
1,514,751																				X								Δ96 bp	coding (521-616/1002)	ccpA ←	catabolite control protein A
1,515,124													X															Δ81 bp	coding (163-243/1002)	ccpA ←	catabolite control protein A
1,526,361																							X					G→A	Q222* (<u>C</u>AA→<u>T</u>AA)	ptsB ←	phosphate ABC transp. ATP-binding protein
1,526,565													X															G→A	Q154* (<u>C</u>AA→<u>T</u>AA)	ptsB ←	phosphate ABC transp. ATP-binding protein
1,526,748																			X									G→A	Q93* (<u>C</u>AA→<u>T</u>AA)	ptsB ←	phosphate ABC transp. ATP-binding protein
1,527,396																									X			G→A	A150V (GCA→GTA)	pstB2 ←	phosphate ABC transp. ATP-binding protein
1,527,657																				X								G→A	T63M (A<u>C</u>G→A<u>T</u>G)	pstB2 ←	phosphate ABC transp. ATP-binding protein
1,528,108																					X							C→G	G213A (G<u>G</u>A→G<u>C</u>A)	pstA ←	phosphate ABC transp. permease
1,529,518																										X		Δ1 bp	coding (148/921)	pstC ←	phosphate ABC transp. permease

Position	Change	Amino acid	Gene	Description
1,548,299	T→G	Q23P (CAA→CCA)	purN ←	phosphoribosylglycinamide formyltransferase
1,811,611	C→A	G88W (GGG→TGG)	RS08840 ←	ABC transporter ATP-binding protein
1,896,891	T→C	intergenic (-93/+306)	RS09205 ← / ← RS09210	xanthine phosphoribosyltransferase/hydrolase
2,060,079	C→T	E284K (GAA→AAA)	xdh ←	selenium-dependent xanthine dehydrogenase

Position	Change	Amino acid	Gene	Description
2,082,421	Δ999 bp		[RS10080]–[RS10085]	catechol 2,3-dioxygenase; multidrug ABC trans
2,114,057	C→T	G18S (GGT→AGT)	RS10245 ←	GtrA family protein
2,123,718	C→T	R53C (CGC→TGC)	RS10295 →	GntR family transcriptional regulator
2,181,891	+A	coding (74/1269)	RS10530 ←	exonuclease SbcCD subunit D
2,306,141	C→T	intergenic (-14/+376)	RS11200 ← / ← RS11205	MarR family regulator/YkuJ family protein
2,354,333	4 bp→GAAA	intergenic (-96/+293)	RS11435 ← / ← tpx	valine--tRNA ligase/thiol peroxidase
2,366,924	T→A	intergenic (-60/+32)	rbsK ← / ← RS11500	ribokinase/substrate-binding protein
2,444,841	Δ1 bp	coding (269/1656)	RS11840 →	peptide ABC transporter substrate-binding protein
2,572,276	G→A	A464V (GCA→GTA)	rny ←	ribonuclease Y
2,627,715	C→A	L1162F (TTG→TTT)	rpoC ←	RNA polymerase subunit beta prime
2,630,992	T→C	Y70C (TAT→TGT)	rpoC ←	RNA polymerase subunit beta prime
2,633,037	G→A	R649C (CGT→TGT)	rpoB ←	RNA polymerase subunit beta
2,633,525	C→T	R486H (CGT→CAT)	rpoB ←	RNA polymerase subunit beta

*Clone cell shading indicates: Pink, evolved at pH 4.8; violet, evolved at pH 7.0; blue, evolved at pH 9.0; cyan, biofilm evolution at pH 9.0. Mutation cell shading: pink, *fusA* mutations; blue, mutations in *pstABC*, *pstB2*, and *pyrR*.

FIGURES

Figure 1. Acid-selected *fusA* **mutants show no growth enhancement.** *E. faecalis* strains were

isolated after 500 generations of planktonic evolution (described under Methods) at pH 4.8. Each

clone containing a single mutation in *fusA* was cultured approximately 20 h at 37°C in BHI

buffered at pH 4.8, at pH 7.0, or at pH 9.2, then 2 µl was inoculated into 200 µl at the same pH.

Four representative curves are shown for each strain: F03 (*fusA*), brown lines (A, B, C); F05

(*fusA*), red lines (D, E, F); OG1RF (ancestor), black lines for all panels.

Figure 2. Evolved clones with *fusA* mutations lose resistance to fusidic acid. For each clone, eight replicate cultures were compared with replicates of the ancestor OG1RF. All were cultured with 400 μg/ml fusidic acid, in BHI buffered at pH 7.0 at 37°C. **A, B, C.** Clones evolved at pH 4.8 (red lines) show lower growth than strain OG1RF (black lines). Endpoint growth was compared at 22 h using t-test ($p < 0.01$). **D, E, F.** Clones evolved at pH 7.0 (purple lines) show growth rates indistinguishable from that of strain OG1RF (black lines).

OG1RF   M08 (*pyrR*)

M02 (*pyrB2*)   M06 (*pstC*)

351 **Figure 3. Small-colony phenotype of biofilm-evolved isolates.** Each biofilm-evolved strain of

352 *E. faecalis* OG1RF was streaked on BHI and cultured approximately 20 h at 25ºC, then

353 approximately 4 h at 37°C.

Figure 4. High pH-selected biofilm mutants show increased growth at high pH and decreased growth at low pH. *E. faecalis* strains M02, M06, M08 were isolated as small colonies from bead-transfer biofilm evolution of *E. faecalis* OG1RF at pH 9.0 (described under Methods). Each clone was cultured approximately 20 h at 37°C in BHI buffered at pH 4.8, at pH 7.0, or at pH 9.2, then 2 μl was inoculated into 200 μl of BHI buffered at the same pH. Four representative curves are shown for each strain: M02 (*pstB2*), cyan lines (A, B, C); M06 (*pstC*),

blue lines (D, E, F); M08 (*pyrR*), violet lines (G, H, I); OG1RF (ancestor), black lines for all

panels. Culture growth was compared at time 6 h by t-test ($p < 0.05$).

REFERENCES

1. Arias CA, Murray BE. 2012. The rise of the *Enterococcus*: Beyond vancomycin
resistance. Nat Rev Microbiol 10:266–278.

2. Bourgogne A, Garsin DA, Qin X, Singh K v., Sillanpaa J, Yerrapragada S, Ding Y,
Dugan-Rocha S, Buhay C, Shen H, Chen G, Williams G, Muzny D, Maadani A, Fox KA,
Gioia J, Chen L, Shang Y, Arias CA, Nallapareddy SR, Zhao M, Prakash VP, Chowdhury
S, Jiang H, Gibbs RA, Murray BE, Highlander SK, Weinstock GM. 2008. Large scale
variation in *Enterococcus faecalis* illustrated by the genome analysis of strain OG1RF.
Genome Biol 9:R110.

3. Lebreton F, Willems RJL, Gilmore MS. 2014. Enterococcus Diversity, Origins in Nature,
and Gut ColonizationEnterococci: From Commensals to Leading Causes of Drug
Resistant Infection. http://www.ncbi.nlm.nih.gov/pubmed/24649513.

4. Dubin K, Pamer EG. 2017. *Enterococci* and their interactions with the intestinal
microbiome. Microbiol Spectr 5:BAD-0014-2016.

5. Švec P, Vandamme P, Bryndová H, Holochová P, Kosina M, Mašlaňová I, Sedláček I.
2012. Enterococcus plantarum sp. nov., isolated from plants. Int J Syst Evol Microbiol
62:1499–1505.

6. García-Solache M, Rice LB. 2019. The enterococcus: A model of adaptability to its
environment. Clin Microbiol Rev. American Society for Microbiology
https://doi.org/10.1128/CMR.00058-18.

7. Gardini F, Martuscelli M, Caruso MC, Galgano F, Crudele MA, Favati F, Guerzoni ME,
Suzzi G. 2001. Effects of pH, temperature and NaCl concentration on the growth kinetics,
proteolytic activity and biogenic amine production of Enterococcus faecalisInternational
Journal of Food Microbiology.

8. Morandi S, Brasca M, Alfieri P, Lodi R, Tamburini A. 2005. Influence of pH and
temperature on the growth of Enterococcus faecium and Enterococcus faecalis. Lait
85:181–192.

9. Gaca AO, Lemos JA. 2019. Adaptation to Adversity: the Intermingling of Stress
Tolerance and Pathogenesis in Enterococci. Microbiology and Molecular Biology
Reviews 83.

10. Mchugh CP, Zhang P, Michalek S, Eleazer PD. 2004. pH Required to Kill Enterococcus
faecalis in Vitro.

11. Weckwerth PH, Zapata RO, Vivan RR, Tanomaru-Filho M, Maliza AGA, Duarte MAH.
2013. In Vitro alkaline pH resistance of Enterococcus faecalis. Braz Dent J 24:474–476.

12. dos Santos LDR, Furlan JPR, Gallo IFL, Ramos MS, Savazzi EA, Stehling EG. 2021.
Occurrence of multidrug-resistant Enterococcus faecium isolated from environmental
samples. Lett Appl Microbiol 73:237–246.

13. Ramos S, Silva V, Dapkevicius M de LE, Igrejas G, Poeta P. 2020. Enterococci, from
harmless bacteria to a pathogen. Microorganisms. MDPI AG
https://doi.org/10.3390/microorganisms8081118.

14.   Sen S, Saha T, Bhattacharya S, Mondal N, Ghosh W, Chakraborty R. 2020. Draft Genome
Sequences of Two Boron-Tolerant, Arsenic-Resistant, Gram-Positive Bacterial Strains,
Lysinibacillus sp. OL1 and Enterococcus sp. OL5, Isolated from Boron-Fortified
Cauliflower-Growing Field Soils of Northern West Bengal, India
https://doi.org/10.1128/MRA.

15.   Singhal N, Maurya AK, Mohanty S, Kumar M, Virdi JS. 2019. Evaluation of bile salt
hydrolases, cholesterol-lowering capabilities, and probiotic potential of enterococcus
faecium isolated from rhizosphere. Front Microbiol 10.

16.   Yu Z, Shi D, Liu W, Meng Y, Meng F. 2020. Metabolome responses of Enterococcus
faecium to acid shock and nitrite stress. Biotechnol Bioeng 117:3559–3571.

17.   Flahaut S, Hartke A, Giard J, Benachour A, Boutibonnes P, Auffray Y. 1996. Relationship
between stress response towards bile salts, acid and heat treatment in Enterococcus
faecalis . FEMS Microbiol Lett 138:49–54.

18.   Benachour A, Muller C, Dabrowski-Coton M, le Breton Y, Giard JC, Rincé A, Auffray Y,
Hartke A. 2005. The Enterococcus faecalis SigV protein is an extracytoplasmic function
sigma factor contributing to survival following heat, acid, and ethanol treatments. J
Bacteriol 187:1022–1035.

19.   Salze M, Giard JC, Riboulet-Bisson E, Hain T, Rincé A, Muller C. 2020. Identification of
the general stress stimulon related to colonization in Enterococcus faecalis. Arch
Microbiol 202:233–246.

20.   Evans M, Davies JK, Sundqvist G, Figdor D. 2002. Mechanisms involved in the resistance
of Enterococcus faecalis to calcium hydroxide. Int Endod J 35:221–228.

21.   Ran S, He Z, Liang J. 2013. Survival of Enterococcus faecalis during alkaline stress:
Changes in morphology, ultrastructure, physiochemical properties of the cell wall and
specific gene transcripts. Arch Oral Biol 58:1667–1676.

22.   Ran S, Liu B, Jiang W, Sun Z, Liang J. 2015. Transcriptome analysis of Enterococcus
faecalis in response to alkaline stress. Front Microbiol 6.

23.   Flahaut S, Hartke A, Giard J-C, Auffray Y. 1997. Alkaline Stress Response in
Enterococcus faecalis: Adaptation, Cross-Protection, and Changes in Protein
SynthesisAPPLIED AND ENVIRONMENTAL MICROBIOLOGY.

24.   Harden MM, He A, Creamer K, Clark MW, Hamdallah I, Martinez KA, Kresslein RL,
Bush SP, Slonczewski JL. 2015. Acid-adapted strains of *Escherichia coli* K-12 obtained
by experimental evolution. Appl Environ Microbiol 81:1932–1941.

25.   He A, Penix SR, Basting PJ, Griffith JM, Creamer KE, Camperchioli D, Clark MW,
Gonzales AS, Erazo JSC, George NS, Bhagwat AA, Slonczewski JL. 2017. Acid
evolution of *Escherichia coli* K-12 eliminates amino acid decarboxylases and reregulates
catabolism. Appl Environ Microbiol 83:e00442-17.

26.   Creamer KE, Ditmars FS, Basting PJ, Kunka KS, Hamdallah IN, Bush SP, Scott Z, He A,
Penix SR, Gonzales AS, Eder EK, Camperchioli DW, Berndt A, Clark MW, Rouhier KA,
Slonczewski JL. 2017. Benzoate- and salicylate-tolerant strains of *Escherichia coli* K-12
lose antibiotic resistance during laboratory evolution. Appl Environ Microbiol 83:e02736.

27.   Moore JP, Li H, Engmann ML, Bischof KM, Kunka KS, Harris ME, Tancredi AC,
448   Ditmars FS, Basting PJ, George NS, Bhagwat AA, Slonczewski L. 2019. Inverted
449   regulation of multidrug efflux pumps, acid resistance, and porins in benzoate-evolved
450   Escherichia coli K-12. Appl Environ Microbiol 85:e00966-19.

451   28.   Hamdallah I, Torok N, Bischof KM, Majdalani N, Chadalavada S, Mdluli N, Creamer KE,
452   Clark M, Holdener C, Basting PJ, Gottesman S, Slonczewski JL. 2018. Experimental
453   evolution of *Escherichia coli* K-12 at high pH and RpoS induction. Appl Environ
454   Microbiol 84:e00520.

455   29.   Johnson MD, Bell J, Clarke K, Chandler R, Pathak P, Xia Y, Marshall RL, Weinstock
456   GM, Loman NJ, Winn PJ, Lund PA. 2014. Characterization of mutations in the PAS
457   domain of the EvgS sensor kinase selected by laboratory evolution for acid resistance in
458   Escherichia coli. Mol Microbiol 93:911–927.

459   30.   Nakajo K, Komori R, Ishikawa S, Ueno T, Suzuki Y, Iwami Y, Takahashi N. 2006.
460   Resistance to acidic and alkaline environments in the endodontic pathogen Enterococcus
461   faecalis. Oral Microbiol Immunol 21:283–288.

462   31.   Martin M, Hölscher T, Dragoš A, Cooper VS, Kovács ÁT. 2016. Laboratory evolution of
463   microbial interactions in bacterial biofilms. J Bacteriol. American Society for
464   Microbiology https://doi.org/10.1128/JB.01018-15.

465   32.   Poltak SR, Cooper VS. 2011. Ecological succession in long-term experimentally evolved
466   biofilms produces synergistic communities. ISME Journal 5:369–378.

467   33.   Khan A, Davlieva M, Panesso D, Rincon S, Miller WR, Diaz L, Reyes J, Cruz MR,
468   Pemberton O, Nguyen AH, Siegel SD, Planet PJ, Narechania A, Latorre M, Rios R, Singh
469   K v, Ton-That H, Garsin DA, Tran TT, Shamoo Y, Arias CA. Antimicrobial sensing
470   coupled with cell membrane remodeling mediates antibiotic resistance and virulence in
471   Enterococcus faecalis https://doi.org/10.1073/pnas.1916037116/-/DCSupplemental.

472   34.   Sun Y, Lu H, Zhang X, Wu Q, Bi W, Liu H, Cao J, Zhou T. 2018. Phenotype and
473   genotype alteration during adaptive evolution of Enterococcus faecalis to antimicrobials.
474   Infection, Genetics and Evolution 62:80–85.

475   35.   Guo X, Peisker K, Bäckbro K, Chen Y, Koripella RK, Mandava CS, Sanyal S, Selmer M.
476   2012. Structure and function of FusB: An elongation factor G-binding fusidic acid
477   resistance protein active in ribosomal translocation and recycling. Open Biol 2.

478   36.   Gupta SK, Pfeltz RF, Wilkinson BJ, Gustafson JE. 2022. Transcriptomic and
479   Metabolomic Analysis of a Fusidic Acid-Selected fusA Mutant of Staphylococcus aureus.
480   Antibiotics 11.

481   37.   Durango H, Morales H, Murillo E. 2021. Evaluation of the Virulence Factor Prole of
482   Avian Pathogenic Escherichia Coli in Clinical Isolates of Avian Samples in Caloto,
483   Cauca, Colombia https://doi.org/10.21203/rs.3.rs-1184472/v1.

484   38.   Lieberman MT, van Tyne D, Dzink-Fox JA, Ma EJ, Gilmore MS, Fox JG. 2018. Long-
485   term colonization dynamics of Enterococcus faecalis in implanted devices in research
486   macaques. Appl Environ Microbiol 84.

487   39.   Höring S, Sobotta K, Schneider S, Löffler B, Rödel J. 2019. Dwarfs in disguise: multiple
488   spinal abscesses and spondylodiscitis caused by an Enterococcus faecium small-colony
489   variant. Access Microbiol 1.

490   40.   Proctor RA, von Eiff C, Kahl BC, Becker K, McNamara P, Herrmann M, Peters G. 2006.
491   Small colony variants: A pathogenic form of bacteria that facilitates persistent and
492   recurrent infections. Nat Rev Microbiol https://doi.org/10.1038/nrmicro1384.

493   41.   Deatherage DE, Barrick JE. 2014. Identification of mutations in laboratory-evolved
494   microbes from next-generation sequencing data using *breseq*. Methods in Molecular
495   Biology 1151:165–188.

496   42.   Barrick JE, Colburn G, Deatherage DE, Traverse CC, Strand MD, Borges JJ, Knoester
497   DB, Reba A, Meyer AG. 2014. Identifying structural variation in haploid microbial
498   genomes from short-read resequencing data using *breseq*. BMC Genomics 15:1039.

499   43.   Yang Z, Bielawski JP. 2000. Statistical methods for detecting molecular adaptation. TREE
500   15:496–503.

501   44.   Gupta SK, Pfeltz RF, Wilkinson BJ, Gustafson JE. 2022. Transcriptomic and
502   Metabolomic Analysis of a Fusidic Acid-Selected fusA Mutant of Staphylococcus aureus.
503   Antibiotics 11.

504   45.   Tokmakova IP, Ryabchenko LE, Gerasimova T v., Kameneva S v., Yanenko AS. 2017.
505   Mutations in the fusA Gene Encoding Elongation Factor G in the Coryneform Bacterium
506   Lead to Increased Lysine Production. Appl Biochem Microbiol 53:781–785.

507   46.   Qayyum S, Sharma D, Bisht D, Khan AU. 2019. Identification of factors involved in
508   Enterococcus faecalis biofilm under quercetin stress. Microb Pathog 126:205–211.

509   47.   Maddalo G, Chovanec P, Stenberg-Bruzell F, Nielsen H v., Jensen-Seaman MI, Ilag LL,
510   Kline KA, Daley DO. 2011. A reference map of the membrane proteome of Enterococcus
511   faecalis. Proteomics 11:3935–3941.

512   48.   Bhardwaj P, Islam MZ, Kim C, Nguyen UT, Palmer KL. 2021. ddcP, pstB, and excess D-
513   lactate impact synergism between vancomycin and chlorhexidine against Enterococcus
514   faecium 1,231,410. PLoS One 16.

515   49.   Copin R, Sause WE, Fulmer Y, Balasubramanian D, Dyzenhaus S, Ahmed JM, Kumar K,
516   Lees J, Stachel A, Fisher JC, Drlica K, Phillips M, Weiser JN, Planet PJ, Uhlemann AC,
517   Altman DR, Sebra R, van Bakel H, Lighter J, Torres VJ, Shopsin B. 2019. Sequential
518   evolution of virulence and resistance during clonal spread of community-acquired
519   methicillin-resistant Staphylococcus aureus. Proc Natl Acad Sci U S A 116:1745–1754.

520   50.   Ghim S-Y, Kim CC, Bonner ER, D'elia JN, Grabner GK, Switzer RL. 1999. The
521   Enterococcus faecalis pyr Operon Is Regulated by Autogenous Transcriptional
522   Attenuation at a Single Site in the 5 LeaderJOURNAL OF BACTERIOLOGY.

523   51.   Wang Q, Sun FJ, Liu Y, Xiong LR, Xie LL, Xia PY. 2010. Enhancement of biofilm
524   formation by subinhibitory concentrations of macrolides in icaADBC-positive and -
525   negative clinical isolates of Staphylococcus epidermidis. Antimicrob Agents Chemother
526   54:2707–2711.

52. Belli WA, Marquis RE. 1991. Adaptation of Streptococcus mutans and Enterococcus hirae
to Acid Stress in Continuous CultureAPPLIED AND ENVIRONMENTAL
MICROBIOLOGY.

53. Miller C, Kong J, Tran TT, Arias CA, Saxer G, Shamoo Y. 2013. Adaptation of
Enterococcus faecalis to daptomycin reveals an ordered progression to resistance.
Antimicrob Agents Chemother 57:5373–5383.

54. Lemaire S, Van Bambeke F, Pierard D, Appelbaum PC, Tulkens PM. 2011. Activity of
fusidic acid against extracellular and intracellular staphylococcus aureus: Influence of ph
and comparison with linezolid and clindamycin. Clinical Infectious Diseases 52.

55. Thieme L, Klinger-Strobel M, Hartung A, Stein C, Makarewicz O, Pletz MW. 2018. In
vitro synergism and anti-biofilm activity of ampicillin, gentamicin, ceftaroline and
ceftriaxone against Enterococcus faecalis. Journal of Antimicrobial Chemotherapy
73:1553–1561.

56. Penn CJ, Camberato JJ. 2019. A critical review on soil chemical processes that control
how soil ph affects phosphorus availability to plants. Agriculture (Switzerland) 9.

57. Arai Y, Sparks D. 2007. Phosphate Reaction Dynamics in Soils and Soil Components: A
Multiscale Approach. Advances in Agronomy 94:135–179.

58. Bond PL, Keller J, Blackall LL. 1999. Anaerobic phosphate release from activated sludge
with enhanced biological phosphorus removal. A possible mechanism of intracellular pH
control. Biotechnol Bioeng 63:507–515.

59. Beenken KE, Dunman PM, McAleese F, Macapagal D, Murphy E, Projan SJ, Blevins JS,
Smeltzer MS. 2004. Global gene expression in Staphylococcus aureus biofilms. J
Bacteriol 186:4665–4684.

60. Kakinuma Y. 1998. Inorganic Cation Transport and Energy Transduction in Enterococcus
hirae and Other StreptococciMICROBIOLOGY AND MOLECULAR BIOLOGY
REVIEWS.

553