Role of Lactate in the Regulation of Transcriptional Activity of Breast Cancer-Related Genes and Epithelial-to-Mesenchymal Transition Proteins: A Comparison of MCF7 and MDA-MB-231 Cancer Cell Lines

[1,2,3]Inigo San-Millan*, [2]Janel L. Martinez, [1]Shivaun Lueke Pickard, [1]Hui Yu, [5]Fred R. Hirsch, [6]George A. Brooks and [1]Christopher J. Rivard

[1]Department of Medicine, Division of Medical Oncology, University of Colorado Anschutz Medical Campus, Aurora, CO, USA
[2]Department of Medicine, Division of Endocrinology, Metabolism and Diabetes, University of Colorado Anschutz Medical Campus, Aurora, CO, USA
[3]Department of Human Physiology and Nutrition, University of Colorado, Colorado Springs, CO, USA
[5]Tisch Cancer Institute, Center for Thoracic Oncology, Mount Sinai Health System, New York, NY, USA
[6]Department of Integrative Biology, University of California, Berkeley, Berkeley, CA, USA
*Correspondence: Inigo.sanmillan@cuanschutz.edu

Simple Summary: This study shows how endogenous lactate is a major regulator of key genes involved in two main breast cancer cell types: estrogen receptor positive (ER[+]) and triple negative breast cancer cells (TPBC). Lactate regulates both gene and protein expression in these cells. Furthermore, lactate is also an important player in the regulation of epithelial-to-mesenchymal transition (EMT), a process involved in metastasis. Targeting lactate production and exchange within and among cancer cells should offer novel therapeutics opportunities.

Abstract:
The Warburg Effect is characterized by accelerated glycolytic metabolism and lactate production and is a hallmark of cancer cells. Recently, we have demonstrated the role of endogenous, glucose-derived lactate as an oncometabolite which regulates gene expression in the estrogen receptor positive (ER+) MCF7 cell line cultivated in glucose media (San-Millan, Julian et al. 2019). Presently, with the addition of a triple negative breast cancer (TNBC) cell line, MDA-MB-231, we further confirm the effect of lactate on gene expression patterns, but extend results to include lactate effects on protein expression. As well, we report effects of lactate on the expression of E-cadherin and vimentin, proteins associated with epithelial-to-mesenchymal transition (EMT). Endogenous lactate regulates the expression of multiple genes involved in carcinogenesis. In MCF7 cells, lactate increased the expression of *EGFR, VEGF HIF-1a, KRAS, MIF, mTOR, PIK3CA, TP53, and CDK4* (The *Ital* is used for genes) as well as decreased the expression of *ATM, BRCA1, BRCA2, E2F1, MET, MYC,* and *RAF* mainly at 48h of exposure. On the other hand, in the MDA-MB-

231 cell line, lactate increased the expression of *PIK3CA, VEGF, EGFR, mTOR, HIF-1α, ATM, E2F1, TP53* and decreased the expression of *BRCA1, BRCA2, CDK4, CDK6, MET, MIF, MYC,* and *RAF* after 48h of exposure. Protein expression of representative genes corroborated mRNA expressions. Finally, lactate decreased E-cadherin protein expression in MCF7 cells and increased vimentin expression in MDA-MB-231 cells.

In this study, we demonstrate that endogenous lactate produced under aerobic conditions (Warburg Effect) is capable of eliciting important regulation of gene and protein expression in both ER+ and TNBC cell lines. The regulation of multiple genes by lactate is widespread and involves those involved in carcinogenesis including DNA repair, cell growth, proliferation, angiogenesis, and metastasis. Furthermore, both cell lines demonstrated changes in the expression of EMT biomarkers representing a shift to a more mesenchymal phenotype with exposure to endogenous lactate.

Keywords: lactate; glycolysis, carcinogenesis, gene expression; protein expression; epithelial-mesenchymal transition

1. Introduction

Described by Otto Warburg a century ago (Warburg and Minami 1923) accelerated glucose consumption and lactate production under fully aerobic conditions is ubiquitous in solid tumors. With discovery of DNA by Watson and Crick in 1953 (Watson and Crick 1953), the landscape of cancer research completely switched from emphasis on metabolism to genetics. At present, while it is recognized that cancer is characterized by mutations of multiple genes involved in cell growth, proliferation, angiogenesis or DNA repair (Vogelstein, Papadopoulos et al. 2013), regrettably despite decades of efforts, the cure of through gene science has not been realized. Hence, others and we, have decided to re-open the gates of cancer metabolism. This reopening including studies of cell metabolism has led to a renaissance in cancer biology.

Observations of accelerated glucose uptake and lactate secretion under fully aerobic conditions led Warburg to posit that cancer could be attributed to an injury to the cellular respiratory system (mitochondria) (Warburg 1925), although that idea could not be quite correct (Hussien, Rihn et

al. 2013). Still, for many decades lactate was considered a "waste" product resulting from lack of oxygen. However, pioneering work and discoveries by George Brooks since the 1980's have demonstrated that lactate is the preferred fuel for most cells, the main gluconeogenic precursor in the body, and a signaling molecule (a "lactormone," "exerkine and "myokine") capable of inducing gene expression (Brooks 1986, Brooks 2018, Brooks, Osmond et al. 2022). In concert, Zhang et al. have reported recently, that the key role of lactate in histone acetylation ("lactylation") is the acetylation of lysine histone residues, acting as an epigenetic modulator and increasing gene transcription from exposed chromatin in both human and mouse cells (Di Zhang 2019). More recently, Jiang et al. have shown that lactate modulates cellular metabolism through histone lactylation-mediated genes in non-small cell lung cancer (Jiang, Huang et al. 2021). As well, Verma and colleagues have also recently demonstrated that tumor-derived lactate can elicit epigenomic reprograming in cancer-associated fibroblasts from pancreatic ductal adenocarcinoma (Bhagat, Von Ahrens et al. 2019). Moreover, W. Liu and colleagues have recently demonstrated that lactate regulates cell cycle by remodeling the anaphase promoting complex (Liu, Wang et al. 2023).

In 2017, we developed the "lactagenesis hypothesis" (San-Millan and Brooks 2017) by which we posited that accelerated lactate production in tumors was not for bioenergetics (Brooks 2009, Brooks 2018), but rather owing to its powerful signaling properties (Brooks 2018, Brooks 2020), the role of lactate in tumors was to stimulate carcinogenesis. Posteriorly, in 2019 we demonstrated that lactate functions as an "oncometabolite" and regulates the expression of major genes related to carcinogenesis in MCF7 breast cancer cells (San-Millan, Julian et al. 2019). Recognition of the importance of lactate as a hallmark of cancer led several groups to block lactate transport and exchange among and cancer cells and within tumors as a therapeutic approach (Sonveaux, Vegran et al. 2008, Sonveaux, Copetti et al. 2012, Draoui, Schicke et al. 2013, Doherty, Yang et al. 2014, Draoui, Schicke et al. 2014, Mishra and Banerjee 2019, Kuo, Huang et al. 2020).

The first aim of the present study was to investigate the role of endogenous (glucose-derived) lactate as an oncometabolite on important oncogene expression in two representative breast

cancer cell lines. Herein, we sought to extend our findings by studying the representative breast

cancer cells lines MCF7, an estrogen-receptor positive (ER+) cell line, and MDA-MB-231, a triple-

negative cell line (TNBC). To further confirm the importance of lactate in carcinogenesis, we also

assessed protein expression for some of the important genes in both breast cancer cells.

The second aim of our study was to assess the potential role of lactate in the regulation of the

epithelial-to-mesenchymal transition (EMT). EMT is a key process in cancer progression

characterized by the loss of epithelial towards mesenchymal properties, thus allowing cells to

acquire enhanced capacity for motility, migration, invasiveness, and resistance to apoptosis. This

increased migratory capacity from a primary tumor site to a secondary, tumor-free site, involves

EMT as an important hallmark of metastasis (Thiery 2002, Mittal 2018). In general, MCF7 cells

are characterized by possessing an epithelial phenotype, while MDA-MB-231 a mesenchymal one

(Wang, Xu et al. 2019).

EMT is a complex multifactorial process with two major proteins representative of the process:

E-cadherin and vimentin. E-cadherin is a transmembrane protein which facilitates intercellular

(cell-to-cell) adhesions as well as morphogenesis, limiting cell motility (Chao, Shepard et al. 2010).

The decrease of E-cadherin expression disrupts cellular adhesion as well as elicits loss of cellular

architecture, facilitating motility and metastasis (Wells, Yates et al. 2008, Devlin and Verschuren

2018). Furthermore, autocrine signaling reinforces the mesenchymal phenotype and stimulates

endothelial growth factor receptor (EGFR), downregulating E-cadherin (Kopfstein and Christofori

2006).

Vimentin is a structural protein (type III intermediate filament protein) expressed in

mesenchymal cells. Vimentin is attached to the nucleus, endoplasmic reticulum, and

mitochondria, anchoring organelles in cytosol (Katsumoto, Mitsushima et al. 1990, Cheng, Shen

et al. 2016). Vimentin induces cell migration by inhibiting focal-adhesion proteins (Gregor,

Osmanagic-Myers et al. 2014). Moreover, it has been shown that vimentin is a key player in

metastasis by regulating cell migration and invasion as well as tumor angiogenesis. (Chen, Fang

et al. 2021). Furthermore, it has been shown that TNBC cells possess significantly higher amount

of vimentin than ER+ cancer cells (Su, Hopfinger et al. 2018).

2. Materials and Methods

2.1 Cancer cell lines

MCF7 and MDA-MB-231, both adenocarcinoma breast cancer cell lines were obtained from the ATCC (Manassas, VA) and stored in liquid nitrogen until use. Cells were grown on tissue culture treated dishes (Genessee Scientific, San Diego, CA) in DMEM 1x media with 1g/L glucose (10-014-CV, Corning, Manassas, VA), 10% FBS, and 1X pen/strep. In experiments requiring high glucose media, cells were grown with DMEM 1X media with either 1 or 4.5 g/L glucose (10-013-CV, Corning). In these experiments, the normal 10% FBS was changed to 10% Nu-Serum (#51000, Corning Life Sciences, Glendale, AZ) to reduce the amount of lactate in the initial medium. Media was also amended with 10 and 20 mM added L-lactate both for 6h and 48h exposures. All cell cultures were grown at 37°C in a laboratory incubator with 5% CO_2 atmosphere (Forma Isotemp 3530, ThermoFisher, Marietta, OH). Trypsin (0.25% trypsin, 25-053-CI, Corning) was utilized during culture expansion.

2.2. QPCR analysis

Cell lines were trypsinized and resuspended in media, followed by low-speed centrifugation to a soft pellet. The cell pellet was then harvested for RNA using the RNeasy Plus Mini Kit (#74134, Qiagen, Hilden, Germany) as per the manufacturer's protocol. The concentration of purified RNA was determined using a Nanodrop 2000 spectrophotometer (Thermo Scientific, Waltham, MA). cDNA was prepared from 500ng RNA per reaction using the iScript cDNA Synthesis Kit (#1708891, Bio-Rad, Hercules, CA). QPCR was performed using probe-based primer sets and master mix (Prime Time, Integrated DNA Technologies, Coralville, IA) on a CFX96 real time platform (Bio-Rad). Data was evaluated using Maestro software (Bio-Rad) and compared to a standard dilution curve from cell line cDNA. The appropriate housekeeping genes to use for data normalization were determined employing the Human Reference Gene Panel (Bio-Rad). Data was normalized with TBP and HPRT1 analysis.

2.3. Cell Protein Isolation and Western Blot Protocol

Western blots for cell line protein expression were performed by harvesting cell cultures

from growth plates by removal of spent media, washing plates 3X with cold 1X PBS, followed by

addition of cell lysis buffer (#9803, Cell Signaling Technologies, Danvers, MA) amended with

AEBSF (1 uM, SBR00015, Sigma) rather than PMSF. Cells were scraped and transferred to a

microcentrifuge tube and frozen at -80°C until analyzed. Samples were thawed, sonicated for 5

seconds at 50% power with a pencil type sonicator (model 450, Branson, Danbury, CT). Lysate

samples were centrifuged at maximum speed in a microfuge at 4°C for 15 minutes and the soluble

protein supernatant removed to a new microcentrifuge tube. Protein concentrations of samples

were determined using the BCA protein assay (#23225, Pierce) as per the manufacturers

protocol. Protein specimens were aliquoted in 50 uL amounts and frozen at -80°C until analysis.

Western blots were loaded at 2.5 to 50 ug total protein per lane in 4-20% gradient pre-cast

polyacrylamide gels (#3450033, Bio-Rad). Gels were also loaded with the Dual pre-stained

protein standards (#1610374, Bio-Rad) to evaluate protein separation during electrophoresis and

transfer to membranes. Gels were transferred to PVDF membranes (#1620177, Bio-Rad) for 2

hours at 250 mAmps followed by blocking in 5% w/v milk in TTBS buffer for 1 hour. Blocked

membranes were probed with primary antibodies overnight on a laboratory rocker at 4°C.

Membranes were washed 5X with cold TTBS and incubated with anti-Rabbit or anti-mouse

secondary antibodies conjugated to HRP (#7074 and #7076, Cell Signaling). After washing,

membranes were visualized with ECL reagent (Clarity, #1705060, Bio-Rad) on radiographic film

in the dark room. Western blot films were imaged with an Epson 310 scanner (Los Alamitos, CA).

Primary antibodies were from Cell Signaling including ATM (#2873), beta Actin (#4970), CDK4

(#12790), E cadherin (#14472), EGFR (#4267), HIF1 alpha (#35169), LDHa (#3582), MET (#8198),

MIF (#87501), mTOR (#2983), PIK3CA (#4249), RAS (#67648), and vimentin (#5741), from R&D

Systems (Minneapolis, MN) including LDHb (MAB9205), and from Abcam (Cambridge, UK)

including Citrate Synthase (AB129095).

2.4. Analysis of lactate and glucose concentrations in spent growth media

Lactate in spent growth media was determined employing the L-lactate Assay Kit I from Eton

Biosciences (#120001400A, San Diego, CA). Sample lactate analysis was performed in the 96-will

plate format with comparison to a standard curve in the range of 30 – 3000uM. Assay plates were

measured for the formazan product at 490 nM using a Synergy 2 plate reader (BioTek, Winooski,

VT). Net lactate production was determined by subtracting lactate in the initial media from

cancer cell's lactate production at each time point during the experiments.

2.5. Seahorse XFe real-time ATP rate assay

For Seahorse experiments, cell lines were maintained in either 1 g/L or 4.5 g/L glucose

growth medium for at least 72 hours prior to analysis. Cells were plated at the concentration of

about 14,000 cells/well in XFe 96 microplate prior to washing and changing to assay-specific

medium sans phenol red and sodium bicarbonate. Brightfield visualization was performed prior

to the run utilizing a Cytation Plate Imaging Scanner (BioTek). The real-time measurement of

oxygen consumption rate (OCR) and glycolytic extra-cellular acidification rate (ECAR) was

performed using the Seahorse XFe 96 Extracellular Flux Analyzer (Agilent Technologies, Santa

Clara, CA) with the Seahorse XF real-time ATP rate assay Kit (#103592-100, Agilent Technologies)

according to the manufacturer's protocol. At the conclusion of the assay run, cells are counted

via fluorescence in the Cytation reader following exposure to Hoechest 33342 cell-permeable

DNA stain (2 µM final, #62249, Thermo Scientific). OCR values were normalized to cell number.

2.6. Statistical analyses

Independent sample t-tests and analyses of variance (ANOVA) were utilized for group-mean

comparisons via GraphPad Prism software (version 9.2.1, GraphPad, San Diego, CA).

3. Results

3.1. Glucose-derived endogenous lactate (Warburg Effect)

Because lactate was not present in treatment media, the lactate present in the spent media was

endogenously-derived in both cell lines from glycolysis (glucose-derived lactate). Glycolysis

progressed readily in both cell lines with total and net lactate releases significantly higher after

48h than 6h of incubation (p<0.001) (Fig-1A and 1B, respectively). Notably, lactate release from

MDA-MB-231 is higher than from MCF7 cells (69%, p<0.01 at 6 hours, 72%, p<0.0001 at 48 hours)

(Fig-1A and Fig-1B). Further, as discussed below, exogenous lactate additions of 10mM and

20mM to incubation media did not affect net lactate production (Fig-1B).

The higher glycolytic phenotype of MDA-MB-231 was confirmed via our Seahorse XFe Real-Time

ATP Rate analysis. Overall, the total basal ATP production rates of both low-glucose and high-

glucose adapted MDA-MB-231 cells were significantly higher than for MCF7 cells (Fig-2A). In

MDA-MB-231 cells, more than half of ATP was derived from glycolysis. Once again, by that

measure, MDA-MB-231 cells displayed a more glycolytic phenotype than MCF7 cells (Fig-2B).

Consistent with Seahorse analyses, expression showed increased levels of the "lactagenic"

enzyme isoform, LDHA, was higher in MDA-MB-231 cells than in MCF7 cells (Fig-3). Conversely,

in terms of oxidative phenotype, MCF7 cells showed increased protein expression of the

mitochondrial marker citrate synthase (CS) as compared to MDA-MB-231 cells (Fig-3).

226
227   **Figure 1.** Comparison of lactate production in MCF7 and MDA-MB-231 cell lines. Lactate concentration in the spent
228   media at 6 and 48 hours, normalized to total cell protein concentration (**A**). Comparison of net lactate production in
229   spent media in response to 10 mM and 20 mM added exogenous lactate in 4.5 g/L glucose medium with increasing
230   incubation time (**B**).

231
232
233

234
235
236 **Figure 2.** Total basal ATP production rate (**A**) and basal ATP production distribution (**B**) in MCF7 and MDA-MB-231
237 breast cancer cell lines. Cell lines were maintained in either 1 g/L or 4.5 g/L glucose growth medium for 72 hours
238 prior to analysis.
239
240

241
242 **Figure 3.** Differences in citrate synthase (CS) and lactate dehydrogenase A (LDHA) protein expression between MCF7
243 and MDA-MB-231 cells, representing the higher glycolytic nature of MDA-MB-231 compared to the higher OXPHOS
244 capacity of MCF7.
245
246

3.2. Gene expression regulation by lactate

Glycolysis in the cancer cell lines we studied readily resulted in lactate excretion and

accumulation (Fig-1). Glucose-stimulated lactate regulated the expression of multiple

representative cancer-related genes in both MCF7 and MDA-MB-231 cell lines (Fig-4). However,

this regulation was heterogeneous and varied between cell lines. MCF7 cell lines showed

increased expressions of *EGFR, KRAS, PIK3CA, mTOR, VEGF, HIF-1α, TP53, MIF,* and as well,

decreased expressions of *BRCA1, BRCA2, ATM, E2F1, CDK6, MET, MYC, and RAF* when exposed

to endogenously produced lactate, mainly at 48h (Fig-1, Fig-4a).

In response to glucose-stimulated lactate, MDA-MB-231 cells showed increased expressions of

VEGF, HIF-1α, PIK3CA, mTOR E2F1, EGFR and TP53 (Fig-4b). On the other hand, lactate exposure

decreased the expression of *BRCA1, BRCA2, CDK6, MET, MIF, MYC, RA*F, mainly at 48h. However,

CDK4 expression increased at 6h, but decreased at 48h (Fig-4b).

The addition of 10mM and 20mM exogenous lactate to glucose-containing media did not

increase gene expression compared to endogenously-derived lactate and is further explained in

Discussion. Noteworthy, lactate increased the expression monocarboxylate transporters (MCTs)

(Fig-4b). Genes encoding for *MCT1*, increased significantly in MCT7 cells, while *MCT4* expression

increased in both cell lines in response to increased lactate exposure.

265
Figure 4. Gene expression after 6 and 48h of endogenous (glucose-derived lactate) and exogenous lactate exposure
in both MCF7 (**A**) and MDA-MB-231 cell lines (**B**).

3.3. Protein expression regulation by lactate

Next, we measured protein expression of several representative genes that were altered in

expression by lactate. Western blots confirmed that the protein expression patterns of

representative genes mirror the expression patterns of the genes identified as having changed

from mRNA analyses (Fig-5). In MCF7 cells, 48h lactate exposure increased protein expression of

EGFR, PIK3CA, RAS, HIF-1α, mTOR and CDK4 and decreased the expression of *ATM* (Fig-5A),

confirming data from mRNA expression analysis. In MDA-MB-231 cells, 48h lactate exposure

increased protein expression of *PIK3CA, mTOR, HIF-1α* and *ATM*, did not affect *RAS* expression,

and decreased the expression of *MET* and *CDK4* (which increased at 6h) (Fig-5B), reflecting mRNA

expressions. Like in the case of gene expression, the addition of 10mM and 20mM exogenous

lactate did not further increase protein expression compared to endogenous lactate that is

further addressed in Discussion.

281
Figure 5. Gene and protein expression of representative cancer genes in MCF7 (**A**) and MDA-MB-231 cell lines (**B**)
following exposure to endogenous lactate (glucose-derived lactate) plus exogenous lactate at 6 and 48 hours as
determined via qPCR analysis and Western blot, respectively.

3.4. Possible role of Lactate in the regulation of epithelial-to-mesenchymal transition (EMT)

Our results can be interpreted to reveal an important role of lactate as a regulator in the

EMT process in both cancer cell lines. E-cadherin protein expression was decreased by lactate in

MCF-7 cells (p=0.058), (Fig-6). Moreover, while vimentin was significantly increased by lactate

exposure in MDA-231 cells (p<0.01), we could not detect vimentin in MCF-7 cells (Fig-6).

Like in the case of gene expression experiments, the addition of 10mM and 20mM exogenous

lactate did not affect protein expression compared to endogenous lactate production in either

cell line (see Discussion).

296
297 **Figure 6.** Protein expression of key EMT-related proteins, E-cadherin and vimentin, in MCF7 and MDA-MB-231
298 breast cancer cell lines in response to 0, 10 mM, and 20 mM exogenous lactate additions at 6 and 48 hours.
299
300
4. Discussion

The role of lactate in regulating the expression of multiple genes is not new. Brooks and

colleagues already demonstrated that lactate was able to regulate the expression of 630 genes

in multinucleated myotubes and striated fibers-L6 cells (Hashimoto, Hussien et al. 2007). Further,

Zhang et al. have demonstrated that lactate-derived lactylation of histone-associated lysine

residues serves as an epigenetic modification that directly stimulates gene transcription by

exposing chromatin and identified 28 lactylation sites on core histones in human and mouse cells

(Di Zhang 2019). As well, Jiang et al. have also recently demonstrated that lactate regulates

cellular metabolism in non-small lung cancer cells through histone lactylation (Jiang, Huang et al.

2021). Moreover, we recently demonstrated the role of lactate as an oncometabolite, regulating

the expression of important cancer-related genes in MCF7 cells (San-Millan, Julian et al. 2019). In

the present study, we show that lactate is a major regulator of multiple genes and dependent

proteins involved in both ER+ and TNBC cells. Furthermore, we show that lactate could play an

important role in the epithelial-to-mesenchymal transition and hence, *vice versa* (mesenchymal-

to-epithelial transition) by regulating the expression of important proteins, specifically E-

cadherin and vimentin, involved in these transformations.

As depicted in Figure 1A, lactate is readily produced by both cancer cell lines studied, with the

largest lactate accumulation after 48h of exposure to glucose media. Therefore, this lactate is

endogenous lactate, or glucose-derived lactate, as a result of the Warburg Effect (Warburg and

Minami 1923, Warburg 1925) We also showed this phenomenon in our recent study with MCF7

cells (San-Millan, Julian et al. 2019). In the study herein, we observed that MDA-MD-231 cells

produce significantly higher quantities of lactate compared to MCF7 cells, with lactate being

produced nearly twice as fast as in MCF7 cells per total cellular protein after 48h (p<0.001) (Fig-

1). This finding is in line with the higher glycolytic nature of MDA-MD-231 cells reported by others

(Lunetti, Di Giacomo et al. 2019, Reda, Refaat et al. 2019, Ocana, Martinez-Poveda et al. 2020),

which we confirmed via Seahorse cellular respiration analysis (Fig-2). *In vivo*, the glycolytic

phenotype is associated with higher cancer aggressiveness and patient mortality (Jang, Han et al.

2012, Vlassenko, McConathy et al. 2015, Wang, Jiang et al. 2020, Zhang, Zheng et al. 2020).

Similar to what is observed *in vivo*, we found net lactate production at 48h accumulated to 20-

40mM in both cell lines (Fig-1B), which is consistent with lactate concentrations found in a variety

of tumor microenvironments (Brizel, Schroeder et al. 2001).

In our experiments, we added two different lactate concentrations (10mM and 20mM) to our

experimental medium in order to assess whether the addition of exogenous lactate would

amplify the expression of multiple genes in a dose-dependent manner vs endogenously-derived

lactate. Consequently, we concluded that endogenously produced lactate achieved the

maximum lactate-stimulated responses to gen and protein expression. The maximum

endogenous lactate simulated-response was also true for EMT proteins, vimentin and E-cadherin.

Our data shows that endogenous lactate is sufficient to alter genetic and proteomic responses in

both cancer cell lines. A main reason for a lack of physiological effect by the additional exogenous

lactate in the treatment media could reside in lactate transport kinetics. Lactate is transported

into and out of cells by monocarboxylate transporters (MCTs), which are bidirectional,

symporters (Brown and Brooks 1994, Halestrap 2012). Lactate is transported through cell

membranes following proton and monocarboxylate concentration gradients (Roth and Brooks

1990, Roth and Brooks 1990). Hence, metabolic activity and physiological processes such as

circulatory transport determine the gradients influencing lactate monocarboxylate (mainly

lactate, the physiological Lactate/Pyruvate ranging from 10 to 500, (Bergman, Wolfel et al. 1999,

Henderson, Horning et al. 2004) release from lactate net producers (driver cells) and consumers

(recipient cells) (Brooks, Osmond et al. 2023). Thus, cellular lactate efflux as a result of lactate

production from glucose (i.e., Warburg Effect) could prevent net influx of exogenous lactate

which could probably be the case of our experiments (Figs 4 and 5). Related, is that we observed

a significant increase in MCT4 expression in both cell lines at 48h (Fig-4), suggesting a feed-

forward stimulus by lactate on MCT4 transporter expression in order to export lactate to the

extracellular space where it then becomes a main player in acidosis of the tumor

microenvironment (TME) and regulator of multiple TME functions (de la Cruz-López, Castro-

Muñoz et al. 2019, Pérez-Tomás and Pérez-Guillén 2020).

Solid tumors accumulate multiple genetic mutations and in the case of breast cancers, the

average number of nonsynonymous mutations is 33, where about 95% of them are single-base

substitutions (Vogelstein, Papadopoulos et al. 2013). Within the number of genes mutated,

"driver genes" are the key ones conferring cancer cells a selective growth advantage (Haber and

Settleman 2007, Vogelstein, Papadopoulos et al. 2013). Within these driver genes, it is thought

that breast tumors require mutations in three to six driver genes (Vogelstein, Papadopoulos et

al. 2013). In both cells studied herein, lactate regulated the expression of multiple important

genes involved in all major processes of carcinogenesis including cell proliferation, growth,

angiogenesis, metastasis, and tumor suppression (Fig-4). Most of the genes prevalent in results

of our study are considered "driver genes" which are the key genes conferring selective growth

advantage to the cancer, as opposed to untransformed cells (Bozic, Antal et al. 2010),.

367 In MCF7 cells, we observed that lactate elicits significant increases in the expression of genes

368 involved in cell signaling, cell growth, and proliferation like *EGFR, PIK3CA, VEGF, mTOR, KRAS,*

369 *MIF*, and *CDK4*. Lactate also elicited increased expression in genes involved in angiogenesis like

370 *HIF1α* and *VEGF*. It is known that lactate stimulates angiogenesis through regulation of the

371 expression of vascular endothelial growth factor (VEGF), a signaling protein that has been known

372 to stimulate the formation and growth of blood vessels (Beckert, Farrahi et al. 2006, Kumar, Viji

373 et al. 2007, Dhup, Dadhich et al. 2012, Polet and Feron 2013). Furthermore, Hypoxia-inducible

374 factor 1 (HIF-1) is a master regulator of angiogenesis including the transcription of VEGF

375 (Forsythe, Jiang et al. 1996, Hirota and Semenza 2006). According to Semenza, HIF-1 is

376 fundamental for activating the transcription of genes encoding glucose transporters and

377 glycolytic enzymes which elicit metabolic reprogramming from OXPHOS to glycolysis and lactate

378 formation (Semenza 2010). In our recent study with MCF7 cells (San-Millan, Julian et al. 2019)

379 and in the study herein, we show that lactate activates the transcription of *HIF-1α* (Figs-4-5).

380 Hence, there appears to be a reciprocal relationship between HIF-1 and lactate and HIF

381 expression.

382 Importantly, we found that lactate elicits overexpression of a key tumor suppressor gene, *TP53*

383 in both cancer cell lines. Although P53 is typically considered a "tumor suppressor" protein that

384 binds to and repairs damaged DNA or signals for apoptosis in irreparable cells, missense

385 mutations in *TP53* elicit mutant p53 proteins with potential gain-of-function properties that

386 stimulate tumor cell proliferation, migration, invasion, survival, chemoresistance, cancer

387 metabolism, and/or tissue architecture disruption (Lee, Park et al. 2019). Consequently,

388 overexpression of P53 may be associated with increased mortality and poor prognosis (Isola,

389 Visakorpi et al. 1992, Friedrichs, Gluba et al. 1993, Wang, Yu et al. 2016). Furthermore, we

390 observed that lactate decreased the expression of two other main tumor suppressor factors,

391 BRCA1 and BRCA2, as well as the expression of ATM, a gene involved in DNA repair (Lavin, Delia

392 et al. 2006).

393 In MDA-MB231 cells, lactate increased the expression of genes involved in cell signaling, cell

394 growth, and proliferation including *VEGF, PI3CKA, HIF-1α, EGFR, mTOR* and *CDK4*, although the

latter only at 6h. As in the case with MCF7 cells, we also observed an increase in the expression
of tumor suppressor factor, *TP53* which is mutated in ~80% of TNBCs (Li, Zhang et al. 2019). The
well-known *TP53* missense mutation of MDA-MB-231 specifically leads to increased cancer cell
survival, migration, and invasiveness. Additionally, like in the case of MCF7, lactate decreased
gene expression for *BRCA1* and *BRCA2.*

To corroborate the findings on gene expression obtained through qPCR analysis, we performed
Western blots on a representative number of dependent proteins in both cell lines showing
lactate's involvement in protein expression at 6 and 48h of lactate exposure (Fig-5). Overall, it is
interesting to observe the differences in gene and protein expression produced by lactate
between the two unique breast cancer cell lines. While some key genes and proteins such as
PIK3CA, VEGF, mTOR and *EGFR* were upregulated by endogenous lactate in , there were
differences between cell lines in the expression of other genes such as *ATM, CDK4, and E2F1.*
This could be expected given the heterogeneous and different genetic characteristics of both cell
lines.

We believe it is noteworthy to observe the role of lactate in regulating EGFR in MCF7 cells. EGFR
is activated via the binding of its specific ligand at the cell surface, epithelial growth factor, and
elicits receptor dimerization and phosphorylation, leading to complex signal transduction
necessary for cellular proliferation, differentiation, growth, and migration as well as inhibition of
apoptosis within the cell. Therapeutically, EGFR is a frontline target in EGFR-subfamily-positive
tumors through a variety of approaches including monoclonal antibodies as well as tyrosine
kinase inhibitors (TKI's) (Huang, Armstrong et al. 2004, Lee, Hahn et al. 2014). These direct
inhibitors of EGFR bind to and block the ATP-binding pocket of the receptor, resulting in inhibition
of downstream signal transduction. EGFR is a "mother" kinase for two main signaling pathways
that are often dysregulated in cancer, which are the RAS/RAF/MEK/ERK and PI3K/Akt/mTOR
pathways, which are also regulated by lactate in our study herein.

In our study, we showed a significant increase in EGFR protein expression in MCF7 cells in
response to endogenous lactate production. This is important in ER+ cancer cells as a decrease
in drug response over time and eventual resistance to tamoxifen is a major problem in this largely

effective cancer therapy. In these type of ER+ cancers, EGFR overexpression can overrule the

signaling inhibition of 17β-estradiol (E_2) by tamoxifen, increasing carcinogenic signaling through

the PI3K/Akt pathway (Moerkens, Zhang et al. 2014); therefore, anti-EGFR therapy in conjunction

with tamoxifen could be preferentially employed in ER+ cancers overexpressing EGFR(Jeong, Bae

et al. 2019).

Finally, regarding EMT, we observed that lactate exposure elicits a decrease in E-cadherin in

MCF7 cells that accompanied by a significant increase in EGFR elicited by lactate. That effect

could explain some synergies in the decrease of expression of E-cadherin contributing to the loss

of epithelial characteristics and positioning lactate as a possible important regulator in this

process. On the other hand, MD-MBA-131 cells possess significantly higher amounts of vimentin

than MCF7 cells. Lactate exposure significantly increased the protein expression of vimentin at

48h, thus, facilitating a shift to a more mesenchymal, and therefore, metastatic phenotype.

Hence, lactate appears to be an important regulator of vimentin and E-cadherin expression in

MD-MBA-131 and MCF7 cells, respectively. Therefore, lactate may be an important regulator of

EMT in tumor cell lines.

5. Conclusions

Endogenous lactate as a result of the Warburg Effect is capable of eliciting important changes of

both gene and protein expression in both MCF7 (ER+) and MDA-MB-231 (TNBC) cell lines. The

regulation of multiple genes by lactate is widespread and involves main genes involved in the

carcinogenesis process including DNA repair, cell growth, proliferation, apoptosis evasion,

angiogenesis, and metastasis. Furthermore, a key step in carcinogenesis and metastasis is the

epithelial-to-mesenchymal transition (EMT) and herein we show that lactate probably plays an

important role in the regulation of EMT in breast cancer.

Blocking cellular lactate production as well as the exchange between cancer cells and surrounding

stroma cells, affecting the tumor microenvironment, should be a priority in cancer research.

Limitations and Challenges

We believe that due to the potent epigenetic-effector properties of lactate on multiple genes and the major metabolic implications of cancer studies there are a variety of elements that need to be better explored and understood. As an example and comparable to many other studies, our previous study employed glycolytic enzyme, GAPDH, as a traditional housekeeping gene (San-Millan, Julian et al. 2019) . In our current study herein, we used other traditional housekeeping genes, TBP and HPRT1, as these genes were determined to be less affected by experimental conditions via a reference gene panel of 15 validated housekeeping genes. Furthermore, the actual methods of glucose media exposure are not physiological as they don't obey the exact metabolic kinetics of glucose uptake and oxidation by normal cells. Furthermore, it is also noteworthy that the accelerated glycolysis and metabolic reprogramming of cancer cells obey different kinetics respect to normal cells. A true physiological glucose delivery to cancer cells mimicking glucose kinetics in cancer should be important to accurately understanding cancer metabolism and bioenergetics as well as lactate production for carcinogenesis.

Author Contributions: Conceptualization, I.S., C.R. and J.M..; methodology, C.R., J.M. and S.L.P.; formal analysis, C.R., J.M, S.L.P, H.Y., G.S. and I.S.; investigation, I.S., C.R., J.M and S.L.P.; resources, C.R., F.H .and I.S..; data curation, C.R., J.M., S.L.P and I.S.; writing—original draft preparation, I.S., C.R. and JM; writing—review and editing, I.S., C.R., J.M. and G.B.; funding acquisition, I.S. All authors have read and agreed to the published version of the manuscript.

Funding: Funding for this study came from IS-M Cellular Metabolism Laboratory funds, The Tadej Pogačar Cancer Research Foundation and supplementary support from NIH 1 RO1 AG059715-01 to GB.

Data Availability Statement: All data from this study are available in the manuscript text and figures. Further information can be obtained from the corresponding author upon reasonable request.

Acknowledgements: The authors thank Christina Coughlan, Adriana Solano and Raleigh Jonscher for meaningful discussions of the research results. Use of the Seahorse XFe96 analyzer was

provided by Matthew Jackman in the NORC Molecular and Cellular Analytical Core at the

University of Colorado Cancer Center.

Conflicts of Interest: The authors declare no conflict of interest

References

1.   Beckert, S., F. Farrahi, R. S. Aslam, H. Scheuenstuhl, A. Konigsrainer, M. Z. Hussain and T. K. Hunt
(2006). "Lactate stimulates endothelial cell migration." Wound Repair Regen **14**(3): 321-324.
2.   Bergman, B. C., E. E. Wolfel, G. E. Butterfield, G. D. Lopaschuk, G. A. Casazza, M. A. Horning and
G. A. Brooks (1999). "Active muscle and whole body lactate kinetics after endurance training in
men." Journal of applied physiology **87**(5): 1684-1696.
3.   Bhagat, T. D., D. Von Ahrens, M. Dawlaty, Y. Zou, J. Baddour, A. Achreja, H. Zhao, L. Yang, B.
Patel, C. Kwak, G. S. Choudhary, S. Gordon-Mitchell, S. Aluri, S. Bhattacharyya, S. Sahu, P. Bhagat,
Y. Yu, M. Bartenstein, O. Giricz, M. Suzuki, D. Sohal, S. Gupta, P. A. Guerrero, S. Batra, M. Goggins,
U. Steidl, J. Greally, B. Agarwal, K. Pradhan, D. Banerjee, D. Nagrath, A. Maitra and A. Verma
(2019). "Lactate-mediated epigenetic reprogramming regulates formation of human pancreatic
cancer-associated fibroblasts." Elife **8**.
4.   Bozic, I., T. Antal, H. Ohtsuki, H. Carter, D. Kim, S. Chen, R. Karchin, K. W. Kinzler, B. Vogelstein
and M. A. Nowak (2010). "Accumulation of driver and passenger mutations during tumor
progression." Proceedings of the National Academy of Sciences **107**(43): 18545-18550.
5.   Brizel, D. M., T. Schroeder, R. L. Scher, S. Walenta, R. W. Clough, M. W. Dewhirst and W. Mueller-
Klieser (2001). "Elevated tumor lactate concentrations predict for an increased risk of metastases in
head-and-neck cancer." Int J Radiat Oncol Biol Phys **51**(2): 349-353.
6.   Brooks, G. A. (1986). "The lactate shuttle during exercise and recovery." Medicine and science in
sports and exercise **18**(3): 360-368.
7.   Brooks, G. A. (2009). "Cell-cell and intracellular lactate shuttles." J Physiol **587**(Pt 23): 5591-5600.
8.   Brooks, G. A. (2018). "The Science and Translation of Lactate Shuttle Theory." Cell metabolism
27(4): 757-785.
9.   Brooks, G. A. (2020). "The tortuous path of lactate shuttle discovery: From cinders and boards to
the lab and ICU." Journal of Sport and Health Science(21).
10. Brooks, G. A., A. D. Osmond, J. A. Arevalo, C. C. Curl, J. J. Duong, M. A. Horning, D. D. Moreno
Santillan and R. G. Leija (2022). "Lactate as a major myokine and exerkine."
11. Brooks, G. A., A. D. Osmond, J. A. Arevalo, J. J. Duong, C. C. Curl, D. D. Moreno-Santillan and R.
G. Leija (2023). "Lactate as a myokine and exerkine: drivers and signals of physiology and
metabolism." J Appl Physiol (1985) **134**(3): 529-548.
12. Brown, M. A. and G. A. Brooks (1994). "Trans-stimulation of lactate transport from rat sarcolemmal
membrane vesicles." Arch Biochem Biophys **313**(1): 22-28.
13. Chao, Y. L., C. R. Shepard and A. Wells (2010). "Breast carcinoma cells re-express E-cadherin during
mesenchymal to epithelial reverting transition." Mol Cancer **9**: 179.
14. Chen, Z., Z. Fang and J. Ma (2021). "Regulatory mechanisms and clinical significance of vimentin
in breast cancer." Biomed Pharmacother **133**: 111068.
15. Cheng, F., Y. Shen, P. Mohanasundaram, M. Lindström, J. Ivaska, T. Ny and J. E. Eriksson (2016).
"Vimentin coordinates fibroblast proliferation and keratinocyte differentiation in wound healing
via TGF-β–Slug signaling." Proceedings of the National Academy of Sciences **113**(30): E4320-E4327.

16. de la Cruz-López, K. G., L. J. Castro-Muñoz, D. O. Reyes-Hernández, A. García-Carrancá and J. Manzo-Merino (2019). "Lactate in the Regulation of Tumor Microenvironment and Therapeutic Approaches." Front Oncol **9**: 1143.

17. Devlin, J. R. and E. W. Verschuren (2018). "More than a Tumor Suppressor: E-Cadherin Loss Drives Lung Cancer Metastasis." Am J Respir Cell Mol Biol **59**(2): 141-142.

18. Dhup, S., R. K. Dadhich, P. E. Porporato and P. Sonveaux (2012). "Multiple biological activities of lactic acid in cancer: influences on tumor growth, angiogenesis and metastasis." Curr Pharm Des **18**(10): 1319-1330.

19. Di Zhang, Z. T., , He Huang, Guolin Zhou, Chang Cui1 Yejing Weng, Wenchao Liu, Sunjoo Kim, Sangkyu Lee, Mathew Perez-Neut, Jun Ding, Daniel Czyz, Rong Hu, Zhen Ye, Maomao He, Y. George Zheng, Howard A. Shuman, Lunzhi Dai, Bing Ren, Robert G. Roeder, Lev Becker, & Yingming Zhao (2019). "Metabolic regulation of gene expression by histone lactylation." Nature(574): 575-580.

20. Doherty, J. R., C. Yang, K. E. Scott, M. D. Cameron, M. Fallahi, W. Li, M. A. Hall, A. L. Amelio, J. K. Mishra, F. Li, M. Tortosa, H. M. Genau, R. J. Rounbehler, Y. Lu, C. V. Dang, K. G. Kumar, A. A. Butler, T. D. Bannister, A. T. Hooper, K. Unsal-Kacmaz, W. R. Roush and J. L. Cleveland (2014). "Blocking lactate export by inhibiting the Myc target MCT1 Disables glycolysis and glutathione synthesis." Cancer Res **74**(3): 908-920.

21. Draoui, N., O. Schicke, A. Fernandes, X. Drozak, F. Nahra, A. Dumont, J. Douxfils, E. Hermans, J. M. Dogne, R. Corbau, A. Marchand, P. Chaltin, P. Sonveaux, O. Feron and O. Riant (2013). "Synthesis and pharmacological evaluation of carboxycoumarins as a new antitumor treatment targeting lactate transport in cancer cells." Bioorg Med Chem **21**(22): 7107-7117.

22. Draoui, N., O. Schicke, E. Seront, C. Bouzin, P. Sonveaux, O. Riant and O. Feron (2014). "Antitumor activity of 7-aminocarboxycoumarin derivatives, a new class of potent inhibitors of lactate influx but not efflux." Mol Cancer Ther **13**(6): 1410-1418.

23. Forsythe, J. A., B.-H. Jiang, N. V. Iyer, F. Agani, S. W. Leung, R. D. Koos and G. L. Semenza (1996). "Activation of vascular endothelial growth factor gene transcription by hypoxia-inducible factor 1." Molecular and cellular biology **16**(9): 4604-4613.

24. Friedrichs, K., S. Gluba, H. Eidtmann and W. Jonat (1993). "Overexpression of p53 and prognosis in breast cancer." Cancer **72**(12): 3641-3647.

25. Gregor, M., S. Osmanagic-Myers, G. Burgstaller, M. Wolfram, I. Fischer, G. Walko, G. P. Resch, A. Jörgl, H. Herrmann and G. Wiche (2014). "Mechanosensing through focal adhesion-anchored intermediate filaments." The FASEB Journal **28**(2): 715-729.

26. Haber, D. A. and J. Settleman (2007). "Drivers and passengers." Nature **446**(7132): 145-146.

27. Halestrap, A. P. (2012). "The monocarboxylate transporter family--Structure and functional characterization." IUBMB Life **64**(1): 1-9.

28. Hashimoto, T., R. Hussien, S. Oommen, K. Gohil and G. A. Brooks (2007). "Lactate sensitive transcription factor network in L6 cells: activation of MCT1 and mitochondrial biogenesis." FASEB J **21**(10): 2602-2612.

29. Henderson, G. C., M. A. Horning, S. L. Lehman, E. E. Wolfel, B. C. Bergman and G. A. Brooks (2004). "Pyruvate shuttling during rest and exercise before and after endurance training in men." J Appl Physiol **97**(1): 317-325.

30. Hirota, K. and G. L. Semenza (2006). "Regulation of angiogenesis by hypoxia-inducible factor 1." Critical Reviews in Oncology/Hematology **59**(1): 15-26.

31. Huang, S., E. A. Armstrong, S. Benavente, P. Chinnaiyan and P. M. Harari (2004). "Dual-agent molecular targeting of the epidermal growth factor receptor (EGFR) combining anti-EGFR antibody with tyrosine kinase inhibitor." Cancer research **64**(15): 5355-5362.

32. Hussien, R., B. H. Rihn, H. Eidi, C. Ronzani, O. Joubert, L. Ferrari, O. Vazquez, D. Kaufer and G. A. Brooks (2013). "Unique growth pattern of human mammary epithelial cells induced by polymeric nanoparticles." Physiol Rep **1**(4): e00027.

33. Isola, J., T. Visakorpi, K. Holli and O. P. Kallioniemi (1992). "Association of overexpression of tumor suppressor protein p53 with rapid cell proliferation and poor prognosis in node-negative breast cancer patients." J Natl Cancer Inst **84**(14): 1109-1114.

34. Jang, S. M., H. Han, K. S. Jang, Y. J. Jun, S. H. Jang, K. W. Min, M. S. Chung and S. S. Paik (2012). "The Glycolytic Phenotype is Correlated with Aggressiveness and Poor Prognosis in Invasive Ductal Carcinomas." J Breast Cancer **15**(2): 172-180.

35. Jeong, Y., S. Y. Bae, D. You, S. P. Jung, H. J. Choi, I. Kim, S. K. Lee, J. Yu, S. W. Kim and J. E. Lee (2019). "EGFR is a therapeutic target in hormone receptor-positive breast cancer." Cell Physiol Biochem **53**(5): 805-819.

36. Jiang, J., D. Huang, Y. Jiang, J. Hou, M. Tian, J. Li, L. Sun, Y. Zhang, T. Zhang, Z. Li, Z. Li, S. Tong and Y. Ma (2021). "Lactate Modulates Cellular Metabolism Through Histone Lactylation-Mediated Gene Expression in Non-Small Cell Lung Cancer." Front Oncol **11**: 647559.

37. Katsumoto, T., A. Mitsushima and T. Kurimura (1990). "The role of the vimentin intermediate filaments in rat 3Y1 cells elucidated by immunoelectron microscopy and computer-graphic reconstruction." Biol Cell **68**(2): 139-146.

38. Kopfstein, L. and G. Christofori (2006). "Metastasis: cell-autonomous mechanisms versus contributions by the tumor microenvironment." Cell Mol Life Sci **63**(4): 449-468.

39. Kumar, V. B., R. I. Viji, M. S. Kiran and P. R. Sudhakaran (2007). "Endothelial cell response to lactate: implication of PAR modification of VEGF." J Cell Physiol **211**(2): 477-485.

40. Kuo, T.-C., K.-Y. Huang, S.-C. Yang, S. Wu, W.-C. Chung, Y.-L. Chang, T.-M. Hong, S.-P. Wang, H.-Y. Chen, T.-H. Hsiao and P.-C. Yang (2020). "Monocarboxylate Transporter 4 Is a Therapeutic Target in Non-small Cell Lung Cancer with Aerobic Glycolysis Preference." Molecular Therapy - Oncolytics **18**: 189-201.

41. Lavin, M. F., D. Delia and L. Chessa (2006). "ATM and the DNA damage response. Workshop on ataxia-telangiectasia and related syndromes." EMBO Rep **7**(2): 154-160.

42. Lee, J.-K., S. Hahn, D.-W. Kim, K. J. Suh, B. Keam, T. M. Kim, S.-H. Lee and D. S. Heo (2014). "Epidermal growth factor receptor tyrosine kinase inhibitors vs conventional chemotherapy in non–small cell lung cancer harboring wild-type epidermal growth factor receptor: a meta-analysis." Jama **311**(14): 1430-1437.

43. Lee, M., I. A. Park, S. H. Heo, Y. A. Kim, G. Gong and H. J. Lee (2019). "Association between p53 Expression and Amount of Tumor-Infiltrating Lymphocytes in Triple-Negative Breast Cancer." J Pathol Transl Med **53**(3): 180-187.

44. Li, J. P., X. M. Zhang, Z. Zhang, L. H. Zheng, S. Jindal and Y. J. Liu (2019). "Association of p53 expression with poor prognosis in patients with triple-negative breast invasive ductal carcinoma." Medicine (Baltimore) **98**(18): e15449.

45. Liu, W., Y. Wang, L. H. Bozi, P. Fischer, M. P. Jedrychowski, H. Xiao, T. Wu, N. Darabedian, X. He and E. L. Mills (2023). "Lactate regulates cell cycle by remodeling the anaphase promoting complex." Nature: 1-3.

46. Lunetti, P., M. Di Giacomo, D. Vergara, S. De Domenico, M. Maffia, V. Zara, L. Capobianco and A. Ferramosca (2019). "Metabolic reprogramming in breast cancer results in distinct mitochondrial bioenergetics between luminal and basal subtypes." FEBS J **286**(4): 688-709.

47. Mishra, D. and D. Banerjee (2019). "Lactate Dehydrogenases as Metabolic Links between Tumor and Stroma in the Tumor Microenvironment." Cancers (Basel) **11**(6).

48. Mittal, V. (2018). "Epithelial Mesenchymal Transition in Tumor Metastasis." Annual Review of Pathology: Mechanisms of Disease **13**(1): 395-412.

49. Moerkens, M., Y. Zhang, L. Wester, B. van de Water and J. H. Meerman (2014). "Epidermal growth factor receptor signalling in human breast cancer cells operates parallel to estrogen receptor alpha signalling and results in tamoxifen insensitive proliferation." <u>BMC Cancer</u> **14**: 283.

50. Ocana, M. C., B. Martinez-Poveda, A. R. Quesada and M. A. Medina (2020). "Glucose Favors Lipid Anabolic Metabolism in the Invasive Breast Cancer Cell Line MDA-MB-231." <u>Biology (Basel)</u> **9**(1).

51. Pérez-Tomás, R. and I. Pérez-Guillén (2020). "Lactate in the Tumor Microenvironment: An Essential Molecule in Cancer Progression and Treatment." <u>Cancers (Basel)</u> **12**(11).

52. Polet, F. and O. Feron (2013). "Endothelial cell metabolism and tumour angiogenesis: glucose and glutamine as essential fuels and lactate as the driving force." <u>J Intern Med</u> **273**(2): 156-165.

53. Reda, A., A. Refaat, A. A. Abd-Rabou, A. M. Mahmoud, M. Adel, S. Sabet and S. S. Ali (2019). "Role of mitochondria in rescuing glycolytically inhibited subpopulation of triple negative but not hormone-responsive breast cancer cells." <u>Scientific Reports</u> **9**(1): 13748.

54. Roth, D. A. and G. A. Brooks (1990). "Lactate and pyruvate transport is dominated by a pH gradient-sensitive carrier in rat skeletal muscle sarcolemmal vesicles." <u>Archives of biochemistry and biophysics</u> **279**(2): 386-394.

55. Roth, D. A. and G. A. Brooks (1990). "Lactate transport is mediated by a membrane-bound carrier in rat skeletal muscle sarcolemmal vesicles." <u>Archives of biochemistry and biophysics</u> **279**(2): 377-385.

56. San-Millan, I. and G. A. Brooks (2017). "Reexamining cancer metabolism: lactate production for carcinogenesis could be the purpose and explanation of the Warburg Effect." <u>Carcinogenesis</u> **38**(2): 119-133.

57. San-Millan, I., C. G. Julian, C. Matarazzo, J. Martinez and G. A. Brooks (2019). "Is Lactate an Oncometabolite? Evidence Supporting a Role for Lactate in the Regulation of Transcriptional Activity of Cancer-Related Genes in MCF7 Breast Cancer Cells." <u>Front Oncol</u> **9**: 1536.

58. Semenza, G. L. (2010). "HIF-1: upstream and downstream of cancer metabolism." <u>Curr Opin Genet Dev</u> **20**(1): 51-56.

59. Sonveaux, P., T. Copetti, C. J. De Saedeleer, F. Vegran, J. Verrax, K. M. Kennedy, E. J. Moon, S. Dhup, P. Danhier, F. Frerart, B. Gallez, A. Ribeiro, C. Michiels, M. W. Dewhirst and O. Feron (2012). "Targeting the lactate transporter MCT1 in endothelial cells inhibits lactate-induced HIF-1 activation and tumor angiogenesis." <u>PLoS One</u> **7**(3): e33418.

60. Sonveaux, P., F. Vegran, T. Schroeder, M. C. Wergin, J. Verrax, Z. N. Rabbani, C. J. De Saedeleer, K. M. Kennedy, C. Diepart, B. F. Jordan, M. J. Kelley, B. Gallez, M. L. Wahl, O. Feron and M. W. Dewhirst (2008). "Targeting lactate-fueled respiration selectively kills hypoxic tumor cells in mice." <u>J Clin Invest</u> **118**(12): 3930-3942.

61. Su, Y., N. R. Hopfinger, T. D. Nguyen, T. J. Pogash, J. Santucci-Pereira and J. Russo (2018). "Epigenetic reprogramming of epithelial mesenchymal transition in triple negative breast cancer cells with DNA methyltransferase and histone deacetylase inhibitors." <u>J Exp Clin Cancer Res</u> **37**(1): 314.

62. Thiery, J. P. (2002). "Epithelial-mesenchymal transitions in tumour progression." <u>Nat Rev Cancer</u> **2**(6): 442-454.

63. Vlassenko, A. G., J. McConathy, L. E. Couture, Y. Su, P. Massoumzadeh, H. S. Leeds, M. R. Chicoine, D. D. Tran, J. Huang, S. Dahiya, D. S. Marcus, S. J. Fouke, K. M. Rich, M. E. Raichle and T. L. Benzinger (2015). "Aerobic Glycolysis as a Marker of Tumor Aggressiveness: Preliminary Data in High Grade Human Brain Tumors." <u>Dis Markers</u> **2015**: 874904.

64. Vogelstein, B., N. Papadopoulos, V. E. Velculescu, S. Zhou, L. A. Diaz, Jr. and K. W. Kinzler (2013). "Cancer genome landscapes." <u>Science</u> **339**(6127): 1546-1558.

65. Wang, L., C. Xu, X. Liu, Y. Yang, L. Cao, G. Xiang, F. Liu, S. Wang, J. Liu, Q. Meng, J. Jiao and Y. Niu (2019). "TGF-β1 stimulates epithelial–mesenchymal transition and cancer-associated

myoepithelial cell during the progression from in situ to invasive breast cancer." <u>Cancer Cell International</u> **19**(1): 343.

66. Wang, L., X. Yu, J. Li, Z. Zhang, J. Hou and F. Li (2016). "Prognostic significance of p53 expression in patients with esophageal cancer: a meta-analysis." <u>BMC Cancer</u> **16**(1): 373.

67. Wang, Z., Q. Jiang and C. Dong (2020). "Metabolic reprogramming in triple-negative breast cancer." <u>Cancer Biol Med</u> **17**(1): 44-59.

68. Warburg, O. (1925). "The Metabolism of Carcinoma Cells." <u>The Journal of Cancer Research</u> **9**(1): 148-163.

69. Warburg, O. and S. Minami (1923). "Versuche an überlebendem carcinom-gewebe." <u>Journal of Molecular Medicine</u> **2**(17): 766-777.

70. Watson, J. D. and F. H. Crick (1953). "Molecular structure of nucleic acids: a structure for deoxyribose nucleic acid." <u>Nature</u> **171**(4356): 737-738.

71. Wells, A., C. Yates and C. R. Shepard (2008). "E-cadherin as an indicator of mesenchymal to epithelial reverting transitions during the metastatic seeding of disseminated carcinomas." <u>Clin Exp Metastasis</u> **25**(6): 621-628.

72. Zhang, D., Y. Zheng, S. Yang, Y. Li, M. Wang, J. Yao, Y. Deng, N. Li, B. Wei, Y. Wu, Y. Zhu, H. Li and Z. Dai (2020). "Identification of a Novel Glycolysis-Related Gene Signature for Predicting Breast Cancer Survival." <u>Front Oncol</u> **10**: 596087.