1

Adverse outcomes in SARS-CoV-2 infected pregnant mice are gestational age-dependent

and resolve with antiviral treatment

Patrick S. Creisher,[1,*] Jamie L. Perry,[1,*] Weizhi Zhong,[1] Jun Lei, [2,3] Kathleen R Mulka,[4] Hurley

Ryan,[5] Ruifeng Zhou,[1] Elgin H. Akin,[1] Anguo Liu, [2,3] Wayne Mitzner, [5] Irina Burd,[2,3,†] Andrew

Pekosz,[1,5,†] Sabra L. Klein[1,†]

[1]W. Harry Feinstone Department of Molecular Microbiology and Immunology, The Johns

Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA

[2]Integrated Research Center for Fetal Medicine, Department of Gynecology and Obstetrics,

Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

[3]Department of Obstetrics, Gynecology & Reproductive Sciences, University of Maryland School

of Medicine, Baltimore, Maryland, USA

[4]Department of Molecular and Comparative Pathobiology, The Johns Hopkins School of

Medicine, Baltimore, Maryland.

[5]Department of Environmental Health and Engineering, The Johns Hopkins Bloomberg School

of Public Health, Baltimore, MD, USA.

[†]To whom correspondence should be addressed: Sabra L. Klein (sklein2@jhu.edu), Andrew

Pekosz (apekosz1@jhu.edu), & Irina Burd (iburd@som.umaryland.edu)

*co-first authors

Short title: SARS-CoV-2 during mouse pregnancy

Key words: coronavirus, gestation, anti-viral, morbidity, development, Paxlovid

Manuscript statistics:

      Abstract: 195 words
      Text: 10029 words
      Figures: 7
      References: 68

Conflict of interest statement:

The authors have declared that no conflict of interest exists.

Abstract:

SARS-CoV-2 infection during pregnancy is associated with severe COVID-19 and

adverse fetal outcomes, but the underlying mechanisms remain poorly understood. Moreover,

clinical studies assessing therapeutics against SARS-CoV-2 in pregnancy are limited. To

address these gaps, we developed a mouse model of SARS-CoV-2 infection during pregnancy.

Outbred CD1 mice were infected at embryonic day (E) 6, E10, or E16 with a mouse adapted

SARS-CoV-2 (maSCV2) virus. Outcomes were gestational age-dependent with greater

morbidity, reduced pulmonary function, reduced anti-viral immunity, greater viral titers, and more

adverse fetal outcomes occurring with infection at E16 (3rd trimester-equivalent) than with

infection at either E6 (1st trimester-equivalent) or E10 (2nd trimester-equivalent). To assess the

efficacy of ritonavir-boosted nirmatrelvir (recommended for pregnant individuals with COVID-

19), we treated E16-infected dams with mouse equivalent doses of nirmatrelvir and ritonavir.

Treatment reduced pulmonary viral titers, decreased maternal morbidity, and prevented adverse

offspring outcomes. Our results highlight that severe COVID-19 during pregnancy and adverse

fetal outcomes are associated with heightened virus replication in maternal lungs. Ritonavir-

boosted nirmatrelvir mitigated adverse maternal and fetal outcomes of SARS-CoV-2 infection.

These findings prompt the need for further consideration of pregnancy in preclinical and clinical

studies of therapeutics against viral infections.

1. Introduction:

Pregnancy is a risk factor for developing severe COVID-19, with pregnant individuals at increased risk of hospitalization, mechanical ventilation, and mortality compared to non-pregnant patients (1, 2). The time of infection during gestation contributes to increased severity, with hospitalization and intensive care unit admission being greater in the third than either the second or first trimester (3). While the specific mechanisms that contribute to the increased risk of severe outcomes during pregnancy are not specified, both immunological and physiological changes are likely involved. The immune system undergoes unique shifts as pregnancy progresses, including increased regulatory T and B lymphocytes as well as reduced cytotoxic and cellular immunity, to protect the developing semi-allogenic fetus (4, 5). The general anti-inflammatory shift during the second and third trimesters also may increase the risk of severe outcomes from viruses, including SARS-CoV-2, by blunting anti-viral immune responses (5). Moreover, physiological changes associated with pregnancy including cardiovascular, respiratory, endocrine, and metabolic alterations may further contribute to disease severity (6). While these pregnancy-associated factors are hypothesized to contribute to severe disease and death following infection with SARS-CoV-2, the exact mechanisms contributing to severe COVID-19 disease during pregnancy in humans remain unknown.

In addition to causing severe outcomes in pregnant patients, SARS-CoV-2 infection during pregnancy also can result in adverse fetal outcomes including preterm birth, stillbirth, small size for gestational age, and reduced birth weight (7), as well as increased risks of neurobehavioral deficits and delayed motor skills in infants born to infected mothers (8). Like maternal disease, adverse perinatal and fetal outcomes appear to be influenced by gestational age, with greater risk observed after infection in the third as compared with either the second or first trimester (3, 8, 9). Direct placental infection or vertical transmission of SARS-CoV-2 is exceedingly rare (10-12), and thus is unlikely to be the source of adverse fetal and neonatal outcomes. The exact mechanisms underlying these adverse outcomes remain unknown.

Because of their risk for severe COVID-19 and adverse pregnancy outcomes, pregnant

individuals are prioritized for receipt of available emergency use authorized antivirals and

vaccines (13), despite being excluded from clinical trials of SARS-CoV-2 vaccines and antivirals

(14). SARS-CoV-2 mRNA vaccines have been proven to be safe and effective during pregnancy

(15-17), and the United States Centers for Disease Control and Prevention recommends

vaccination for women who are pregnant, recently pregnant, or trying to become pregnant (15).

The safety and efficacy of SARS-CoV-2 antivirals during pregnancy has not been as well

studied. Pregnant people are recommended to receive the antivirals remdesivir (brand name

Veklury) and ritonavir-boosted nirmatrelvir (brand name Paxlovid) when indicated (18). While

neither antiviral included pregnant people in their clinical trials (19), observational studies of

remdesivir indicate its safety and efficacy in pregnant populations (20). Nirmatrelvir is an oral

antiviral that inhibits the SARS-CoV-2 M^{PRO} protease and is packaged with ritonavir, a

previously established HIV protease inhibitor and pharmacologic booster, which does not have

direct antiviral effects on SARS-CoV-2 but instead works to prolong the bioavailability of

nirmatrelvir through the inhibition of the hepatic cytochrome P-450 (CYP) 3A4 enzyme (21, 22).

Ritonavir-boosted nirmatrelvir treatment during pregnancy appears safe, with no adverse

obstetric outcomes reported in small observational studies (23, 24). The efficacy of ritonavir-

boosted nirmatrelvir in preventing SARS-CoV-2 infection or disease during pregnancy remains

an open question, in part because most studies to date were not designed to evaluate efficacy

(23, 24).

Animal models of microbial infections during pregnancy provide mechanistic insight into

adverse maternal and fetal outcomes by enabling deeper analysis of vertical transmission and

maternal and fetal immune responses. Animal models have elucidated the pathogenesis of

infections such Zika virus, influenza A virus, *Plasmodium falciparum,* and Group B

Streptococcus infections during pregnancy (13). In the absence of human clinical trials, animal

models of infection during pregnancy can be used to characterize the safety and efficacy of

therapeutics in this high-risk population. To date, published animal models of SARS-CoV-2

infection during pregnancy have been limited (12, 25), which has hindered investigation into

both host and viral factors that may underlie the severe outcomes observed in humans. Animal

models have only been used to study the potential reproductive toxicity of nirmatrelvir in rats

and rabbits, with no evidence of fetal abnormalities, maternal toxicity, or other adverse

outcomes (26). Whether equivalent dosing of nirmatrelvir administered during pregnancy is

equally efficacious against SARS-CoV-2 infection in pregnancy as in nonpregnant animals has

not been reported.

In the current study, we developed a mouse model of SARS-CoV-2 infection during

pregnancy to investigate maternal and offspring outcomes and elucidate the contribution of

gestational age, pulmonary and placental involvement in adverse outcomes, and control of virus

replication. Further, we sought to assess the efficacy of ritonavir-boosted nirmatrelvir in limiting

virus replication, preventing maternal disease, and mitigating adverse offspring outcomes. Our

results demonstrate that SARS-CoV-2 infection during late gestation causes more severe

maternal disease and adverse offspring outcomes than infections earlier during gestion, which

is associated with reduced pulmonary anti-viral type 1 interferon (IFN) responses, greater viral

replication in the lungs, and placental insufficiencies. Treatment with ritonavir-boosted

nirmatrelvir not only reduced maSCV2 replication but prevented severe disease and adverse

fetal outcomes, highlighting additional benefits of antiviral treatment during pregnancy.

2 Materials and Methods:

2.1 Viruses and cells

A mouse adapted strain of SARS-CoV-2 (maSCV2), originally generated by Dr. Ralph

Baric (27) was obtained from the Biodefense and Emerging Infections Research Resources

Repository (BEI Resources #NR-55329). Working stocks of maSCV2 virus were generated by

infecting Vero-E6-TMPRSS2 cells at a multiplicity of infection (MOI) of 0.01 tissue culture

infectious dose 50 ($TCID_{50}$) per cell in infection media [Dulbecco's Modified Eagle Medium

(DMEM; Sigma # D5796) supplemented with 2.5% filter-sterilized fetal bovine serum (Gibco#

10-437-028), 100 U/ml penicillin and 100 μg/ml streptomycin (Gibco #15149-122), 1 mM l-

glutamine (Gibco #2503081), and 1-mM sodium pyruvate (Gibco #11-360-070)]. Approximately

72 hours post infection, the supernatant fluids were collected, clarified by centrifugation (400g

for 10 minutes), and stored in aliquots at -70°C.

2.2 Experimental mice

Adult (8-12 weeks of age) timed pregnant and nonpregnant female CD-1 IGS mice were

purchased from Charles River Laboratories. Pregnant mice arrived on embryonic day (E) 4, E8,

and E14 and were singly housed, and nonpregnant female mice were housed at 5 per cage.

Mice were housed under standard animal biosafety level three (ABSL3) housing conditions with

ad libitum food and water. Mice were given at least 24 hours to acclimate to the ABSL3 facility

prior to infections (28). All monitoring and experimental procedures were performed at the same

time each day.

2.3 SCV2 infections and monitoring

All animal experiments and procedures took place in an ABSL3 facility at the Johns

Hopkins School of Medicine. Experimental pregnant mice were intranasally infected at E6, E10,

or E16 with $TCID_{50}$ 10^5 of maSCV2 in 30 μl of DMEM (Sigma #D5796) or mock inoculated with

30 μl of media. Prior to intranasal infection, mice were anesthetized via intraperitoneal

ketamine/xylazine cocktail (80 mg/kg ketamine, 5 mg/kg xylazine). Following intranasal

infections, body mass and clinical signs of disease were monitored for 14 days or until tissue

collection. Clinical scores given to mice on a scale of 0-4, with a point given for piloerection,

dyspnea, hunched posture, and absence of an escape response on each day (29, 30). Clinical

scores over the course of 14 days for each animal were summed to give a cumulative clinical

disease score.

2.4 Antiviral treatment

Experimental animals were administered vehicle alone [1% (w/v) Soluplus (BASF

#50539897), 1% (w/v) Tween 80 (Sigma #59924), 0.5% (w/v) methylcellulose (Sigma #94378)

in purified water], high dose nirmatrelvir alone (300mg/kg; MedChem Express #HY-138687), or

an animal equivalent dose of nirmatrelvir boosted with ritonavir [1.7 mg nirmatrelvir/dose

(MedChem Express #HY-138687), 0.6 mg ritonavir/dose (Sigma #(#155213-67-5)]. Animal

equivalent doses were calculated as described (31) by converting the standard human dose of

nirmatrelvir and ritonavir (32) to a body-surface-area equivalent for mice (31) and using an

assumed mass of 30g for all calculations so that pregnant and nonpregnant animals receive the

same amount per dose. Mice were administered treatment via oral gavage twice daily for 5 days

or until tissue collection, starting 4 hours after infection as described in a published pre-clinical

study of nirmatrelvir (33).

2.5 Offspring measurements and behavior

Offspring from mock inoculated dams and maSCV2 infected dams were measured at

postnatal day (PND) 0, within 12 hours of birth. Body mass (g), length measured from nose to

anus (mm), and head diameter measured from ear to ear (mm) were recorded for each pup,

and the average for each independent litter was calculated and reported to avoid confounding

litter effects. Pups at PND5 were subjected to developmental neurobehavioral assays of surface

righting, cliff aversion, and negative geotaxis as described (34, 35). For each test, 1–2 male and

1–2 female offspring from at least 5 independent litters were used per condition to avoid

confounding litter effects. Pups were subjected to 3 attempts at each test, with the time to

complete each test recorded on a stopwatch. The pups' best trial for each test was used for

analysis.

2.6 Diffusion capacity of carbon monoxide

To measure the lung function of experimental mice, diffusion capacity for carbon

monoxide (DF_{CO}) was measured. Modifications to previously a published protocol (36) were

made for application in ABSL3. Mice were anesthetized with ketamine/xylazine cocktail (80

mg/kg ketamine, 5 mg/kg xylazine). Mice were tracheostomized with an 18-G stub needle. For

each mouse, two 3 ml syringes containing 0.8 ml of ~0.5% neon (Ne, an insoluble inert tracer

gas), ~0.5% carbon monoxide (CO), and balanced air were pre-filled and sealed with a 4-way

stop cock. Following tracheostomy, the gas was injected into the tracheostomy stub-needle to

inflate the lungs for two seconds and held for eight seconds. After eight seconds, the 0.8 ml

volume was withdrawn back into the syringe in two seconds and the syringe's stop cock closed,

then the gas in the syringe was diluted to 2 ml with room air and resealed. This was repeated

using the second syringe for each mouse. Mice were euthanized via cervical dislocation. The

closed syringes were decontaminated in an oven at 75°C for 15 minutes within the ABSL3 to

inactivate any virus in the gas sample. DF_{CO} was measured using gas chromatography as

previously described (36).

2.7 Tissue collection

Experimental dams (infected at E6, E10, or E16) or nonpregnant female mice were

euthanized at 3 days post-infection. Mice were anesthetized via isoflurane and exsanguination

was preformed via cardiac puncture. At the time of euthanasia, the total number of viable and

nonviable fetuses was quantified for each pregnant dam. Fetal viability was determined as the

percentage of fetuses within uterine horns that were viable. Fetuses were counted as nonviable

if they were smaller or discolored compared to gestational age-matched live fetuses or if a fetus

was absent at an implantation site (35, 37, 38). Maternal lungs were collected, separated by

lobe, and flash frozen on dry ice for homogenization. The left lung was inflated and fixed in zinc

buffered formalin (Thermo Fisher Scientific #NC9351419) for at least 72 hours in preparation for

histology. Fetuses and placenta were flash frozen in dry ice or fixed in 4% paraformaldehyde

(Thermo Fisher Scientific #J19943.K2) for 72- hours at 4°C for immunohistochemistry. A subset

of uninfected pregnant (E16) and nonpregnant adult mice were euthanized and the median liver

lobe was collected, and flash frozen in dry ice for Western blot.

2.8 Pulmonary histopathology

Fixed lungs were sliced into 3-mm blocks, embedded in paraffin, sectioned to 5 μm,

mounted on glass slides, and stained with hematoxylin and eosin (H&E) solution to evaluate

lung inflammation. Semiquantitative histopathological scoring was performed by a board-

certified veterinary pathologist to measure both severity of inflammation and the extent of

inflammation (39-41). Severity of perivascular and peribronchiolar mononuclear inflammation

was scored on a scale of 0- to 4 (0, no inflammation; 1, 1 cell layer; 2, 2-3 cell layers; 3, 4-5 cell

layers; 4, > 5 cell layers). Severity of alveolar inflammation was scored on a scale of 0-4 (0, no

inflammation; 1- increased inflammatory cells in alveoli, septa clearly distinguished; 2 –

inflammatory cells fill alveoli, septa clearly distinguished; 3 – inflammatory cells fill multiple

adjacent alveoli, septa difficult to distinguish; 4 – inflammatory cells fill multiple adjacent alveoli

with septal necrosis). Extent of inflammation was scored separately for perivascular,

peribronchiolar, and alveolar areas on a scale of 0 to 4 (0, no inflammation; 1, 2-25% tissue

affected; 2, up to 50% tissue affected; 3, up to 75% tissue affected; 4, >75% of tissue affected).

Individual scores were summed to give a cumulative inflammation score.

2.9 Infectious virus quantification and tissue inactivation

Frozen right cranial lungs, nasal turbinates, placentas, and fetuses were homogenized in

lysing matrix D bead tubes (MP Biomedicals #6913100). Homogenization media [500ml DMEM

(Sigma # D5796), 5ml penicillin/streptomycin (Gibco #15149-122)] was added to bead tubes

containing tissue at a minimum volume of 400μl and maximum volume of 1200μl (10% w/v) and

homogenized at 4.0m/s for 45s in a MP Fast-prep 24 5G instrument. After homogenization, the

supernatant was divided in half and transferred to two new microcentrifuge tube. Triton X-100

was added to one of the transferred supernatants to a final concentration of 0.5% to inactivate

maSCV2 and incubated at room temp for 30 minutes. Infectious and inactivated homogenates

were stored at -80°C. Infectious virus titers in tissue homogenate were determined by $TCID_{50}$

assay. Tissue homogenates were serially diluted in infection media in sextuplicate into 96-well

plates confluent with Vero-E6-TMPRSS2 cells, incubated at 37°C for six days. After incubation,

10% neutral buffered formalin was added to all wells to fix cells prior to staining and left

overnight. Formalin was discarded and the plates were stained with naphthol blue black stain for

visualization. Infectious virus titers were determined via the Reed and Muench method.

2.10 Placental histology and immunohistochemistry

Placentas were fixed for 72 hours at 4°C in 4% PFA in the ABSL3. Placentas were

washed five times with PBS and immersed in 30% sucrose until saturation. Using a Leica

CM1950 cryostat, the specimens were cut at 20-μm thickness and mounted on positively

charged slides (Fisher Scientific #12-550-15). Routine H&E staining was performed to evaluate

the morphological change of the placentas. For immunohistochemical staining, slides were

washed with PBS, which was followed by permeabilization in PBS solution containing 0.05%

Triton X-100 and 10% normal goat serum (Invitrogen #50197Z) for 30 min. Placentas were

incubated with rabbit anti-vimentin (1:200, Abcam # ab92547), or rabbit anti-cytokeratin (1:200,

Dako #Z0622) overnight at 4°C. The next day, sections were rinsed with PBS and then

incubated with donkey anti-rabbit (ThermoFisher #R37119) fluorescent secondary antibodies

(ThermoFisher #R37115) diluted 1:500 for 3 h at room temperature. DAPI (Roche

#10236276001) was applied for counterstaining, followed by mounting with Fluoromount-G

(eBioscience #00-4958-02). Images were taken using a Zeiss Axioplan 2 microscope (Jena,

Germany) under 20x magnification. Cell density of vimentin and cytokeratin positive cell

quantification was performed using Image J (1.47v). For each placenta, six random images in

labyrinth at the middle level (thickest) of placenta were taken, and the average fluorescent area

calculated for that placenta. One placenta per dam was used and 4–5 dams per group were

analyzed.

2.11 Cortical thickness measurement

A subset of offspring was randomly selected to be euthanized via decapitation at PND 0

and heads were fixed for 72 hours at 4°C in 4% PFA in the ABSL3. Fetal heads were washed

five times with PBS and immersed in 30% sucrose until saturation. Using a Leica CM1950

cryostat, the specimens were cut at 20-µm thickness and mounted on positively charged slides

(Fisher Scientific #12-550-15). Nissl staining was performed, and images were taken under × 5

magnification using a Canon EOS Rebel (Tokyo, Japan). Coronal cortical thickness was

measured from five random sections at the striatum level of each neonatal brain, as previously

described (38). Cortical thickness was measured from both brain hemispheres in each section

using ImageJ software, and the average of 10 measurements per specimen presented.

Quantification shown represents the average measurement from a single randomly chosen pup

for each dam.(38)

2.12 Interferon β ELISA

Interferon β in inactivated right cranial lung homogenate was measured by ELISA

according to the manufacturer's protocol (PBL Assay Science # 42410-1).

2.13 Western Blot

Flash frozen median liver lobes were homogenized in 1X Cell lysis Buffer (Cell Signaling

Technology #9803) with 1X Protease Inhibitor cocktail (Sigma-Aldrich #P8340) and sodium

fluoride (Fisher Scientific #S299 100) (20µl lysis buffer per mg tissue). Protein lysates were

stored at -80°C until analysis. Protein concentration of each lysate was measured using the

Pierce BCA Protein Assay Kit (Thermo Fisher Scientific #23225). For each sample, 20µg of

protein was subjected to sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-

PAGE) on NuPAGE 4-12% Bis-Tris gels (Thermo Fisher Scientific #NP0329). The gel was

blotted onto Immobilon-FL PVDF Membrane (Millipore #IPFL00010) and the membranes were

blocked with a 1:1 mixture of 1XPBS/Tween-20 solution (Sigma-Aldrich #P3563) and Intercept

blocking buffer (LI-COR Biosciences #927-70001) for 30 minutes at room temperature.

Membranes were treated with a primary antibody diluted in blocking solution at 4°C overnight on

a rocker. Membranes were then washed with PBS-Tween three times and incubated in

secondary antibody solutions for one hour at room temperature on a rocker. Membranes were

washed three times in PBS-Tween and then imaged on a ProteinSimple FluoroChem Q imager.

Individual bands were quantified using Image Studio software (LI-COR Biosciences; version

3.1.4) The signal from each band was normalized against the GAPDH signal and graphed as

arbitrary units. Primary antibodies used were rabbit anti-P450 3A4/CYP3A4 (abcam #ab3572)

and mouse anti-GAPDH (abcam #ab82450). Secondary antibodies included goat anti- mouse

Alexa Fluor 488 (Thermo Fisher Scientific #A11001) and donkey anti-rabbit Alexa Fluor Plus

647 (Thermo Fisher Scientific #A32795).

2.14 Viral RNA extraction, sequencing, and analysis

For each sample, 200µl of right cranial lung or nasal turbinate homogenate was mixed

with 1mL of TRIzol (Invitrogen #15596026), followed by 200µl of chloroform (Fisher Scientific

#C298-500) to extract RNA, and centrifuged at 12,000 g for 15 minutes at 4°C. The clear portion

of the supernatant was then pelleted at 12,000 g for 10 minutes along with 500µl of 100%

isopropyl alcohol (Fisher #A416) at 4°C. Pelleted RNA was then washed with 75% of ethanol

(Fisher Scientific #BP2818-500), air dried and resuspended in 20µl of nuclease-free

water. Reverse transcription was carried out using ProtoScript® II First Strand cDNA Synthesis

Kit (New England Biolabs #E6560S) with random hexamer mix. The Mpro region was then

amplified using forward primer 5' ACAAAGATAGCACTTAAGGGTGG 3' and reverse primer 5'

GCGAGCTCTATTCTTTGCACTAA 3' and Oxford Nanopore sequenced by Plasmidsaurus

(SNPsaurus LLC). Consensus sequences for lung and turbinate virus isolates were imported

and aligned to Mpro ORF (NC_045512.2) using ClustalO v1.2.3 in Geneious Prime v2023.0.4.

Alignments were imported into R v4.1.1, visualized, and annotated using seqvisR v0.2.5.

ORF1a and nonstructural protein annotation was visualized using BioRender. Raw FASTQ files

for Mpro sequencing has been deposited through SRA under BioProject PRJNA940500 at

accession numbers SRR23689223 - SRR23689259.

2.15 Statistical analyses

Post infection body mass changes were plotted and the area under the curve (AUC) was

calculated to provide individual data points that captured change over time, with AUCs

compared with either two tailed unpaired t-test or two-way analysis of variance (ANOVA)

followed by *post-hoc* Bonferroni multiple comparisons tests. To compare body mass changes

across gestational ages, individual AUCs were subtracted from the average AUC of mock mice

at the same gestational age, with the difference from mock AUC compared with two-way

ANOVAs followed by *post-hoc* Bonferroni multiple comparisons test. Cumulative clinical scores

were analyzed using the Kruskal-Wallis test. Viral titers in lungs from infected dams were

analyzed using one-way or two-way ANOVAs followed by *post-hoc* Bonferroni multiple

comparisons test. Western blot quantification, IHC quantification, and cortical thickness

measurements were analyzed with two-tailed unpaired t tests. Cumulative inflammation scoring,

DFCO, IFN-β, and fetal measurements were analyzed with two-way ANOVAs followed by *post-*

hoc Bonferroni multiple comparisons test. Pup neurodevelopment results were analyzed with

two-way or three-way ANOVAs followed by *post-hoc* Bonferroni multiple comparisons test. Fetal

viability data were analyzed with a χ^2 test. Data are presented as mean ± SEM or as the

median (cumulative clinical score). Mean or median differences were considered statistically

significant at $p < 0.05$. Statistical analyses were performed using GraphPad Prism v9.5

(GraphPad Software).

2.16 Study approval

All animal procedures were approved by the Johns Hopkins University Animal Care and Use

Committee (MO21H246). SARS-CoV-2 was handled in a BSL-3 containment facility using

institution approved biosafety protocols (P2003120104).

3 Results:

3.1 Mouse adapted SARS-CoV-2 causes morbidity in pregnant mice, which increases with

gestational age

To evaluate if SARS-CoV-2 caused greater disease in pregnant than nonpregnant mice

and if maternal morbidity was impacted by gestational age, we intranasally inoculated outbred

pregnant CD1 dams at E6, E10, or E16, roughly corresponding developmentally to human first,

second, or third trimesters, respectively (42), or age-matched nonpregnant females with mouse

adapted SARS-CoV-2 (maSCV2)(27) or media and measured body mass change as a

measure of morbidity. Nonpregnant females (**Figure 1A**) and dams infected at E6 (**Figure 1B**)

experienced mild morbidity, losing significant body mass [i.e., approximately 10% of their initial

body mass by 4 days post infection (dpi)], but then appearing indistinguishable from mock-

inoculated females by 7 dpi. In contrast, dams infected at E10 (**Figure 1C**) or E16 (**Figure 1D**)

experienced prolonged maternal morbidity, with E10-infected dams gaining less body mass for

the remainder of gestation than mock inoculated dams (**Figure 1C**), and E16-infected dams

failing to regain body mass for the remainder of gestation or during lactation as compared to

mock-inoculated dams (**Figure 1D**). To compare the impact of gestation on maternal morbidity,

the change in body mass relative to gestational-age matched mock-inoculated animals (**Figure

1E**) and cumulative clinical scores of disease (**Figure 1F**) were analyzed. maSCV2 infection of

pregnant dams at E16 resulted in greater body mass loss and clinical disease than infection of

either nonpregnant females or dams at either E6 or E10. Pregnancy and gestational age

increase the severity of SARS-CoV-2 outcomes in mice, consistent with human COVID-19 data

(1, 3).

3.2 Pregnant dams infected late in gestation have reduced IFN-β responses, increased viral

load, and reduced pulmonary function after infection

Deficits in type 1 interferon (IFN) signaling are associated with severe COVID-19 in

nonpregnant people (43) and mice (44). Pregnancy is associated with downregulation of

prototypical cytolytic and anti-viral pathways, including type I IFNs, and upregulation of anti-

inflammatory pathways toward mid to late gestation (45). We hypothesized that E16-infected

dams would have a reduced type I IFN response after SARS-CoV-2 infection compared to E6-

infected dams and nonpregnant females. To test this, we infected pregnant dams at E6 and E16

as well as age-matched nonpregnant females with maSCV2 or media and collected lungs at 3

dpi. Infected nonpregnant females as well as E6-infected dams had significantly increased

concentrations of pulmonary IFN-β as compared with matched mock-inoculated females

(**Figure 2A**). In contrast, maSCV2 infection at E16 resulted in pulmonary concentrations of IFN-

β that were indistinguishable from vehicle-inoculated dams and suppressed relative to either

non-pregnant females or E6 dams infected with maSCV2 (**Figure 2A**). To determine if reduced

anti-viral IFN-β concentrations were associated with worse pulmonary outcomes, at 3dpi, we

evaluated infectious viral titers (**Figure 2B**), pulmonary histopathology (**Figure 2C-D**,

Supplemental Figure 1), and diffusion capacity (DF_{CO}; **Figure 2E**) in the lungs of nonpregnant,

E6-, or E16- pregnant females there were either maSCV2 or mock-infected. E16-infected dams

had significantly greater pulmonary titers of infectious virus than either E6-infected dams or

infected nonpregnant females (**Figure 2B**). maSCV2 infection induced pulmonary

histopathological changes, including intra-alveolar necrosis and inflammatory cell debris, and

peribronchiolar, and perivascular mononuclear inflammatory infiltrates that were observed in

nonpregnant (**Supplementary Figure 1A**, representative image), E6- (**Supplemental Figure**

1B, representative image), and E16- (**Figure 2C**, representative image) infected mice (**Figure**

2D, scoring) to equivalent levels. maSCV2 infection at E16 significantly reduced pulmonary

function, as measured by DF_{CO}, which was not observed following maSCV2 infection at E6 or in

nonpregnant females. These data suggest that late gestation is associated with reduced

antiviral responses, greater virus replication, and reduced pulmonary function.

3.3 SARS-CoV-2 infection late in gestation induces placental labyrinth congestion and

disruption of trophoblast cell layer

As placental pathology has been observed during COVID-19 (46-48), we next

investigated if maSCV2 could infect or cause damage to the placenta. Dams were mock- or

maSCV2-infected at E10 [when the placenta is formed (42)] or E16 and euthanized at 3 dpi, and

placentas and fetal tissues were analyzed for infectious virus and tissue damage. All placentas

and fetal tissues were negative for infectious virus (**Supplemental Figure 2**), consistent with

human reports that direct placental infection and vertical transmission during COVID-19 is rare

(10, 11). Despite no detectable infectious virus, placentas from E16-infected dams showed

evidence of labyrinth congestion, which can result in placental insufficiency (49), and reduced

number of mononuclear trophoblast giant cells (**Figure 3A**, representative images), suggestive

of damage to the trophoblast-endothelial cell barrier, which separates maternal and fetal blood

in the labyrinth of the murine placenta (50). Staining for cytokeratin (trophoblasts, **Figure 3B**

and **Supplemental Figure 3A**) and vimentin (endothelial cells, **Figure 3C** and **Supplemental**

Figure 3B) was performed and revealed a significant loss of trophoblasts, but not endothelial

cells, in placentas from E16-infected compared with mock-infected dams (**Figure 3B-C**). These

data illustrate disruption of the maternal and fetal barrier in the absence of direct viral infection

or vertical transmission. Moreover, cell numbers in placentas of E10-infected dams did not differ

from placentas of mock-inoculated dams (**Supplemental Figure 3**), suggesting that placental

damage may be associated with the more severe maternal disease seen with infection at E16.

3.4 SARS-CoV-2 infection late in gestation causes intrauterine growth restriction

COVID-19 during human pregnancy is associated with adverse pregnancy and fetal

outcomes including preterm birth, stillbirth, small size for gestational age, and reduced birth

weight (7). To evaluate if the maSCV2-induced maternal morbidity and placental damage

observed after infection at E16 was associated with adverse pregnancy or fetal outcomes, we

inoculated dams with maSCV2 or media at E6, E10, or E16, with a subset of dams euthanized

at 3 dpi to evaluate fetal viability and the remainder followed to evaluate birth outcomes. Neither

fetal viability (**Figure 4A**) nor litter size (**Figure 4B**) was affected by maSCV2 infection during

pregnancy at any gestational age. maSCV2 infection at E6 or E10 resulted in modest reductions

in fetal growth relative to fetuses from mock-inoculated dams (**Figure 4C-E**). In contrast,

maSCV2 infection at E16 led to significantly smaller pups in terms of mass, length, and head

size relative to fetuses from mock-infected dams (**Figure 4C-E**). Collectively, fetuses from E16-

infected dams had greater growth restriction than fetuses from either E6- or E10-infected dams

(**Figure 4C-E**). Reduced birth size was not mediated by pre-term birth as all dams, regardless of

infection, delivered at approximately E20 (51). These data indicate that maSCV2 infection

during the third trimester-equivalent of pregnancy results intrauterine growth restriction, which

was not observed when infection occurs earlier during gestation.

3.5 Offspring of SARS-CoV-2 infection late in gestation display cortical thinning and reduced

neurodevelopmental behaviors

In addition to adverse perinatal outcomes, COVID-19 during pregnancy also has been

associated with an increased risk of neurodevelopmental disorders in infants within their first

year of life (8). As such, we evaluated offspring of mock-inoculated or maSCV2-infected dams

at E16 for reduced cortical thickness at postnatal day (PND) 0 and delayed neurobehavioral

function at PND 5. Offspring of E16-infected dams had significant cortical thinning in

comparison to offspring from mock-inoculated dams (**Figure 5A-B**), consistent with their

reduced head diameter (**Figure 4E**). Offspring of E16-infected dams displayed delayed surface

righting (**Figure 4C**), cliff aversion (**Figure 4D**), and negative geotaxis (**Figure 4E**) as compared

with offspring from mock-infected dams. Male offspring were more affected by maternal

infection at E16 than female offspring, consistent with literature indicating that males are more

severely impacted by *in utero* insults (52, 53). Offspring of dams that were either maSCV2- or

mock-infected at E6 were also subjected to neurobehavioral testing, and no effect of either

maternal infection or sex of offspring was observed (**Supplemental Figure 4**). These data

highlight that infection with maSCV2 during the third trimester-equivalent of pregnancy causes

both short and long-term adverse fetal outcomes, in the absence of vertical transmission and

consistent with human literature (2, 8).

3.6 Ritonavir-boosted nirmatrelvir treatment prevents morbidity and reduces pulmonary viral

titers following SARS-CoV-2 infection late in gestation

Because of the increased risk of severe COVID-19 and adverse fetal outcomes,

pregnant individuals are recommended to receive the antiviral ritonavir-boosted nirmatrelvir (18,

19). There is, however, limited data on its efficacy during pregnancy, with human and animal

studies primarily focused on evaluating safety and toxicity (23, 26). Additionally, studies

evaluating nirmatrelvir's efficacy in nonpregnant animals utilized high doses of nirmatrelvir alone

in lieu of boosting with ritonavir (33, 54). To better reflect the doses administered to pregnant

individuals, we first evaluated the efficacy of nirmatrelvir and ritonavir at doses calculated to be

the mouse equivalent to a human doses (31) in nonpregnant females compared to high dose

nirmatrelvir alone. Mouse equivalent dosing of nirmatrelvir and ritonavir was equivalent to high

dose nirmatrelvir alone at preventing maSCV2 induced morbidity (**Supplemental Figure 5A**)

and reducing pulmonary viral loads (**Supplemental Figure 5B**) in nonpregnant females. As

CYP3A enzymes are responsible for the metabolism of nirmatrelvir (55), we evaluated liver

CYP3A in pregnant dams at E16 and age-matched non-pregnant females and found no

difference in total expression (**Figure 6A**), further supporting the use of mouse-equivalent doses

of nirmatrelvir and ritonavir during pregnancy.

To evaluate the efficacy of ritonavir-boosted nirmatrelvir during pregnancy, we treated

maSCV2 and mock-infected dams twice daily with mouse-equivalent doses of nirmatrelvir and

ritonavir or vehicle for 5 days (32), starting at 4 hours after infection. maSCV2-infected dams

treated with vehicle failed to gain mass during the remainder of pregnancy and had reduced

mass compared to mock-inoculated dams through lactation (**Figure 6B**). In contrast, treatment

of maSCV2-infected dams with ritonavir-boosted nirmatrelvir prevented maternal morbidity and

resulted in morbidity curve AUCs that were equivalent to those of mock-inoculated dams

(**Figure 6B**). Ritonavir-boosted nirmatrelvir did not significantly reduce infectious viral loads in

the nasal turbinates of pregnant or nonpregnant females (**Figure 6C**). In the lungs, however,

ritonavir-boosted nirmatrelvir reduced viral loads in pregnant, but not nonpregnant, females

compared to vehicle-treated comparators, likely because infected vehicle-treated nonpregnant

females already had lower viral loads than infected pregnant vehicle-treated dams (**Figure 6D**).

We next determined if treatment with ritonavir-boosted nirmatrelvir selected for mutations in the

coding region corresponding to the gene that encodes for the SARS-CoV-2 MPRO protease.

The sequences encoding M^{PRO} did not differ between viral RNA obtained from ritonavir-boosted

nirmatrelvir treated mice and vehicle treated mice, regardless of either pregnancy status or

tissue type (**Figure 6E**, **Supplemental Figure 6**), suggesting that ritonavir-boosted nirmatrelvir

is not selecting for mutations that would potentially reduce its efficacy, at least by 3 dpi.

3.7 Ritonavir-boosted nirmatrelvir treatment prevents adverse offspring outcomes induced by

SARS-CoV-2 infection late in gestation

To evaluate if ritonavir-boosted nirmatrelvir prevented adverse fetal and offspring

outcomes, offspring of E16-infected and mock-inoculated dams treated with ritonavir-boosted

nirmatrelvir, or vehicle were evaluated at birth and PND5. Offspring of maSCV2-infected dams

treated with vehicle were significantly smaller than offspring of mock-inoculated dams in mass

(**Figure 7A**), length (**Figure 7B**), and head diameter (**Figure 7C**) at birth and demonstrated

significant delays in surface righting (**Figure 7D**), cliff aversion (**Figure 7E**), and negative

geotaxis (**Figure 7F**) at PND5, with greater neurobehavioral delays in males than females

(**Figure 7D-F**). Offspring of maSCV2-infected dams treated with ritonavir-boosted nirmatrelvir,

however, did not differ from offspring of mock-inoculated dams in any size measures at birth

(**Figure 7A-C**) or neurobehaviors at PND5 (**Figure 7D-F**). Ritonavir-boosted nirmatrelvir

treatment prevented maSCV2-induced intrauterine growth restriction and neurobehavioral

deficits in both males and females. Offspring of mock-inoculated dams treated with ritonavir-

boosted nirmatrelvir did not differ from offspring of mock-inoculated dams in any offspring

measure (**Figure 7A-F**), consistent with reproductive studies in rabbits which did not find toxicity

during pregnancy (26) Overall, these findings suggest that treatment with ritonavir-boosted

nirmatrelvir during pregnancy can not only reduce maternal pulmonary viral load, but prevents

maternal morbidity, and mitigates adverse fetal and offspring outcomes.

4 Discussion:

Animal models of COVID-19 are powerful tools to study pathogenesis, consider risk-

altering conditions such as pregnancy, and evaluate therapeutic interventions (56, 57). In the

current study, we established a mouse model of SARS-CoV-2 infection during pregnancy that

recapitulates many of the clinical findings of COVID-19 during human pregnancy. Pregnant

dams infected with maSCV2 in late gestation experienced the most severe disease, exhibiting

reduced pulmonary function and increased viral titers, while their offspring were small for

gestational age and had neurodevelopmental delays. These findings are consistent with

observations in humans where pregnant individuals with COVID-19, especially in mid to late

gestation, have greater risk of severe disease, resulting in increased hospitalization and critical

care admission (3), as well as adverse fetal outcomes, such as small size for gestational age at

birth, and neurodevelopmental deficits in the first year of life (7, 8, 58).

At the maternal-fetal interface, intranasal maSCV2 infection resulted in placental

alterations without direct virus infection, which is in accordance with the hallmarks of placental

damage, inflammation, and maternal immune cell infiltration observed in placentas from

mothers with COVID-19 during pregnancy (46-48). After characterizing the negative outcomes

of maSCV2 infection in pregnancy, we used our model to assess the efficacy of ritonavir-

boosted nirmatrelvir at a mouse-equivalent dose to what pregnant humans receive. This

antiviral regimen was well tolerated by pregnant dams, reduced pulmonary virus titers, mitigated

maternal morbidity, and prevented adverse offspring outcomes. Observational studies in human

pregnancies indicate that ritonavir boosted-nirmatrelvir does not pose safety or toxicity risk to

pregnant individuals (23), and may reduce COVID-19 symptoms without requiring additional

medical interventions (24).

In addition to recapitulating aspects of human COVID-19 during pregnancy, our model

identified reduction in pulmonary IFN-β secretion after infection late in gestation and a

corresponding increase in pulmonary viral titer as critical mediators of worse outcomes in late

compared with early gestation. As deficits in type 1 IFN signaling have been associated with

severe COVID-19 in both nonpregnant individuals (43) and mice (44), our data suggest that

maternal morbidity may, in part, be due to an inability of pregnant dams to control viral

replication because of a reduced type I IFN responses, particularly during late gestation. This

potential mechanism of severe disease is consistent with immunological alterations of mouse

and human pregnancy where the maternal immune response shifts to an anti-inflammatory

profile to support the semi-allogenic fetus and diverts from anti-viral and cytotoxic activity (4).

Moreover, in human pregnancy, there is a documented decline in early anti-viral effector cells

and products including natural killer cells and type I IFNs (45).

The adverse maternal and fetal outcomes of SARS-CoV-2 infection during pregnancy

are like those observed in other mouse models of viral pathogenesis during pregnancy including

Zika virus (ZIKV) and influenza A virus (IAV) infection. Mouse models of ZIKV infection during

pregnancy have shown that adverse fetal and neonatal outcomes including congenital

abnormalities, reduced cortical thickness, and neurobehavioral deficits (35, 38), are mediated in

part by transplacental virus transmission and acute placental inflammation (35, 37). While

vertical transmission during ZIKV infection contributes to adverse outcomes, we and others

have shown that the maternal immune response, including elevated production of IL-1β, also

plays a key role in pathogenesis (35, 59). Vertical transmission of virus during COVID-19 is

largely unseen in humans (10, 11, 47), and in mice the placental pathology following maSCV2

infection occurred without vertical transmission. These data further highlight that adverse

neonatal outcomes are not exclusive to vertical transmission of viruses, but by maternal immune

activation and damage at the maternal-fetal interface. Mouse models of IAV infection during

pregnancy further demonstrate maternal morbidity and mortality which is more severe in

pregnant than nonpregnant animals (50, 60). Reduced type I IFN responses and greater viral

loads in the lungs in pregnant dams late in gestation also have been observed in IAV infection

(61), further supporting that pregnancy-associated suppression of type I IFNs is a mechanism of

severe maternal disease after respiratory virus infection.

Mouse models of viral infection during pregnancy are a valuable tool to assess the

safety and efficacy of therapeutics to prevent adverse maternal and fetal outcomes. Our results

support the efficacy of ritonavir-boosted nirmatrelvir for COVID-19 during pregnancy. While

human studies of ritonavir-boosted nirmatrelvir during pregnancy are still needed, these findings

provide a foundation for future human clinical trial design to include pregnant patients. Current

approaches to assessing antiviral therapeutics in preclinical animal models include reproductive

toxicity studies using supraphysiological doses but neglect to evaluate if pregnancy alters

efficacy (26, 62, 63). Therefore, future preclinical models of antiviral therapies must be designed

carefully to consider the complex interactions between pregnancy, viral pathogenesis, and drug

pharmacokinetics. Mouse models of ZIKV antiviral treatment during pregnancy have illustrated

the ability of maternal antiviral treatment to prevent vertical transmission to fetuses (64), a major

adverse outcome associated with ZIKV infection during pregnancy. Pregnant individuals are

largely excluded from clinical trials (14), which has contributed to a reduced uptake of antivirals

and vaccines in pregnant populations, including COVID-19 therapeutics (65, 66). This exclusion

is concerning because pregnant individuals and their neonates are highly vulnerable to many

pathogens (67, 68). With further development of mouse models of viral infection and novel

therapeutics in pregnancy, however, preclinical studies can guide clinical trial design and

promote the inclusion of pregnant populations. By considering pregnancy in clinical trials,

access and uptake of protective therapeutics during pregnancy can be improved.

5 Acknowledgements:

The authors would like to thank Dr. Ralph Baric as well as the Klein, Pekosz, Davis, and

Baumgarth laboratories for discussions about these data, and Ariana Campbell for early

assistance with animal studies. We would also like to thank Dr. Jason Villano and the expert

animal care staff at the Johns Hopkins School of Medicine for assistance with maintenance of

SARS-CoV-2-infected dams.

6 Funding:

Funding provided by NIH/NICHD R01HD097608 (I.B. and S.L.K), NIH/NIAID training grant

T32AI007417-26 (P.C.), and NIAID N7593021C00045 (A.P.).

7 Contributions:

SK, AP, IB, PC, and JP conceptualized and designed the experiments. PC and JP performed

animal experiments. AP, WZ, and RZ grew and quantified viruses and homogenized and

inactivated tissue. PC and JP performed ELISA and Western blot. KM imaged and scored lung

histology slides. PC, JP, HR, and WM performed DFco analysis. JL and AL stained, imaged,

and analyzed placental and fetal head tissue. EA analyzed sequencing data. PC and JP

statistically analyzed and graphed data. PC, JP, and SK wrote the manuscript with input from all

authors. All authors read and provided edits to drafts and approved the final submission. The

order of authors was determined based on contributions to the overall design, experimentation,

analyses, and writing.

8 References:

1.   Lokken EM, Huebner EM, Taylor GG, Hendrickson S, Vanderhoeven J, Kachikis A, et al.
Disease severity, pregnancy outcomes, and maternal deaths among pregnant patients

with severe acute respiratory syndrome coronavirus 2 infection in Washington State.
Am J Obstet Gynecol. 2021;225(1):77 e1- e14.

2.   Woodworth KR, Olsen EO, Neelam V, Lewis EL, Galang RR, Oduyebo T, et al. Birth and
Infant Outcomes Following Laboratory-Confirmed SARS-CoV-2 Infection in Pregnancy -
SET-NET, 16 Jurisdictions, March 29-October 14, 2020. *MMWR Morb Mortal Wkly Rep.*
2020;69(44):1635-40.

3.   Stock SJ, Carruthers J, Calvert C, Denny C, Donaghy J, Goulding A, et al. SARS-CoV-2
infection and COVID-19 vaccination rates in pregnant women in Scotland. *Nat Med.*
2022;28(3):504-12.

4.   Mor G, Cardenas I, Abrahams V, and Guller S. Inflammation and pregnancy: the role of
the immune system at the implantation site. *Ann N Y Acad Sci.* 2011;1221:80-7.

5.   Abu-Raya B, Michalski C, Sadarangani M, and Lavoie PM. Maternal Immunological
Adaptation During Normal Pregnancy. *Front Immunol.* 2020;11:575197.

6.   Yu W, Hu X, and Cao B. Viral Infections During Pregnancy: The Big Challenge Threatening
Maternal and Fetal Health. *Matern Fetal Med.* 2022;4(1):72-86.

7.   Piekos SN, Price ND, Hood L, and Hadlock JJ. The impact of maternal SARS-CoV-2
infection and COVID-19 vaccination on maternal-fetal outcomes. *Reprod Toxicol.*
2022;114:33-43.

8.   Edlow AG, Castro VM, Shook LL, Kaimal AJ, and Perlis RH. Neurodevelopmental
Outcomes at 1 Year in Infants of Mothers Who Tested Positive for SARS-CoV-2 During
Pregnancy. *JAMA Netw Open.* 2022;5(6):e2215787.

9.   Piekos SN, Roper RT, Hwang YM, Sorensen T, Price ND, Hood L, et al. The effect of
maternal SARS-CoV-2 infection timing on birth outcomes: a retrospective multicentre
cohort study. *Lancet Digit Health.* 2022;4(2):e95-e104.

10.   Simbar M, Nazarpour S, and Sheidaei A. Evaluation of pregnancy outcomes in mothers
with COVID-19 infection: a systematic review and meta-analysis. *J Obstet Gynaecol.*
2023;43(1):2162867.

11.   Vigil-Vazquez S, Carrasco-Garcia I, Hernanz-Lobo A, Manzanares A, Perez-Perez A,
Toledano-Revenga J, et al. Impact of Gestational COVID-19 on Neonatal Outcomes: Is
Vertical Infection Possible? *Pediatr Infect Dis J.* 2022;41(6):466-72.

12.   Kim B, Park KH, Lee OH, Lee G, Kim H, Lee S, et al. Effect of severe acute respiratory
syndrome coronavirus 2 infection during pregnancy in K18-hACE2 transgenic mice. *Anim*
Biosci. 2023;36(1):43-52.

13.   Vermillion MS, and Klein SL. Pregnancy and infection: using disease pathogenesis to
inform vaccine strategy. *NPJ Vaccines.* 2018;3:6.

14.   Klein SL, Creisher PS, and Burd I. COVID-19 vaccine testing in pregnant females is
necessary. *J Clin Invest.* 2021;131(5).

15.   LipKind HS V-BG, DeSilva M, et al. . Receipt of COVID-19 Vaccine During Pregnancy and
Preterm of Small-for-Gestationa-Age at Birth- Eight Integrated Health Care
Organizations, United States, December 15, 2020-July 22, 2021. *MMWR Morb Mortal*
Wkly Rep. 2022;71:26-30.

16.   Watanabe A, Yasuhara J, Iwagami M, Miyamoto Y, Yamada Y, Suzuki Y, et al. Peripartum
Outcomes Associated With COVID-19 Vaccination During Pregnancy: A Systematic
Review and Meta-analysis. *JAMA Pediatr.* 2022;176(11):1098-106.

17.   Villar J, Soto Conti CP, Gunier RB, Ariff S, Craik R, Cavoretto PI, et al. Pregnancy
outcomes and vaccine effectiveness during the period of omicron as the variant of
concern, INTERCOVID-2022: a multinational, observational study. *Lancet.*
2023;401(10375):447-57.
18.   NIH. COVID-19 Treatment Guidelines.
https://www.covid19treatmentguidelines.nih.gov/therapies/antiviral-therapy/ritonavir-
boosted-nirmatrelvir--paxlovid-/. Accessed July 14th 2022.
19.   Arco-Torres A, Cortes-Martin J, Tovar-Galvez MI, Montiel-Troya M, Riquelme-Gallego B,
and Rodriguez-Blanque R. Pharmacological Treatments against COVID-19 in Pregnant
Women. *J Clin Med.* 2021;10(21).
20.   Burwick RM, Yawetz S, Stephenson KE, Collier AY, Sen P, Blackburn BG, et al.
Compassionate Use of Remdesivir in Pregnant Women With Severe Coronavirus Disease
2019. *Clin Infect Dis.* 2021;73(11):e3996-e4004.
21.   Hsu A, Granneman GR, and Bertz RJ. Ritonavir. Clinical pharmacokinetics and
interactions with other anti-HIV agents. *Clin Pharmacokinet.* 1998;35(4):275-91.
22.   Society for Maternal Fetal Medicine. FDA Issues EUA for the Treatment of Mild-to-
Moderate COVID-19
Maternal-Fetal Medicine Subspecialists Support Use in Pregnant Patients.
https://s3.amazonaws.com/cdn.smfm.org/media/3287/Treatment_1.10.pdf. Accessed
July 14th 2022.
23.   William M. Garneau MD MPH KJ-BD, MSN CNM,2 Michelle O. Ufua BS,2 Heba H.
Mostafa MBBCh PhD,3 Sabra L. Klein PhD,4, 5 Irina Burd MD PhD, *2 Kelly A. Gebo MD
MPH*. Clinical outcomes of pregnant patients treated with Nirmatrelvir/Ritonavir for
acute COVID-19: a case series of twenty-five patients from the Johns Hopkins CROWN
registry. *JAMA Open.* 2022;in press.
24.   Loza A, Farias R, Gavin N, Wagner R, Hammer E, and Shields A. Short-term Pregnancy
Outcomes After Nirmatrelvir-Ritonavir Treatment for Mild-to-Moderate Coronavirus
Disease 2019 (COVID-19). *Obstet Gynecol.* 2022;140(3):447-9.
25.   Lin K, Liu M, Bao L, Lv Q, Zhu H, Li D, et al. Safety and protective capability of an
inactivated SARS-CoV-2 vaccine on pregnancy, lactation and the growth of offspring in
hACE2 mice. *Vaccine.* 2022;40(32):4609-16.
26.   Catlin NR, Bowman CJ, Campion SN, Cheung JR, Nowland WS, Sathish JG, et al.
Reproductive and developmental safety of nirmatrelvir (PF-07321332), an oral SARS-
CoV-2 M(pro) inhibitor in animal models. *Reprod Toxicol.* 2022;108:56-61.
27.   Leist SR, Dinnon KH, 3rd, Schafer A, Tse LV, Okuda K, Hou YJ, et al. A Mouse-Adapted
SARS-CoV-2 Induces Acute Lung Injury and Mortality in Standard Laboratory Mice. *Cell.*
2020;183(4):1070-85 e12.
28.   Creisher PS, Campbell AD, Perry JL, Roznik K, Burd I, and Klein SL. Influenza subtype-
specific maternal antibodies protect offspring against infection but inhibit vaccine-
induced immunity and protection in mice. *Vaccine.* 2022;40(47):6818-29.
29.   Vom Steeg LG, Dhakal S, Woldetsadik YA, Park HS, Mulka KR, Reilly EC, et al. Androgen
receptor signaling in the lungs mitigates inflammation and improves the outcome of
influenza in mice. *PLoS Pathog.* 2020;16(7):e1008506.

30.   Vom Steeg LG, Vermillion MS, Hall OJ, Alam O, McFarland R, Chen H, et al. Age and
676   testosterone mediate influenza pathogenesis in male mice. *Am J Physiol Lung Cell Mol*
677   *Physiol.* 2016;311(6):L1234-L44.
678   31.   Nair AB, and Jacob S. A simple practice guide for dose conversion between animals and
679   human. *J Basic Clin Pharm.* 2016;7(2):27-31.
680   32.   FDA. 2021.
681   33.   Owen DR, Allerton CMN, Anderson AS, Aschenbrenner L, Avery M, Berritt S, et al. An
682   oral SARS-CoV-2 M(pro) inhibitor clinical candidate for the treatment of COVID-19.
683   *Science.* 2021;374(6575):1586-93.
684   34.   Feather-Schussler DN, and Ferguson TS. A Battery of Motor Tests in a Neonatal Mouse
685   Model of Cerebral Palsy. *J Vis Exp.* 2016(117).
686   35.   Lei J, Vermillion MS, Jia B, Xie H, Xie L, McLane MW, et al. IL-1 receptor antagonist
687   therapy mitigates placental dysfunction and perinatal injury following Zika virus
688   infection. *JCI Insight.* 2019;4(7).
689   36.   Limjunyawong N, Fallica J, Ramakrishnan A, Datta K, Gabrielson M, Horton M, et al.
690   Phenotyping mouse pulmonary function in vivo with the lung diffusing capacity. *J Vis*
691   *Exp.* 2015(95):e52216.
692   37.   Creisher PS, Lei J, Sherer ML, Dziedzic A, Jedlicka AE, Narasimhan H, et al.
693   Downregulation of transcriptional activity, increased inflammation, and damage in the
694   placenta following in utero Zika virus infection is associated with adverse pregnancy
695   outcomes. *Front Virol.* 2022;2.
696   38.   Vermillion MS, Lei J, Shabi Y, Baxter VK, Crilly NP, McLane M, et al. Intrauterine Zika
697   virus infection of pregnant immunocompetent mice models transplacental transmission
698   and adverse perinatal outcomes. *Nat Commun.* 2017;8:14575.
699   39.   Meyerholz DK, and Beck AP. Histopathologic Evaluation and Scoring of Viral Lung
700   Infection. *Methods Mol Biol.* 2020;2099:205-20.
701   40.   Meyerholz DK, Sieren JC, Beck AP, and Flaherty HA. Approaches to Evaluate Lung
702   Inflammation in Translational Research. *Vet Pathol.* 2018;55(1):42-52.
703   41.   Armando F, Beythien G, Kaiser FK, Allnoch L, Heydemann L, Rosiak M, et al. SARS-CoV-2
704   Omicron variant causes mild pathology in the upper and lower respiratory tract of
705   hamsters. *Nat Commun.* 2022;13(1):3519.
706   42.   Sones JL, and Davisson RL. Preeclampsia, of mice and women. *Physiol Genomics.*
707   2016;48(8):565-72.
708   43.   Hadjadj J, Yatim N, Barnabei L, Corneau A, Boussier J, Smith N, et al. Impaired type I
709   interferon activity and inflammatory responses in severe COVID-19 patients. *Science.*
710   2020;369(6504):718-24.
711   44.   Ogger PP, Garcia Martin M, Michalaki C, Zhou J, Brown JC, Du Y, et al. Type I interferon
712   receptor signalling deficiency results in dysregulated innate immune responses to SARS-
713   CoV-2 in mice. *Eur J Immunol.* 2022;52(11):1768-75.
714   45.   Alberca RW, Pereira NZ, Oliveira L, Gozzi-Silva SC, and Sato MN. Pregnancy, Viral
715   Infection, and COVID-19. *Front Immunol.* 2020;11:1672.
716   46.   Rad HS, Rohl J, Stylianou N, Allenby MC, Bazaz SR, Warkiani ME, et al. The Effects of
717   COVID-19 on the Placenta During Pregnancy. *Front Immunol.* 2021;12:743022.

718   47.   Argueta LB, Lacko LA, Bram Y, Tada T, Carrau L, Rendeiro AF, et al. Inflammatory
719   responses in the placenta upon SARS-CoV-2 infection late in pregnancy. *iScience.*
720   2022;25(5):104223.
721   48.   Huynh A, Sehn JK, Goldfarb IT, Watkins J, Torous V, Heerema-McKenney A, et al. SARS-
722   CoV-2 Placentitis and Intraparenchymal Thrombohematomas Among COVID-19
723   Infections in Pregnancy. *JAMA Netw Open.* 2022;5(3):e225345.
724   49.   Elmore SA, Cochran RZ, Bolon B, Lubeck B, Mahler B, Sabio D, et al. Histology Atlas of
725   the Developing Mouse Placenta. *Toxicol Pathol.* 2022;50(1):60-117.
726   50.   Kim HM, Kang YM, Song BM, Kim HS, and Seo SH. The 2009 pandemic H1N1 influenza
727   virus is more pathogenic in pregnant mice than seasonal H1N1 influenza virus. *Viral
728   Immunol.* 2012;25(5):402-10.
729   51.   Elovitz MA, Wang Z, Chien EK, Rychlik DF, and Phillippe M. A new model for
730   inflammation-induced preterm birth: the role of platelet-activating factor and Toll-like
731   receptor-4. *Am J Pathol.* 2003;163(5):2103-11.
732   52.   Hunter SK, Hoffman MC, D'Alessandro A, Noonan K, Wyrwa A, Freedman R, et al. Male
733   fetus susceptibility to maternal inflammation: C-reactive protein and brain
734   development. *Psychol Med.* 2021;51(3):450-9.
735   53.   Sutherland S, and Brunwasser SM. Sex Differences in Vulnerability to Prenatal Stress: a
736   Review of the Recent Literature. *Curr Psychiatry Rep.* 2018;20(11):102.
737   54.   Uraki R, Kiso M, Iida S, Imai M, Takashita E, Kuroda M, et al. Characterization and
738   antiviral susceptibility of SARS-CoV-2 Omicron BA.2. *Nature.* 2022;607(7917):119-27.
739   55.   Costantine MM. Physiologic and pharmacokinetic changes in pregnancy. *Front
740   Pharmacol.* 2014;5:65.
741   56.   Munoz-Fontela C, Widerspick L, Albrecht RA, Beer M, Carroll MW, de Wit E, et al.
742   Advances and gaps in SARS-CoV-2 infection models. *PLoS Pathog.* 2022;18(1):e1010161.
743   57.   Chu H, Chan JF, and Yuen KY. Animal models in SARS-CoV-2 research. *Nat Methods.*
744   2022;19(4):392-4.
745   58.   Fallach N, Segal Y, Agassy J, Perez G, Peretz A, Chodick G, et al. Pregnancy outcomes
746   after SARS-CoV-2 infection by trimester: A large, population-based cohort study. *PLoS
747   One.* 2022;17(7):e0270893.
748   59.   Casazza RL, Philip DT, and Lazear HM. Interferon Lambda Signals in Maternal Tissues to
749   Exert Protective and Pathogenic Effects in a Gestational Stage-Dependent Manner.
750   *mBio.* 2022;13(3):e0385721.
751   60.   Littauer EQ, Esser ES, Antao OQ, Vassilieva EV, Compans RW, and Skountzou I. H1N1
752   influenza virus infection results in adverse pregnancy outcomes by disrupting tissue-
753   specific hormonal regulation. *PLoS Pathog.* 2017;13(11):e1006757.
754   61.   Engels G, Hierweger AM, Hoffmann J, Thieme R, Thiele S, Bertram S, et al. Pregnancy-
755   Related Immune Adaptation Promotes the Emergence of Highly Virulent H1N1 Influenza
756   Virus Strains in Allogenically Pregnant Mice. *Cell Host Microbe.* 2017;21(3):321-33.
757   62.   Kala S, Watson B, Zhang JG, Papp E, Guzman Lenis M, Dennehy M, et al. Improving the
758   clinical relevance of a mouse pregnancy model of antiretroviral toxicity; a
759   pharmacokinetic dosing-optimization study of current HIV antiretroviral regimens.
760   *Antiviral Res.* 2018;159:45-54.

761 63.   Donner B, Niranjan V, and Hoffmann G. Safety of oseltamivir in pregnancy: a review of
762   preclinical and clinical data. *Drug Saf.* 2010;33(8):631-42.
763 64.   Watanabe S, Tan NWW, Chan KWK, and Vasudevan SG. Assessing the utility of antivirals
764   for preventing maternal-fetal transmission of zika virus in pregnant mice. *Antiviral Res.*
765   2019;167:104-9.
766 65.   Sebghati M, and Khalil A. Uptake of vaccination in pregnancy. *Best Pract Res Clin Obstet*
767   *Gynaecol.* 2021;76:53-65.
768 66.   Firouzbakht M, Sharif Nia H, Kazeminavaei F, and Rashidian P. Hesitancy about COVID-
769   19 vaccination among pregnant women: a cross-sectional study based on the health
770   belief model. *BMC Pregnancy Childbirth.* 2022;22(1):611.
771 67.   Kourtis AP, Read JS, and Jamieson DJ. Pregnancy and infection. *N Engl J Med.*
772   2014;370(23):2211-8.
773 68.   Neu N, Duchon J, and Zachariah P. TORCH infections. *Clin Perinatol.* 2015;42(1):77-103,
774   viii.

Figure Legends

Figure 1. maSCV2 infection of pregnant dams results in gestation-dependent morbidity.

Nonpregnant adult females (A) or dams at embryonic day (E) 6 (B), E10 (C), and E16 (D) were

intranasally inoculated with 10^5 TCID$_{50}$ of a mouse adapted SARS-CoV-2 (maSCV2) or mock

inoculated with media. Following infection, mice were monitored for change in body mass and

clinical signs of disease over fourteen days (A-D). Area under the curve (AUC) of body mass

change curves for infected and uninfected animals were calculated, and then the AUC of

infected animals was subtracted from the average AUC of mock animals of the same

reproductive status and gestational age (E). Clinical scores given to animals included dyspnea,

piloerection, hunched posture, and absence of an escape response and are quantified on a

score of 0-4. The cumulative clinical score over the 14-day monitoring period is reported for

each animal (F). Individual shapes (A-D) or bars (E-F) represent the mean (A-E) or median (F) ±

standard error of the mean (A-E) from two independent replications (n =7-13/group) with

individual mice indicated by shapes (E-F). Significant differences ($p < 0.05$) were determined by

two tailed unpaired t-test of AUCs (A-F), one-way ANOVA with Bonferroni post hoc test (E), or

Kruskal-Wallis test (F) and are indicated by an asterisk (*).

Figure 2. Pregnant dams infected during the third trimester-equivalent have reduced IFN-

β responses, increased viral load, and reduced pulmonary function after infection.

Nonpregnant adult females or dams at embryonic day (E) 6 and E16 were intranasally

inoculated with 10^5 TCID$_{50}$ maSCV2 or mock inoculated with media and euthanized at three

days post infection (DPI) to collect maternal and fetal tissues. IFNβ and viral titers in the right

cranial lungs were measured using ELISA (A) and TCID$_{50}$ assay (B), respectively. Sections of

fixed left lungs were stained by hematoxylin & eosin (H&E) to evaluate lung inflammation and

images were taken at 20x magnification, with representative images of lungs maSCV2 or mock-

inoculated at E16 shown (C). Asterisks (*) indicate intra-alveolar necrosis and inflammatory

infiltrates, and arrows indicate peribronchiolar inflammatory infiltrates. Histopathological scoring

was performed by a board-certified veterinary pathologist to measure cumulative inflammation

scores (D). A subset of mice were tracheostomized at three DPI to measure pulmonary function

through the diffusion capacity for carbon monoxide (DF$_{CO}$) prior to euthanasia (E). Bars

represent the mean (A-E) ± standard error of the mean from at least two independent

replications (n =5-11/group) with individual mice indicated by shapes. Significant differences (*p*

< 0.05) were determined by two-way ANOVA with Bonferroni post hoc test (A,D,E) or one-way

ANOVA with Bonferroni post hoc test (B) and are indicated by an asterisk (*). Scale bar: 100

μm. LOD indicates the limit of detection.

Figure 3. Third trimester-equivalent maSCV2 infection induces congestion and disrupts

the trophoblast layer of the placental labyrinth zone.

At embryonic day (E) 16, pregnant dams were intranasally inoculated with 10^5 TCID$_{50}$ of

maSCV2 or mock inoculated with media and euthanized at 3 dpi to collect placentas.

Representative H&E images were taken at 20x magnification (A). Within H&E stained placentas,

arrows indicate trophoblast giant cells and asterisks (*) indicate congestion. Placentas were

immunostained for cytokeratin (B, red) to mark trophoblasts or vimentin (C, red) to mark

endothelial cells and DAPI (blue) to label nuclei. Representative images were taken at 20x

magnification. Quantification of the percentage positive area for each marker is shown. Bars

represent the mean ± standard error of the mean (n =5/group) with each shape indicating 1

placenta and is the mean quantification of 6 fields of view. Significant differences ($p < 0.05$)

were determined by unpaired two tailed t-test and are indicated by an asterisk (*). Scale bar:

100 μm.

Figure 4. Third trimester-equivalent maSCV2 infection causes intrauterine growth

restriction.

At embryonic day (E) 6, E10, or E16, pregnant dams were intranasally inoculated with 10^5

$TCID_{50}$ of maSCV2 or mock inoculated with media. At 3 dpi, a subset of dams were euthanized,

and fetal viability was determined as the percentage of fetuses within the uterus (A, n= total

number of fetuses from 8-12 dams per group from two independent replicates). A subset of

dams were followed into the postnatal period to characterize adverse birth outcomes. At

postnatal day 0 (PND0) overall litter size (B), pup mass (C), pup body length (D), and pup head

diameter (E) were measured. Average measurements of each independent litter were graphed

to account for litter effects (B-E). Bars represent the mean ± standard error of the mean from

two independent replicates (n =5-14/group) with the average of individual litters indicated by

shapes. Significant differences ($p < 0.05$) were determined by χ2 (A) or two-way ANOVA with

Bonferroni post hoc test (B-E) and are indicated by an asterisk (*).

Figure 5. Offspring of dams infected with maSCV2 during the third trimester-equivalent

display cortical thinning and reduced neurodevelopmental function.

At embryonic day (E) 16, pregnant dams were intranasally inoculated with 10^5 $TCID_{50}$ of

maSCV2 or mock inoculated with media. At PND0, a randomly selected subset of pups were

euthanized via decapitation to collect fetal heads, which were fixed, sliced, and Nissl stained.

Cortical thickness (A, red arrows) was measured from both brain hemispheres per pup and

quantified as the average of 10 measurements per pup, with a single pup randomly chosen per

dam (B, n=9-10 independent litters/group). A subset of offspring were followed to PND5, sexed,

and the neurobehavioral assays of surface righting (C), cliff aversion (D), and negative geotaxis

(E) were performed to measure neurological development. 1-2 pups per sex per dam were

subjected to each test subsequently, with 3 trials given per test, and each pup's best trial for

each test was reported (C-E, n=9-10 independent litters/group). Bars represent the mean ±

standard error of the mean with each shape indicating 1 pup. Significant differences ($p < 0.05$)

were determined by unpaired two tailed t-test (B) or two-way ANOVA with Bonferroni post hoc

test (C-E) and are indicated by an asterisk (*). Graphics built with Biorender.com

Figure 6. Ritonavir-boosted nirmatrelvir mitigates maternal morbidity and reduced viral

titers in the lungs of pregnant dams.

Uninfected adult nonpregnant and pregnant (i.e., embryonic day (E)16) females were

euthanized, liver tissue collected, and western blots performed to quantify the amount of overall

CYP3A expression (A, n=4-5/group). At E16, pregnant dams or age-matched nonpregnant

females were intranasally infected with 10^5 TCID$_{50}$ of maSCV2 or mock inoculated with media.

Starting at 4 hours post infection and continuing twice daily for 5 days or until tissue collection,

mice were treated with 1.7 mg nirmatrelvir and 0.6 mg ritonavir per dose or vehicle and were

monitored for changes in body mass for fourteen days (B, n=6/group). A subset of dams were

euthanized at 3 dpi, and nasal turbinate and lung tissue were collected, and viral titers were

measured by TCID$_{50}$ assay (C-D, n=5-11/group). RNA was extracted from lung homogenate,

reverse transcribed using ProtoScript® II First Strand cDNA Synthesis Kit, the M^{pro} region

amplified, and Oxford Nanopore sequenced by Plasmidaurus. Consensus sequences were

imported and aligned to Mpro using ClustalO v1.2.3 in Geneious Prime v2023.0.4. Alignments

were imported into R v4.1.1., visualized, and annotated using seqvisR v0.2.5 (E, n=4/group).

Bars represent the mean ± standard error of the mean from two independent replications with

individual mice indicated by shapes (A,C,D). Significant differences ($p < 0.05$) were determined

by two tailed unpaired t-test (A), two-way ANOVA with Bonferroni post hoc test of AUCs (B), or

two-way ANOVA with Bonferroni post hoc test (C-D) and are indicated by an asterisk (*).

Sequence graphic built using Biorender.com. LOD indicates the limit of detection.

Figure 7. Ritonavir-boosted nirmatrelvir prevents adverse offspring birth outcomes and

neurodevelopmental deficits associated with maternal maSCV2 infection.

At embryonic day (E)16, pregnant dams were intranasally inoculated with 10^5 TCID$_{50}$ of

maSCV2 or mock inoculated with media. Starting at 4 hours post infection and continuing twice

daily for 5 days or until tissue collection, mice were treated with 1.7 mg nirmatrelvir and 0.6 mg

ritonavir per dose or vehicle. At PND0, a subset of pups were measured for pup mass (A), pup

length (B), and pup head diameter (C). Average measurements of each litter were graphed to

account for litter effects (A-C, n=6 independent litters/group). A subset of offspring were

followed to PND5, sexed, and the neurobehavioral assays of surface righting (D), cliff aversion

(E), and negative geotaxis (F) were performed to measure neurological development. 1-2 pups

per sex per dam were subjected to each test subsequently, with 3 trials given per test, and each

pup's best trial for each test was reported (D-F, n=6-8 independent litters/group). Bars represent

the mean ± standard error of the mean with each shape indicating 1 pup. Significant differences

($p < 0.05$) were determined by two-way ANOVA with Bonferroni post hoc test (A-C) or three-way

ANOVA with Bonferroni post hoc test (D-F) and are indicated by an asterisk (*).

Figure 1.

maSCV2 infection of pregnant dams results in gestation-dependent morbidity.
Nonpregnant adult females (A) or dams at embryonic day (E)6 (B), E10 (C), and E16 (D) were intranasally inoculated with 10^5 $TCID_{50}$ of a mouse adapted SARS-CoV-2 (maSCV2) or mock inoculated with media. Following infection, mice were monitored for change in body mass and clinical signs of disease over fourteen days (A-D). Area under the curve (AUC) of body mass change curves for infected and uninfected animals were calculated, and then the AUC of infected animals was subtracted from the average AUC of mock animals of the same reproductive status and gestational age (E). Clinical scores given to animals included dyspnea, piloerection, hunched posture, and absence of an escape response and are quantified on a score of 0-4. The cumulative clinical score over the 14-day monitoring period is reported for each animal (F). Individual shapes (A-D) or bars (E-F) represent the mean (A-E) or median (F) ± standard error of the mean (A-E) from two independent replications (n =7-13/group) with individual mice indicated by shapes (E-F). Significant differences ($p < 0.05$) were determined by two tailed unpaired t-test of AUCs (A-F), one-way ANOVA with Bonferroni post hoc test (E), or Kruskal-Wallis test (F) and are indicated by an asterisk (*).

Figure 2.

Pregnant dams infected during the third trimester-equivalent have reduced IFN-β responses, increased viral load, and reduced pulmonary function after infection.

Nonpregnant adult females or dams at embryonic day (E)6 and E16 were intranasally inoculated with 105 TCID50 maSCV2 or mock inoculated with media and euthanized at three days post infection (DPI) to collect maternal and fetal tissues. IFN☐ and viral titers in the right cranial lungs were measured using ELISA (A) and TCID50 assay (B), respectively. Sections of fixed left lungs were stained by hematoxylin & eosin (H&E) to evaluate lung inflammation and images were taken at 20x magnification, with representative images of lungs maSCV2 or mock-inoculated at E16 shown(C). Asterisks (*) indicate intra-alveolar necrosis and inflammatory infiltrates, and arrows indicate peribronchiolar inflammatory infiltrates. Histopathological scoring was performed by a board-certified veterinary pathologist to measure cumulative inflammation scores (D). A subset of mice were tracheostomized at three DPI to measure pulmonary function through the diffusion capacity for carbon monoxide (DFCO) prior to euthanasia (E). Bars represent the mean (A-E) ± standard error of the mean from at least two independent replications (n =5-11/group) with individual mice indicated by shapes. Significant differences (p < 0.05) were determined by two-way ANOVA with Bonferroni post hoc test (A,D,E) or one-way ANOVA with Bonferroni post hoc test (B) and are indicated by an asterisk (*). Scale bar: 100 μm. LOD indicates the limit of detection.

Figure 3.

Third trimester-equivalent maSCV2 infection induces congestion and disrupts the trophoblast layer of the placental labyrinth zone.
At embryonic day (E)16, pregnant dams were intranasally inoculated with 10^5 TCID$_{50}$ of maSCV2 or mock inoculated with media and euthanized at 3 dpi to collect placentas. Representative H&E images were taken at 20x magnification (A). Within H&E stained placentas, arrows indicate trophoblast giant cells and asterisks (*) indicate congestion. Placentas were immunostained for cytokeratin (B, red) to mark trophoblasts or vimentin (C, red) to mark endothelial cells and DAPI (blue) to label nuclei. Representative images were taken at 20x magnification. Quantification of the percentage positive area for each marker is shown. Bars represent the mean ± standard error of the mean (n =5/group) with each shape indicating 1 placenta and is the mean quantification of 6 fields of view. Significant differences ($p < 0.05$) were determined by unpaired two tailed t-test and are indicated by an asterisk (*). Scale bar: 100 µm.

Figure 4.

Third trimester-equivalent maSCV2 infection causes intrauterine growth restriction.
At embryonic day (E)6, E10, or E16, pregnant dams were intranasally inoculated with 10^5 $TCID_{50}$ of maSCV2 or mock inoculated with media. At 3 dpi, a subset of dams were euthanized and fetal viability was determined as the percentage of fetuses within the uterus (A, n= total number of fetuses from 8-12 dams per group from two independent replicates). A subset of dams were followed into the postnatal period to characterize adverse birth outcomes. At postnatal day 0 (PND0) overall litter size (B), pup mass (C), pup body length (D), and pup head diameter (E) were measured. Average measurements of each independent litter were graphed to account for litter effects (B-E). Bars represent the mean ± standard error of the mean from two independent replicates (n =5-14/group) with the average of individual litters indicated by shapes. Significant differences ($p < 0.05$) were determined by χ2 (A) or two-way ANOVA with Bonferroni post hoc test (B-E) and are indicated by an asterisk (*).

Figure 5.

Offspring of dams infected with maSCV2 during the third trimester-equivalent display cortical thinning and reduced neurodevelopmental function.
At embryonic day (E)16, pregnant dams were intranasally inoculated with 10^5 TCID$_{50}$ of maSCV2 or mock inoculated with media. At PND0, a randomly selected subset of pups were euthanized via decapitation to collect fetal heads, which were fixed, sliced, and Nissl stained. Cortical thickness (A, red arrows) was measured from both brain hemispheres per pup and quantified as the average of 10 measurements per pup, with a single pup randomly chosen per dam (B, n=9-10 independent litters/group). A subset of offspring were followed to PND5, sexed, and the neurobehavioral assays of surface righting (C), cliff aversion (D), and negative geotaxis (E) were performed to measure neurological development. 1-2 pups per sex per dam were subjected to each test subsequently, with 3 trials given per test, and each pup's best trial for each test was reported (C-E, n=9-10 independent litters/group). Bars represent the mean ± standard error of the mean with each shape indicating 1 pup. Significant differences ($p < 0.05$) were determined by unpaired two tailed t-test (B) or two-way ANOVA with Bonferroni post hoc test (C-E) and are indicated by an asterisk (*). Graphics built with Biorender.com

Figure 6.

Ritonavir-boosted nirmatrelvir mitigates maternal morbidity and reduced viral titers in the lungs of pregnant dams.

Uninfected adult nonpregnant and pregnant (i.e., embryonic day (E)16) females were euthanized, liver tissue collected, and western blots performed to quantify the amount of overall CYP3A expression (A, n=4-5/group). At E16, pregnant dams or age-matched nonpregnant females were intranasally infected with 10^5 $TCID_{50}$ of maSCV2 or mock inoculated with media. Starting at 4 hours post infection and continuing twice daily for 5 days or until tissue collection, mice were treated with 1.7 mg nirmatrelvir and 0.6 mg ritonavir per dose or vehicle and were monitored for changes in body mass for fourteen days (B, n=6/group). A subset of dams were euthanized at 3 dpi, and nasal turbinate and lung tissue were collected, and viral titers were measured by $TCID_{50}$ assay (C-D, n=5-11/group). RNA was extracted from lung homogenate, reverse transcribed using ProtoScript® II First Strand cDNA Synthesis Kit, the M^{pro} region amplified, and Oxford Nanopore sequenced by Plasmidaurus. Consensus sequences were imported and aligned to Mpro using ClustalO v1.2.3 in Geneious Prime v2023.0.4. Alignments were imported into R v4.1.1., visualized, and annotated using seqvisR v0.2.5 (E, n=4/group). Bars represent the mean ± standard error of the mean from two independent replications with individual mice indicated by shapes (A,C,D). Significant differences ($p < 0.05$) were determined by two tailed unpaired t-test (A), two-way ANOVA with Bonferroni post hoc test of AUCs (B), or two-way ANOVA with Bonferroni post hoc test (C-D) and are indicated by an asterisk (*). Sequence graphic built using Biorender.com. LOD indicates the limit of detection.

Figure 7.

Ritonavir-boosted nirmatrelvir prevents adverse offspring birth outcomes and neurodevelopmental deficits associated with maternal maSCV2 infection.

At embryonic day (E)16, pregnant dams were intranasally inoculated with 10^5 TCID$_{50}$ of maSCV2 or mock inoculated with media. Starting at 4 hours post infection and continuing twice daily for 5 days or until tissue collection, mice were treated with 1.7 mg nirmatrelvir and 0.6 mg ritonavir per dose or vehicle. At PND0, a subset of pups were measured for pup mass (A), pup length (B), and pup head diameter (C). Average measurements of each litter were graphed to account for litter effects (A-C, n=6 independent litters/group). A subset of offspring were followed to PND5, sexed, and the neurobehavioral assays of surface righting (D), cliff aversion (E), and negative geotaxis (F) were performed to measure neurological development. 1-2 pups per sex per dam were subjected to each test subsequently, with 3 trials given per test, and each pup's best trial for each test was reported (D-F, n=6-8 independent litters/group). Bars represent the mean ± standard error of the mean with each shape indicating 1 pup. Significant differences ($p < 0.05$) were determined by two-way ANOVA with Bonferroni post hoc test (A-C) or three-way ANOVA with Bonferroni post hoc test (D-F) and are indicated by an asterisk (*).