Inoviridae **prophage and bacterial host dynamics during diversification, succession and**

Atlantic invasion of Pacific-native *Vibrio parahaemolyticus*.

Randi L. Foxall[a,b]*, Jillian Means[a,c,d]*, Ashley L. Marcinkiewicz[a,c,e]*, Christopher Schillaci[a,f,g],

Kristin DeRosia-Banick[a,h], Feng Xu[a,b,i], Jeffrey A. Hall[a,j], Stephen H. Jones[a,f], Vaughn S.

Cooper[a,b,k], Cheryl A. Whistler[a,b,#]

[a] Northeast Center for Vibrio Disease and Ecology, University of New Hampshire, Durham, New

Hampshire, USA

[b] Department of Molecular, Cellular, and Biomedical Sciences, University of New Hampshire,

USA

[c] Graduate program in Microbiology, University of New Hampshire, Durham, New Hampshire,

USA

[d] Current address: Synlogic Inc., Cambridge, MA, USA

[e] Current address: Division of Infectious Disease, Wadsworth Center, New York State,

Department of Health, Albany, New York, USA

[f] Department of Natural Resources and the Environment, University of New Hampshire,

Durham, New Hampshire, USA

[g] Current address: NOAA Fisheries Service, Greater Atlantic Regional Fisheries Office,

Gloucester Massachusetts

[h] Current address: Milken School of Public Health, George Washington University, Washington

D.C., USA

[i] Current address: Children's Hospital of Philadelphia, Philadelphia, PA, USA

[j] Hubbard Center for Genome Studies, University of New Hampshire, Durham, NH, USA

[k] Department of Microbiology and Molecular Genetics, University of Pittsburgh School of

Medicine, Pittsburgh, PA, USA

[#]Corresponding Author: Cheryl.Whistler@unh.edu

* Contributed equally to this work. Ms. Marcinkiewicz initiated the study and documented the

phage distribution in clinical and environmental samples, associated phage with aquaculture

areas, and conducted initial phylogeographic analyses. Ms. Means extended the research and

assembled inovirus genomes, conducted phylogenetic and bioinformatics analyses of phage,

produced visuals and tables for the manuscript, and completed the first draft of the manuscript.

Ms. Foxall conducted the final genome analyses of both host and inoviruses, completed

population structure analyses, identified inoviruses from public databases, conducted

comprehensive analyses of global patterns of phage association and relatedness, conducted all

analyses requested by reviewers, and produced both visuals and methods for final publication.

Running Title: Inovirus mediated evolution of *Vibrio parahaemolyticus*

Abstract word count: 190

Importance word count: 120

Manuscript word count: 4987

Abstract: The epidemiology of *Vibrio parahaemolyticus,* the leading cause of seafood-borne

bacterial gastroenteritis of humans world-wide, dramatically changed in the United States

following the establishment of a Pacific native lineage called sequence type (ST) 36 in the

Atlantic. In this study we used phylogeography based on traceback to environmental source

locations and comparative genomics to identify features that promoted evolution, dispersal, and

competitive dominance of ST36. The major genomic differentiation and competitive success of

ST36 was associated with a striking succession of filamentous prophage in the family *Inoviridae*

(inoviruses), including loss of an inovirus prophage that had been maintained for decades in the

endemic north Pacific population. Subsequently, at least five distinct progenitors arising from

this diversification translocated from the Pacific into the Atlantic and established four

geographically defined clonal subpopulations with remarkably low migration or mixing. Founders

of two prevailing Atlantic subpopulations each acquired new stable and diagnostic inoviruses

while other subpopulations that apparently declined did not. Broader surveys indicate inoviruses

are common and active among the global population of *V. parahaemolyticus* and though

inovirus replacements, such as in ST36, appear to be infrequent, they are notable in pathogenic

lineages that dispersed.

Importance: An understanding of the processes that contribute to emergence of pathogens

from environmental reservoirs is critical as changing climate precipitates pathogen evolution and

population expansion. Phylogeographic analysis of *Vibrio parahaemolyticus* hosts combined

with analysis of their *Inoviridae* phage resolved ambiguities of diversification dynamics which

preceded successful Atlantic invasion by the epidemiologically predominant ST36 lineage. It has

been established experimentally that filamentous phage can limit host recombination, but here

we show that phage loss is linked to rapid bacterial host diversification during epidemic spread

in natural ecosystems alluding to a potential role for ubiquitous inoviruses in the adaptability of

pathogens. This work paves the way for functional analyses to define the contribution of

inoviruses in the evolutionary dynamics of environmentally transmitted pathogens.

Introduction

Cases of *V. parahaemolyticus*, the leading cause of bacterial seafood-borne

gastroenteritis of humans world-wide recently increased along the United States (US) Atlantic

coast (1-3). Rising seasonal illnesses were tied to warming ocean temperatures, a growing

aquaculture industry, and the emergence of endemic pathogens (3-6). But incursion of a Pacific

Northwest (PNW) lineage called sequence type (ST) 36 into the Atlantic following an

anomalously mild winter is the most important driver of this shift (3, 5, 7). ST36 began causing

infections in 1979 (8) and was originally limited to the Pacific Northwest (PNW) (9). Although

sporadic infections were occasionally reported outside the PNW, local sources were rarely

implicated (7, 9, 10). This changed in 2012 when ST36 began causing illnesses traced to

Atlantic sources and in 2013 it caused a 13-state outbreak traced to multiple northeast US

locations (5, 7). Unlike previous non-endemic strains causing outbreaks from Atlantic sources,

including the pandemic ST3 (11), and ST8 (12), ST36 continues to cause sporadic disease from

a few northeast US locations. A better understanding of how and where ST36 established

populations is needed to aid in management to limit illness. Furthermore, this expansion of

ST36 provides a unique opportunity to broaden our understanding of the population and

environmental context by which pandemic strains arise and spread.

Spatiotemporal analyses of the epidemiology of ST36 identified that a new population

arose through recombination and replaced the original PNW population by 2000 (9). Disease

patterns implied potential reciprocal transfer between both US coasts, though inference was

informed by a few cases where infection occurred from shellfish consumed on the opposite

coast from where illnesses were reported and further complicated by modeling of a mode of

spread through human populations (9), whereas seasonal illnesses by *V. parahaemolyticus* are

typically vectored by environmentally contaminated raw seafood. This highlights a continuing

challenge where a lack of environmental traceback in clinical reporting that uncouples strains

from their source bodies of water obscures environmental distribution (5, 9). As with other

human pathogens originating from the natural environment, this leaves a critical gap of

knowledge of the ecological context of pathogen evolution and expansion.

Bacteriophages, the viruses that infect bacteria, can shape, and quell environmentally

vectored human pathogen populations (13-19). Lytic phages alter the dynamics of bacterial

competition though selective predation, sometimes of the most numerous bacteria, thereby

maintaining population diversity (20-22). Phages also impact diversity through transfer of novel

DNA including toxin-encoding genes and lysogenic conversion (23-28). Chromosomally

integrated prophages are double-edged swords in that they can benefit their host by excluding

superinfections by related phage and by attacking their hosts' competitors, but they can also

harm their hosts by diverting resources for virion production or by host death upon lytic induction

(25-27, 29, 30). Lytic phages are abundant in habitats of *V. parahaemolyticus* (31), and also

under consideration for detection and biological control (32-37). Though the roles of phage in

mediating bacterial competition among *Vibrio cholerae* is well appreciated (14, 38), with the

exception of a few descriptive reports (39-44), the contribution of prophage to *V.*

parahaemolyticus population structure is undetermined.

Prior studies have produced valuable insight into the growing problem of the ST36

lineage's increasing clinical prevalence and potential to spread globally (9, 45), but the

adaptations that prompted lineage replacement and dispersal are not evident nor are the

ecological factors that spurred diversification. Here, we examined the phylogenomics of the

Atlantic ST36 incursion—accompanied by interactions with filamentous *Inoviridae* phage—by

capturing a broad geographic and temporal distribution of the population. By curating strains

with environmental traceback, we determined the phylogeography of ST36 wherein multiple

strains that first diversified in the Pacific, subsequently translocated to the Atlantic, and

established spatially distinct and non-mixing subpopulations therein. This also revealed that

changes in inovirus content accompanied diversification and lineage replacement in the PNW

and preceded clonal expansion by distinct founders of two persisting Atlantic ST36 populations

that continue to cause sporadic disease.

Results

Multiple distinct ST36 lineages clonally expanded to form geographically stable Atlantic

subpopulations

To investigate the evolution of ST36 as it translocated from the PNW to the Atlantic, we

curated a collection of genomes and mapped their environmental sources onto their constructed

whole genome phylogenies. Locations include two Pacific (PNW and California [CA]), and five

Atlantic (Galicia Spain, the Gulf of Maine [GOM], an Island south of Cape Cod, [SCC], Long

Island Sound [LIS] and the mid-Atlantic coast [MAC] [Table S1]). The phylogenies exemplify

how multiple clades emerged from the old-PNW population (Fig. 1) and gave rise to the new-

PNW population (9). A single relic of the old-PNW population and 103 isolates from seven

lineages arising from the modern ST36 diversification were traced to environmental locations

outside the PNW (Fig. 1, clades I-VII). Notably, most translocated clades either contain a PNW-

traced isolate or are most closely related to clades dominated by PNW isolates suggesting

these differentiated in the PNW prior to translocation rather than after arrival in the Atlantic.

Assignment of isolates to environmental sources revealed striking geographic structure

of the Atlantic subpopulations in GOM, SCC, and LIS, indicating that each was founded by a

genetically unique individual (Fig. 1). All but one isolate traced to the GOM are clonal (clade II)

and share ancestry with PNW isolates (9, 46-48), and the 2006 New York isolate VpG-1 (9)(Fig.

1, Table S1). Most isolates traced to other Atlantic locations and those from CA share common

ancestry with isolates reported from British Columbia Canada in 2005, and later in WA in 2011-

2012 (Fig. 1 and Table S1). All but one SCC-traced isolate (clade III) are clonal and only two

clade members trace to other nearby locations. LIS-traced isolates are also mostly clonal (76%)

(clade VII), though two distinct 2018 isolates group within a mixed-location clade (clade V) with

isolates trace to the PNW in 2011, and from oysters consumed in CA in 2015. Phylogenies built

with core non-recombining variation mostly agreed with whole genome phylogenies, though a

change in clonal assignment of two strains (CTVP25C and MAVP-48) alluded to the possibility

that horizontally acquired variation these shared in common with the GOM and SCC clades

respectively may have obscured their distinct heritage (Fig. 1, S1 & S2; Tables S2- S4).

Given the geographic linkage of clonal subpopulations (Fig. 1), we examined the

population genetic structure of isolates from known locations to identify patterns of coancestry

and admixture (49). The conservative use of core variation and eight ancestral populations

(which best explains the data, see Fig. S3A & B) identified two old-PNW populations that were

replaced by a single modern-day PNW population, comprised of two geographically distinct

lineages: one related to GOM isolates and the other to most SCC/LIS/MAC isolates (Fig. 1 &

Fig. 2). Six PNW isolates were exceptions in that they belong to the same population and form

clades with other translocated isolates, including those traced to Spain. Although no PNW

isolates belong to the population that translocated to CA, two PNW isolates exhibit a shared

history of admixture with the CA isolates providing clues to this lineage's origin. The clonal

clades from SCC and LIS, which are in very close proximity to each other and without

geographic barriers, also belong to the same population, whereas two MAC isolates form a

distinct population. Atlantic-derived isolates from locations south of the GOM exhibit little

coancestry with other populations, whereas the GOM clade displays near equal mixed

coancestry between the prevalent modern PNW population and the other Atlantic populations

(Fig. 2). Five PNW isolates also exhibited this same pattern of mixed heritage.

Analysis of admixture that incorporated horizontally acquired content separated the two

Atlantic populations on opposite sides of Cape Cod, MA (GOM and SCC) from the rest of those

in the Atlantic which grouped with strains from CA (Fig. S4). This also relocated a single GOM

clade isolate (MAVP-20) into the CA/MAC/LIS population. This suggested that horizontally

acquired variation in the SCC and GOM lineages (that MAVP-20 lacks) may define their

uniqueness. In contrast to the GOM and SCC populations that exhibit little admixture, the

modern PNW ST36 isolates exhibit complex admixture as do members of the LIS population

(Fig. S4). Relatively few individuals within the GOM and SCC populations display some

admixture with each other. Thus, population structures inferred from horizontal variation better

aligned most strains to their geographic locations even as horizontal content overshadowed

some interesting ancestral patterns, specifically, the shared mixed heritage of the GOM clade

with modern PNW strains (Fig. 2).

Inovirus loss and reacquisition accompanied ST36 population replacement and expansion

One explanation for the differences in core versus whole genome population structure is

that differential genetic gains or losses preceded clonal expansion. To identify such features, we

compared high-quality genomes of four Atlantic clade members, (MAVP-1, MAVP-23, MAVP-

36, and MAVP-26), to PNW strains of the new-PNW (12310) and old-PNW (10290) clades. This

revealed MAVP-26, MAVP-36, and 10290 each contain a unique prophage integrated into the

dif site at the replication terminus of chromosome I, similar to prophage f237 found in pandemic

O3:K6 strains (39, 41) (Fig.3). In contrast, the other two Atlantic clade members and the new-

PNW isolate lack a phage in this location (Fig. 3). These are classified in the family *Inoviridae*

(hereafter inoviruses) and we assigned unique names, including "vB" for <u>v</u>irus infecting <u>B</u>acteria,

and "Vipa" in reference to the host _Vibrio parahaemolyticus_ and isolate name hereafter Vipa26

in MAVP-26, Vipa36 in MAVP-36, and Vipa10290 in strain 10290 (50).

These inoviruses have a conserved central core (ORF 1-7) and ORF9 and ORF10 (Fig.

3A), and two variable regions. Whereas core phage gene functions were identifiable (Table 1),

the functions encoded in variable regions were not discernable. Nucleotide variation in the

inoviruses represented up to 46% of their host's genome variation helping to explain differences

between whole and core genome phylogenies (Fig. 1, S1 & S2) and population structures (Fig.

2 and Fig. S4). Though this content would be credited to recombination, surprisingly, genomes

that lack inoviruses had a higher proportion of gene content assigned to blocks of

recombination: 2.7% in 12310 and MAVP-1, and 2.0% in MAVP-23, compared to 1.3% in

10290, 1.4% in MAVP-26, and 1.8% in MAVP-36 (Fig. S1 and Table S3). Though divergent

from one another, inovirus variation likely did not alter function: the 74 variant sites in the core of

Vipa26 generated only three non-synonymous mutations, whereas the 78 variant sites in Vipa36

produced 13 non-synonymous mutations (Fig. 3B, Table S6). ORF1-7 are under purifying

selection (codon-based Z test =9.425, p < 0.001) implying these are essential genes, and that

their function is preserved. This also suggests the infections not cryptic, and the phage are likely

still functional. Comparisons of core inovirus content from additional ST36 lysogens revealed

100% identity within the Vipa26 and Vipa36 lineages, whereas Vipa10290 had 18 non-

conserved variable sites, perhaps reflecting its long history with a sizeable ST36 population

(Fig. 3B).

With only one exception (A5Z905), all old-PNW clade members, including CTVP44C,

contained Vipa10290 or related phage (Fig. 1, square). ST39, a member of the same clonal

complex as ST36, also contains Vipa10290 whereas other clonal complex members (ST59 and

ST21), share a distinct inovirus (Fig. S5, Tables 2, S9, & S10). Despite this history of lysogeny

among the endemic population, no modern-day and definitively traced PNW ST36 isolates

harbor inoviruses (Fig. 1 & Table S1 & S7) and this loss coincides with population diversification

(Fig. 1). However, like GOM and SCC clade members, a total of nine other modern ST36

isolates collected between 2007-2016, some that dispersed to new locations and all others

untraced, acquired eight other inoviruses (Fig. 1, Table 2).

The distribution of community-acquired phage offers clues of residency and dispersal.

For the Atlantic ST36 populations, inovirus content was diagnostic of lineage and

geography. Vipa26 was absent from only one member (MAVP-20) of the ST36 GOM clonal

clade and present in only one distinctive isolate from a different location (CTVP25C)

representing either basal or independent acquisition (Fig. 1 circles). Vipa36 (Fig. 1 triangles) is

in all members of the SCC clade and only two isolates from this clade were traced to nearby

locations. Because these inoviruses, like f237, are not widespread and have been maintained

by these lineages tied to two locations, they are diagnostic of these ST36 subpopulations. In

contrast, and like the modern PNW traced-ST36 isolates, no isolates of the LIS or MAC clonal

clades harbor inoviruses.

The association of inoviruses with two Atlantic clades, and their absence in others,

suggests that the founding subpopulation progenitors acquired phage upon Atlantic invasion

only in some locations perhaps reflecting phage predation pressure. Environmental surveys

revealed that although 22% of Northeast US *V. parahaemolyticus* isolates harbor inoviruses,

this proportion varied by location (50). Proportions were lower in LIS (10%), than in GOM and

SCC (30% and 42% respectively). Whereas no environmental isolates other than ST36 harbor

Vipa36, multiple non-ST36 isolates collected in NH as early as 2008 contain Vipa26 (Table S1,

Fig. 4, & S6). This suggests Vipa26 was acquired locally as supported by its absence from VpG-

1 (Fig. 1, Table S1 & S7).

Because comparative phage phylogeny could elucidate the history of ST36 as members

moved through other populations, we expanded our search for inoviruses among available

genomes of *V. parahaemolyticus.* These prophage genomes added to the *Inoviridae* diversity

(Fig. & S5) but display little discernable phylogeographic population structure in that related

phage are found in both the Atlantic and Pacific *V. parahaemolyticus* populations, and phage in

Asian isolates group among most branches, with the exception of Vipa10290. Two ST36

isolates harbor two different inoviruses maintained by members of the PNW co-resident but

unrelated ST43 (Table 2, Fig. S5). Though most members of the two ST36 lineages reported in

Peru also lack inoviruses, including a 2011 environmental isolate related to the Spain lineage

(45), one clinical isolate of this lineage harbors a Vipa10290-related inovirus, whereas a second

clinical isolate (1.146-15), harbors f237 (Fig. 4).

The phylogenomic distribution of inoviruses in all publicly available *V. parahaemolyticus*

genomes indicates they are common, with 46% of genomes containing one or more inovirus

(Fig. 5, Fig. S6, and Table S10) yet distributed unevenly. Many lineages harbor a persisting

inovirus (Fig. 5 and Fig. S6). A few lineages harbor two evolutionary distinct inoviruses

concurrently suggesting that protection from superinfection, a common attribute of both

temperate phage and inoviruses (51, 52), is not absolute (Fig. S5). In addition to Vipa26, a

second inovirus is present in multiple, unrelated STs (Fig. S6). Even though most STs have an

inovirus prophage that they propagate vertically, the inoviruses of some isolates were replaced

by another, like ST36 (Table 3, Fig. S6). Notable among these are emergent pathogenic

lineages including ST43 and the prevalent Atlantic endemic ST631 lineage that has caused

illnesses along the North American Atlantic coast (4), the pandemic complex (ST3) and a

diverse and far-spreading Asian lineage that is now an Atlantic resident (ST8).

Discussion

Here we demonstrate that changes in inovirus prophage accompanied diversification,

competitive replacement, dispersal, and Atlantic invasion of *V. parahaemolyticus* ST36. This

suggests that inoviruses could countribute to the emergence of new lineages with the potential

to spread more broadly. The misleading nature of disease reporting that does not align illnesses

with environmental reservoirs and declining availability of recent ST36 genomes from the US

PNW have been obstacles to deciphering the ecological context of ST36 evolution and spread.

We addressed these limitations using genomes from isolates reported in Canada and other US

states that were publicly available and by augmenting available genomes with a collection of

ST36 clinical isolates from the Northeast US. We carefully curated environmental traceback to

advance our understanding of ST36 ecological expansion into the Atlantic Ocean.

Our analysis indicates five or more unique ST36 progenitors diversified in the Pacific and

then translocated into the Atlantic to found subpopulations (Fig. 1, S2, 2, and S4). The earliest

sequenced ST36 isolate traced to the Atlantic in 2006 (VpG-1) and several PNW strains share a

similar pattern of mixed ancestry with this isolate and group with the GOM clade suggesting its

progenitor may have arrived prior to 2006 with little effect on epidemiology until after the

anomalously mild winter of 2011-2012 (1, 2). Isolates traced to SCC, LIS and MAC are more

related to each other, and multiple Pacific-traced isolates group within or adjacent to their clades

(Fig. 1, 2, & S2). This pattern is most simply explained by divergence of these populations'

ancestors in the Pacific prior to introduction into the Atlantic (Fig. 1, 2 & S2). Clade V's shared

ancestry with CA and PNW isolates further implicates the PNW as the source population. And

although imperfect trace back or mislabeled genomes could explain some unexpected patterns,

nearly every Pacific-traced relative of Atlantic residential populations were isolated, reported,

and sequenced independently and prior to Atlantic isolation (Table S1). Although our

interpretations of population evolution and dispersal contrast with previous reports (9), this is

mostly explained by differences in geographic assignment of isolates to environmental sources

instead of the readily available location of disease reporting as our goal was to understand

dynamics affecting reservoirs of emergent pathogenic lineages (See details of Table S1).

Traceback confirmation from CA (Table S1), disease reporting in Mexico (53) and

environmental isolation in Peru (9, 45) further corroborate that these lineages diversified in the

Pacific. And though missing intermediate populations could suggest unknown reservoirs exist, it

is more likely that sampling bias, a decrease in availability of PNW genomes, decreased clinical

prevalence of ST36 in the PNW, and lack of surveillance there have compounded challenges

with deciphering this pathogen's evolution and spread. Based on mounting evidence that most if

not all lineages arose in the PNW, this warrants further examination of the PNW environmental

reservoir to manage further spread.

Even as increasing ocean temperatures and weather events have been tied to both

seasonal illnesses by *V. parahaemolyticus* and its spread to new locations (9, 54-57), the stable

and non-mixing distribution of Atlantic ST36 lineages suggests that environmental fitness that

promotes persistence may be their key to success. It is notable that in BC Canada where many

diversified ST36 isolates were first reported, rising infection rates were not fully explained by

temperature models but rather attributed to new pathogenic strains (58, 59). Like the Northeast

US, the MAC experienced a warming trend in 2012 (9), nonetheless this location has not

experienced persistent problems with ST36 (1, 10). Combined, we believe warming ocean

temperatures are only part of the picture and that underexplored ecological interactions are at

play in driving *V. parahaemolyticus* invasion and persistence (60). Even in locations in the

Atlantic with recurrent illness, environmental prevalence of ST36 is extremely rare (Table S1)

illustrating how low-level persisting populations can change epidemiology. Identification of

unique ecological and population contexts of the MAC compared to more northern locations

could provide useful insight.

Phage loss and later sequential phage replacement is one of the most striking and

common features of ST36 lineage succession in the PNW and population expansion into the

Atlantic. Nearly all old PNW lineage members harbor Vipa10290 (Fig. 1 and Table S1), whereas

strains of the new ST36 lineages isolated in the PNW after 2002 lack inoviruses, though other

phage are not uncommon (Table S3). The presence of Vipa10290 in other clonal complex

members could indicate a basally acquired phage was lost during lineage diversification.

Though inovirus excision is infrequently reported (61, 62), laboratory passage of ST36 can lead

to both phage curing and loss of protection from superinfection (63). The lack of inovirus-

harboring PNW ST36 population members indicates they ultimately were at a disadvantage in

the context of a changing population even though other residents concurrently maintained

inoviruses (e.g., ST43, Table 2) and potentially transmitted these to a few ST36 members. The

subsequent acquisition of inoviruses by diverging ST36 lineages (Table 2) indicates ST36

remained susceptible to infection, though the persisting inovirus-free state of modern-PNW

ST36 lineages suggests that resistance may have evolved during their diversification, as has

been described in other sympatric *Vibrio* populations (64). In contrast, that the progenitors of

two clinically prevailing Atlantic ST36 lineages acquired inoviruses prior to clonal expansion,

while ST36 populations in the LIS where inoviruses are less abundant did not, indicates phage

acquisition was not requisite for success, perhaps reflecting differences in phage predation and

other population dynamics. Even so, whereas resident ST36 from SCC and GOM have

continued to cause sporadic disease, infections from LIS have precipitously declined, in part

reflective of successful management.

Inovirus prophage confer obvious costs through persistent non-lethal virion production

(61), suggesting their loss would be favored; and yet inoviruses have been maintained by

prevalent lineages for decades (Table 2 & 4, Fig. 5). This could reflect their fastidious nature,

but the success of the two northernmost Atlantic populations harboring inoviruses contrasting

with the unsuccessful lineages that lacked them signals alternatives are possible: that

inoviruses may confer advantages in some as yet undefined context. For example, shed

inovirus virions contribute to biofilm matrix (65) thereby promoting virulence and antimicrobial

resistance, but also potentially enhancing survival during long-distance movement on particles

(52, 54, 66-68). Shed inoviruses can mislead human immune responses and decrease host

ability to clear bacterial infections, thereby enhancing virulence (69). Furthermore, variable

content in some inoviruses is linked to anti-predation (17, 70) and virulence (22, 27, 39). We find

it intriguing that succession in the PNW and loss of inoviruses by ST36 accompanied its clinical

decline there. Future functional studies with these inoviruses are essential to discern whether

any of these possibilities are at play.

Prophage can also modify their hosts in ways that alter their capacity for horizontal gene

transfer (HGT). Prophage can protect against superinfection, including by unrelated lytic phages

(27, 61, 71) thereby blocking transduction. They also modulate, and concurrently block

conjugative pili (72). The concept that loss of prophage-conferred immunity contributed to ST36

evolution is attractive considering DNA release by predation of non-lysogen susceptible hosts

could promote transformation through natural competence, though the conditions of natural

competence by *V. parahaemolyticus* are unknown (73-75). Type IV pili which are the receptor

for some inoviruses also facilitate transformation mediated HGT (51, 52). If the newly inovirus-

less ST36 lineages were more vulnerable to phage infections, phage transduction, and conjugal

transfer alike this could have contributed to its rapid diversification (Fig. 1) whereas

translocating lineages would likely not have evolved mechanisms to resist resident Atlantic

phage facilitating re-acquisition (31, 52). In keeping with this premise, many of the most highly

divergent ST36 individuals lacking inoviruses evolved by multiple recombination events

including an isolate of the MAC clade (VP30). Accumulation of non-inovirus prophage was

notable in the early diverging Spain lineage (clade I) (9). And whereas the inovirus-harboring

SCC and the GOM clades remain remarkably stable, as do their phage, Atlantic isolates lacking

inoviruses, including the single GOM clade member (MAVP-20) and many members of the LIS

population exhibit complex admixture (Fig. S4). In contrast, the inovirus-harboring old-PNW relic

CTVP44C isolated in LIS in 2013 is virtually unchanged from the last known members of this

lineage from the PNW that were isolated in 2002 (Fig. 1, Table S1). Furthermore, a 2005 isolate

of this same lineage lacking Vipa10290, A5Z905, is among the most highly divergent ST36

strains (Fig. 1, 2, S2). Similarly, pandemic O3:K6 strains, which stably maintain f237, reported in

the Americas between 1996-2012 display remarkably little genetic variation from the type strain

for this lineage RIMD 2210366 (76). Though these data only establish correlation of phage

absence and diversification, it alludes to the possibility that inovirus prophage that provide some

level of resistance to superinfection and block conjugation could thwart HGT, and by contrast,

loss of inoviruses could remove barriers to recombination and promote rapid diversification. The

selective conditions that favor inovirus acquisition and maintenance or loss would thereby

influence some mechanisms that could promote rapid evolution, potentially of more or even less

pathogenic potential.

These analyses suggest an intriguing connection of inovirus prophage to pathogen

evolutionary dynamics, alluding to the possibility that inoviruses could be gatekeepers of HGT in

natural ecosystems driving changes in human disease epidemiology. With broader population

analysis and functional studies, this could lay the foundation for a new evolutionary paradigm

where inovirus prophage-mediated immunity is a major governing force beyond the appreciated

roles of phage in selective predation (20-22) and phage conversion (23-27). The apparent ability

of ST36 and some other lineages (Table 2 & 3) to transition between phage infection states or

replace their phage could generate flexibility in the balancing of fitness tradeoffs under different

selective regimes that are not yet understood (29, 64). The high prevalence of inoviruses not

only among the global population of *V. parahaemolyticus* (Fig. 5) but universally in other

bacteria (27) with little clarity of how inoviruses impact the ecology of their hosts calls for more

mechanistic analyses which could reveal varied roles for inoviruses in promoting evolution and

competitive dominance of emerging pathogens.

Materials and Methods

Vibrio parahaemolyticus genomes

Regional clinical strains spanning 2010-2018, and trace back were acquired from public

health laboratories (2). Only isolates with single source traces were assigned a location.

Environmental isolates were collected between 2007-2015 (1, 2). Isolates were identified as *V.*

parahaemolyticus by *tlh* gene amplification (77, 78) and identity confirmed by genome

sequencing. Publicly available genomes were acquired from NCBI

(https://www.ncbi.nlm.nih.gov/genome/681; August 2019).

DNA was extracted using the Wizard Genomic DNA purification Kit (Promega, Madison

WI USA) or by organic extraction (2). Sequencing libraries were prepared as described (79).

Genomic DNA was sequenced using an Illumina – HiSeq2500 device at the Hubbard Center for

Genome Studies at the University of New Hampshire, using a 150bp paired-end reads with de-

multiplexing and quality filtering prior to analysis. The *de novo* genome assembly were

performed using the A5 pipeline (80), and annotations assigned with Prokka1.9 using "*Vibrio*"

for the reference database (81). The sequence types were determined using the SRST2

pipeline (82) or using assemblies (83) referencing https://pubmlst.org/vparahaemolyticus/(84).

Reference inoviruses were extracted from genomes sequenced using the Pacific Biosciences

RSII technology and with Illumina short read error correction as described (4).

V. parahaemolyticus *Phylogenetic Relationships*.

Because all ST36 strains are closely related (clonal) and recombination contributed to

the recent lineage divergence, ST36 strain relationships (Fig. 1) and all *V. parahaemolyticus*

relationships (Fig. 5) were determined using reference-free, whole genome alignments using

kSNP 3.1 (85) and maximum likelihood phylogenetic trees built by RAxML (86) and visualized

with iTOL(87). The optional Kchooser script provided in kSNP3.1 determined the optional kmer

size of 19, for each alignment dataset. Whole genome trees were rooted with *Vibrio*

alginolyticus ARGOS_108 (hidden after rooting in iTOL). Maximum likelihood trees were

inferred using the GTR-GAMMA model of nucleotide substitution, and the convergence criteria

autoMRE option for automatically determining a sufficient number of rapid bootstrap replicates.

RAxML completed a thorough ML search after optimization on every 5[th] bootstrapped tree,

which were mapped on the ML tree with the best likelihood ("raxmlHPC-PTHREADS-AVX -m

GTRGAMMA -f a -N autoMRE -x 12345 -p 12345" options). Gene/features were visualized with

EasyFig 2.2.0 (88, 89).

In parallel, homoplastic regions resulting from recombination were identified in genomes

by alignment to reference strain 10296 (90) (summary table S2) using Gubbins 2.3.4 (91) and

removed, and the relationships of strains determined based on nucleotide SNP alignments

generated with Gubbins 2.3.4 (91) (Fig. 2). Phylogenies were inferred as above, except using

the GTR-CAT nucleotide substitution model (because the alpha parameter was >10). ML search

and optimization, rapid bootstrap criterion, and mapping of bootstraps were done as described

above. Unmapped reads were assembled using SPAdes 3.13.1 (92) and annotated using

Prokka 1.14(81) (Table S4).

Population structure of ST36 was performed using whole and core SNPs genetic

matrixes generated from SNP alignments in the R statistical package LEA (49, 93, 94) (Fig. 2).

To determine the number of genetic clusters best explained by the distribution of genomic

variation, we explored ancestral populations (*K*) of between 1-20 and used the entropy criterion

to evaluate the quality of fit of the statistical model to the data and selected the probable number

of populations based on minimal entropy. Once strains were assigned to populations, the least-

squares estimates of ancestry proportions were plotted using ggplot2 (95).

Inovirus Identification

ST36 strains were compared to reference 10290 (5) using Breseq (96). Unique

prophage were classified as belonging to the family *Inoviridae* proposed subfamily

Protoinoviridae (27) and assigned unique names (97, 98) (Table 2). All *V. parahaemolyticus*

genomes were independently searched against a BLASTn (99) database constructed using the

central core (ORF1 – ORF7) of vB Vipa-26. Nucleotide sequences of 745 phage were extracted

and to determine if they were unique, maximum likelihood phylogenies using the GTR-GAMMA

model for substitution and rapid bootstraps were constructed via RAxML on nucleotide SNPs

generated by kSNP (Fig. S6). Rapid bootstraps reached criterion after 350 replicates and were

mapped on the tree as described above. The complete prophage assemblies of unique ST36

inoviruses were named from the highest quality assembly as were other prevalent inoviruses

(Table S9). Relatedness of select whole phage genomes was visualized using the Genome-

BLAST Distance Phylogeny (GBDP) method (100) to conduct pairwise comparisons of the

nucleotide sequences under settings recommended for prokaryotic viruses (101). Trees were

rooted at the midpoint (28) and visualized using iTOL (87) (Fig. S5).

To determine if phage evolved under selection, a Nei-Gojobori codon-based Z test (102)

was performed in MEGA 6 (103). Protein sequences for the phages Vf33 (NC_005948.1) (104,

105), CTXΦ (MF155889.1)(27), and the type strain for *Inoviridae* M13, which infects *Escherichia*

coli (GCF_000845205.1) were compared using BLASTp (99).

Oligonucleotide primers were designed for multiplex amplification with the species-

specific *tlh* primers (77, 78) where primers annealing within ORF3 within ORF5 identified

inovirus presence, and primers annealing within HypD and ORF9 produced different size

amplicons diagnostic of Vipa26 and Vipa36 (Table 4). Isolates were screened with 0.2µM primer

annealed at 55°C with 1.5 min extension and phage identity confirmed by genome sequencing.

Data Availability

Genome accessions are listed in Tables S1, S7, S8 and Table 2.

Acknowledgments: Partial funding was provided by the National Sea Grant College Program

and projects R/SFA-4, R-HCE-3, and R/SFA-13, and by the New Hampshire Agricultural

Experimentation Station. This work (scientific contribution number 2995) was supported by the

USDA National Institute of Food and Agriculture Hatch projects NH00609 [accession number

233555], NH00625 [accession number 1004199], NH00658 [accession number 1013479], and

NH00698 [accession number 1023165]. We thank Narjol Gonzalez-Escalona for helpful insight

and feedback; Kanwit Kohl, Stephen Combes and Heather Grieser from the Maine Center for

Disease Control and Prevention for strains and information; Tracy Stiles, Eric Hickey, Coleen

Murphy, Michael Moore, and Jana Ferguson from the MA Department of Public Health for

helpful insight, strains and epidemiological information; Jeff Kennedy and Christian Pepitas from

the MA Division of Marine Fisheries for insight and guidance; Christopher Nash and Robert

Atwood from the New Hampshire Department of Environmental Services and the Fish and

Game Department, and Colleen Smith from the New Hampshire Department of Health and

Human Services for information and discussion. We also thank Zoe Pavlik for curation of

genome accessions.

Role of authors: JM, AM, RF, FX, CS, KD, JH, VC, SJ, CW collected samples, performed and

interpreted analyses, curated databases, and constructed visuals. JM and AM drafted the

manuscript and CW and RF completed writing the manuscript. SJ, VC, and CW designed and

directed the studies. All coauthors reviewed and edited the manuscript for accuracy and

completeness.

Figure 1. Maximum-likelihood phylogeny of ST36 *Vibrio parahaemolyticus*. ML-

phylogenies were built with 1,025,281 aligned variable nucleotide sites identified in quality ST36

isolates. Isolates are colored to correspond to geographic environmental traceback, where

available (Table S1) and where no color indicates unknown or ambiguous origin. Isolates

acquired from four Northeast US States as part of this study are named uniquely by a

combination of state (Maine [ME], New Hampshire [NH], Massachusetts [MA], and Connecticut

[CT]) of reporting and sequential numbers. Symbols next to isolates indicate identity of inovirus

content. The ancestral PNW ST36 population is identified by a black bar, whereas seven

different lineages associated with translocation events (I – VII) are identified by greyscale bars.

Bootstraps are from 250 replicates (criterion reached).

Figure 2. Population structure of ST36 strains. Coancestry estimates were inferred from

SNP matrices using 741 SNPs identified from core non-recombining genomes of ST36 isolates

with known environmental source and at least one close relative (see Table S5 for excluded

genomes) by LEA (49). Colored bars represent proportion of genetic variation derived from 8

ancestral populations (see Fig S3A). Geographic locations are marked by colors below strain

name, followed by phage content (symbol) and year of isolation.

Figure 3. Comparisons of inoviruses in *V. parahaemolyticus* ST36. (A) Alignments of

reference inovirus genomes depicting unique and shared content, and % nucleotide identity

between Vipa36 (9,721 bp), Vipa26 (10,893 bp) and Vipa10290 (9,414 bp) as compared to f237

(8,784 bp). Orthologous ORFs are depicted by the same color except for non-homologous light

blue hypothetical proteins. (B) Core inovirus genome (ORFs 1-7, and ORF9-10) single

nucleotide polymorphisms (SNPs) were identified by comparison of quality ST36 genomes to

the alignment consensus of reference inovirus genomes for Vipa36 (21 genomes), Vipa26 (24

genomes), and Vipa10290 (23 genomes) and used to assess spatial distribution of variation

between the inoviruses and within each inovirus lineage. SNPs are mapped by location where

ORF identities are labeled and colored to match those in (A), and SNP type designated by

colored blocks (see Table S6). This indicates few non-synonymous mutations (see text for

details) and uneven distribution of variation with ORF3, an intergenic region between ORF4 and

ORF5, and the 5' end of ORF5 being identical in all three phage. Only Vipa10290 displayed

non-conserved core genome variation among ST36 prophage.

Table 1. Inferred functions of inovirus genes [1].

ORF	Vf33 (E-value/%Identity)	CTXΦ (E-value/%Identity)	M13 (E-value/%Identity)	Predicted Functions[a]
ORF1	Vpf402 (0.0/99%)	RstA (7×10^{-66}/36%)	gII (size and location)	Replication initiation protein
ORF2 Integrase	Vpf117 (1×10^{-85}/92.5%)	RstB (0.04/13.3%)	-	Integration
ORF3	Vpf81 (0.004/7.8%)	RstC (size and location)	gV (size and location)	ssDNA binding protein, helix destabilizing
ORF4	Vpf77 (5.3/3.7%)	Cep (7.0/7.3%)	gVIII (0.45/4.9%)	Major coat protein
ORF5	Vpf491 (0.0/59%)	OrfU (0.54/11.9%)	gIII (0.6/11.3%)	Adsorption, termination of assembly, tail protein
ORF6	Vpf104 (0.2/19%)	Ace (5×10^{-10}/18.3%)	gVI (4.1/8.7%)	Minor coat protein, termination of assembly
ORF7	Vpf380 (1×10^{-4}/13.9%)	Zot (2×10^{-27}/17.8%)	gI (5×10^{-4}/8.5%)	Assembly protein, maturation
ORF9	Vpf122 (4×10^{-66}/68%)	RstR (0.17/5.7%)	-	Transcriptional repressor, regulator
ORF10	Vpf152 (1^{-44}/88-86%)	-	-	Function unknown

524   [1]ORF number corresponds to numbers designated for f237, and comparative analyses between

525   homologous ORFs completed with Vipa26 as a reference.

526   [a] Functions summarized from Mai-Prochnow et al., 2015 (36) and Chang et al., 1998 (37)

Table 2. Summary of *Inoviridae* prophage in ST36 isolates

Strain	Inovirus name	NCBI Accession (Genome coordinates)	Additional ST36 genomes with inovirus	Other STs with related inoviruses
10290	vB_Vipa10290	AVOH01000008.1 (1370845..1378511)	Many	38, 39
MAVP-26	vB_Vipa26	MT188662	Many	1574, NF
MAVP-36	vB_Vipa36	MT188663	Many	None
CDC_K5323G	vB_Vipa5323	MIUF01000019.1 (188973..190181) MIUF01000072.1 MIUF01000065.1 MIUF01000002.1 (4141..6171)	None	752, 1132, NF
CDC_K5308	vB_Vipa5308	MIUE01000114.1 (7376..8729) MIUE01000031.1 (5201..8319)	None	43
10-4255	vB_Vipa71[1]	MT193890 NIXZ01000007.1 (181573..191867)	None	43
CDC_A8962	vB_Vipa8962	LHRO01000017.1 (70737..72090)	None	327
CDC_K5512	vB_Vipa5512	MIUZ01000070.1 (628..7658)	CDC_K5345G, CDC_K5280	1713
MEVP-10	vB_Vipa10	MT188666	None	None
1.146-15	f237	WSRX01000192.1, WSRX01000239.1, WSRX01000194.1, WSRX01000150.1	None	3, 152, 809, 1473, NF
1.220-16	other	AAXNNA010000036.1	None	None

NF: Sequence type that is not yet assigned; None: no strains harboring phage.

[1]Named for ST43 strain MAVP-71

Figure 4. Whole genome phylogenies of representative inoviruses from diverse locations and sequence types. Whole genome phylogenies of select inoviruses from environmental (E) and clinical (C) isolates of *V. parahaemolyticus* labeled by name and year of isolation. Bootstraps criterion met after 350 replicates. Environmental traceback color coding and symbols, representing named phage corresponds to those in Figure 1 where circle is Vipa26,

square is Vipa10290, triangle is Vipa36 and grey square is other (Vipa8296). ST36 isolates are

in bold.

Figure 5. Maximum-likelihood tree of diverse *V. parahaemolyticus* isolates with their

inovirus content. A maximum-likelihood (ML) phylogeny was built on 1,025,281 genome SNPs

where 250 bootstraps (criterion met) were mapped onto the best scoring ML tree. Isolates are

colored to correspond to geographic region as in Fig. 1, where no color indicates unknown or

ambiguous origin. Symbols next to strains indicate unique inovirus content. Outermost label

indicates the sequence type (ST) where NF or unlabled = sequence type not found or not

known, followed by year of isolation, if known.

Table 3. Lineages other than ST36 with diverse phage genotypes.

Sequence type	Number of unique Inovirus prophage in lineage	Sequenced strains without prophage
631	4	Yes
674	4	No
43	5	Yes
12	4	No
3	2	Yes
8	2	Yes
308	2	Yes
120	2	Yes

549   **Table 4. Multiplex PCR for detection of inoviruses**

Primer	Sequence (5'-3')	Amplicon Size (bp)	
		Host (*V.p.*)	Inovirus
TLH-F2 TLH-R	AGAACTTCATCTTGATGACACTGC GCTACTTTCTAGCATTTTCTCTGC	401	N/A
ST36Phage F2 ST36Phage R2	AGCAACGAAAACGCCTGT ACCGTATCACCAATGGACTGT	N/A	~1000
PhHypDF3 PhORF9R1	AAGTGCTACATGAATGAAAGTGCT TCAATGAAGTATCACGAAATGACTA	N/A	Vipa26: 1440 Vipa36: 854

N/A: not an amplification template

Literature Cited:

1.   Urquhart EA, Jones SH, Jong WY, Schuster BM, Marcinkiewicz AL, Whistler CA, Cooper
VS. 2016. Environmental conditions associated with elevated *Vibrio parahaemolyticus*
concentrations in Great Bay Estuary, New Hampshire. PloS ONE 11:e0155018.
2.   Xu F, Ilyas S, Hall JA, Jones SH, Cooper VS, Whistler CA. 2015. Genetic
characterization of clinical and environmental *Vibrio parahaemolyticus* from the
Northeast USA reveals emerging resident and non-indigenous pathogen lineages. Front
Microbiol 6:272.
3.   Baker-Austin C, Trinanes JA, Taylor NG, Hartnell R, Siitonen A, Martinez-Urtaza J.
2013. Emerging *Vibrio* risk at high latitudes in response to ocean warming. Nature
Climate Change 3:73-77.
4.   Xu F, Gonzalez-Escalona N, Sebra RP, Dress KP, Cooper VS, Jones SH, Whistler CA.
2017. Parallel evolution of two clades of a major Atlantic endemic *Vibrio*
parahaemolyticus pathogen lineage by independent acquisition of related pathogenicity
islands. Appl Environ Microbiol doi:https://doi.org/10.1101/155234.
5.   Martinez-Urtaza J, Baker-Austin C, Jones JL, Newton AE, Gonzalez-Aviles GD, DePaola
A. 2013. Spread of Pacific Northwest *Vibrio parahaemolyticus* strain. N Engl J Med
369:1573-1574.
6.   Vezzulli L, Grande C, Reid PC, Hélaouët P, Edwards M, Höfle MG, Brettar I, Colwell RR,
Pruzzo C. 2016. Climate influence on *Vibrio* and associated human diseases during the
past half-century in the coastal North Atlantic. Proceedings of the National Academy of
Sciences:201609157.
7.   Newton AE, Garrett N, Stroika SG, Halpin JL, Turnsek M, Mody RK, Division of
Foodborne W, Environmental D. 2014. Notes from the field: Increase in *Vibrio*
parahaemolyticus infections associated with consumption of Atlantic coast shellfish—
2013. MMWR Morbidity Mortality Weekly Report 63:335-336.
8.   Abbott SL, Powers C, Kaysner CA, Takeda Y, Ishibashi M, Joseph SW, Janda JM. 1989.
Emergence of a restricted bioserovar of *Vibrio parahaemolyticus* as the predominant
cause of *Vibrio*-associated gastroenteritis on the West Coast of the United States and
Mexico. J Clin Microbiol 27:2891-3.
9.   Martinez-Urtaza J, van Aerle R, Abanto M, Haendiges J, Myers RA, Trinanes J, Baker-
Austin C, Gonzalez-Escalona N. 2017. Genomic variation and evolution of *Vibrio*
parahaemolyticus ST36 over the course of a transcontinental epidemic expansion. MBio
8.
10.   Martinez-Urtaza J, Powell A, Jansa J, Rey JL, Montero OP, Campello MG, Lopez MJ,
Pousa A, Valles MJ, Trinanes J, Hervio-Heath D, Keay W, Bayley A, Hartnell R, Baker-
Austin C. 2016. Epidemiological investigation of a foodborne outbreak in Spain
associated with U.S. West Coast genotypes of *Vibrio parahaemolyticus*. Springerplus
5:87.
11.   CDC. 1999. Outbreak of *Vibrio parahaemolyticus* infection associated with eating raw
oysters and clams harvested from Long Island Sound--Connecticut, New Jersey, and
New York, 1998. MMWR Morb Mortal Wkly Rep 48:48-51.
12.   Haendiges J, Jones J, Myers RA, Mitchell CS, Butler E, Toro M, Gonzalez-Escalona N.
2016. A nonautochthonous U.S. strain of *Vibrio parahaemolyticus* isolated from
Chesapeake Bay oysters caused the outbreak in Maryland in 2010. Appl Environ
Microbiol 82:3208-3216.
13.   Zahid MS, Udden SM, Faruque AS, Calderwood SB, Mekalanos JJ, Faruque SM. 2008.
Effect of phage on the infectivity of *Vibrio cholerae* and emergence of genetic variants.
Infect Immun 76:5266-73.
14.   Jensen MA, Faruque SM, Mekalanos JJ, Levin BR. 2006. Modeling the role of
bacteriophage in the control of cholera outbreaks. Proc Natl Acad Sci U S A 103:4652-7.

15. Faruque SM, Bin Naser I, Fujihara K, Diraphat P, Chowdhury N, Kamruzzaman M, Qadri
F, Yamasaki S, Ghosh AN, Mekalanos JJ. 2005. Genomic sequence and receptor for
the *Vibrio cholerae* phage KSF-1phi: evolutionary divergence among filamentous
vibriophages mediating lateral gene transfer. J Bacteriol 187:4095-103.
16. Nanda AM, Thormann K, Frunzke J. 2015. Impact of spontaneous prophage induction
on the fitness of bacterial populations and host-microbe interactions. J Bacteriol
197:410-9.
17. Brussow H. 2007. Bacteria between protists and phages: from antipredation strategies to
the evolution of pathogenicity. Mol Microbiol 65:583-9.
18. Molina-Quiroz RC, Camilli A, Silva-Valenzuela CA. 2023. Role of bacteriophages in the
evolution of pathogenic *Vibrios* and lessons for phage therapy. Adv Exp Med Biol
1404:149-173.
19. Hampton HG, Watson BNJ, Fineran PC. 2020. The arms race between bacteria and
their phage foes. Nature 577:327-336.
20. Bohannan BJM, Lenski RE. 2000. The relative importance of competition and predation
varies with productivity in a model community. Am Nat 156:329-340.
21. Joo J, Gunny M, Cases M, Hudson P, Albert R, Harvill E. 2006. Bacteriophage-mediated
competition in *Bordetella* bacteria. Proc Biol Sci 273:1843-8.
22. Ritchie JM, Wagner PL, Acheson DW, Waldor MK. 2003. Comparison of Shiga toxin
production by hemolytic-uremic syndrome-associated and bovine-associated Shiga
toxin-producing *Escherichia* coli isolates. Appl Environ Microbiol 69:1059-66.
23. Taylor MJ, Thorne CB. 1963. Transduction of *Bacillus licheniformis* and *Bacillus subtilis*
by each of two phages. J Bacteriol 86:452-61.
24. Romig WR, Brodetsky AM. 1961. Isolation and preliminary characterization of
bacteriophages for *Bacillus subtilis*. J Bacteriol 82:135-41.
25. Buckling A, Rainey PB. 2002. Antagonistic coevolution between a bacterium and a
bacteriophage. Proc Biol Sci 269:931-6.
26. Rodriguez-Valera F, Martin-Cuadrado AB, Rodriguez-Brito B, Pasic L, Thingstad TF,
Rohwer F, Mira A. 2009. Explaining microbial population genomics through phage
predation. Nat Rev Microbiol 7:828-36.
27. Roux S, Krupovic M, Daly RA, Borges AL, Nayfach S, Schulz F, Sharrar A, Matheus
Carnevali PB, Cheng JF, Ivanova NN, Bondy-Denomy J, Wrighton KC, Woyke T, Visel
A, Kyrpides NC, Eloe-Fadrosh EA. 2019. Cryptic inoviruses revealed as pervasive in
bacteria and archaea across Earth's biomes. Nat Microbiol doi:10.1038/s41564-019-
0510-x.
28. Breitbart M, Rohwer F. 2005. Here a virus, there a virus, everywhere the same virus?
Trends Microbiol 13:278-84.
29. Askora A, Yamada T. 2015. Two different evolutionary lines of filamentous phages in
Ralstonia solanacearum: their effects on bacterial virulence. Front Genet 6:217.
30. Yen M, Camilli A. 2017. Mechanisms of the evolutionary arms race between *Vibrio*
cholerae and Vibriophage clinical isolates. Int Microbiol 20:116-120.
31. Richards GP, Chintapenta LK, Watson MA, Abbott AG, Ozbay G, Uknalis J, Oyelade
AA, Parveen S. 2019. Bacteriophages against pathogenic *Vibrios* in Delaware Bay
oysters (*Crassostrea virginica*) during a period of high levels of pathogenic *Vibrio*
parahaemolyticus. Food Environ Virol 11:101-112.
32. Onarinde BA, Dixon RA. 2018. Prospects for biocontrol of *Vibrio parahaemolyticus*
contamination in blue mussels (*Mytilus edulus*)-a year-long study. Front Microbiol
9:1043.
33. Zhang H, Yang Z, Zhou Y, Bao H, Wang R, Li T, Pang M, Sun L, Zhou X. 2018.
Application of a phage in decontaminating *Vibrio parahaemolyticus* in oysters. Int J Food
Microbiol 275:24-31.

34. Wang W, Li M, Li Y. 2015. Intervention strategies for reducing *Vibrio parahaemolyticus*
in seafood: a review. J Food Sci 80:R10-9.
35. Cao Y, Zhang Y, Lan W, Sun X. 2021. Characterization of vB_VpaP_MGD2, a newly
isolated bacteriophage with biocontrol potential against multidrug-resistant *Vibrio*
parahaemolyticus. Arch Virol 166:413-426.
36. Brossard Stoos KA, Ren J, Shields-Cutler RR, Sams KL, Caldwell S, Ho MB, Rivara G,
Whistler CA, Jones SH, Wiedmann M, DeMent J, Getchell RG, Marquis H. 2022.
Coastal water bacteriophages infect various sets of *Vibrio parahaemolyticus* sequence
types. Front Microbiol 13:1041942.
37. Ding T, Sun H, Pan Q, Zhao F, Zhang Z, Ren H. 2020. Isolation and characterization of
Vibrio parahaemolyticus bacteriophage vB_VpaS_PG07. Virus Res 286:198080.
38. Faruque SM, Naser IB, Islam MJ, Faruque AS, Ghosh AN, Nair GB, Sack DA,
Mekalanos JJ. 2005. Seasonal epidemics of cholera inversely correlate with the
prevalence of environmental cholera phages. Proc Natl Acad Sci U S A 102:1702-7.
39. Nasu H, Iida T, Sugahara T, Yamaichi Y, Park KS, Yokoyama K, Makino K, Shinagawa
H, Honda T. 2000. A filamentous phage associated with recent pandemic *Vibrio*
parahaemolyticus O3:K6 strains. J Clin Microbiol 38:2156-61.
40. Iida T, Hattori A, Tagomori K, Nasu H, Naim R, Honda T. 2001. Filamentous phage
associated with recent pandemic strains of *Vibrio parahaemolyticus*. Emerg Infect Dis
7:477-8.
41. Iida T, Makino K, Nasu H, Yokoyama K, Tagomori K, Hattori A, Okuno T, Shinagawa H,
Honda T. 2002. Filamentous bacteriophages of vibrios are integrated into the dif-like site
of the host chromosome. J Bacteriol 184:4933-5.
42. Chan B, Miyamoto H, Taniguchi H, Yoshida S. 2002. Isolation and genetic
characterization of a novel filamentous bacteriophage, a deleted form of phage f237,
from a pandemic *Vibrio parahaemolyticus* O4:K68 strain. Microbiol Immunol 46:565-9.
43. Garin-Fernandez A, Wichels A. 2020. Looking for the hidden: characterization of
lysogenic phages in potential pathogenic *Vibrio* species from the North Sea. Mar
Genomics 51:100725.
44. Garin-Fernandez A, Glockner FO, Wichels A. 2020. Genomic characterization of
filamentous phage vB_VpaI_VP-3218, an inducible prophage of *Vibrio*
parahaemolyticus. Mar Genomics 53:100767.
45. Abanto M, Gavilan RG, Baker-Austin C, Gonzalez-Escalona N, Martinez-Urtaza J. 2020.
Global expansion of Pacific Northwest *Vibrio parahaemolyticus* sequence type 36.
Emerg Infect Dis 26:323-326.
46. Banerjee SK, Kearney AK, Nadon CA, Peterson C-L, Tyler K, Bakouche L, Clark CG,
Hoang L, Gilmour MW, Farber JM. 2014. Phenotypic and genotypic characterization of
Canadian clinical isolates of *Vibrio parahaemolyticus* collected from 2000 to 2009.
Journal of Clinical Microbiology 52:1081-1088.
47. Ronholm J, Petronella N, Kenwell R, Banerjee S. 2015. Draft whole-genome sequences
of 14 *Vibrio parahaemolyticus* clinical isolates with an ambiguous K serogroup. Genome
Announc 3.
48. Ronholm J, Petronella N, Chew Leung C, Pightling AW, Banerjee SK. 2016. Genomic
features of environmental and clinical *Vibrio parahaemolyticus* isolates lacking
recognized virulence factor are dissimilar. Appl Environ Microbiol 82:1102-1113.
49. Frichot E, François O. 2015. LEA: an R package for landscape and ecological
association studies. Methods in Ecology and Evolution 6:925-929.
50. Marcinkiewicz AL. 2016. Bacterial and phage interactions influencing *Vibrio*
parahaemolyticus ecology. Master of Science in Microbiology. University of New
Hampshire.

51. Schmidt AK, Fitzpatrick AD, Schwartzkopf CM, Faith DR, Jennings LK, Coluccio A, Hunt
DJ, Michaels LA, Hargil A, Chen Q, Bollyky PL, Dorward DW, Wachter J, Rosa PA,
Maxwell KL, Secor PR. 2022. A filamentous bacteriophage protein inhibits Type IV pili to
prevent superinfection of *Pseudomonas aeruginosa*. mBio 13:e0244121.
52. Ismail MH, Michie KA, Goh YF, Noorian P, Kjelleberg S, Duggin IG, McDougald D, Rice
SA. 2021. The repressor C protein, Pf4r, controls superinfection of *Pseudomonas*
aeruginosa PAO1 by the Pf4 filamentous phage and regulates host gene expression.
Viruses 13.
53. Leon-Sicairos N, Zatarain-Lopez R, Angulo-Zamudio UA, Velazquez-Roman J, Flores-
Villasenor H, Martinez-Garcia JJ, Moreno-Perez MA, Buelna-Romero A, Hernandez-
Monroy I, Lopez-Martinez I, Cuen-Diaz HM, Diaz-Quinonez JA, Canizalez-Roman A.
2022. *Vibrio parahaemolyticus* is associated with diarrhea cases in Mexico, with a
dominance of pandemic O3:K6 clones. Int J Environ Res Public Health 19.
54. Rosales D, Ellett A, Jacobs J, Ozbay G, Parveen S, Pitula J. 2022. Investigating the
relationship between nitrate, total dissolved nitrogen, and phosphate with abundance of
pathogenic *Vibrios* and harmful algal blooms in Rehoboth Bay, Delaware. Appl Environ
Microbiol 88:e0035622.
55. Martinez-Urtaza J, Huapaya B, Gavilan RG, Blanco-Abad V, Ansede-Bermejo J,
Cadarso-Suarez C, Figueiras A, Trinanes J. 2008. Emergence of Asiatic *Vibrio* diseases
in South America in phase with El Nino. Epidemiology 19:829-37.
56. McLaughlin JB, DePaola A, Bopp CA, Martinek KA, Napolilli NP, Allison CG, Murray SL,
Thompson EC, Bird MM, Middaugh JP. 2005. Outbreak of Vibrio parahaemolyticus
gastroenteritis associated with Alaskan oysters. N Engl J Med 353:1463-70.
57. Konrad S, Paduraru P, Romero-Barrios P, Henderson SB, Galanis E. 2017. Remote
sensing measurements of sea surface temperature as an indicator of *Vibrio*
parahaemolyticus in oyster meat and human illnesses. Environ Health 16:92.
58. Galanis E, Otterstatter M, Taylor M. 2020. Measuring the impact of sea surface
temperature on the human incidence of *Vibrio* sp. infection in British Columbia, Canada,
1992-2017. Environ Health 19:58.
59. Taylor M, Cheng J, Sharma D, Bitzikos O, Gustafson R, Fyfe M, Greve R, Murti M,
Stone J, Honish L, Mah V, Punja N, Hexemer A, McIntyre L, Henry B, Kendall P,
Atkinson R, Buenaventura E, Martinez-Perez A, Galanis E, Team TOI. 2018. Outbreak
of *Vibrio parahaemolyticus* associated with consumption of raw oysters in Canada, 2015.
Foodborne Pathog Dis 15:554-559.
60. Hartwick MA, Urquhart EA, Whistler CA, Cooper VS, Naumova EN, Jones SH. 2019.
Forecasting seasonal *Vibrio parahaemolyticus* concentrations in New England shellfish.
Int J Environ Res Public Health 16.
61. Mai-Prochnow A, Hui JG, Kjelleberg S, Rakonjac J, McDougald D, Rice SA. 2015. 'Big
things in small packages: the genetics of filamentous phage and effects on fitness of
their host'. FEMS Microbiol Rev 39:465-87.
62. Das B, Bischerour J, Barre FX. 2011. VGJphi integration and excision mechanisms
contribute to the genetic diversity of *Vibrio cholerae* epidemic strains. Proc Natl Acad Sci
U S A 108:2516-21.
63. Means J. 2019. Filamentous bacteriophage associated with shaping community struture
and fitness of invasive *Vibrio parahaemolyticus* ST36 Master's Theses and Capstones.
University of New Hampshire.
64. Hussain FA, Dubert J, Elsherbini J, Murphy M, VanInsberghe D, Arevalo P, Kauffman K,
Rodino-Janeiro BK, Gavin H, Gomez A, Lopatina A, Le Roux F, Polz MF. 2021. Rapid
evolutionary turnover of mobile genetic elements drives bacterial resistance to phages.
Science 374:488-492.

65. Secor PR, Sweere JM, Michaels LA, Malkovskiy AV, Lazzareschi D, Katznelson E,
Rajadas J, Birnbaum ME, Arrigoni A, Braun KR, Evanko SP, Stevens DA, Kaminsky W,
Singh PK, Parks WC, Bollyky PL. 2015. Filamentous bacteriophage promote biofilm
assembly and function. Cell Host Microbe 18:549-59.
66. Martinez-Urtaza J, Blanco-Abad V, Rodriguez-Castro A, Ansede-Bermejo J, Miranda A,
Rodriguez-Alvarez MX. 2012. Ecological determinants of the occurrence and dynamics
of Vibrio parahaemolyticus in offshore areas. ISME J 6:994-1006.
67. Doni L, Oliveri C, Lasa A, Di Cesare A, Petrin S, Martinez-Urtaza J, Coman F,
Richardson A, Vezzulli L. 2023. Large-scale impact of the 2016 marine heatwave on the
plankton-associated microbial communities of the Great Barrier Reef (Australia). Mar
Pollut Bull 188:114685.
68. Xu Y, Yang L, Wang Y, Zhu Z, Yan J, Qin S, Chen L. 2022. Prophage-encoded gene
VpaChn25_0734 amplifies ecological persistence of *Vibrio parahaemolyticus* CHN25.
Curr Genet 68:267-287.
69. Sweere JM, Van Belleghem JD, Ishak H, Bach MS, Popescu M, Sunkari V, Kaber G,
Manasherob R, Suh GA, Cao X, de Vries CR, Lam DN, Marshall PL, Birukova M,
Katznelson E, Lazzareschi DV, Balaji S, Keswani SG, Hawn TR, Secor PR, Bollyky PL.
2019. Bacteriophage trigger antiviral immunity and prevent clearance of bacterial
infection. Science 363.
70. Erken M, Weitere M, Kjelleberg S, McDougald D. 2011. In situ grazing resistance of
Vibrio cholerae in the marine environment. FEMS Microbiol Ecol 76:504-12.
71. Paff ML, Nuismer SL, Ellington A, Molineux IJ, Bull JJ. 2016. Virus wars: using one virus
to block the spread of another. PeerJ 4:e2166.
72. Lin A, Jimenez J, Derr J, Vera P, Manapat ML, Esvelt KM, Villanueva L, Liu DR, Chen
IA. 2011. Inhibition of bacterial conjugation by phage M13 and its protein g3p:
quantitative analysis and model. PLoS One 6:e19991.
73. Molina-Quiroz RC, Dalia TN, Camilli A, Dalia AB, Silva-Valenzuela CA. 2020. Prophage-
dependent neighbor predation fosters horizontal gene transfer by natural transformation.
mSphere 5.
74. Matthey N, Stutzmann S, Stoudmann C, Guex N, Iseli C, Blokesch M. 2019. Neighbor
predation linked to natural competence fosters the transfer of large genomic regions in
Vibrio cholerae. Elife 8.
75. Dalia AB, McDonough E, Camilli A. 2014. Multiplex genome editing by natural
transformation. Proc Natl Acad Sci U S A 111:8937-42.
76. Guerrero A, Gomez-Gil B, Lizarraga-Partida ML. 2021. Genomic stability among O3:K6
V. parahaemolyticus pandemic strains isolated between 1996 to 2012 in American
countries. BMC Genom Data 22:38.
77. Panicker G, Call DR, Krug MJ, Bej AK. 2004. Detection of pathogenic *Vibrio spp.* in
shellfish by using multiplex PCR and DNA microarrays. Appl Environ Microbiol 70:7436-
7444.
78. Whistler CA, Hall JA, Xu F, Ilyas S, Siwakoti P, Cooper VS, Jones SH. 2015. Use of
whole-genome phylogeny and comparisons for development of a multiplex PCR assay
to identify sequence type 36 *Vibrio parahaemolyticus*. Journal of clinical microbiology
53:1864-1872.
79. Baym M, Kryazhimskiy S, Lieberman TD, Chung H, Desai MM, Kishony R. 2015.
Inexpensive multiplexed library preparation for megabase-sized genomes. PloS one
10:e0128036.
80. Tritt A, Eisen JA, Facciotti MT, Darling AE. 2012. A5. An integrated pipeline for *de novo*
assembly of microbial genomes. PLoS ONE 7:e42304 DOI:
10.1371/journal.pone.0042304.

81.   Seemann T. 2014. Prokka: rapid prokaryotic genome annotation.
805   Bioinformatics:DOI:10.1093/bioinformatics/btu153.
806   82.   Inouye M, Conway TC, Zobel J, Holt KE. 2012. Short read sequence typing (SRST):
807   multi-locus sequence types from short reads. BMC Genomics 13:338.
808   83.   Jolley KA, Maiden MC. 2014. Using MLST to study bacterial variation: prospects in the
809   genomic era. Future microbiology 9:623-630.
810   84.   Inouye M, Dashnow H, Raven LA, Schultz MB, Pope BJ, Tomita T, Zobel J, Holt KE.
811   2014. SRST2: Rapid genomic surveillance for public health and hospital microbiology
812   labs. Genome Med 6:90.
813   85.   Gardner SN, Slezak T, Hall BG. 2015. kSNP3.0: SNP detection and phylogenetic
814   analysis of genomes without genome alignment or reference genome. Bioinformatics
815   31:2877-8.
816   86.   Stamatakis A. 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses
817   with thousands of taxa and mixed models. Bioinformatics 22:2688-2690.
818   87.   Letunic I, Bork P. 2019. Interactive Tree Of Life (iTOL) v4: recent updates and new
819   developments. Nucleic Acids Res 47:W256-W259.
820   88.   Darling AE, Mau B, Perna NT. 2010. progressiveMauve: multiple genome alignment with
821   gene gain, loss and rearrangement. PLoS One 5:e11147.
822   89.   Sullivan MJ, Petty NK, Beatson SA. 2011. Easyfig: a genome comparison visualizer.
823   Bioinformatics 27:1009-10.
824   90.   T. S. 2015. Snippy: fast bacterial variant calling from NGS reads [Internet]. Available
825   from https://github.com/tseemann/snippy Accessed
826   91.   Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, Parkhill J,
827   Harris SR. 2014. Rapid phylogenetic analysis of large samples of recombinant bacterial
828   whole genome sequences using Gubbins. Nucleic acids research:gku1196.
829   92.   Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. 2020. Using SPAdes
830   De Novo Assembler. Curr Protoc Bioinformatics 70:e102.
831   93.   Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane
832   T, McCarthy SA, Davies RM, Li H. 2021. Twelve years of SAMtools and BCFtools.
833   Gigascience 10.
834   94.   Page AJ, Taylor B, Delaney AJ, Soares J, Seemann T, Keane JA, Harris SR. 2016.
835   SNP-sites: rapid efficient extraction of SNPs from multi-FASTA alignments. Microb
836   Genom 2:e000056.
837   95.   Wickham H. 2016. ggplot2: elegant graphics for data analysis. Springer.
838   96.   Deatherage DE, Barrick JE. 2014. Identification of mutations in laboratory-evolved
839   microbes from next-generation sequencing data using breseq. Methods Mol Biol
840   1151:165-88.
841   97.   Krupovic M, Dutilh BE, Adriaenssens EM, Wittmann J, Vogensen FK, Sullivan MB,
842   Rumnieks J, Prangishvili D, Lavigne R, Kropinski AM, Klumpp J, Gillis A, Enault F,
843   Edwards RA, Duffy S, Clokie MR, Barylski J, Ackermann HW, Kuhn JH. 2016.
844   Taxonomy of prokaryotic viruses: update from the ICTV bacterial and archaeal viruses
845   subcommittee. Arch Virol 161:1095-9.
846   98.   Adriaenssens E, Brister JR. 2017. How to Name and Classify Your Phage: An Informal
847   Guide. Viruses 9.
848   99.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment
849   search tool. J Mol Biol 215:403-10.
850   100.   Meier-Kolthoff JP, Auch AF, Klenk HP, Goker M. 2013. Genome sequence-based
851   species delimitation with confidence intervals and improved distance functions. BMC
852   Bioinformatics 14:60.
853   101.   Meier-Kolthoff JP, Goker M. 2017. VICTOR: genome-based phylogeny and classification
854   of prokaryotic viruses. Bioinformatics 33:3396-3404.

855   102.   Nei M, Gojobori T. 1986. Simple methods for estimating the numbers of synonymous
856   and nonsynonymous nucleotide substitutions. Mol Biol Evol 3:418-26.
857   103.   Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013. MEGA6: Molecular
858   Evolutionary Genetics Analysis version 6.0. Mol Biol Evol 30:2725-9.
859   104.   Taniguchi H, Sato K, Ogawa M, Udou T, Mizuguchi Y. 1984. Isolation and
860   characterization of a filamentous phage, Vf33, specific for *Vibrio parahaemolyticus*.
861   Microbiology and immunology 28:327-337.
862   105.   Chang B, Yoshida S, Miyamoto H, Ogawa M, Horikawa K, Ogata K, Nishibuchi M,
863   Taniguchi H. 2000. A unique and common restriction fragment pattern of the nucleotide
864   sequences homologous to the genome of vf33, a filamentous bacteriophage, in
865   pandemic strains of *Vibrio parahaemolyticus* O3:K6 O4:K68, and O1:K untypeable.
866   FEMS Microbiol Lett 192:231-6.

Figure1

Figure2

Figure 3

Figure4

Figure5