Title: Long-term citizen science data reveals environmental correlates of tropical tree

flowering at the regional scale

Authors: Krishna Anujan[1,2]*, Jacob Mardian[3], Carina Luo[3], Ramraj Rajakumar[3], Hana Tasic[3],

Nadia Akseer[3], Geetha Ramaswami[1]

Affiliations:

[1] SeasonWatch, Nature Conservation Foundation, 1311, "Amritha", 12th Main, Vijayanagar 1st

Stage, Mysore 570017, India

[2] Department of Ecology, Evolution and Environmental Biology, Columbia University in the

City of New York, 1110 Amsterdam Ave, New York, NY 10025, United States

[3] Modern Scientist Global, 110 James Street, Suite 101, St. Catharines, Ontario L2R 7E8,

Canada

* corresponding author. Email: krishna.anujan@gmail.com

OPEN RESEARCH STATEMENT

ABSTRACT

Tropical tree reproductive phenology is sensitive to changing climate, but inter-individual and

interannual variability at the regional scale is poorly understood. While large-scale and long-term

datasets of environmental variables are available, reproductive phenology needs to be measured

in-site, limiting the spatiotemporal scales of the data. We leveraged a unique dataset assembled

by SeasonWatch, a citizen-science phenology monitoring programme in India to assess the

environmental correlates of flowering in three ubiquitous and economically important tree

species - jackfruit, mango and tamarind - in the south-western Indian state of Kerala. We

explored (i) seasonal patterns in the flowering status of trees (ii) environmental correlates of

flowering onset considering only trees with consecutive observations and (iii) spatiotemporal

patterns in these environmental correlates to aid future hypotheses for changing phenology

patterns. We used 165006 phenology observations spread over 19591 individual trees over 9

years. We first used bootstrapped circular statistics that accounts for observation biases in time to

examine consistency in seasonality of flowering status over the whole season, and flowering

onset across years and trees. Similar to results from cohort-based tree monitoring, we

demonstrate seasonality in flowering status and onset across species, but also report large

interannual and inter-individual variability. We then used used generalized linear mixed models

with remotely sensed observations (ERA5-LAND) to show that some of the interannual variation

in flowering onset across individuals was associated with environmental variables. Soil moisture,

minimum temperature and solar radiation had significant associations with the onset of flowering

but these effects were heterogeneous across species and habitats across Kerala. Our results

become increasingly important in the face of large spatiotemporal change in the climate of this

landscape and other tropical regions. We demonstrate the potential and limitations of citizen-

science observations in making and testing predictions at scale for predictive climate science in

tropical landscapes.

INTRODUCTION

Tree reproductive phenology is altered by a changing climate, affecting species interactions and ecosystem functions and services (Cleland et al. 2007, Mora et al. 2015). Flowering in trees is triggered by environmental cues like rainfall events, dry spells, and photoperiod, and is therefore sensitive to a changing environment (Chapman et al. 2005, Chang-Yang et al. 2016, Pau et al. 2018). Changes in onset, persistence, and quantity of reproductive phenophases can alter flower and fruit availabilty in the landscape, affecting interactions with other species like pollinators, frugivores, and seed dispersers (Visser and Both 2005, CaraDonna et al. 2014, Morellato et al. 2016). Consequently, for wild populations of trees, changes in phenology can influence regeneration dynamics and population persistence. Changing phenology can also have socio-cultural impacts, through altering food production and creating mismatches in cultural understanding of tree behaviours (Mora et al. 2015, Morellato et al. 2016). While landscape-level phenological shifts in temperate regions have been studied over decades, large gaps remain in ecological documentation and predictions for shifts in the reproductive phenology of tropical trees.

In tropical landscapes where trees grow and reproduce year-round, environmental cues for phenological change can be spatially and temporally diffuse (Parmesan 2007, Chen et al. 2018, Abernethy et al. 2018).   There is a growing understanding of the influence of environmental factors on leaf phenology of tropical species, but information on reproductive phenology is lacking (Chaturvedi et al. 2021, Devi et al. 2023). Cohort monitoring of trees has shown that

temperature, precipitation, and solar radiation (e.g. cold days, dry spells, total accumulated

rainfall and soil moisture) affect the onset and persistence of flowering phenology in tropical

trees (Singh and Singh 1992, Brearley et al. 2007, Pau et al. 2013, Diez et al. 2014, Wright and

Calderón 2018). Interannual rainfall variability - high rainfall years and drought years including

cycles like the El Niño Southern Oscillation and the Indian Ocean Dipole - can result in

interannual variability in tree phenophase onset and intensity (Medway 1972, Zimmerman et al.

2007, Chang-Yang et al. 2016, Chapman et al. 2018, Dunham et al. 2018, Pau et al. 2018).

Furthermore, climate change has resulted in altered rainfall and temperature regimes as well as

extreme events in tropical environments, increasing inter-annual variability (Mora et al. 2013).

To predict future phenological shifts in the tropics, an important first step is to understand the

extent of environmental controls on tropical tree reproductive phenology.

Environmental cues could affect phenology with a lag time, interact with habitat factors and

affect individuals differently, necessitating large-scale and long-term data. In tropical species, at

least six years of data is necessary to accurately detect phenology in regularly flowering species,

with much more for noisy species (Bush et al. 2017). Species across habitats may have different

responses to the same environmental cues, with differences in timing due to urbanisation (Neil

and Wu 2006), variation in synchrony with latitude (Borchert et al. 2004) or with distance to

coast (Vanderplank and Ezcurra 2016). Tropical tree phenology also exhibits large variation

among individuals, necessitating large spatial samples besides long-term data for a process-based

understanding(Satake et al. 2013, CaraDonna et al. 2014, Li et al. 2021). A recent review of

reproductive phenology studies from the Neotropics showed that across 218 datasets, the

majority of datasets lasted 2 years or less, and only 10.4% monitored > 15 individuals per species

(Mendoza et al. 2017). Across the tropics, phenology of ubiquitous species have rarely been

recorded systematically, hindering the potential to make large-scale predictions of climate

impacts of tree-associated ecosystem services (Piao et al. 2019, McDonough MacKenzie et al.

2020). In the absence of robust datasets, potentially high intraspecific variation in reproductive

phenology makes it difficult to parametrise models that span spatial scales of environmental

change.

Citizen science allows access to longitudinal data across large spatial scales that is difficult to

achieve through other sampling techniques. Citizen science data has led to novel understanding

of phenological patterns in other taxa - observations of migratory bird populations across North

America have been useful in documenting phenological shifts, spatial differences in sensitivities

to changing climate, as well as disentangling drivers of change (Hurlbert and Liang 2012, Kelly

et al. 2016, Haas et al. 2022). Other efforts, like the USA National Phenology Network, have

helped quantify advancing spring behaviours of plant species in North America (Schwartz et al.

2012, Brunsdon and Comber 2012). SeasonWatch, a citizen science phenology monitoring

programme has been collecting data on several phenophases for common, widespread tree

species across India (Ramaswami et al. 2019, 2021). Each week, observers record flowering,

along with other phenophases, on specific trees consistently using a uniform protocol that

records both presence and intensity of a phenophase (Fig 1a). This dataset, with substantial

contributions since 2014, provides a unique opportunity to build baselines for phenology in the

Indian subcontinent. In the past, some analyses of these data have revealed patterns of change in

at least one species - Cassia fistula - over the time period of observation (Ramaswami and

Quader 2018, Ramaswami et al. 2021). These common species, spread across ecoregions and

habitats, can add to phenological understanding at large scales, complementing focussed,

localised studies.

Here, we investigated the environmental correlates of flowering in common tree species in India

at the individual tree scale using SeasonWatch data. We used 165006 phenology observations

across 19591 individual trees over 9 years from tree species in the southwestern Indian state of

Kerala, and correlated these with remotely-sensed environmental variables using machine

learning and generalised linear mixed model approaches. We asked (i) what are the seasonal

patterns in flowering onset and flowering status of common trees across Kerala and what is the

extent of interannual and interindividual variation? (ii) What are the environmental correlates

associated with tree-level flowering onset? (iv) How are relevant environmental cues changing

spatiotemporally across the landscape? We focussed on flowering of three species with the

greatest number of individuals and observations across Kerala - jackfruit (*Artocarpus*

heterophyllus), mango (*Mangifera indica*) and tamarind (*Tamarindus indica*).   We expected (i)

interannual and interindividual variation in flowering status and onset to varying degrees across

the three species, (ii) based on research using cohort sampling of trees, we expected temperature

and precipitation patterns to influence the onset of flowering. (iii) However, given the large

spatial extent of our dataset, we also expected that tree flowering responses to environmental

correlates would vary with elevation, aspect, and degree of urbanisation. (iv) Finally, through

exploratory analysis, we expected to observe and quantify spatiotemporal variation in

environmental correlates of flowering phenology in Kerala.

METHODS

STUDY AREA, SPECIES AND SEASONWATCH DATA

We analyzed phenology and environmental data from the south-western Indian coastal state of

Kerala. Bounded between 8° 18' N, 12° 48' N, 74° 52' E and 72° 22' E, Kerala lies along the

Arabian Sea with a tropical monsoon climate. The total population of the state is over 3 million,

spread over an area of 38,863 km^2. The state receives an average annual rainfall of over 3000mm

via two major branches of the Indian monsoon system - the southwest (June-September) and

northeast (October-December) monsoon. Citizen scientists from the state have contributed

substantially to phenology data; across all SeasonWatch data (SeasonWatch Citizen Scientist

Network, 2022, Ramaswami (2022)), 263,288 observations across species and categories, and

including 93.23% of observations of top three species are from Kerala. These data span a large

range of the existing environmental variability in the state in terms of rainfall, temperature,

elevation.

We analyzed flowering phenology in three frequently observed species in the SeasonWatch

dataset - *Artocarpus heterophyllus* (henceforth "jackfruit"), *Mangifera indica* (henceforth

"mango") and *Tamarindus indica* (henceforth "tamarind") from January 2014 - July 2022. All

three tree species are grown across the state in orchards, homes, and also occur wild. As part of

SeasonWatch, observers register one or more "regular" trees from a list of suggested species on a

mobile phone application or web browser and record phenology on a weekly basis. The project

also allows for "casual" or one-time observations of trees (outside an observer's "regular" list),

often during coordinated events, expanding the spatial coverage of the data. All observations to

SeasonWatch comprise estimates of quantity ("none", "few", "many") of leaf (young, mature,

dry), flower (bud, open flower) and fruit (unripe, ripe) phenophases of a tree. For each of the 7

phenophases, the standard protocol for reporting involves dividing the canopy of a tree into

thirds, reporting as 'none' if it is not observed, as 'few' if a phenophase is observed only in 1/3rd

of the canopy, and as 'many' if a phenophase is observed on more than 1/3rd of the canopy (Fig

1a). SeasonWatch data is contributed by voluntary participants; the programme recruits citizen

scientists by first identifying regional partner organisations, training the trainers in these

organisations who then recruit and train participants (e.g. teachers, students and community

members), thus amplifying the network reach.   The project minimises inter-observer biases and

errors in observation through workshops and training as well as one-on-one follow-up and

troubleshooting. Pertinently, the flowering phases of the common trees chosen for analysis are

distinctly observable with the naked eye, and familiar to participants, lending confidence to

community data. The data distribution across species and years is provided in Table 1, Appendix

S1: Fig S1 and Appendix S1: Fig S2.

165 **DATA CLEANING AND PROCESSING**

166 We cleaned and processed SeasonWatch data for jackfruit, mango and tamarind from 2014-2022

167 to create two different data subsets - flowering status and flowering onset. To maximise inclusion

168 of data points, we included all "regular" and "causal" observations that we were able to reliably

169 assign and validate locations (see Appendix S1: Section S2). We recoded the none/few/many

170 observations of bud and flowers into presence/absence of flowering by coding an observation as

171 "1" if it was recorded as "few" or "many" for buds or flowers, and "0" otherwise. We chose to

172 combine the bud and the flowering phenophases to minimise observation errors in these species

173 where flowers are small and/or greenish and can be difficult to distinguish from buds, especially

174 in taller canopies. Our choice of binary variable also reduces potential user subjectivity in the

175 categorising of "few" and "many".

176 *Flowering status*

177 Flowering status included all observations across trees and years in the dataset where presence or

178 absence of flowering (bud or flower) was recorded. We assembled a final flowering status dataset

179 of 165006 observations across 19596 trees; 7927 trees of jackfruit, 9919 of mango and 1750 of

180 tamarind across the state of Kerala (Fig 1, Appendix S1: Table S1). The number of observations

181 varied by week and year, with numbers generally increasing in the later part of the study period

182 because of increased participation in the programme. (Appendix S1: Fig S1) Most trees in this

183 dataset were observed few times, but several trees had long, uninterrupted flowering status

184 timeseries for one or multiple years (Appendix S1: Fig S4).

Flowering onset

The onset of flowering is defined as the first week in a year when a tree switches in status from

non-flowering to flowering. To estimate true flowering onset at the tree level, we filtered the

flowering status dataset to exclude all observations for which there was no flowering status

observation in the preceding week, thus minimising observation bias. Since we were interested in

the drivers of the switch from non-flowering to flowering, we further filtered this dataset to only

include observations when the tree was observed to be non-flowering in the previous week. For

trees that were observed to have multiple onsets in a year (starting from June, the beginning of

the wet season), we retained only the first onset observation. This reduced the dataset to 62721

observations with an immediate predecessor, across 2442 trees, out of which, 2638 were true

onset observations (0, 1) and the remaining were status quo observations (0, 0) (Table 1,

Appendix S1: Fig S2). By filtering to only observations with a non-flowering predecessor, the

flowering onset dataset has minimal temporal autocorrelation among observations, allowing the

application of mixed modelling frameworks.

ENVIRONMENTAL DATASETS

We derived seven potential spatiotemporal predictors of flowering from remotely sensed data -

three related to temperature (minimum, maximum and mean), three to precipitation (total

precipitation, soil moisture and the number of consecutive dry days), and solar radiation. We

used daily measurements of precipitation, temperature, solar radiation and soil moisture for the

entire soil column from the ERA5-Land dataset at the 11.1 km spatial resolution and

summarised these to the fortnight (14 days) preceding each observation using Google Earth

Engine (Table 2). Monthly and quarterly summaries were highly correlated to the fortnightly

value (Appendix S1: Fig S3) and hence we used this finest temporal resolution (fortnight) for the

analysis. Along with spatiotemporal predictors, we calculated spatial factors - elevation, aspect

and degree of urbanisation - at each tree location that might potentially act as modifiers for the

temporal predictors (Table 2). We used a Digital Elevation Model created from the Shuttle Radio

Topography Mission (SRTM) at the 30 m resolution to calculate elevation, slope and aspect at

the tree location. Further, we used the NASA nightlights layer from 2016 at the 500 m resolution

as a proxy for the degree of urbanisation (Sutton et al. 2001).

DATA ANALYSIS

We analysed seasonality in flowering status and flowering onset at the population level using

circular statistics and then modelled predictors of individual-level flowering onset across years of

observations using generalised linear mixed models. We modelled flowering seasonality and the

ecological predictors of flowering onset separately for each of the three species, as these species

are physiologically and phylogenetically distinct (belonging to families - Moraceae,

Anacardiaceae, and Fabaceae) and may therefore show distinct flowering seasonality and

heterogeneous responses to the same environmental cues. Finally we examined spatiotemporal

patterns in environmental predictors across Kerala using hotspot analyses. Analyses were

performed using R version 4.1.3 and ArcGIS Pro version 3.0.2 (R Core Team 2022).

CIRCULAR STATISTICS TO ESTIMATE POPULATION FLOWERING SEASONALITY AND ONSET

SEASONALITY

We used a newly developed bootstrap method to assess seasonality in flowering in the three

species across the state of Kerala (Willig et al. 2024). Seasonality in flowering is best described

using circular statistics like circular mean and the Rayleigh statistic which describe the clustering

of the data in circular axes (here, week of year) (Morellato et al. 2010). Unlike long-term field

monitoring datasets which have fixed cohorts that are repeatedly monitored during the entire

monitoring period, citizen science datasets like SeasonWatch can have a) different start and end

dates for different individuals b) different timeseries lengths and c) larger proportion of

missingness, despite greater spatial spread. This unique data structure leads to non-uniform data

distribution across the temporal extent and makes this analogous to capture sampling of random

individuals in a population at a fixed frequency, violating the uniformity assumptions of

traditional circular statistics (Willig et al. 2024). Using a bootstrapping method, we calculated

the circular mean of flowering status and flowering onset for each year of study. We sampled

observations with replacement for each week of study to a constant N, equal to the average

weekly observation for that species, making the sample distribution uniform across the year.

With this uniform sample, we calculated circular mean, Rayleigh statistic (a measure of temporal

clustering) and the p-value of the statistic. We repeated the sampling for 1000 iterations and

calculated the bootstrap mean and deviation. To examine interannual consistency in flowering

243 onset across individuals, we calculated mean week of flowering onset and standard deviation for

244 each tree across years all years of onset observations.

245 **GENERALISED LINEAR MIXED MODELS FOR FLOWERING ONSET AT THE INDIVIDUAL LEVEL**

246 We examined the effects of environmental variables on the onset of flowering in the three species

247 using generalised linear mixed effects models for interactions between variables and variable

248 collinearity. For each of the three species modelled, the time-varying predictors were identified

249 by picking the most influential variables from random forest (machine learning) models using all

250 environmental variables as predictors (Appendix S1: Section S4). For each species, we picked

251 most important variable in the temperature group (among minimum, maximum and mean), the

252 most important variable in the precipitation group (among total precipitation, number of dry days

253 and soil moisture), and solar radiation. For all species, the most influential temperature variable

254 was minimum temperature and precipitation variable was soil moisture. This grouping allowed

255 us to test the effects of temperature, precipitation and solar radiation, while avoiding including

256 highly correlated variables that can violate mixed modelling assumptions.

257 We accounted for interactions between time-varying and static predictors, excluding known,

258 correlated variable combinations. Specifically, we excluded interactions between elevation and

259 minimum temperature, aspect and precipitation (the Western Ghats range in Kerala affects

260 rainfall patterns), degree of urbanisation and minimum temperature (known heat island effects in

261 urban areas), degree of urbanisation and soil moisture (potentially associated with heat island

262   effects). Further, to account for data structure in space and time, we included random effects of

263   tree and the spatial grid in which it was located.

264                          $$phenophase_{l,t}$$

265              $$\sim minimum.temperature_{l,t} + soil.moisture_{l,t} + solar.radiation_{l,t}$$

266              $$+ elevation_l * soil.moisture_{l,t} + elevation_l * solar.radiation_{l,t} + aspect_l$$

267              $$* minimum.temperature_{l,t} + aspect_l * soilmoisture_{l,t} + +urbanisation_l$$

268              $$* solar.radiation_{l,t} + (1|GridID)$$

Where minimum temperature, soil moisture and solar radiation are measured at the location l for

the fortnight before the observation t, elevation, aspect and urbanisation are spatial predictors

tied to the tree location l. The model includes two random effects on the intercept; $TreeID$ to

account for repeated measures on the same tree and $GridID$ to account for spatial

autocorrelation.

We modelled the onset of the phenophase using a generalized linear mixed model with a

binomial family of errors using the *package* lme4 (Bates et al. 2015). For all models, we mean-

centred all predictor variables at 0 and scaled it to a distribution with standard deviation 1 for

ease of comparing the effect sizes. We tested the fit of these models using conditional and

marginal R^2 using packages *performance* and *sjPlot* (Lüdecke et al. 2021, Lüdecke 2023). We

used a cross-validation technique, leave-one-out cross-validation (LOOCV), to test the predictive

power of our model for "out-of-bag" observations. In this method, we ran each model 9 times,

systematically leaving one year of observations out each time. We then used the model

coefficients to predict these "new" observations and tested their accuracy, and then tested for

reliability of predicted values using F1 scores. F1 scores are good indicators of the distribution of

true and false positives and negatives by taking the harmonic mean of precision and recall

(Hossin and Sulaiman 2015, Grandini et al. 2020).

HOTSPOT ANALYSIS

We analysed spatiotemporal patterns of environmental variables that affect flowering onset in

Kerala, with an intention to identify directional changes or spatial shifts. Hotspot analysis allows

the identification of such spatiotemporal patterns using the entire timeseries of environmental

data across the years measured. A hot spot/cold spot is defined as a location which has high (or

low) value for an environmental variable, like temperature, precipitation or soil moisture, and is

surrounded by other areas with similar values; a spatial cluster. Using fortnightly aggregated

environmental variables from 2014-2022, we ran emerging hotspot analysis in ArcGIS Pro for

minimum temperature, mean soil moisture and total solar radiation, the three variables used for

the GLMM analysis, across Kerala. We used a combination of two statistical measures 1) the

Getis-Ord Gi* statistic to identify hot spots and cold spots of climate measurements for each

time step (i.e., fortnightly), and 2) the Mann Kendall trend test to examine how hot spots and

cold spots have evolved over time (Mann 1945, Sen 1968, Kendall et al. 1987, Getis and Ord

1992). For further details on methods, refer Appendix S1: Section S7.

RESULTS

Jackfruit, mango and tamarind trees exhibit differing patterns of flowering seasonality and

substantial variation among years in mean flowering and flowering onset in the state of Kerala.

Across 9 years of data (2014 to 2022), bootstrapped circular mean of flowering status showed

that mean flowering week was week 4.88 for jackfruit, 4.68 mango and 25.4 for tamarind.

Temporal clustering in flowering across all years was low, but jackfruit and mango showed

higher clustering in flowering across the population than tamarind; bootstrapped Rayleigh

statistic, $z = 0.29$ (95% CI = 0.29, 0.3) for jackfruit, 0.41 (95% CI = 0.41, 0.42) for mango and

0.1(95% CI = 0.09, 0.11) for tamarind. Correspondingly, mean week of flowering onset was

week 0.57 for jackfruit, 0.4 for mango and 19.45 for tamarind. Patterns of temporal clustering in

flowering onset mirrored flowering status; Rayleigh statistic, $z = 0.5$ (95% CI = 0.48, 0.52) for

jackfruit, 0.52 (95% CI = 0.5, 0.54) for mango and 0.19 (95% CI = 0.14, 0.24) for tamarind.

Across all three species, trees were observed to be flowering at all times of the year; no week had

a zero proportion of trees flowering (Fig 1).

Across 8 full years of observations (excluding the first half of 2014), mean week of flowering

status and flowering onset showed interannual variation, but there was low variation in temporal

clustering of flowering (Fig 2). Mean flowering week ranged from week 2.73 in 2019 to 7.8 in

2016 for jackfruit, 2.39 in 2021 to 7.35 in 2016 for mango and -25.15 in 2019 to 25.47 in 2015

for tamarind (Fig 2a). Clustering in flowering status was relatively similar across the years (Fig

2a). For jackfruit, bootstrapped Rayleigh statistic of flowering status ranged from $z = 0.27$ (95%

CI = 0.26, 0.28) in 2019 to z = 0.25 (95% CI = 0.23, 0.27) in 2016. Values for mango ranged

from z = 0.43 (95% CI = 0.42, 0.45) in 2021 to z = 0.36 (95% CI = 0.35, 0.38) in 2016. Values

for tamarind ranged from z = 0.1 (95% CI = 0.08, 0.12) in 2019 to z = 0.14 (95% CI = 0.08, 0.2)

in 2015. Flowering onset typically occurred 4-6 weeks before mean flowering week and showed

more clustering within species than flowering status (Fig 2). Mean flowering onset across the

years ranged from week -3.27 in 2017 to 3.96 in 2019 in jackfruit, -0.7 in 2019 to 3.57 in 2016 in

mango, -23.49 in 2018 to 25.37 in 2015 in tamarind (Fig 2b). Temporal clustering in flowering

onset was less variable across the years; with tamarind having lower onset clustering than

jackfruit and mango, during most years. Clustering within species ranged had a smaller range,

lowest clustering in jackfruit was observed in 2017 with z = 0.49 (95% CI = 0.42, 0.07) and

highest in 2019 with z = 0.51 (95% CI = 0.46, 0.38) (Fig 2b). For mango, lowest and highest

clustering of onset was observed in 2019 with z = 0.53 (95% CI = 0.47, 0.1) and 2016 with z =

0.55 (95% CI = 0.44, 0.47) respectively (Fig 2b). For tamarind, lowest clustering was observed

in 2018 with z = 0.24 (95% CI = 0.08, 0.12) and highest in 2015 with z = 0.44 (95% CI = 0.08,

0.2) (Fig 2b). There were no observable directional trends with time in timing or clustering of

flowering.

Individual trees across all three species exhibited large variation in mean flowering onset week

across the years as well as low consistency in the date of onset (Fig 3). Across all trees with

observed true onset across the years, mean flowering onset week was distributed widely across

the year for all three species (Fig 3a). Most trees of jackfruit showed a mean flowering onset

week in the calendar year of -6, mango in week -12 and tamarind in week -20 (Fig 3a). However,

trees showed large variation in onset week across years (Fig 3b). For trees with at least three

years of onset observations, median standard deviation of flowering onset week was 6.91 for

jackfruit, 6.39 for mango and 15.94 for tamarind.

PROXIMATE ENVIRONMENTAL CUES FOR FLOWERING ONSET ON A TREE

Flowering onset at the tree level across Kerala was predicted by minimum temperature, soil

moisture and solar radiation but in varying magnitudes and directions across species (Fig 4). For

jackfruit and mango, flowering onset was associated with lower minimum temperature and drier

soils while for tamarind with higher minimum temperatures and wetter soils (Fig 4). Solar

radiation had a weak positive effect on the probability of flowering in mango (Fig 4). GLMMs

models of flowering onset had substantially better fits for jackfruit and mango than tamarind; the

conditional R^2 for the models, a metric of model fit including random effects, for jackfruit,

mango and tamarind were 0.228, 0.242 and 0.131. Marginal R^2, that only includes predictions

using fixed effects, was 0.079, 0.102 and 0.026 for jackfruit, mango and tamarind, suggesting

poor predictability from environmental variables alone, especially for tamarind. This suggests

that fixed effects of environmental variation alone explained only a small proportion of the

variation across flowering onset observations.

The ICC of the models were 0.162, 0.156 and 0.107 suggesting that random effects were not

very clustered. To assess the predictive power of the model for out-of-bag data, we assessed the

agreement between the predicted and observed values from the onset model for one year of the

data that was withheld from modelling across 9 LOOCV runs. Coefficient across these runs were

similar in magnitude and direction (Appendix S1: Section S6). Pearson's correlation coefficients

for predicted vs observed for jackfruit and mango across these runs 0.011, 0.031. The average

mean square errors, respectively were 0.043, 0.041. F1 scores were 0.002, 0.007. The model was

unable to predict flowering onset in tamarind with all predictions tending to non-flowering.

Elevation and degree of urbnaisation modified the influence of environmental factors on the

onset of flowering in jackfruit and mango trees (Fig 4). Drier soils were associated with higher

probability of flowering onset in mango, but this effect was stronger in lower elevations. In

parallel, solar radiation increased the probability of flowering onset in mango trees, but this

effect was dampened in more urbanised areas (Fig 4). In jackfruit trees, the influence of solar

radiation as a cue for flowering onset was greater in lower elevation and more urban areas (Fig

4). However, main effects of solar radiation were not significant at the 90% confidence level,

making interpretations of these interactions difficult. No interactions were significant for onset of

flowering in tamarind trees. However, the model was a poor fit, and therefore, the possibility of

significant interactions with more data cannot be ruled out.

SPATIOTEMPORAL PATTERNS IN ENVIRONMENTAL VARIABLES

Locations across Kerala are experiencing heterogenous change in climatic conditions. Emerging

hotspot analysis revealed spatiotemporal patterns in the three variables assessed, minimum

temperature, soil moisture and solar radiation, in the time period when SeasonWatch data was

collected (Fig 5). Aggregated across the state, these environmental variables did not show

significant overall directional trend of change, defined as new or intensifying hotspot/coldspot, at

the spatial scale assessed, from 2014-2022. Along the west to east elevational gradient in Kerala,

minimum temperature had persistent hotspots in coastal locations, while inland sites towards the

east were persistent cold spots, potentially due to higher elevation (Fig 5). Emerging hotspot

analysis for soil moisture revealed a persistent hotspot in the south centre of the state. This

hotspot corresponds to the eastern parts of the districts of Ernakulam and Kottayam, and the

western parts of Idukki. For minimum temperature and soil moisture, pixels around the

persistent hotspots, show varying and heterogenous trends in hotspots or coldspots. Mid-

elevation sites in Kerala between the persistent coldspot and hotspot for minimum temperature

have been classified as oscillating, sporadic or historical hot/coldspots, suggesting heterogenous

change. Similarly for soil moisture, areas around the persistent hotspot show sporadic or

oscillating hotspots. No directional patterns were discernible for solar radiation.

DISCUSSION

Altered temperature and precipitation patterns are expected to affect phenological responses of

tropical tree species, but their variability across large spatial scales is rarely quantified and poorly

understood. We used a unique citizen science dataset from the southwestern state of Kerala in

India (Fig 1), unparalleled in scale for the tropics, to examine seasonality in flowering in

common tropical tree species and the extent and correlates of interannual and interindividual

variability. Our results show that over 9 years of weekly observations, timing of flowering status

and flowering onset across three species shows 4-6 weeks of variation across years at the

population level (Fig 2). This interannual variation in the timing of flowering was not mirrored in

temporal clustering flowering observed at the population level, which remained relatively

constant, but was reflected in individual trees that showed substantial interannual variation in

timing of flowering onset(Fig 3). We show that some of this individual-level variation in

phenology of flowering onset across the years is associated with recent, local climatic conditions,

such as soil moisture and minimum temperature, the effects of which can be different along

elevation and urbanisation gradients (Fig 4). These environmental cues associated with

phenology of these common trees, however, are changing across Kerala (Fig 5), with potential

consequences for phenology and associated ecosystem services. Our results show the potential of

citizen science datasets to provide crucial insights at spatial and temporal scales relevant for

adaptive ecosystem management.

Our use of citizen science data and associated temporal data gaps limits the use of some common

methods like Fourier transformation to disentangle seasonal patterns from long-term drivers at

the individual tree level. Cohort-based long-term monitoring of tree populations has led to

important understanding of the influence of seasonal precipitation and interannual variability

through ENSO events, droughts etc. on reproductive phenology of trees in tropical sites (Wright

et al. 1999, Harrison 2000, Sakai and Kitajima 2019). Given the lack in uninterrupted timeseries

at the individual tree level, we were unable to disentangle individual variation in phenology from

temporal patterns for long timescale. However, our study spans nine years of observations, more

than the minimum recommended 6 years for tropical species (Bush et al. 2018), and the spatial

extent of the dataset is much larger compared to the limitations of long-term cohort monitoring

(often restricted to a site) allowing us to extend inferences to large spatial scales. Examining

flowering observations across a wide variation in habitats also revealed that trees in high

elevation or urban areas respond differently to environmental cues, adding to prevalent

understanding of the contextual nature of flowering cues (Neil and Wu 2006, Babweteera et al.

2018, Kabano et al. 2020). Such accumulating broad-scale datasets led by citizen scientists could

be important complements to finer scale long-term expert-led monitoring to understand shifting

phenological baselines across species ranges.

Extending our results across tropical trees and landscapes poses certain limitations because of

some broad ecological and functional similarities in the three species studied. All three species

studied were dioecious, evergreen, widespread, insect-pollinated, bear fleshy fruits, and

cultivated in home gardens (Table S1). Notably, jackfruit trees in the dataset showed high

tendencies for multiple instances of flowering onset within a year, and tamarind showed low

seasonality in flowering overall (Fig 2). Mixed models of flowering onset showed that

environmental predictors explained only ~10% of variation in probability of switch to flowering

status, while including random effects of sites explained ~20% of the variation across all three

species. Since these are cultivated species, the differences in varieties among individuals has a

known influence on phenological variation (Makhmale et al. 2016, Singh et al. 2021).

Applicability of these inferences across diverse tropical species with different leaf and

reproductive phenology could pose challenges. Reproductive strategies in tropical trees are often

diverse and may have long-term evolutionary function in addition to short-term ecological

response (Singh and Kushwaha 2006). Borchert et al. (2004) showed that flowering seasonality

in tropical species can be plastic across biome ranges, with greater periodicity in the drier forests

across species. We show that even in low latitudes with low seasonality, flowering status and

flowering onset within species show strong periodicity and interannual variation associated with

climatic variables. Given the phylogenetic distinctiveness among the three species, we are

encouraged to conclude that these conclusions could apply over a broader set of species.

Our results become pertinent under existing and future climatic variability in the tropics, and the

potential impacts on the reproductive phenology of commercially important tropical trees.

Beyond the timescales of our study, much of the global tropics, is projected to move into a

temperature and precipitation regime that has no analog globally, making it difficult to predict

and plan for ecosystem responses (Dahinden et al. 2017). Across the Indian subcontinent,

precipitation and temperature variability has increased in the past century and is projected to

continue to increase (Power et al. 2013, Dahinden et al. 2017, Zhang and Fueglistaler 2019).

Moreover, altered climatic processes such as the El Nino Southern Oscillation and Indian Ocean

dipole are affecting weather with consequences for Indian ecosystems (Revadekar et al. 2009,

Wang and Wang 2014, Gadgil and Francis 2016). However, climate change predictions and

action for the tropics are still largely at coarse spatial scales, making local forecasting and

mitigation difficult (Dessai et al. 2009, Shepherd and Sobel 2020). Across Kerala, our results

from emerging hotspot analyses show spatiotemporal patterning in climate variables associated

with flowering phenology in common tree species (Fig 5). An important future direction for

phenological studies would be to understand intraspecific variation across space and forecast

phenological responses in these novel climates.

Our study using repeat citizen-science data and individual based modelling at spatial and

temporal scales of climate modelling, represents a significant first step in combining tropical

ecology with climate science beyond broad generalisations.   Despite recent remote sensing

advances that aid large-scale and repeat measurement of ecosystem structure and change,

measurement of tropical reproductive phenology remains challenging (Davis et al. 2022, Lee et

al. 2023). Species may have inconspicuous flowers and fruits or these phenophases may coexist

with leaves, requiring repeat on-ground measurements that are extremely difficult to organise at

the large-scale. Citizen science may be one of the few ways to acquire such data, and we have

leveraged such a unique dataset compiled by SeasonWatch to predict the onset of a reproductive

phenophase (Abernethy et al. 2018, McDonough MacKenzie et al. 2020, Ramaswami et al.

2021). Broadly, we demonstrate the potential of citizen science observations in improving the

state of our knowledge in climate science and infusing local ecology into climate-adaptive

management. Predictive modelling informed by variation among individuals and sites is a crucial

next step to phenology research as spatiotemporal patterns in climate variables change across

tropical landscapes. Our study thus demonstrates the potential of citizen science in filling data

gaps from expert-led site-specific monitoring and providing predictions at the large-scale to aid

locally relevant climate policies.

ACKNOWLEDGEMENTS

The authors would like to acknowledge the SeasonWatch team for support during this work,

especially Mohammed Nizar, who helped in the manual validation of parts of the data. We also

thank Karthik T, Anand M O, Suhel Quader and the NCF EPE team for discussions on data

management, analyses and interpretations. SW is grateful for the support of the Wipro

Foundation for their continued efforts.

RESEARCH ETHICS

The study did not deal with human subjects or data with personal identifiers. For SeasonWatch

data, we followed ethical protocol that was agreed on by the SeasonWatch Citizen Scientist

Network.

AUTHOR CONTRIBUTIONS

Funding for this work was acquired by GR and NA, while the study was conceptualised by GR

and KA. Phenology data was collected by the SeasonWatch Citizen Scientist Network. KA

cleaned and validated the phenology data while JM accessed and collated the geospatial data.

KA, JM, CL and RR performed the analysis led by KA and supervised by GR and NA, with

administrative support from HT. KA visualised the data and wrote the first draft; KA, JM and GR

edited the manuscript with support from all authors.

FUNDING SOURCES

The analysis was funded by the Booster Programme at the Patrick J McGovern Foundation.

SeasonWatch data collection was supported by the Wipro Foundation.

CONFLICT OF INTEREST

The authors have no conflict of interest to declare.

TABLES

Table 1: Data distributions. *Description of the SeasonWatch dataset for jackfruit, mango and*

tamarind in Kerala from January 2014 to July 2022

	Artocarpus heterophyllus	Mangifera indica	Tamarindus indica
Total number of observations	72,097	68,696	24,213
Total number of trees	7,927	9,919	1,750
Number of trees with single observations	6,270	8,128	1,277
Number of observations with no flowering at t-1	25,695	26,212	12,426
Number of trees in observations with no flowering at t-1	1,028	1,105	320

 Table 2: Environmental data. *Variables, data sources and transformations*

Datasets	Data Source	Resolution	Transformations used	Variables derived
Flowering observations	SeasonWatch	temporal: ~once every two weeks		Flowering binary and ordinal
	https://www.seasonwatch.in/	spatial: tree-level observations		
Precipitation	ERA5-Land	temporal: hourly, aggregated to fortnightly	Sum	Total precipitation, DryDays
	https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-land?tab=overview	spatial: 11.1 km grid		
Temperature	ERA5-Land	temporal: hourly, aggregated to fortnightly	Max, min, mean	Max temp, min temp, mean temp
	https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-land?tab=overview	spatial: 11.1 km grid		
Soil Moisture	ERA5-Land	temporal: hourly, aggregated to fortnightly	Mean (of total column depth at each time step), Mean (hourly -> Daily -> Fortnightly)	Mean SM
	https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-land?tab=overview	spatial: 11.1 km grid		
Solar radiation	ERA5-Land	temporal: hourly, aggregated to fortnightly	Sum (hourly -> daily -> Fortnightly)	SolarRad
	https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-land?tab=overview	spatial: 11.1 km grid		
elevation	SRTM	temporal: no change	terrain functions	Elevation, slope, aspect
	https://www.usgs.gov/centers/eros/science/usgs-eros-archive-	spatial: 30 m grid		

Datasets	Data Source	Resolution	Transformations used	Variables derived
	digital-elevation-shuttle-radar-topography-mission-srtm-1			
urbanisation	NASA nightlights	temporal: no change		urbanisation
	https://earthobservatory.nasa.gov/features/NightLights	spatial: 0.5 km grid		

FIGURE CAPTIONS

Figure 1: SeasonWatch protocol, species and map of observations. a) SeasonWatch

observation protocol for tree phenology - phenophase quantities 'none', 'few', 'many' roughly

correspond to the presence of a phenophase in 0, 30%, and >30% of the tree canopy b) Outline of

India with the state of Kerala marked in grey. c), d) and e) - map of observations of jackfruit,

mango and tamarind trees across Kerala, considered in this study along with illustrations of

flowering and fruiting phenophases in jackfruit, mango and tamarind respectively. Panels f), g)

and h) show seasonal patterns in flowering of the three species in Kerala from 2014-2022. Dots

represent mean proportion of trees flowering in that week and error bars represent standard

deviation across the years of study. Image credits: a) Sayee Girdhari for SeasonWatch;

illustration in c) and e) Neelam Modi for SeasonWatch; illustration in d) Mango (Mangifera

indica L.): fruiting branch with numbered sections of flower and seed. Chromolithograph by P.

Depannemaeker, c.1885, after B. Hoola van Nooten. Wellcome Collection. Source: Wellcome

Collection Licence: Public Domain Mark.

Figure 2: Mean date of flowering presence and flowering onset from 2014-2022 for the

species of interest in Kerala. Bootstrapped estimates of flowering presence from weekly

observations each year for a) jackfruit, b) mango, and c) tamarind. Bootstrapped estimates of

flowering onset (first definitive observation of flowering on any tree) from weekly observations

each year for d) jackfruit, e) mango, and f) tamarind. For a given year, arrow direction represents

the mean week and arrow length represents z-score, a measure of degree of clustering from a

bootstrapped Rayleigh test. See methods for details on how these were calculated.

Figure 3: Variability in flowering onset across years for the same tree from 2014-2022.

Distribution of mean week of flowering onset across years for trees with multiple onset

observations for a) jackfruit, b) mango, and c) tamarind. Distribution of the standard deviation of

flowering onset weeks for trees of a) jackfruit, b) mango, and c) tamarind.

Figure 4: Model coefficients for flowering onset from GLMM models Generalised linear

mixed effect models (with a binomial family of errors) modelling flowering onset; i.e. the

probability of observing flowering given no flowering at the previous timestep. Model includes

most influential temperature variable, precipitation variable from the ML models and solar

radiation. Model also includes all interactions barring known correlated combinations. Dots

reporesent mean effect and the lines represent 95% confidence intervals of the estimate. The

vertical grey line represents no-effect. Predictors with confidence intervals that do not overlap

with the no-effect line are considered significant; others are represented with grey dots.

Figure 5 - Hotspot analysis of environmental data using spatiotemporal data from 2014-

2022. Hotspot analyses for a) minimum temperature within a fortnight, b) soil moisture within

the fortnight and c) solar radiation within the fortnight within each grid of 11 km x 11 km spatial

resolution

REFERENCES

Abernethy, K., E. R. Bush, P.-M. Forget, I. Mendoza, and L. P. C. Morellato. 2018. Current

issues in tropical phenology: A synthesis. Biotropica 50:477–482.

Babweteera, F., A. J. Plumptre, G. S. Adamescu, L. P. Shoo, C. M. Beale, V. Reynolds, P. Nyeko,

and G. Muhanguzi. 2018. The ecology of tree reproduction in an African medium altitude rain

forest. Biotropica 50:405–417.

Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting Linear Mixed-Effects Models

Using **Lme4**. Journal of Statistical Software 67.

Borchert, R., S. A. Meyer, R. S. Felger, and L. Porter-Bolland. 2004. Environmental control of

flowering periodicity in Costa Rican and Mexican tropical dry forests. Global Ecology and

Biogeography 13:409–425.

Brearley, F. Q., J. Proctor, J. Proctor, Laszlo Nagy, L. L. Nagy, L. Nagy, G. Dalrymple, and B. C.

Voysey. 2007. Reproductive phenology over a 10-year period in a lowland evergreen rain forest

of central Borneo. Journal of Ecology 95:828–839.

Brunsdon, C., and L. Comber. 2012. Assessing the changing flowering date of the common lilac

in North America: A random coefficient model approach. GeoInformatica 16:675–690.

Bush, E. R., K. Abernethy, K. J. Jeffery, C. E. G. Tutin, L. J. T. White, E. Dimoto, J.-T.

Dikangadissi, A. S. Jump, and N. Bunnefeld. 2017. Fourier analysis to detect phenological cycles

using long-term tropical field data and simulations. Methods in Ecology and Evolution 8:530–

540.

Bush, E. R., N. Bunnefeld, E. Dimoto, J.-T. Dikangadissi, K. Jeffery, C. Tutin, L. White, and K.

A. Abernethy. 2018. Towards effective monitoring of tropical phenology: Maximizing returns

and reducing uncertainty in long-term studies. Biotropica 50:455–464.

CaraDonna, P. J., A. M. Iler, and D. W. Inouye. 2014. Shifts in flowering phenology reshape a

subalpine plant community. Proceedings of the National Academy of Sciences of the United

States of America 111:4916–4921.

Chang-Yang, C.-H., I.-F. Sun, C. H. Tsai, C.-L. Lu, and C.-F. Hsieh. 2016. ENSO and frost

codetermine decade-long temporal variation in flower and seed production in a subtropical rain

forest. Journal of Ecology 104:44–54.

Chapman, C. A., L. J. Chapman, T. T. Struhsaker, A. E. Zanne, C. J. Clark, C. J. Clark, and J. R.

Poulsen. 2005. A long-term evaluation of fruiting phenology: Importance of climate change.

Journal of Tropical Ecology 21:31–45.

Chapman, C. A., K. Valenta, T. R. Bonnell, K. A. Brown, and L. J. Chapman. 2018. Solar

radiation and ENSO predict fruiting phenology patterns in a 15-year record from Kibale National

Park, Uganda. Biotropica 50:384–395.

Chaturvedi, R. K., A. Tripathi, A. S. Raghubanshi, and J. S. Singh. 2021. Functional traits

indicate a continuum of tree drought strategies across a soil water availability gradient in a

tropical dry forest. Forest Ecology and Management 482:118740.

Chen, Y.-Y., A. Satake, I.-F. Sun, Y. Kosugi, M. Tani, S. Numata, S. P. Hubbell, C. Fletcher, N. N.

Supardi, and S. J. Wright. 2018. Species-specific flowering cues among general flowering

Shorea species at the Pasoh Research Forest, Malaysia. Journal of Ecology 106:586–598.

Cleland, E. E., I. Chuine, A. Menzel, H. A. Mooney, and M. D. Schwartz. 2007. Shifting plant

phenology in response to global change. Trends in Ecology & Evolution 22:357–365.

Dahinden, F., E. M. Fischer, and R. Knutti. 2017. Future local climate unlike currently observed

anywhere. Environmental Research Letters 12:084004.

Davis, C. C., G. M. Lyra, D. S. Park, R. Asprino, R. Maruyama, D. Torquato, B. I. Cook, and A.

M. Ellison. 2022. New directions in tropical phenology. Trends in Ecology & Evolution 37:683–

693.

Dessai, S., M. Hulme, R. Lempert, and R. Pielke. 2009. Do We Need Better Predictions to Adapt

to a Changing Climate? Eos, Transactions American Geophysical Union 90:111–112.

Devi, N. L., F. Q. Brearley, and S. K. Tripathi. 2023. Phenological diversity among sub-tropical

moist forest trees of north-eastern India. Journal of Tropical Ecology 39:e29.

Diez, J. M., I. Ibáñez, J. A. Silander, R. Primack, H. Higuchi, H. Kobori, A. Sen, and T. Y. James.

2014. Beyond seasonal climate: Statistical estimation of phenological responses to weather.

Ecological Applications 24:1793–1802.

Dunham, A. E., O. H. Razafindratsima, P. Rakotonirina, and P. C. Wright. 2018. Fruiting

phenology is linked to rainfall variability in a tropical rain forest. Biotropica 50:396–404.

Gadgil, S., and P. A. Francis. 2016. El Niño and the Indian rainfall in June. Current Science

110:1010–1022.

Getis, A., and J. K. Ord. 1992. The Analysis of Spatial Association by Use of Distance Statistics.

Geographical Analysis 24:189–206.

Grandini, M., E. Bagli, and G. Visani. 2020. Metrics for Multi-Class Classification: An

Overview.

Haas, E., F. A. La Sorte, Hanna M. McCaslin, M. C. Belotti, and K. G. Horton. 2022. The

correlation between <scp>eBird</scp> community science and weather surveillance radar-

based estimates of migration phenology. Global Ecology and Biogeography.

Harrison, R. D. 2000. Repercussions of El Niño: Drought causes extinction and the breakdown

of mutualism in Borneo. Proceedings of the Royal Society of London. Series B: Biological

Sciences 267:911–915.

Hossin, M., and M. N. Sulaiman. 2015. A Review on Evaluation Metrics for Data Classification

Evaluations. International Journal of Data Mining & Knowledge Management Process 5:01–11.

Hurlbert, A. H., and Z. Liang. 2012. Spatiotemporal variation in avian migration phenology:

Citizen science reveals effects of climate change. PLOS ONE 7:31662.

Kabano, P., A. Harris, and S. Lindley. 2020. Sensitivity of Canopy Phenology to Local Urban

Environmental Characteristics in a Tropical City. Ecosystems 24:1–15.

Kelly, J. F., K. G. Horton, P. M. Stepanian, K. M. de Beurs, T. D. Fagin, E. S. Bridge, and P. B.

Chilson. 2016. Novel measures of continental-scale avian migration phenology related to

proximate environmental cues. Ecosphere 7.

Kendall, D. E., T. L. S. M. Kendall, P. S. A. Stuart, and P. M. S. S. J. K. Ord. 1987. The advanced

theory of statistics: Volume 3: Design and analysis, and time-series. Oxford University Press,

Incorporated.

Lee, C. K. F., G. Song, H. C. Muller-Landau, S. Wu, S. J. Wright, K. C. Cushman, R. F. Araujo,

S. Bohlman, Y. Zhao, Z. Lin, Z. Sun, P. C. Y. Cheng, M. K.-P. Ng, and J. Wu. 2023. Cost-

effective and accurate monitoring of flowering across multiple tropical tree species over two

years with a time series of high-resolution drone imagery and deep learning. ISPRS Journal of

Photogrammetry and Remote Sensing 201:92–103.

Li, W., Q. Xin, X. Zhou, Z. Zhang, and Y. Ruan. 2021. Comparisons of numerical phenology

models and machine learning methods on predicting the spring onset of natural vegetation across

the Northern Hemisphere. Ecological Indicators 131:108126.

Lüdecke, D. 2023. sjPlot: Data visualization for statistics in social science. Manual.

Lüdecke, D., M. S. Ben-Shachar, I. Patil, P. Waggoner, and D. Makowski. 2021. performance:

An R package for assessment, comparison and testing of statistical models. Journal of Open

Source Software 6:3139.

Makhmale, S., P. Bhutada, L. Yadav, and B. K. Yadav. 2016. Impact of climate change on

phenology of Mango– The case study. Ecology Environment & Conservation.

Mann, H. B. 1945. Nonparametric Tests Against Trend. Econometrica 13:245.

McDonough MacKenzie, C., A. S. Gallinat, and L. Zipf. 2020. Low-cost observations and

experiments return a high value in plant phenology research. Applications in Plant Sciences 8.

Medway, L. 1972. Phenology of a tropical rain forest in Malaya. Biological Journal of The

Linnean Society 4:117–146.

Mendoza, I., C. A. Peres, and L. P. C. Morellato. 2017. Continental-scale patterns and climatic

drivers of fruiting phenology: A quantitative Neotropical review. Global and Planetary Change

148:227–241.

Mora, C., I. R. Caldwell, E. S. McBryde, J. M. Caldwell, M. R. Fisher, B. M. Genco, and S. W.

Running. 2015. Suitable Days for Plant Growth Disappear under Projected Climate Change:

Potential Human and Biotic Vulnerability. PLOS Biology 13:1.

Mora, C., A. G. Frazier, R. J. Longman, R. Dacks, M. M. Walton, E. J. Tong, J. J. Sanchez, L. R.

Kaiser, Y. O. Stender, J. M. Anderson, C. M. Ambrosino, I. Fernandez-Silva, L. M. Giuseffi, and

T. W. Giambelluca. 2013. The projected timing of climate departure from recent variability.

Nature 502:183–187.

Morellato, L. P. C., L. F. Alberti, and I. L. Hudson. 2010. Applications of Circular Statistics in

Plant Phenology: A Case Studies Approach. Pages 339–359 Phenological Research. Springer

Netherlands, Dordrecht.

Morellato, L. P. C., B. Alberton, S. T. Alvarado, B. D. Borges, E. Buisson, M. G. G. Camargo, L.

F. Cancian, D. W. Carstensen, D. F. E. Escobar, P. T. P. Leite, I. Mendoza, N. M. W. B. Rocha, N.

C. Soares, T. S. F. Silva, V. G. Staggemeier, A. S. Streher, B. da Cunha Vargas, and C. A. Peres.

2016. Linking plant phenology to conservation biology. Biological Conservation 195:60–72.

Neil, K., and J. Wu. 2006. Effects of urbanization on plant flowering phenology: A review. Urban

Ecosystems 9:243–257.

Parmesan, C. 2007. Influences of species, latitudes and methodologies on estimates of

phenological response to global warming. Global Change Biology 13:1860–1872.

Pau, S., D. K. Okamoto, O. Calderón, and S. J. Wright. 2018. Long-term increases in tropical

flowering activity across growth forms in response to rising CO2 and climate change. Global

Change Biology 24:2105–2116.

Pau, S., E. M. Wolkovich, B. I. Cook, C. J. Nytch, J. Regetz, J. K. Zimmerman, and S. J. Wright.

2013. Clouds and temperature drive dynamic changes in tropical flower production. Nature

Climate Change 3:838–842.

Piao, S., Q. Liu, A. Chen, I. A. Janssens, Y. Fu, J. Dai, L. Liu, X. Lian, M. Shen, and X. Zhu.

2019. Plant phenology and global climate change: Current progresses and challenges. Global

Change Biology 25:1922–1940.

Power, S. B., F. Delage, C. T. Y. Chung, G. Kociuba, Kevin Keay, and K. Keay. 2013. Robust

twenty-first-century projections of El Niño and related precipitation variability. Nature 502:541–

545.

R Core Team. 2022. R: A Language and Environment for Statistical Computing. R Foundation

for Statistical Computing, Vienna, Austria.

Ramaswami, G. 2022. SeasonWatch - Tree Phenology Dataset (India). [object Object].

Ramaswami, G., A. Datta, A. Reddy, and S. Quader. 2019. Tracking Phenology in the Tropics

and in India: The Impacts of Climate Change. Pages 45–59 The Implications of Climate Change

for Indian Biodiversity: An Overview.

Ramaswami, G., and S. Quader. 2018. The Case of the Confusing Kanikonna Trees. The Wire.

Ramaswami, G., S. Sidhu, and S. Quader. 2021. Using citizen science to build baseline data on

tropical tree phenology. CURRENT SCIENCE 121.

Revadekar, J. V., D. R. Kothawale, and K. Rupa Kumar. 2009. Role of El Niño/La Niña in

temperature extremes over India. International Journal of Climatology 29:2121–2129.

Sakai, S., and K. Kitajima. 2019. Tropical phenology: Recent advances and perspectives.

Ecological Research 34:50–54.

Satake, A., T. Kawagoe, Y. Saburi, Y. Chiba, Gen Sakurai, and H. Kudoh. 2013. Forecasting

flowering phenology under climate warming by modelling the regulatory dynamics of flowering-

time genes. Nature Communications 4:2303–2303.

Schwartz, M. D., J. L. Betancourt, and J. F. Weltzin. 2012. From Caprio's lilacs to the USA

National Phenology Network. Frontiers in Ecology and the Environment 10:324–327.

Sen, P. K. 1968. Estimates of the Regression Coefficient Based on Kendall's Tau. Journal of the

American statistical association.

Shepherd, T. G., and A. H. Sobel. 2020. Localness in Climate Change. Comparative Studies of

South Asia, Africa and the Middle East 40:7–16.

Singh, J. S., and V. K. Singh. 1992. Phenology of seasonally dry tropical forest. CURRENT

SCIENCE 63.

Singh, K. P., and C. P. Kushwaha. 2006. Diversity of Flowering and Fruiting Phenology of Trees

in a Tropical Deciduous Forest in India. Annals of Botany 97:265–276.

Singh, S. N., S. Sengupta, M. Kundu, S. Sahay, and D. Kumar. 2021. Phenological

characterization of different mango genotype under Sabour condition. The Pharma Innovation

Journal.

Sutton, P., D. Roberts, C. Elvidge, and K. Baugh. 2001. Census from Heaven: An estimate of the

global human population using night-time satellite imagery. International Journal of Remote

Sensing 22:3061–3076.

Vanderplank, S. E., and E. Ezcurra. 2016. Marine influence controls plant phenological dynamics

in Mediterranean Mexico. Journal of Plant Ecology 9:410–420.

Visser, M. E., and C. Both. 2005. Shifts in phenology due to global climate change: The need for

a yardstick. Proceedings of The Royal Society B: Biological Sciences 272:2561–2569.

Wang, X., and C. Wang. 2014. Different impacts of various El Niño events on the Indian Ocean

Dipole. Climate Dynamics 42:991–1005.

Willig, M. R., J. Rojas-Sandoval, and S. J. Presley. 2024. Phenological patterns in ecology:

Problems using circular statistics and solutions based on simulations. Methods in Ecology and

Evolution 15:868–885.

Wright, S. J., and O. Calderón. 2018. Solar irradiance as the proximate cue for flowering in a

tropical moist forest. Biotropica 50:374–383.

Wright, S. J., C. Carrasco, O. Calderón, and S. Paton. 1999. The El Niño Southern Oscillation,

Variable Fruit Production, and Famine in a Tropical Forest. Ecology 80:1632–1647.

Zhang, Y., and S. Fueglistaler. 2019. Mechanism for Increasing Tropical Rainfall Unevenness

With Global Warming. Geophysical Research Letters 46:14836–14843.

Zimmerman, J. K., S. J. Wright, O. Calderón, M. A. Pagan, and S. R. Paton. 2007. Flowering and

fruiting phenologies of seasonal and aseasonal neotropical forests: The role of annual changes in

irradiance. Journal of Tropical Ecology 23:231–251.

a

Circular mean of flowering status

b

Circular mean of flowering onset

a Distribution of tree-level mean flowering onset week

jackfruit mango tamarind

Number of trees

Mean flowering week

b

jackfruit mango tamarind

median = 6.91 median = 6.39 median = 15.94

Number of trees

Standard deviation (weeks) of tree-level flowering onset across years

a) onset of jackfruit flowering
b) onset of mango flowering
c) onset of tamarind flowering