Berberine dimers as chemical probes of *Pseudomonas aeruginosa* MexXY-OprM efflux function and inhibition

Logan G. Kavanaugh[1#], Andrew R. Mahoney[2#], Debayan Dey[1], William M. Wuest[2,3*] and Graeme L. Conn[1,3*]

[1]Department of Biochemistry, Emory University School of Medicine, Atlanta, GA

[2]Department of Chemistry, Emory University, Atlanta, GA

[3]Emory Antibiotic Resistance Center, Emory University, Atlanta, GA

[#]Co-first authors

*Correspondence to: Graeme L. Conn, gconn@emory.edu or William M. Wuest, wwuest@emory.edu

Keywords

Pseudomonas, antibiotic, efflux, Resistance-Nodulation-Division (RND), berberine, efflux pump inhibitor, MexXY-OprM.

Abbreviations

4,6-DOS, 4,6-deoxysteptamine; **CA-MHB**, cation-adjusted Mueller Hinton broth; **DPB**, Distal binding pocket; **EPI**, efflux pump inhibitor; **FIC**, fractional inhibitory concentration; **HTVS**, high-throughput virtual screening; **PBP**, proximal binding pocket; PEG, polyethylene glycol; **RND**, Resistance-Nodulation-Division. *Antibiotics*: **Ami**, amikacin; **Cef**, cefepime; **Cip**, ciprofloxacin; **Gen**, gentamicin; **Kan**, kanamycin; **Imi**, imipenem; **Tig**, tigecycline; **Tob**, tobramycin.

Abstract

The Resistance-Nodulation-Division (RND) efflux pump superfamily is pervasive among Gram-negative pathogens and contributes extensively to clinical antibiotic resistance. The opportunistic pathogen *Pseudomonas aeruginosa* contains 12 RND-type efflux systems, with four contributing to resistance including MexXY-OprM which is uniquely able to export aminoglycosides. At the site of initial substrate recognition, small molecule probes of the inner membrane transporter (e.g., MexY) have potential as important functional tools to understand substrate selectivity and a foundation for developing adjuvant efflux pump inhibitors (EPIs). Here, we optimized the scaffold of berberine, a known but weak MexY EPI, using an *in-silico* high-throughput screen to identify berberine dimers with enhanced synergistic action with aminoglycosides. Further, docking and molecular dynamics simulations of berberine dimers reveal unique contact residues and thus sensitivities of MexY from distinct *P. aeruginosa* strains. This work thereby reveals berberine dimers to be useful probes of MexY transporter function and potential leads for EPI development.

Introduction

Multi-drug resistant bacteria pose a global health threat with escalating infection cases resulting in increased morbidity and mortality, longer hospital stays, and increased financial burden[1,2]. In patients with cystic fibrosis, chronic lung infections caused by *Pseudomonas aeruginosa* are common and eradication is challenging[3-5]. In *P. aeruginosa* and other Gram-negative pathogens, expression of the chromosomally encoded Resistance-Nodulation-Division (RND) superfamily efflux pumps contributes extensively to multidrug resistance[6-11]. However, development of efflux pump inhibitors (EPIs) as therapeutics has proven challenging and is further impacted by the more broadly slowed progression of new antibiotic discovery and development[12,13].

P. aeruginosa encodes 12 hydrophobe/ amphiphile efflux-1 (HAE-1) family efflux pumps within the RND superfamily, of which four contribute to clinical antibiotic resistance: MexAB-OprM, MexCD-OprJ, MexEF-OprN, and MexXY-OprM[7-10]. These four pumps have overlapping, yet distinct antibiotic substrate profiles with MexXY-OprM having a unique ability to recognize and efflux aminoglycoside antibiotics[14-16]. For therapeutic regimes requiring aminoglycoside use, such as inhaled tobramycin for treatment of cystic fibrosis-associated *P. aeruginosa* infections, the MexXY-OprM efflux system plays a crucial role in resistance development and treatment failure[17-19]. A deeper understanding of the mechanisms governing antibiotic recognition by RND systems and the development of small molecule EPIs are urgently needed to negate efflux-mediated antibiotic resistance.

RND efflux pumps are tripartite protein complexes comprising a homotrimeric outer membrane channel (e.g., OprM), homohexameric periplasm adaptor protein (e.g., MexX), and a homotrimeric inner membrane transporter (e.g., MexY; **Fig. 1a**)[20-22]. The inner membrane transporter is a substrate/ H$^+$ antiporter and is the initial site for substrate recognition. Each protomer of the trimeric inner membrane

transporter contains 12 transmembrane helices (TM1-TM12) and a periplasmic domain with six subdomains that comprise the pore (PN1, PN2, PC1, PC2) and the MexX docking interface (DN, DC) (**Fig. 1b**)[23-26]. Two main predicted entry channels from the periplasmic space into the transporter are the cleft, between PC1 and PC2 of a single protomer, and the vestibule, between PN2 of one protomer and PC2 of a second protomer (**Fig. 1b**)[18,24]. Each protomer cycles through three conformational states that are required for substrate entry into the transporter and subsequent translocation to the channel of the periplasmic adapter: access, binding, and extrusion[24,25,27]. Potential efflux substrates first enter the proximal binding pocket (PBP) of the access state located within PC1 and PC2 and activate a conformational change at the glycine-rich "switch loop" (**Fig. 1b**)[28]. Substrates then move to the distal binding pocket (DBP) as the transporter cycles to the binding and then extrusion states for release of substrate to the open efflux tunnel. However, specific residues within the PBP and DBP required for substrate recognition and conformational state change have yet to be elucidated.

To provide insight into substrate recognition and/ or potential EPI binding, we aimed to identify potential molecular probes with distinct binding sites in the PBP /DBP region of MexY. Using the natural isoquinoline alkaloid berberine, a specific but weak MexY EPI[29,30], as the starting scaffold in virtual screening against a homology model of MexY, we identified four novel berberine analogs. Through iterative rounds of structure-guided docking, molecular dynamics simulations, organic synthesis, and microbiological assays, we identified berberine dimers that displayed increased synergy with aminoglycosides compared to natural berberine in *P. aeruginosa* PAO1, PA7, PA14, and pan-aminoglycoside resistant clinical isolates (**Supplemental Table S1**). Through alteration of the dimer linker properties, we also demonstrate the applicability of these new berberine derivatives to identify binding pocket residues and substrate conformational properties which may play a role in specific substrate recognition by MexY.

Results

Virtual screening predicts four diverse berberine scaffolds as MexY chemical probes.

In silico high-throughput virtual screening (HTVS) in the Schrödinger software was used to predict berberine analogs with enhanced binding scores and/ or distinct binding sites within the MexY PBP and DBP. Berberine was first used as a query on PubChem to generate a custom compound set comprising ~10,000 potential ligands with a Tanimoto coefficient threshold >0.8. This ligand set was next docked using HTVS Glide (Schrödinger) into a binding pocket comprising the PBP and DBP of a *P. aeruginosa* PAO1 MexY (MexY[PAO1]) homology model in its binding conformational state[31]. Ligands were ranked and the top scoring 1000 were subsequently re-docked with increased precision using Glide SP (Schrödinger). The top 1000 ligands were also clustered using a K-means algorithm based on chemical similarity to ensure selection of diverse scaffolds for further experimental investigation. This process produced four chemically

diverse ligands (**Scheme 1**, *Generation 1*): berberine conjugated with piperazine (Ber-Pip [**1**] and Ber-
Carb [**2**]) or propanamine moieties (Ber-Prop [**3**]), and a berberine dimer with a hexane linker (Ber-C6 [**4**]).

Our docking model predicted berberine binding to MexY^{PAO1} through multiple π-stacking
interactions, including F610 in the switch loop and W177 in the DBP, as previously seen[29] (**Fig. 2a**).
Furthermore, docking predicted three of the berberine analogs (Ber-Pip [1], Ber-Prop [3], Ber-C6 [4]) to
have comparable or increased binding scores compared to berberine (**Supplemental Table S2**). In the
DBP, Ber-Pip [**1**] and Ber-Carb [**2**] were predicted to have π-π stacking and π-cation interactions,
respectively, while the terminal amino group of Ber-Prop [**3**] was positioned to interact with D124
(**Supplemental Fig. S1a-c** and **Supplemental Table S3**). Ber-C6 [**4**] was predicted to bind in the DBP at
the PC1/DN interface through multiple interactions, including π-stacking with Y752, π-cation interaction
with K764, hydrogen bonding with the backbone of T176, and a salt bridge formed with E273
(**Supplemental Fig. S1d** and **Supplemental Table S3**).

Berberine dimer has increased synergy with aminoglycosides.
To test the validity of our computational modeling, a semisynthetic strategy was developed to obtain the
top four ligands Ber-Pip [**1**], Ber-Carb [**2**], Ber-Prop [**3**] and Ber-C6 [**4**] (**Scheme 1**–*Generation 1* and see
Supplementary Methods). Synthesis began with selective C9-demethylation of commercial berberine
chloride via vacuum pyrolysis to generate the natural product berberrubine chloride in high yield. The
resulting phenol was appended to a commercially available tertiary amine scaffold via a carbamate linkage
to give Ber-Carb [**2**]. Next, the electron-rich nature of the berberrubine ring system allowed for
derivatization of C13 by double condensation of berberrubine chloride and N-methylpiperazine onto
formaldehyde under acidic conditions, generating Ber-Pip [**1**]. Heating berberrubine chloride with 1,6-
diiodohexane for 72 hours successfully generated the berberine dimer Ber-C6 [**4**], as well as a small
amount of monohaloalkylated berberine. Because of the small amount of this isolated byproduct as well
as its low reactivity, the propanamine derivative required an alternate approach in which berberrubine
chloride was instead alkylated using 3-(Boc-amino)propyl bromide. Deprotection with hydrochloric acid
then successfully afforded the desired ligand Ber-Prop [**3**] as the hydrochloride salt.

Checkerboard synergy assays were performed for berberine and the four synthesized analogs in
P. aeruginosa PAO1 in the presence of two 4,6-deoxysteptamine (4,6-DOS) aminoglycosides, kanamycin
(Kan) and gentamicin (Gen) (**Fig. 1c**). Checkerboard assays confirmed berberine (4 µg/mL) is a weak
inhibitor of the MexXY-OprM efflux pump, reducing the minimum inhibitory concentration (MIC) by two-fold
and with a fractional inhibitory concentration (FIC) of 0.5, where FIC values < 0.5 are considered synergistic
(**Fig. 1c**, **Table 1**, and **Supplemental Table S4**). As MICs for berberine and analogs could not be
experimentally determined, FIC value calculations used estimated MICs obtained via nonlinear regression
of final culture OD$_{600}$ (see Methods). Ber-Prop [**3**] and Ber-C6 [**4**] showed increased synergy with Kan and
Gen, with Ber-C6 [**4**] reducing the MIC two- to six-fold (FIC = 0.28-0.51). However, growth inhibition was
observed in the presence of Ber-C6 [**4**] alone suggesting aminoglycoside-independent growth inhibition

(**Fig. 1c**, **Table 1**, and **Supplemental Table S4**). Piperazine ligands, Ber-Pip [**1**] and Ber-Carb [**2**] showed weaker or no synergy with the two aminoglycosides tested (**Fig. 1c**, **Table 1**, and **Supplemental Table S4**). As a result, we selected the berberine dimer, Ber-C6 [**4**], as our focus for further development (**Scheme 1**–*Generation 2*) due to its increased aminoglycoside synergism, higher predicted binding score, and more extensive and diverse predicted interactions within the DBP of MexY[PAO1].

Berberine alkane linker length defines MexXY-OprM inhibition efficacy and specificity.

Berberine dimers were designed with systematic variation in alkyl chain length connecting the two berberine groups to assess the role of linker length, flexibility, and overall size of the dimeric scaffold in predicted binding and MexXY-OprM inhibition. These *Generation 2* ligands with propane (Ber-C3 [**5**]), butane (Ber-C4 [**6**]), octane (Ber-C8 [**7**]), decane (Ber-C10 [**8**]), or dodecane (Ber-C12 [**9**]) linkers (**Scheme 1**) were docked in MexY[PAO1] binding pocket as before (**Fig. 2b-c**, **Supplemental Fig. S1e-g** and **Supplemental Table S2**).

An overall trend between longer linker length and increased predicted binding score was observed with the exception of the shortest alkane-linked dimer, Ber-C3 [**5**] (**Supplemental Table S2**).The predicted lowest energy docked pose for Ber-C4 [**6**] differs from Ber-C6 [**4**], with loss of hydrophobic interactions but gain of alternative stabilization via a salt bridge interaction with E175 in the DBP (**Supplemental Fig. S1e** and **Supplemental Table S3**). In contrast, the dimer with the shortest linker, Ber-C3 [**5**], is predicted to make a greater number of stabilizing interactions in the DBP through hydrogen bonding (to S46, T176 and S272), π-stacking/cation contacts (to Y127 and K291), and a salt bridge (with E273) (**Fig. 2b** and **Supplemental Table S3**). Ber-C8 [**7**] was predicted to be stabilized through hydrogen bonding interactions with S46 and K79 in the PBP, while Ber-C10 [**8**] and Ber-C12 [**9**] had predicted interactions in the DBP with overlapping (K291), and distinct residues (E175 and Y127, respectively; **Fig. 2d**, **Supplemental Fig. S1f,g** and **Supplemental Table S3**). These docking results suggest that increased linker flexibility may allow the two berberine groups to independently adopt favorable conformations resulting in optimal interactions within the binding pocket, as no direct interactions were predicted with any alkane linker. Conversely, the Ber-C3 [**5**] dimer can fortuitously adopt numerous stabilizing interactions within the DBP despite being more conformationally constrained.

Each of these new berberine dimer analogs was obtained via reaction of berberrubine chloride with commercially available alkyl dihalides (**Scheme 1**–*Generation 2* and see Supplementary Methods). Yields were moderate and increased consistently with alkyl chain length, suggesting that the dimerization reaction is slowed either by the steric repulsion of the bulky polycyclic rings or the electronic repulsion of the charged quaternary nitrogen atoms as they approach one another. This hypothesis is additionally supported by the observation that addition of a single berberrubine monomer to the alkyl halide was relatively rapid, leading to isolable monoalkylated product in less than one hour, but complete addition of a second berberrubine molecule took up to 72 hours.

Checkerboard synergy assays were performed using *P. aeruginosa* PAO1 with each dimer in

combination with Kan and Gen (**Fig. 3a** and **Supplemental Fig. S2a**). Shortening the alkane linker in Ber-

C3 [**5**] and Ber-C4 [**6**] reduced the MIC of Kan up to four-fold (FIC=0.30 and 0.38, respectively; **Fig. 3a**,

Table 1, **Supplemental Fig. S2a** and **Supplemental Table S4**). Additionally, Ber-C4 reduced the MIC of

Gen up to four-fold (FIC=0.38) (**Supplemental Fig. S2a** and **Supplemental Table S4**). Ber-C3 at 64 µg/mL

was able to reduce the MIC of Gen four-fold (**Table 1**), but at the highest concentration (128 µg/mL),

reduced the Gen MIC six-fold (FIC=0.30, **Fig. 3a**). Increasing alkane linker length resulted in increased

synergistic killing of PAO1 in combination with aminoglycosides, but with increased aminoglycoside-

independent growth inhibition. Ber-C8 [**7**] reduced the MIC of Kan and Gen four-fold (FIC=0.28) and six-

fold (FIC=0.38), respectively, while Ber-C10 [**8**] and Ber-C12 [**9**] reduced the MICs of both aminoglycosides

by six- to eight-fold (**Fig. 3a**, **Table 1**, **Supplemental Fig. S2a** and **Supplemental Table S4**).

Next, to assess each berberine dimer's effect on growth phase and the potential for aminoglycoside

specific interactions, we performed time-kill assays using *P. aeruginosa* PAO1 with two additional,

structurally distinct 4,6-DOS aminoglycoside substrates, amikacin (Ami) and tobramycin (Tob; **Fig. 3b**,

Supplemental Fig. S3 and **Supplemental S4a,c**). These assays were done in the presence of the

berberine dimers with the two shortest and two longest alkane linkers at 64 µg/mL as this concentration

was predicted by checkerboard assays to allow differences to be observed across the range of linker

lengths. As for Kan and Gen, berberine reduced the MIC two-fold for both Ami and Tob, indicating that its

weak effect is independent of aminoglycoside substrate (**Table 1**, **Fig. 3b**, and **Supplemental Fig. S4a,c**).

Consistent with its modestly tighter predicted binding, Ber-C3 [**5**] reduced the MIC for Ami four-fold (to 0.25

µg/mL) and Tob two-fold (to 0.25 µg/mL), while Ber-C4 [**6**] MIC reduction was equivalent to berberine

(**Table 1**, **Fig. 3b**,and **Supplemental Fig. S4a,c**). For the longer alkyl-linked dimers, Ber-C10 [**8**] reduced

the MIC six-fold for both Ami and Tob, whereas Ber-C12 [**9**] reduced the MIC of Ami six-fold but the Tob

MIC only four-fold (**Table 1**, **Fig. 3b**, and **Supplemental Fig. S4a,c**). Confirming the aminoglycoside-

independent growth inhibition observed in previous checkerboard assays, growth curves of *P. aeruginosa*

PAO1 in the presence of Ber-C10 [**8**] and Ber-C12 [**9**] showed a considerable extension of lag-phase

growth switching to log-phase at 12 hours (compared to four hours) and a slower log-phase growth rate

(**Fig. 3b**, and **Supplemental Fig. S4a,c**).

The observation of altered growth patterns in the time-kill assays with PAO1 next prompted further

investigation of Ber-C3 [**5**], Ber-C4 [**6**], Ber-C10 [**8**], and Ber-C12 [**9**] specificity for the MexXY-OprM efflux

pump using equivalent experiments in a *P. aeruginosa* Δ*mexXY* strain (**Fig.3b**, **Supplementary Table S1**,

and **Supplementary Figure S4b,d**). Loss of functional MexXY-OprM in the absence of berberine ligands,

resulted in a four-fold MIC reduction for Ami and two-fold reduction for Tob (**Table 1**). The MICs for Ami

and Tob in the presence of Ber-C3 [**5**] or Ber-C4 [**6**] do not exceed those observed in the Δ*mexXY*

background suggesting these analogs with shorter alkane linkers are specific probes for the MexXY efflux

system (**Table 1**). This conclusion was further validated in time-kill assays where addition of either

compound in the Δ*mexXY* background did not alter growth patterns (**Fig. 3b** and **Supplementary Fig.**

S4b,d). Of note, the Ber-C3 [**5**] compound reduced the MIC of both aminoglycosides to *mexXY* knockout
levels, suggesting it may serve as a useful starting scaffold for MexXY-OprM EPIs. Consistent with
previous checkerboard assays, Ber-C10 [**8**] and Ber-C12 [**9**] further reduced the MIC of Ami (6-fold) and
Tob (6- and 4-fold, respectively) to below those observed in the absence of MexXY-OprM (**Fig.3b, Table**
1, and **Supplementary Figure S4b,d**). Additionally, inactivation of MexXY-OprM did not influence the
observed lag-phase extension or slower log-phase growth rate, suggesting the longer alkane dimers are
sensitizing the cell to aminoglycosides in both MexXY-dependent and -independent manners.

We considered the possibility that the longer alkane-linked dimers might be disrupting membrane
integrity, similar to amphipathic quaternary ammonium ligands with long hydrophobic chains[32,33]. However,
using a sheep erythrocyte hemolysis assay, we found no evidence that berberine, Ber-C4 [**6**], or either
long alkane dimer, Ber-C10 [**8**] or Ber-C12 [**9**], is hemolytic at 128 µg/mL, a concentration at which MexXY-
independent activity is clearly observed for the latter two analogs (**Supplemental Fig. S5a**). Further, using
a vancomycin susceptibility assay as a probe for membrane permeability[34] in *P. aeruginosa* PAO1 and
PAO1Δ*mexXY* resulted in identical vancomycin MICs with and without Ber-C12 [**9**] at 64 µg/mL, again
suggesting that the off-target effect(s) of this analog are not related to membrane perturbation
(**Supplemental Fig. S5b,c**). The true additional target of the longer alkane dimers is currently unknown
and an area on-going investigation.

Berberine and Ber-C3 [5] compete for binding in the MexY^PAO1 DBP
Our structure-guided docking and checkerboard assays suggest that berberine and Ber-C3 [**5**] are specific
inhibitors of the MexXY-OprM efflux system and bind at distinct but overlapping sites within the DBP (**Fig.**
2a-b). If correct, berberine and Ber-C3 [**5**] should compete for binding in the DBP and would therefore not
show synergistic activity in combination. Therefore, to provide an initial experimental validation of the ligand
docking, this idea was tested in a three-way synergy assay using a fixed concentration of berberine (64
µg/mL) and a range of concentrations of Ber-C3 [**5**] (128-1 µg/mL) and Ami (4-0.125 µg/mL). Consistent
with competition for binding, a weak antagonistic effect (two-fold increase in Ami MIC) or the same MIC as
berberine alone was observed at most concentrations of Ber-C3 [**5**] (1-16 and 32-64 µg/mL, respectively;
Supplemental Table S5). Only at 128 µg/mL was Ber-C3 [**5**] able to fully outcompete berberine for the
overlapping binding pocket and resulting in an Ami MIC equivalent to analog the alone (0.25 µg/mL;
Supplemental Table S5).

Linker structure plays a role in MexXY-OprM inhibition efficacy and specificity.
To further define how linker length, flexibility and chemical structure impact berberine dimer activity and
their potential as probes of MexXY function, four additional dimers were designed with alternative linkers:
para-berberine substituents linked by one or two aryl groups, Ber-pAr [**10**] and Ber-Biph [**11**], respectively,
or by polyethylene glycol (PEG) linkers of different lengths, Ber-PEG5 [**12**] and Ber-PEG8 [**13**] (**Scheme**
1–*Generation 3*). For example, if dimer linker length is the primary determinant of efflux inhibition then

dimers of a similar size with different linker types should maintain similar activity (Ber-PEG5 [**10**] *vs.* Ber-

C4 [**6**], Ber-pAr [**12**] *vs.* Ber-C6 [**4**], Ber-PEG8 [**11**] *vs.* Ber-C8 [**7**], and Ber-Biph [**13**] *vs.* Ber-C10 [**8**].

However, should flexibility or physico-chemical properties be critical, then conformationally more rigid aryl

rings and/or more hydrophilic PEG moieties should show activity differences between the alkane, aryl, and

PEG linkers of comparable length.

Computational docking of Ber-pAr [**10**], Ber-Biph [**11**], Ber-PEG5 [**12**], and Ber-PEG8 [**13**] showed

increased predicted binding scores for MexY^{PAO1} compared to their corresponding length alkane-linked

dimers (**Supplemental Table S2**). Inspection of the lowest energy docking poses for the similarly sized

analogs revealed Ber-pAr [**10**], Ber-PEG5 [**12**], and Ber-PEG8 [**13**] have unique predicted interactions in

the binding pocket compared to their respective length-compared alkane-linked dimers. Ber-pAr [**10**] and

Ber-PEG5 [**12**] were both predicted to be stabilized by interactions spanning the PBP (via hydrogen

bonding to S671) and DBP (via π-cation interaction with Y127) (compare **Fig. 2d** and **Supplementary**

Fig.S1d, and **Supplementary Fig.S1** panels **e** and **i**, respectively; and see **Supplemental Table S3**).

Additionally, Ber-pAr [**10**] formed an additional linker-mediated π-cation interaction with K173 (**Fig. 2d** and

Supplementary Table S3). The longer PEG linker dimer, Ber-PEG8 [**13**], was predicted to be stabilized

only in the DBP through a π-cation interaction (with Y127) and salt bridge (with E175) (compare

Supplemental Fig. S1 panels **f** and **j**; and see **Supplemental Table S3**). In contrast, Ber-Biph [**11**] shared

predicted interactions (via salt bridge with E175 and hydrogen bonding to K291) with its corresponding

length alkane-linked dimer, Ber-C10 [**8**], while gaining a linker-mediated π-cation interaction with K173 and

an additional salt bridge with E273 (compare **Supplemental Fig. S1** panels **g** and **h**; and see

Supplemental Table S3). To complement these computational predictions with experimental assays of

dimer activity, the four aryl- or PEG-linked dimers were generated by chemical synthesis as before

(**Scheme 1**–*Generation* 3 and see Supplementary Methods). Synthesis of these *Generation 3* PEG-ylated

analogs and aryl analogs proceeded similarly to the alkyl linkers over 72 hours despite the more activated

benzylic electrophiles in the latter.

Checkerboard synergy assays were performed as before using *P. aeruginosa* PAO1 in combination

with Kan and Gen (**Fig. 3a** and **Supplemental Fig. S2b**). Notably, increased predicted binding scores did

not correlate with increased aminoglycoside synergy as previously observed for *Generation 1* and

Generation 2 analogs (**Table 1** and **Supplemental Table S2**). The Ber-pAr [**10**] dimer showed synergy

with only Gen, reducing the MIC up to four-fold (FIC = 0.33-0.52), and predicted FICs for both tested

aminoglycosides were greater than the comparable Ber-C6 [**4**] dimer (**Fig. 3a** and **Supplemental Table**

S4). Ber-Biph [**11**], Ber-PEG5 [**12**], and Ber-PEG8 [**13**] showed similar or decreased synergy with Kan and

Gen compared to berberine, resulting in FIC values substantially higher than each corresponding length

alkane-linked dimer (**Supplemental Fig. S2a,b** and **Supplemental Table S4**).

Time-kill assays in *P. aeruginosa* PAO1 showed only two-fold reduction in MIC for both Ami and

Tob for Ber-pAr [**10**], Ber-PEG5 [**12**], and Ber-PEG8 [**13**] which is lower than the corresponding alkyl-linked

dimer in each case and comparable to berberine (**Fig. 3b**, **Table 1**, and **Supplemental Fig. S4a,c**). Ber-

pAr [**10**] alone resulted a reduction in log-phase growth rate but neither Ber-PEG5 [**12**] nor Ber-PEG8 [**13**] affected overall growth rate (**Fig. 3b** and **Supplemental Fig. S4a,c**, respectively). To assess the specificity for the MexXY-OprM efflux pump, time-kill assays were performed in the isogenic Δ*mexXY* strain. While no difference in growth rate was again observed in the presence of the PEG-ylated dimers, the reduced log-phase growth rate in the presence of Ber-pAr [**10**] observed in PAO1 was also observed in the Δ*mexXY* background (**Supplemental Fig. S4b,d** and **Fig. 3b**, respectively), suggesting this compound may exhibit off-target effects. However, as for the longer alkane-linked dimers, we found no evidence for an effect on membrane integrity (**Supplementary Fig. S5a**).

Overall, computational docking predicted increased binding affinities for *Generation 3* compounds compared to their similarly sized alkane dimer analogs, but this did not result in the anticipated increased synergy with aminoglycosides. We therefore propose the alkane linker flexibility may play an important role in ligand uptake into MexY and subsequent entry into the PBP/ DBP.

Molecular dynamics (MD) simulations reveal important berberine probe characteristics.

From the studies thus far, three ligands were selected as lead probes – Ber-C3 [**5**], Ber-pAr [**10**], and Ber-C12 [**9**] – reflecting the highest activity identified but with distinct linker flexibility, predicted binding pocket, and/ or specificity for MexXY-OprM. First, to more precisely define the nature and stability of predicted interactions made by berberine and these dimeric analogs with the MexY[PAO1] DBP/ PBP, we used a 50 ns MD simulation in the Schrödinger Desmond module followed by MM/GBSA calculations on selected structures along the trajectory. Interestingly, berberine and Ber-C3 [**5**] showed greater overall conformational flexibility over the simulation, compared to Ber-pAr [**10**] and Ber-C12 [**9**] (**Fig. 4a**). However, while similar total numbers of interactions were identified for each ligand pre- and post-simulation, only Ber-C3 [**5**] and Ber-pAr [**10**] maintained specific interactions over their trajectories (**Fig. 4a** and **Supplemental Table S6**).

MM/GBSA calculations were performed on representative structures in each 10 ns window (100 frames) over the 50 ns simulation to obtain the relative free energy of each ligand in its binding pocket and thus reveal potentially important characteristics of a MexY inhibitor (**Fig. 4b-d** and **Supplemental Fig. S6a-c**). As expected, the overall average binding scores for Ber-C3 [**5**] (-116.0 kcal/mol), Ber-pAr [**9**] (-110.8 kcal/mol), and Ber-C12 [**10**] (-128.9 kcal/mol) were significantly greater than berberine (-65.23 kcal/mol; **Fig. 4b**) where the total binding score is calculated from individual electrostatic, hydrogen bond, π-packing (π-mediated interactions), lipophilic, Van der Waals and solvation energetic scores.

The dimeric berberine ligands had significantly higher electrostatic interaction contributions than berberine, but there were no significant differences in hydrogen bonding contributions (**Supplemental Fig. S6a-b**). Ber-C12 [**9**] was predicted to make significantly more lipophilic interactions in the binding pocket than Ber-C3 [**5**], Ber-pAr [**10**], and berberine, which contributed to its high predicted binding affinity (**Supplemental Fig. S6c**). Reflecting its increased interactions with hydrophobic residues within the DBP during the MD simulation, Ber-C3 [**5**] has a significantly more favorable π-stacking mediated interactions,

potentially due to enhanced flexibility seen within the simulation allowing it to position optimally to interact

with the binding pocket (**Fig. 4a,c**). This was further validated in the ligand strain energy calculation (the

difference between the ligand's free energy when bound compared to free in solution), where Ber-C3 [**5**]

showed the lowest ligand strain energy (**Fig. 4b,d**). MD simulations of berberine and lead dimer probes

thus highlight characteristics, such as increased binding affinity, ability for multiple hydrophobic

interactions, while maintaining flexibility and overall low ligand strain, as important for the development of

MexXY-OprM specific efflux pump inhibitors.

Top berberine alkane dimers are active against P. aeruginosa PA7 and PA14.

To further investigate the efficacy and utility of our top lead probes, Ber-C3 [**5**], Ber-pAr [**10**], and Ber-C12

[**9**], we evaluated synergy with Gen in two additional *P. aeruginosa* strains, PA7 and PA14 (**Supplemental**

Table S1 and **Table 2**). Alignment of the MexY protein sequence from *P. aeruginosa* PA7 (GenBank

ABR84278; MexY^{PA7}) and PA14 (GenBank ABJ11205; MexY^{PA14}) with MexY^{PAO1} (GenBank OP868818)

revealed 96.56% and 99.04% amino acid identity, respectively. This high conservation suggests that

berberine and all analogs should maintain comparable predicted binding to the three MexY proteins. This

was confirmed using SP Glide which produced binding scores that were broadly comparable to MexY^{PAO1}

for both MexY^{PA7} and MexY^{PA14}, with only a few larger differences (> 1 kcal/mol) and generally moderately

tighter and weaker binding for MexY^{PA7} and MexY^{PA14}, respectively (**Supplemental Table S2**). However,

mapping the unique residues between the three *P. aeruginosa* strains on our MexY^{PAO1} homology model

reveals that while most differences reside in the docking and transmembrane domains, four are located

within the PBP/DBP: A589G (MexY^{PAO1}), D149G (MexY^{PA7}), S144A (MexY^{PA7}) and E175Q (MexY^{PA14}) (**Fig.**

5a). Whether these differences result in differential activity of the berberine analogs was therefore explored

with berberine and the three top selected dimers, Ber-C3 [**5**], Ber-pAr [**10**], and Ber-C12 [**9**].

Time-kill assays were performed for berberine, Ber-C3 [**5**], Ber-C12 [**9**], and Ber-pAr [**10**], at 64

µg/mL in combination with Gen against *P. aeruginosa* PA7 and PA14 (**Fig. 5b**). Berberine reduced the

MIC for Gen in PA7 two to four-fold and up to two-fold in PA14 (**Table 2** and **Fig. 5b**). Alkane-linked dimers,

Ber-C3 [**5**] and Ber-C12 [**9**], exhibited similar activity in each strain, reducing the Gen MIC six-fold in PA7

and two-fold in PA14 (**Table 2** and **Fig. 5b**). These results contrast to PAO1 where Ber-C3 [**5**] and Ber-

C12 [**9**] differentially reduced the MIC of Gen four-fold and eight-fold, respectively. For the aryl-linker dimer,

Ber-pAr [**10**] reduced the MIC of Gen four-fold in PA7, comparable to PAO1, but reduced the MIC up to 2-

fold in PA14. (**Table 2** and **Fig. 5b**). Overall, changes in Gen MICs in PAO1 were comparable to those

observed in PA7, but all ligands were less effective in PA14 (**Table 2**).

We next used computational docking to investigate whether any of the binding pocket residue

changes identified in the MexY sequence alignment (**Fig. 5a**) might be responsible for the observed

differences in synergism in PAO1/ PA7 compared to PA14. Docking of Ber-C3 [**5**], Ber-pAr [**10**], and Ber-

C12 [**9**] in homology models of MexY^{PAO1}, MexY^{PA7}, and MexY^{PA14} elucidated important differences in

predicted binding poses of the berberine dimers, suggesting that inherent differences between each MexY

binding pocket may indeed exist (**Supplemental Fig. S7** and **S8**). In particular, a predicted interaction with

MexY from three strains was observed at residue 175, i.e. MexY^{PAO1}/ MexY^{PA7} E175 and MexY^{PA14} Q175

(**Fig. 5c** and **Supplemental Fig. S7b-d**). In MexY^{PAO1}/ MexY^{PA7}, E175 extends into the binding pocket,

whereas Q175 of MexY^{PA14} adopts a distinct conformation resulting in potential clash with the berberine

analogs (**Fig. 5c**). This observation suggests that the identity and thus positioning of the residue 175 side

chain may influence the preferred ligand conformation and subsequent binding pocket interactions. For

example, Ber-C3 [**5**] adopts an 'extended' conformation in MexY^{PAO1}/ MexY^{PA7} but a 'compact'

conformation in MexY^{PA14}, resulting in loss of important π-cation/ π- π stacking interactions (with Y127 and

K291) and salt bridges (to E273 in MexY^{PAO1} and E175 in MexY^{PA7}). These differences in interactions

resulting from the change at residue 175 can thus explain the decreased fold-change in Gen MIC observed

for PA14 (**Fig. 5c**, **Table 2**, and **Supplemental Fig. S7b**-**S8**). Thus, these results exemplify how berberine

dimers can be used as functional probes of the MexY binding pocket to elucidate important residues

involved in ligand recognition.

Berberine dimers are active against pan-aminoglycoside resistant clinical isolates and show

preferential synergy with aminoglycoside substrates of MexXY-OprM

Having defined activity in *P. aeruginosa* PAO1, PA7 and PA14, we extended these studies to further

explore the potential utility of the top three berberine dimer analogs by assessing efflux inhibition activity

in two pan-aminoglycoside resistant *P. aeruginosa* clinical isolates, K2156 and K2161 [35] (**Supplemental**

Table S1). Alignment of the MexY protein sequence from *P. aeruginosa* K2156 (GenBank OP868819;

MexY^{K2156}) and K2161 (GenBank OP868820; MexY^{K2161}) with MexY^{PAO1} (GenBank OP868818) revealed

99.52% and 99.62% amino acid identity, respectively. Unique residues from both clinical isolates were

primarily located in the transmembrane region of the MexY^{PAO1} homology model, with no observed changes

in the PBP/ DBP binding pocket (**Supplemental Fig. S9**). However, both MexY^{K2156} and MexY^{K2161} have

Q840E (cleft) and T543A (transmembrane) amino acid substitutions, with additional unique V980I

(transmembrane) changes in MexY^{K2156} and A33V (vestibule) and D428N (central cavity) in MexY^{K2161}

(**Supplemental Fig. S9**).

MIC assays were performed for clinical isolates K2156 and K2161 in the presence of 4,6-DOS

aminoglycosides (Kan, Ami, Gen, and Tob) and 64 µg/mL of berberine or berberine dimer (**Table 3**). To

compare berberine ligand synergy with aminoglycosides, MICs for the isogenic Δ*mexXY* knockout for each

clinical isolate were also determined (**Table 3** and **Table S1**). Berberine, Ber-C3 [**5**], and Ber-pAr [**10**]

exhibited increased synergy against K2156, with a six- to eight-fold reduction in all MICs, where eight-fold

reduction represented complete MexXY-OprM inhibition. Ber-C12 [**9**] reduced the MIC of Kan below

Δ*mexXY* knockout levels (ten-fold reduction; ; **Table 3**). Additionally, berberine and the tested berberine

dimers all showed increased synergy of four to six-fold MIC reduction with Tob, which were below the two-

fold reduction observed in the Δ*mexXY* background (**Table 3**). In contrast, a low to moderate change (zero

to four-fold) reduction in Kan, Ami and Gen MICs was observed for K2161 in the presence of berberine

and berberine dimers, of which Ber-C3 [**5**] and Ber-C12 [**9**] showed increased synergy compared to berberine and Ber-pAr [**10**]. Interestingly, Ber-C3 [**5**] and Ber-C12 [**9**] inhibited Tob to Δ*mexXY* knockout levels (four-fold reduction), but none of the berberine compounds inhibited below observed knockout levels, suggesting off-target effects may also be strain dependent (**Table 3**). The variable synergies exhibited with pan-aminoglycoside clinical isolates in the presence of berberine, Ber-C3 [5], Ber-C12 [9], and Ber-pAr [10] in combination with homology modeling suggests these chemical probes could potentially be used to further understand efflux substrate interactions at entry channels (e.g., cleft, vestibule, central cavity) into MexY.

Finally, to assess the ability of these dimeric berberine probes to provide insight into binding preferences of other, non-aminoglycoside substrates and their routes through the MexY transporter, we determined MICs for a representative panel of additional antibiotics: tigecycline (Tig), cefepime (Cef), ciprofloxacin (Cip); and the non-MexY-substrate imipenem (Imi) (**Table 3**). Ber-C3 [**5**] and Ber-C12 [**9**] reduced the MIC of Tig two- to four-fold in PAO1 and K2156, whereas Ber-C12 [**9**] produced only a two-fold reduction in K2161 (**Table 3**). Additionally, Ber-C3 [**5**] reduced the MIC two-fold for the substrate Cef in PAO1. All other interactions between non-aminoglycoside substrates (Cef or Cip), *P. aeruginosa* strains (PAO1, K2156, or K2161), and ligands (berberine, Ber-C3 [**5**], Ber-C12 [**9**], or Ber-pAr [**10**]) resulted in either no difference in observed MIC or an increase in resistance (**Table 3**). Such increase in resistance is most likely due to increased efflux of these substrates by another RND-efflux system in *P.* aeruginosa. Additionally, a two-fold decrease in the MIC of non-substrate Imi was observed in the presence of Ber-C12 [**9**] (PAO1 and K2161) and Ber-pAr (PAO1) (**Table 3**). These data suggest that the berberine alkyl dimers can be used as aminoglycoside-specific probes for MexY efflux function across different *P. aeruginosa* strains. Additionally, our berberine dimers may serve as useful tool for understanding crosstalk between multiple RND systems in *Pseudomonas*.

Discussion

The development of efflux pump inhibitors (EPIs) remains challenging due to limitations in our understanding of substrate recognition and efflux mechanisms, as well as the resource-intensive nature of screening chemical libraries *in vitro* for potential hits[36]. Although no EPIs have progressed to clinical approval, those discovered such as phenylalanine arginyl β-napthylamide (PAβN) and carbonyl cyanide-m-chlorophenylhydrazone (CCCP) have provided valuable tools to study efflux function [37-41]. However, these general inhibitors suffer significant limitations such as being substrates for multiple RND systems within the same organism or having off-target effects, both of which limit their use as tools to study specific RND efflux transporter function[41]. Therefore, we aimed to identify a pathway to new chemical probes, specific for the MexXY-OprM efflux system, to elucidate efflux function and inhibition.

Here, based on its known MexXY-OprM EPI activity, we used the naturally occurring and well-studied alkaloid berberine as a starting scaffold for a structure-guided computational screen[29,30,42-47]. HTVS

identified four berberine analogs (*Generation 1*) as potential leads with diverse predicted binding preferences in the MexY PBP/ DPB. Among the *Generation 1* ligands, addition of piperazine moieties to monomeric berberine did not enhance inhibition as previously shown for other pyridylpiperazine-based allosteric inhibitors of RND efflux systems[48]. However, our studies identified an alkane-linked berberine dimer, Ber-C6 [**4**], as an alternative promising lead for MexY. Berberine dimers have been shown to bind preferentially to G-quadruplex DNA but, to our knowledge, have yet to be characterized as protein inhibitors[49,50].

Using novel berberine dimers of variable linker length (*Generation 2*) and chemical nature (*Generation 3*), we identified linker characteristics that affect inhibition of the MexXY-OprM efflux system. Generally, alkyl linker length correlated with increased binding scores, and these predictions were corroborated by biological data showing improved synergy with aminoglycosides. However, the exception to this trend was observed for the dimer with the shortest linker, Ber-C3 [**5**]), which had higher predicted binding scores, primarily due increased hydrophobic interactions. Again, however, biological assays corroborated these computational predictions, revealing that this ligand had increased synergy with aminoglycosides and reduced the MIC to Δ*mexXY* knockout levels. Importantly, Ber-C3 [**5**] is non-toxic to pseudomonal cells and appears specific for the MexXY-OprM system, unlike previously identified berberine analogs[46,47] Of note, longer alkyl dimers did exhibit clear aminoglycoside-independent growth inhibition, but contrary to expectation our data suggest that this phenomenon is not attributable to membrane disruption. Berberine was identified to negatively impact DNA topoisomerase I/II and DNA, causing cell cycle arrest and DNA damage[51-55]. In the absence of membrane effects, we therefore speculate that the longer alkyl-linked berberine dimers may act similarly and studies are currently ongoing to define the basis of these off-target effects.

Initial docking studies of berberine in the MexY[PAO1] binding pocket identified interactions with the switch loop as has previously been observed[29]. Our docking of additional berberine ligands was also validated through a three-way synergy assay (berberine, Ber-C3 [5], and amikacin) in PAO1 which supported the predicted overlapping binding sites for these ligands. Docking analyses with berberine alkyl-linked dimer analogs predicted multiple binding sites within the substrate binding region of MexY, including the PBP, switch loop, DBP, and at the DBP-porter interface, highlighting the potential utility of these berberine dimers as chemical probes for understanding the ligand translocation pathway and its kinetics. Although binding scores for dimers with alternative aryl or PEG linkers were higher compared to the comparably sized alkyl-linked dimers, these predictions were not borne out in the biological analyses for these ligands. As docking analysis does not consider the process of ligand entry into MexY, we propose that the dimers with more rigid and/ or hydrophobic linkers are less readily able to reach the MexY PBP/ DBD. Furthermore, molecular dynamics simulations suggest that possessing a short, but slightly flexible linker that allows for optimal π-stacking interactions in the binding pocket is an important characteristic for ligand binding to MexY.

Antimicrobial susceptibility testing (e.g., checkerboard assays, MICs, and time-kill assays) also
revealed preferential inhibition by berberine dimers for different drugs of the same class. For example, Ber-
pAr [**10**] showed increased synergy with gentamicin compared to other aminoglycosides tested. Further
development toward these specificities could thus allow berberine analogs to be used to probe specific
binding regions for substrates the MexXY-OprM efflux system. Additionally, reduction in MIC for non-
aminoglycoside substrates was possible, but inhibiting to the level of MexXY knockout in *P. aeruginosa*
strains was inconsistent, suggesting that these substrates can more efficiently outcompete berberine
analogs for binding or have alternative binding site(s), or that they are also effluxed by other RND
systems[56-59]. Further, we found that Ber-pAr [**10**] and Ber-C12 [**9**] had a strain-dependent effect on
enhanced susceptibility to non-efflux substrate imipenem, a phenomenon also previously observed in *P.*
aeruginosa PAO12 with berberine[60]. Importantly, our top berberine dimer analogs, Ber-C3 [**5**], Ber-C12 [**9**],
and Ber-pAr [**10**], showed promising activity against two pan-aminoglycoside resistant clinical *P.*
aeruginosa isolates with variable MexXY-OprM dependencies, K2156 and K2161, as well as *P. aeruginosa*
PA7 and PA14 strains. As proof-of-concept for the utility of the berberine dimers as chemical probes, we
were able to identify a *P. aeruginosa* strain specific residue (MexY^{PAO1}/MexY^{PA7} E175 *vs*. MexY^{PA14} Q175)
which may influence inherent properties of the MexY binding pocket and therefore subsequent efflux
inhibition. Through checkerboard synergy assays, we identified that Ber-C3 [**5**] has reduced synergy with
aminoglycosides in PA14 compared to PAO1 and PA7. Docking of Ber-C3 [**5**] into all three strains MexY
homology models identified ligand conformational changes due to the position of the 175 residue for
glutamic acid versus glutamine. These data highlight the functionality of berberine dimers as probes to
identify putative important residues involved in MexXY-OprM efflux inhibition in multiple *P. aeruginosa*
strains.

In summary, using chemical biology and small-molecule discovery, we generated a panel of novel
potential probes of the transporter component of the MexXY-OprM efflux pump in *P. aeruginosa*. As RND
efflux pumps are ubiquitous in Gram-negative pathogens and contribute extensively to multidrug
resistance, understanding the chemical and mechanical properties of efflux pumps and their respective
substrates is imperative. Further development of chemical probes, such as our lead berberine dimers, can
shed light on mechanisms that govern efflux pump substrate recognition in RND efflux pumps and help
drive new strategies for specific EPI development.

Methods

Homology modeling of MexY and MexY-Per
The amino acid sequences of the MexY from *P. aeruginosa* PAO1, PA7 and PA14 were retrieved from
The *Pseudomonas* Genome Database[61] (accession codes GCF_000006765.1, GCF_000017205.1, and
GCF_000404265.1, respectively) and a homology model of an asymmetric MexY homotrimer was
constructed using the cryo-EM structure of MexB as the template (PDB 6IOL) on the Swiss Model server

(https://swissmodel.expasy.org). Models built with the Swiss Model server were minimized in MacroModel (Schrödinger LLC 2022-3) using the OPLS4 force field and Polak-Ribier Conjugate Gradient (PRCG) minimization method with 2500 iterations. A MexY^{PAO1} model was also generated with the transmembrane regions removed and the remaining periplasmic components joined with Gly/Ser linker ("MexY-per", comprising residues: MexY$^{1-38}$-GSGSGGSGGS (linker)-MexY$^{558-863}$). The MexY-per model was constructed using the full *P. aeruginosa* PAO1 MexY homology model as the starting template in Biologics (Schrödinger LLC 2022-3) and with enhanced loop sampling for the linker region.

High-throughput virtual screening (HTVS) and molecular docking

The PyMol Caver plugin was used to identify potential functional pockets in the three conformationally distinct protomers of the trimeric MexY^{PAO1} homology model and the predicted binding pocket in the pore domain of the binding conformational state protomer (chain B) was selected for HTVS and docking studies. The MexY^{PAO1} model from *P. aeruginosa* PAO1 was prepared for docking in Protein Preparation Wizard and energy minimized prior to grid creation with OPLS4 force field (Schrodinger LLC 2022-3). For HTVS, a ligand set (~10,000 berberine analogs) was created using berberine as the query in PubChem. To ensure chemical diversity within this ligand set, we applied a Tanimoto coefficient cutoff of 0.8 and resultant ligands were prepared in LigPrep with the OPLS4 force field (Schrödinger LLC 2022-3). A region comprising the combined PBP and DBP of MexY^{PAO1} was targeted using a docking grid centered between MexY residues F610 and F623 with a 30 Å^3 outer grid (where the ligands can dock) and 15 Å^3 inner grid (where the ligand centroid can be placed). HTVS Glide docking module (Schrödinger LLC 2022-3) was used for screening of the ligand set and the top 1000 scoring ligands were re-docked with SP Glide (Schrödinger LLC 2022-3). The top 100 scoring ligands from SP Glide were selected based on chemical similarity (Tanimoto coefficient) using K-means clustering algorithm in the Discovery Informatics module (Schrödinger LLC 2022-3).

For additional docking studies using SP Glide, models of MexY^{PA7} and MexY^{PA14} were prepared as for MexY^{PAO1}, above. Due to the flexibility of Generation 2 and Generation 3 berberine dimer linkers, conventional docking could not sample all potential ligand conformations. Therefore, we used ConfGen module with mixed torsional/ low mode sampling method with 100 steps/ rotatable bond and 1000 iterations (Schrödinger LLC 2022-3) to generate all potential conformers of these ligands. These conformers were subsequently used as the input for SP Glide docking (Schrödinger LLC 2022-3) in MexY^{PAO1}, MexY^{PA7} and MexY^{PA14}.

Berberine analog synthesis

Instrumentation, chemicals, and general notes–Nuclear magnetic resonance (NMR) spectra were recorded on either a Varian INOVA 600 (600/150 MHz) or Varian INOVA 500 (500/125 MHz) spectrometer. Chemical shifts are quoted in parts per million (ppm) relative to tetramethylsilane standard and with the indicated solvent as an internal reference. The following abbreviations are used to describe signal

multiplicities: s (singlet), d (doublet), t (triplet), q (quartet), m (multiplet), br (broad), dd (doublet of doublets), dt (doublet of triplets), etc. Accurate mass spectra were recorded on an Agilent 6520 Accurate-Mass Q-TOF LC/MS.

Non-aqueous reactions were performed under an atmosphere of argon, in flame-dried glassware, with HPLC-grade solvents dried by passage through activated alumina. Brine refers to a saturated aqueous solution of sodium chloride, sat. NaHCO$_3$ refers to a saturated aqueous solution of sodium bicarbonate, sat. NH$_4$Cl refers to a saturated aqueous solution of ammonium chloride, etc. "Column chromatography," unless otherwise indicated, refers to purification in a gradient of increasing ethyl acetate concentration in hexanes or methanol concentration in dichloromethane on a Biotage® flash chromatography purification system. All chemicals were used as received from Oakwood, TCI America, Sigma-Aldrich, Alfa Aesar, or CombiBlocks.

Experimental procedures and characterization data–Berberrubine chloride (2.4 eq.) was combined with either A. diiodide (1 eq.) or B. dibromide (1 eq.)/sodium iodide (2 eq.) in N,N-dimethylformamide (0.1 M) in a sealed reaction vessel and heated to 70°C. This solution was stirred under a static argon atmosphere until thin layer chromatography (TLC) analysis showed reaction completion (48-72 hours). Solvent was removed _in vacuo_ and product was purified via silica gel column chromatography. All synthesized analogs used in biological analysis were purified to >95% purity by high-performance liquid chromatography (HPLC). Additional details of all synthetic procedures and results of chemical analyses are provided in the Supplementary Information.

Bacterial strains and growth conditions
Strains used in this study (**Supplemental Table S1**) were previously reported _P._ aeruginosa PAO1 (K767), PAO1Δ_mexXY_ (K1525), and pan-aminoglycoside resistant clinical isolates K2156 and K2161[14,19,62]. _P. aeruginosa_ PA7 and PA14 were kindly provided by Dr. Joanna Goldberg (Department of Pediatrics, Emory University, Georgia, USA). Bacterial strains were maintained on Lysogeny Broth (LB)-Miller or LB-Miller agar (Merck) and grown aerobically at 37°C overnight. Strains were stored in LB-Miller broth supplemented with 16% glycerol at -80°C.

DNA methods
Chromosomal DNA was extracted from _P. aeruginosa_ PAO1 (K767), PAO1ΔmexXY (K1525), or clinical isolates K2156 and K2161 using Qiagen DNeasy Blood and Tissue Kit according to manufacturer protocol. The _mexY_ gene from each strain was subsequently amplified using five primers that encompass the 5' and 3' ends of the gene and three internal primers to encompass all 3.1 kb. Purified PCR products were sent for Sanger Sequencing (GeneWiz). Resultant FASTA sequences were aligned to _P. aeruginosa_ PAO1 (accession GCF_000006765.1) and deposited to NCBI GenBank under the accessions OP868818 (K767),

OP868819 (K2156), and OP868820 (K2161). Disruption of *mexXY* in *P. aeruginosa* PAO1Δ*mexXY*

(K1525) was validated by colony PCR.

Antibiotic susceptibility assays

Growth conditions–For all antimicrobial susceptibility assays, *P. aeruginosa* strains were grown in cation-

adjusted Mueller-Hinton broth (CA-MHB) prepared using premixed MHB (Sigma) supplemented with 20

mg/L calcium chloride (OmniPur) and 10 mg/L magnesium chloride (Sigma). Strains were cultured

aerobically, shaking vigorously, at 37°C for 18 hours.

Minimum inhibitory concentration (MIC)–All MICs were determined using serial two-fold microdilution

method in 96-well microtiter plates as done previously[63]. Antibiotics used in this study were: kanamycin

(Kan; OmniPur), amikacin (Ami; ChemImpex), gentamicin (Gen; Sigma), tobramycin (Tob; Alfa Aesar),

cefepime (Cef; ChemImpex), ciprofloxacin (Cip; Enzo), tigecycline (Tig; Astatech), imipenem (Imi; Combi-

Blocks), and vancomycin (Van; Sigma). MIC values were determined at 18 hours post-inoculation by

reading optical density at 600 nm (OD_{600}) on a Synergy Neo2 Multi-Mode Plate Reader (Agilent BioTek),

and inhibition of growth defined as $OD_{600}<0.5$ above background. All MIC assays were performed at least

twice starting from fresh streak plates on separate days and each assay was performed as two technical

replicates. Assays were performed for all *P. aeruginosa* strains to determine starting MICs in the absence

of berberine or berberine analogs. All MIC assays with berberine ligands (Ber, Ber-C3 [**5**], Ber-C12 [**9**], or

Ber-pAr [**10**]) were adjusted for a final concentration of 1% DMSO in each well.

Checkerboard–Checkerboard assays were performed with *P. aeruginosa* PAO1 (K767) using Kan or Gen,

and all berberine ligands. Two-fold serial dilutions for a final concentration of Kan [256-4 µg/mL] or Gen[4-

0.06 µg/mL] in CA-MHB were dispensed across the rows of a 96-well microtiter plate. Berberine and

berberine analogs were first resuspended in 100% DMSO and then diluted with water to yield stocks [1280-

10 µg/mL] in 20% DMSO. Finally, the berberine and berberine analogs were dispensed down the columns

of the 96-well microtiter plate to give final two-fold decreasing concentrations (128-1 µg/mL) and 1% DMSO

final concentration. *P. aeruginosa* PAO1 (K767) was grown to OD_{600} of 0.1 in CA-MHB and was diluted to

a final concentration of 10^5 cells. MIC values were determined as above at 18 hours post-inoculation by

reading OD_{600}, where inhibition of growth defined as $OD_{600}<0.5$ above background. All checkerboard

assays were performed at least twice starting from fresh streak plates on separate days. OD_{600} readings

each replicate between antibiotic and berberine analog interaction were averaged and normalized from 0-

1.

For analysis of antibiotic-berberine interactions, fractional inhibitory concentration (FIC) was

determined as previously described[64], where a FIC value < 0.5 indicates synergy between antibiotic and

ligand. For berberine and berberine analogs without an experimentally measurable MIC value (i.e., greater

than 128 µg/mL), the final OD_{600} values from cultures at 18 hours after exposure to a range of ligand

concentrations (128-0 µg/mL) were plotted and nonlinear regression of the log transformed OD_{600} values

was used to extrapolate a calculated MIC value.

Time-kill–Growth curves for *P. aeruginosa* PAO1 (K767) and PAO1ΔmexXY (K1525) were measured in

the absence or presence of amikacin or tobramycin at concentrations ranging from ¼ to 2× the MIC and

with select berberine ligands (64 µg/mL). Berberine ligands were prepared and added to 96-well microtiter

plate as described above for the checkerboard assays. Microtiter plates containing technical replicates for

each antibiotic-berberine pair were incubated for 18 hours with vigorous shaking using in Cytation5

Imaging Reader (Agilent BioTek). OD_{600} values were recorded every 15 minutes and select time points (0,

2, 4, 6, 8, 10, 12, 18, and 24 hours) were plotted. Growth curves were performed in duplicate at least twice

from freshly streaked plates on separate days.

Three-way synergy–A three-way synergy assay was performed using *P. aeruginosa* PAO1 (K767) in the

presence of berberine (64 µg/m), Ber-C3 [**5**] (128-1 µg/mL) and Ami (4-0.06 µg/mL). The 96-well microtiter

plate for Ber-C3 [**5**] and amikacin was set up as a checkerboard synergy assay (as described above).

Berberine was added to all wells to a final concentration of 64 µg/mL and the total concentration of

berberine and Ber-C3 [**5**] was adjusted for a final concentration of 1% DMSO. MIC values were determined

as above at 18 hours post-inoculation by reading OD_{600}. The assay was performed twice using two fresh

streak plates for starter cultures on two different days.

Hemolysis assay

Hemolysis was performed on mechanically defibrinated sheep blood (Hemostat Labs). One mL of

erythrocytes was centrifuged for 10 minutes at 3,800 rpm. Supernatant was discarded, cells were washed

in 1 mL of 1× phosphate-buffered saline (PBS), and pelleted. This washing step was repeated five times.

The final pellet was diluted 20-fold in 1× PBS. Berberine and berberine ligands were resuspended in 100%

DMSO and the final starting concentration in each well was 2.5%. Berberine ligands were two-fold serially

diluted in DMSO from 128-0.5 µg/mL in a 96-well microtiter plate. 100 µL of sheep blood suspension was

added to 100 µL of berberine ligand, 100 µL 1% Triton-X (positive control), or 100 µL 1× PBS (negative

control). Samples were incubated for one hour at 37°C with shaking at 200 rpm, and then pelleted. The

OD_{540} of the resulting supernatant was recorded using Synergy Neo2 Multi-Mode Plate Reader (Agilent

BioTek). The concentration inducing 20% red blood cell lysis was calculated for each ligand based on

positive and negative control absorbances. Hemolysis assays were performed in triplicate.

MD simulation and MM/GBSA calculation

619   We used the same strategy and rationale for using a reduced model of *P. aeruginosa* PAO1 MexY-per as

620   previously described for AcrB[65,66]. The MexY-per SP Glide docking model with chemically diverse

621   berberine probes bound to the PBP and DBP binding pocket was used as a starting structure for MD

simulations. Simulations were run in duplicate with each pair of runs initiated using a different initial velocity with OPLS4 force field, producing essentially identical results. Each system was first neutralized by adding sodium ions around the protein using the System Builder module. The neutralized protein was placed in TIP3P water, and random water molecules were substituted to obtain an ionic strength of 150 mM. Each solvated system was relaxed using a series of restrained minimization stages each of 1 ns duration: 1) all heavy atoms with force constant 50 kcal/molÅ^2, all protein atoms with 2) 10 kcal/molÅ^2 and 3) 5 kcal mol kcal/molÅ^2, and finally 4) with no constraints. Unrestrained MD simulations were then performed for 60 ns, comprising 10 ns for equilibration followed by a 50 ns production run in the isothermal-isobaric (NPT) ensemble using Nose-Hoover chain thermostat and Martyna-Tobias-Klein barostat with relaxation times of 1 and 2 ps, respectively. The equations of motion were integrated using multiple time steps for short-range (2 fs) and long-range (6 fs) interactions with a 10 Å cutoff applied for non-bonded interactions. MM/GBSA calculations were performed in the Prime module (Schrödinger LLC 2022-3) with the OPLS4 force field and allowing protein side chain flexibility within a 5 Å radius of the bound ligand.

Acknowledgements

We thank Dr. Keith Poole in the Department of Biomedical and Molecular Sciences, Queens University, Ontario, Canada for providing *P. aeruginosa* PAO1, clinical isolates, and respective *mexXY* knockout strains. We thank Dr. Joanna Goldberg in the Department of Pediatrics, Emory University, Atlanta, Georgia, USA, for providing *P. aeruginosa* PA7 and PA14 isolates. This work was supported by the National Institute of General Medical Sciences (R35 GM119426 to W.M.W), National Institute of General Medical Sciences NRSA Pre-Doctoral Fellowship (F31 GM143891 to L.G.K), Cystic Fibrosis Foundation Student Traineeship (Kavana21H0 to L.G.K), an ACS MEDI Pre-doctoral Fellowship (A.R.M.), and Emory University through an Accelerator Grant from the Biological Discovery through Chemical Innovation (BDCI) initiative (to G.L.C. and W.M.W.). L.G.K. also gratefully acknowledges support from Dr. John and Linda McGowan from the Atlanta Chapter ARCS Foundation.

Author Contributions

L.G.K. and **A.R.M.** designed and performed experiments, analyzed data, and wrote the original draft and edited the manuscript; **D.D.** performed computational analyses; **W.M.W.** and **G.L.C.** conceived the study, supervised the project, and edited the manuscript. All authors jointly discussed the results and implications at all stages of the work and edited the final manuscript.

Competing Interests

The authors declare no competing interests.

Data Availability

Datasets generated during this analysis are available on NCBI GenBank under accession number BankIt2645695 OP868818-20.

References

1. Centers for Disease C, Prevention. Antibiotic Resistance Threats in the United States, 2019. Centers for Disease Control and Prevention; 2019.

2. Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. Lancet. 2022;399(10325):629-55. Epub 20220119. doi: 10.1016/s0140-6736(21)02724-0. PubMed PMID: 35065702; PMCID: PMC8841637.

3. Jurado-Martín I, Sainz-Mejías M, McClean S. Pseudomonas aeruginosa: An Audacious Pathogen with an Adaptable Arsenal of Virulence Factors. Int J Mol Sci. 2021;22(6). Epub 20210318. doi: 10.3390/ijms22063128. PubMed PMID: 33803907; PMCID: PMC8003266.

4. King J, Murphy R, Davies JC. Pseudomonas aeruginosa in the Cystic Fibrosis Lung. Adv Exp Med Biol. 2022;1386:347-69. doi: 10.1007/978-3-031-08491-1_13. PubMed PMID: 36258079.

5. Reynolds D, Kollef M. The Epidemiology and Pathogenesis and Treatment of Pseudomonas aeruginosa Infections: An Update. Drugs. 2021;81(18):2117-31. Epub 20211107. doi: 10.1007/s40265-021-01635-6. PubMed PMID: 34743315; PMCID: PMC8572145.

6. Blair JM, Richmond GE, Piddock LJ. Multidrug efflux pumps in Gram-negative bacteria and their role in antibiotic resistance. Future Microbiol. 2014;9(10):1165-77. doi: 10.2217/fmb.14.66. PubMed PMID: 25405886.

7. Kishk RM, Abdalla MO, Hashish AA, Nemr NA, El Nahhas N, Alkahtani S, Abdel-Daim MM, Kishk SM. Efflux MexAB-Mediated Resistance in P. aeruginosa Isolated from Patients with Healthcare Associated Infections. Pathogens. 2020;9(6). Epub 20200615. doi: 10.3390/pathogens9060471. PubMed PMID: 32549303; PMCID: PMC7350317.

8. Gomis-Font MA, Pitart C, Del Barrio-Tofiño E, Zboromyrska Y, Cortes-Lara S, Mulet X, Marco F, Vila J, López-Causapé C, Oliver A. Emergence of Resistance to Novel Cephalosporin-β-Lactamase Inhibitor Combinations through the Modification of the Pseudomonas aeruginosa MexCD-OprJ Efflux Pump. Antimicrob Agents Chemother. 2021;65(8):e0008921. Epub 20210716. doi: 10.1128/aac.00089-21. PubMed PMID: 34060900; PMCID: PMC8284450.

9. Xu C, Liu H, Pan X, Ma Z, Wang D, Zhang X, Zhu G, Bai F, Cheng Z, Wu W, Jin Y. Mechanisms for Development of Ciprofloxacin Resistance in a Clinical Isolate of Pseudomonas aeruginosa. Front Microbiol. 2020;11:598291. Epub 20210108. doi: 10.3389/fmicb.2020.598291. PubMed PMID: 33488544; PMCID: PMC7819972.

10. Singh M, Yau YCW, Wang S, Waters V, Kumar A. MexXY efflux pump overexpression and aminoglycoside resistance in cystic fibrosis isolates of Pseudomonas aeruginosa from chronic infections. Can J Microbiol. 2017;63(12):929-38. Epub 20170918. doi: 10.1139/cjm-2017-0380. PubMed PMID: 28922614.

11. Davin-Regli A, Pages JM, Ferrand A. Clinical Status of Efflux Resistance Mechanisms in Gram-Negative Bacteria. Antibiotics (Basel). 2021;10(9). Epub 20210916. doi: 10.3390/antibiotics10091117. PubMed PMID: 34572699; PMCID: PMC8467137.

12. Ventola CL. The Antibiotic Resistance Crisis. Pharmacy and Therapeutics. 2015;40(4):277-83.

13. Bhardwaj AK, Mohanty P. Bacterial efflux pumps involved in multidrug resistance and their inhibitors: rejuvinating the antimicrobial chemotherapy. Recent Pat Antiinfect Drug Discov. 2012;7(1):73-89. doi: 10.2174/157489112799829710. PubMed PMID: 22353004.

14. Masuda N, Ohya S. Cross-resistance to meropenem, cephems, and quinolones in Pseudomonas aeruginosa. Antimicrob Agents Chemother. 1992;36(9):1847-51. doi: 10.1128/aac.36.9.1847. PubMed PMID: 1416876; PMCID: PMC192198.

15. Okamoto K, Gotoh N, Nishino T. Alterations of susceptibility of Pseudomonas aeruginosa by overproduction of multidrug efflux systems, MexAB-OprM, MexCD-OprJ, and MexXY/OprM to carbapenems: Substrate specificities of the efflux systems. Journal of Infection and Chemotherapy. 2002;8(4):371-3. doi: 10.1007/s10156-002-0193-7.

16. Morita Y, Kimura N, Mima T, Mizushima T, Tsuchiya T. Roles of MexXY- and MexAB-multidrug efflux pumps in intrinsic multidrug resistance of Pseudomonas aeruginosa PAO1. J Gen Appl Microbiol. 2001;47(1):27-32. Epub 2002/12/17. doi: 10.2323/jgam.47.27. PubMed PMID: 12483565.

17. Vogne C, Aires JR, Bailly C, Hocquet D, Plésiat P. Role of the Multidrug Efflux System MexXY in the Emergence of Moderate Resistance to Aminoglycosides among Pseudomonas aeruginosa Isolates from Patients with Cystic Fibrosis. Antimicrobial Agents and Chemotherapy. 2004;48(5):1676-80. doi: 10.1128/AAC.48.5.1676-1680.2004.

18. Lau CH, Hughes D, Poole K. MexY-promoted aminoglycoside resistance in Pseudomonas aeruginosa: involvement of a putative proximal binding pocket in aminoglycoside recognition. mBio. 2014;5(2):e01068. Epub 2014/04/24. doi: 10.1128/mBio.01068-14. PubMed PMID: 24757215; PMCID: PMC3994515.

19. Sobel ML, McKay GA, Poole K. Contribution of the MexXY Multidrug Transporter to Aminoglycoside Resistance in Pseudomonas aeruginosa Clinical Isolates. Antimicrobial Agents and Chemotherapy. 2003;47(10):3202-7. doi: 10.1128/AAC.47.10.3202-3207.2003.

20. Shi X, Chen M, Yu Z, Bell JM, Wang H, Forrester I, Villarreal H, Jakana J, Du D, Luisi BF, Ludtke SJ, Wang Z. In situ structure and assembly of the multidrug efflux pump AcrAB-TolC. Nat Commun.

2019;10(1):2635. Epub 20190614. doi: 10.1038/s41467-019-10512-6. PubMed PMID: 31201302; PMCID: PMC6570770.

21. Du D, Wang Z, James NR, Voss JE, Klimont E, Ohene-Agyei T, Venter H, Chiu W, Luisi BF. Structure of the AcrAB-TolC multidrug efflux pump. Nature. 2014;509(7501):512-5. doi: 10.1038/nature13205.

22. Tsutsumi K, Yonehara R, Ishizaka-Ikeda E, Miyazaki N, Maeda S, Iwasaki K, Nakagawa A, Yamashita E. Structures of the wild-type MexAB–OprM tripartite pump reveal its complex formation and drug efflux mechanism. Nature Communications. 2019;10(1):1520. doi: 10.1038/s41467-019-09463-9.

23. Nakashima R, Sakurai K, Yamasaki S, Hayashi K, Nagata C, Hoshino K, Onodera Y, Nishino K, Yamaguchi A. Structural basis for the inhibition of bacterial multidrug exporters. Nature. 2013;500(7460):102-6. doi: 10.1038/nature12300.

24. Murakami S, Nakashima R, Yamashita E, Matsumoto T, Yamaguchi A. Crystal structures of a multidrug transporter reveal a functionally rotating mechanism. Nature. 2006;443(7108):173-9. doi: 10.1038/nature05076.

25. Seeger MA, Schiefner A, Eicher T, Verrey F, Diederichs K, Pos KM. Structural asymmetry of AcrB trimer suggests a peristaltic pump mechanism. Science. 2006;313(5791):1295-8. doi: 10.1126/science.1131542. PubMed PMID: 16946072.

26. Sennhauser G, Amstutz P, Briand C, Storchenegger O, Grütter MG. Drug Export Pathway of Multidrug Exporter AcrB Revealed by DARPin Inhibitors. PLoS Biology. 2007;5(1). doi: 10.1371/journal.pbio.0050007.

27. Seeger MA, Diederichs K, Eicher T, Brandstätter L, Schiefner A, Verrey F, Pos KM. The AcrB efflux pump: conformational cycling and peristalsis lead to multidrug resistance. Curr Drug Targets. 2008;9(9):729-49. doi: 10.2174/138945008785747789. PubMed PMID: 18781920.

28. Müller RT, Travers T, Cha HJ, Phillips JL, Gnanakaran S, Pos KM. Switch Loop Flexibility Affects Substrate Transport of the AcrB Efflux Pump. J Mol Biol. 2017;429(24):3863-74. Epub 20171005. doi: 10.1016/j.jmb.2017.09.018. PubMed PMID: 28987732.

29. Laudadio E, Cedraro N, Mangiaterra G, Citterio B, Mobbili G, Minnelli C, Bizzaro D, Biavasco F, Galeazzi R. Natural Alkaloid Berberine Activity against Pseudomonas aeruginosa MexXY-Mediated Aminoglycoside Resistance: In Silico and in Vitro Studies. J Nat Prod. 2019;82(7):1935-44. Epub 20190705. doi: 10.1021/acs.jnatprod.9b00317. PubMed PMID: 31274312.

30. Morita Y, Nakashima K, Nishino K, Kotani K, Tomida J, Inoue M, Kawamura Y. Berberine Is a Novel Type Efflux Inhibitor Which Attenuates the MexXY-Mediated Aminoglycoside Resistance in

Pseudomonas aeruginosa. Front Microbiol. 2016;7:1223. Epub 2016/08/23. doi: 10.3389/fmicb.2016.01223. PubMed PMID: 27547203; PMCID: PMC4975076.

31. Dey D, Kavanaugh LG, Conn GL. Antibiotic Substrate Selectivity of MexY and MexB Efflux Systems Is Determined by a Goldilocks Affinity. Antimicrobial Agents and Chemotherapy. 2020;64(8):e00496-20, /aac/64/8/AAC.-20.atom. doi: 10.1128/AAC.00496-20.

32. Jennings MC, Ator LE, Paniak TJ, Minbiole KP, Wuest WM. Biofilm-eradicating properties of quaternary ammonium amphiphiles: simple mimics of antimicrobial peptides. Chembiochem. 2014;15(15):2211-5. Epub 20140821. doi: 10.1002/cbic.201402254. PubMed PMID: 25147134.

33. Alkhalifa S, Jennings MC, Granata D, Klein M, Wuest WM, Minbiole KPC, Carnevale V. Analysis of the Destabilization of Bacterial Membranes by Quaternary Ammonium Compounds: A Combined Experimental and Computational Study. Chembiochem. 2020;21(10):1510-6. Epub 20200218. doi: 10.1002/cbic.201900698. PubMed PMID: 31859426; PMCID: PMC7237276.

34. Muheim C, Götzke H, Eriksson AU, Lindberg S, Lauritsen I, Nørholm MHH, Daley DO. Increasing the permeability of Escherichia coli using MAC13243. Sci Rep. 2017;7(1):17629. Epub 20171215. doi: 10.1038/s41598-017-17772-6. PubMed PMID: 29247166; PMCID: PMC5732295.

35. Sobel ML, McKay GA, Poole K. Contribution of the MexXY multidrug transporter to aminoglycoside resistance in Pseudomonas aeruginosa clinical isolates. Antimicrob Agents Chemother. 2003;47(10):3202-7. Epub 2003/09/25. doi: 10.1128/aac.47.10.3202-3207.2003. PubMed PMID: 14506031; PMCID: PMC201159.

36. Lomovskaya O, Bostian KA. Practical applications and feasibility of efflux pump inhibitors in the clinic—A vision for applied use. Biochemical Pharmacology. 2006;71(7):910-8. doi: https://doi.org/10.1016/j.bcp.2005.12.008.

37. Mesaros N, Glupczynski Y, Avrain L, Caceres NE, Tulkens PM, Van Bambeke F. A combined phenotypic and genotypic method for the detection of Mex efflux pumps in Pseudomonas aeruginosa. J Antimicrob Chemother. 2007;59(3):378-86. Epub 2007/02/10. doi: 10.1093/jac/dkl504. PubMed PMID: 17289770.

38. Bhattacharyya T, Sharma A, Akhter J, Pathania R. The small molecule IITR08027 restores the antibacterial activity of fluoroquinolones against multidrug-resistant Acinetobacter baumannii by efflux inhibition. Int J Antimicrob Agents. 2017;50(2):219-26. Epub 2017/05/24. doi: 10.1016/j.ijantimicag.2017.03.005. PubMed PMID: 28533185.

39. Osei Sekyere J, Amoako DG. Carbonyl Cyanide m-Chlorophenylhydrazine (CCCP) Reverses Resistance to Colistin, but Not to Carbapenems and Tigecycline in Multidrug-Resistant Enterobacteriaceae. Front Microbiol. 2017;8:228. Epub 2017/03/07. doi: 10.3389/fmicb.2017.00228. PubMed PMID: 28261184; PMCID: PMC5306282.

40. Whalen KE, Poulson-Ellestad KL, Deering RW, Rowley DC, Mincer TJ. Enhancement of antibiotic
activity against multidrug-resistant bacteria by the efflux pump inhibitor 3,4-dibromopyrrole-2,5-dione
isolated from a Pseudoalteromonas sp. J Nat Prod. 2015;78(3):402-12. Epub 2015/02/04. doi:
10.1021/np500775e. PubMed PMID: 25646964.

41. Li XZ, Plésiat P, Nikaido H. The challenge of efflux-mediated antibiotic resistance in Gram-negative
bacteria. Clin Microbiol Rev. 2015;28(2):337-418. doi: 10.1128/cmr.00117-14. PubMed PMID:
25788514; PMCID: PMC4402952.

42. Imanshahidi M, Hosseinzadeh H. Pharmacological and therapeutic effects of Berberis vulgaris and its
active constituent, berberine. Phytother Res. 2008;22(8):999-1012. doi: 10.1002/ptr.2399. PubMed
PMID: 18618524.

43. Kulkarni SK, Dhir A. Berberine: a plant alkaloid with therapeutic potential for central nervous system
disorders. Phytother Res. 2010;24(3):317-24. doi: 10.1002/ptr.2968. PubMed PMID: 19998323.

44. Vuddanda PR, Chakraborty S, Singh S. Berberine: a potential phytochemical with multispectrum
therapeutic activities. Expert Opin Investig Drugs. 2010;19(10):1297-307. doi:
10.1517/13543784.2010.517745. PubMed PMID: 20836620.

45. Liang Y, Xu X, Yin M, Zhang Y, Huang L, Chen R, Ni J. Effects of berberine on blood glucose in
patients with type 2 diabetes mellitus: a systematic literature review and a meta-analysis. Endocr J.
2019;66(1):51-63. Epub 20181103. doi: 10.1507/endocrj.EJ18-0109. PubMed PMID: 30393248.

46. Giorgini G, Mangiaterra G, Cedraro N, Laudadio E, Sabbatini G, Cantarini M, Minnelli C, Mobbili G,
Frangipani E, Biavasco F, Galeazzi R. Berberine Derivatives as Pseudomonas aeruginosa MexXY-
OprM Inhibitors: Activity and In Silico Insights. Molecules. 2021;26(21). Epub 20211102. doi:
10.3390/molecules26216644. PubMed PMID: 34771051; PMCID: PMC8587913.

47. Kotani K, Matsumura M, Morita Y, Tomida J, Kutsuna R, Nishino K, Yasuike S, Kawamura Y. 13-(2-
Methylbenzyl) Berberine Is a More Potent Inhibitor of MexXY-Dependent Aminoglycoside Resistance
than Berberine. Antibiotics (Basel). 2019;8(4). Epub 2019/11/09. doi: 10.3390/antibiotics8040212.
PubMed PMID: 31698691; PMCID: PMC6963850.

48. Plé C, Tam HK, Vieira Da Cruz A, Compagne N, Jiménez-Castellanos JC, Müller RT, Pradel E,
Foong WE, Malloci G, Ballée A, Kirchner MA, Moshfegh P, Herledan A, Herrmann A, Deprez B,
Willand N, Vargiu AV, Pos KM, Flipo M, Hartkoorn RC. Pyridylpiperazine-based allosteric inhibitors of
RND-type multidrug efflux pumps. Nat Commun. 2022;13(1):115. Epub 20220110. doi:
10.1038/s41467-021-27726-2. PubMed PMID: 35013254; PMCID: PMC8749003.

49. Giri P, Kumar GS. Isoquinoline alkaloids and their binding with polyadenylic acid: potential basis of
therapeutic action. Mini Rev Med Chem. 2010;10(7):568-77. doi: 10.2174/138955710791384009.
PubMed PMID: 20500148.

50. Tera M, Hirokawa T, Okabe S, Sugahara K, Seimiya H, Shimamoto K. Design and synthesis of a berberine dimer: a fluorescent ligand with high affinity towards G-quadruplexes. Chemistry. 2015;21(41):14519-28. Epub 20150813. doi: 10.1002/chem.201501693. PubMed PMID: 26272465.

51. Krishnan P, Bastow KF. The 9-position in berberine analogs is an important determinant of DNA topoisomerase II inhibition. Anticancer Drug Des. 2000;15(4):255-64. PubMed PMID: 11200501.

52. Qin Y, Pang JY, Chen WH, Zhao ZZ, Liu L, Jiang ZH. Inhibition of DNA topoisomerase I by natural and synthetic mono- and dimeric protoberberine alkaloids. Chem Biodivers. 2007;4(3):481-7. doi: 10.1002/cbdv.200790040. PubMed PMID: 17372950.

53. Kim JB, Yu JH, Ko E, Lee KW, Song AK, Park SY, Shin I, Han W, Noh DY. The alkaloid Berberine inhibits the growth of Anoikis-resistant MCF-7 and MDA-MB-231 breast cancer cell lines by inducing cell cycle arrest. Phytomedicine. 2010;17(6):436-40. Epub 20091002. doi: 10.1016/j.phymed.2009.08.012. PubMed PMID: 19800775.

54. Li DD, Xu Y, Zhang DZ, Quan H, Mylonakis E, Hu DD, Li MB, Zhao LX, Zhu LH, Wang Y, Jiang YY. Fluconazole assists berberine to kill fluconazole-resistant Candida albicans. Antimicrob Agents Chemother. 2013;57(12):6016-27. Epub 20130923. doi: 10.1128/aac.00499-13. PubMed PMID: 24060867; PMCID: PMC3837902.

55. Domadia PN, Bhunia A, Sivaraman J, Swarup S, Dasgupta D. Berberine targets assembly of Escherichia coli cell division protein FtsZ. Biochemistry. 2008;47(10):3225-34. Epub 20080215. doi: 10.1021/bi7018546. PubMed PMID: 18275156.

56. Dean CR, Visalli MA, Projan SJ, Sum PE, Bradford PA. Efflux-mediated resistance to tigecycline (GAR-936) in Pseudomonas aeruginosa PAO1. Antimicrob Agents Chemother. 2003;47(3):972-8. doi: 10.1128/aac.47.3.972-978.2003. PubMed PMID: 12604529; PMCID: PMC149306.

57. Jeannot K, Elsen S, Köhler T, Attree I, van Delden C, Plésiat P. Resistance and virulence of Pseudomonas aeruginosa clinical strains overproducing the MexCD-OprJ efflux pump. Antimicrob Agents Chemother. 2008;52(7):2455-62. Epub 20080512. doi: 10.1128/aac.01107-07. PubMed PMID: 18474583; PMCID: PMC2443911.

58. Zhao L, Wang S, Li X, He X, Jian L. Development of in vitro resistance to fluoroquinolones in Pseudomonas aeruginosa. Antimicrob Resist Infect Control. 2020;9(1):124. Epub 20200805. doi: 10.1186/s13756-020-00793-8. PubMed PMID: 32758289; PMCID: PMC7405379.

59. Lee A, Mao W, Warren MS, Mistry A, Hoshino K, Okumura R, Ishida H, Lomovskaya O. Interplay between efflux pumps may provide either additive or multiplicative effects on drug resistance. J Bacteriol. 2000;182(11):3142-50. doi: 10.1128/jb.182.11.3142-3150.2000. PubMed PMID: 10809693; PMCID: PMC94500.

60. Su F, Wang J. Berberine inhibits the MexXY-OprM efflux pump to reverse imipenem resistance in a
clinical carbapenem-resistant Pseudomonas aeruginosa isolate in a planktonic state. Exp Ther Med.
2018;15(1):467-72. Epub 20171101. doi: 10.3892/etm.2017.5431. PubMed PMID: 29387199;
PMCID: PMC5769289.

61. Winsor GL, Griffiths EJ, Lo R, Dhillon BK, Shay JA, Brinkman FS. Enhanced annotations and
features for comparing thousands of Pseudomonas genomes in the Pseudomonas genome
database. Nucleic Acids Res. 2016;44(D1):D646-53. Epub 20151117. doi: 10.1093/nar/gkv1227.
PubMed PMID: 26578582; PMCID: PMC4702867.

62. De Kievit TR, Parkins MD, Gillis RJ, Srikumar R, Ceri H, Poole K, Iglewski BH, Storey DG. Multidrug
efflux pumps: expression patterns and contribution to antibiotic resistance in Pseudomonas
aeruginosa biofilms. Antimicrob Agents Chemother. 2001;45(6):1761-70. doi:
10.1128/aac.45.6.1761-1770.2001. PubMed PMID: 11353623; PMCID: PMC90543.

63. Wiegand I, Hilpert K, Hancock RE. Agar and broth dilution methods to determine the minimal
inhibitory concentration (MIC) of antimicrobial substances. Nat Protoc. 2008;3(2):163-75. doi:
10.1038/nprot.2007.521. PubMed PMID: 18274517.

64. Meletiadis J, Pournaras S, Roilides E, Walsh TJ. Defining fractional inhibitory concentration index
cutoffs for additive interactions based on self-drug additive combinations, Monte Carlo simulation
analysis, and in vitro-in vivo correlation data for antifungal drug combinations against Aspergillus
fumigatus. Antimicrob Agents Chemother. 2010;54(2):602-9. Epub 20091207. doi:
10.1128/aac.00999-09. PubMed PMID: 19995928; PMCID: PMC2812160.

65. Ruggerone P, Vargiu AV, Collu F, Fischer N, Kandt C. Molecular Dynamics Computer Simulations of
Multidrug RND Efflux Pumps. Comput Struct Biotechnol J. 2013;5:e201302008. Epub 20130303. doi:
10.5936/csbj.201302008. PubMed PMID: 24688701; PMCID: PMC3962194.

66. Vargiu AV, Nikaido H. Multidrug binding properties of the AcrB efflux pump characterized by
molecular dynamics simulations. Proc Natl Acad Sci U S A. 2012;109(50):20637-42. Epub 20121121.
doi: 10.1073/pnas.1218348109. PubMed PMID: 23175790; PMCID: PMC3528587.

Tables

Table 1. Aminoglycoside minimum inhibitory concentration (MIC) against *P. aeruginosa* PAO1 in the presence of berberine and berberine analogs (64 µg/mL).

Strain	Berberine analog	MIC; µg/mL[a] (fold change)[b]			
		Kan	Gen	Ami	Tob
PAO1	-	64	1	1	0.5
PAO1Δ*mexXY*	-	16 (4)	0.25 (4)	0.25 (4)	0.25 (2)
PAO1	Berberine	32 (2)	0.5 (2)	0.5 (2)	0.25 (2)
	Ber-Pip [**1**]	64 (-)	1 (-)	*nd*	*nd*
	Ber-Carb [**2**]	32 (2)	0.5 (2)	*nd*	*nd*
	Ber-Prop [**3**]	32 (2)	0.5 (2)	*nd*	*nd*
	Ber-C3 [**5**]	16 (4)	0.25 (4)	0.25 (4)	0.25 (2)
	Ber-C4 [**6**]	16 (4)	0.25 (4)	0.5 (2)	0.25 (2)
	Ber-C6 [**4**]	16 (4)	0.25 (4)	*nd*	*nd*
	Ber-C8 [**7**]	16 (4)	0.125 (6)	*nd*	*nd*
	Ber-C10 [**8**]	8 (6)	0.0625 (8)	0.0125 (6)	0.0625 (6)
	Ber-C12 [**9**]	8 (6)	0.125 (6)	0.125 (6)	0.125 (4)
	Ber-pAr [**10**]	32 (2)	0.25 (4)	0.5 (2)	0.25 (2)
	Ber-Biph [**11**]	32 (2)	0.5 (2)	*nd*	*nd*
	Ber-PEG5 [**12**]	32 (2)	0.5 (2)	0.5 (2)	0.25 (2)
	Ber-PEG8 [**13**]	32 (2)	0.5 (2)	0.5 (2)	0.25 (2)

[a]Determined from checkerboard and time-kill assays. Aminoglycosides: kanamycin (Kan), amikacin (Ami), gentamicin (Gen), and tobramycin (Tob); *nd*, not determined.

[b]fold change relative to wild-type PAO1. (-)no fold change.

Table 2. Gentamicin MIC against *P. aeruginosa* strains PA7 and PA14 in the absence and presence of berberine or select analogs.

Strain	Gentamicin MIC, µg/mL (fold change)[a,b,c]				
	-	Berberine	Ber-C3 [**5**]	Ber-C12 [**9**]	Ber-pAr [**10**]
PAO1	1	0.5 (2)	0.25 (4)	0.125 (8)	0.25 (4)
PA7	256	128/64 (2/4)	32 (6)	32 (6)	64 (4)
PA14	2	2/1 (-/2)	1 (2)	1 (2)	2/1 (-/2)

[a]Berberine or analog included at 64 µg/mL.
[b]MIC data for PAO1 are the same as in Table S4 and are shown for ease of comparison.
[c]fold change is relative to respective to wild-type PAO1, PA7, or PA14, (-) represents no fold change

Table 3. Antibiotic minimum inhibitory concentration (MIC) in absence or presence of select berberine analogs for *P. aeruginosa* PAO1 and two clinical pan-aminoglycoside resistant strains *(*K2156 and K2161).

Genotype	WT	ΔXY[b]	WT			
Analog[a]	-	-	Berberine	Ber-C3 [5]	Ber C12 [9]	Ber-pAr [10]
P. aeruginosa PAO1						
Kan	64	16 (4)	32 (2)	16 (4)	8 (6)	32 (2)
Ami	1	0.25 (4)	0.5 (2)	0.25 (4)	0.125 (6)	0.5 (2)
Gen	1	0.25 (4)	0.5 (2)	0.25 (4)	0.125 (6)	0.25 (4)
Tob	0.5	0.25 (2)	0.25 (2)	0.25 (2)	0.125 (4)	0.25 (2)
Tig	4	0.5 (6)	4 (-)	2 (2)	2 (2)	4 (-)
Cef	1	0.5 (2)	1 (-)	0.5 (2)	2 (+2)	1 (-)
Cip	0.125	0.125 (-)	0.125 (-)	0.125 (-)	0.5 (+4)	0.25 (+2)
Imi	4	4 (-)	4 (-)	4 (-)	2 (2)	2 (2)
Pan-aminoglycoside resistant *P. aeruginosa* clinical isolate K2156						
Kan	256	16 (8)	16 (8)	16 (8)	8 (10)	16 (8)
Ami	8	0.5 (8)	0.5 (8)	0.5 (8)	0.5 (8)	0.5 (8)
Gen	8	0.5 (8)	1 (6)	0.5 (8)	0.5 (8)	1 (6)
Tob	2	1 (2)	0.5 (4)	0.25 (6)	0.5 (4)	0.5 (4)
Tig	4	0.5 (6)	4 (-)	2 (2)	1 (4)	4 (-)
Cef	2	1 (2)	2 (-)	2 (-)	2 (-)	2 (-)
Cip	1	1 (-)	1 (-)	1 (-)	4 (+4)	2 (+2)
Imi	2 (-)	2 (-)	2 (-)	2 (-)	2 (-)	2 (-)
Pan-aminoglycoside resistant *P. aeruginosa* clinical isolate K2161						
Kan	256	64 (4)	256 (-)	128 (2)	128 (2)	256 (-)
Ami	16	2 (6)	8 (2)	4 (4)	8 (2)	8 (2)
Gen	16	2 (6)	8 (2)	4 (4)	4 (4)	8 (2)
Tob	8	2 (4)	4 (2)	2 (4)	2 (4)	4 (2)
Tig	4	1 (4)	4 (-)	4 (-)	2 (2)	4 (-)
Cef	0.25	0.125 (2)	0.5 (+2)	0.25 (-)	1 (+4)	1 (+4)
Cip	0.125	0.06 (2)	0.25 (+2)	0.25 (+2)	0.5 (+4)	0.25 (+2)
Imi	2	2 (-)	2 (-)	2 (-)	1 (2)	2 (-)

[a]Berberine or dimeric analog included at a fixed concentration of 64 µg/mL where indicated.
[b]Isogenic MexXY knockout of the corresponding parental strain, PAO1 or clinical isolate.
[c]Antibiotics include MexXY-OprM substrates kanamycin (Kan), amikacin (Ami), gentamicin (Gen), tobramycin (Tob), tigecycline (Tig), cefepime (Cef), and ciprofloxacin (Cip), and the nonsubstrate imipenem (Imi).
[d]fold decrease relative to respective wild type PAO1, K2156, or K2161. (-) indicates no change. (+) indicates an increase in MIC.

Schemes

Scheme 1. Overview of berberine analog synthesis.

Figures and Legends

Fig. 1. HTVS identifies berberine analogs as potential probes of MexY. (**a**) Homology model of the MexXY-OprM RND system comprising outer membrane (OM) protein OprM (blue), periplasmic (P) adaptor protein MexX (gold), and inner membrane (IM) transporter MexY (red). (**b**) Schematic of the MexY homotrimer in its three conformational states: access (dark grey), binding (red), and extrusion (light grey). The two putative main substrate entry channels (cleft and vestibule), primary substrate binding region (DBP and PBP), and path to MexX are indicated. A zoomed in view of the binding pocket structure (boxed) also highlights the MexX docking domain, DC (light green) and DN (pink), the porter domain, PN1 (blue), PN2 (teal), PC1 (purple), PC2 (orange), and the switch loop (lime green). (**c**) Checkerboard synergy assays in *P. aeruginosa* PAO1 with berberine and selected analogs from HTVS (*Generation 1*). Increased synergy is observed for Ber-Prop [**3**] and Ber-C6 [**4**] with Kan (top, orange) and Gen (bottom, teal) compared to berberine. Data are shown as the normalized mean of the optical density (OD_{600}) of two biological replicates (0 is no growth, and 1 is maximum growth). The lowest Fractional Inhibitory Concentration (FIC) score for each compound-antibiotic pair is given in the upper right corner.

Fig. 2. Predicted docking of select berberine analogs in the MexY[PAO1] binding pocket reveals potential ligand-specific interactions. Docking of **(a)** berberine and the top three ligands of interest **(b)** Ber-C3 [**5**] , **(c)** Ber-C12 [**9**], and **(d)** Ber-pAr [**10**] in the MexY[PAO1] binding pocket spanning from the PBP to the DBP. Berberine docks in the DBP at the switch loop, Ber-C3 [**5**] and Ber-C12 [**9**] show preference for docking in the DBP, and Ber-pAr [**10**] spans the PBP and DBP.

Fig. 3. Top berberine dimers exhibit increased synergy with aminoglycosides compared to berberine. (a) Checkerboard synergy assays with berberine and top selected berberine dimers in *P. aeruginosa* PAO1. Increased synergy is observed with all ligands and kanamycin (top, orange) and gentamicin (bottom, teal) compared to natural berberine. The lowest fractional inhibitory concentration (FIC) score obtained is shown in the top right corner of each plot. Data are shown as the normalized mean of the optical density (OD_{600}) of two biological replicates (0 is no growth, and 1 is maximum growth). **(b)** Time-kill assays for berberine and dimer analogs at 64 µg/mL performed over 18 hours in both *P. aeruginosa* PAO1 (top) and its *mexXY* isogenic knockout (bottom). Synergy was observed with amikacin for all ligands tested, and Ber-C3 and Ber-C12 reduced the MIC four-fold (red open triangle) compared to no berberine dimer (red filled triangle). Berberine and Ber-pAr reduce the MIC by only two-fold (teal open square) at the tested concentration compared to no berberine dimer (teal filled square). Significant aminoglycoside-independent growth inhibition is observed for Ber-C12 (dashed grey line) compared to the no antibiotic growth control (grey line). Two-fold reduction of the MIC (black open square) for Ber-C12 in *P. aeruginosa* PAO1Δ*mexXY* also suggests off-target activity. Data are shown as the average OD_{600} of biological replicates, each with two technical replicates, with error bars representing the standard deviation.

Fig. 4. Molecular dynamics simulations of berberine and berberine dimers reveal important characteristics for effective chemical probes of efflux function by MexY. (a) 50 ns molecular dynamics simulations of berberine (top left) and berberine dimers, Ber-C3 [**5**] (top right), and Ber-C12 [**9**] (bottom left), and Ber-pAr [**10**] (bottom right) in the MexY[PAO1] DBP and PBP. Conformations at the start (0 ns) and every 10 ns are shown as semi-transparent sticks (0 and 30 ns frames are indicated) and the final conformation (at 50 ns) is shown as solid sticks (orange). Parameters derived from MM/GBSA calculations: **(b)** binding affinity, **(c)** π-packing, and **(d)** ligand strain. Data shown are the individual calculated values for six frames (0 ns and one per 10 ns segment between 0-50 ns) with the average shown as a bar plot. Ber-C3 [**5**] was found to be an effective and specific efflux pump inhibitor for the MexXY-OprM efflux system compared to berberine, Ber-pAr [**9**] and Ber-C12 [**10**]. Statistical analysis was performed using one-way ANOVA on the mean for each ligand and a post-hoc Turkey test was used for pairwise comparison. Statistically significant results are shown (* p≤0.05, ** p≤0.01, *** p≤0.001, and **** p≤0.0001).

Fig. 5. Berberine and berberine dimers are active against *P. aeruginosa* PA7 and PA14. (a) Unique residues identified from alignment of MexY sequences from *P. aeruginosa* PAO1 (orange), PA7 (pink), and PA14 (green) mapped on a cartoon schematic of MexY[PAO1] in the binding state. The three domains of the MexY transporter (docking, porter, and transmembrane) are indicated. **(b)** Time-kill assays for berberine and dimer analogs at 64 µg/mL performed over 18 hours in both *P. aeruginosa* PA14 (top) and PA7 (bottom). Synergy was observed in *P. aeruginosa* PA14 with Gen for Ber-C3 [**5**] and Ber-C12 [**9**] which reduced the MIC two-fold (teal open square) compared to no berberine dimer present (teal filled square). In *P. aeruginosa* PA7, all dimer ligands tested reduced the MIC, at minimum, four-fold (red open triangle) at the tested concentration compared to no berberine dimer (red filled triangle). No significant aminoglycoside-independent growth inhibition was observed for berberine or any berberine dimer (dashed grey line) compared to the no antibiotic growth control (grey line). Data shown are the average OD$_{600}$ measurement with error bars representing the standard deviation. **(c)** A zoomed in view of E175 conformation in MexY[PAO1] (orange) and MexY[PA7] (pink) which permits binding of berberine (top left), Ber-C3 [**5**] (top right), Ber-pAr [**10**] (bottom left), and Ber-C12 [**9**] (bottom right) in the DBP. In contrast, steric hinderance due to the Q175 conformation in MexX[PA14] results in altered predicted ligand to bind in this region.