Rhizosphere and detritusphere habitats modulate expression of soil N-cycling genes during

plant development

Ella T. Sieradzki[1,*], Erin E. Nuccio[2], Jennifer Pett-Ridge[2,3,4], Mary K. Firestone[1,#]

[1] Department of Environmental Science, Policy and Management, University of California

Berkeley, Berkeley, CA 94704, USA

[2] Physical and Life Sciences Directorate, Lawrence Livermore National Laboratory, Livermore,

CA 94551, USA

[3] Life & Environmental Sciences Department, UC Merced

[4] Innovative Genomics Institute, UC Berkeley

[#] Corresponding author: mkfstone@berkeley.edu

* Current address: Laboratoire Ampère, École centrale de Lyon, France

Keywords: soil microbiome, soil nitrogen, rhizosphere, gene expression, metatranscriptomics,

nitrification, plant litter, detritusphere

Abstract

Interactions between plant roots and rhizosphere bacteria mediate nitrogen (N)-cycling

processes and create habitats rich in low molecular weight (growing roots, rhizosphere) and

complex organic molecules (decaying root litter, detritusphere) compared to bulk soil. Microbial

N-cycling is regulated by a diverse suite of genes from many interconnected metabolic

pathways; but most studies of soil N-cycling gene expression have focused on single pathways.

Currently, we lack a comprehensive understanding of the interplay between soil N-cycling gene

regulation, spatial habitat and time. Here we present an analysis of a replicated time series of

soil metatranscriptomes; we followed multiple N transformations in four soil habitats

(rhizosphere, detritusphere, mixed rhizo-/detriusphere, bulk soil) over a period of active root

growth for the annual grass, *Avena fatua*. The presence of root litter and living roots

significantly altered the trajectory of N-cycling gene expression. Across soil habitats, the most

highly expressed N-transformation genes were related to extracellular proteases, ammonium

assimilation into microbial biomass via glutamate synthase, and ammonium oxidation.

Upregulation of bacterial assimilatory nitrate reduction in the rhizosphere suggests that

rhizosphere bacteria were actively competing with roots for nitrate. Simultaneously, bacterial

ammonium assimilatory pathways were upregulated in both rhizosphere and detritusphere soil,

which could have limited N availability to plants. The detritusphere supported dissimilatory

processes DNRA and denitrification. Expression of ammonium oxidation genes was almost

exclusively performed by three phylotypes of *Thaumarchaeota* and was upregulated in

unamended bulk soil. Unidirectional ammonium assimilation and its regulatory genes

(glutamine synthetase/glutamate synthase, or GS/GOGAT) were upregulated in soil surrounding

relatively young roots and more highly decayed root litter, suggesting N may have been limiting

in these habitats (the GS/GOGAT pathway is known to be activated under low N availability).

We did not detect expression of N-fixation or anammox genes. Our comprehensive

metatranscriptomic time-series of organic and inorganic N-cycling in rhizosphere,

detritusphere, and bulk soil, indicates that differences in C and inorganic N availability control

contemporaneous transcription of N-cycling pathways in soil microhabitats that exist in close

spatial proximity.

Contribution to the field

Plant roots modulate microbial nitrogen cycling by regulating the supply of root-derived carbon

and nitrogen uptake. These differences in resource availability cause distinct micro-habitats to

develop: soil near living roots (rhizosphere), decaying roots (detritusphere), near both

(rhizo/detritusphere), or outside the direct influence of roots (bulk). While many genes control

the microbial processes involved in the nitrogen cycle, most research has focused on single

genes and pathways, neglecting the interactive effects these pathways have on each other. The

processes controlled by these pathways determine consumption and production of N by soil

microorganisms. We followed the expression of N-cycling genes in the primary four soil

microhabitats over a period of active root growth for an annual grass. We found that the

presence of root litter and living roots significantly altered gene expression involving in multiple

nitrogen pathways. We also found populations with genes for multiple pathways, where

expression was likely shaped by available forms of carbon and by competition with plants for

inorganic nitrogen. Phylogenetic differences in spatial and temporal expression of the soil

microbial N-pathway genes ultimately regulate N-availability to plants.

Introduction

Nitrogen (N) is a key limiting nutrient for plant growth, since in soil, N mostly occurs in

organic forms within large, complex molecules that are unavailable to plants. Soil

microorganisms have suites of extracellular enzymes capable of degrading these molecules,

freeing small organic and inorganic N to the surrounding soil and benefitting nearby plants.

Other bacteria and archaea transform N to nitrate—an inorganic form that is more available to

many plants. Meanwhile, microorganisms also take up inorganic N for a variety of growth-

supporting assimilatory and dissimilatory metabolic processes. This can lead to some

competition for N between soil bacteria and plants. All of these processes likely occur

simultaneously in soil, but we have an incomplete understanding of how they vary with time

and in distinct soil habitats.

Soils contain numerous microhabitats with heterogeneous distributions of substrates and

resources (Sokol et al., 2022). The influence of growing roots can dominate (rhizosphere) or

dead root litter may dominate (detritusphere); alternatively these two habitats may co-occur

when live roots regrow into previously colonized areas (rhizo/detritusphere). In previous

analyses, we have shown that distinct bacterial guilds operate in rhizosphere, detritusphere,

and combined rhizosphere-detritusphere soil habitats (Nuccio et al., 2020). The spatial

organization of these soil habitats may be particularly important for nutrient transfers that

create distinct N microhabitats that enable or limit N-cycling. For example: (1) in the

rhizosphere, exudate-driven blooms of microbial growth are followed by predation that

liberates the nutrient capital held in bacterial, fungal, and metazoan bodies, this redistribution

of microbial biomass causes conversion to lower molecular weight compounds (*e.g.*, amino

acids) that are deaminated to NH_4^+ by diverse bacteria and fungi (Clarholm, 1985; Koller et al.,

2013); (2) in the detritusphere, organic N in root litter (e.g., lignoproteins (Myrold, 2021)) is

mineralized to make inorganic N available soil microbes, and also serves as a carbon source for

soil microbes (Pepe-Ranney et al., 2016), creating a demand for N to maintain cellular

stoichiometry; (3) in bulk soils, microbes may express more macromolecule degradation genes

relative to more resource rich sites near roots, where degradation enzymes for low molecular

weight compounds are more prevalent (Shi et al., 2018).

Rates of microbial N transformations in soil can be significantly impacted by the presence of

both growing and decaying roots. Gross N-mineralization, immobilization, and nitrification rates

vary as a function of proximity to plant roots, root age and concentration of organic material in

grassland soil (Personeni and Loiseau, 2005; Herman et al., 2006). Roots can significantly

deplete ammonium in the rhizosphere within days of root introduction into sterile soil

(Trofymow et al., 1987). However, in wild soils with intact microbial communities, rhizosphere-

associated bacteria can transiently compete with plants (e.g., *A. barbata* roots) for inorganic

nitrogen (Jackson et al., 1989a). As part of this competition, plants can inhibit rhizosphere

nitrogen transformations such as biological nitrification inhibition (BNI) via production of

certain root exudate compounds (Nardi et al., 2020). Plants have also been shown to inhibit

both nitrification and denitrification processes in the rhizosphere—both processes lead to loss

of N from the plant-available pool (Moreau et al., 2019)—and litter amendment has been

shown to increase denitrification (Che et al., 2018). Understanding the N-cycling tradeoffs

between different soil habitats is key for our ability to better model and predict the controls of

soil N cycling.

Soil metatranscriptome analysis is a powerful tool for linking nitrogen cycling genes to

specific microbial populations, and correlating transcription levels to environmental conditions

(Nicol et al., 2008; Liu et al., 2010; Placella and Firestone, 2013; Nuccio et al., 2020; Sieradzki et

al., 2020; Tosi et al., 2020). In comparison, qPCR can target only single genes, and primer biases

may leave some taxa undetected. A recent metatranscriptomics study demonstrated a short-

term coupling of nitrogen cycling pathways and availability of simple carbons simulating

priming by root exudates (Chuckran et al., 2021). Otherwise, there is little knowledge on the

contemporaneous feedback through gene expression between carbon availability and nitrogen

cycling in naturally complex soil habitats and over annual plant-relevant time frames (Yergeau

et al., 2014, 2018; Žifčáková et al., 2016).

Here, we used soil metatranscriptomics analysis to build a comprehensive time-resolved

representation of both organic and inorganic nitrogen transformations in an annual grassland

soil. We characterized a period of active root growth, as well as root decomposition, of wild oat

grass (*Avena fatua*), a common species in Mediterranean grasslands. We hypothesized that

different N cycle pathways would be prevalent in different soil microhabitats that vary in

available C and N resources. We analyzed 48 metatranscriptomes from rhizosphere and bulk

soil collected over a 3-week time series in the presence or absence of root litter; this allowed us

to identify the dominant N-cycling pathways associated with each habitat and untangle the

effects of carbon supply by live or dead roots from competition with the plant for N. We

previously analyzed a different subset of this data to assess expression of genes coding for

carbohydrate degradation (Nuccio et al., 2020) and organic nitrogen degrading enzymes

(Sieradzki et al., 2020). While these prior studies provide important context, our analysis here is

unique in its comprehensive exploration of how N transformation gene expression is influenced

by the distinct environmental conditions of different soil habitats. The use of

metatranscriptomics (as opposed to qPCR) also facilitated identification of specific microbial

populations involved in each pathway, and helped us test for multiple N transforming pathways

expressed within populations found in mutiple soil habitats. Finally, we measured changes in N-

pathway gene expression over time as the rhizosphere aged and as root litter was degraded.

Methods

Experimental design, sample collection and sequencing

The experimental procedures and initial data processing for our 48 soil

metatranscriptome dataset are described in detail by (Nuccio et al., 2020). Briefly, *Avena fatua*

was grown in a fine loam Alfisol complex (Ultic Haploxeralf mixed with a Mollic Palexeralf) from

the Hopland Research and Extension Center (Hopland, CA); pH 5.6, 2% total C) in microcosms

with a sidecar with transparent walls. After six weeks, roughly halfway through the plant life

span, the solid divider to the sidecar was removed and replaced with a slotted divider so that

roots could grow into the sidecar. Root growth was marked on the sidecar wall. All sidecars

contained bulk soil bags that were inaccessible to roots. Half of the bulk soil bags as well as the

soil in half of the sidecars were amended with dried *A. fatua* root detritus (litter C:N = 13.4).

Once the sidecar was opened, paired rhizosphere and bulk soil triplicates were harvested

destructively after 3, 6, 12 and 22 days from amended and unamended microcosms for a total

of 48 samples. Lifeguard Soil Preservation Reagent (MoBio) was added to 1g subsample of the

harvested soil. The supernatant and roots were removed, and the samples were stored at -

80°C.

DNA/RNA co-extraction was performed with phenol-chloroform. DNA and RNA were

separated with an AllPrep kit (Qiagen) and RNA was DNase treated (TURBO DNase, Thermo-

Fisher Scientific). Ribosomal RNA was depleted (RiboZero, Illumina) and the mRNA was

retrotranscribed. cDNA was sequenced with an Illumina HiSeq 2000 2x150 (TruSeq SBS v3)

protocol at the Joint Genome Institute. In addition, the V4 hypervariable region of the 16S gene

(primers 515F and 806R) was amplified from the extracted DNA and sequenced via Illumina

MiSeq v3 2x300. Amplicons were analyzed by the Joint Genome Institute. Operational

taxonomic units (OTUs) were clustered at 97% identity with USEARCH (Edgar, 2010) and

taxonomy was assigned at 95% id by RDP (Wang et al., 2007).

Expression of genes from a curated collection of Hopland-soil genomes

As described previously (Nuccio et al., 2020), raw reads were quality-trimmed (Q20) and

rRNA and tDNA were removed. Library size, evenness, richness and Shannon diversity were

comparable between experimental groups, with a mean library size of 43 M paired end reads.

Reads were mapped at 80% identity with BBsplit (Bushnell, 2014; Nuccio et al., 2020) against a

dereplicated reference set of 282 Hopland CA soil genomes including stable isotope probing

metagenome assembled genomes (SIP-MAGs) (Starr et al., 2018), metagenomic assembled

genomes (MAGs, NCBI PRJNA517182), soil isolates (Starr et al., 2018; Zhalnina et al., 2018) and

single amplified genomes (SAGs) (Nuccio et al., 2020). A mean of 12.3% of the reads per sample

mapped unambiguously to the reference set.

Prodigal (Hyatt et al., 2010; Nuccio et al., 2020) was used to predict open reading frames

(ORFs) in the genomes. The ORFs were annotated using KEGG (Kanehisa and Goto, 2000) and

ggKbase (http://ggkbase.berkeley.edu). Gene counts were calculated with R package Rsubread

function featureCounts (Liao et al., 2014). Three samples were removed from further analysis

due to an exceptionally low number of features: H3_Rhizo_NoLitter_39, H1_Rhizo_Litter_2 and

H2_Rhizo_Litter_9.

All features and their metatranscriptomic read recruitment were analyzed with the R

package DESeq2 (Liao et al., 2014; Love et al., 2014) requiring an adjusted p-value < 0.05.

Ordination and visualization were conducted using R package ggplot2 (Wickham, 2016) and

vegan (Oksanen et al., 2008; Wickham, 2016); the vegan function Adonis was used to detect

significant factors affecting expression of nitrogen cycling genes.

Expression of nitrogen cycling genes identified in assembled metatranscriptomes

After quality control, the metatranscriptomes were assembled into contigs within

samples. Only contigs larger than 200 bp were clustered at 99% ID with cd-hit-est (Huang et al.,

2010). Prodigal (Huang et al., 2010; Hyatt et al., 2010) was used to predict ORFs from the

cluster representatives, and KEGG models (Kanehisa and Goto, 2000) were used to locate

nitrogen cycling genes based on the KEGG nitrogen cycle module (see Supplemental Table S1

for a list of KEGG profiles, gene lengths and thresholds used). Some ORFs were identified by

several HMMs due to homology (e.g. nitrate reductases *nxrA*, *narG* and *napA*). In those cases,

the highest scoring (by bitscore) hit from all HMMs was selected. Extracellular proteases were

called by reciprocal blast to proteases from the MEROPS database that contained signal

peptides (Rawlings et al., 2016; Nguyen et al., 2019). Reads were mapped to nitrogen cycling

ORFs at a minimum identity 95% and minimum breadth 75% using bbmap (Bushnell, 2014;

Nuccio et al., 2020). Normalization to sequencing depth was done using DESeq2, and when

comparing between different genes expression was also divided by the relevant mean

assembled gene length to control for gene-length biases in recruitment. Heat maps were

generated in R using gplots (Warnes et al., 2015).

Phylogeny of the ammonia monooxygenase subunit A (*amoA*) was determined by

aligning *amoA* assembled transcripts to curated full amoA gene sequences from RefSeq (Jun 1,

2019) using mafft-linsi (sensitive mode) (Katoh and Standley, 2013) and removing positions

represented by less than 50% of the aligned sequences with GBlocks version 0.91 (Dereeper et

al., 2008) with parameters -b3=50 -b4=5 -b5=h. RAxML version 8.2.12 (Kozlov et al., 2019) with

parameters set to matrix LG and seed=13 was used to build a Newick tree which was visualized

in iTOL (Letunic and Bork, 2019). The phylogeny of *nirK* transcripts was determined in the same

manner. The aggregated coverage of *amoA*/*nirK* variants from all samples was superimposed

onto the tree in iTOL. Additionally, since nitrate reductases that participate in multiple

pathways are homologous, we wanted to make sure they were correctly identified by the

HMMs. Therefore, we placed the assembled sequences on a protein reference tree (Matheus

Carnevali et al., 2019) and corrected the annotations according to tree clustering.

Normalized expression per gene per time point was compared between groups using

ANOVA and Tukey HSD tests, and p-values were corrected for multiple comparisons (time,

location, amendment and all interactions between them). Upregulation was determined as

log2-fold values calculated by DESeq with corrected *p* values < 0.05.

Results

To assess whether expression of organic and inorganic nitrogen cycling genes was

affected by the presence of root litter in the rhizosphere or in nearby bulk soil, we tested for

significant differences in N transformation gene expression over a three-week time series (3, 6,

12, 22 days). The PCoA based on Bray-Curtis dissimilarity of expression of N cycling genes

normalized to sequencing depth revealed clustering by location (rhizosphere/bulk soil)

throughout the whole time-series; the last time point (T4; 22 days) notably distinct from the

others (fig. 1A). Analysis of the first three time points only (3, 6, and 12 days) revealed an

additional clustering by litter amendment (fig. 1B). In both analyses, the first two principal

components explained 40% of the variation. PERMANOVA analysis using all time points

indicated a significant effect of location (rhizosphere vs. bulk soil), time and litter amendment

(p-value = 10^{-4}) (see sup. table 1 for all values). The combined R^2 of all significant factors was

63%. When we conducted the same analysis but excluded the last time point (T4), the effects of

single factors remained significant (p-value = 10^{-4}). The combined R^2 of all significant factors

was 61%, and the variability explained by time decreased from 28% with T4 to 10% without T4,

whereas the variability explained by location and amendment increased by 5-8% each.

Figure 1: Principal coordinates analysis (PCoA) of nitrogen cycling gene expression. Influence of

living roots and root detritus on soil microbial gene expression during 3 weeks of *Avena fatua*

root growth (harvests at 3, 6, 12, 22 days) in a CA annual grassland soil. Input data are Bray-Curtis

dissimilarity of expression of N cycling genes normalized to sequencing depth, determined by

mapping reads to ORFs assembled from metatranscriptomes followed by log ratio

transformation, in (A) all time points with the last time point circled and (B) excluding the last

time point (22 days). Groups are indicated in the ellipses: B=Bulk, BL=Bulk Litter, R=Rhizosphere,

RL=Rhizosphere Litter. n=3 for each habitat and timepoint. Ellipses represent a 90% confidence

interval of the ordination coordinates (calculated by stat_ellipse).

Expression of nitrogen cycling genes by pathway

To identify the dominant N-cycling pathways in four soil microhabitats (rhizosphere,

detritusphere, rhizo/detritusphere, bulk soil), we compared differential expression of each

nitrogen cycling gene in the presence or absence of roots and litter amendment. Expression and

upregulation trends of N cycling genes (defined in Table S1) are shown in fig. 2. Several genes

involved in assimilatory pathways were significantly upregulated in the rhizosphere:

assimilatory nitrate reduction genes *nasA*, *NR* and *nirA*, glutamate dehydrogenase *gdh2* and

GS/GOGAT genes *glt1* and *gltB* (fig. 2). The first step of GS/GOGAT, *glnA*, was upregulated both

in the rhizosphere and in the presence of litter (fig. 2), and the GS/GOGAT regulatory proteins

glnD and *glnG*, triggered under N limitation, were upregulated in the rhizosphere, whereas *glnB*

was upregulated in the presence of litter (sup. fig. 1 B,C). In the detritusphere, the presence of

litter also triggered macromolecular organic nitrogen degradation (extracellular protease and

chitinase), as well as dissimilatory pathways; dissimilatory nitrate reduction (DNRA) and

denitrification (fig. 2). However, not all genes in these pathways were upregulated and some

were not detected at all (*nirS* and *norBC* in denitrification), whereas others were not

significantly upregulated (*nirK*, *narGHI*, *napB*) (fig. 2). Finally, nitrification, DNRA and the first

step of denitrification were upregulated in bulk soil compared to the rhizosphere (sup. fig. S1).

Several genes were particularly highly expressed in our dataset (e.g., ammonia

monooxygenase subunits *amoA* and *amoC,* and nitrate reductase *nirK* (sup. fig. S1 A,B)) but did

not show clear patterns of differential expression by either soil habitat treatment or time. We

did not detect any genes involved in nitrogen fixation or anaerobic ammonia oxidation.

Expression of fungal N cycle genes such as *cyp55* (nitric oxide to nitrous oxide) and *nit-6* (nitrite

to ammonium) was very low (sup. fig. 1C).

Figure 2: Expression level and upregulation by soil habitat of nitrogen cycling genes from assembled transcripts. Bold font indicates high expression levels (mean counts in each time point >5 after normalization to gene length and sequencing depth). Significant upregulation was determined by ANOVA and Tukey post hoc test (adjusted p<0.05). Numbers denote pathways: 1=Ammonia oxidation, 2=Nitrite oxidation, 3=Assimilatory nitrate reduction,

4=Denitrification, 5=Anammox, 6=Nitrogen fixation, 7=DNRA, 8=Ammonium assimilation (GDH

and GS/GOGAT), 9=Macromolecular N mineralization, amt = Ammonium transporter. The

functional role of each enzyme is explained in Table S2. The nirK analog is an archaeal enzyme

which participates in archaeal nitrification, but unlike its bacterial analog, it does not generate

nitrate (Stein, 2019).

Archaeal nitrification

Two of the most highly expressed genes we identified were subunits of ammonium

monooxygenase, the first enzyme in the nitrification pathway. Nitrification genes were not

affected by the presence of litter, but *nirK*, *amoB*, *hao*, *nxrA* and *nxrB* were significantly

upregulated in bulk soil compared to rhizosphere soil (ANOVA: p<0.05, Tukey HSD test:

adjusted p<0.05) (sup. table S2).

Ammonium oxidation can be performed either by ammonium oxidizing archaea (AOA)

or ammonium oxidizing bacteria (AOB). The *amoA* subunit of ammonia monooxygenase is

commonly used as a taxonomic marker to differentiate between AOA and AOB. To determine

which taxa were responsible for the high expression of ammonia oxidation genes, we placed

the 13 *amoA* ORFs identified in assembled transcripts into a phylogenetic tree with full-length

amoA reference sequences from the RefSeq database (fig. 3A). The cumulative coverage per

amoA variant in all samples was superimposed onto the tree. Read recruitment to AOA was

orders of magnitude higher than to AOB, accounting for 98% ± 0.007% (N=16) of the reads

mapped to *amoA* per sample. While there were some ORFs placed into comammox clade B,

their expression was negligible compared to archaeal *amoA* (fig. 3A).

The expression of hydroxylamine oxidoreductase (*hao*), the next step in nitrification

after ammonium oxidation, was extremely low across all samples. Similarly, expression of

nitrite oxidase (*nxrAB*) was very low. However, the expression of *nirK* was very high (fig. 2). A

phylogenetic analysis placed 96 out of 158 *nirK* assembled ORFs in the archaeal clade and, as in

the case of *amoA*, revealed that the archaeal variants recruited more than 95% ± 0.07% (N=16)

of the reads that mapped to *nirK* (fig. 3B). Three archaeal phylotypes dominated *nirK*

expression, with 76% ± 0.07% of all reads mapped to *nirK*.

To determine whether the high expression of archaeal nitrification genes can be

explained by abundance of AOA compared to AOB, we searched our *16S rRNA* gene amplicons

(Nuccio et al., 2020) for known AOA and AOB. The relative abundance of AOA *16S rRNA* genes

was 2- to 5-fold higher than for AOB, and the relative abundance of AOB was consistently

higher than that of nitrite-oxidizing bacteria (NOB) (fig. 3C). The aggregate expression of *amoA*

and *nirK* was well correlated with a ratio of *nirK*:amoA = 1.6 (R^2=0.7, *p*<0.0001) (fig. 3D). In

addition, the aggregated expression of amoA was also highly correlated to that of

Thaumarchaeal extracellular proteases (Sieradzki et al., 2020) (adjusted R^2=0.61, *p*<0.0001).

Figure 3: Phylogenetic determination of nitrifiers by *amoA*, *nirK* and *16S-rRNA*. Phylogeny of

assembled ORFs (black) (A) *amoA* and (B) *nirK* placed into phylogenetic trees containing curated

reference sequences from RefSeq (grey). Colors represent phylogenetic clades. Bars around the

trees represent cumulative coverage from all samples to specific assembled transcripts. (C)

Relative abundance of AOA, AOB and nitrite oxidizing bacteria (NOB) of the entire microbial

community by *16S-rRNA* gene abundance. (D) Positive correlation between expression of *amoA*

and *nirK* transcripts by location and amendment (R^2=0.7). (E) Positive correlation between

expression of *amoA* and *Thaumarchaeota* extracellular protease transcripts by location and

amendment (R^2=0.61).

Ammonium assimilation and transport pathways

The most highly expressed ammonia acquisition pathway was glutamate synthase

(GS/GOGAT), which can be activated by a cascade of four enzymes under nitrogen limitation

(fig. 4A) (MacNeil et al., 1982; Bueno et al., 1985; Leigh and Dodsworth, 2007). As opposed to

single-gene dependent transporters (*amt*) and glutamate dehydrogenase (GDH), GS/GOGAT is a

two-step pathway: the first step is encoded by the gene *glnA*, and the second step either by

glt1, Glu or the heterodimer *gltBD*. Both ammonium transporters (sup. fig. 1) and GS/GOGAT

(*glnA*, fig. 4B) were significantly upregulated in detritusphere microhabitats within the young

rhizosphere (Rhizo Litter, 3 days) and bulk soil at the final timepoint (Bulk Litter, 22 days) (fig.

4b). *glnA* was also significantly upregulated in the early rhizosphere compared to the bulk

control (Rhizo, fig 4B), and three out of four detected proteins involved in GS/GOGAT were also

upregulated in the rhizosphere after 3, 6 and 12 days (sup. fig. S3). We found that expression of

gltBD was generally much higher than that of *glt1*, whereas *Glu* was not detected (sup. fig. 1).

Expression of an additional ammonium assimilation pathway, glutamate dehydrogenase (GDH),

was low (sup. fig. S1). GDH consists of a single step performed by one of two enzymes: *gdh2* or

gudB. *Gdh2* was upregulated in the rhizosphere, whereas *gudB* was upregulated in the

presence of litter (sup. fig. S1, sup. table S2).

In theory, there is an advantage to using GS/GOGAT for ammonia assimilation over GDH

under N limitation, as GS/GOGAT in a one-way pathway, whereas GDH can also catalyze

deamination of glutamate which might lead to loss of ammonium via diffusion. Expression of

GS/GOGAT was an order of magnitude higher than that of GDH under all experimental

conditions (fig. 4C). The ratio of expression (GS/GOGAT to GDH) significantly increased over

time (Permanova ratio~group*time, p(time)=0.002). In the young rhizosphere the ratio

appeared to be higher in the rhizosphere compared to bulk soil, whereas in the mature

rhizosphere, litter seemed to trigger a higher ratio (fig. 4D). Expression of the GS/GOGAT

regulatory proteins *glnD, glnG and glnL* was significantly upregulated in the rhizosphere

compared to bulk soil, particularly in the young rhizosphere (Table S3). Most regulatory genes

were also upregulated in the detritusphere compared to unamended soil at the last time point.

Figure 4: Expression of ammonium assimilation pathways and their regulatory genes. (A)

Simplified conceptual representation of *glnA* activation via the regulatory cascade (B)

Normalized expression of *glnA* in amended/unamended rhizosphere and bulk soil. Asterisks

denote a significant increase compared to the same litter treatment in bulk soil (e.g.,

unamended rhizosphere vs. unamended bulk) (C) ratio of GS/GOGAT to GDH (*glnA*:*gudB)*

expression in the metatranscriptomes. T. (D) Mean differential expression compared to

unamended bulk soil of regulatory genes that activate GS/GOGAT under N limitation as well as

glnA.

Discussion

In this study we used metatranscriptomic gene expression (% of reads mapped to a

gene) and upregulation (fold change of expression in a habitat compared to unamended bulk

soil) to identify contemporaneous N transformations in the main soil habitats: rhizosphere

(near live roots), detritusphere (near dead roots), rhizo-detritusphere (live and dead roots) and

bulk soil (neither). We found that expression of genes coding for N-cycling processes varied

between soil habitats as well as over time. The use of metatranscriptomic analyses allowed us

to track multiple N cycling pathways occurring at the same time. Moreover, as N pools in this

grassland soil are relatively constant but fluxes are high, it has been suggested that studying N

cycling parameters may be a better proxy than pools (Sudderth et al., 2012).

Habitat and temporal patterns of N cycling gene expression were interrelated. Effects of

habitat on gene expression were detected at the first three sampling times, but not after 22

days. At this time point, the soil was much drier due to plant transpiration (Nuccio et al., 2020).

This decrease in soil moisture likely affected rates of substrate diffusion as well as microbial

mobility, leading to changes in the environmental milieu experienced by the microbes.

Removing the last time point from analyses, revealed strong effects of roots and litter as well as

the interaction between the two, implying that rhizosphere bacteria respond to litter

amendment differently than bulk soil bacteria. At 22 days, the roots present in the rhizosphere

treatment may have begun senescence and degradation processes more similar to that of root

litter. . Moreover, root exudation decreases as the root senesces, thus likely reducing the effect

of simple carbon sources originating from exudation as well as driving usage of litter, which is a

less labile C source (Shi et al., 2018).

Several N cycling pathway genes were differentially expressed in different soil habitats.

In the rhizosphere we identified upregulation of assimilatory nitrate reduction and ammonium

assimilation through glutamate dehydrogenase (GDH; *gdh2*) and glutamine synthetase

GS/GOGAT; *glnA, glt1, gltB*). An additional GDH enzyme, *gudB*, was upregulated in the litter-

amended rhizosphere. Three of four regulatory enzymes that activate GS/GOGAT under N

limitation were also upregulated in the rhizosphere, and the fourth was upregulated in the

litter-amended rhizosphere. The upregulation of these pathways likely indicate that the

rhizosphere microbial community experiences inorganic N limitation. N limitation can stem

from competition over inorganic N with the plant (Jackson et al., 1989b; Schimel et al., 1989), as

well as from a higher demand for N as microbial density is ca. 10X higher in the rhizosphere

compared to bulk soil (Pett-Ridge et al., 2021). *Avena* plants have been shown to be better

competitors than their rhizosphere community for nitrate, potentially explaining why

rhizosphere microorganisms would upregulate nitrate reduction to ammonium.

Pathways upregulated in the detritusphere compared to bulk soil were dissimilatory

nitrate reduction (DNRA), denitrification, ammonium transporters, mineralization of organic N

and ammonium assimilation via GDH (*gudB*) and GS/GOGAT (*glnA*, *gltD*). Dissimilatory

processes, often overlooked in soil N cycling studies (Moreau et al., 2019), can eventually lead

to loss of gaseous nitrogen from the system either as N_2, N_2O or NO. As N is commonly a

limiting nutrient in the rhizosphere, coveted by both plants and microbes, dissimilatory

processes might be disadvantageous in this habitat. Indeed, plants have been demonstrated to

outcompete nitrate reducing bacteria for nitrate (Moreau et al., 2015) and there is some

evidence that root exudates can inhibit denitrification (Dassonville et al., 2011). Some of the

genes involved in dissimilatory processes were also significantly upregulated in litter-amended

bulk soil compared to litter-amended rhizosphere, also suggesting the inhibition of N-loss

processes in rhizosphere soil. Expression of assimilatory processes that were upregulated in the

detritusphere compared to bulk soil (*gudB, gltD*) was higher than that of any DNRA or

denitrification genes. Assimilatory processes are possibly a more common strategy for N

acquisition in the detritusphere, and their upregulation compared to unamended soil is likely

driven by carbon availability from root litter to maintain C:N cellular stoichiometry. Compared

to bulk soil, both detritusphere and rhizosphere have a much higher oxygen consumption due

to C utilization and biomass increase. However, in the rhizosphere there is extreme competition

over nitrate against the plant, whereas in the detritusphere the driving force is only C and O

consumption which may create anaerobic or microaerobic niches that would support DNRA and

denitrification.

Bulk soil was characterized by upregulation of genes coding for nitrification, another

process that leads to loss of N from the system through generation of nitrous oxide and

leaching of nitrate from soil (Prosser et al., 2020). Ammonia monooxygenase subunit C (*amoC*)

was also one of the most highly expressed genes in this soil. Interestingly, the expression of

subsequent steps in bacterial nitrification (*hao, nxrA, nxrB*) was extremely low. However,

expression of nirK, traditionally considered a part of the denitrification process, was one of the

highest we detected. When we explored the phylogeny of detected amoA and nirK variants, we

discovered that all highly transcribed variants were archaeal. nirK has been detected in archaeal

genomes and the current leading hypothesis is that archaea oxidize hydroxylamine by

combining it with nitric oxide (NO), yielding two nitrite molecules, one of which is reduced back

to NO by *nirK* (Prosser et al., 2020).

While it is possible that conditions are more favorable for nitrification in bulk soil as

there is less competition over ammonium in the absence of plant roots, it is also possible that

nitrification was downregulated or inhibited in the rhizosphere. Biological nitrification inhibition

(BNI) by the exudation of secondary metabolites from plant roots that can suppress ammonia

oxidation by nitrifying microorganisms (Subbarao et al., 2007, 2013), thus increasing the

availability of ammonium to plants. Hence it is proposed as one strategy to increase crop yields

and reduce fertilizer loss (Coskun et al., 2017). Root exudates of *A. fatua* have been shown to

inhibit nitrification by ammonia oxidizing bacterium *Nitrosomonas europea* (O'Sullivan et al.,

2017). If *A. fatua* inhibited ammonia oxidizing archaea in order to increase ammonium

availability to its roots, it would likely have appeared as upregulation in bulk soil. Given that

microbes in the rhizosphere appear to be experiencing N-limitation, our observations from the

rhizosphere and bulk soil may be partially explained by competition over ammonium between

rhizosphere microorganisms and the plant.

Despite the low relative abundance of AOA, archaeal genes for ammonia

monooxygenase (*amoABC*) were among the most highly expressed out of all N-cycling

transcripts in this study across time, location (bulk soil vs. rhizosphere) and presence or absence

of litter amendment. The relative abundance of AOA in our study based on 16S-rRNA gene

abundance, while never exceeding 1% of the microbial community, was several-fold higher than

that of ammonia oxidizing bacteria (AOB) and matches previous results from other grasslands

(Leininger et al., 2006; Nicol et al., 2008; Zeglin et al., 2011). While they may be very slow

growers, AOA can be highly metabolically active in soil (Pratscher et al., 2011). Moreover,

transcription of archaeal *amoA* has also been shown to be correlated to nitrification rates

(Zhang et al., 2010; Pratscher et al., 2011; Orellana et al., 2019). A similar experiment using a

different species of *Avena* demonstrated that gross nitrification rates decreased with

rhizosphere age (Herman et al., 2006). This pattern further supports correlation between *amoA*

expression and nitrification rates, and could reflect increasing competition over labile N as well

as reduced root exudation by the plant in aging rhizosphere (Jackson et al., 1988; Hu et al.,

2001; Zhalnina et al., 2018).

Expression of *amoA* has been previously shown to be dominated by few archaeal

phylotypes (Tourna et al., 2008; Offre et al., 2009; Gubry-Rangin et al., 2011). Not only do our

results support this, but they quantitatively show that on average 98% of the reads mapped to

amoA were assigned to three variants, all archaeal. All three phylotypes were within the

Nitrososphaera clade, commonly found in soil (Tolar et al., 2017). Similarly, *nirK* expression was

also dominated by three variants, but surprisingly these were most closely related to another

soil AOA genus – *Nitrosocosmicus*. Additionally, expression of *amoA* and *nirK* is well correlated.

It is possible that Nitrososphaera are performing ammonium oxidation and *Nitrosocosimus*

oxidizing the product. While there are ammonia oxidizing and nitrite oxidizing bacteria, such a

cooperation is not known in archaea. However, it is also possible that the taxonomic placement

of one of the genes is wrong. Both *amoA* and *nirK* have been identified in curated genomes of

AOA (Jung et al., 2014; Bayer et al., 2016; Carini et al., 2018) which raises the possibility that the

same AOA cell can enact both steps of nitrification (Kozlowski et al., 2016; Reji et al., 2019), but

dual expression of amoA and nirK by AOA has not yet been shown experimentally. Our strong

correlation suggests that there may be a relationship between these two processes, but further

experimentation is required to determine if they are mechanistically linked.

Thaumarchaeal extracellular proteases identified in the same dataset were previously

found to be upregulated in bulk soil (Sieradzki et al. 2022) and their temporal expression

patterns highly correlated with *amoA* expression. Thaumarchaeota have been shown to prefer

ammonium from organic nitrogen sources such as urea, amino acids and peptides in forest soil

(Levičnik-Höfferle et al., 2012). As AOA have to compete with the plant over ammonium in the

rhizosphere, they may turn to organic N instead (Craine et al., 2007; Blagodatskaya and

Kuzyakov, 2008), explaining the tight correlation between *amoA* and AOA proteases.

Ammonium assimilation via glutamate synthase (GS/GOGAT) is generally thought to be

preferred to glutamate dehydrogenase (GDH) under N limitation due to the K_m of GS/GOGAT

which is an order of magnitude lower than that of GDH (Sakamoto et al., 1975; Alibhai and

Villafranca, 1994). In our study GS/GOGAT gene expression was widespread across treatments

and over time, with significant upregulation in response to both roots and litter. Moreover, the

expression of GS/GOGAT was consistently at least an order of magnitude higher than that of

GDH. The first step of the GS/GOGAT pathway, *glnA* (glutamine synthetase), was one of the

most highly expressed genes in our study. The P_{II} regulatory protein *glnB* which is activated

during N limitation and activates GS/GOGAT over GDH (Huergo et al., 2013) was also highly

expressed under all experimental conditions and upregulated in the presence of live or dead

roots. In contrast, expression of single-enzyme pathway glutamate dehydrogenase (GDH) was

much lower. Assuming comparable downstream post-transcriptional regulation, a preference

for multi-step unidirectional pathway GS/GOGAT (Hochman et al., 1988) as opposed to the

bidirectional GDH, which could also perform glutamate deamination (Melo-Oliveira et al.,

1996), could result from nitrogen limitation, as it prevents nitrogen loss from the organism

(Yuan et al., 2009).

We identified differences in expression of GS/GOGAT and GDH between habitats.

Expression of *glnA* (GS/GOGAT) was higher in the rhizosphere compared to bulk soil at 3, 6 and

12 days, and was higher in litter amended soil compared to unamended both in bulk soil and in

the rhizosphere. Its expression was highest in the presence of both live and dead roots (litter-

amended rhizosphere) at 3 and 6 days, indicating an additive effect of live roots and root litter.

This effect may be due to high resource demand when C is available by both litter and

exudates, which exacerbated the demand for N to maintain cellular stoichimetry (Zhu et al.,

2014; Moreau et al., 2019). Expression of carbohydrate active enzymes (CAZy) in the same soil

was also higher in the young rhizosphere and decreased over time and expression of

extracellular protease genes was highest at 3 days (Nuccio et al., 2020; Sieradzki et al., 2020).

This additive effect, generally accepted to be short-term (Kuzyakov et al., 2000), disappears by

6 days of rhizosphere maturation.

Temporal effects of ammonium assimilation gene expression varied by gene, pathway

and habitat. For example, the two GDH genes we identified were upregulated in different

habitats: rhizosphere (*gdh2*) and detritusphere (*gudB*). The expression of both genes decreased

over time, possibly because ammonium concentrations dropped to a concentration too low for

the affinity of this enzyme (Alibhai and Villafranca, 1994). The expression of *glnA* (GS/GOGAT)

tells a more complicated story. In the litter-amended rhizosphere its expression decreases over

time, as does the expression of its regulatory genes. However, in the litter-amended bulk soil

the expression of *glnA* increases from day 6 on. The consistently lower expression of this

enzyme in unamended bulk soil implies that any C input, be it from rhizodeposition of root

litter, creates a demand for N that is at least somewhat alleviated via the GS/GOGAT pathway.

Finally, we posited that there is a trade-off between the GDH and GS/GOGAT pathways that

corresponds to nitrogen limitation. Therefore, we examined the ratio of gene expression

between *glnA* (GS/GOGAT) and the more highly expressed of the GDH genes, *gudB*. This ratio

was consistently higher than 10, implying a general preference for GS/GOGAT, possibly due to

low N availability at all times with or without the presence of roots or root litter. This ratio also

increased over time under all experimental conditions, but this increase began earlier in the

litter-amended rhizosphere. However, by 22 days the highest ratio was in litter-amended bulk

soil, possibly because growth rates are slower in bulk soil compared to the rhizosphere (Pett-

Ridge et al., 2021) and therefore N limitation developed slower.

Conclusions

Here we present a comprehensive analysis of expression of all nitrogen cycling pathways

over time and in the main soil habitats: rhizosphere, litter-amended rhizosphere, litter-

amended bulk soil and unamended bulk soil. We propose several classes of controllers on the

expression of nitrogen cycling genes in soil with or without roots and/or root litter. Processes

that lead to loss of N from the system were downregulated near live roots. Availability of

carbon (C) and the quality of that C , whether from root exudates or root litter, could drive N

uptake to maintain cellular stoichiometry, and proximity to live roots can lead to competition

for inorganic nitrogen. The use of metatranscriptomics allowed us to track multiple pathways

and identify trade-offs between them which are likely based on N availability and limitation.

Acknowledgements

The authors would like to thank Dr. Graeme Nicol, Dr. Christina Hazard, Dr. Kataryna

Zhalnina, Dr. Rachel Neurath, Dr. Rachel Hestrin, Dr. Nameer Baker, Alexa Nicholas, Katerina

Estera-Molina, Dr. Spencer Diamond and Dr. Jillian Banfield for insightful discussions and

emotional support during the covid-19 pandemic. This research was supported by the U.S.

Department of Energy Office of Science, Office of Biological and Environmental Research

Genomic Science program under Awards DE-SC0020163 and DE-SC0016247 to M.K.F at UC

Berkeley and awards SCW1589 and SCW1678 to J.P-R. at Lawrence Livermore National

Laboratory. Work conducted at Lawrence Livermore National Laboratory was supported under

the auspices of the U.S. DOE under Contract DE-AC52-07NA27344. Sequencing was conducted

as part of Community Sequencing Awards 1487 to J.P.R. and 1472 to M.K.F..

E.T.S was partially supported by Marie Sklodowska-Curie postdoctoral fellowship "DIVOBIS".

References

Alibhai, M., and Villafranca, J. J. (1994). Kinetic and mutagenic studies of the role of the active site
residues Asp-50 and Glu-327 of Escherichia coli glutamine synthetase. *Biochemistry* 33, 682–686.

Bayer, B., Vojvoda, J., Offre, P., Alves, R. J. E., Elisabeth, N. H., Garcia, J. A., et al. (2016). Physiological
and genomic characterization of two novel marine thaumarchaeal strains indicates niche
differentiation. *ISME J.* 10, 1051–1063.

Blagodatskaya, E., and Kuzyakov, Y. (2008). Mechanisms of real and apparent priming effects and their
dependence on soil microbial biomass and community structure: critical review. *Biol. Fertil. Soils* 45,
115–131.

Bueno, R., Pahel, G., and Magasanik, B. (1985). Role of glnB and glnD gene products in regulation of the
glnALG operon of Escherichia coli. *J. Bacteriol.* 164, 816–822.

Bushnell, B. (2014). BBTools software package. *URL http://sourceforge. net/projects/bbmap*.

Carini, P., Dupont, C. L., and Santoro, A. E. (2018). Patterns of thaumarchaeal gene expression in culture
and diverse marine environments. *Environ. Microbiol.* 20, 2112–2124.

Che, R., Qin, J., Tahmasbian, I., Wang, F., Zhou, S., Xu, Z., et al. (2018). Litter amendment rather than
phosphorus can dramatically change inorganic nitrogen pools in a degraded grassland soil by
affecting nitrogen-cycling microbes. *Soil Biol. Biochem.* 120, 145–152.

Chuckran, P. F., Fofanov, V., Hungate, B. A., Morrissey, E. M., Schwartz, E., Walkup, J., et al. (2021). Rapid
Response of Nitrogen Cycling Gene Transcription to Labile Carbon Amendments in a Soil Microbial
Community. *mSystems* 6. doi: 10.1128/mSystems.00161-21.

Clarholm, M. (1985). Interactions of bacteria, protozoa and plants leading to mineralization of soil
nitrogen. *Soil Biol. Biochem.* 17, 181–187.

Coskun, D., Britto, D. T., Shi, W., and Kronzucker, H. J. (2017). Nitrogen transformations in modern
agriculture and the role of biological nitrification inhibition. *Nature Plants* 3. doi:
10.1038/nplants.2017.74.

591 Craine, J. M., Morrow, C., and Fierer, N. (2007). Microbial nitrogen limitation increases decomposition.
592    *Ecology* 88, 2105–2113.

593 Dassonville, N., Guillaumaud, N., Piola, F., Meerts, P., and Poly, F. (2011). Niche construction by the
594    invasive Asian knotweeds (species complex Fallopia): impact on activity, abundance and community
595    structure of denitrifiers and nitrifiers. *Biol. Invasions* 13, 1115–1133.

596 Dereeper, A., Guignon, V., Blanc, G., Audic, S., Buffet, S., Chevenet, F., et al. (2008). Phylogeny.fr: robust
597    phylogenetic analysis for the non-specialist. *Nucleic Acids Research* 36, W465–W469. doi:
598    10.1093/nar/gkn180.

599 Edgar, R. C. (2010). Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 26,
600    2460–2461.

601 Gubry-Rangin, C., Hai, B., Quince, C., Engel, M., Thomson, B. C., James, P., et al. (2011). Niche
602    specialization of terrestrial archaeal ammonia oxidizers. *Proc. Natl. Acad. Sci. U. S. A.* 108, 21206–
603    21211.

604 Herman, D. J., Johnson, K. K., Jaeger, C. H., Schwartz, E., and Firestone, M. K. (2006). Root Influence on
605    Nitrogen Mineralization and Nitrification in Avena barbata Rhizosphere Soil. *Soil Sci. Soc. Am. J.* 70,
606    1504–1511.

607 Hochman, A., Nissany, A., and Amizur, M. (1988). Nitrate reduction and assimilation by a moderately
608    halophilic, halotolerant bacterium Ba1. *Biochimica et Biophysica Acta (BBA) - General Subjects* 965,
609    82–89.

610 Huang, Y., Niu, B., Gao, Y., Fu, L., and Li, W. (2010). CD-HIT Suite: a web server for clustering and
611    comparing biological sequences. *Bioinformatics* 26, 680–682.

612 Huergo, L. F., Chandra, G., and Merrick, M. (2013). PIIsignal transduction proteins: nitrogen regulation
613    and beyond. *FEMS Microbiology Reviews* 37, 251–283. doi: 10.1111/j.1574-6976.2012.00351.x.

614 Hu, S., Chapin, F. S., Firestone, M. K., Field, C. B., and Chiariello, N. R. (2001). Nitrogen limitation of
615    microbial decomposition in a grassland under elevated CO_2. *Nature* 409, 188–191. doi:
616    10.1038/35051576.

617 Hyatt, D., Chen, G.-L., Locascio, P. F., Land, M. L., Larimer, F. W., and Hauser, L. J. (2010). Prodigal:
618    prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11,
619    119.

620 Jackson, L. E., Schimel, J. P., and Firestone, M. K. (1989a). Short-term partitioning of ammonium and
621    nitrate between plants and microbes in an annual grassland. *Soil Biology and Biochemistry* 21, 409–
622    415. doi: 10.1016/0038-0717(89)90152-1.

623 Jackson, L. E., Schimel, J. P., and Firestone, M. K. (1989b). Short-term partitioning of ammonium and
624    nitrate between plants and microbes in an annual grassland. *Soil Biol. Biochem.* 21, 409–415.

625 Jackson, L. E., Strauss, R. B., Firestone, M. K., and Bartolome, J. W. (1988). Plant and soil nitrogen
626    dynamics in California annual grassland. *Plant Soil* 110, 9–17.

Jung, M.-Y., Park, S.-J., Kim, S.-J., Kim, J.-G., Sinninghe Damsté, J. S., Jeon, C. O., et al. (2014). A
Mesophilic, Autotrophic, Ammonia-Oxidizing Archaeon of Thaumarchaeal Group I.1a Cultivated
from a Deep Oligotrophic Soil Horizon. *Applied and Environmental Microbiology* 80, 3645–3655.
doi: 10.1128/aem.03730-13.

Kanehisa, M., and Goto, S. (2000). KEGG: kyoto encyclopedia of genes and genomes. *Nucleic Acids Res.*
28, 27–30.

Katoh, K., and Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol. Biol. Evol.* 30, 772–780.

Koller, R., Scheu, S., Bonkowski, M., and Robin, C. (2013). Protozoa stimulate N uptake and growth of
arbuscular mycorrhizal plants. *Soil Biol. Biochem.* 65, 204–210.

Kozlov, A. M., Darriba, D., Flouri, T., Morel, B., and Stamatakis, A. (2019). RAxML-NG: a fast, scalable and
user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* 35, 4453–4455.

Kozlowski, J. A., Stieglmeier, M., Schleper, C., Klotz, M. G., and Stein, L. Y. (2016). Pathways and key
intermediates required for obligate aerobic ammonia-dependent chemolithotrophy in bacteria and
Thaumarchaeota. *ISME J.* 10, 1836–1845.

Kuzyakov, Y., Friedel, J. K., and Stahr, K. (2000). Review of mechanisms and quantification of priming
effects. *Soil Biol. Biochem.* 32, 1485–1498.

Leigh, J. A., and Dodsworth, J. A. (2007). Nitrogen regulation in bacteria and archaea. *Annu. Rev.*
Microbiol. 61, 349–377.

Leininger, S., Urich, T., Schloter, M., Schwark, L., Qi, J., Nicol, G. W., et al. (2006). Archaea predominate
among ammonia-oxidizing prokaryotes in soils. *Nature* 442, 806–809.

Letunic, I., and Bork, P. (2019). Interactive Tree Of Life (iTOL) v4: recent updates and new developments.
Nucleic Acids Res. 47, W256–W259.

Levičnik-Höfferle, S., Nicol, G. W., Ausec, L., Mandić-Mulec, I., and Prosser, J. I. (2012). Stimulation of
thaumarchaeal ammonia oxidation by ammonia derived from organic nitrogen but not added
inorganic nitrogen. *FEMS Microbiol. Ecol.* 80, 114–123.

Liao, Y., Smyth, G. K., and Shi, W. (2014). featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* 30, 923–930.

Liu, B., Mørkved, P. T., Frostegård, Å., and Bakken, L. R. (2010). Denitrification gene pools, transcription
and kinetics of NO, N2O and N2 production as affected by soil pH. *FEMS Microbiology Ecology* 72,
407–417. doi: 10.1111/j.1574-6941.2010.00856.x.

Love, M. I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol.* 15, 550.

MacNeil, T., Roberts, G. P., MacNeil, D., and Tyler, B. (1982). The products of glnL and glnG are
bifunctional regulatory proteins. *Mol. Gen. Genet.* 188, 325–333.

Matheus Carnevali, P. B., Schulz, F., Castelle, C. J., Kantor, R. S., Shih, P. M., Sharon, I., et al. (2019).
Hydrogen-based metabolism as an ancestral trait in lineages sibling to the Cyanobacteria. *Nat.*
Commun. 10, 463.

Melo-Oliveira, R., Oliveira, I. C., and Coruzzi, G. M. (1996). Arabidopsis mutant analysis and gene
regulation define a nonredundant role for glutamate dehydrogenase in nitrogen assimilation. *Proc.*
Natl. Acad. Sci. U. S. A. 93, 4718–4723.

Moreau, D., Bardgett, R. D., Finlay, R. D., Jones, D. L., and Philippot, L. (2019). A plant perspective on
nitrogen cycling in the rhizosphere. *Funct. Ecol.* 33, 540–552.

Moreau, D., Pivato, B., Bru, D., Busset, H., Deau, F., Faivre, C., et al. (2015). Plant traits related to
nitrogen uptake influence plant-microbe competition. *Ecology* 96, 2300–2310.

Myrold, D. D. (2021). "15 - Transformations of nitrogen," in *Principles and Applications of Soil*
Microbiology (Third Edition), eds. T. J. Gentry, J. J. Fuhrmann, and D. A. Zuberer (Elsevier), 385–421.

Nardi, P., Laanbroek, H. J., Nicol, G. W., Renella, G., Cardinale, M., Pietramellara, G., et al. (2020).
Biological nitrification inhibition in the rhizosphere: determining interactions and impact on
microbially mediated processes and potential applications. *FEMS Microbiol. Rev.* 44, 874–908.

Nguyen, T. T. H., Myrold, D. D., and Mueller, R. S. (2019). Distributions of Extracellular Peptidases Across
Prokaryotic Genomes Reflect Phylogeny and Habitat. *Front. Microbiol.* 10, 413.

Nicol, G. W., Leininger, S., Schleper, C., and Prosser, J. I. (2008). The influence of soil pH on the diversity,
abundance and transcriptional activity of ammonia oxidizing archaea and bacteria. *Environ.*
Microbiol. 10, 2966–2978.

Nuccio, E. E., Starr, E., Karaoz, U., Brodie, E. L., Zhou, J., Tringe, S. G., et al. (2020). Niche differentiation is
spatially and temporally regulated in the rhizosphere. *ISME J.* doi: 10.1038/s41396-019-0582-x.

Offre, P., Prosser, J. I., and Nicol, G. W. (2009). Growth of ammonia-oxidizing archaea in soil microcosms
is inhibited by acetylene. *FEMS Microbiol. Ecol.* 70, 99–108.

Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Simpson, G. L., Solymos, P., et al. (2008). The vegan
package. *Community ecology package. [http://r-forge. r-project. org/projects/vegan/]*. Available at:
https://www.researchgate.net/profile/Gavin_Simpson/publication/228339454_The_vegan_Packag
e/links/0912f50be86bc29a7f000000/The-vegan-Package.pdf.

Orellana, L. H., Hatt, J. K., Iyer, R., Chourey, K., Hettich, R. L., Spain, J. C., et al. (2019). Comparing DNA,
RNA and protein levels for measuring microbial dynamics in soil microcosms amended with
nitrogen fertilizer. *Sci. Rep.* 9, 17630.

O'Sullivan, C. A., Whisson, K., Treble, K., Roper, M. M., Micin, S. F., and Ward, P. R. (2017). Biological
nitrification inhibition by weeds: wild radish, brome grass, wild oats and annual ryegrass decrease
nitrification rates in their rhizospheres. *Crop and Pasture Science* 68, 798. doi: 10.1071/cp17243.

Pepe-Ranney, C., Campbell, A. N., Koechli, C. N., Berthrong, S., and Buckley, D. H. (2016). Unearthing the
Ecology of Soil Microorganisms Using a High Resolution DNA-SIP Approach to Explore Cellulose and
Xylose Metabolism in Soil. *Front. Microbiol.* 7, 703.

Personeni, E., and Loiseau, P. (2005). Species strategy and N fluxes in grassland soil. *European Journal of*
Agronomy 22, 217–229. doi: 10.1016/j.eja.2004.02.007.

Pett-Ridge, J., Shi, S., Estera-Molina, K., Nuccio, E., Yuan, M., Rijkers, R., et al. (2021). "Rhizosphere
Carbon Turnover from Cradle to Grave: The Role of Microbe–Plant Interactions," in *Rhizosphere*
Biology: Interactions Between Microbes and Plants, eds. V. V. S. R. Gupta and A. K. Sharma
(Singapore: Springer Singapore), 51–73.

Placella, S. A., and Firestone, M. K. (2013). Transcriptional response of nitrifying communities to wetting
of dry soil. *Appl. Environ. Microbiol.* 79, 3294–3302.

Pratscher, J., Dumont, M. G., and Conrad, R. (2011). Ammonia oxidation coupled to CO2 fixation by
archaea and bacteria in an agricultural soil. *Proc. Natl. Acad. Sci. U. S. A.* 108, 4170–4175.

Prosser, J. I., Hink, L., Gubry-Rangin, C., and Nicol, G. W. (2020). Nitrous oxide production by ammonia
oxidizers: Physiological diversity, niche differentiation and potential mitigation strategies. *Glob.*
Chang. Biol. 26, 103–118.

Rawlings, N. D., Barrett, A. J., and Finn, R. (2016). Twenty years of the MEROPS database of proteolytic
enzymes, their substrates and inhibitors. *Nucleic Acids Res.* 44, D343–50.

Reji, L., Tolar, B. B., Smith, J. M., Chavez, F. P., and Francis, C. A. (2019). Depth distributions of nitrite
reductase (nirK) gene variants reveal spatial dynamics of thaumarchaeal ecotype populations in
coastal Monterey Bay. *Environ. Microbiol.* 21, 4032–4045.

Sakamoto, N., Kotre, A. M., and Savageau, M. A. (1975). Glutamate dehydrogenase from Escherichia coli:
purification and properties. *J. Bacteriol.* 124, 775–783.

Schimel, J. P., Jackson, L. E., and Firestone, M. K. (1989). Spatial and temporal effects on plant-microbial
competition for inorganic nitrogen in a california annual grassland. *Soil Biol. Biochem.* 21, 1059–
1066.

Shi, S., Herman, D. J., He, Z., Pett-Ridge, J., Wu, L., Zhou, J., et al. (2018). Plant roots alter microbial
functional genes supporting root litter decomposition. *Soil Biol. Biochem.* 127, 90–99.

Sieradzki, E. T., Nuccio, E. E., Pett-Ridge, J., and Firestone, M. K. (2020). Transcription of protease and
chitinase genes provides a window onto macromolecular organic nitrogen decomposition in soil.
bioRxiv, 2020.12.14.422732. doi: 10.1101/2020.12.14.422732.

Sokol, N. W., Slessarev, E., Marschmann, G. L., Nicolas, A., Blazewicz, S. J., Brodie, E. L., et al. (2022). Life
and death in the soil microbiome: how ecological processes influence biogeochemistry. *Nat. Rev.*
Microbiol. 20, 415–430.

Starr, E. P., Shi, S., Blazewicz, S. J., Probst, A. J., Herman, D. J., Firestone, M. K., et al. (2018). Stable
isotope informed genome-resolved metagenomics reveals that Saccharibacteria utilize microbially-
processed plant-derived carbon. *Microbiome* 6, 122.

Stein, L. Y. (2019). Insights into the physiology of ammonia-oxidizing microorganisms. *Curr. Opin. Chem.*
Biol. 49, 9–15.

Subbarao, G. V., Rondon, M., Ito, O., Ishikawa, T., Rao, I. M., Nakahara, K., et al. (2007). Biological
nitrification inhibition (BNI)—is it a widespread phenomenon? *Plant and Soil* 294, 5–18. doi:
10.1007/s11104-006-9159-3.

Subbarao, G. V., Sahrawat, K. L., Nakahara, K., Rao, I. M., Ishitani, M., Hash, C. T., et al. (2013). A
paradigm shift towards low-nitrifying production systems: the role of biological nitrification
inhibition (BNI). *Ann. Bot.* 112, 297–316.

Sudderth, E. A., St. Clair, S. B., Placella, S. A., Swarbreck, S. M., Castanha, C., Herman, D. J., et al. (2012).
Annual grassland resource pools and fluxes: sensitivity to precipitation and dry periods on two
contrasting soils. *Ecosphere* 3, art70–art70.

Tolar, B. B., Herrmann, J., Bargar, J. R., van den Bedem, H., Wakatsuki, S., and Francis, C. A. (2017).
Integrated structural biology and molecular ecology of N-cycling enzymes from ammonia-oxidizing
archaea. *Environ. Microbiol. Rep.* 9, 484–491.

Tosi, M., Brown, S., Ferrari Machado, P. V., Wagner-Riddle, C., and Dunfield, K. (2020). Short-term
response of soil N-cycling genes and transcripts to fertilization with nitrification and urease
inhibitors, and relationship with field-scale N2O emissions. *Soil Biol. Biochem.* 142, 107703.

Tourna, M., Freitag, T. E., Nicol, G. W., and Prosser, J. I. (2008). Growth, activity and temperature
responses of ammonia-oxidizing archaea and bacteria in soil microcosms. *Environmental
Microbiology* 10, 1357–1364. doi: 10.1111/j.1462-2920.2007.01563.x.

Trofymow, J. A., Coleman, D. C., and Cambardella, C. (1987). Rates of rhizodeposition and ammonium
depletion in the rhizosphere of axenic oat roots. *Plant and Soil* 97, 333–344. doi:
10.1007/bf02383223.

Wang, Q., Garrity, G. M., Tiedje, J. M., and Cole, J. R. (2007). Naive Bayesian classifier for rapid
assignment of rRNA sequences into the new bacterial taxonomy. *Appl. Environ. Microbiol.* 73,
5261–5267.

Warnes, G. R., Bolker, B., Bonebakker, L., Gentleman, R., Liaw, W. H. A., Lumley, T., et al. (2015). gplots:
Various R programming tools for plotting data. Available at:
https://www.scienceopen.com/document?vid=0e5d8e31-1fe4-492f-a3d8-8cd71b2b8ad9.

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer.

Yergeau, E., Sanschagrin, S., Maynard, C., St-Arnaud, M., and Greer, C. W. (2014). Microbial expression
profiles in the rhizosphere of willows depend on soil contamination. *ISME J.* 8, 344–358.

Yergeau, E., Tremblay, J., Joly, S., Labrecque, M., Maynard, C., Pitre, F. E., et al. (2018). Soil
contamination alters the willow root and rhizosphere metatranscriptome and the root-rhizosphere
interactome. *ISME J.* 12, 869–884.

Yuan, J., Doucette, C. D., Fowler, W. U., Feng, X.-J., Piazza, M., Rabitz, H. A., et al. (2009). Metabolomics-
driven quantitative analysis of ammonia assimilation in E. coli. *Mol. Syst. Biol.* 5, 302.

Zeglin, L. H., Taylor, A. E., Myrold, D. D., and Bottomley, P. J. (2011). Bacterial and archaeal amoA gene
distribution covaries with soil nitrification properties across a range of land uses. *Environ.*

Microbiol. Rep. 3, 717–726.

Zhalnina, K., Louie, K. B., Hao, Z., Mansoori, N., da Rocha, U. N., Shi, S., et al. (2018). Dynamic root
exudate chemistry and microbial substrate preferences drive patterns in rhizosphere microbial
community assembly. *Nat Microbiol* 3, 470–480.

Zhang, L.-M., Offre, P. R., He, J.-Z., Verhamme, D. T., Nicol, G. W., and Prosser, J. I. (2010). Autotrophic
ammonia oxidation by soil thaumarchaea. *Proc. Natl. Acad. Sci. U. S. A.* 107, 17240–17245.

Zhu, B., Gutknecht, J. L. M., Herman, D. J., Keck, D. C., Firestone, M. K., and Cheng, W. (2014).
Rhizosphere priming effects on soil carbon and nitrogen mineralization. *Soil Biol. Biochem.* 76, 183–
192.

Žifčáková, L., Větrovský, T., Howe, A., and Baldrian, P. (2016). Microbial activity in forest soil reflects the
changes in ecosystem properties between summer and winter. *Environ. Microbiol.* 18, 288–301.