Predicting future conservation areas while avoiding competition in two alpine urodele amphibians

severely threatened by climate change

Headings: Accounting for competition in Alpine salamander SDM

Nicolas Dubos[1, 2], Antoine Havard[1], Angelica Crottini[1, 2, 3], Daniele Seglie[4], Franco Andreone[5]

[1] CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, *InBIO* Laboratório Associado,

Campus de Vairão, Universidade do Porto, 4485-661 Vairão, Portugal

[2] BIOPOLIS Program in Genomics, Biodiversity and Land Planning, CIBIO, Campus de Vairão, 4485-661

Vairão, Portugal

[3] Departamento de Biologia, Faculdade de Ciências, Universidade do Porto, rua do Campo Alegre s/n, 4169–

007 Porto, Portugal

[4] ELEADE Società Cooperativa, 10010 Chiaverano (TO), Italy

[5] Museo Regionale di Scienze Naturali, Via G. Giolitti, 36, I-10123 Torino, Italy

Abstract

Climate change will cause important declines in species distributions, especially when living at high

altitudes. The Critically Endangered *Salamandra lanzai* from SW Alps may be severely exposed to future

climate change effects and its suitable climate may shrink or shift. Another Alpine salamander (*S. atra*) is

present in the region, which in case of spatial overlap may represent a competitor for *S. lanzai*. It is urgent to

estimate the effect of future climate change on these species and identify priority areas for conservation

while accounting for competition between both species. With a Species Distribution Modelling (SDM)

approach, we projected the current and future climate suitability of both salamander species. We accounted

for uncertainty related to the methods (model replicates) and climate projections (data source, global

circulation model and scenario) to provide a consensus map for practitioners. This map also takes into

account potential competition with *S. atra* by penalizing the suitability scores of S. *lanzai* by the scores of *S.*

atra. We predict a severe effect of climate change on both species. Most of the current habitats are projected

to become largely unsuitable by 2070, regardless of the climatology and scenario. We identified important

spatial disagreements between projections based on different data sources, mostly due to precipitation

projections and daily temperature variation. This highlights the need to account for multiple climatologies in

mountainous environments. Both species' habitats are highly fragmented, which is expected to prevent

distributional shifts through natural dispersion. We suggest to explore the possibility of translocation for the

most threatened populations and simultaneously develop captive breeding programs. Biotic interactions are

rarely accounted for in SDMs, and we encourage the documentation of species with similar ecological

requirements to improve the relevance of SDMs for future conservation planning.

Keywords: Biotic interactions, Climate change, Mountains, Priority areas, Species distribution models,

Salamandra lanzai.

Introduction

Climate change is one of the main threats to montane species, especially amphibians (Raxworthy et al., 2008; Mccain & Colwell, 2011). Warming temperatures will likely cause upslope shifts in species distributions, potentially leading to extinction (Freeman & Class Freeman, 2014; Rana et al., 2021). The effect of climate change varies along latitude (Tewksbury et al. 2008; Dubos et al. 2019), which particularly applies to montane species, with temperate species less likely to respond negatively than tropical ones (Freeman et al. 2021). Upslope shifts may not always reduce the amount of suitable area in some systems but also bring newly available regions, thus increasing the availability of suitable habitats (Elsen & Tingley, 2015). Hence there is a need to compile climate change predictions for mountainous environments and quantify the amount of suitable environments in the future.

Species distribution models (SDMs) can be used to identify the most suitable conditions for a given species, and forecast potential future geographical shifts in these conditions. SDMs are commonly used for testing ecological hypotheses (Anderson et al., 2009; Dubos, Augros, et al., 2022), assessing alien species invasion risks (Bellard et al., 2013; Dubos, Fieldsend, et al., 2022; Lanner et al., 2022), forecasting the potential effect of environmental change (Araújo et al., 2005), and supporting conservation and management efforts (Schwartz, 2012; Leroy et al., 2014; Dubos, Lenormand, et al., 2022). In conservation application, SDMs can provide maps of projected future environmental suitability at the resolution of environmental data. These maps can be used as a tool for practitioners to seek suitable habitats or microhabitats within the identified suitable pixels and propose priority areas for conservation actions (e.g. for the identification of potential translocation sites, or for protection status).

Biotic interactions are rarely accounted for in SDMs because they are highly demanding in term of data and analysis. For a given species, suitable climatic conditions may shift with climate change, and these conditions may overlap with that of another species sharing similar ecological requirements. The sites identified as suitable in the future may represent candidate sites for conservation action for both species, but would expose them to competition. Hence, there is a need to identify the best candidate sites for either species while mutually excluding them.

We studied the effect of climate change on the potential distribution of two alpine salamanders with similar ecological adaptations, Lanza's salamander (*Salamandra lanzai*) and alpine salamander (*S. atra*). Both these species are typical of montane environments and evolved an aplacental viviparity (with full embryonic

development occurring within their body), that allowed them to complete their reproductive cycle

independently from the occurrence of water bodies. This is likely an adaptation to cold climate at high

temperatures (Ma et al., 2018). In both species, several populations are highly isolated due to both their past

biogeographic histories and more recent anthropogenic landscape modification (Tessa, Crottini, & Andreone,

2007). Current patchy distribution suggests that individuals from these isolated populations will likely not be

able to follow the shifting climatic conditions through dispersal (Andreone et al. 2004). In this study we

identified the climatic predictors explaining the current distribution of *S. lanzai* and *S. atra* using an SDM

approach. We expect a strong effect of climate change on both species, especially for *S. lanzai* due to its

narrow distribution (Platts et al., 2014; Dubos, et al., 2022). Due to its conservation status (Critically

Endangered; IUCN SSC Amphibian Specialist Group, 2022), we focus on *S. lanzai* for conservation

considerations. We projected both species climate suitability on current and future (2070) conditions in the

Alps in order to (1) quantify the effect of climate change on both species' distributions and (2) identify

priority areas for the conservation of *S. lanzai* while accounting for competition.

Methods

Occurrence data

We obtained 1991 and 841 unique locality records for *S. lanzai* and *S. atra*, respectively. These have been

retrieved from iNaturalist (https://www.inaturalist.org/) and Parc Naturel Régional du Queyras

(https://www.pnr-queyras.fr/). To avoid pseudo-replication, we resampled one locality record per pixel of

climate data (i.e. data thinning; Steen et al. 2021). At the resolution of 1×1 km, this led to a final sample of

n = 104 or n = 102 presence points for *S. lanzai* after resampling on CHELSA data, or with Worldclim,

respectively. The final sample for *S. atra* was n = 711 presence points based on CHELSA (710 with

Worldclim).

Climate data

We used six uncorrelated bioclimatic variables (Pearson's r < 0.7; Dormann et al. 2013) for which we had

hypotheses of causal relationship with both species' biology (Fourcade, Besnard, & Secondi, 2018; Dubos,

Préau, et al., 2022) at 30 arc seconds resolution (approximately 1 km) for the current and future (2070)

climate from two sources: CHELSA (Karger et al., 2017) and Worldclim global climate data (Fick &

Hijmans, 2017). These data sources used different methods to compute the climatologies. Worldclim is based

on interpolated data with elevation and distance to the coast as predictors in addition to satellite data, while

CHELSA is based on statistical downscaling for temperature, and precipitation estimations incorporating

orographic factors (i.e., wind fields, valley exposition, boundary layer height). We describe the causality of

the relationship between predictors and species' ecology in Table 1. The six variables retained were annual

mean temperature (Bio1, which was collinear with summer and winter mean temperatures), autumn mean

temperature (Bio8), isothermality (Bio3), temperature seasonality (Bio4), precipitation of the driest month

(Bio14), and summer precipitation (Bio18). We used a statistical process to select the most relevant

predictors (elbow criterion for the relative importance, as measured by the randomForest R package;

Breiman 2001). We did not include precipitation seasonality (Bio15) despite this variable is often selected in

mountainous areas (e.g. De Marchi et al. 2020), because its causal relationship with the biology of alpine

salamanders is not clear. To consider the uncertainty related to future climate projections, we used three

Global Circulation Models (GCMs; i.e., BCC-CSM1-1, MIROC5, and HadGEM2-AO) and two greenhouse

gas emission scenarios (the most optimistic RCP26 and the most pessimistic RCP85).

Table 1 Selected variables and ecological hypotheses of causality for the presence of *Salamandra*

lanzai and *S. atra* in the Alps. We give the relative importance of each variable as measured by the Random

Forest algorithm (mean decrease in Gini index). Predictors with the highest relative importance were

included in the models (in bold for each species, selected on the basis of elbow criterion).

Bioclimatic variable	Relative importance		Ecological interpretation	Reference
	S. lanzai	*S. atra*		
Bio1 (annual mean T)	**Worldclim: 0.24** **CHELSA: 0.23**	**Worldclim: 138** **CHELSA: 134**	Summer activity	Ribéron and Miaud 1999; Kissel et al. 2019
Bio3 (isothermality)	**Worldclim: 0.15** **CHELSA: 0.16**	Worldclim: 105; CHELSA: 78	Higher metabolic cost in very cold nights	Authors' hypothesis
Bio4 (T seasonality)	**Worldclim: 0.07** **CHELSA: 0.12**	**Worldclim: 124** **CHELSA: 134**	Higher metabolic cost in highly variable years, driven by hot winters	Kissel et al. 2019
Bio18 (P of warmest quarter)	Worldclim: 0.12 CHELSA: 0.05	**Worldclim: 106** **CHELSA: 117**	Summer activity, food availability, larval dessication risk	Ribéron and Miaud 1999; Kissel et al. 2019; (Romano et al. 2021)
Bio19 (P of coldest quarter)	**Worldclim: 0.24** **CHELSA: 0.26**	**Worldclim: 96** **CHELSA: 108**	Overall habitat moisture	Miaud et al. 2001; Andreone et al. 2004; Romano et al. 2021

Land-use data

We accounted for habitat requirements by filtering all unsuitable habitats from our model projections. We used a filter instead of integrating habitat predictors to our models because (1) the species are strongly dependent on given habitat types (hence, removing unsuitable habitats is realistic), (2) this improves statistical parsimony and (3) final projections benefit from finer resolution (climate projections were resampled to fit land-use data). We obtained high-resolution land cover categories from Corine Land Cover 2018 (100 m resolution) derived from remote sensing (Copernicus Service Information, 2022). We assume agricultural and urbanized areas to be unsuitable to the presence of salamanders (Andreone et al., 2004; Geiger, 2006; Reinthaler-Lottermoser et al., 2010; Romano et al., 2018). These habitat types were removed from final projections, only to keep habitats that are used by both species, i.e. coniferous forests, natural open areas and bare rock (Andreone, De Michelis, & Clima, 1999; Ribéron & Miaud, 1999; Miaud et al., 2001; Bonato & Fracasso, 2003, 2014; Andreone et al., 2004).

Ecological niche modelling

We modelled and projected climatic niches using highly performing single tuned algorithm (Valavi et al. 2021), i.e. Random Forest down-sampled (hereafter RF down-sampled, i.e. RF parametrised to deal with a large number of background samples and few presence records; Prasad et al. 2006). We set RF down-sampled to run for 1000 bootstrap samples/trees, with 5 cross validation runs and 5 sets of 10,000 pseudo-absences. Since data were strongly spatially aggregated, we did not use a spatial partitioning for cross validation runs (Valavi et al. 2019), but a random partitioning instead. We applied a sample bias correction technique consisting in a selection of pseudo-absence points that imitates the spatial bias of presence points (Phillips et al. 2009). To do so, we used a null geographic model, i.e. a map of the density of unthinned presence points, which we use as a probability weight for pseudo-absence selection.

We determined the background extent following a hypothesis of past dispersal, accounting for the Equilibrium hypothesis, and minimizing the surface to better account for sample bias (Vollering et al., 2019). We assume both alpine salamanders were able to disperse in the whole Alps in the past, hence both backgrounds encompassed the Alps and their surroundings. To account for Equilibrium, we mutually excluded the regions where the putative competitor is present, i.e. the Central and Eastern Alps for *S. lanzai*

and South-Western Alps for *S. atra*. The fire salamander *Salamandra salamandra* is present at the foot of the

Alps. However, we did not exclude the non-mountainous areas at the foot of the Alps because *S. atra* and *S.*

salamandra appear not to be in competition, or competition does not lead to spatial segregation (Werner,

Lötters, & Schmidt, 2014).

We evaluated model performance using the Boyce Index (Hirzel et al., 2006). An index of 1 suggests that

suitability models were perfectly able to predict presence points, a null value suggests that models are no

better than random and a negative value implies a counter prediction. All models for which Boyce index <

0.5 were discarded. We present projections of the mean suitability obtained across all highly performing

models.

Conservation application: Accounting for uncertainty and competition

To provide sound conservation guidelines, we built a consensus map of the mean predictions across all

highly performing models, discounting the standard deviation (mean – SD; Kujala et al. 2013; Dubos et al.

2022a). Final 'consensus' maps represent the areas that were identified as suitable with the lowest

disagreement between model replicates, hence the best candidate sites for conservation action.

Priority areas for the conservation of *S. lanzai* may fall into areas occupied by *S. atra* in Central and Eastern

Alps, which may expose both species to competition and reduce success probability for possible

conservation efforts (e.g. translocation). To avoid potential sympatry and hence, competition between *S.*

lanzai and *S. atra*, we excluded areas that are suitable to *S. atra* from the consensus map of *S. lanzai*.

Similarly, we subtracted the consensus suitability of *S. atra* to the consensus suitability of *S. lanzai*, with the

obtained result to be interpreted as a priority map for conservation efforts for *S. lanzai* in Central and Eastern

Alps.

Results

Ecological niche models

We selected four climatic predictors for either species. For *S. lanzai*, we retained bio1, bio3, bio4 and bio19.

For *S. atra*, we kept bio1, bio4, bio18 and bio19 (Fig. S1—S4). For *S. lanzai*, models performed well (i.e.

suitability scores are high at presence points), with an average Boyce index of 0.67 (CHELSA; 5 models

discarded) and 0.84 (Worldclim; no model discarded) for each climate data, respectively. For *S. atra*, model

performance was very high (CHELSA: mean Boyce = 0.98; Worldclim: mean Boyce = 0.99; no model

discarded).

For *S. lanzai*, current predictions identified different areas with suitable conditions depending on the source

of climate data. With Worldclim, suitable conditions can be found further south of its current distribution

(Fig. 1). On the contrary with CHELSA, suitable conditions extend to the north of its current distribution

(Fig. 2). For *S. atra*, suitable conditions are consistently found across the northern and eastern part of the

Alps with both climate data sources (Fig. 3, 4).

Fig. 1 Environmental suitability for *Salamandra lanzai* based on Worldclim climatologies. Left: current;

Right: future (2070). Future projections include two emission scenarios (SSP126 and SSP585) and three

GCMs.

Fig. 2 Environmental suitability for *Salamandra lanzai* based on CHELSA climatologies. Left, current;

Right, future (2070). Future projections include two emission scenarios (SSP126 and SSP585) and three

GCMs.

Fig. 3 Environmental suitability for *Salamandra atra* based on Worldclim climatologies. Left, current; Right, future (2070). Future projections include two emission scenarios (SSP126 and SSP585) and three GCMs.

Fig. 4 Environmental suitability for *Salamandra atra* based on CHELSA climatologies. Left, current; Right, future (2070). Future projections include two emission scenarios (SSP126 and SSP585) and three GCMs.

Future climate suitability

At known populations (i.e. presence points), climate suitability will decline for both species according to

both climate data sources, and for all GCMs and scenarios. The predicted decline is generally more severe

when based on Worldclim data as compared to CHELSA (Fig. 5).

For *S. lanzai*, models based on Worldclim predicted that the most suitable areas will be located in the eastern

Alps. According to CHELSA, the most suitable areas will be located at the current distribution and in the

northern part of central Alps. For *S. atra*, the most suitable areas will remain similar to the current ones, but

with suboptimal conditions throughout the Alps.

Accounting for uncertainty and competition

When taking into account the uncertainty related to model replicates and climate data (source, GCMs and

scenarios), the most suitable areas for *S. lanzai* will be located in central Alps / eastern Switzerland, between

the Swiss National Park and the Park Ela (Fig. 6). For *S. atra*, the most suitable areas also will be in central

Alps, although further west to *S. lanzai*'s projected suitable area (i.e. around the Beverin Nature Park).

When accounting for potential competition with *S. atra*, the most suitable areas by 2070 will be located in

the Italian Alps, in the Parco Nazionale dello Stelvio and the Parco delle Orobie Bergamasche. Our

projections did not include the area currently occupied by *S. lanzai* because it is also projected to be suitable

to *S. atra*. However, it is unlikely that *S. atra* would reach this region through natural dispersion. As such,

the current distribution of *S. lanzai* may still be included in the priority areas for the conservation of this

species.

Fig. 5 Current and future (2070) environmental suitability for two alpine salamanders (Top: *Salamandra*

lanzai; Bottom: *S. atra*) according to two climate data sources (Worldclim and CHELSA) and a range of

scenarios and GCMs.

Fig. 6 Consensus predictions for current climate suitability of *Salamandra lanzai* and *S. atra* (mean

prediction across all models penalised by uncertainty), with the most suitable areas for *S. lanzai* excluding

suitable areas for *S. atra*.

Discussion

We projected the future climate suitability of two Alpine salamanders to inform possible conservation guidelines which takes into account climate projection uncertainty and competition. Our results suggest a severe effect of climate change on both species. We provide a single conservation priority map as a practical tool for practitioners.

Ecological niche models

Salamandra lanzai is found in cold areas. We were not able to identify a minimum temperature for activity, presumably because our presence data encompassed the coldest pixels of the selected background. Suitable areas are characterized by low temperature variation throughout the year and the day. Hot or short winters may increase metabolic costs and affect adult survival (Kissel et al., 2019), while extremely cold nights may affect larval and adult survival (Madison, 1997). Paradoxically, hot winters may expose salamanders to extreme cold because of lower snow cover in winter. Indeed, salamanders often use mammal shelters (including marmot and other rodents burrows; Ribéron and Miaud 1999) and consequently may suffer from thinner snow cover (due to less effective thermal isolation) as temperature increases (Tafani et al. 2013). According to Worldclim projections, *S. lanzai* occupies the wettest areas of the western Alps. In this case, winter precipitation explained better than summer precipitation the current distribution. This effect may constitute a proxy for the available amount of water stored in the soil for the rest of the year. Surprisingly, projections based on CHELSA identified areas that are among the driest of the study area. CHELSA takes into account orographic factors and wind exposure. *Salamandra lanzai* usually occupies microhabitats with little wind exposure, presumably because wind may exacerbate the effect of cold on salamanders, as well as increase the chances of desiccation through evapotranspiration. We interpret the effect of summer precipitation for CHELSA as a proxy for wind exposure, which may explain the shape of the relationship. Our data also enabled us to identify an optimal temperature window for *S. atra* (1–7°C annual mean) which is consistent with our expectations regarding activity. The wetter the habitat during summer, the more likely it is to be occupied by the species. Results are less consistent regarding winter precipitation, with a notable disagreement between Worldclim and CHELSA projections. Differences may be related to the different treatment of topography between both climatologies. We call for further investigation on the interpretation of precipitation predictors in the Alps.

On the use of 1 km-resolution climate data in mountains

The use of downscaled climate data is usually recommended in mountainous environments (Lembrechts et al., 2019). Recent studies of mountain-dwelling salamanders, including *S. atra*, used downscaled climatic predictors to better take into account the heterogeneity of mountain environments (Feldmeier et al., 2020; Stickley & Fraterrigo, 2023). Both studies found that accounting for micro-climate reduces the predicted impact of climate change on salamanders, notably due to buffering effects of forest vegetation and micro-climate availability in valleys non-exposed to the sun. In our case, both species live in forested areas (apart from the highest altitude populations of *S. lanzai* living in saxicolous habitats), suggesting that forest management might not be a relevant avenue for the conservation of alpine salamanders. The approach of Feldmeier et al. (2020) highlights the consideration of habitat management on north-exposed valleys for *S. atra*. Our results based CHELSA data—which take into account valley exposure (Karger et al., 2017)— enable us to examine the orientation of suitable valleys at 1 km-resolution, and fully support their conclusions since most of the future suitable areas are predicted in north-exposed valleys (Fig. 4). Nonetheless, we stress that our future projections suggest a strong decline in climate suitability for *S. atra* also in these areas (Fig. 5). Very high resolution climate data are rarely available at large scale, and future projections often implies statistical downscaling based on transposed and interpolated anomaly between coarse and fine scale data (e.g., Feldmeier et al., 2020; Lembrechts et al., 2019). Despite the use of 1 km resolution omits the presence of microhabitats, our future projections are still meaningful when considered relative to predictions of current suitability.

The future of Alpine salamanders

We predict an important decline in climate suitability for both species, with only suboptimal climates left available by 2070. Winter temperature rising may increase metabolic costs, which would affect survival rates. Warmer summers likely reduce activity for *S. atra* (Geiger 2006), which supports a lower foraging efficiency, reproduction and overall survival. Precipitations are projected to decrease, which may delay age at maturity, reduce body length and growth rate, and increase gestation duration (Miaud et al. 2001; Andreone et al. 2004). According to the source of climate data, *S. lanzai* may either persist within its current distribution area or go fully extinct. We recommend to maintain or improve conservation efforts within the current range, and consider translocation to the identified areas of central Alps as a possible avenue to be

considered in the future for the most threatened populations. The results and efficiency of translocation

action is variable (Seddon, Strauss, & Innes, 2012; Thévenin, 2019), and since both *S. lanzai* and *S. atra* are

aplacental viviparous species, these should be carefully evaluated and their cost and benefits assessed. While,

viviparity would assure a better isolation of foetuses within mother's body, it is evident that the evolution of

this K-oriented breeding strategy require habitat stability (i.e. caused either by climate warming or by

translocation). In parallel to these evaluations we encourage to consider developing captive breeding and

gametes cryopreservation as possible actions for conservation.

Concluding remarks

This study showed the importance of considering multiple climatologies when projecting climate change

effects on species distributions in a mountainous environment. We provide a high-resolution map accounting

for uncertainty related to climate data, habitat requirements and biotic interactions (competition), which is

here made available to practitioners to support the decisions-making process for the conservation actions

targeting a Critically Endangered species potentially on the brink of extinction. We encourage the

documentation of species with similar ecological requirements to improve the consideration of competition

in species distribution models.

Acknowledgements

We thank everyone that contributed to the data collection in the field. Portuguese National Funds through

FCT (Fundação para a Ciência e a Tecnologia) support the research contract

(2020.00823.CEECIND/CP1601/CT0003) and research project (PTDC/BIA-EVL/31254/2017) to Angelica

Crottini and the research contract to Nicolas Dubos [ICETA_2021_26]. We are grateful to Lucio Bonato,

Gentile Francesco Ficetola, Claude Miaud and Benedikt R. Schmidt for useful discussions. We thank Matteo

Quartesan for data provision and useful exchanges.

Author contributions

ND led the writing of the manuscript, designed and conducted the analyses. AH contributed to data analysis

and writing. DS and FA have made substantial contribution to acquisition of the data. AC and FA have made

a substantial contribution to data interpretation, conception of the study and manuscript revision. All authors

have approved the final version for submission, have agreed to be accountable for the accuracy and integrity

of the work that they conducted, have agreed to resolve queries relating to the work that they conducted and

contributed to the study.

References

Anderson, B. J., Arroyo, B. E., Collingham, Y. C., Etheridge, B., Fernandez-De-Simon, J., Gillings, S., Gregory, R. D., Leckie, F. M., Sim, I. M. W., Thomas, C. D., Travis, J., & Redpath, S. M. (2009). Using distribution models to test alternative hypotheses about a species' environmental limits and recovery prospects. *Biol. Conserv.* **142**, 488–499.

Andreone, F., De Michelis, S., & Clima, V. (1999). A montane amphibian and its feeding habits: Salamandra lanzai (caudata, salamandridae) in the alps of northwestern Italy. *Ital. J. Zool.* **66**, 45–49.

Andreone, F., Miaud, C., Bergò, P. E., Doglio, S., Stocco, P., Riberon, A., & Gautier, P. (2004). Living at high altitude: Testing the effects of life history traits upon the conservation of salamandra lanzai (amphibia, salamandridae). *Ital. J. Zool.* **71**, 35–43.

Araújo, M. B., Pearson, R. G., Thuiller, W., & Erhard, M. (2005). Validation of species–climate impact models under climate change. *Glob. Chang. Biol.* **11**, 1504–1513.

Bellard, C., Thuiller, W., Leroy, B., Genovesi, P., Bakkenes, M., & Courchamp, F. (2013). Will climate change promote future invasions? *Glob. Chang. Biol.* **19**, 3740–3748.

Bonato, L., & Fracasso, G. (2003). Movements, distribution pattern and density in a population of Salamandra atra aurorae (Caudata: Salamandridae). *Amphib. Reptil.* **24**, 251–260.

Bonato, L., & Fracasso, G. (2014). Epigean habitat of a population of Salamandra atra aurorae: a preliminary analysis. *Atti X Congr. Naz. Soc. Herpetol. Ital.* 47–55.

Breiman, L. (2001). Random forests. *Ensemble Mach. Learn. Methods Appl.* **45**, 5–32.

Copernicus Service Information. (2022). No Title. *Available at http://land.copernicus.eu.*

De Marchi, G., Bombieri, G., Boz, B., Leandri, F., & Richard, J. (2020). Has the west been won? A field survey and a species distribution model of iberolacerta horvathi in the alps. *Acta Herpetol.* **15**, 3–14.

Dubos, N., Augros, S., Deso, G., Probst, J., Notter, J., & Roesch, M. (2022). Here be dragons: important spatial uncertainty driven by climate data in forecasted distribution of an endangered insular reptile. *Anim. Conserv.*

Dubos, N., Dehorter, O., Yves, P., & Viol, I. Le. (2019). Thermal constraints on body size depend on the population's position within the species' thermal range in temperate songbirds. *Glob. Ecol. Biogeogr.* **28**, 96–106.

Dubos, N., Fieldsend, T. W., Roesch, M. A., Stephane, A., Aurélien, B., Arthur, C., Ineich, I., Kenneth, K., Boris, L., Malone, S. L., Probst, J., Raxworthy, C. J., & Crottini, A. (2022). Choice of climate data influences current and future global invasion risks for two Phelsuma geckos. *BioRxiv.*

Dubos, N., Lenormand, M., Castello, L., Oberdorff, T., & Guisan, A. (2022). Protection gaps in Amazon floodplains will increase with climate change: insight from the world's largest scaled freshwater fish. *Aquat. Conserv. Mar. Freshw. Ecosyst.* 1–29.

Dubos, N., Montfort, F., Grinand, C., Nourtier, M., Deso, G., Probst, J.-M., Razafimanahaka, J. H., Andriantsimanarilafy, R. R., Rakotondrasoa, E. F., Razafindraibe, P., Jenkins, R., & Crottini, A. (2022). Are narrow-ranging species doomed to extinction? Projected dramatic decline in future climate suitability of two highly threatened species. *Perspect. Ecol. Conserv.* **20**, 18–28.

Dubos, N., Préau, C., Lenormand, M., Papuga, G., Montsarrat, S., Denelle, P., Louarn, M. Le, Heremans, S., Roel, M., Roche, P., & Luque, S. (2022). Assessing the effect of sample bias correction in species distribution models. *Ecol. Indic.* **145**, 109487.

Elsen, P. R., & Tingley, M. W. (2015). Global mountain topography and the fate of montane species under climate change. *Nat. Clim. Chang.* **5**, 772–776.

Feldmeier, S., Schmidt, B. R., Zimmermann, N. E., Veith, M., Ficetola, G. F., & Lötters, S. (2020). Shifting aspect or elevation? The climate change response of ectotherms in a complex mountain topography. *Divers. Distrib.* **26**, 1483–1495.

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *Int. J. Climatol.* **37**, 4302–4315.

Fourcade, Y., Besnard, A. G., & Secondi, J. (2018). Paintings predict the distribution of species, or the challenge of selecting environmental predictors and evaluation statistics. *Glob. Ecol. Biogeogr.* **27**, 245–256.

Freeman, B. G., & Class Freeman, A. M. (2014). Rapid upslope shifts in New Guinean birds illustrate strong distributional responses of tropical montane species to global warming. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 4490–4494.

Freeman, B. G., Song, Y., Feeley, K. J., & Zhu, K. (2021). Montane species track rising temperatures better in the tropics than in the temperate zone. *Ecol. Lett.* **24**, 1697–1708.

346 Geiger, C. (2006). Ecological requirements of the Alpine Salamander Salamandra atra : assessing the effects
347   of current habitat structure and landscape dynamics on local distribution Table of contents.

348 Hirzel, A. H., Le Lay, G., Helfer, V., Randin, C., & Guisan, A. (2006). Evaluating the ability of habitat
349   suitability models to predict species presences. *Ecol. Modell.* **199**, 142–152.

350 IUCN SSC Amphibian Specialist Group. (2022). Salamandra lanzai. *IUCN Red List Threat. Species 2022*.

351 Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., Zimmermann, N. E., Linder,
352   H. P., & Kessler, M. (2017). Climatologies at high resolution for the earth's land surface areas. *Sci.*
353   *Data* **4**, 1–20.

354 Kissel, A. M., Palen, W. J., Ryan, M. E., & Adams, M. J. (2019). Compounding effects of climate change
355   reduce population viability of a montane amphibian. *Ecol. Appl.* **29**, 1–12.

356 Kujala, H., Moilanen, A., Araújo, M. B., & Cabeza, M. (2013). Conservation planning with uncertain climate
357   change projections. *PLoS One* **8**.

358 Lanner, J., Dubos, N., Geslin, B., Leroy, B., Hernández-Castellano, C., Dubaić, J. B., Bortolotti, L., Calafat,
359   J. D., Ćetković, A., Flaminio, S., Le Féon, V., Margalef-Marrase, J., Orr, M., Pachinger, B., Ruzzier, E.,
360   Smagghe, G., Tuerlings, T., Vereecken, N., & Meimberg, H. (2022). On the road: Anthropogenic factors
361   drive the invasion risk of a wild solitary bee species. *Sci. Total Environ.* **827**, 154246.

362 Lembrechts, J. J., Lenoir, J., Roth, N., Hattab, T., Milbau, A., Haider, S., Pellissier, L., Pauchard, A., Ratier
363   Backes, A., Dimarco, R. D., Nuñez, M. A., Aalto, J., & Nijs, I. (2019). Comparing temperature data
364   sources for use in species distribution models: From in-situ logging to remote sensing. *Glob. Ecol.*
365   *Biogeogr.* **28**, 1578–1596.

366 Leroy, B., Bellard, C., Dubos, N., Colliot, A., Vasseur, M., Courtial, C., Bakkenes, M., Canard, A., & Ysnel,
367   F. (2014). Forecasted climate and land use changes, and protected areas: the contrasting case of spiders.
368   (B. Bradley, Ed.)*Divers. Distrib.* **20**, 686–697.

369 Ma, L., Buckley, L. B., Huey, R. B., & Du, W. G. (2018). A global test of the cold-climate hypothesis for the
370   evolution of viviparity of squamate reptiles. *Glob. Ecol. Biogeogr.* **27**, 679–689.

371 Madison, D. M. (1997). The Emigration of Radio-Implanted Spotted Salamanders, Ambystoma maculatum.
372   *J. Herpetol.* **31**, 542–551.

373 Mccain, C. M., & Colwell, R. K. (2011). Assessing the threat to montane biodiversity from discordant shifts
374   in temperature and precipitation in a changing climate. *Ecol. Lett.* **14**, 1236–1245.

375 Miaud, C., Andreone, F., Riberon, A., Michelis, S. De, Clima, V., Castanet, J., Francillon-Vieillot, H., &
376   Guyétant, R. (2001). Variations in age , size at maturity and gestation duration among two neighbouring
377   populations of the alpine salamander (Salamandra lanzai). *J. Zool.* **254**, 251–260.

378 Phillips, S. J., Dudík, M., Elith, J., Graham, C. H., Lehmann, A., Leathwick, J., & Ferrier, S. (2009). Sample
379   selection bias and presence-only distribution models: Implications for background and pseudo-absence
380   data. *Ecol. Appl.* **19**, 181–197.

381 Platts, P. J., Garcia, R. A., Hof, C., Foden, W., Hansen, L. A., Rahbek, C., & Burgess, N. D. (2014).
382   Conservation implications of omitting narrow-ranging taxa from species distribution models, now and
383   in the future. *Divers. Distrib.* **20**, 1307–1320.

384 Prasad, A. M., Iverson, L. R., & Liaw, A. (2006). Newer classification and regression tree techniques:
385   Bagging and random forests for ecological prediction. *Ecosystems* **9**, 181–199.

Rana, S. K., Rana, H. K., Luo, D., & Sun, H. (2021). Estimating climate-induced 'Nowhere to go' range shifts of the Himalayan Incarvillea Juss. using multi-model median ensemble species distribution models. *Ecol. Indic.* **121**, 107127.

Raxworthy, C. J., Pearson, R. G., Rabibisoa, N., Rakotondrazafy, A. M., Ramanamanjato, J.-B., Raselimanana, A. P., Wu, S., Nussbaum, R. a., & Stone, D. a. (2008). Extinction vulnerability of tropical montane endemism from warming and upslope displacement: a preliminary appraisal for the highest massif in Madagascar. *Glob. Chang. Biol.* **14**, 1703–1720.

Reinthaler-Lottermoser, U., Meikl, M., Gimeno, A., Weinke, E., & Schwarzenbacher, R. (2010). A new approach for surveying the Alpine Salamander (Salamandra atra) in Austria. *Acta Herpetol.* **5**, 249–253.

Ribéron, A., & Miaud, C. (1999). Home range and shelter use in Salamandra lanzai (Caudata, Salamandridae). *Amphibia-Reptilia* **21**, 225–260.

Romano, A., Costa, A., Salvidio, S., Menegon, M., Garollo, E., Tabarelli de Fatis, K., Miserocchi, D., Matteucci, G., & Pedrini, P. (2018). Forest management and conservation of an elusive amphibian in the Alps: Habitat selection by the Golden Alpine Salamander reveals the importance of fine woody debris. *For. Ecol. Manage.* **424**, 338–344.

Schwartz, M. W. (2012). Using niche models with climate projections to inform conservation management decisions. *Biol. Conserv.* **155**, 149–156.

Seddon, P. J., Strauss, W. M., & Innes, J. (2012). Animal Translocations: What are they and why do we do them? *Reintroduction Biol. Integr. Sci. Manag.* 1–32.

Steen, V. A., Tingley, M. W., Paton, P. W. C., & Elphick, C. S. (2021). Spatial thinning and class balancing: Key choices lead to variation in the performance of species distribution models with citizen science data. *Methods Ecol. Evol.* **12**, 216–226.

Stickley, S. F., & Fraterrigo, J. M. (2023). Microclimate species distribution models estimate lower levels of climate-related habitat loss for salamanders. *J. Nat. Conserv.* **72**, 126333.

Tafani, M., Cohas, A., Bonenfant, C., Gaillard, J. M., & Allainé, D. (2013). Decreasing litter size of marmots over time: A life history response to climate change? *Ecology* **94**, 580–586.

Tessa, G., Crottini, A., & Andreone, F. (2007). A new finding of Salamandra lanzai in the Upper Sangone Valley (NW Italy) marks the species ' most disjunct population (Amphibia : Urodela : Salamandridae) **2**, 53–58.

Tewksbury, J. J., Huey, R. B., & Deutsch, C. A. (2008). Putting the Heat on Tropical Animals The Scale of Prediction. *Science* **320**, 1296–1297.

Thévenin, C. (2019). Reintroduction efficiency: a stepping stone approach to reintroduction success? *Anim. Conserv.* **22**, 116–117.

Valavi, R., Elith, J., Lahoz-Monfort, J. J., & Guillera-Arroita, G. (2019). blockCV: An r package for generating spatially or environmentally separated folds for k-fold cross-validation of species distribution models. *Methods Ecol. Evol.* **10**, 225–232.

Valavi, R., Guillera-Arroita, G., Lahoz-Monfort, J. J., & Elith, J. (2021). Predictive performance of presence-only species distribution models: a benchmark study with reproducible code. *Ecol. Monogr.* **0**, 1–27.

Vollering, J., Halvorsen, R., Auestad, I., & Rydgren, K. (2019). Bunching up the background betters bias in species distribution models. *Ecography (Cop.).* **42**, 1717–1727.

426 Werner, P., Lötters, S., & Schmidt, B. R. (2014). Analysis of habitat determinants in contact zones of
427 parapatric European salamanders. *J. Zool.* **292**, 31–38.
428

429 **Supporting information**

430 Fig. S1 Predicted values for the climate suitability of *S. lanzai* based on Worldclim predictors

431   Fig. S2 Predicted values for the climate suitability of *S. lanzai* based on CHELSA predictors

432   Fig. S3 Predicted values for the climate suitability of *S. atra* based on Worldclim predictors

433   Fig. S4 Predicted values for the climate suitability of *S. atra* based on CHELSA predictors