Main Manuscript

Wandering albatrosses exert high take-off effort only when both wind and waves are gentle

Leo Uesaka *[a,b], Yusuke Goto [a,c,d], Masaru Naruoka [e], Henri Weimerskirch [d], Katsufumi Sato [a], Kentaro Q. Sakamoto [a].

[a] Atmosphere and Ocean Research Institute, The University of Tokyo, Kashiwa, Chiba 277-8564, Japan; [b] Information and Technology Center, The University of Tokyo, Kashiwa, Chiba, 277-0882, Japan; [c] Graduate School of Environmental Studies, Nagoya University, Furo, Chikusa, Nagoya, 464-8601, Japan; [d] Centre d'Etudes Biologiques de Chize (CEBC), UMR 7372 CNRS, Université de La Rochelle, 79360 Villiers-en-Bois, France; [e] Aeronautical Technology Directorate, Japan Aerospace Exploration Agency (JAXA), Chofu, Japan

Corresponding Author: Leo Uesaka, Information and Technology Center, The University of Tokyo, Kashiwa, Chiba, 277-0882, Japan, +81 90 2831 5573

Contact info: leo-u@g.ecc.u-tokyo.ac.jp

Declaration of interests: The authors declare no conflict of interest.

Abstract

The relationship between the environment and marine animal small-scale behavior is
not fully understood. This is largely due to the difficulty in obtaining environmental datasets
with a high spatiotemporal precision. The problem is particularly pertinent in assessing the
influence of environmental factors in rapid, high energy consuming behavior such as seabird
take-off. To fill the gaps in the existing environmental datasets, we employed novel techniques
using animal-borne sensors with motion records to estimate wind and ocean wave parameters
and evaluated their influence on wandering albatross take-off patterns. Measurements
revealed that wind speed and wave heights experienced by wandering albatrosses during
take-off ranged from 0.7 ~ 15.4 m/s and 1.6 ~ 6.4 m, respectively. The four indices measured
(flapping number, frequency, sea surface running speed, and duration) also varied with the
environmental conditions (e.g., flapping number varied from 0 to over 20). Importantly, take-
off was easier under higher wave conditions than under lower wave conditions at a constant
wind speed, and take-off effort increased only when both wind and waves were gentle. Our
data suggests that both ocean waves and winds play important roles for albatross take-off and
advances our current understanding of albatross flight mechanisms.

Impact statement: Wind and ocean wave conditions experienced by albatrosses were
estimated using an animal-borne recorder and revealed that take-off was easier under higher
wave conditions.

Keywords: Seabird, Biologging, Ocean environment, Take-off

Introduction

Various oceanic environmental factors affect the ecology of marine animals. Predicted climate changes suggest increases in extreme climatic events (such as cyclones). Thus, evaluating individual relationships between each environmental factor and marine animal behaviors is urgent for marine ecological conservation, especially for top predators that significantly impact the entire ecosystem. However, there are potential limitations: direct measures of marine animals empirical environmental data is nearly impossible due to the spatiotemporal gaps in the observation network of the open ocean (Ardhuin et al., 2019; Villas Bôas et al., 2019). Various environmental parameters (such as ocean wind, waves, and sea surface temperature) are assumed to be important factors affecting the movement and foraging of flying seabirds (Dunn, 1973; Haney et al., 1992; Adams and Navarro, 2005; Nevitt et al., 2008; Suryan et al., 2008). Previous research has revealed that many interesting seabird behaviors correlate with the ocean environment. However, the environmental data largely relies on ocean climatic models (Padget et al., 2019; Weimerskirch and Prudor, 2019; Clay et al., 2020; Clairbaux et al., 2021), as in situ observation data is limited and often collected a long distance from the bird. For example, records are collected at the colony island or using the nearest government observation point (Kogure et al., 2016; Yamamoto et al., 2017). Therefore, interpreting the data is difficult when doubts exist on whether birds actually experienced the same environmental conditions, making any conclusions conservative estimates only (Clay et al., 2020; Clairbaux et al., 2021). For instance, although winter cyclones in the North Atlantic can induce mass seabird mortality, revealing the small-scale behavioral responses which lead to mortality is almost impossible with the spatiotemporal limits of thermodynamic modelling data (Clairbaux et al., 2021).

Seabird take-off may be affected by the surrounding environment (Clay et al., 2020) but has never been effectively investigated. Notably, behaviors with short timeframes (such as take-offs) require localized environmental data on spatiotemporally small scales, which is difficult to obtain, even using mathematical weather models. Many procellariiformes have special flight techniques that use vertical wind shear, called dynamic soaring (Rayleigh, 1883; Richardson, 2011), while take-off requires a large amount of energy (Sakamoto et al., 2013) owing to vigorous flapping (Sato et al., 2009; Sakamoto et al., 2013) and sea surface running to reach the velocity to initiate take-off (Sato et al., 2009). Previous research revealed the heart rate of the largest seabird, wandering albatross (*Diomedea exulans*), drastically increases at the moment of take-off reaching 3 – 4 times the basal heart rate (Weimerskirch et al., 2000). After take-off, the tachycardia progressively decreases during flight (Weimerskirch et al., 2000),

the flying heart rate is close to the basal rate of a resting bird on the nest. Therefore, the high energy expenditure associated with take-off strongly influences the total energy expenditure of wandering albatross during the foraging trip, unlike flight duration or distance (Shaffer et al., 2001a). Thus, take-off is one of the most important behaviors in the daily energy budget of flying seabirds in the open ocean. Understanding the relationship between take-off and the ocean environment is critical for estimating the future climate change effects on the life history of seabirds (Weimerskirch et al., 2012).

Previous research has partially identified the role of wind conditions on take-off when investigating general flight tactics of seabirds (Kogure et al., 2016; Clay et al., 2020). For example, the flapping effort of the European shag (*Gulosus aristotelis*) at take-off decreases as wind speed increases (Kogure et al., 2016). However, a comprehensive understanding of take-off has not been achieved as other environmental parameters, such as waves, have not been investigated. Ocean waves potentially affect take-off efforts because seabirds usually run on the ocean surface as they take-off (Norberg and Norberg, 1971; Sato et al., 2009). Additionally, the ocean surface slope is a key factor in creating complicated wind patterns immediately above the surface, which may affect the flight tactics of procellariiformes (Bousquet et al., 2017).

In this study, we devised a new approach to estimate the empirical local environmental conditions using seabird dynamic motion records without the aid of either mathematical weather models or observational data. The recent development of animal-borne recorders has been remarkable (Wilmers et al., 2015). It is now possible to deploy various sensors on animal-borne recorders, to generate a new field of oceanography: ocean observations using animal-borne sensors (Harcourt et al., 2019; McMahon et al., 2021). Many studies have reported that marine environmental data can be collected using highly mobile marine animals such as pinnipeds, sea turtles, and seabirds (Charrassin et al., 2002, 2008; Roquet et al., 2014; Doi et al., 2019). Furthermore, unlike direct measurement by deploying sensors (e.g., thermometers), indirect techniques to observe the physical environment via the dynamic animal motion records (Yoda et al., 2014; Yonehara et al., 2016; Goto et al., 2017; Sánchez-Román et al., 2019; Uesaka et al., 2022) generated using the Global Navigation Satellite System (GNSS) have been developed recently. GNSS is now able to record animal position and movement every second (or even in sub-second scales) using very small animal-borne recorders. These newly developed techniques using animal-borne recorders should compensate for previous observational gaps in oceanic data, especially in the open ocean and polar regions where our access and deployment of observation equipment is complicated. The environmental variables

obtained directly from free ranging animals provide the localized environmental conditions they
experience. Seabirds are one of the most enthusiastically studied oceanic species because of
their high mobility and adaptability to both air and water. Methods involving wind and wave
observation, using GNSS data regarding the flight paths and floating motions of seabirds on
the sea surface, are well-developed and have potential applicability in future studies (Yonehara
et al., 2016; Goto et al., 2017; Uesaka et al., 2022).
Wandering albatrosses were investigated because their habitat includes the
Subantarctic (30 °S – 60 °S), where the ocean is annually rough (Suryan et al., 2008) causing
their flight behaviors to be largely influenced by the ocean conditions (Richardson, 2011;
Weimerskirch et al., 2012). Furthermore, previous studies have revealed the foraging area has
shifted southward annually with the polar shift of the westerly wind pattern (Weimerskirch et
al., 2012). Considering the enormous cost of take-off (Weimerskirch et al., 2000; Shaffer et al.,
2001a), studying their response to various environmental conditions is essential for us to
estimate the impacts of climate change on the life history of seabirds. We aim to estimate the
physical environmental conditions (ocean winds and wave heights) experienced by wandering
albatrosses as they take-off by utilizing the dynamic motion records to evaluate the effects of
wind and wave conditions on take-off dynamics (Fig. 1). Procellariiformes, like many seabird
species, require extensive limb motion for take-off which is not limited to flapping behavior.
Therefore, the evaluation of take-off effort involves both surface running and flapping
behaviors. The wandering albatross individuals were tagged using recorders that include both
GPS and acceleration sensors with high time resolutions.
113

Results

Trip data

We obtained 1477 h from 44 wandering albatrosses in 2019 (N = 21, 623 h) and 2020 (N = 23, 854 h). Two types of recorders with different battery sizes were used. The mean recording time of the trip data and the standard deviation (SD) was 9.5 ± 1.3 h for the small battery recorders and 59.7 ± 9.6 h for the large battery recorders. The albatross sex ratio was balanced between years and recorder type (Table. S1).

The absolute value of the GPS horizontal velocity revealed 703 take-offs from 1477 h of trips. A total of 453 out of 703 take-offs were followed by more than 5 min of flight. For each flight, the wind speed and direction were estimated using the flight path (Yonehara et al., 2016). A total of 299 take-offs occurred after more than 15 min of floating time. Wave heights were estimated for each of the 299 take-offs using the floating motions (Uesaka et al., 2022). For 185 take-offs, we estimated the wind and wave conditions in combination.

Environmental conditions at the take-off moment

Of the 453 estimated wind parameters, 26 were unreliable based on the Akaike information criterion (AIC) comparison and were not included in the analysis. The remaining 427 results revealed wind speeds of 6 – 8 m/s were most frequently experienced by taking-off wandering albatrosses (Fig. S1A). Mean ± SD of the estimated wind speed was 6.5 ± 2.7 m/s, and the maximum and minimum wind speeds were 15.4 and 0.7 m/s, respectively. Winds blowing from west to east were frequently observed (Fig. S1B). This result is consistent with the prevalence of westerlies around the wandering albatross breeding colony (Nicholls et al., 1997; Weimerskirch et al., 2015).

Ocean waves were estimated using all 299 take-offs after more than 15 min of floating time to calculate the significant wave height. The most frequently experienced wave heights ranged from 2.5 ~ 3.0 m at the take-off moment (Fig. S1C) and the mean ± SD was 3.0 ± 0.8 m. The minimum and maximum wave heights were 1.6 and 6.4 m, respectively. Like wind direction, the wave direction (coming from) had a west bias due to the westerlies (Fig. S1D).

Take-off properties

To quantify the take-off effort, we calculated four parameters: running duration, running speed, flapping number, and flapping frequency from the acceleration records obtained at the moment of take-off. Mean ± SD running duration of wandering albatross was 5.1 ± 1.5 s with a range from 1.1 to 11.7 s (Fig. S2A). The mean value for males was slightly lower than that

for females (Fig. S3A), however, the difference was not significant (M: 5.0 ± 1.5 s, F:5.2 ± 1.5

s, p = 0.10, Mann-Whitney U test). The albatross running speed mean value ± SD was 6.5 ±

1.6 m/s (Fig. S2B). Male birds had slightly higher speeds than females (M: 6.7 ± 1.5 m/s,

F:6.3 ± 1.6, p < 0.01, Fig. S3B). Running duration and speed significantly correlated

(Pearson's r = 0.57, p < 0.01). The linear regression slope (with a fixed intercept of zero) was

1.23 m/s^2 (Fig. S4). We interpret the slope as the running wandering albatross acceleration.

The flapping number, i.e., the number of wing flaps after the running phase, was

estimated using the dorsoventral acceleration. The mean flapping number was 4.3 times with

a range from zero to over 20 times (Fig. S2C). Take-offs without flapping after the running

phase were frequently observed (33.3%). Conversely, continuous flapping above 20 times

were also occasionally observed, which corresponds to a lengthy flapping duration (8 s ~) after

take-off, considering the flapping frequency of wandering albatross (2.5 ~ 3.0 Hz). There was

no significant difference in flapping number between the sexes (p = 0.22, Mann-Whitney U test,

Fig. S3C). The mean ± SD flapping frequency was 2.55 ± 0.29 Hz, and most ranged from 2 to

3 Hz (Fig. S2D). However, some flapping frequency results were outside the detection range

(1.8 ~ 4 Hz) and not included in our analysis. Therefore, the sample size of the flapping

frequency used in our analysis was 669. There was no significant difference in flapping

frequency between the sexes (p = 0.18, Mann-Whitney U test, Fig. S3D).

Environmental effects on take-off parameters

The take-off directions were compared with the wind direction estimated from the flight

path after take-off. Wandering albatrosses tended to take-off with headwinds (p < 0.01, v-test)

(Fig. 2). However, the cruising direction (moving direction from the take-off point to the bird

location after 5 min) did not correlate with headwind direction. The mean ± SD air speed of

wandering albatrosses at the end of the running phase (lift-off moment from the sea surface)

calculated using the running speed, wind speed, and relative take-off direction was 12.2 ± 3.1

m/s.

The relationships between each take-off parameter (running duration, running speed,

flapping number, and flapping frequency) with environmental conditions (wind speed and wave

height) were tested using linear mixed models (LMMs). The running duration required for

wandering albatross take-off significantly decreased as wind speed and wave height increased

(Fig. 3). Similarly, the running speed was significantly lower under stronger wind and higher

wave conditions. Wandering albatrosses tend to flap fewer times under stronger wind

conditions. Conversely, wandering albatrosses can flap over 20 times in weak wind conditions,

although the flapping number in weak wind conditions varies greatly. There is also a declining

trend in the flapping number with wave height. Albatross take-offs in wave heights below 2 m

always require flapping. The flapping frequency was lower as the wind speed and wave height

increased, however, the trend with higher wave heights remains unclear. The LMMs results

are provided in Table S2 of the supplementary information.

Independent effect of wind and waves on take-off

Although some ocean wave components are generated by ocean winds, the

correlation between the wind speed and wave height is not consistent. Some of the albatross

take-offs involved information on both wind speed and wave height. Therefore, we evaluated

the respective effects of wind and waves on wandering albatross take-offs. The correlation

between wind speed and wave height was not strong ($r = 0.27$, $p < 0.01$). Some take-offs were

performed in weak winds but high wave conditions or the opposite conditions (Fig. 4A). Take-

off conditions were divided into four environmental categories using the peak value, which

were 6.0 m/s (wind speed) and 2.8 m (wave height). The categories comprised: 48 samples

(weak wind low wave: WL), 33 samples (weak wind high wave: WH), 27 samples (strong wind

low wave: SL), and 77 samples (strong wind high wave: SH). The running duration varied

significantly between the four categories ($p < 0.01$, Kruskal-Wallis test). The mean running

duration in the WL conditions was 6.0, which was the longest of the four categories (Fig. 4B).

Relatively long running (of over 6 s) mainly occurred in WL conditions, and the running duration

decreased with the wind speed or wave height (Fig. S5). Similar results were obtained for both

running speed and flapping number. However, flapping frequency did not significantly vary

between the four categories ($p = 0.06$). Take-offs involving over 30 flaps mainly occurred in

WL conditions.

The variance inflation factor (VIF) of wind speed and wave height was 6.86, which did

not exceed the general threshold of 10 (Dormann et al., 2013). Among the LMMs, models

including wind speed, wave height, and the interaction used the smallest AIC for all take-off

parameters (Table S3). However, the difference between the lowest and the second lowest

AIC was below two for running speed, flapping number, and flapping frequency. The running

duration simulation using the estimated coefficient shows that even under weak wind

conditions (2 m/s), running duration decreases from 8 to 4 s as the wave height increases.

Conversely, low values were maintained under strong wind conditions (8 m/s) regardless of

the wave height (Fig. 5). Similarly, the running speed decreased from 9 to 6 m/s as the wave

height increased, regardless of the wind strength. The flapping number followed the same

trend. Conversely, the flapping frequency did not decrease as the wave height increased.

Discussion

Although observational networks in the ocean are under development and mathematical weather modelling accuracy is increasing, they remain unable to accurately estimate the surrounding environment of marine animals at small scales. Here, we demonstrated that environmental variables estimated using individual animal recorders provide valuable new insight into locomotor behavior when spatiotemporal scale and accuracy of mathematical weather models and observational networks are too broad for the research. In this study, we provided details on how seabird take-offs are affected by wind and waves.

Seabird take-offs using accelerometers

We quantified the running behavior of seabirds at the moment of take-off, which is the most energy-consuming behavior for soaring seabirds (Weimerskirch et al., 2000; Shaffer et al., 2001a; Sakamoto et al., 2013). Previous studies have ascribed this large energy expenditure to the vigorous flapping required for take-off (Shaffer et al., 2001a; Sato et al., 2009; Clay et al., 2020). Indeed, the continuous flapping behavior, which is rare in cruising flight, was recorded even after the running phase of take-off in this study. However, we suggest that the running behavior should also entail a large cost in take-off because albatrosses have to reach a fast initial speed to lift off the sea surface by rapidly moving their hindlimbs for up to ~ 10 s in unfavorable conditions (as demonstrated in this study).

We provide a first attempt at detecting the running signal of seabird take-offs and construct a relatively simple algorithm (which can be easily applied to other species) using lateral acceleration. The running duration may increase or decrease by approximately 0.5 s depending on the algorithm configuration, such as smoothing parameters and threshold values. However, we focused on the relative changes in the running behavior in association with wind speed and wave height, and absolute value error is not a serious problem.

The flapping characteristics of wandering albatross during the running phase were also researched. However, the dorsoventral acceleration signal fluctuates during the running phase making it difficult to identify each flapping signal, even after applying the band-pass filter. Therefore, counting the number of flaps immediately after the running phase was the only reliable parameter for evaluating flapping effort. Sato et al. (2009) reported that the flapping frequency of wandering albatrosses at the moment of take-off is higher (2.9 ~ 3.4 Hz) than that of cruising flight (2.5 ~ 2.7 Hz). In our study, the flapping frequency after the running phase was not as high as Sato et al. (2009) reported. Therefore, it is likely that wandering albatrosses undertake high frequency flapping only during the running phase, as they lift off the sea surface.

After lift-off (i.e., the running phase is completed), wandering albatrosses continue with a
moderate flapping frequency until they reach a certain degree of flight stability. Simultaneous
video records of flapping and running motions with acceleration records are required to
separate the parameter estimates.

In-depth studies on seabird take-offs are just beginning with the aid of miniaturized
animal-borne recorders with the main aim of understanding how seabirds flap their wings.
However, in land birds (e.g., finches and doves) take-off requires a large contribution by
hindlimbs (Provini et al., 2012) and the role of the hindlimb in take-off kinematics is as important
as that of the wing (Provini and Abourachid, 2018). Therefore, it is highly likely that seabird
take-offs also require a substantial contribution by the hindlimbs, and thus, further seabird
hindlimb research is required. Our study provides the basic characteristics of wandering
albatross running behavior, including running duration and speed.

Take-off effort with environmental conditions

Our results demonstrate that wandering albatrosses can take-off in a variety of
environmental conditions (wind speed: 0.7 ~ 15.4 m/s, wave height: 1.6 ~ 6.4 m). A previous
study on wandering albatrosses identified the transition state from resting to flying tended to
increase as the wind speed increased (Clay et al., 2020). Our results found some take-offs
were performed under weak wind (2 – 4 m/s) conditions, suggesting wind speed is not the only
parameter influencing flight decisions of wandering albatross, and that wave height should be
included in future studies.

The results showed that the running and flapping behavior tended to decrease as the
wind or wave conditions increased. Running duration decreased as either the wind speed or
the wave height increased and peaked when both the wind and wave conditions were weak.
The same trends existed in running speed and flapping number. Although optimum statistical
models for each take-off parameter were determined using the AIC value, some models
provided similar results to this model. For instance, the AIC difference in running speed
between the best model and the second-lowest AIC model was only 0.27. However, both
models included wind speed and wave height as the explanatory variables, similar to the other
take-off parameters, except flapping frequency. The purpose of constructing a linear model
was to clarify whether the effects of wind and waves are independent. As long as both wind
speed and wave height were included as explanatory variables in the model, they reduced the
running and flapping behavior requirement. Therefore, we can conclude that both strong winds
and high waves aid wandering albatross take-offs. The flapping frequency after the running

phase was the only parameter that did not correlate with wave height (as identified using the

LMMs model). However, we assume the flapping frequency during the running phase is more

important. Future research needs to investigate the effects of wave height as wandering

albatrosses need to climb up or run down the wave slope. Therefore, the flapping frequency

during the running phase should be highly influenced by wave height.

Contribution of strong wind and high waves to seabird take-off

The reduced running behavior and flapping times under strong wind conditions is

simply described by the lift force mechanism which has been predicted by previous studies

(Kogure et al., 2016; Clay et al., 2020). Seabirds need to gain lift force before take-off, and the

magnitude of force is proportional to the square of the relative speed of the wings to the

surrounding air (air speed) (Vogel, 1983). It has been anecdotally suggested that seabirds

take-off into the wind (i.e., headwind), because stronger winds can produce a sufficiently large

lift even before the ground speed of the seabird reaches the value required for flight. As a

partial demonstration of this theory, a study on the European shag (*Gulosus aristotelis*)

(Kogure et al., 2016) found the take-off direction was significantly biased toward headwinds.

Regarding soaring seabirds, only one study (Clay et al., 2020) on wandering albatrosses has

confirmed a bias in take-off direction with wind direction. However, the authors acknowledge

the limits in the mathematical weather model and GPS sampling resolutions and recognize the

unreliability of small-scale responses to in situ variation in the atmosphere. Our study reveals

wandering albatrosses significantly tend to take-off into the wind, using robust fine scale data

estimated from the flight records of wandering albatross. Moreover, there was no correlation

between cruising and headwind direction, indicating that wandering albatrosses face the wind

on take-off regardless of their destination. Our data is reliable as the empirical value provided

is actually experienced by the albatross. Furthermore, by quantitatively evaluating the flapping

and running effort, we demonstrate the theory of effortless take-offs by soaring seabirds in

stronger wind conditions.

The mean air speed of wandering albatrosses at the end of the running phase was

close to the average flight speed (approximately 15 m/s) (Weimerskirch et al., 2002), and

similar to predicted best glide speeds, (Shaffer et al., 2001b) indicating that wandering

albatrosses gain sufficient lift at the end of the running phase and efficiently utilize ocean wind.

Wind speed varies with altitude, therefore the wind blowing on the ocean surface must be

smaller than the values estimated from the flight records of wandering albatross as they usually

fly 3 – 12 m above the ocean surface (Pennycuick, 1982). Therefore, the calculated air speed

is probably an overestimate when compared with the ocean surface. To compensate for the
insufficient lift force gained during the running phase, wandering albatrosses flap their wings
several times after the running phase. Therefore, the flapping number in weak wind conditions
can exceed dozens before reaching stable flight.

The most important finding of our study is that the take-off effort estimated by the
running behavior and number of flaps decreased not only with stronger winds but also with
higher waves. While the role of ocean wind on flying seabirds has been well described
(Pennycuick, 2008), how ocean waves influence the flight of seabirds remains largely unknown.
However, many studies have reported the characteristic flight of soaring seabirds by tracking
the ocean wave surface over long distances (Pennycuick, 1982, 2008; Richardson, 2011;
Stokes and Lucas, 2021), which even occurs in weak or no wind conditions (Pennycuick, 1982).
Seabirds seem to be aided by atmospheric forces above the slope-like wave topography; the
flight method using the shape of wave is called wave-slope soaring (Richardson, 2011). It is
well recognized that air flows occur above ocean waves (Buckley and Veron, 2016; Bousquet
et al., 2017). Richardson (2011) described the theoretical model of wave-slope soaring, where
the flight mechanism of albatross is a combination of both dynamic soaring, which uses vertical
wind shear above the ocean surface (~ 15 m), and wave-slope soaring, which uses the updraft
caused by the wave topography. Thus, seabirds can continue to soar in weak wind conditions.
Further, mathematical analysis has revealed that the wave-induced updraft (even in windless
conditions) can provide 60% of the transportation cost of a brown pelican (*Pelecanus*
occidentalis, 2 ~ 3 kg), which is a wave-slope soarer (Stokes and Lucas, 2021). Thus, it is
possible that the take-off effort by wandering albatross is also reduced by high waves. While
qualitative field observations and mathematical demonstrations provide the only previous
research on the role of waves on soaring seabirds, we experimentally demonstrated that ocean
waves aid the most energy-consuming behavior, take-off. This finding helps future discussions
on ocean topographical mechanisms affecting seabird flight.

The mechanism by which high waves aid wandering albatross take-off is not entirely
clear. It is difficult to conclude a certain updraft is producing additional lift for wandering
albatrosses, and it is also possible that there are other unresolved mechanisms. For example,
a rough topographic surface can provide a favorable bump, like a slope or cliff to jump off into
the air. Our results were restricted to wave height as the parameter of the ocean surface.
Future research involving ocean surface steepness or wave frequency components will reveal
the detailed mechanism of how waves facilitate seabird take-off behavior. In particular, ocean
surface topography relies heavily on whether the dominant wave component is due to a swell

(low-frequency waves propagated from a distance) or wind waves (high-frequency waves
generated by local wind); moreover, this topography affects the wind pattern on the sea surface.

In conclusion, we revealed how the take-off effort of wandering albatross changes in
various oceanic conditions. As take-off is one of the most energy-consuming behaviors that
can dominate the total energy expenditure of a wandering albatross journey, this data will be
of great value for considering how climate changes can alter the life of albatrosses. Future
research, especially on albatrosses, should quantitatively evaluate the energy consumption of
take-off with the wind and wave conditions. Currently, there is no major barrier to
accomplishing this goal, it would require utilizing motion records to estimate the surrounding
environment with additional methods to estimate energy consumption, such as cardiograms.
Recognizing the negative effect of the changing oceanic environment on seabirds (Sydeman
et al., 2015), revealing the direct small-scale mechanisms of environmental factors (such as
wind, wave, tide, current, and sea surface temperature) effects on animal behavior, especially
in take-off is urgently required. The concept of estimating the surrounding environment using
motion records is a novel solution with great potential to unravel the small spatiotemporal
uncertainties in seabird research.

Materials and Methods

Field experiment

The recorders, Ninja-scan (Little Leonardo, Tokyo, Japan), record triaxial acceleration at a very high time resolution (100 Hz). Ninja-scan also records 3D GPS positions (5 Hz), Doppler velocity (5 Hz), temperature (6 Hz), pressure (6 Hz), geomagnetism (6 Hz), and angular velocity (100 Hz). There are two types of Ninja-scans with different battery masses (Naruoka et al., 2021). Small Ninja-scans weighed 28 g, which is 0.3 ~ 0.4% of wandering albatross body mass, and are expected to record for 7 h. Large Ninja-scans weighed 91 g, which corresponds to 0.8 ~ 1.3% of wandering albatross body mass, and are expected to record for 65 h.

Ninja scans were attached to breeding wandering albatrosses at Possession Island, Crozet archipelago (46°25 S, 51°44 E) in the South Indian Ocean in 2019 and 2020. In 2019, 12 small Ninja scans were attached (in tandem) to 6 individuals. On each bird, one recorder had a delay timer so that the two recording periods did not overlap. Additionally, 15 birds had individual Ninja-scans attached, of which 8 were small Ninja-scans and 7 were large Ninja scans. In 2020, 10 small Ninja-scans were attached in tandem to 5 individuals. Additionally, 19 birds had individual Ninja-scans attached, of which 7 were small Ninja-scans and 12 were large Ninja-scans. In summary, 21 and 24 wandering albatrosses were tagged in 2019 and 2020, respectively. All experiments were performed from late January to early March of each year, which corresponds to the incubation period of wandering albatrosses. Recorders were attached to the back of each bird with waterproof tape (Tesa, Hamburg, Germany) and glue (Loctite; Henkel, Dusseldorf, Germany). All recorders were retrieved within 35 days. One small Ninja-scan which had been attached in isolation in 2020 did not work correctly. The effects of the attached recorders on wandering albatrosses were previously assessed (Phillips et al., 2003; Barbraud and Weimerskirch, 2012) and revealed that small recorders (less than 3% of their body mass) do not negatively impact breeding or foraging behaviors. The experiment was conducted as part of Program 109 of the Institut Polaire Paul Emile Victor with permission from the Préfet des Terrs Australes et Antarctiques Françaises, France.

Take-off identification

First, data recorded on the colony island was eliminated based on the GPS position. Then, take-off was determined using the absolute value of the GPS horizontal velocity. When wandering albatrosses float on the sea surface (i.e., before take-off), a relatively low speed which is generally below 2.5 m/s, is recorded, while the flying speed exceeds 5 m/s

(Weimerskirch et al., 2002). Take-off was defined as the moment when the horizontal speed
exceeds 4 m/s and rises to a higher speed. The soaring (flying) speed occasionally meets this
criteria. Therefore, the horizontal speed was smoothed using the moving average (20 points :
0.4 s). If the horizontal speed crossed the 4 m/s line several times within a short period, they
were classed as take-offs for very short flights and were not used in our investigation.
Therefore, we selected only take-offs that included over 30 s of floating followed by over 30 s
of flying.

Wind estimation

Yonehara et al., (2016) proposed estimating the wind speed and direction of seabird
flight paths using the sinusoidal curve relationship between flight speed and flight direction.
When seabirds fly in the air, their flight speed against the ground (ground speed) is mainly
affected by the wind speed, which is maximized in tail winds and minimized in head winds.
The maximum speed is the sum of the flight speed against air (air speed) and wind speed,
whereas the minimum speed is the difference between the air speed and wind speed. The
relationship between the flight speed (ground speed) and flight direction recorded by the GPS
are fitted using a sinusoidal curve (Shimatani et al., 2012). We followed the methodology in
Yonehara et al. (2016) to collate the flight speed V and flight direction θ data for 5 min after
take-off and the curve was fitted using the following equation:

$$V = V_a + V_w \cos(\theta + \phi_w)$$

where V_a is the air speed, V_w is the wind speed, and ϕ_w is the wind direction. Ten seconds
immediately after the take-off moment was not included in the estimation. Following Yonehara
et al. (2016), the AIC of the sinusoidal fitting was compared to the linear fitting with a fixed
slope of zero. When the AIC difference between the linear and sinusoidal fitting was below two,
the estimated results were considered unreliable and discarded. Wind speeds and directions
were not calculated when take-offs were not followed by over five minutes of flight. The
sinusoidal fitting was performed using Igor Pro version 8.04 (Wavemetrics, Portland, OR, USA).

Wave estimation

The ocean wave properties experienced by seabirds before take-off were estimated
by analyzing the floating motion at the sea surface (Uesaka et al., 2022). The wave height was
estimated from the vertical GPS displacement records before take-off. The estimate requires
sufficiently long records of vertical displacement. Therefore, the wave height was not
calculated for take-offs that did not follow a surface floating time of over 15 min. The sampling

period of 15 min ensured the reliability of the wave statistics (Whitford et al., 2001) and

provided a large volume of estimated wave data. The estimate did not include the 10 s before

the detected take-off moment. We followed the methodology of Uesaka et al. (2022). The

vertical GPS displacement records were high-pass filtered using a cut-off frequency of 0.07

Hz to eliminate the GPS derived error (Olynik et al., 2002). We separated the time series record

of the vertical displacement into individual waves by applying the zero-up-crossing method.

The mean wave height of the highest third of all individual waves was calculated to provide the

significant wave height, which is the most widely used statistical wave parameter (Whitford et

al., 2001).

Sea surface running by seabirds

Many procellariiformes require a running phase before take-off from the sea surface

(Sato et al., 2009). However, studies using accelerometers have not focused on the

acceleration signal of this behavior. Surface running involves asymmetrical leg movements.

Therefore, the lateral acceleration obtained from the recorder (attached to the back of the

seabirds) provided signals derived from the running motion (Fig. 6). We confirmed that running

signals appear in the lateral acceleration records at the moment of take-off by streaked

shearwaters (*Calonectris leucomelas*), which are phylogenetically similar to wandering

albatrosses (see Supplementary Information Text and Fig. S6).

To explore the running duration of wandering albatross, we constructed an algorithm

to detect the running phase from the lateral acceleration around take-off. The lateral

acceleration signal is composed of a dominant component (0.25 ~ 0.4 s) and a high-frequency

fluctuation component (< 0.2 s period). Although the dominant component is the lateral

movement derived from surface running, the flapping period of wandering albatross appears

around this period (0.3 ~ 0.4 s). The flapping behavior is laterally symmetrical and does not

appear in the lateral acceleration records. However, this is not always the case, when

(occasionally) recorders are attached to the back of the seabird in a slightly tilted position. To

avoid confusion between running and flapping behavior, a high-frequency fluctuation

component in the lateral acceleration signal was used to detect the running phase. A band-

pass filter was designed to extract the high-frequency fluctuation component from the

acceleration records, and then the variance per unit time (0.6 s) was calculated at each point.

Running phase was defined as when the acceleration variance exceeded the threshold value

(2% of the peak value). This algorithm reasonably detects the running phase regardless of the

running duration. If there is a signal gap in the middle of the running phase, the algorithm

regards the gap as the end of the running phase, underestimating the running duration.

However, these cases are rare, and we assume it does not affect our evaluation of the running

characteristics of wandering albatross. The horizontal speed at the end of the running phase

and take-off direction were calculated using the GPS velocity. The take-off direction was

defined as the vectoral average direction during the running phase. We also calculated the

cruising direction which was defined as the moving direction 5 min after take-off. All procedures

were performed using Igor Pro version 8.04 (Wavemetrics, Portland, OR, USA).

Flapping behavior after the running phase

Dorsoventral acceleration records include signals derived from seabird wing flapping

behavior (Fig. 6). The flapping signals during the running phase fluctuate, which is assumed

to be caused by the leg-derived dorsoventral motion. This caused the flapping data to be

unclear in identifying the flapping number and frequency. Therefore, we only focused on the

wing flapping signals after the running phase. A bandpass filter extracted the clearest flapping

signals (1.8 ~ 4.0 Hz). The number of continuous flapping signals after the running phase were

counted. The flapping period of the wandering albatross is approximately 0.3 ~ 0.4 s, therefore

we defined the end point when the flapping interval exceeded 0.5 s. The flapping frequency

after the running phase was calculated using the spectral peak value of the continuous

wavelet-transformed dorsoventral acceleration. All procedures were performed using Igor Pro

version 8.04 (Wavemetrics, Portland, OR, USA).

Comparison of the take-off parameters with environmental conditions

The wind directional bias of the take-off direction was tested using the v-test (modified

Rayleigh test). The air speed V_a at the end of the running phase was estimated using the

following equation based on the parameters obtained from this study:

$$V_a = V_r + V_w \cos(\theta_t - \phi_w)$$

where V_r is the running speed at the end of the running phase, V_w is the wind speed, θ_t is the

take-off direction, and ϕ_w is the wind direction. The effects of wind speed and wave height on

each take-off parameter (running duration, running speed, flapping number, and flapping

frequency) were evaluated using linear mixed models (LMMs) with individuals treated as

random effects. To identify significance levels, the models were compared to null models

based on the AIC value.

To evaluate the combined effect of wind and waves, we categorized take-off conditions

into four groups, "WL conditions," "WH conditions," "SL conditions," and "SH conditions".

Threshold values were decided based on the peak in the curve of the fitted probability density

distribution (wind speed: 6.0 m/s, wave height 2.8 m). Weibull distribution and log normal

distribution were used as the fitting function for wind speed and wave height, respectively

(Ferreira and Guedes Soares, 2000; Carta et al., 2009). The values of each take-off parameter

were compared between the four groups by Kruskal-Wallis test. Furthermore, the independent

effects of wind and waves on take-off parameters were evaluated using LMMs, including wind

speed, wave height, and their interaction as explanatory parameters with individuals as

random effects. VIF was also calculated before the LMMs analysis to assess whether the

multicollinearity effect could be dismissed. The V-test was performed using Igor Pro version

8.04 (Wavemetrics, Portland, OR, USA). Statistical test and LMMs calculations were

performed using the Python 3.0 and PypeR package.

Acknowledgments

We thank Yoshinari Yonehara, Julien Collet, Timothée Poupart, and all the members of the research station in the Crozet Islands for their support during the fieldwork. We appreciate the fruitful comments of Kagari Aoki, Chihiro Kinoshita, Laxmi Kumar Parajuli, and Aran Garrod. We are grateful to Taichi Sakamoto and Takashi Mukai (ATTACCATO) for providing customized rechargeable Li-ion batteries for the Ninja-scans. We also thank Michihiko Suzuki and Koichiro Ikeda (Little Leonardo) for molding the Ninja-scans. The field studies were financially supported by a research project entitled "Cyber ocean: next generation navigation system on the sea" funded by the CREST program (JPMJCR1685) of Japan Science and Technology Agency; Grants-in-Aid for Scientific Research from JSPS (17H00776 to K. Sato); the Tohoku Ecosystem-Associated Marine Science; and Institut Polaire Français Paul-Émile Victor Program 109.

References

Adams NJ, Navarro RA. 2005. Foraging of a coastal seabird: flight patterns and movements of breeding cape gannets *Morus capensis*. *African J Mar Sci* **27**:239–248. doi:10.2989/18142320509504082

Ardhuin F, Stopa JE, Chapron B, Collard F, Husson R, Jensen RE, Johannessen J, Mouche A, Passaro M, Quartly GD, Swail V, Young I. 2019. Observing sea states. *Front Mar Sci* **6**:1–29. doi:10.3389/fmars.2019.00124

Barbraud C, Weimerskirch H. 2012. Assessing the effect of satellite transmitters on the demography of the wandering albatross *Diomedea exulans*. *J Ornithol* **153**:375–383. doi:10.1007/s10336-011-0752-8

Bousquet GD, Triantafyllou MS, Slotine JJE. 2017. Optimal dynamic soaring consists of successive shallow arcs. *J R Soc Interface* **14**. doi:10.1098/rsif.2017.0496

Buckley MP, Veron F. 2016. Structure of the airflow above surface waves. *J Phys Oceanogr* **46**:1377–1397. doi:10.1175/JPO-D-15-0135.1

Carta JA, Ramírez P, Velázquez S. 2009. A review of wind speed probability distributions used in wind energy analysis. Case studies in the canary islands. *Renew Sustain Energy Rev* **13**:933–955. doi:10.1016/j.rser.2008.05.005

Charrassin JB, Hindell M, Rintoul SR, Roquet F, Sokolov S, Biuw M, Costa D, Boehme L, Lovell P, Coleman R, Timmermann R, Meijers A, Meredith M, Park YH, Bailleul F, Goebel M, Tremblay Y, Bost CA, McMahon CR, Field IC, Fedak MA, Guinet C. 2008. Southern ocean

frontal structure and sea-ice formation rates revealed by elephant seals. *Proc Natl Acad Sci*
U S A **105**:11634–11639. doi:10.1073/pnas.0800790105

Charrassin JB, Park YH, Maho Y Le, Bost CA. 2002. Penguins as oceanographers unravel
hidden mechanisms of marine productivity. *Ecol Lett* **5**:317–319. doi:10.1046/j.1461-
0248.2002.00341.x

Clairbaux M, Mathewson P, Porter W, Fort J, Strøm H, Moe B, Fauchald P, Descamps S,
Helgason HH, Bråthen VS, Merkel B, Anker-Nilssen T, Bringsvor IS, Chastel O,
Christensen-Dalsgaard S, Danielsen J, Daunt F, Dehnhard N, Erikstad KE, Ezhov A, Gavrilo
M, Krasnov Y, Langset M, Lorentsen SH, Newell M, Olsen B, Reiertsen TK, Systad GH,
Thórarinsson TL, Baran M, Diamond T, Fayet AL, Fitzsimmons MG, Frederiksen M, Gilchrist
HG, Guilford T, Huffeldt NP, Jessopp M, Johansen KL, Kouwenberg AL, Linnebjerg JF,
Major HL, Tranquilla LMF, Mallory M, Merkel FR, Montevecchi W, Mosbech A, Petersen A,
Grémillet D. 2021. North atlantic winter cyclones starve seabirds. *Curr Biol* **31**:3964-
3971.e3. doi:10.1016/j.cub.2021.06.059

Clay TA, Joo R, Weimerskirch H, Phillips RA, den Ouden O, Basille M, Clusella-Trullas S, Assink
JD, Patrick SC. 2020. Sex-specific effects of wind on the flight decisions of a sexually
dimorphic soaring bird. *J Anim Ecol* **89**:1811–1823. doi:10.1111/1365-2656.13267

Doi T, Storto A, Fukuoka T, Suganuma H, Sato K. 2019. Impacts of temperature measurements
from sea turtles on seasonal prediction around the arafura sea. *Front Mar Sci* **6**:1–11.
doi:10.3389/fmars.2019.00719

Dormann CF, Elith J, Bacher S, Buchmann C, Carl G, Carré G, Marquéz JRG, Gruber B,
Lafourcade B, Leitão PJ, Münkemüller T, Mcclean C, Osborne PE, Reineking B, Schröder
B, Skidmore AK, Zurell D, Lautenbach S. 2013. Collinearity: a review of methods to deal
with it and a simulation study evaluating their performance. *Ecography (Cop)* **36**:27–46.
doi:10.1111/j.1600-0587.2012.07348.x

Dunn EK. 1973. Changes in fishing ability of terns associated with windspeed and sea surface
conditions. *Nature* **244**:520–521. doi:10.1038/244520a0

Ferreira JA, Guedes Soares C. 2000. Modelling distributions of significant wave height. *Coast
Eng* **40**:361–374. doi:10.1016/S0378-3839(00)00018-1

Goto Y, Yoda K, Sato K. 2017. Asymmetry hidden in birds' tracks reveals wind, heading, and
orientation ability over the ocean. *Sci Adv* **3**:1–10. doi:10.1126/sciadv.1700097

Haney JC, Fristrup KM, Lee DS. 1992. Geometry of visual recruitment by seabirds to ephemeral
foraging flocks. *Ornis Scand* **23**:49–62. doi:10.2307/3676427

Harcourt R, Sequeira AMM, Zhang X, Roquet F, Komatsu K, Heupel M, McMahon C, Whoriskey
F, Meekan M, Carroll G, Brodie S, Simpfendorfer C, Hindell M, Jonsen I, Costa DP, Block B,
Muelbert M, Woodward B, Weise M, Aarestrup K, Biuw M, Boehme L, Bograd SJ, Cazau D,

Charrassin JB, Cooke SJ, Cowley P, de Bruyn PJN, Jeanniard du Dot T, Duarte C, Eguíluz VM, Ferreira LC, Fernández-Gracia J, Goetz K, Goto Y, Guinet C, Hammill M, Hays GC, Hazen EL, Hückstädt LA, Huveneers C, Iverson S, Jaaman SA, Kittiwattanawong K, Kovacs KM, Lydersen C, Moltmann T, Naruoka M, Phillips L, Picard B, Queiroz N, Reverdin G, Sato K, Sims DW, Thorstad EB, Thums M, Treasure AM, Trites AW, Williams GD, Yonehara Y, Fedak MA. 2019. Animal-borne telemetry: an integral component of the ocean observing toolkit. *Front Mar Sci* **6**. doi:10.3389/fmars.2019.00326

Kogure Y, Sato K, Watanuki Y, Wanless S, Daunt F. 2016. European shags optimize their flight behavior according to wind conditions. *J Exp Biol* **219**:311–318. doi:10.1242/jeb.131441

McMahon CR, Roquet F, Baudel S, Belbeoch M, Bestley S, Blight C, Boehme L, Carse F, Costa DP, Fedak MA, Guinet C, Harcourt R, Heslop E, Hindell MA, Hoenner X, Holland K, Holland M, Jaine FRA, Jeanniard du Dot T, Jonsen I, Keates TR, Kovacs KM, Labrousse S, Lovell P, Lydersen C, March D, Mazloff M, McKinzie MK, Muelbert MMC, O'Brien K, Phillips L, Portela E, Pye J, Rintoul S, Sato K, Sequeira AMM, Simmons SE, Tsontos VM, Turpin V, van Wijk E, Vo D, Wege M, Whoriskey FG, Wilson K, Woodward B. 2021. Animal Borne Ocean Sensors – AniBOS – an essential component of the global ocean observing system. *Front Mar Sci* **8**:1–21. doi:10.3389/fmars.2021.751840

Naruoka M, Goto Y, Weimerskirch H, Mukai T, Sakamoto T, Sakamoto KQ, Sato K. 2021. Application of inertial and GNSS integrated navigation to seabird biologging. *J Robot Mechatronics* **33**:526–536. doi:10.20965/jrm.2021.p0526

Nevitt GA, Losekoot M, Weimerskirch H. 2008. Evidence for olfactory search in wandering albatross, *Diomedea exulans*. *Proc Natl Acad Sci U S A* **105**:4576–4581. doi:10.1073/pnas.0709047105

Nicholls DG, Murray MD, Butcher E, Moors P. 1997. Weather systems determine the non-breeding distribution of wandering albatrosses over southern oceans. *Emu* **97**:240–244. doi:10.1071/mu97031

Norberg RA, Norberg UM. 1971. Take-off, landing, and flight speed during fishing flights of *Gavia stellata* (pont.). *Ornis Scand* **2**:55–67. doi:10.2307/3676239

Olynik M, Petovello MG, Cannon ME, Lachapelle G. 2002. Temporal variability of GPS error sources and their effect on relative positioning accuracy. Proceedings of the 2002 National Technical Meeting of The Institute of Navigation. 2002. pp. 877–888.

Padget O, Stanley G, Willis JK, Fayet AL, Bond S, Maurice L, Shoji A, Dean B, Kirk H, Juarez-Martinez I, Freeman R, Bolton M, Guilford T. 2019. Shearwaters know the direction and distance home but fail to encode intervening obstacles after free-ranging foraging trips. *Proc Natl Acad Sci U S A* **116**:21629–21633. doi:10.1073/pnas.1903829116

Pennycuick C. 2008. Modelling the flying bird, Academic Press. New York.

Pennycuick CJ. 1982. The flight of petrels and albatrosses (procellariiformes), observed in south
georgia and its vicinity. *Philos Trans R Soc London B, Biol Sci* **300**:75–106.
doi:10.1098/rstb.1982.0158

Phillips RA, Xavier JC, Croxall JP. 2003. Effects of satellite transmitters on albatrosses and
petrels. *Auk* **120**:1082–1090. doi:10.1093/auk/120.4.1082

Provini P, Abourachid A. 2018. Whole-body 3D kinematics of bird take-off: key role of the legs to
propel the trunk. *Sci Nat* **105**. doi:10.1007/s00114-017-1535-8

Provini P, Tobalske BW, Crandell KE, Abourachid A. 2012. Transition from leg to wing forces
during take-off in birds. *J Exp Biol* **215**:4115–4124. doi:10.1242/jeb.074484

Rayleigh. 1883. The soaring of birds. *Nature* **27**:534–535. doi:10.1038/027534a0

Richardson PL. 2011. How do albatrosses fly around the world without flapping their wings?. *Prog
Oceanogr* **88**:46–58. doi:10.1016/j.pocean.2010.08.001

Roquet F, Williams G, Hindell MA, Harcourt R, McMahon C, Guinet C, Charrassin JB, Reverdin
G, Boehme L, Lovell P, Fedak M. 2014. A southern indian ocean database of hydrographic
profiles obtained with instrumented elephant seals. *Sci Data* **1**:1–10.
doi:10.1038/sdata.2014.28

Sakamoto KQ, Takahashi A, Iwata T, Yamamoto T, Yamamoto M, Trathan PN. 2013. Heart rate
and estimated energy expenditure of flapping and gliding in black-browed albatrosses. *J
Exp Biol* **216**:3175–3182. doi:10.1242/jeb.079905

Sánchez-Román A, Gómez-Navarro L, Fablet R, Oro D, Mason E, Arcos JM, Ruiz S, Pascual A.
2019. Rafting behaviour of seabirds as a proxy to describe surface ocean currents in the
balearic sea. *Sci Rep* **9**:1–11. doi:10.1038/s41598-018-36819-w

Sato K, Sakamoto KQ, Watanuki Y, Takahashi A, Katsumata N, Bost CA, Weimerskirch H. 2009.
Scaling of soaring seabirds and implications for flight abilities of giant pterosaurs. *PLoS One*
4:1–6. doi:10.1371/journal.pone.0005400

Shaffer SA, Costa DP, Weimerskirch H. 2001a. Behavioural factors affecting foraging effort of
breeding wandering albatrosses. *J Anim Ecol* **70**:864–874. doi:10.1046/j.0021-
8790.2001.00548.x

Shaffer SA, Weimerskirch H, Costa DP. 2001b. Functional significance of sexual dimorphism in
wandering albatrosses, *Diomedea exulans. Funct Ecol* **15**:203–210. doi:10.1046/j.1365-
2435.2001.00514.x

Shimatani IK, Yoda K, Katsumata N, Sato K. 2012. Toward the quantification of a conceptual
framework for movement ecology using circular statistical modeling. *PLoS One* **7**:e50309.
doi:10.1371/journal.pone.0050309

Stokes IA, Lucas AJ. 2021. Wave-slope soaring of the brown pelican. *Mov Ecol* **9**:1–13.
doi:10.1186/s40462-021-00247-9

Suryan RM, Anderson DJ, Shaffer SA, Roby DD, Tremblay Y, Costa DP, Sievert PR, Sato F,

Ozaki K, Balogh GR, Nakamura N. 2008. Wind, waves, and wing loading: morphological

specialization may limit range expansion of endangered albatrosses. *PLoS One* **3**:e4016.

doi:10.1371/journal.pone.0004016

Sydeman WJ, Poloczanska E, Reed TE, Thompson SA. 2015. Climate change and marine

vertebrates. *Science* **350**:772–777. doi:10.1126/science.aac9874

Uesaka L, Goto Y, Yonehara Y, Komatsu K, Naruoka M, Weimerskirch H, Sato K, Sakamoto KQ.

2022. Ocean wave observation utilizing motion records of seabirds. *Prog Oceanogr*

200:102713. doi:10.1016/j.pocean.2021.102713

Villas Bôas AB, Ardhuin F, Ayet A, Bourassa MA, Brandt P, Chapron B, Cornuelle BD, Farrar JT,

Fewings MR, Fox-Kemper B, Gille ST, Gommenginger C, Heimbach P, Hell MC, Li Q,

Mazloff MR, Merrifield ST, Mouche A, Rio MH, Rodriguez E, Shutler JD, Subramanian AC,

Terrill EJ, Tsamados M, Ubelmann C, Sebille E van. 2019. Integrated observations of global

surface winds, currents, and waves: requirements and challenges for the next decade. *Front*

Mar Sci **6**:1–34. doi:10.3389/fmars.2019.00425

Vogel S. 1983. Life in moving fluids. Princeton University Press. doi:10.1515/9780691212975

Weimerskirch H, Bonadonna F, Bailleul F, Mabille G, Dell'Omo G, Lipp H-P. 2002. GPS tracking

of foraging albatrosses. *Science* **295**:1259–1259. doi:10.1126/science.1068034

Weimerskirch H, Delord K, Guitteaud A, Phillips RA, Pinet P. 2015. Extreme variation in migration

strategies between and within wandering albatross populations during their sabbatical year

and their fitness consequences. *Sci Rep* **5**:8853. doi:10.1038/srep08853

Weimerskirch H, Guionnet T, Martin J, Shaffer SA, Costa DP. 2000. Fast and fuel efficient?

Optimal use of wind by flying albatrosses. *Proc R Soc B Biol Sci* **267**:1869–1874.

doi:10.1098/rspb.2000.1223

Weimerskirch H, Louzao M, de Grissac S, Delord K. 2012. Changes in wind pattern alter

albatross distribution and life-history traits. *Science* **335**:211–214.

doi:10.1126/science.1210270

Weimerskirch H, Prudor A. 2019. Cyclone avoidance behaviour by foraging seabirds. *Sci Rep*

9:1–9. doi:10.1038/s41598-019-41481-x

Whitford DJ, Waters JK, Vieira MEC. 2001. Teaching time-series analysis. ii. wave height and

water surface elevation probability distributions. *Am J Phys* **69**:497–504.

doi:10.1119/1.1331301

Wilmers CC, Nickel B, Bryce CM, Smith JA, Wheat RE, Yovovich V, Hebblewhite M. 2015. The

golden age of bio-logging: how animal-borne sensors are advancing the frontiers of ecology.

Ecology **96**:1741–1753. doi:10.1890/14-1401.1

Yamamoto T, Kohno H, Mizutani A, Sato H, Yamagishi H, Fujii Y, Murakoshi M, Yoda K. 2017.

Effect of wind on the flight of brown booby fledglings. *Ornithol Sci* **16**:17–22.

doi:10.2326/osj.16.17

Yoda K, Shiomi K, Sato K. 2014. Foraging spots of streaked shearwaters in relation to ocean

surface currents as identified using their drift movements. *Prog Oceanogr* **122**:54–64.

doi:10.1016/j.pocean.2013.12.002

Yonehara Y, Goto Y, Yoda K, Watanuki Y, Young LC, Weimerskirch H, Bost CA, Sato K. 2016.

Flight paths of seabirds soaring over the ocean surface enable measurement of fine-scale

wind speed and direction. *Proc Natl Acad Sci U S A* **113**:9039–9044.

doi:10.1073/pnas.1523853113

 Figures

Fig. 1. Conceptual framework of the study estimating environmental conditions experienced by studied individual.

Fig. 2. Effect of wind direction on wandering albatross take-off. Relative take-off direction to wind direction (black circles, n = 427) significantly distributed around 0° (head-wind), in contrast to cruising direction relative to the wind (gray x-mark, n = 427). The radial axis represents the wind speed.

Fig. 3. Environmental effect on take-off. Effort for the take-off (running duration, running speed, flapping number, and flapping frequency) significantly decreased as wave height and wind speed increased (p < 0.01) except the relationship between flapping frequency and wave height (p = 0.026). Solid line shows the linear regression line determined from the LMMs and the number at the right top corner on each graph shows the sample sizes.

Fig. 4. (A) Correlation between wind speed and wave height was weak (r = 0.27, n = 185). Bar charts and solid lines written above and right of the scatter plot are normed histograms of wind speed, wave height and curve fitted lines. Based on the peak value of fitted lines scatter plots were divided into four groups, WL: weak wind low wave (open square), WH: weak wind high wave (filled square), SL: strong wind low wave (open circle), and SH: strong wind high wave (filled circle). (B) Take-off effort comparison among four groups (a: running duration, b: running speed, c: flapping number, and d: flapping frequency). Cross mark indicates the mean value.

Fig. 5. (A) running duration, (B) running speed, (C) flapping number, and (D) flapping frequency in response to the wave height change under weak wind (dashed line, 2 m/s) and strong wind (solid line, 8 m/s) conditions estimated from the LMM results. Gray area represents 99% CI.

Fig. 6. Time series data of horizontal speed (top), lateral acceleration (middle), and dorsoventral acceleration (bottom) signals of the wandering albatross at the moment of take-off. Horizontal speed starts increasing from the beginning of the take-off. Red square shows the detected running phase based on the variance of the lateral acceleration signal. Red bars show the detected flapping behavior after the running phase based on the dorsoventral acceleration signal. Dorsoventral signal during the running phase fluctuates, probably due to the shaking body derived from the running motion, and thus it is not easy to judge the existence of flapping behavior.