Main Manuscript

Wandering albatross exert high take-off effort in weak wind with low wave conditions.

Leo Uesaka [*a,b], Yusuke Goto [a,c,d], Masaru Naruoka [e], Henri Weimerskirch [d], Katsufumi Sato [a], Kentaro Q. Sakamoto [a].

[a] Atmosphere and Ocean Research Institute, The University of Tokyo, Kashiwa, Chiba 277-8564, Japan; [b] Information and Technology Center, The University of Tokyo, Kashiwa, Chiba, 277-0882, Japan; [c] Graduate School of Environmental Studies, Nagoya University, Furo, Chikusa, Nagoya, 464-8601, Japan; [d] Centre d'Etudes Biologiques de Chize (CEBC), UMR 7372 CNRS, Université de La Rochelle, 79360 Villiers-en-Bois, France; [e] Aeronautical Technology Directorate, Japan Aerospace Exploration Agency (JAXA), Chofu, Japan

Corresponding Author: Leo Uesaka, Information and Technology Center, The University of Tokyo, Kashiwa, Chiba, 277-0882, Japan, +81 90 2831 5573

Contact info: leo-u@g.ecc.u-tokyo.ac.jp

Declaration of interests: The authors declare no conflict of interest.

Abstract

The relationship between the environment and the small-scaled behavior of marine animals is not fully understood. This is largely due to the difficulty in obtaining environmental datasets with a high spatiotemporal precision. The problem is particularly pertinent in assessing the influence of environmental factors in rapid high energy consuming behavior such as seabirds take-off. Here, to fill the gaps in existing database, we employed novel techniques using animal-borne sensors with motion records to estimate parameters on wind and ocean waves, and evaluated their influence on wandering albatrosses take-off. The measurement revealed that the wind speed and the wave height that the wandering albatrosses experienced during take-off ranged from 0.7 ~ 15.4 m/s and 1.6 ~ 6.4 m, respectively. The four indices that were measured (flapping number, frequency, running speed, and duration on the sea surface) also varied with the environmental conditions (i.e., flapping number varied from 0 to over 20). Importantly, taking-off was easier under higher wave condition in the constant wind speed, and take-off effort increased only when both wind and waves were gentle. Our data suggests that both ocean waves and winds play important roles in albatross take-off, and advances our current understanding of albatross flight mechanisms.

Impact statement: Winds and ocean waves condition experienced by albatrosses were estimated using animal-borne recorder and revealed that taking-off was easier under higher wave condition in the constant wind speed.

Keywords: Seabird, Biologging, ocean environment, Take-off

Introduction

Various oceanic environmental factors affect the ecology of marine animals.
Predicted climate changes suggest increases in extreme climatic events (such as cyclones).
Evaluating individual relationships between each environmental factor and marine animal
behaviors is urgent for marine ecological conservation, especially for top predators that
significantly impact the whole ecosystem. However, there are potential limitations: direct
measures of empirical environmental data of marine animals is nearly impossible due to the
spatiotemporal gaps in the observation network of the open ocean (Ardhuin et al., 2019;
Villas Bôas et al., 2019). Various environmental parameters (such as ocean wind, waves,
and sea surface temperature) are assumed to be important factors affecting the movement
and foraging of flying seabirds (Adams and Navarro, 2005; Dunn, 1973; Haney et al., 1992;
Nevitt et al., 2008; Suryan et al., 2008). Previous research has revealed many interesting
seabird behaviors correlate with the ocean environment. However, the environmental data
largely relies on ocean climatic modelling (Clairbaux et al., 2021; Clay et al., 2020; Padget et
al., 2019; Weimerskirch and Prudor, 2019), as in situ observation data is limited and often
collected a long distance from the bird. For example, records are collected at the colony
island or using the nearest government observation point (Kogure et al., 2016; Yamamoto et
al., 2017). Therefore, interpreting the data is difficult when doubts exist on whether the birds
actually experienced the environmental conditions, making any conclusions conservative
estimates only (Clairbaux et al., 2021; Clay et al., 2020). For example, although winter
cyclones in the North Atlantic can induce mass seabird mortality, revealing the small-scale
behavioral responses which lead to mortality is almost impossible with the spatiotemporal
limits of thermodynamic modelling data (Clairbaux et al., 2021).

Seabird take-off may be affected by the surrounding environment (Clay et al., 2020)
but has never been effectively investigated. Notably, behaviors with short timeframes (such
as take-offs) require environmental data on spatiotemporally small scales, which is difficult to
obtain, even using climatic models. Many procellariiformes have special flight techniques that
use vertical wind shear, called dynamic soaring (Rayleigh, 1883; Richardson, 2011), and
take-off requires a large amount of energy (Sakamoto et al., 2013) owing to vigorous flapping
(Sakamoto et al., 2013; Sato et al., 2009) and sea surface running to reach the velocity to
initiate take-off (Sato et al., 2009). Previous research revealed the heart rate of the largest
seabird, wandering albatross (*Diomedea exulans*), drastically increases at the moment of
take-off reaching 3 – 4 times the basal heart rate (Weimerskirch et al., 2000). After take-off,
the tachycardia progressively decreases during flight (Weimerskirch et al., 2000), the flying

heart rate is close to the basal rate of a resting bird on the nest. Therefore, the highest

proportion of the total energy expenditure of wandering albatross during the foraging trip is

the take-off rate, not the duration or distance of the flight (Shaffer et al., 2001). Thus, take-off

is one of the most important behaviors in the daily energy budget of flying seabirds in the

open ocean. Understanding the relationship between take-off and the ocean environment is

critical for estimating the future climate change effects on the life history of seabirds

(Weimerskirch et al., 2012).

Previous research has partially identified the role of wind conditions on take-off when

investigating general flight tactics of seabirds (Clay et al., 2020; Kogure et al., 2016). For

example, the flapping effort of the European shag (*Gulosus aristotelis*) at take-off decreases

as wind speed increases (Kogure et al., 2016). However, a comprehensive understanding of

take-off has not been achieved as other environmental parameters, such as waves, have not

been investigated. Ocean waves potentially affect take-off efforts because seabirds usually

run on the ocean surface as they take-off (Norberg and Norberg, 1971; Sato et al., 2009).

Additionally, the ocean surface slope is a key factor in creating complicated wind patterns

immediately above the surface, which may affect the flight tactics of procellariiformes

(Bousquet et al., 2017).

In this study, we devised a new approach to estimate the empirical environmental

conditions using seabird dynamic motion records without the aid of either climate models or

observational data. The recent development of animal-borne recorders has been remarkable

(Wilmers et al., 2015). It is now possible to deploy various sensors on animal-borne

recorders, to generate a new field of oceanography: ocean observations using animal-borne

sensors (Harcourt et al., 2019; McMahon et al., 2021). Many studies have reported that

marine environmental data can be collected using highly mobile marine animals such as

pinnipeds, sea turtles, and seabirds (Charrassin et al., 2008, 2002; Doi et al., 2019; Roquet

et al., 2014). Furthermore, unlike direct measurement by deploying sensors (e.g.,

thermometers), indirect techniques to observe the physical environment via the dynamic

animal motion records (Goto et al., 2017; Sánchez-Román et al., 2019; Uesaka et al., 2022;

Yoda et al., 2014; Yonehara et al., 2016) generated using the Global Navigation Satellite

System (GNSS) have been developed recently. GNSS is now able to record their position

and movement every second (or even in sub-second scales) using very small animal-borne

recorders. These newly developed techniques using animal-borne recorders should

compensate for previous observational gaps in oceanic data, especially in the open ocean

and polar regions where our access and deployment of observation equipment is

complicated. The environmental variables obtained directly from animals provide the
environmental conditions they experience. Seabirds are one of the most enthusiastically
studied oceanic species because of their high mobility and adaptability to both air and water.
Methods involving the observation of wind and waves, using GNSS data regarding the flight
paths and floating motions of seabirds on the sea surface, are well-developed and have
potential applicability in future studies (Goto et al., 2017; Uesaka et al., 2022; Yonehara et al.,
2016).

Wandering albatrosses were investigated because their habitat includes the
Subantarctic (30 °S – 60 °S), where the ocean is annually rough (Suryan et al., 2008)
causing their flight style to be largely influenced by the ocean conditions (Richardson, 2011;
Weimerskirch et al., 2012). Furthermore, previous studies have revealed the foraging area
has annually shifted southward with the polar shift of the westerly wind pattern
(Weimerskirch et al., 2012). Considering the enormous cost of take-off (Shaffer et al., 2001;
Weimerskirch et al., 2000), studying their response to various environmental conditions is
essential for us to estimate the impacts of climate change on the life history of seabirds. We
aim to estimate the physical environmental conditions (ocean winds and wave heights)
experienced by wandering albatross as they take-off by utilizing the dynamic motion records
to evaluate the effects of wind and wave conditions on take-off dynamics (Fig. 1).
Procellariiformes, like many seabird species, require extensive limb motion for take-off which
is not limited to flapping behavior. Therefore, the evaluation of take-off effort involves both
surface running and flapping behaviors. The wandering albatross individuals were tagged
using recorders that include both GPS and acceleration sensors with high time resolutions.
115

Results

Trip data

We obtained 1477 h from 44 wandering albatross in 2019 (N = 21, 623 h) and 2020 (N = 23, 854 h). Two types of recorders with different battery sizes were used. The mean recording time of the trip data and the standard deviation (SD) was 9.5 ± 1.3 h for the small battery recorders and 59.7 ± 9.6 h for the large battery recorders. The albatross sex ratio was balanced between years and recorder type (Table. S1).

The absolute value of the GPS horizontal velocity revealed 703 take-offs from 1477 h of trips. A total of 453 out of 703 take-offs were followed by more than 5 min of flight. For each flight, the wind speed and direction were estimated using the flight path (Yonehara et al., 2016). A total of 299 take-offs occurred after more than 15 min of floating time. The wave height were estimated for each 299 take-offs using the floating motions (Uesaka et al., 2022). For 185 take-offs, we estimated the wind and wave conditions in combination.

Environmental conditions at the take-off moment

Of the 453 estimated wind parameters, 26 were unreliable based on the Akaike information criterion (AIC) comparison and were not included in the analysis. The remaining 427 results revealed wind speeds of 6 – 8 m/s were most frequently experienced by taking-off wandering albatross (Fig. S1A). Mean ± SD of the estimated wind speed was 6.5 ± 2.7 m/s, and the maximum and minimum wind speeds were 15.4 and 0.7 m/s, respectively. Winds blowing from west to east were frequently observed (Fig. S1B). This result is consistent with the prevalence of westerlies around the wandering albatross breeding colony (Nicholls et al., 1997; Weimerskirch et al., 2015).

Ocean waves were estimated using all 299 take-offs after more than 15 min of floating time to calculate the significant wave height. The most frequently experienced wave heights ranged from 2.5 ~ 3.0 m at the take-off moment (Fig. S1C) and the mean ± SD was 3.0 ± 0.8 m. The minimum and maximum wave heights were 1.6 and 6.4 m, respectively. Like wind direction, the wave direction (coming from) had a west bias due to the westerlies (Fig. S1D).

Take-off properties

To quantify the take-off effort, we calculated four parameters: running duration, running speed, flapping number, and flapping frequency from the acceleration records obtained at the moment of take-off. Mean ± SD running duration of wandering albatross was

5.1 ± 1.5 s with a range from 1.1 s to 11.7 s (Fig. S2A). The mean value for males was
slightly lower than that for females (Fig. S3A), but the difference was not significant (M:
5.0 ± 1.5 s, F:5.2 ± 1.5 s, p = 0.10, Mann-Whitney U test). The albatross running speed
mean value ± SD was 6.5 ± 1.6 m/s (Fig. S2B). Male birds had slightly higher speeds than
females (M: 6.7 ± 1.5 m/s, F:6.3 ± 1.6, p < 0.01, Fig. S3B). Running duration and speed
significantly correlated (Pearson's r = 0.57, p < 0.01). The linear regression slope (with a
fixed intercept of zero) was 1.23 m/s^2 (Fig. S4). We interpret the slope as the acceleration of
running wandering albatross.

The flapping number, i.e., the number of wing flaps after the running phase, was
estimated using the dorsoventral acceleration. The mean flapping number was 4.3 times with
a range from zero to over 20 times (Fig. S2C). Take-offs without flapping after the running
phase were frequently observed (33.3%). Conversely, continuous flapping above 20 times
were also occasionally observed, which corresponds to a long time flapping (8 s ~) after
take-off considering the flapping frequency of wandering albatross (2.5 ~ 3.0 Hz). There was
no significant difference in flapping number between the sexes (p = 0.22, Mann-Whitney U
test, Fig. S3C). The mean ± SD flapping frequency was 2.55 ± 0.29 Hz, and most ranged
from 2 to 3 Hz (Fig. S2D). However, some flapping frequency results were outside the
detection range (1.8 ~ 4 Hz) and not included in our analysis. Therefore, the sample size of
the flapping frequency used in our analysis was 669. There was no significant differences in
flapping frequency between the sexes (p = 0.18, Mann-Whitney U test, Fig. S3D).

Environmental effects on take-off parameters

The take-off directions were compared with the wind direction estimated from the
flight path after take-off. Wandering albatross tended to take-off with headwinds (p < 0.01, v-
test) (Fig. 2). However, the cruising direction did not correlate with headwind direction,
indicating that wandering albatrosses face the wind in take-off regardless of their destination.
The mean ± SD air speed of wandering albatrosses at the end of the running phase (lift-off
moment from the sea surface) calculated using the running speed, wind speed, and relative
take-off direction was 12.2 ± 3.1 m/s.

The relationships between each take-off parameter (running duration, running speed,
flapping number, and flapping frequency) with environmental conditions (wind speed and
wave height) were tested using linear mixed models (LMMs). The running duration required
for wandering albatross take-off significantly decreased as wind speed and wave height
increased (Fig. 3). Similarly, the running speed was significantly lower under stronger wind

and higher wave conditions. Wandering albatross tend to flap fewer times under stronger
wind conditions. Conversely, wandering albatross can flap over 20 times in weak wind
conditions, although the flapping number in weak wind conditions varies greatly. There is
also a declining trend in the flapping number with wave height. Albatross taking-offs in wave
heights below 2 m always require flapping. The flapping frequency was lower as the wind
speed and wave height increased, but the trend with higher wave heights remains unclear.
The LMMs results are provided in Table S2 of the supplementary information.
191
Independent effect of wind and waves on take-off
Although some components of ocean waves are generated by ocean winds, the
correlation between the wind speed and wave height is not consistent. Some of the albatross
take-offs involved information on both wind speed and wave height. Therefore, we evaluated
the respective effects of wind and waves on wandering albatross take-offs. The correlation
between wind speed and wave height was not strong ($r = 0.27$, $p < 0.01$). Some take-offs
were performed in weak winds but high wave conditions or the opposite conditions (Fig. 4A).
The take-off conditions were divided into four environmental categories using the peak in the
curve of the fitted probability density distribution, which were 6.0 m/s (wind speed) and 2.8 m
(wave height) (wind speed: Weibull distribution (Carta et al., 2009), wave height: log normal
distribution (Ferreira and Guedes Soares, 2000)). The categories comprised: 48 samples
(weak wind low wave: WL), 33 samples (weak wind high wave: WH), 27 samples (strong
wind low wave: SL), and 77 samples (strong wind high wave: SH). The running duration
varied significantly between the four categories ($p < 0.01$, Kruskal-Wallis test). The mean
running duration in the WL conditions was 6.0, which was the longest of the four categories
(Fig. 4B). Relatively long running (of over 6 s) mainly occurred in WL conditions, and the
running duration decreased with the wind speed or wave height (Fig. S5). Similar results
were obtained for both running speed and flapping number. However, flapping frequency did
not significantly vary between the four categories ($p = 0.06$). Take-offs involving over 30 flaps
mainly occurred in WL conditions.
The variance inflation factor (VIF) of wind speed and wave height was 6.86, which
did not exceed the general threshold of 10 (Dormann et al., 2013). The LMMs, models
including wind speed, wave height, and the interaction used the smallest AIC for all take-off
parameters (Table S3). However, the difference between the lowest and the second lowest
AIC was below two for running speed, flapping number, and flapping frequency. The
simulation of the running duration using the estimated coefficient shows that even under

weak wind conditions (2 m/s), the running duration decreases from 8 to 4 s as the wave

height increases. Conversely, low values were maintained under strong wind conditions

(8 m/s) regardless of the wave height (Fig. 5). Similarly, the running speed decreased from 9

to 6 m/s as the wave height increased, regardless of the wind strength. The flapping number

followed the same trend. Conversely, the flapping frequency did not decrease as the wave

height increased.

Discussion

Although observation networks in the ocean are under development and climate modelling accuracy is increasing, they remain unable to accurately estimate the surrounding environment of marine animals at small scales. We demonstrate the environmental variables estimated using individual animal recorders are highly valuable when the spatiotemporal scale and accuracy of climate models and observation networks are too broad for the research. We provide details on how seabird take-offs are affected by wind and waves.

Seabird take-offs using accelerometers

We quantified the running behavior of seabirds at the moment of take-off, which is the most energy-consuming behavior for soaring seabirds (Sakamoto et al., 2013; Shaffer et al., 2001; Weimerskirch et al., 2000). Previous studies have ascribed this large energy expenditure to the vigorous flapping required for take-off (Clay et al., 2020; Sato et al., 2009; Shaffer et al., 2001). Indeed, the continuous flapping behavior is rare in cruising flight and was even recorded after the running phase of take-off in this study. However, we suggest the running behavior is also a large cost in take-off because albatross have to reach a fast initial speed to lift off the sea surface by rapidly moving their hindlimbs for up to ~ 10 s in unfavorable conditions (as demonstrated in this study).

We provide a first attempt at detecting the running signal of taking-off seabirds and build a relatively simple algorithm (which can be easily applied to other species) using lateral acceleration. The running duration may increase or decrease by approximately 0.5 s depending on the configure of the algorithm, such as smoothing parameters and threshold values. However, we focused on the relative changes in the running behavior in association with wind speed and wave height, and the error of the absolute value is not a serious problem.

The flapping characteristics during the running phase were also researched. The dorsoventral acceleration signal fluctuates during the running phase making it difficult to identify each flapping signal, even after applying the band-pass filter. Therefore, counting the number of flaps immediately after the running phase was the only reliable parameter for evaluating flapping effort. It was difficult to confirm the frequency of the running phase was derived from flapping behavior. Sato et al. (2009) (Sato et al., 2009) reported that the flapping frequency of wandering albatrosses at the moment of take-off is higher (2.9 ~ 3.4 Hz) than that of cruising flight (2.5 ~ 2.7 Hz). In our study, the flapping frequency after the running phase was not as high as Sato et al. (2009) reported. Therefore, it is likely that the

259   wandering albatross undertake high frequency flapping only during the running phase, as

260   they lift off the sea surface. After lift-off (i.e., the running phase is finished), the wandering

261   albatross continue with a moderate flapping frequency until they reach a certain degree of

262   flight stability. Simultaneous video records of flapping and running motions with acceleration

263   records are required to separate the parameter estimates.

264   In-depth studies on seabird take-offs are just beginning with the aid of miniaturized

animal-borne recorders with the main aim of understanding how seabirds flap their wings (12,

15). However, in land birds (e.g., finches and doves) take-off requires a large contribution by

hindlimbs (Provini et al., 2012) and the role of the hindlimb in take-off kinematics is as

important as that of the wing (Provini and Abourachid, 2018). Therefore, it is highly likely that

seabird take-offs also require a great contribution by the hindlimbs. Further seabird hindlimb

research is required. This study provides the basic characteristics of the running behavior of

wandering albatross, running duration, and running speed.

Take-off effort with environmental conditions

274   The first finding is that wandering albatross can take-off in a variety of environmental

conditions (wind speed: 0.7 ~ 15.4 m/s, wave height: 1.6 ~ 6.4 m). A previous study on

wandering albatross identified the transition state from resting to flying tended to increase as

the wind speed increased (Clay et al., 2020). Our results found some take-offs were

performed under weak wind (2 – 4 m/s) conditions, suggesting wind speed is not the only

parameter influencing flight decisions of wandering albatross, and wave height should be

included in future studies.

281   The results showed that the running and flapping behavior tends to decrease as the

wind or wave conditions increase. Running duration decreases as either the wind speed or

the wave height increases and peaks when both the wind and waves are weak. The same

trends existed in running speed and flapping number. Although optimum statistical models

for each take-off parameter are determined using the AIC value, some models provide

similar results to this model. For instance, the AIC variation in running speed between the

best model and the second-lowest AIC model was only 0.27. However, both models

contained wind speed and wave height as the explanatory variables, similar to the other

take-off parameters, except flapping frequency. The purpose of building a linear model was

to clarify whether the effects of wind and waves are independent. As long as both wind

speed and wave height were included as explanatory variable in the model it reduced the

volume of running and flapping behavior. Therefore, we can conclude that both strong winds

and high waves aid wandering albatross take-offs. The flapping frequency after the running

phase was the only parameter that did not correlate with wave height (as identified using the

LMMs model). However, we assume the flapping frequency during the running phase is

more important. Future research needs to investigate the effects of wave height as

wandering albatross need to climb up or run down the wave slope. Therefore, the flapping

frequency during the running phase should be highly influenced by wave height.

Contribution of strong wind and high waves to seabird take-off

The reduced running behavior and flapping times under strong wind conditions is

simply described by the mechanism of the lift force which has been predicted by previous

studies (Clay et al., 2020; Kogure et al., 2016). Seabirds need to gain lift force before take-off,

and the volume of force is proportional to the square of the relative speed of the wings to the

surrounding air (air speed) (Vogel, 1983). It has been anecdotally suggested that seabirds

take-off into the wind (i.e., headwind), and stronger winds can produce a sufficiently large lift

even before the ground speed of the seabird reaches the value required for flight. As a

partial demonstration of this theory, a study on the European shag (*Gulosus aristotelis*)

(Kogure et al., 2016) found the take-off direction was significantly biased towards headwinds.

Regarding soaring seabirds, only one study (Clay et al., 2020) on wandering albatrosses has

confirmed a bias in take-off direction with wind direction. However, the authors acknowledge

the limits in the climate model and GPS sampling resolutions and recognized the unreliability

of small-scale responses to in situ variation in the atmosphere. Our study reveals wandering

albatross significantly tend to take-off into the wind, using robust fine scale data estimated

from the flight records of wandering albatross. Our data is reliable as the empirical value

provided is actually experienced by the albatross. Furthermore, by quantitatively evaluating

the flapping and running effort, we demonstrate the theory of effortless take-offs by soaring

seabirds in stronger wind conditions.

The mean air speed of wandering albatrosses at the end of the running phase was

close to the average flight speed (approximately 15 m/s) (Weimerskirch, 2002), indicating

that wandering albatross gain sufficient lift at the end of the running phase and efficiently

utilize ocean wind. Wind speed varies with altitude, therefore the wind blowing on the ocean

surface must be smaller than the values estimated from the flight records of wandering

albatross as albatross usually fly 3 – 12 m above the ocean surface (Pennycuick, 1982).

Therefore, the calculated air speed is probably an overestimate when compared with the

ocean surface. To compensate for the insufficient lift force gained during the running phase,

wandering albatross flap their wings several times after the running phase. Therefore, the
flapping number in weak wind conditions can exceed dozens before reaching stable flight.

The most important finding of this study is that the take-off effort estimated by the
running behavior and number of flaps decreased not only with stronger winds but also with
higher waves. While the role of ocean wind on flying seabirds has been well described
(Pennycuick, 2008), how ocean waves influence the flight of seabirds remains largely
unknown. However, many studies have reported the characteristic flight of soaring seabirds
by tracking the ocean wave surface over long distances (Pennycuick, 2008, 1982;
Richardson, 2011; Stokes and Lucas, 2021), which even occurs in weak or no wind
conditions. Seabirds seem to be aided by atmospheric forces above the slope-like wave
topography, the flight method using the shape of waves is called wave-slope soaring
(Richardson, 2011). It is well recognized that air flows occur above ocean waves (Bousquet
et al., 2017; Buckley and Veron, 2016). Richardson (2011) (Richardson, 2011) described the
theoretical model of wave-slope soaring, where the flight mechanism of albatross are a
combination of both dynamic soaring, which uses vertical wind shear above the ocean
surface (~ 15 m), and wave-slope soaring, which uses the updraft caused by the wave
topography. Thus, seabirds can continue to soar in weak wind conditions. Further
mathematical analysis reveal the wave-induced updraft (even in windless conditions) can
provide 60% of the transportation cost of a brown pelican (*Pelecanus occidentalis*, 2 ~ 3 kg),
which is a wave-slope soarer (Stokes and Lucas, 2021). Thus, it is possible that the take-off
effort by wandering albatross is also reduced by high waves. While qualitative field
observations and mathematical demonstrations provide the only previous research on the
role of waves on soaring seabirds, we experimentally demonstrated that ocean waves aid
the most energy-consuming behavior, take-off. This finding helps future discussions on the
oceanic topography mechanisms affecting seabird flight.

The mechanism by which high waves aid wandering albatross take-off is not clear. It
is difficult to conclude a certain updraft is producing additional lift for wandering albatrosses,
it is possible there are other unresolved mechanisms. It is also possible that a rough
topographic surface provides a favorable bump, like a slope or cliff to jump off into the air.
Our results were restricted to wave height as the parameter of the ocean surface. Future
research involving ocean surface steepness or wave frequency components will reveal the
detailed mechanism of how waves make seabird take-off easier. In particular, ocean surface
topography heavily relies on whether the dominant wave component is due to a swell (low-
frequency waves propagated from a distance) or wind waves (high-frequency waves

generated by local wind); moreover, this topography affects the wind pattern on the sea

surface.

We revealed how the take-off effort of wandering albatross change in various

oceanic conditions. As take-off is one of the most energy-consuming behaviors that can

dominate the total energy expenditure of a wandering albatross journey, this data will be a

great help for considering how climate changes can alter the life of albatrosses. Future

research, especially on albatross, should quantitatively evaluate the energy consumption of

take-off with the wind and wave conditions. Currently, there is no major barrier to

accomplishing this goal, it would require utilizing motion records to estimate the surrounding

environment with additional methods to estimate energy consumption, such as cardiograms.

Recognizing the negative effect of the changing oceanic environment on seabirds (Sydeman

et al., 2015), revealing the direct small-scale mechanisms of environmental factors (such as

wind, wave, tide, current, and sea surface temperature) effects on animal behavior,

especially in take-off is urgently required. The concept of estimating the surrounding

environment using motion records is a novel solution with great potential to unravel the small

spatiotemporal uncertainties in seabird research.

Materials and Methods

Field experiment

The recorders, Ninja-scan (Little Leonardo, Tokyo, Japan), record triaxial

acceleration at a very high time resolution (100 Hz). Ninja-scan also records 3D GPS

positions (5 Hz), Doppler velocity (5 Hz), temperature (6 Hz), pressure (6 Hz),

geomagnetism (6 Hz), and angular velocity (100 Hz). There are two types of Ninja-scans

with different battery masses (Naruoka et al., 2021). Small Ninja-scans weighed 28 g,

which is below 1% of the body mass of wandering albatross are expected to record for 7

h. Large Ninja-scans weighed 91 g, which corresponds to approximately 1% of the body

mas of a wandering albatross are expected to record for 65 h.

Ninja scans were attached to breeding wandering albatross at Possession Island,

Crozet archipelago (46°25 S, 51°44 E) in the South Indian Ocean in 2019 and 2020. In

2019, 12 small Ninja scans were attached (in tandem) to 6 individuals. On each bird, one

recorder had a delay timer so that the recording period of the two recorders did not

overlap. Additionally, 15 birds had individual Ninja-scans attached, of which 8 were small

Ninja-scans and 7 were large Ninja scans. In 2020, 10 small Ninja-scans were attached

in tandem to 5 individuals. Additionally, 19 birds had individual Ninja-scans attached, of

which 7 were small Ninja-scans and 12 were large Ninja-scans. In summary, 21 and 24

wandering albatrosses were tagged in 2019 and 2020, respectively. All experiments

were performed from late January to early March of each year, which corresponds to the

incubation period of wandering albatross. Recorders were attached to the back of each

bird with waterproof tape (Tesa, Hamburg, Germany) and glue (Loctite; Henkel,

Dusseldorf, Germany). All recorders were retrieved within 35 days. One small Ninja-scan

which had been attached in isolation in 2020 did not work correctly. The effects of the

attached recorders on wandering albatross were previously assessed (Barbraud and

Weimerskirch, 2012; Phillips et al., 2003) and revealed that small recorders (less than

3% of their body mass) do not negatively impact breeding or foraging behaviors. The

experiment was conducted as part of Program 109 of the Institut Polaire Paul Emile

Victor with permission from the Préfet des Terrs Australes et Antarctiques Françaises,

France.

Take-off identification

First, data recorded on the colony island was eliminated based on the GPS
position. Then, take-off was determined using the absolute value of the GPS horizontal
velocity. When wandering albatross float on the sea surface (i.e., before take-off), a
relatively low speed which is generally below 2.5 m/s, is recorded, while the flying speed
exceeds 5 m/s (Weimerskirch, 2002). Take-off was defined as the moment when the
horizontal speed exceeds 4 m/s and rises to a higher speed. The soaring (flying) speed
occasionally meets this criteria. Therefore, the horizontal speed was smoothed using the
moving average (20 points : 0.4 s). If the horizontal speed crossed the 4 m/s line several
times within a short period, they were classed as take-offs for very short flights and were
not used in our investigation. Therefore, we selected only take-offs that included over 30
s of floating followed by over 30 s of flying.
421
Wind estimation
Yonehara et al. (2016) (Yonehara et al., 2016) proposed estimating the wind
speed and direction of seabird flight paths using the sinusoidal curve relationship
between the flight speed and flight direction. When seabirds fly in the air, their flight
speed against the ground (ground speed) is mainly affected by the wind speed, which is
maximized in tail winds and minimized in head winds. The maximum speed is the sum of
the flight speed against air (air speed) and wind speed, whereas the minimum speed is
the difference between the air speed and wind speed. The relationship between the flight
speed (ground speed) and flight direction recorded by the GPS are fitted using a
sinusoidal curve (Shimatani et al., 2012). We followed the methodology in Yonehara et
al. (2016) to collate the flight speed V and flight direction θ data for 5 min after take-off
and the curve was fitted using the following equation:

$$V = V_a + V_w \cos(\theta + \phi_w)$$

where V_a is the air speed, V_w is the wind speed, and ϕ_w is the wind direction. Ten
seconds immediately after the take-off moment was not included in the estimation.
Following Yonehara et al. (2016), the AIC of the sinusoidal fitting was compared to the
linear fitting with a fixed slope of zero. When the AIC difference between the linear and
sinusoidal fitting was below two, the estimated results were considered unreliable and
discarded. Wind speeds and directions were not calculated when take-offs were not
followed by over five minutes of flight. The sinusoidal curves were performed using Igor
Pro version 8.04 (Wavemetrics, Portland, OR, USA).

Wave estimation

The ocean wave properties experienced by seabirds before take-off were estimated by analyzing the floating motion at the sea surface (Uesaka et al., 2022). The wave height was estimated from the vertical GPS displacement records before take-off. The estimate requires sufficiently long records of vertical displacement. Therefore, the wave height was not calculated for take-offs that did not follow a surface floating time of over 15 min. The sampling period of 15 min ensured the reliability of the wave statistics (Whitford et al., 2001) and provided a large volume of estimated wave data. The estimate did not include the ten seconds before the detected take-off moment. We followed the methodology of Uesaka et al. (2022) (Uesaka et al., 2022). The vertical GPS displacement records were high-pass filtered using a cutting-off frequency of 0.07 Hz to eliminate the GPS derived error (Olynik et al., 2002). We separated the time series record of the vertical displacement into individual waves by applying the zero-up-crossing method. The mean wave height of the highest third of all individual waves was calculated to provide the significant wave height, which is the most widely used statistical wave parameter (Whitford et al., 2001).

Sea surface running by seabirds

Many procellariiformes require a running phase before take-off from the sea surface (Sato et al., 2009). However, studies using accelerometers have not focused on the acceleration signal of this behavior. Surface running involves asymmetrical leg movements. Therefore, the lateral acceleration obtained from the recorder (attached to the back of the seabirds) provides signals derived from the running motion (Fig. 6). We confirmed that running signals appear in the lateral acceleration records at the moment of take-off by streaked shearwaters (*Calonectris leucomelas*), which are phylogenetically like wandering albatross (see Supplementary Information Text and Fig. S6).

To explore the running duration of wandering albatross, we built an algorithm to detect the running phase from the lateral acceleration around take-off. The lateral acceleration signal is composed of a dominant component (0.25 ~ 0.4 s) and a high-frequency fluctuation component (< 0.2 s period). Although the dominant component is the lateral movement derived from surface running, the flapping period of wandering albatross appears around this period (0.3 ~ 0.4 s). The flapping behavior is laterally

symmetrical and does not appear in the lateral acceleration records. However, this is not

always the case, when (occasionally) recorders are attached to the back of the seabird

in a slightly tilted position. To avoid confusion between running and flapping behavior, a

high-frequency fluctuation component in the lateral acceleration signal was used to

detect the running phase. A band-pass filter was designed to extract the high-frequency

fluctuation component from the acceleration records, and then the variance per unit time

(0.6 s) was calculated at each point. Running phase was defined as when the variance

of the acceleration exceeded the threshold value (2% of the peak value). This algorithm

reasonably detects the running phase regardless of the running duration. However, if

there is a signal gap in the middle of the running phase, the algorithm regards the gap as

the end of the running phase, underestimating the running duration. However, these

cases are rare, and we assume it does not affect our evaluation of the running

characteristics of wandering albatross. We also calculated the horizontal speed at the

end of the running phase and take-off direction using the GPS velocity. The take-off

direction was defined as the vectoral average direction during the running phase. We

also calculated the cruising direction which was defined as the moving direction 5 min

after take-off. All procedures were performed using Igor Pro version 8.04 (Wavemetrics,

Portland, OR, USA).

Flapping behavior after the running phase

Dorsoventral acceleration records include signals derived from seabird wing

flapping behavior (Fig. 6). The flapping signals during the running phase fluctuate, which

is assumed to be caused by the leg-derived dorsoventral motion. This caused the

flapping data to be unclear in identifying the flapping number and frequency. Therefore,

we only focused on the wing flapping signals after the running phase. A bandpass filter

extracted the clearest flapping signals (1.8 ~ 4.0 Hz). The number of continuous flapping

signals after the running phase were counted. The flapping period of the wandering

albatross is approximately 0.3 ~ 0.4 s, therefore we defined the end point when the

interval of the flapping exceeded 0.5 s. The flapping frequency after the running phase

was calculated using the spectral peak value of the continuous wavelet-transformed

dorsoventral acceleration. All procedures were performed using Igor Pro version 8.04

(Wavemetrics, Portland, OR, USA).

Comparison of the take-off parameters with environmental conditions

The wind directional bias of the take-off direction was tested using the v-test

(modified Rayleigh test). The air speed V_a at the end of the running phase was estimated

using the following equation based on the parameters obtained from this study:

$$V_a = V_r + V_w \cos(\theta_t - \phi_w)$$

Where V_r is the running speed at the end of the running phase, V_w is the wind speed, θ_t

is the take-off direction, and ϕ_w is the wind direction. The effects of wind speed and

wave height on each take-off parameter (running duration, running speed, flapping

number, and flapping frequency) were evaluated using linear mixed models (LMMs) with

individuals treated as random effects. To identify significance levels, the models were

compared to null models based on the AIC value.

To evaluate the combined effect of wind and waves, we categorized take-off

conditions into four groups, "WL conditions," "WH conditions," "SL conditions," and "SH

conditions". The mean values of each take-off parameter were compared between the

four groups. Furthermore, the independent effects of wind and waves on take-off

parameters were evaluated using LMMs, including wind speed, wave height, and their

interaction as explanatory parameters with individuals as random effects. VIF was also

calculated before the LMMs analysis to assess whether the multicollinearity effect could

be dismissed. The V-test was performed using Igor Pro version 8.04 (Wavemetrics,

Portland, OR, USA). The LMMs calculations were performed using the PypeR package

in Python 3.0.

Acknowledgments

We thank Yoshinari Yonehara, Julien Collet, Timothée Poupart, and all the members of the research station in the Crozet Islands for their support during the fieldwork. We appreciate the fruitful comments of Kagari Aoki, Chihiro Kinoshita, and Aran Garrod. We are grateful to Taichi Sakamoto and Takashi Mukai (ATTACCATO) for providing customized rechargeable Li-ion batteries for the Ninja-scans. We also thank Michihiko Suzuki and Koichiro Ikeda (Little Leonardo) for molding the Ninja-scans. The field studies were financially supported by a research project entitled "Cyber ocean: next generation navigation system on the sea" funded by the CREST program (JPMJCR1685) of Japan Science and Technology Agency; Grants-in-Aid for Scientific Research from JSPS (17H00776 to K. Sato); the Tohoku Ecosystem-Associated Marine Science; and Institut Polaire Français Paul-Émile Victor Program 109.

References

Adams NJ, Navarro RA. 2005. Foraging of a coastal seabird: Flight patterns and movements of breeding Cape gannets Morus capensis. *African J Mar Sci* **27**:239–248. doi:10.2989/18142320509504082

Ardhuin F, Stopa JE, Chapron B, Collard F, Husson R, Jensen RE, Johannessen J, Mouche A, Passaro M, Quartly GD, Swail V, Young I. 2019. Observing sea states. *Front Mar Sci* **6**:1–29. doi:10.3389/fmars.2019.00124

Barbraud C, Weimerskirch H. 2012. Assessing the effect of satellite transmitters on the demography of the Wandering Albatross Diomedea exulans. *J Ornithol* **153**:375–383. doi:10.1007/s10336-011-0752-8

Bousquet GD, Triantafyllou MS, Slotine JJE. 2017. Optimal dynamic soaring consists of successive shallow arcs. *J R Soc Interface* **14**. doi:10.1098/rsif.2017.0496

Buckley MP, Veron F. 2016. Structure of the Airflow above Surface Waves. *J Phys Oceanogr* **46**:1377–1397. doi:10.1175/JPO-D-15-0135.1

Carta JA, Ramírez P, Velázquez S. 2009. A review of wind speed probability distributions used in wind energy analysis. Case studies in the Canary Islands. *Renew Sustain Energy Rev* **13**:933–955. doi:10.1016/j.rser.2008.05.005

Charrassin JB, Hindell M, Rintoul SR, Roquet F, Sokolov S, Biuw M, Costa D, Boehme L, Lovell P, Coleman R, Timmermann R, Meijers A, Meredith M, Park YH, Bailleul F, Goebel M, Tremblay Y, Bost CA, McMahon CR, Field IC, Fedak MA, Guinet C. 2008. Southern Ocean

frontal structure and sea-ice formation rates revealed by elephant seals. *Proc Natl Acad Sci*

U S A **105**:11634–11639. doi:10.1073/pnas.0800790105

Charrassin JB, Park YH, Maho Y Le, Bost CA. 2002. Penguins as oceanographers unravel

hidden mechanisms of marine productivity. *Ecol Lett* **5**:317–319. doi:10.1046/j.1461-

0248.2002.00341.x

Clairbaux M, Mathewson P, Porter W, Fort J, Strøm H, Moe B, Fauchald P, Descamps S,

Helgason HH, Bråthen VS, Merkel B, Anker-Nilssen T, Bringsvor IS, Chastel O,

Christensen-Dalsgaard S, Danielsen J, Daunt F, Dehnhard N, Erikstad KE, Ezhov A, Gavrilo

M, Krasnov Y, Langset M, Lorentsen SH, Newell M, Olsen B, Reiertsen TK, Systad GH,

Thórarinsson TL, Baran M, Diamond T, Fayet AL, Fitzsimmons MG, Frederiksen M, Gilchrist

HG, Guilford T, Huffeldt NP, Jessopp M, Johansen KL, Kouwenberg AL, Linnebjerg JF,

Major HL, Tranquilla LMF, Mallory M, Merkel FR, Montevecchi W, Mosbech A, Petersen A,

Grémillet D. 2021. North Atlantic winter cyclones starve seabirds. *Curr Biol* **31**:3964-

3971.e3. doi:10.1016/j.cub.2021.06.059

Clay TA, Joo R, Weimerskirch H, Phillips RA, den Ouden O, Basille M, Clusella-Trullas S, Assink

JD, Patrick SC. 2020. Sex-specific effects of wind on the flight decisions of a sexually

dimorphic soaring bird. *J Anim Ecol* **89**:1811–1823. doi:10.1111/1365-2656.13267

Doi T, Storto A, Fukuoka T, Suganuma H, Sato K. 2019. Impacts of Temperature Measurements

From Sea Turtles on Seasonal Prediction Around the Arafura Sea. *Front Mar Sci* **6**:1–11.

doi:10.3389/fmars.2019.00719

Dormann CF, Elith J, Bacher S, Buchmann C, Carl G, Carré G, Marquéz JRG, Gruber B,

Lafourcade B, Leitão PJ, Münkemüller T, Mcclean C, Osborne PE, Reineking B, Schröder

B, Skidmore AK, Zurell D, Lautenbach S. 2013. Collinearity: A review of methods to deal

with it and a simulation study evaluating their performance. *Ecography (Cop)* **36**:27–46.

doi:10.1111/j.1600-0587.2012.07348.x

Dunn EK. 1973. Changes in Fishing Ability of Terns associated with Windspeed and Sea Surface

Conditions. *Nature* **244**:520–521. doi:10.1038/244520a0

Ferreira JA, Guedes Soares C. 2000. Modelling distributions of significant wave height. *Coast*

Eng **40**:361–374. doi:10.1016/S0378-3839(00)00018-1

Goto Y, Yoda K, Sato K. 2017. Asymmetry hidden in birds' tracks reveals wind, heading, and

orientation ability over the ocean. *Sci Adv* **3**:1–10. doi:10.1126/sciadv.1700097

Haney JC, Fristrup KM, Lee DS. 1992. Geometry of visual recruitment by seabirds to ephemeral

foraging flocks. *Ornis Scand* **23**:49–62. doi:10.2307/3676427

Harcourt R, Sequeira AMM, Zhang X, Roquet F, Komatsu K, Heupel M, McMahon C, Whoriskey

F, Meekan M, Carroll G, Brodie S, Simpfendorfer C, Hindell M, Jonsen I, Costa DP, Block B,

Muelbert M, Woodward B, Weise M, Aarestrup K, Biuw M, Boehme L, Bograd SJ, Cazau D,

Charrassin JB, Cooke SJ, Cowley P, de Bruyn PJN, Jeanniard du Dot T, Duarte C, Eguíluz

VM, Ferreira LC, Fernández-Gracia J, Goetz K, Goto Y, Guinet C, Hammill M, Hays GC,

Hazen EL, Hückstädt LA, Huveneers C, Iverson S, Jaaman SA, Kittiwattanawong K, Kovacs

KM, Lydersen C, Moltmann T, Naruoka M, Phillips L, Picard B, Queiroz N, Reverdin G, Sato

K, Sims DW, Thorstad EB, Thums M, Treasure AM, Trites AW, Williams GD, Yonehara Y,

Fedak MA. 2019. Animal-borne telemetry: An integral component of the ocean observing

toolkit. *Front Mar Sci* **6**. doi:10.3389/fmars.2019.00326

Kogure Y, Sato K, Watanuki Y, Wanless S, Daunt F. 2016. European shags optimize their flight

behavior according to wind conditions. *J Exp Biol* **219**:311–318. doi:10.1242/jeb.131441

McMahon CR, Roquet F, Baudel S, Belbeoch M, Bestley S, Blight C, Boehme L, Carse F, Costa

DP, Fedak MA, Guinet C, Harcourt R, Heslop E, Hindell MA, Hoenner X, Holland K, Holland

M, Jaine FRA, Jeanniard du Dot T, Jonsen I, Keates TR, Kovacs KM, Labrousse S, Lovell

P, Lydersen C, March D, Mazloff M, McKinzie MK, Muelbert MMC, O'Brien K, Phillips L,

Portela E, Pye J, Rintoul S, Sato K, Sequeira AMM, Simmons SE, Tsontos VM, Turpin V,

van Wijk E, Vo D, Wege M, Whoriskey FG, Wilson K, Woodward B. 2021. Animal Borne

Ocean Sensors – AniBOS – An Essential Component of the Global Ocean Observing

System. *Front Mar Sci* **8**:1–21. doi:10.3389/fmars.2021.751840

Naruoka M, Goto Y, Weimerskirch H, Mukai T, Sakamoto T, Sakamoto KQ, Sato K. 2021.

Application of Inertial and GNSS Integrated Navigation to Seabird Biologging. *J Robot*

Mechatronics **33**:526–536. doi:10.20965/jrm.2021.p0526

Nevitt GA, Losekoot M, Weimerskirch H. 2008. Evidence for olfactory search in wandering

albatross, Diomedea exulans. *Proc Natl Acad Sci U S A* **105**:4576–4581.

doi:10.1073/pnas.0709047105

Nicholls DG, Murray MD, Butcher E, Moors P. 1997. Weather systems determine the non-

breeding distribution of Wandering Albatrosses over southern oceans. *Emu* **97**:240–244.

doi:10.1071/mu97031

Norberg RA, Norberg UM. 1971. Take-Off, Landing, and Flight Speed during Fishing Flights of

Gavia stellata (Pont.). *Ornis Scand* **2**:55–67. doi:10.2307/3676239

Olynik M, Petovello MG, Cannon ME, Lachapelle G. 2002. Temporal Variability of GPS Error

Sources and Their Effect on Relative Positioning AccuracyProceedings of the 2002 National

Technical Meeting of The Institute of Navigation. 2002. pp. 877–888.

Padget O, Stanley G, Willis JK, Fayet AL, Bond S, Maurice L, Shoji A, Dean B, Kirk H, Juarez-

Martinez I, Freeman R, Bolton M, Guilford T. 2019. Shearwaters know the direction and

distance home but fail to encode intervening obstacles after free-ranging foraging trips. *Proc*

Natl Acad Sci U S A **116**:21629–21633. doi:10.1073/pnas.1903829116

Pennycuick C. 2008. Modelling the Flying Bird, Academic Press. New York.

Pennycuick CJ. 1982. The flight of petrels and albatrosses (procellariiformes), observed in South
Georgia and its vicinity. *Philos Trans R Soc London B, Biol Sci* **300**:75–106.
doi:10.1098/rstb.1982.0158
Phillips RA, Xavier JC, Croxall JP. 2003. Effects of Satellite Transmitters on Albatrosses and
Petrels. *Auk* **120**:1082–1090. doi:10.1093/auk/120.4.1082
Provini P, Abourachid A. 2018. Whole-body 3D kinematics of bird take-off: key role of the legs to
propel the trunk. *Sci Nat* **105**. doi:10.1007/s00114-017-1535-8
Provini P, Tobalske BW, Crandell KE, Abourachid A. 2012. Transition from leg to wing forces
during take-off in birds. *J Exp Biol* **215**:4115–4124. doi:10.1242/jeb.074484
Rayleigh. 1883. THE SOARING OF BIRDS. *Nature* **27**:534–535. doi:10.1038/027534a0
Richardson PL. 2011. How do albatrosses fly around the world without flapping their wings? *Prog
Oceanogr* **88**:46–58. doi:10.1016/j.pocean.2010.08.001
Roquet F, Williams G, Hindell MA, Harcourt R, McMahon C, Guinet C, Charrassin JB, Reverdin
G, Boehme L, Lovell P, Fedak M. 2014. A Southern Indian Ocean database of hydrographic
profiles obtained with instrumented elephant seals. *Sci Data* **1**:1–10.
doi:10.1038/sdata.2014.28
Sakamoto KQ, Takahashi A, Iwata T, Yamamoto T, Yamamoto M, Trathan PN. 2013. Heart rate
and estimated energy expenditure of flapping and gliding in black-browed albatrosses. *J
Exp Biol* **216**:3175–3182. doi:10.1242/jeb.079905
Sánchez-Román A, Gómez-Navarro L, Fablet R, Oro D, Mason E, Arcos JM, Ruiz S, Pascual A.
2019. Rafting behaviour of seabirds as a proxy to describe surface ocean currents in the
Balearic Sea. *Sci Rep* **9**:1–11. doi:10.1038/s41598-018-36819-w
Sato K, Sakamoto KQ, Watanuki Y, Takahashi A, Katsumata N, Bost CA, Weimerskirch H. 2009.
Scaling of soaring seabirds and implications for flight abilities of giant pterosaurs. *PLoS One*
4:1–6. doi:10.1371/journal.pone.0005400
Shaffer SA, Costa DP, Weimerskirch H. 2001. Behavioural factors affecting foraging effort of
breeding wandering albatrosses. *J Anim Ecol* **70**:864–874. doi:10.1046/j.0021-
8790.2001.00548.x
Shimatani IK, Yoda K, Katsumata N, Sato K. 2012. Toward the Quantification of a Conceptual
Framework for Movement Ecology Using Circular Statistical Modeling. *PLoS One* **7**:e50309.
doi:10.1371/journal.pone.0050309
Stokes IA, Lucas AJ. 2021. Wave-slope soaring of the brown pelican. *Mov Ecol* **9**:1–13.
doi:10.1186/s40462-021-00247-9
Suryan RM, Anderson DJ, Shaffer SA, Roby DD, Tremblay Y, Costa DP, Sievert PR, Sato F,
Ozaki K, Balogh GR, Nakamura N. 2008. Wind, waves, and wing loading: Morphological

specialization may limit range expansion of endangered albatrosses. *PLoS One* **3**:e4016.

doi:10.1371/journal.pone.0004016

Sydeman WJ, Poloczanska E, Reed TE, Thompson SA. 2015. Climate change and marine

vertebrates. *Science* **350**:772–777. doi:10.1126/science.aac9874

Uesaka L, Goto Y, Yonehara Y, Komatsu K, Naruoka M, Weimerskirch H, Sato K, Sakamoto KQ.

2022. Ocean wave observation utilizing motion records of seabirds. *Prog Oceanogr*

200:102713. doi:10.1016/j.pocean.2021.102713

Villas Bôas AB, Ardhuin F, Ayet A, Bourassa MA, Brandt P, Chapron B, Cornuelle BD, Farrar JT,

Fewings MR, Fox-Kemper B, Gille ST, Gommenginger C, Heimbach P, Hell MC, Li Q,

Mazloff MR, Merrifield ST, Mouche A, Rio MH, Rodriguez E, Shutler JD, Subramanian AC,

Terrill EJ, Tsamados M, Ubelmann C, Sebille E van. 2019. Integrated observations of global

surface winds, currents, and waves: Requirements and challenges for the next decade.

Front Mar Sci **6**:1–34. doi:10.3389/fmars.2019.00425

Vogel S. 1983. Life in Moving Fluids. Princeton University Press. doi:10.1515/9780691212975

Weimerskirch H. 2002. GPS Tracking of Foraging Albatrosses. *Science* **295**:1259–1259.

doi:10.1126/science.1068034

Weimerskirch H, Delord K, Guitteaud A, Phillips RA, Pinet P. 2015. Extreme variation in migration

strategies between and within wandering albatross populations during their sabbatical year

and their fitness consequences. *Sci Rep* **5**:8853. doi:10.1038/srep08853

Weimerskirch H, Guionnet T, Martin J, Shaffer SA, Costa DP. 2000. Fast and fuel efficient?

Optimal use of wind by flying albatrosses. *Proc R Soc B Biol Sci* **267**:1869–1874.

doi:10.1098/rspb.2000.1223

Weimerskirch H, Louzao M, de Grissac S, Delord K. 2012. Changes in Wind Pattern Alter

Albatross Distribution and Life-History Traits. *Science* **335**:211–214.

doi:10.1126/science.1210270

Weimerskirch H, Prudor A. 2019. Cyclone avoidance behaviour by foraging seabirds. *Sci Rep*

9:1–9. doi:10.1038/s41598-019-41481-x

Whitford DJ, Waters JK, Vieira MEC. 2001. Teaching time-series analysis. II. Wave height and

water surface elevation probability distributions. *Am J Phys* **69**:497–504.

doi:10.1119/1.1331301

Wilmers CC, Nickel B, Bryce CM, Smith JA, Wheat RE, Yovovich V, Hebblewhite M. 2015. The

golden age of bio-logging: How animal-borne sensors are advancing the frontiers of

ecology. *Ecology* **96**:1741–1753. doi:10.1890/14-1401.1

Yamamoto T, Kohno H, Mizutani A, Sato H, Yamagishi H, Fujii Y, Murakoshi M, Yoda K. 2017.

Effect of wind on the flight of brown booby fledglings. *Ornithol Sci* **16**:17–22.

doi:10.2326/osj.16.17

Yoda K, Shiomi K, Sato K. 2014. Foraging spots of streaked shearwaters in relation to ocean
surface currents as identified using their drift movements. *Prog Oceanogr* **122**:54–64.
doi:10.1016/j.pocean.2013.12.002
Yonehara Y, Goto Y, Yoda K, Watanuki Y, Young LC, Weimerskirch H, Bost CA, Sato K. 2016.
Flight paths of seabirds soaring over the ocean surface enable measurement of fine-scale
wind speed and direction. *Proc Natl Acad Sci U S A* **113**:9039–9044.
doi:10.1073/pnas.1523853113
713
714

Fig. 1. Conceptual framework of the study estimating environmental condition experienced by studied individual.

Fig. 2. Effect of wind direction on take-off of wandering albatrosses. Relative take-off direction to the wind direction (black circles, n = 427) significantly distributed around 0° (head-wind), whereas the cruising direction relative to the wind (gray x-mark, n = 427) is not. The radial axis represents the wind speed.

Fig. 3. Environmental effect on take-off. Effort for the take-off (Running duration, Running speed, Flapping number, and Flapping frequency) significantly decreased as wave height and wind speed increased (p < 0.01) except the relationship between flapping frequency and wave height (p = 0.026). Solid line shows the linear regression line determined from the LMMs and the number at the right top corner on each graphs shows the sample sizes.

Fig. 4. (A) Correlation between wind speed and wave height was weak (r = 0.27, n = 185). Bar charts and solid lines written above and right of the scatter plot are normed histograms of wind speed and wave height and curve fitted lines. Based on the peak value of fitted lines scatter plots were divided into 4 groups, WL: weak wind low wave (open square), WH: weak wind high wave (filled square), SL: strong wind low wave (open circle), and SH: strong wind high wave (filled circle). (B) Take-off effort comparison among four groups (a: running duration, b: running speed, c: flapping number, and d: flapping frequency). Cross mark indicate the mean value.

Fig. 5. (A) running duration, (B) running speed, (C) flapping number, and (D) flapping frequency in response to the wave height change under weak wind (dashed line, 2 m/s) and strong wind (solid line, 8 m/s) condition estimated from the result of LMMs. Gray area represents 99% CI.

Fig. 6. Time series data of horizontal speed (top), lateral acceleration (middle), and dorsoventral acceleration (bottom) signals of the wandering albatross at the moment of taking off. Horizontal speed start increasing from the beginning of the take-off. Red square shows the detected running phase based on the variance of the lateral acceleration signal. Red bars show the detected flapping behavior after the running phase based on the dorsoventral acceleration signal. Dorsoventral signal during the running phase is fluctuated, probably due to the shaking body derived from the running motion, and thus it is not easy to judge the existence of flapping behavior.

GPS record

GPS record

Animal-borne recorder

Acceleration record

Wave estimation from drifting motion

Wind estimation from flight path

Take-off effort