Proteomic survey of the DNA damage response in *Caulobacter crescentus* reveals

evidence of post-transcriptional control

Running Title: Survey of post-transcription control upon DNA damage

Tommy F. Tashjian[#], Rilee D. Zeinert[*], Stephen J. Eyles, Peter Chien[#]

Department of Biochemistry and Molecular Biology, University of Massachusetts

Amherst, Amherst, MA, USA

*Current address: Cell Biology and Metabolism Program, Eunice Kennedy Shriver

National Institute of Child Health and Human Development, Bethesda, MD 20892-5430,

USA

#co-corresponding authors:

Tommy Tashjian, Department of Biochemistry and Molecular Biology, University of

Massachusetts Amherst, Life Science Laboratories Room N325, 240 Thatcher Rd,

Amherst MA 01003-9364, (413)545-2310, ttashjian@umass.edu

Peter Chien, Department of Biochemistry and Molecular Biology, University of

Massachusetts Amherst, Life Science Laboratories Room N325, 240 Thatcher Rd,

Amherst MA 01003-9364, (413)545-2310, pchien@umass.edu;

ABSTRACT

The bacterial DNA damage response is a critical, coordinated response to DNA

replication stress. The canonical bacterial DNA damage response, first characterized in

Escherichia coli, is controlled by the global transcriptional regulator LexA and the

recombinase RecA. While genome-wide studies have described how the DNA damage

response is regulated at a transcriptional level, relatively little is known about post-

transcriptional regulation of this response. Here, we perform a proteome-wide survey of

the DNA damage response in *Caulobacter crescentus*. We find that not all changes in

protein abundance during the response to DNA damage are predicted by changes in

transcription. We validate one of these post-transcriptionally regulated candidates to

show its importance to survival of DNA damage. To investigate post-translational control

of the DNA damage response, we perform a similar survey in cells lacking the Lon

protease. This reveals that induction of the DNA damage response at the protein level is

dampened in these strains, consistent with their reduced tolerance to DNA damage.

Finally, proteome-wide stability measurements following damage nominate candidate

Lon substrates that suggest post-translational regulation of the DNA damage response.

IMPORTANCE

The DNA damage response helps bacteria to respond to and potentially survive DNA

damage. The mutagenesis that is induced as part of this response plays a role in bacterial

evolution and is essential to the development and spread of antibiotic resistance.

Understanding how bacteria coordinate their response to DNA damage could help us to

combat this growing threat to human health. While the transcriptional regulation of the

bacterial DNA damage response has been characterized, this study is the first to our

knowledge to compare changes in RNA and protein levels to identify potential targets of

post-transcriptional regulation in response to DNA damage.

INTRODUCTION

The bacterial DNA damage response is tightly linked to the mutagenesis that drives

bacterial evolution, antibiotic resistance development, and the horizontal gene transfer

that spreads resistance from cell to cell. The canonical DNA damage response was first

characterized in *Escherichia coli*. During DNA damage ssDNA accumulates in the cell

and is bound by the cell's main recombinase RecA to form a nucleoprotein filament that

induces the autocleavage of the global transcriptional repressor LexA (1, 2). Cleavage of

the repressor LexA promotes transcription of more than 40 genes related to DNA damage

tolerance and repair (3, 4).

In addition to LexA-mediated transcriptional regulation, the DNA damage response is also

controlled by sRNA, protein-protein interactions, and proteolysis. For example, in *E. coli*,

the toxin TisB is regulated by LexA and by the antitoxin sRNA IstR1 (5, 6). Similarly,

during the *Caulobacter crescentus* DNA damage response expression of the TonB

receptor ChvT is inhibited by the sRNA ChvR (7). Alternatively, the activity of several DNA

damage proteins are regulated by protein-protein interactions (e.g. DinB (8, 9), MutS and

MutL reviewed in (10)). Finally, proteolysis plays a key role in the induction (through

degradation of remnants of cleaved LexA (11)) and shut down of the DNA damage

response (by degrading key proteins at the end of the response, e.g. UmuD' (12) and

UvrA (13)). Given these examples of post-transcriptional regulation for DNA damage, we

reasoned that a proteomics survey of this same stress would be valuable.

Compared to *E. coli*, we have limited knowledge of the DNA damage response in

Caulobacter. This alphaproteobacteria also contains a coordinated DNA damage

response that is controlled by the global transcriptional repressor LexA (14–16). A second

DNA damage-controlled transcriptional repressor DriD that, like LexA, responds to ssDNA

(16, 17), is also involved in the induction of another set of DNA damage response genes

(16). Other than the discovery of ChvR (7) described above, little is known about the role

of post-transcriptional regulation in *Caulobacter's* DNA damage response.

In the first stage of our proteomics study, we surveyed post-transcriptional regulation of

the DNA damage response in *Caulobacter*. As expected, we found a high level of synergy

between the transcriptional regulation of the response and the changes in protein

abundance upon DNA damage. We were also able to identify proteins that appear to be

regulated post-transcriptionally during the DNA damage response and explored their

physiological impact. In the second stage of this study, we examined protein stability

upon DNA damage. As expected after translational shutoff, we detected protein level

decreases in several known proteolysis substrates. In addition, some of the proteins that

were induced upon DNA damage are also candidate proteolysis substrates.

Our data show that not all protein abundance changes during the DNA damage response

are driven by transcriptional control and identify novel *Caulobacter* DNA damage

response factors that are likely regulated post-transcriptionally.

RESULTS

Proteomics identify upregulated DNA damage response factors

To characterize protein abundance changes during the DNA damage response, we first used the sentinel protein RecA to determine the timescale of an acute mitomycin C (MMC) treatment in *Caulobacter* (Figure 1A). Based on this characterization, we compared whole cell proteome abundances using mass spectrometry before and after MMC treatment (timeline described in Figure 1A, full list of results in Table S1 and S2). The post-MMC timepoint was taken at the timepoint that showed the highest RecA concentration by western blot (Figure 1A). As expected, we found several DNA damage response proteins to be more abundant after MMC treatment (RecA, UvrA, RuvB, ImuA, and MmcA; Figure 1B). To characterize our dataset further, we compared changes in protein abundance observed by our proteomics approach to published changes in RNA levels upon MMC treatment (15), to those genes predicted to be regulated by LexA (14), and to potential fitness effects observed by transposon sequencing (Table S3).

Past work has explored transcriptional responses to DNA damage in Caulobacter, including bioinformatic predictions of LexA binding sites (14), RNA-seq during extended chronic damaging conditions (7), and microarray experiments with a variety of damaging conditions (15). Based on consistency with our experimental design we used Modell, et al. as a comparison (15). In this study, transcript levels were monitored using microarrays over several time points after MMC treatment in rich medium (15). To identify genes that are transcriptionally upregulated upon MMC treatment, we used the maximum $\log_2$(Fold Change of RNA) from these timepoints for each gene and treated genes that were induced more than one standard deviation above the mean as "upregulated". This threshold is intentionally inclusive to capture even weakly induced genes, as we are

particularly interested in identifying genes that are not upregulated transcriptionally in our

next analysis step.

This strategy yielded in 615 upregulated genes of which 146 could be identified as

peptides in our proteomics survey. Of these, 43 show significant increase in protein

abundance upon DNA damage (Table S4, $\log_2$(3 hrs/0 hrs) > 0, FDR < 0.05). As

expected, the known DNA damage proteins mentioned above that showed increased

protein abundances following MMC treatment were also significantly upregulated at the

RNA level (Figure 1B). These findings validate that our proteomics approach and data

analysis methods can be used to successfully identify genes upregulated during the DNA

damage response. There are some limitations, as no peptides were identified for a

number of DNA damage response proteins predicted to be upregulated by microarray

((15), e.g. SidA, DidA, DnaE2).

Post-transcriptional control of DNA damage response proteins

We identified 81 proteins that significantly increased in abundance upon DNA damage

($\log_2$(Fold Change in Protein) > 0, FDR < 0.05) and did not show a corresponding

significant increase in RNA level ((15), Table S5). Some of these are known to be involved

in DNA replication (*dnaA*, *dnaQ*) or repair (*radA*) and one is a homolog of the DNA repair

helicase *dinG* (CCNA_02118). In addition, three proteins involved in response to oxidative

stress were found in this list (two superoxide dismutase proteins: CCNA_01650,

CCNA_03672, and a hyperoxide resistance protein: CCNA_00963). None of these genes

are predicted to be transcriptionally regulated by LexA (14) or *Caulobacter's* LexA-

independent DNA damage response [16] but protein abundances indicate that they are

upregulated upon DNA damage.

Two of the 81 genes (CCNA_03496: *yaaA* and CCNA_00276: hypothetical protein) are

of particular interest, as our Tn-seq experiments identified them as potentially important

for survival of MMC (Table S3). CCNA_03496 is a homolog of *E. coli*'s *yaaA* ([18], we will

refer to this gene as *yaaA* henceforth), which is important for regulating iron levels during

oxidative stress conditions [19]. CCNA_00276 is a hypothetical cupin superfamily protein.

Based on published RB-Tnseq results, strains carrying transposon insertions in *yaaA* or

CCNA_00276 have a sensitivity to the DNA damaging agent cisplatin [20]. For these

reasons, we decided to further characterize the connection between these genes and

MMC survival.

Contrary to our expectations, deletion of these genes had no effect on growth of isolated

cultures with MMC (Figure S2A and S2B). To focus on fitness differences, we turned to

a competition assay. We observed that a Δ*yaaA* strain expressing the mKate2 fluorescent

protein is significantly less able than the wild type mKate2-expressing strain to compete

in co-culture with a wild type control expressing the Venus fluorescent protein (Figure

2C). This competitive disadvantage becomes significantly worse in the presence of MMC

(Figure 2C). By contrast, the ΔCCNA_00276 (denoted as Δ276 throughout) does not

show a disadvantage (Figure 2C). To confirm that these results are not an artifact of the

fluorescent strains, we flipped the markers to test Venus-expressing deletion strains

against mKate2-expressing controls and saw similar results (Figure S2C). These data

support our hypothesis that YaaA plays a role in the DNA damage response.

Finally, we note that of the 146 genes that are transcriptionally upregulated upon DNA

damage and were identified in our proteomics, 103 showed no significant increase in

protein level (Table S6). Two of these proteins have known roles in DNA repair (UvrD and

RecN) and one is a homolog of the DNA recombination protein RmuC (CCNA_00272).

An increase in RNA levels without a corresponding increase in protein levels suggests

some level of post-transcriptional regulation.

Identification of factors that are downregulated and important for MMC survival

We next looked at those genes downregulated following MMC damage. We used the

minimum $\log_2$(Fold Change of RNA) from the Laub lab microarray timepoints for each

gene for the transcriptional response (15). To ensure inclusion for even weakly regulated

genes, a transcript that is more than one standard deviation below the mean of these

values was considered "downregulated". With these criteria, we identified 614 genes that

are downregulated upon DNA damage. Of these, 292 were identified in our proteomics

(Figure 3A).

We identified 99 proteins that became significantly less abundant upon DNA damage and

did not show a decrease in RNA abundance (Table S7). These are proteins that may be

post-transcriptionally downregulated during the DNA damage response, and include

known DNA replication factors DNA gyrase A, DNA gyrase B, and topoisomerase I. Our

Tn-seq results identified six of these genes as more likely to contain transposon insertions

upon MMC damage: CCNA_03681: ABC transporter ATP-binding protein, CCNA_00299:

PurD, CCNA_00099: FzlC, CCNA_01034: TonB-dependent outer membrane receptor,

and CCNA_01817: NtrX. We decided to focus on PurD and FzlC for further

characterization as membrane proteins may alter MMC import and NtrX might have

confounding effects because of its role in cell cycle regulation (21, 22).

The gene *purD* (CCNA_00299) encodes phosphoribosylamine-glycine ligase, which

catalyzes the second step of *de novo* purine biosynthesis. *Caulobacter* RB-Tnseq-based

fitness results suggest that transposon insertions in this gene can lead to increased

survival of nalidixic acid (20). While we did not observe any benefit to MMC survival in the

Δ*purD* strain (Figure 3B), we found that a Δ*purD* strain of *Caulobacter* has a relative

stationary phase defect in rich medium not observed during MMC treatment (Figure 3B).

The mutant reaches a maximum density earlier than the wildtype with a final optical

density 14% lower than that of the wild type strain. In the presence of MMC, this difference

is lost, resulting in Δ*purD* strains showing a relative fitness improvement during damage

compared to undamaged conditions, consistent with published Tn-seq results (20) and

our own results (Table S3).

FzlC (CCNA_00099) is a component of the *Caulobacter* cell division machinery that

anchors FtsZ to the membrane (23). Because of the connection between cell division and

DNA repair, we chose to investigate this gene further but found that deletion of *fzlC* did

not have any impact on MMC-sensitivity by growth curve (Figure 3C) or competition

experiment (Figure S3).

Identification of potential Lon substrates induced by the DNA damage response

To investigate the role of post-translational regulation during the DNA damage response,

we performed proteomics in a strain lacking the dispensable Lon protease. By performing

the same proteomics experiment as above in a Δ*lon* strain (full dataset in Table S1 and

S2), we can observe how loss of Lon proteolysis affects protein induction during DNA

damage. Similar to the wild type strain, we monitored RecA levels through induction and

recovery after MMC treatment in Δlon (Figure 4A). We find that the peak induction occurs

at a similar time point in the wild type (Figure 1A) and Δ*lon* strains (Figure 4A) but the

fold-induction of RecA in the Δ*lon* strain is less than half that in the wild type strain. We

reasoned that this was due to upregulation of DNA damage response genes in a Δ*lon*

strain even in the absence of exogenous DNA damaging agents (24). Most of the detected

known DNA damage-related proteins upregulated upon MMC treatment in the wild type

strain are also upregulated in the Δ*lon* strain (See RecA, RuvB, and MmcA in Figure 4B).

The exceptions here are UvrA, which is not significantly induced at the protein level in the

Δ*lon* strain (Figure 4B) and ImuA, which was not identified in the proteomics of the Δ*lon*

strain. In general, the fold change in protein abundance upon DNA damage in the Δ*lon*

strain was reduced compared to the wild type strain (Figure 4C).

To monitor post-translational control during DNA damage, we measured protein

abundances after a translational shutoff. Specifically, we added chloramphenicol to inhibit

translation at the post-MMC treatment timepoint (3 hours in Figure 1A and 5A) and

measured protein abundances 2 hours post-translational shutoff (timeline Figure 5A, full

dataset in Table S1 and S2). For proteins induced as part of the DNA damage response

and degraded by a protease, we expected an increase in protein abundance between the

pre- and post-MMC treatment samples (0 and 3 hours, Figure 5A, left) and a decrease in

protein abundance between the pre- and post-translational shutoff samples (3 and 5

hours, Figure 5A, left). In wild type cells, we see this pattern for the replication initiator

DnaA and IbpA, known substrates of the Lon protease ((25, 26) Figure 5A, right).

In total, we identified 320 proteins that significantly decreased in protein abundance upon

translational shutoff in the wild type strain ($\log_2$(5 hrs/3 hrs) < 0, FDR < 0.05; Table S8).

Twenty-one of these proteins have previously been identified as ClpP substrates by a

ClpP trap in *Caulobacter* (27), including NrdA, DnaX, and CtrA. In addition, other known

ClpXP substrates (e.g., FtsZ, UvrA, LexA) and ClpAP-dependent substrates (e.g., FtsA,

Lon) were identified by these criteria (all mentioned proteins are highlighted in red in

Figure 5B).

We performed the same proteomics experiment during translational shutoff in MMC-

treated Δ*lon* cells to identify Lon substrates induced during DNA damage. Similar to

induction of proteins (Figure 4C), we found that protein loss in the Δ*lon* strain was

dampened compared the wild type strain (Figure S4). As expected from known Lon

substrates DnaA (25) and small heat shock protein IbpA (shown in *E. coli* (26)) are not

rapidly degraded upon translational shutoff in the Δ*lon* strain (Figure 5C, Table S8)

compared to the wild type strain (Figure 5B, Table S8). Additionally, the decrease in

protein abundance upon shutoff is largely unchanged for ClpXP and ClpAP substrates

(see examples for FtsZ, FtsA, and DnaX in Figure S5).

To identify potential Lon substrates in this list, we compared the change in protein

abundance after translational shutoff for each protein in the wild type and Δ*lon* strains.

The 10% of proteins with the greatest difference in $\log_2$(5 hrs/3 hrs) between the two

strains were considered potential Lon substrates (29 candidates; Table S8). Two of these

proteins are the known Lon substrates DnaA (25) and a small heat shock protein IbpA

(26). UvrD, a DNA damage repair helicase, was also identified here.

Based on our results, we conclude that not all changes at the protein abundance during

the DNA damage response can be explained by changes in transcript level. This shutoff

stage of our proteomics experiments allows us to nominate candidates for post-

translational control through degradation.

DISCUSSION

<u>Survey of protein abundance changes upon DNA damage</u>

The transcriptional regulation of the DNA damage response by the RecA/LexA system

and the effect of DNA damage treatment on the transcriptome has been studied in the

canonical *E. coli* response (3, 4) and in *Caulobacter* (14, 15). Past work in gamma-

irradiated *E. coli* showed a correlation between proteome upregulation and transcriptional

changes (28). However, our work is the first to our knowledge to compare proteomic and

transcriptomic changes with the intent to identify targets of post-transcriptional regulation.

While we find that protein abundance changes upon DNA damage largely correspond to

changes in transcript levels, not all protein changes during *Caulobacter's* DNA damage

response can be explained by transcriptional effects. Because these differences could be

explained by a regulation at the post-transcriptional or post-translational level, we

augmented our survey with translational shutoff experiments to identify factors that are

potentially regulated by proteolysis during the DNA damage response.

Our observations in the Δ*lon* strain are consistent with its reduced tolerance of DNA

damage compared to wild type (28). While a similar set of proteins is upregulated upon

DNA damage in the Δ*lon* and the wild type strains, fold-inductions of proteins during the

DNA damage response were generally lower in the Δ*lon* strain (Figure 4). We propose

this is due to the fact that many DNA damage genes are already upregulated in the Δ*lon*

strain even in the absence of exogenous DNA damage treatment (24). These data may

also explain the previous observation that the Δ*lon* strain has an elevated sensitivity to

DNA damage (29).

<u>Possible post-transcriptional regulation of known DNA damage genes</u>

We were able to identify known DNA damage proteins that appear to be post-

transcriptionally regulated in our conditions. Some examples are discussed here.

In *E. coli* UvrA is known to be degraded in a DNA damage-controlled manner by ClpXP

(13). In *Caulobacter*, we find an increase in protein levels following DNA damage that is

likely due to increased transcription (Table S4, (15)). Consistent with a role for ClpXP in

proteolysis, we observe a Lon-independent decrease in protein abundance (Table S8)

upon translational shutoff. These data indicate that the proteolytic control of UvrA

observed in *E. coli* is likely conserved in *Caulobacter*.

After LexA autocleavage is induced by the RecA nucleoprotein filament, both halves of

LexA are proteolyzed by ClpXP in *E. coli* (11). Proteolysis of the DNA-binding N-terminus

of LexA is critical for DNA damage survival, likely because it prevents any residual

repression that may be caused by this fragment (11). Our proteomics dataset only

identified peptides that mapped to the N-terminus of LexA (Figure S6), which is highly

conserved between *E. coli* and *Caulobacter*. Upon translational shutoff we find that the

LexA N-terminus decreases in protein abundance upon translational shutoff in a Lon-

independent manner. This suggests that the proteolysis of the N-terminal LexA fragment

observed in *E. coli* is conserved in *Caulobacter*.

E. coli radA plays a role in recombination and survival to DNA damage (30). In the

hyperthermophilic archaeon *Pyrococcus horikoshii* RadA levels has been shown to be

controlled by protein splicing in the (31), supporting a post-translational control

mechanism for this protein. *Caulobacter radA* is part of an eight-gene operon with genes

for integral membrane proteins and amino acid metabolism (32), lacking a predicted

LexA-binding site (14). While transcript levels for genes in *Caulobacter radA*'s operon

remain constant upon DNA damage (15), RadA protein levels increase significantly upon

DNA damage (Table S5). We were able to detect two other proteins encoded upstream

of *radA* in this operon in our proteomics (CCNA_01728 and CCNA_01730) and they were

not upregulated at the protein level (Table S2). These observations suggest that post-

transcriptional regulation may play a role in the selective expression of the *Caulobacter*

RadA protein.

Finally, while DNA repair gene *uvrD* is upregulated at the transcript level (15, 16), we do

not see a significant increase in UvrD protein abundance upon DNA damage (Table S6).

Our observation that UvrD protein abundance significantly decreases after translational

shutoff suggests that proteolysis may play a role in its regulation. UvrD is stable upon

translational shutoff in the Δ*lon* strain, supporting a role for Lon in this process, either

through direct proteolysis of UvrD or of a second factor that regulates UvrD protein levels.

<u>Additional proteins regulated independent of transcription following DNA damage</u>

Through our studies, we discovered post-transcriptionally regulated factors that were not

previously known to play a role in the *Caulobacter* DNA damage response.

Caulobacter YaaA is upregulated at the protein level upon DNA damage with no

corresponding increase in transcript levels (15). In *E.coli, yaaA* is transcriptionally

regulated by OxyR, is expressed in response to oxidative stress (19, 33) and protects

cells from damage from the Fenton reaction by controlling intracellular levels of

unincorporated iron (19). In *Caulobacter*, *yaaA* does not appear to be regulated by OxyR

(34), but published RB-Tnseq experiments indicates that transposon insertions in *yaaA*

cause a sensitivity to cisplatin (20). In *E. coli* cisplatin treatment causes an increase in

iron uptake through downregulation of Fur (35) and *Δfur ΔyaaA* strains show a

synthetically lethal effect (19). These published works and our observation that Δ*yaaA*

Caulobacter has a competitive disadvantage against the wild type strain in MMC suggest

that *Caulobacter yaaA* could play a similar role (Figure 2C).

We also find evidence for post-transcriptional downregulation of proteins such as the

phosphoribosylamine-glycine ligase PurD, whose protein levels decrease upon DNA

damage without a corresponding decrease in RNA levels (Table S7). PurD is the second

enzyme in the bacterial purine biosynthesis pathway. In *Bacillus anthracis*, *purD* and

some other purine biosynthesis genes are controlled by riboswitches that deactivate

translation when levels of guanosine are high (36). Based on this and our proteomic

results, we propose that *purD* is also controlled by post-transcriptional regulation in

Caulobacter, though it is not clear why downregulation of purine biosynthesis would be

beneficial during the DNA damage response. One possibility is that this is a side-effect of

MMC crosslinking, possibly by disrupting function of a cis- or trans-acting element

regulating translation of the *purD* mRNA. Another possibility is that guanosine derivatives

(such as ppGpp (37)), which are upregulated during DNA damage, may be affecting

translation of this gene.

<u>Concluding thoughts</u>

In our proteomics analysis, we identified and quantified protein abundances for about

30% of the *Caulobacter* proteome. While some of the missing proteins are likely not

expressed during the growth conditions of our assays, some key DNA damage response

proteins are missing from our dataset. Some of these are known to be membrane-

associated (e.g. SidA (15) and DidA (16)), which may reduce our detection, and others

are likely have relatively low abundance, even during DNA damage induction. Future work

using protocols designed specifically to observe these proteins could reveal further

instances of post-transcriptional regulation.

Our survey of protein upregulation during the *Caulobacter* DNA damage response led us

to identify post-transcriptional control of factors important for stress tolerance as

described above. Our work nominates putative Lon substrates that could contribute to

DNA damage tolerance and shows that DNA damage induced proteome changes are

dampened in Δlon strains. Together, these data reveal that not all protein abundance

changes during the DNA damage response can be explained solely by canonical

transcriptional changes and highlights contributions from post-transcriptional regulation.

METHODS

Bacterial strains and growth conditions

Bacterial strains and plasmids used in this work are listed in Tables S9 and S10,

respectively. All *Caulobacter* strains used in this work are NA1000 derivatives.

Caulobacter strains were grown in PYE broth (2 g/L peptone, 1 g/L yeast extract, 1 mM

$MgSO_4$, and 0.5 mM $CaCl_2$). Solid media was made with 1.5% agar. Knockout strains

were created by replacing the native gene with a gentamycin resistance cassette using

the pNPTS138 non-replicating vector (38). Venus- and mKate2-expressing strains were

created by integrating pXGFPC-2 (39) based plasmids encoding fluorescent proteins

driven by a *lac* promoter (40) at the xylose locus.

Bacterial growth curves

An overnight culture was diluted into PYE at an OD600 = 0.05. A volume of 200uL of cells

was added to wells of a 96-well plate. Plate was shaken continuously at 30ºC and OD600

was read every 20 minutes in a BioTek Epoch 2 Microplate Spectrophotometer (Agilent,

Santa Clara, CA). MMC (0.5 µg/ml; Millipore Sigma, St. Louis, MO) was used to treat cells

with DNA damage.

Western blot

Western blots were performed using a commercially available rabbit anti-*E. coli* RecA

primary antibody (1:10,000 dilution; Abcam, Cambridge, MA, 16 hours). Blots were

visualized using the goat anti-rabbit Alexa Fluor™ Plus 800 secondary antibody (1:10,000

dilution; Licor, Lincoln, NE, 1 hour). Bands were quantified using ImageJ (41).

Competition assays

Overnight cultures of each test strain and the control strain were diluted to an OD600 =

1.0, mixed 1:1, and serially diluted 30,000-fold. Undiluted cells were analyzed with a

SpectraMax M2 microplate reader (Molecular Devices, San Jose, CA). Measurements

were taken for cell density (OD600), Venus fluorescence (Excitation 490nm, Cut off

515nm, Emission 530nm), and mKate2 fluorescence (Excitation 575nm, Cut off 590nm,

Emission 630nm). Diluted samples were grown to stationary phase (~48 hours, shaking

at 30°C) and optical density and fluorescence measurements were taken again. The

competitive index was calculated as the $\log_2$[((ratio test/control)END / (ratio

test/control))START / ((average ratio wild type test/control)END/[(ratio wild type

test/control))START].

Sample collection and preparation for proteomics

<u>Pre-damage sample</u>

Three biological replicates of overnight growth of each *Caulobacter* strain (wild type and

Δlon) were diluted to an OD600 = 0.0005 using PYE to a volume of 50mL and grown for

approximately 10 generations to an OD600 of 0.5-0.6. All samples were normalized to an

OD600 of 0.5 using PYE to a volume of 25mL. A 5 mL pre-damage sample was taken,

and cells were collected by centrifugation (6,000xg, 5min). Cell pellets were resuspended

in fresh lysis buffer (8M urea, 50mM HEPES pH7.5, 75mM NaCl), flash frozen, and stored

at -80˚C.

<u>Post-damage/pre-shutoff sample</u>

Cells were treated with 0.5 μg/mL MMC for 1 hour before cells were collected by

centrifugation (6,000xg, 10min) and resuspended in fresh PYE broth to an OD600 = 0.5

to a volume of 25mL. These treatment conditions are similar to those of the microarray

from the Laub lab (15) for simple comparison. Previous experiments indicate that the wild

type strain has about 55% survival to this treatment (38). After 2 hours, all samples were

normalized to an OD600 = 0.5 using PYE to a volume of 25mL. A 5 mL post-damage

sample was taken, and cells were collected by centrifugation (6,000xg, 5min). Cell pellets

were resuspended in fresh lysis buffer, flash frozen, and stored at -80˚C.

<u>Post-shutoff sample</u>

Cells were treated with chloramphenicol (30 μg/mL) for 2 hours. A 5 mL post-shutoff

sample was taken, and cells were collected by centrifugation (6,000xg, 5min). Cell pellets

were resuspended in fresh lysis buffer, flash frozen, and stored at -80˚C.

<u>Sample preparation for proteomics</u>

Samples were freeze-thawed three times in liquid nitrogen to aid cell lysis, then

centrifuged at 16,000xg for 10 mins at 4˚C. Supernatant was carefully transferred to a

fresh 1.5mL tube. A Bradford assay was performed to quantify protein concentration.

Because the Bradford assay is sensitive to high urea concentrations, a small fraction of

each sample was diluted 4-fold for the Bradford assay. 50 μg of each sample was

transferred to a fresh 1.5mL tube. The final volume was adjusted to 50μL using 100mM

Triethylammonium bicarbonate (TEAB) buffer.

To reduce disulfide bonds, 5 μL 200mM tris(2-carboxyethyl)phosphine (TCEP) was

added to each sample and incubated at 55°C for 1 hour. To modify cysteine residues,

5uL of 375mM iodoacetamide was added to each sample and incubated 30 min at room

temperature in the dark. To precipitate protein, 6-volumes of cold acetone was added to

each sample, and they were stored overnight at -20°C.

Samples were centrifuged at 8,000 x g for 10 min at 4°C to collect precipitate. Supernatant

was carefully removed and discarded. Pellets were allowed to dry at room temperature

for 1 hour. Protein pelleted were resuspended in 100uL 50mM TEAB. Samples were

incubated overnight at 37°C with 1.25 μg of trypsin (for a 2.5:100 ratio of trypsin to

protein).

Tandem mass tagging for proteomics

Samples were labelled using the TMT10plex™ Isobaric Label Reagent Set

(ThermoFisher Scientific, Waltham, MA, Catalog #90110) according to the product

instructions. Since each sample only contains 50 μg of peptides instead of the suggested

100 μg, half of each label was used per sample.

Proteomics sample fractionation

For each replicate experiment in each strain, equal volumes of each samples were

combined into a single tube and then dried in a SpeedVac. Pooled samples were

resuspended in 300 μL 0.1% TFA. Samples were dried in a SpeedVac a second time and

resuspended in 300 μL 0.1% TFA. The pH of each sample was measured to confirm that

it was between 2.5 - 3.5, which is critical for effective fractionation.

Pooled samples were fractionated using the Pierce™ High pH Reversed-Phase Peptide

Fractionation Kit (ThermoFisher Scientific, Waltham, MA, Catalog #84868) according to

the kit instructions. Fractions 1 and 5, 2 and 6, 3 and 7, as well as 4 and 8 were pooled

together for a final total of 4 fractions per sample. Samples were dried and resuspended

in 0.1% folic acid in water for mass spectrometry.

Mass spectrometry for tandem mass tagging proteomics

LC-MS analysis was performed, modified from (42) using an Easy-nLC 1000 nanoLC

chromatography system interfaced to an Orbitrap Fusion mass spectrometer

(ThermoFisher Scientific, Waltham, MA). Samples were pre-concentrated and desalted

on to a C18 trap column prior to separation over a 180-min gradient from 0% to 50%

mobile phase B (A:0.1% formic acid in water, B:0.1% formic acid in acetonitrile) at 300

nL/min flow rate with a 75 μm x 15 cm PepMap RLSC column (ThermoFisher Scientific,

Waltham, MA). Mass spectrometry parameters were as follows: ion spray voltage 2000V,

survey scan MS1 120k resolution with a 2 s cycle time, interleaved with data-dependent

ion trap MS/MS of highest intensity ions with charge state-dependent ion selection

window (z=2:1.2 m/z, z=3:0.7 m/z, z=4-6:0.5 m/z) and CID at 35% normalized collision

energy. Additionally, the top 5 (z=2) or 10 (z>2) product ions were synchronously selected

for HCD MS^3 at NCE 65% with Orbitrap detection to generate TMT tag intensities.

Proteomics Data Analysis

RAW data files were analyzed in Proteome Discoverer 2.4 (ThermoFisher Scientific,

Waltham, MA) using the SEQUEST search algorithm against *Caulobacter crescentus*

NA1000 (NCBI Reference Sequence: NC_011916.1) database downloaded from

uniprot.org. The search parameters used are as follows: 10 ppm precursor ion tolerance

and 0.4 Da fragment ion tolerance; up to two missed cleavages were allowed; dynamic

modifications of methionine oxidation and N-terminal acetylation. Peptide matches were

filtered to a protein false discovery rate of 5% using the Percolator algorithm. Peptides

were assembled into proteins using maximum parsimony and only unique and razor

peptides were retained for subsequent analysis. Protein quantitation based on TMT ion

abundance was performed using a co-isolation threshold of 75% and SPS match

threshold 65%. Each TMT channel was normalized to total peptide amount and then

abundance scaled to average 100% for all proteins.

Protein abundances were then normalized to the averages of the 0 hour samples (Table

S1). Protein ratios (Table S2) were calculated using edgeR (43).

ACKNOWLEDGEMENTS

460   This work was supported by NIH/NIGMS R35GM130320 to P. Chien. The mass

461   spectrometry in this paper was performed by Dr. Stephen Eyles at the University of

462   Massachusetts Amherst Mass Spectrometry Core Facility in Amherst, MA

463   (RRID:SCR_019063). The *ΔfzlC* strain was a generous gift from Erin Goley's lab at Johns

464   Hopkins University, Baltimore, MD.

REFERENCES

1. Little JW, Edmiston SH, Pacelli LZ, Mount DW. 1980. Cleavage of the *Escherichia*

coli lexA protein by the *recA* protease. Proc Natl Acad Sci U S A 77:3225–3229.

2. Horii T, Ogawa T, Nakatani T, Hase T, Matsubara H, Ogawa H. 1981. Regulation of

SOS functions: purification of *E. coli* LexA protein and determination of its specific

site cleaved by the RecA protein. Cell 27:515–522.

3. Courcelle J, Khodursky A, Peter B, Brown PO, Hanawalt PC. 2001. Comparative

gene expression profiles following UV exposure in wild-type and SOS-deficient

Escherichia coli. Genetics 158:41–64.

4. Khil PP, Camerini-Otero RD. 2002. Over 1000 genes are involved in the DNA

damage response of *Escherichia coli*. Mol Microbiol 44:89–105.

5. Vogel J, Argaman L, Wagner EGH, Altuvia S. 2004. The small RNA IstR inhibits

synthesis of an SOS-induced toxic peptide. Curr Biol 14:2271–2276.

6. Darfeuille F, Unoson C, Vogel J, Wagner EGH. 2007. An antisense RNA inhibits

translation by competing with standby ribosomes. Mol Cell 26:381–392.

7. Fröhlich KS, Förstner KU, Gitai Z. 2018. Post-transcriptional gene regulation by an

Hfq-independent small RNA in *Caulobacter crescentus*. Nucleic Acids Res

46:10969–10982.

8. Godoy VG, Jarosz DF, Simon SM, Abyzov A, Ilyin V, Walker GC. 2007. UmuD and

RecA Directly Modulate the Mutagenic Potential of the Y-family DNA Polymerase

DinB. Mol Cell 28:1058–1070.

9.   Cafarelli TM, Rands TJ, Godoy VG. 2014. The DinB•RecA complex of *Escherichia*

coli mediates an efficient and high-fidelity response to ubiquitous alkylation lesions.

Environ Mol Mutagen 55:92–102.

10.   Friedhoff P, Li P, Gotthardt J. 2016. Protein-protein interactions in DNA mismatch

repair. DNA Repair (Amst) 38:50–57.

11.   Neher SB, Flynn JM, Sauer RT, Baker TA. 2003. Latent ClpX-recognition signals

ensure LexA destruction after DNA damage. Genes Dev 17:1084–1089.

12.   Gonzalez M, Rasulova F, Maurizi MR, Woodgate R. 2000. Subunit-specific

degradation of the UmuD/D' heterodimer by the ClpXP protease: the role of trans

recognition in UmuD' stability. EMBO J 19:5251–5258.

13.   Pruteanu M, Baker TA. 2009. Controlled degradation by ClpXP protease tunes

the levels of the excision repair protein UvrA to the extent of DNA damage. Mol

Microbiol 71:912–924.

14.   da Rocha RP, Paquola AC de M, Marques M do V, Menck CFM, Galhardo RS.

2008. Characterization of the SOS regulon of *Caulobacter crescentus*. J Bacteriol

190:1209–1218.

15.   Modell JW, Hopkins AC, Laub MT. 2011. A DNA damage checkpoint in

Caulobacter crescentus inhibits cell division through a direct interaction with FtsW.

Genes Dev 25:1328–1343.

16.   Modell JW, Kambara TK, Perchuk BS, Laub MT. 2014. A DNA Damage-Induced,

506   SOS-Independent Checkpoint Regulates Cell Division in *Caulobacter crescentus*.

507   PLoS Biol 12:e1001977.

508   17.   Gozzi K, Salinas R, Nguyen VD, Laub MT, Schumacher MA. 2022. ssDNA is an

509   allosteric regulator of the *C. crescentus* SOS-independent DNA damage response

510   transcription activator, DriD. Genes Dev 36:618–633.

511   18.   Prahlad J, Yuan Y, Lin J, Chang C-W, Iwata-Reuyl D, Liu Y, de Crécy-Lagard V,

512   Wilson MA. 2020. The DUF328 family member YaaA is a DNA-binding protein with

513   a novel fold. Journal of Biological Chemistry 295:14236–14247.

514   19.   Liu Y, Bauer SC, Imlay JA. 2011. The YaaA protein of the Escherichia coli OxyR

515   regulon lessens hydrogen peroxide toxicity by diminishing the amount of intracellular

516   unincorporated iron. J Bacteriol 193:2186–2196.

517   20.   Price MN, Wetmore KM, Waters RJ, Callaghan M, Ray J, Liu H, Kuehl JV,

518   Melnyk RA, Lamson JS, Suh Y, Carlson HK, Esquivel Z, Sadeeshkumar H,

519   Chakraborty R, Zane GM, Rubin BE, Wall JD, Visel A, Bristow J, Blow MJ, Arkin AP,

520   Deutschbauer AM. 2018. Mutant phenotypes for thousands of bacterial genes of

521   unknown function. 7706. Nature 557:503–509.

522   21.   Xing S, Zheng W, An F, Huang L, Yang X, Zeng S, Li N, Ouenzar K, Yu L, Luo L.

523   2022. Transcription Regulation of Cell Cycle Regulatory Genes Mediated by NtrX to

524   Affect *Sinorhizobium meliloti* Cell Division. Genes (Basel) 13:1066.

525   22.   Stein BJ, Fiebig A, Crosson S. 2021. The ChvG-ChvI and NtrY-NtrX Two-

526   Component Systems Coordinately Regulate Growth of *Caulobacter crescentus*. J

527   Bacteriol 203:e0019921.

528   23.   Meier EL, Razavi S, Inoue T, Goley ED. 2016. A novel membrane anchor for

529   FtsZ is linked to cell wall hydrolysis in *Caulobacter crescentus*. Mol Microbiol

530   101:265–280.

531   24.   Mahmoud SA, Aldikacti B, Chien P. 2022. ATP hydrolysis tunes specificity of a

532   AAA+ protease. Cell Reports 40:111405.

533   25.   Liu J, Zeinert R, Francis L, Chien P. 2019. Lon recognition of the replication

534   initiator DnaA requires a bipartite degron. Mol Microbiol 111:176–186.

535   26.   Bissonnette SA, Rivera-Rivera I, Sauer RT, Baker TA. 2010. The IbpA and IbpB

536   small heat-shock proteins are substrates of the AAA+ Lon protease. Mol Microbiol

537   75:1539–1549.

538   27.   Bhat NH, Vass RH, Stoddard PR, Shin DK, Chien P. 2013. Identification of ClpP

539   substrates in *Caulobacter crescentus* reveals a role for regulated proteolysis in

540   bacterial development. Molecular Microbiology 88:1083–1092.

541   28.   Gaougaou G, Shankar S, Liot Q, Constant P, Déziel E, Lacroix M. 2020. Gamma

542   irradiation triggers a global stress response in *Escherichia coli* O157:H7 including

543   base and nucleotides excision repair pathways. Microbial Pathogenesis

544   149:104342.

29.   Zeinert RD, Ogdahl JL, Liu J, Yang Q, Du Y, Barros BB, Freddolino PL, Haynes

CM, Chien P. 2022. A legacy role for DNA binding of Lon protects against genotoxic

stress. bioRxiv https://doi.org/10.1101/317677.

30.   Beam CE, Saveson CJ, Lovett ST. 2002. Role for *radA/sms* in recombination

intermediate processing in *Escherichia coli*. J Bacteriol 184:6836–6844.

31.   Topilina NI, Novikova O, Stanger M, Banavali NK, Belfort M. 2015. Post-

translational environmental switch of RadA activity by extein–intein interactions in

protein splicing. Nucleic Acids Res 43:6631–6648.

32.   Zhou B, Schrader JM, Kalogeraki VS, Abeliuk E, Dinh CB, Pham JQ, Cui ZZ, Dill

DL, McAdams HH, Shapiro L. 2015. The global regulatory architecture of

transcription during the *Caulobacter* cell cycle. PLoS Genet 11:e1004831.

33.   Zheng M, Wang X, Templeton LJ, Smulski DR, LaRossa RA, Storz G. 2001.

DNA microarray-mediated transcriptional profiling of the Escherichia coli response to

hydrogen peroxide. J Bacteriol 183:4562–4570.

34.   Silva LG, Lorenzetti APR, Ribeiro RA, Alves IR, Leaden L, Galhardo RS, Koide

T, Marques MV. 2019. OxyR and the hydrogen peroxide stress response in

Caulobacter crescentus. Gene 700:70–84.

35.   Beaufay F, Quarles E, Franz A, Katamanin O, Wholey W-Y, Jakob U. 2020.

Polyphosphate Functions In Vivo as an Iron Chelator and Fenton Reaction Inhibitor.

mBio 11:e01017-20.

36.　　Kirchner M, Schneider S. 2017. Gene expression control by *Bacillus anthracis*

purine riboswitches. RNA 23:762–769.

37.　　Rasouly A, Pani B, Nudler E. 2017. A Magic Spot in Genome Maintenance.

Trends Genet 33:58–67.

38.　　Tashjian TF, Chien P. 2023. Clamp Loader Processing Is Important during DNA

Replication Stress. J Bacteriol 205:e0043722.

39.　　Thanbichler M, Iniesta AA, Shapiro L. 2007. A comprehensive set of plasmids for

vanillate- and xylose-inducible gene expression in Caulobacter crescentus. Nucleic

Acids Res 35:e137.

40.　　Persat A, Stone HA, Gitai Z. 2014. The curved shape of Caulobacter crescentus

enhances surface colonization in flow. Nat Commun 5:3824.

41.　　Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years

of image analysis. Nat Methods 9:671–675.

42.　　McAlister GC, Nusinow DP, Jedrychowski MP, Wühr M, Huttlin EL, Erickson BK,

Rad R, Haas W, Gygi SP. 2014. MultiNotch MS3 enables accurate, sensitive, and

multiplexed detection of differential expression across cancer cell line proteomes.

Anal Chem 86:7150–7158.

43.　　Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package

for differential expression analysis of digital gene expression data. Bioinformatics

26:139–140.

Figure 1. Proteomics reveals protein abundance changes during response to MMC treatment. (A) A western blot for RecA protein levels was used to determine when to sample for protein abundance changes after DNA damage treatment. Quantification of western blot for two replicates is shown, blot shown in Figure S1. Samples for proteomics were taken before treatment (0 hours, pink line). Cells were treated with MMC for 1 hour and recovered in fresh PYE for 2 additional hours before the post-treatment proteomics sample was taken (3 hours, teal line). RecA quantification was normalized to cross-reacting band for loading (Figure S1). (B) A volcano plot of proteomics data compares the statistical confidence ($-\log_{10}$(false discovery rate)) versus the $\log_2$(3 hrs/0 hrs). Several of the proteins that are significantly upregulated after DNA damage treatment are known DNA damage response proteins (highlighted in red). FDR indicates false discovery rate.

Figure 2. A comparison of proteomics and microarray data identifies proteins that are upregulated upon MMC treatment in a transcription-independent manner. A comparison between our proteomics and previously published microarray experiment that uses similar MMC treatment conditions (15) shows proteins that are upregulated upon DNA damage. Proteins highlighted in red are known DNA damage response genes that are upregulated at the RNA and protein levels. Proteins highlighted in blue are upregulated at the protein level, but not significantly upregulated at the RNA level and were identified by Tn-seq as potentially important for DNA damage survival. Graph

cropped to upper right quadrant to highlight proteins that are upregulated. One gene is

excluded (CCNA_03825) due to its extremely high induction to make other points more

visible. (B) A Venn diagram of proteins that are upregulated at the protein level ($\log_2$(3

hrs/0 hrs) > 0, FDR > 0.05) and genes that are upregulated at the RNA level ($\log_2$(post-

MMC/pre-MMC) > 1 standard deviation above the mean for all transcripts). The smaller

circle on the right side of this diagram reflects upregulated transcripts that were not

identified by proteomics. (C) When each mKate2-expressing test strain is competed in

co-culture against a wild type Venus-expressing fluorescent control strain, only the *ΔyaaA*

mutant shows a competitive disadvantage that is exacerbated by the presence of 0.1

µg/mL MMC. Reciprocal experiment in which test strains are marked with Venus and wild

type control is marked with mKate2 is shown in Figure S2D.

Figure 3. A comparison of proteomics and microarray data identifies proteins that

are downregulated upon MMC treatment in a transcription-independent manner.

The Venn diagram shows proteins that are downregulated at the protein level ($\log_2$(3 hrs/0

hrs) < 0, FDR < 0.05) and genes that are downregulated at the RNA level ($\log_2$(post-

MMC/pre-MMC) > 1 standard deviation below the mean for all transcripts). The smaller

circle on the left side of this diagram reflects proteins not identified in the microarray. The

smaller circle on the right side reflects transcripts that were not identified in our

proteomics. (B) The Δ*purD* strain has a stationary phase defect that is not observed in

the presence of MMC. All experiments were performed in triplicate. Mean and standard

deviation are shown. (C) The Δ*fzlC* strain show no difference in MMC sensitivity

compared to the wild type strain. All experiments were performed in triplicate. Mean and

standard deviation are shown. Same data for wild type controls are shown in (B) and (C)

for ease of comparison.

Figure 4. A similar set of DNA damage response proteins are upregulated in a Δ*lon*

strain, but protein abundance changes are overall smaller. (A) A western blot to

determine RecA protein levels was used to determine when to sample for protein

abundance changes after DNA damage treatment in the wild type Δ*lon* strains. Note that

the wild type data here is repeated from Figure 1A for easy comparison. Quantification of

western blot for two replicates is shown, blot shown in Figure S1. Sampling for proteomics

was performed as described in Figure 1A. RecA quantification was normalized to cross-

643 reacting band for loading (see Figure S1). (B) A volcano plot of proteomics data compares

644 the statistical confidence ($-\log_{10}$(false discovery rate)) versus the $\log_2$(3 hrs/0 hrs). Known

645 DNA damage proteins shown in Figure 1B are also highlighted here in red. ImuA was not

646 identified in the proteomics of the Δ*lon* strain and so is absent. FDR indicates false

647 discovery rate. (C) Comparison of the $\log_2$(3 hrs/0 hrs) for the Δ*lon* versus wild type strain

648 shows that fold changes of protein abundance are generally lower in the Δ*lon* strain. Blue

649 dashed line represents an y=x line. The red line represents the trendline of the data with

650 95% CI drawn (slope = 0.45). The inset plots $\log_2$(3 hrs/0 hrs) for all proteins that are

651 significantly upregulated (FDR <0.05).

Figure 5. Proteomics revealed protein abundance changes when translation is shut off during the response to MMC treatment. (A) Proteomics results for changes in DnaA and IbpA protein abundance upon MMC treatment and subsequent translational shutoff. Samples for proteomics were taken before treatment (0 hours, pink line). Cells were treated with MMC for 1 hour and recovered in fresh PYE for 2 additional hours before the post-treatment proteomics sample was taken (3 hours, teal line). At 3 hours, cells were treated with chloramphenicol and final proteomics samples was taken after 2 additional hours (5 hours, purple line). Note that protein abundances are not directly comparable

between the two strains. Volcano plots compare the statistical confidence ($-\log_{10}$(false

discovery rate)) versus the $\log_2$(5 hrs/3 hrs) for (B) wild type and (C) Δ*lon* strains. Several

of the proteins that are significantly downregulated after chloramphenicol treatment are

known proteolysis substrates (examples highlighted in red in (A); see main text). Two

proteins of interest that will be discussed below are highlighted in blue. FDR indicates

false discovery rate.