**Proteomic survey of the DNA damage response in *Caulobacter crescentus***

Running Title: Proteomics survey of *Caulobacter* DNA damage response

Tommy F. Tashjian[#], Rilee D. Zeinert[*], Stephen J. Eyles, Peter Chien[#]

Department of Biochemistry and Molecular Biology, University of Massachusetts

Amherst, Amherst, MA, USA

*Current address: Cell Biology and Metabolism Program, Eunice Kennedy Shriver

National Institute of Child Health and Human Development, Bethesda, MD 20892-5430,

USA

#co-corresponding authors:

Tommy Tashjian, Department of Biochemistry and Molecular Biology, University of

Massachusetts Amherst, Life Science Laboratories Room N325, 240 Thatcher Rd,

Amherst MA 01003-9364, (413)545-2310, ttashjian@amherst.edu

Peter Chien, Department of Biochemistry and Molecular Biology, University of

Massachusetts Amherst, Life Science Laboratories Room N325, 240 Thatcher Rd,

Amherst MA 01003-9364, (413)545-2310, pchien@umass.edu;

**ABSTRACT**

The bacterial DNA damage response is a critical, coordinated response to endogenous

and exogenous sources of DNA damage. Response dynamics are dependent on

coordinated synthesis and loss of relevant proteins. While much is known about its global

transcriptional control, changes in protein abundance that occur upon DNA damage are

less well characterized at the system level. Here, we perform a proteome-wide survey of

the DNA damage response in *Caulobacter crescentus*. We find that while most protein

abundance changes upon DNA damage are readily explained by changes in transcription,

there are exceptions. The survey also allowed us to identify the novel DNA damage

response factor, YaaA, which has been overlooked by previously published, transcription-

focused studies.

A similar survey in a Δ*lon* strain was performed to explore *lon's* role in DNA damage

survival. The Δ*lon* strain had a smaller dynamic range of protein abundance changes in

general upon DNA damage compared to the wild type strain. This system-wide change

to the dynamics of the response may explain this strain's sensitivity to DNA damage. Our

proteome survey of the DNA damage response provides additional insight into the

complex regulation of stress response and nominates a novel response factor that was

overlooked in prior studies.

**IMPORTANCE**

The DNA damage response helps bacteria to react to and potentially survive DNA

damage. The mutagenesis induced during this stress response contributes to the

development of antibiotic resistance. Understanding how bacteria coordinate their

response to DNA damage could help us to combat this growing threat to human health.

While the transcriptional regulation of the bacterial DNA damage response has been

characterized, this study is the first to our knowledge to assess the proteomic response

to DNA damage in *Caulobacter*.

**INTRODUCTION**

The bacterial DNA damage response is a major driver of antibiotic resistance

development (1, 2) and the horizontal gene transfer that spreads resistance and

pathogenicity from cell to cell (3–6). The canonical DNA damage response was first

characterized in *Escherichia coli*. During DNA damage ssDNA accumulates in the cell

and is bound by the cell's main recombinase RecA to form a nucleoprotein filament that

induces the autocleavage of the global transcriptional repressor LexA (7, 8). Cleavage of

LexA allows transcription of more than 40 genes related to DNA damage tolerance and

repair, including *recA* and *lexA* themselves (9, 10). The energy-dependent protease

ClpXP then degrades the DNA binding N-terminal domain of LexA to prevent residual

repression (11).

As DNA damage is repaired, accumulation of uncleaved LexA represses transcription of

DNA damage response genes. Shutdown of the response is also aided by proteolysis of

certain response factors (e.g. UmuD' (12) and UvrA (13) by ClpXP and UmuD (14) and

SulA (15) by Lon). In fact, both ClpXP (11) and Lon (16, 17) contribute to the survival of

DNA damage in *E. coli*.

The LexA/RecA-based transcriptional control of the response is conserved in *Caulobacter*

*crescentus*, as are many of the genes in the *E. coli* LexA regulon (18).  A second, LexA-

independent DNA damage response pathway that is dependent on global transcriptional

regulator DriD has also been identified and characterized in *Caulobacter* (19, 20). While

the energy-dependent protease Lon is conserved in *Caulobacter* and contributes to

survival of DNA damage (21), Lon's confirmed DNA damage-related substrates (UmuD

(14) and SulA (15)) are not. Lon has been shown to contribute to DNA damage survival

in *Caulobacter* but its role in response to DNA damage remains unclear.

Transcriptome measurements upon treatment with various DNA damaging agents and a

bioinformatic search for LexA binding sites have nominated a set of genes that are

upregulated upon damage in *Caulobacter* (18, 19, 22). While transcription and proteolysis

both play roles in DNA damage response induction and shut-off, relatively little is known

about how well transcriptional changes correspond to protein abundance changes at the

proteome level.

Here, we present a proteomic survey to investigate the dynamics of protein abundance

during the *Caulobacter* DNA damage response. First, we measured protein abundance

changes upon DNA damage treatment. Next, we measured changes following translation

shutoff to characterize potential proteolysis substrates. We found that while most changes

in protein abundance upon DNA damage are explained by changes in transcription, there

are exceptions to this rule. By combining our proteomics and previously measured

changes in transcript levels upon mitomycin C (MMC) treatment (19), we were able to

identify the novel DNA damage response factor YaaA, which has previously been

overlooked in surveys that considered transcriptional changes alone. Our validation

experiments showed this factor to be important during DNA damage, demonstrating the

utility of our approach.

In the second stage of this study, we attempted to identify novel Lon substrates to explain

the importance of this protease to DNA damage survival. While we were unable to identify

any novel Lon substrates that sufficiently explain this phenotype, we did characterize

system-wide changes to the DNA damage response in a Δ*lon* strain. The magnitude of

protein abundance changes upon DNA damage are dampened in the Δ*lon* strain

compared to those in the wild type strain. This effect can be attributed in part to the

elevated basal levels of DNA damage proteins in the untreated Δ*lon* strain (23). This

suggests Lon may be of wider importance to the regulation of the DNA damage response

than simply degrading of one or a few key substrates.

This study provides new insights into Lon's broader importance to protein dynamics in

*Caulobacter's* DNA damage response and identifies a novel factor that is important for

survival to DNA damage.

**RESULTS**

**Proteomics yields novel insights into the DNA damage response**

To characterize protein abundance changes during the DNA damage response, we first

used the sentinel protein RecA to determine the timescale of an acute MMC treatment in

*Caulobacter* (Figure 1A and S1). Cells were treated with MMC for 1 hour and RecA levels

were monitored through recovery (Figure 1A and S1). RecA levels peaked 3 hours after

the addition of MMC (Figure 1A and S1), which is similar to what was previously observed

during chronic MMC treatment (24).

This same timeline was used to obtain samples before and after MMC treatment for a

proteomics survey (timescale in Figure 1B, results in Table S1, S2, and Figure 1C). In

order to determine where post-translational regulation (e.g., proteolysis) may be affecting

protein levels during DNA damage recovery, we treated cells with chloramphenicol at 3

hours to shut off translation and sampled cells for proteomics at 5 hours (timeline in Figure

1B, results in Figure 1C).

To validate our proteomics survey, we looked at the protein abundance changes of known

DNA damage proteins after MMC treatment. As expected, several known DNA damage

response proteins were more abundant after MMC treatment (e.g., RecA, UvrA, RuvB,

ImuA, and MmcA, which are highlighted in red in Figure 1C). In the translation shutoff

portion of the experiment, we observed a decrease in protein abundance of several

known proteolysis substrates, including ClpXP substrates DnaX (25), CtrA (25), FtsZ (26),

UvrA (13), and LexA(11), ClpAP substrates FtsA (26) and Lon (27), and Lon substrates

DnaA (28) and IbpA (29) (all highlighted in red in Figure 1D). These data validated that

our proteomics analysis could be used to identify known DNA damage response factors

and proteolysis substrates.

Previous genome-wide surveys of the *Caulobacter* DNA damage response have focused

on transcriptional regulation (18, 19, 22). We sought to determine how well the changes

in protein abundance correlate to changes in RNA levels. To do this, we compared our

proteomics data to previously published microarray data (19). In our western blot, we

found that that RecA levels peak approximately 3 hours after MMC treatment and this

data was consistent with a previous study (24). The same study indicated that RNA levels

for *recA* and several other DNA damage response genes (e.g., *lexA, uvrA, ruvB, bapE*)

peak at 60-80 minutes after MMC treatment. Based on this timeline and the consistency

with our experimental design we used the study performed by Modell, et al. to compare

our measurement of protein abundance changes to their published changes in RNA levels

upon MMC treatment in PYE (19).  In this study, transcript levels were monitored using

microarrays over several time points (20-80 min) after MMC treatment (or hydroxyurea or

ultra-violet light) in rich medium (or minimal medium) (19).

To discern between genes that are transcriptionally upregulated upon MMC treatment

and those that are not, we used the maximum $\log_2$(Fold Change of RNA) from these

timepoints for each gene and treated genes that were induced more than one standard

deviation above the mean of these as "upregulated". Genes that fell below this cutoff were

considered "not transcriptionally upregulated." This strategy highlighted 615 upregulated

genes of which 146 could be identified as peptides in our proteomics survey. Of these,

43 show significant increase in protein abundance upon DNA damage (Table S4, $\log_2$(3

hrs/0 hrs) > 0, FDR < 0.05). As expected, the known DNA damage proteins mentioned

above that showed increased protein abundances are included in this set (e.g., RecA,

UvrA, RuvB, ImuA, and MmcA, which are highlighted in red in Figure 1C, all data in Table

S4).

We identified 81 proteins that significantly increased in abundance upon DNA damage

($\log_2$(Fold Change in Protein) > 0, FDR < 0.05) that did not show a corresponding

significant increase in RNA level by microarray as defined above ((19), Table S5). To

further validate these hits, we compared them to a RNA-seq dataset that measured

expression of genes following chronic MMC treatment (30). Of our list of 81 proteins, 76

showed no significant increase in expression after chronic MMC treatment (this study

defined upregulated as a 3-fold increase in RNA levels after 4 hours (30)).

We followed up on three of the most induced proteins in this list to determine if they were

important for survival to MMC, but deletion of these genes (CCNA_02118, CCNA_00591,

and CCNA_00276; highlighted in purple in Figure 2A) had no effect on survival to MMC

(Figure S2). These data indicate that while there are examples of proteins that are

upregulated upon DNA damage that are not induced at the RNA level, not all of these are

physiologically relevant to MMC survival.

Of this list of 81 proteins (Table S4) CCNA_03496 was also of particular interest. This

gene is not upregulated upon MMC treatment in microarray (19) but does show a 17%

increase in expression during chronic MMC treatment by RNA-seq (30). Due to its minor

transcriptional induction upon DNA damage, YaaA had previously been overlooked as a

potential DNA damage response factor. It has not previously been shown to be important

for DNA damage survival and does not appear to be regulated by the LexA/RecA system

(18, 22). CCNA_03496 is a homolog of *E. coli*'s *yaaA* ((31), we will refer to this gene as

*yaaA* henceforth), which is important for regulating free iron levels during oxidative stress

conditions (32). Our Tn-seq experiments identified it as potentially important for survival

of MMC (Table S3) and published RB-Tn-seq results suggest that it is important for

survival of the DNA damaging agent cisplatin (33). For these reasons, we decided to

further characterize the connection between *yaaA* and MMC survival.

Deletion of *yaaA* had no effect on growth of isolated cultures treated with MMC (Figure

S3A). We turned to more sensitive competition assays to quantify fitness defects. In these

assays, we grew mKate2-expressing test strains with a Venus-expressing control strain

and measured ratios of mKate2 to Venus fluorescence before and after several

generations of growth. A Δ*yaaA* test strain expressing the mKate2 is less able than the

wild type mKate2-expressing strain to compete in co-culture with the wild type Venus-

expressing control (Figure 2C). This competitive disadvantage worsens in the presence

of MMC (Figure 2B). To confirm that these results are not an artifact of the fluorescent

strains, we reversed the markers to observe Venus-expressing test strains competed

against a mKate2-expressing control and saw similar results (Figure S3B).

Because MMC activation can generate hydrogen peroxide ($H_2O_2$) in the cell (34), we

sought to determine whether the sensitivity of the $\Delta yaaA$ strain to MMC was due to a

sensitivity to DNA damage or to a secondary effect of possible $H_2O_2$ generation. We find

that the $\Delta yaaA$ strain is indeed sensitive to $H_2O_2$ (Figure 2C and reciprocal experiment in

Figure S3C). Importantly, the $\Delta yaaA$ strain is also sensitive to ultra-violet (UV) light, a

non-chemical DNA damaging agent (Figure 2D). Taking these results together, the

simplest explanation is that *yaaA* is important for survival to DNA damage resulting from

photochemistry (UV), crosslinking/intercalation (MMC), or oxidative modification ($H_2O_2$).

We next looked at those genes downregulated following MMC treatment. As before, we

considered a change in RNA levels of more than one standard deviation below the mean

of all values as "downregulated." Based on these criteria, we identified 87 proteins that

became significantly less abundant upon DNA damage without a significant decrease in

RNA abundance (Table S7). We performed further experiments on two genes that our

Tn-seq results identified as more likely to contain transposon insertions upon MMC

damage: *fzlC* and *purD*. However, deletion of these genes did not result in any change to

MMC sensitivity (Supplemental Figure S4).

In the second stage of the proteomics survey, we compared protein level changes before

and after translation shutoff (timeline in Figure 1B). We identified 320 proteins that

significantly decreased in protein abundance (Table S8).  Among these are the DNA

damage response factors LexA and UvrA, which are known to be proteolyzed by ClpXP

in *E. coli* (11, 13). These data suggest that this proteolysis may be conserved in

*Caulobacter*. Others in this list may be substrates for proteolysis by ClpXP or other

*Caulobacter* proteases.

These data suggest that while most protein abundance increases during the DNA

damage response are explained by changes in gene expression, there are some

exceptions. These exceptions may be instances where protein levels are controlled by

post-transcriptional or post-translational regulation, but it is unclear how important these

are to MMC survival. In our survey, we were able to identify and characterize a novel DNA

damage response factor, YaaA, which is important to the survival of oxidative damage as

well as the DNA damaging agents MMC and UV light. Finally, we found evidence to

suggest that the proteolytic regulation of LexA and UvrA that was discovered in *E. coli* is

conserved in *Caulobacter*.

**Investigation of the effect of *lon* deletion on the DNA damage response**

The Lon protease is important for DNA damage survival in several bacterial species (16,

17, 35, 36), including *Caulobacter* (21). To investigate *lon's* role in *Caulobacter's* DNA

damage response we performed the same proteomics experiment as above in a *Δlon*

strain (full dataset in Table S1 and S2). Most of the known DNA damage-related proteins

that were found to increase in abundance upon MMC treatment in the wild type strain

also increase in the Δ*lon* strain (See RecA, RuvB, and MmcA in Figure 3A). The

exceptions to this are UvrA, which did not increase in abundance upon MMC treatment

in the Δ*lon* strain (Figure 3A), and ImuA, which was not identified in the Δ*lon* strain

proteomics.

In the translation shutoff portion of the proteomics experiment known Lon substrates

DnaA and IbpA, which were shown to decrease in abundance in the wild type strain upon

shutoff, showed a far smaller decrease in abundance in the Δ*lon* strain (Figure 3B). As

expected, decreases in protein abundance upon shutoff in the Δ*lon* strain are largely

unchanged for ClpXP and ClpAP substrates (see examples for FtsZ, FtsA, and DnaX in

Figure S5).

To identify potential Lon substrates from our list of potential protease substrates (Table

S8), we compared the change in protein abundance after translational shutoff for each

protein in the wild type and Δ*lon* strains. The 10% of proteins with the greatest difference

in $\log_2$(5 hrs/3 hrs) between the two strains were considered potential Lon substrates (29

candidates; Table S8). The only known DNA damage factor identified in this list was UvrD;

however, loss of *uvrD* is synthetic lethal with the deletion of *lon* (21, 37), suggesting that

accumulation of UvrD is unlikely to underlie the Δ*lon* sensitivity to DNA damage. Overall,

we consider it unlikely that stabilization of a single known DNA damage related protein is

the primary driver for the increased sensitivity to DNA damage for the Δ*lon Caulobacter*

strain. However, we did observe global changes to the DNA damage response in the Δ*lon*

strain, which may be responsible for the MMC sensitivity.

Similar to the wild type strain, we monitored RecA levels through induction and recovery

after MMC treatment in the Δ*lon* strain by western blot (Figure 4A and S1). RecA levels

are about 3-fold higher in the Δ*lon* strain before MMC treatment (average RecA

quantification normalized to ClpP quantification in wild type is 0.09±0.02 and in the Δ*lon*

strain is 0.30±0.02, Figure S1). This is consistent with a previous observation that DNA

damage response genes are upregulated in the Δ*lon* strain even in the absence of

exogenous DNA damaging agents (23). In addition, peak RecA induction in the Δ*lon*

strain occurs at a similar time point as the wild type (Figure 4A and S1) and both strains

reach similar maximum RecA concentration at 3 hours (average RecA quantification

normalized to ClpP quantification in wild type is 0.9±0.2 and in the Δ*lon* strain is 0.8±0.3,

Figure S1). Together, this means that the fold-increase of RecA in the Δ*lon* strain is about

3-fold smaller of that observed in the wild type strain (Figure 4A and S1).

We found it to be a general trend that protein changes upon MMC treatment in Δ*lon* strain

were less extreme than in the wild type strain (Figure 4B). Similarly, changes in protein

abundance were dampened in general upon translation shutoff in Δ*lon* strain (Figure S6).

Furthermore, previous studies have found evidence that highly expressed DNA damage

response proteins are most rapidly degraded (38). We see a similar negative correlation

between protein abundance changes upon MMC treatment (0-3 hours) and upon

translational shutoff (3-5 hours) in the wild type strain (Figure 4C) that is absent in the

Δ*lon* strain (Figure 4D). These proteome-level changes that we observe in the Δ*lon* strain

may be related to *lon's* importance to DNA damage survival in *Caulobacter*.

These data indicate that protein changes during the DNA damage response are generally

explained by changes in RNA levels, but that there are some proteins that do not follow

this trend. In our survey, we identified a novel DNA damage response factor, YaaA, which

had been overlooked in previous transcription-focused surveys. The proteomics survey

of the Δ*lon* strain was not able to identify a single critical Lon substrate that sufficiently

explained the protease's importance to MMC survival. However, we did observe global

changes in protein dynamics that may be responsible for the MMC sensitivity of the strain.

**DISCUSSION**

**Novel insights into the *Caulobacter* DNA damage response**

By comparing our proteomics survey to previously published microarray data (19), we

found that most protein abundance changes upon DNA damage are explained by

changes at the RNA level, though there are exceptions (Figure 2A). Not all genes

upregulated at the transcription or protein level were important for survival, which is

consistent with the observations that genes upregulated during a stress response do not

necessarily overlap with those genes important for tolerating that stress (39–41).

In addition, proteins encoded by three essential genes related DNA replication (*dnaQ,*

*gyrA,* and *gyrB*) showed changes in protein abundance that are not explained by changes

in RNA levels according to microarray (19) or RNA-seq (30). The *dnaQ* gene encodes the

proofreading ε subunit of DNA polymerase III and its expression is upregulated upon

MMC treatment in *E. coli* in a *recA*-dependent manner (42). Our proteomics survey

suggests that its regulation may be post-transcriptional or post-translational in

*Caulobacter*. Since its protein abundance decreases after translational shutoff (Table S2),

proteolysis may play a role. Levels of another replisome component, DnaX, is also

regulated differently in *E. coli* and *Caulobacter.* In *E. coli* a truncated form of DnaX (γ)

that is important during DNA damage (43) is created through a ribosomal frameshift (44),

while in *Caulobacter* similar DNA damage-related γ forms (45) are created through partial

proteolysis by ClpXP (46). It is possible that proteolysis regulates the concentrations of

both of these replisome components in *Caulobacter*.

We find that DNA gyrase levels (subunits A and B) decrease upon MMC treatment in

*Caulobacter* (Table S2) but RNA levels do not appear to decrease when measured by

microarray (19, 22) or RNA-seq (30). DNA gyrase levels continue to decrease after

translational shutoff, indicating that the abundance of this protein may be regulated by

proteolysis (Table S2). In *E. coli* overexpression of a chromosomally encoded DNA

gyrase inhibitor *gyrI* increases survival to MMC (47, 48). It's possible that *Caulobacter*'s

downregulation of DNA gyrase is filling a similar role.

We also identified a novel DNA damage response factor, YaaA, which has previously

been shown to be important for survival to oxidative damage in *E. coli* (32). In

*Caulobacter*, we show that *yaaA* is important to the survival of the DNA damaging agents

MMC and UV light (Figure 2B-2D), indicating that its physiological relevance is not limited

to oxidative damaging conditions. It's unclear if *yaaA's* phenotypes in *Caulobacter* are

related to its role in regulating iron levels as seen in *E. coli* (32).

Previous work has suggested a correlation between level of expression during the DNA

damage response and rate of proteolysis (38). This proteomics survey supports this

model, as we see a negative correlation between protein abundance changes upon MMC

treatment and those upon translation shutoff (Figure 4C). Several DNA damage repair

proteins have been identified as proteolysis substrates in *E. coli,* including the LexA N-

terminal domain, UvrA, RecA, RuvB, and RecN (38). Of these, our proteomics data

provides evidence to support this categorization for the LexA N-terminal domain, UvrA,

and RecA but RuvB and RecN do not decrease in protein abundance upon translation

shutoff in our survey (Table S2).

This proteomics survey has identified examples of proteins that are regulated

independently of transcription during the DNA damage response, though it is unclear how

many of these are physiologically relevant to the response. We have also identified

instances where regulation of certain proteins during DNA damage in *E. coli* appears to

be conserved in *Caulobacter*, however the mechanisms of regulation appear to differ in

some of these cases. Finally, we have identified a novel DNA damage response factor

that has been overlooked in previous surveys that have focused on transcription alone.

This highlights the utility of combining transcriptomic and proteomic approaches to screen

for novel factors.

**The Lon protease and the DNA damage response**

Since DNA damage related Lon substrates UmuD (14) and SulA (15) discovered in *E.*

*coli* are not conserved in *Caulobacter*, it is unclear why *lon* is important to MMC survival

(21). We attempted to identify novel Lon substrates but did not find any potential

substrates that sufficiently explain this phenotype (Table S8).

The Δ*lon* strain exhibits global differences from the wild type strain in protein dynamics

that may account for the strain's sensitivity to MMC (Figure 4 and S6). We find that protein

abundance changes during the DNA damage response are dampened in the Δ*lon* strain

(Figure 4B and S6), which is at least partially due to the higher basal level expression of

DNA damage response genes in the strain (23). We also find that the correlation between

changes in protein abundance upon DNA damage treatment and those upon translation

shutoff is absent in the Δ*lon* strain (Figure 4C and 4D). These proteome-wide changes to

the DNA damage response may be at least partially responsible for the strain's sensitivity

to MMC, possibly due to a diminished range of response dynamics.

**Concluding thoughts**

In our proteomics analysis, we identified and quantified protein abundances for about

30% of the *Caulobacter* proteome. While some of the missing proteins are likely not

expressed during the growth conditions of our assays, some key DNA damage response

proteins are missing from our dataset. Some of these are known to be membrane-

associated (e.g. SidA (19) and DidA (22)), which may reduce our detection, and others

likely have relatively low abundance, even during DNA damage induction. Future work

using protocols designed specifically to observe these proteins could reveal additional

novel factors involved in DNA damage survival or reveal novel mechanisms of regulation

for known factors.

Our survey of protein dynamics during the DNA damage response allowed us to confirm

that, in general, protein abundance changes during the response are explained by

changes in RNA level. We also cataloged exceptions to this general rule and characterize

the novel DNA damage response factor, YaaA. Finally, though we did not identify novel

Lon substrates that explain this protease's importance to DNA damage survival, we did

identify systems-level differences in the size of protein abundance changes during the

response in the absence of *lon* which may play a role in this phenotype.

## METHODS

### Bacterial strains and growth conditions

Bacterial strains and plasmids used in this work are listed in Tables S9 and S10, respectively. All *Caulobacter* strains used in this work are NA1000 derivatives. *Caulobacter* strains were grown in PYE broth (2 g/L peptone, 1 g/L yeast extract, 1 mM $MgSO_4$, and 0.5 mM $CaCl_2$). Solid media was made with 1.5% agar. Knockout strains were created by replacing the native gene with a gentamycin resistance cassette using the pNPTS138 non-replicating vector (45). Venus- and mKate2-expressing strains were created by integrating pXGFPC-2 (49) based plasmids encoding constitutively expressed fluorescent proteins driven by a *lac* promoter (50) at the xylose locus.

### Bacterial growth curves

An overnight culture was diluted into PYE at an OD600 = 0.05. A volume of 200 µL of cells was added to wells of a 96-well plate. Plate was shaken continuously at 30ºC and OD600 was read every 20 minutes in a BioTek Epoch 2 Microplate Spectrophotometer (Agilent, Santa Clara, CA). MMC (0.5 µg/ml; Millipore Sigma, St. Louis, MO) was used to treat cells with DNA damage.

### Western blot

Western blots were performed using a commercially available rabbit anti-*E. coli* RecA primary antibody (1:10,000 dilution; Abcam, Cambridge, MA, 16 hours). Blots were visualized using the goat anti-rabbit Alexa Fluor™ Plus 800 secondary antibody (1:10,000 dilution; Licor, Lincoln, NE, 1 hour). Bands were quantified using ImageJ (51).

## Competition assays

Overnight cultures of each test strain and the control strain were diluted to an OD600 = 1.0, mixed 1:1, and serially diluted 30,000-fold. Undiluted cells were analyzed with a SpectraMax M2 microplate reader (Molecular Devices, San Jose, CA). Measurements were taken for cell density (OD600), Venus fluorescence (Excitation 490 nm, Cut off 515 nm, Emission 530 nm), and mKate2 fluorescence (Excitation 575 nm, Cut off 590 nm, Emission 630 nm). Diluted samples were grown to stationary phase (~48 hours, shaking at 30°C) and optical density and fluorescence measurements were taken again. The competitive index was calculated as the $\log_2$[((ratio test/control)END / (ratio test/control))START / ((average ratio wild type test/control)END/[(ratio wild type test/control))START].

## Sensitivity testing to ultra-violet light

Cells were grown to exponential phase and treated with 0 or 100 $J/m^2$ with a UV Stratalinker (Strategene, San Diego, CA, USA). Cells were diluted $10^0$-$10^{-7}$ and dilutions were plated on PYE agar. Plates were incubated for 48 hours at 30°C and then photographed.

## Transposon Sequencing

Transposon libraries were created as previously described (52, 53). A wild-type EZ-Tn5 library was treated in triplicate with mitomycin C (0.5 μg/mL) for 1 hour, outgrown for about 10 generations, and  sequenced. Data was analyzed using edgeR (54). Differential analysis of the counts in these libraries is available in Table S3.

**Sample collection and preparation for proteomics**

Pre-damage sample

Three biological replicates of overnight growth of each *Caulobacter* strain (wild type and

*Δlon)* were diluted to an OD600 = 0.0005 using PYE to a volume of 50 mL and grown for

approximately 10 generations to an OD600 of 0.5-0.6. All samples were normalized to an

OD600 of 0.5 using PYE to a volume of 25mL. A 5 mL pre-damage sample was taken,

and cells were collected by centrifugation (6,000 x g, 5min). Cell pellets were

resuspended in fresh lysis buffer (8 M urea, 50 mM HEPES pH7.5, 75 mM NaCl), flash

frozen, and stored at -80˚C.

Post-damage/pre-shutoff sample

Cells were treated with 0.5 µg/mL MMC for 1 hour before cells were collected by

centrifugation (6,000 x g, 10 min) and resuspended in fresh PYE broth to an OD600 = 0.5

to a volume of 25 mL. These treatment conditions are similar to those of the microarray

from the Laub lab (19) for simple comparison. Previous experiments indicate that the wild

type strain has about 55% survival to this treatment (45). After 2 hours, all samples were

normalized to an OD600 = 0.5 using PYE to a volume of 25 mL. A 5 mL post-damage

sample was taken, and cells were collected by centrifugation (6,000 x g, 5 min). Cell

pellets were resuspended in fresh lysis buffer, flash frozen, and stored at -80ᵒC.

Post-shutoff sample

Cells were treated with chloramphenicol (30 μg/mL) for 2 hours. A 5 mL post-shutoff

sample was taken, and cells were collected by centrifugation (6,000xg, 5 min). Cell pellets

were resuspended in fresh lysis buffer, flash frozen, and stored at -80˚C.

<u>Sample preparation for proteomics</u>

Samples were freeze-thawed three times in liquid nitrogen to aid cell lysis, then

centrifuged at 16,000 x g for 10 min at 4°C. Supernatant was carefully transferred to a

fresh 1.5mL tube. A Bradford assay was performed to quantify protein concentration.

Because the Bradford assay is sensitive to high urea concentrations, a small fraction of

each sample was diluted 4-fold for the Bradford assay. 50 μg of each sample was

transferred to a fresh 1.5mL tube. The final volume was adjusted to 50 μL using 100 mM

Triethylammonium bicarbonate (TEAB) buffer.

To reduce disulfide bonds, 5 μL 200 mM tris(2-carboxyethyl)phosphine (TCEP) was

added to each sample and incubated at 55˚C for 1 hour. To modify cysteine residues, 5

μL of 375 mM iodoacetamide was added to each sample and incubated 30 min at room

temperature in the dark. To precipitate protein, 6-volumes of cold acetone was added to

each sample, and they were stored overnight at -20˚C.

Samples were centrifuged at 8,000 x g for 10 min at 4˚C to collect precipitate. Supernatant

was carefully removed and discarded. Pellets were allowed to dry at room temperature

for 1 hour. Protein pelleted were resuspended in 100 μL 50 mM TEAB. Samples were

incubated overnight at 37˚C with 1.25 μg of trypsin (for a 2.5:100 ratio of trypsin to

protein).

**Tandem mass tagging for proteomics**

Samples were labelled using the TMT10plex™ Isobaric Label Reagent Set (ThermoFisher Scientific, Waltham, MA, Catalog #90110) according to the product instructions. Since each sample only contains 50 µg of peptides instead of the suggested 100 µg, half of each label was used per sample.

**Proteomics sample fractionation**

For each replicate experiment in each strain, equal volumes of each samples were combined into a single tube and then dried in a SpeedVac. Pooled samples were resuspended in 300 µL 0.1% TFA. Samples were dried in a SpeedVac a second time and resuspended in 300 µL 0.1% TFA. The pH of each sample was measured to confirm that it was between 2.5 - 3.5, which is critical for effective fractionation.

Pooled samples were fractionated using the Pierce™ High pH Reversed-Phase Peptide Fractionation Kit (ThermoFisher Scientific, Waltham, MA, Catalog #84868) according to the kit instructions. Fractions 1 and 5, 2 and 6, 3 and 7, as well as 4 and 8 were pooled together for a final total of 4 fractions per sample. Samples were dried and resuspended in 0.1% folic acid in water for mass spectrometry.

**Mass spectrometry for tandem mass tagging proteomics**

LC-MS analysis was performed, modified from (55) using an Easy-nLC 1000 nanoLC chromatography system interfaced to an Orbitrap Fusion mass spectrometer (ThermoFisher Scientific, Waltham, MA). Samples were pre-concentrated and desalted on to a C18 trap column prior to separation over a 180 min gradient from 0% to 50%

mobile phase B (A:0.1% formic acid in water, B:0.1% formic acid in acetonitrile) at 300

nL/min flow rate with a 75 μm x 15 cm PepMap RLSC column (ThermoFisher Scientific,

Waltham, MA).  Mass spectrometry parameters were as follows: ion spray voltage 2000V,

survey scan MS1 120k resolution with a 2 s cycle time, interleaved with data-dependent

ion trap MS/MS of highest intensity ions with charge state-dependent ion selection

window (z=2:1.2 m/z, z=3:0.7 m/z, z=4-6:0.5 m/z) and CID at 35% normalized collision

energy. Additionally, the top 5 (z=2) or 10 (z>2) product ions were synchronously selected

for HCD MS^3 at NCE 65% with Orbitrap detection to generate TMT tag intensities.

**Proteomics Data Analysis**

RAW data files were analyzed in Proteome Discoverer 2.4 (ThermoFisher Scientific,

Waltham, MA) using the SEQUEST search algorithm against *Caulobacter crescentus*

NA1000 (NCBI Reference Sequence: NC_011916.1) database downloaded from

uniprot.org. The search parameters used are as follows: 10 ppm precursor ion tolerance

and 0.4 Da fragment ion tolerance; up to two missed cleavages were allowed; dynamic

modifications of methionine oxidation and N-terminal acetylation. Peptide matches were

filtered to a protein false discovery rate of 5% using the Percolator algorithm. Peptides

were assembled into proteins using maximum parsimony and only unique and razor

peptides were retained for subsequent analysis. Protein quantitation based on TMT ion

abundance was performed using a co-isolation threshold of 75% and SPS match

threshold 65%. Each TMT channel was normalized to total peptide amount and then

abundance scaled to average 100% for all proteins.

Protein abundances were then normalized to the averages of the 0 hour samples (Table

S1). Protein ratios (Table S2) were calculated using edgeR (54).

**ACKNOWLEDGEMENTS**

This work was supported by NIH/NIGMS R35GM130320 to P. Chien. The mass

spectrometry in this paper was performed by Dr. Stephen Eyles at the University of

Massachusetts Amherst Mass Spectrometry Core Facility in Amherst, MA

(RRID:SCR_019063). The *ΔfzlC* strain was a generous gift from Erin Goley's lab at Johns

Hopkins University, Baltimore, MD.

**REFERENCES**

1.  Cirz RT, Chin JK, Andes DR, de Crécy-Lagard V, Craig WA, Romesberg FE. 2005.
    Inhibition of mutation and combating the evolution of antibiotic resistance. PLoS Biol 3:e176.

2.  Norton MD, Spilkia AJ, Godoy VG. 2013. Antibiotic resistance acquired through a DNA
    damage-inducible response in Acinetobacter baumannii. J Bacteriol 195:1335–1345.

3.  Beaber JW, Hochhut B, Waldor MK. 2004. SOS response promotes horizontal
    dissemination of antibiotic resistance genes. Nature 427:72–74.

4.  Ubeda C, Maiques E, Knecht E, Lasa I, Novick RP, Penadés JR. 2005. Antibiotic-induced
    SOS response promotes horizontal dissemination of pathogenicity island-encoded virulence
    factors in staphylococci. Mol Microbiol 56:836–844.

5.  Maiques E, Ubeda C, Campoy S, Salvador N, Lasa I, Novick RP, Barbé J, Penadés JR.
    2006. beta-lactam antibiotics induce the SOS response and horizontal transfer of virulence
    factors in Staphylococcus aureus. J Bacteriol 188:2726–2729.

6.  Hastings PJ, Rosenberg SM, Slack A. 2004. Antibiotic-induced lateral transfer of antibiotic
    resistance. Trends Microbiol 12:401–404.

7.  Little JW, Edmiston SH, Pacelli LZ, Mount DW. 1980. Cleavage of the *Escherichia coli lexA*
    protein by the *recA* protease. Proc Natl Acad Sci U S A 77:3225–3229.

8.  Horii T, Ogawa T, Nakatani T, Hase T, Matsubara H, Ogawa H. 1981. Regulation of SOS
    functions: purification of *E. coli* LexA protein and determination of its specific site cleaved by
    the RecA protein. Cell 27:515–522.

9.  Courcelle J, Khodursky A, Peter B, Brown PO, Hanawalt PC. 2001. Comparative gene

expression profiles following UV exposure in wild-type and SOS-deficient *Escherichia coli*.

Genetics 158:41–64.

10. Khil PP, Camerini-Otero RD. 2002. Over 1000 genes are involved in the DNA damage

response of *Escherichia coli*. Mol Microbiol 44:89–105.

11. Neher SB, Flynn JM, Sauer RT, Baker TA. 2003. Latent ClpX-recognition signals ensure

LexA destruction after DNA damage. Genes Dev 17:1084–1089.

12. Gonzalez M, Rasulova F, Maurizi MR, Woodgate R. 2000. Subunit-specific degradation of

the UmuD/D' heterodimer by the ClpXP protease: the role of trans recognition in UmuD'

stability. EMBO J 19:5251–5258.

13. Pruteanu M, Baker TA. 2009. Controlled degradation by ClpXP protease tunes the levels of

the excision repair protein UvrA to the extent of DNA damage. Mol Microbiol 71:912–924.

14. Gonzalez M, Frank EG, Levine AS, Woodgate R. 1998. Lon-mediated proteolysis of the

*Escherichia coli* UmuD mutagenesis protein: in vitro degradation and identification of

residues required for proteolysis. Genes Dev 12:3889–3899.

15. Canceill D, Dervyn E, Huisman O. 1990. Proteolysis and modulation of the activity of the cell

division inhibitor SulA in Escherichia coli lon mutants. J Bacteriol 172:7297–7300.

16. Kumaran NAM, Karthik M, Kumar V, Jebasingh T, Munavar MH. 2020. Two new mutations

in dnaJ suppress DNA damage hypersensitivity and capsule overproduction phenotypes of

Δlon mutant of Escherichia coli by modulating the expression of clpYQ (hslUV) and rcsA

genes. Gene 726:144135.

17. Howard-Flanders P, Simson E, Theriot L. 1964. A Locus That Controls Filament Formation

and Sensitivity to Radiation in Escherichia Coli K-12. Genetics 49:237–246.

18. da Rocha RP, Paquola AC de M, Marques M do V, Menck CFM, Galhardo RS. 2008.

Characterization of the SOS regulon of *Caulobacter crescentus*. J Bacteriol 190:1209–1218.

19. Modell JW, Hopkins AC, Laub MT. 2011. A DNA damage checkpoint in *Caulobacter*

*crescentus* inhibits cell division through a direct interaction with FtsW. Genes Dev 25:1328–

1343.

20. Gozzi K, Salinas R, Nguyen VD, Laub MT, Schumacher MA. 2022. ssDNA is an allosteric

regulator of the *C. crescentus* SOS-independent DNA damage response transcription

activator, DriD. Genes Dev 36:618–633.

21. Zeinert RD, Ogdahl JL, Liu J, Yang Q, Du Y, Barros BB, Freddolino PL, Haynes CM, Chien

P. 2022. A legacy role for DNA binding of Lon protects against genotoxic stress. bioRxiv

https://doi.org/10.1101/317677.

22. Modell JW, Kambara TK, Perchuk BS, Laub MT. 2014. A DNA Damage-Induced, SOS-

Independent Checkpoint Regulates Cell Division in *Caulobacter crescentus*. PLoS Biol

12:e1001977.

23. Mahmoud SA, Aldikacti B, Chien P. 2022. ATP hydrolysis tunes specificity of a AAA+

protease. Cell Reports 40:111405.

24. Bos J, Yakhnina AA, Gitai Z. 2012. BapE DNA endonuclease induces an apoptotic-like

response to DNA damage in Caulobacter. Proc Natl Acad Sci U S A 109:18096–18101.

25. Bhat NH, Vass RH, Stoddard PR, Shin DK, Chien P. 2013. Identification of ClpP substrates

in *Caulobacter crescentus* reveals a role for regulated proteolysis in bacterial development.

Molecular Microbiology 88:1083–1092.

26. Camberg JL, Hoskins JR, Wickner S. 2009. ClpXP protease degrades the cytoskeletal

protein, FtsZ, and modulates FtsZ polymer dynamics. Proceedings of the National Academy

of Sciences 106:10614–10619.

27. Barros BB, Mahmoud SA, Chien P, Zeinert RD. 2020. Degradation of Lon in Caulobacter

crescentus. J Bacteriol 203:e00344-20.

28. Liu J, Zeinert R, Francis L, Chien P. 2019. Lon recognition of the replication initiator DnaA

requires a bipartite degron. Mol Microbiol 111:176–186.

29. Bissonnette SA, Rivera-Rivera I, Sauer RT, Baker TA. 2010. The IbpA and IbpB small heat-

shock proteins are substrates of the AAA+ Lon protease. Mol Microbiol 75:1539–1549.

30. Fröhlich KS, Förstner KU, Gitai Z. 2018. Post-transcriptional gene regulation by an Hfq-

independent small RNA in *Caulobacter crescentus*. Nucleic Acids Res 46:10969–10982.

31. Prahlad J, Yuan Y, Lin J, Chang C-W, Iwata-Reuyl D, Liu Y, de Crécy-Lagard V, Wilson MA.

2020. The DUF328 family member YaaA is a DNA-binding protein with a novel fold. Journal

of Biological Chemistry 295:14236–14247.

32. Liu Y, Bauer SC, Imlay JA. 2011. The YaaA protein of the Escherichia coli OxyR regulon

lessens hydrogen peroxide toxicity by diminishing the amount of intracellular unincorporated

iron. J Bacteriol 193:2186–2196.

33. Price MN, Wetmore KM, Waters RJ, Callaghan M, Ray J, Liu H, Kuehl JV, Melnyk RA,

Lamson JS, Suh Y, Carlson HK, Esquivel Z, Sadeeshkumar H, Chakraborty R, Zane GM,

Rubin BE, Wall JD, Visel A, Bristow J, Blow MJ, Arkin AP, Deutschbauer AM. 2018. Mutant

phenotypes for thousands of bacterial genes of unknown function. 7706. Nature 557:503–

509.

34. Tomasz M. 1976. H2O2 generation during the redox cycle of mitomycin C and dna-bound

mitomycin C. Chem Biol Interact 13:89–97.

35. Ching C, Yang B, Onwubueke C, Lazinski D, Camilli A, Godoy VG. 2018. Lon Protease Has

Multifaceted Biological Functions in Acinetobacter baumannii. J Bacteriol 201:e00536-18.

36. Breidenstein EBM, Bains M, Hancock REW. 2012. Involvement of the Lon Protease in the

SOS Response Triggered by Ciprofloxacin in Pseudomonas aeruginosa PAO1. Antimicrob

Agents Chemother 56:2879–2887.

37. SaiSree L, Reddy M, Gowrishankar J. 2000. lon incompatibility associated with mutations

causing SOS induction: null uvrD alleles induce an SOS response in Escherichia coli. J

Bacteriol 182:3151–3157.

38. Neher SB, Villén J, Oakes EC, Bakalarski CE, Sauer RT, Gygi SP, Baker TA. 2006.

Proteomic Profiling of ClpXP Substrates after DNA Damage Reveals Extensive Instability

within SOS Regulon. Molecular Cell 22:193–204.

**Figure 1. Survey of *Caulobacter* DNA damage response distinguishes known DNA damage response factors and protease substrates.** (A) A western blot for RecA protein levels was used to determine peak protein induction (3 hours) to use for post-MMC timepoint for proteomics survey. Quantification of western blot for three replicates is shown, representative blot shown in Figure S1. RecA quantification was normalized to ClpP band and then to time 0 (Figure S1). (B) Samples for proteomics were taken before treatment (0 hours, pink line). Cells were treated with MMC for 1 hour and recovered in fresh medium for 2 additional hours before the post-treatment proteomics sample was taken (3 hours, teal line). At 3 hours chloramphenicol was added to shutoff translation and a post-shutoff sample was taken at 5 hours. (C) A volcano plot of proteomics data

compares the statistical confidence ($-\log_{10}$(false discovery rate)) versus the change in

protein abundance after MMC treatment $\log_2$(3 hrs/0 hrs). Several of the proteins that are

significantly upregulated after DNA damage treatment are known DNA damage response

proteins (highlighted in red). (D) A volcano plot of proteomics data compares the statistical

confidence ($-\log_{10}$(false discovery rate)) versus the change in protein abundance after

translation shutoff $\log_2$(5 hrs/3 hrs). Several of the proteins that are significantly

downregulated after chloramphenicol treatment are known protease substrates

(highlighted in red). FDR, false discovery rate.

**Figure 2. YaaA is upregulated upon MMC treatment and is important for survival to MMC, $H_2O_2$, and UV light.** (A) A comparison between our proteomics and previously published microarray experiment that uses similar MMC treatment conditions (19). Graph on the right is an enlargement of the red box in the graph on the left. Proteins highlighted in red are known DNA damage response genes that are upregulated at the RNA and

protein levels. YaaA is highlighted in blue. CCNA_02118, CCNA_00591, and

CCNA_00276 are highlighted in purple. Graph cropped to upper right quadrant to

highlight proteins that are upregulated. One gene is excluded (CCNA_03825) due to its

extremely high induction to make other points more visible. (B and C) When each mKate2-

expressing test strain is competed in co-culture against a wild type Venus-expressing

fluorescent control strain, only the *ΔyaaA* mutant shows a competitive disadvantage that

is exacerbated by the presence of (B) 0.1 µg/mL MMC or (C) 25 µM hydrogen peroxide.

A reciprocal experiment in which test strains are marked with Venus and wild type control

is marked with mKate2 is shown in Figure S3B and S3C. (D) The wild type and *ΔyaaA*

strains were exposed to 0 or 100 J/m$^2$ of ultra-violet light and 10-fold dilutions were plated.

The *ΔyaaA* strain is more sensitive to ultra-violet light than the wild type strain.

**Figure 3. Proteomics Survey in _Δlon_ strain distinguishes known Lon substrates.**

(A) A volcano plot of proteomics data compares the statistical confidence ($-\log_{10}$(false discovery rate)) versus the $\log_2$(3 hrs/0 hrs). Known DNA damage proteins shown in Figure 1B are highlighted here in red. ImuA was not identified in the proteomics of the _Δlon_ strain and so is absent. (B) A volcano plot compares the statistical confidence ($-\log_{10}$(false discovery rate)) versus the $\log_2$(5 hrs/3 hrs) for the _Δlon_ strain. Known Lon substrates DnaA and IbpA show a smaller decrease in protein abundance after translation shutoff than in the wild type strain (compare to Figure 1D). (C) Protein abundance of known Lon substrates DnaA and IpbA significantly increases upon MMC treatment (0-3

630    hours, pink and teal lines) and then decreases upon translation shutoff in a *lon*-dependent

631    manner (3-5 hours, teal and purple lines). Note that protein abundances are not directly

632    comparable between the two strains. FDR, false discovery rate.

**Figure 4. Protein abundance changes upon DNA damage are generally smaller in the Δ*lon* strain compared to in the wild type strain.** (A) A western blot to determine RecA protein levels was used to determine when to sample for protein abundance changes after DNA damage treatment in the Δ*lon* strain. Note that the wild type data here is repeated from Figure 1A for easy comparison. Quantification of western blot for three replicates is shown, representative blot shown in Figure S1. Sampling for proteomics was performed as described in Figure 1B. RecA quantification was normalized to ClpP band

and then to time 0 (see Figure S1). (B) Comparison of the $\log_2$(3 hrs/0 hrs) for the $\Delta lon$

versus wild type strain shows that fold changes of protein abundance are generally lower

in the $\Delta lon$ strain. Blue dashed line represents an y=x line. The red line represents the

trendline of the data with 95% CI drawn (slope = 0.45). The inset plots $\log_2$(3 hrs/0 hrs)

for all proteins that are significantly upregulated (FDR < 0.05). Comparison of protein

changes upon MMC treatment and translational shutoff indicates a negative correlation

in the wild type strain (C) that is absent in the (D) $\Delta lon$ strain.