A dynamical computational model of theta generation in hippocampal circuits to study theta-gamma oscillations during neurostimulation

Nikolaos Vardalakis[1,2]**, Amélie Aussel**[1,2,3†]**, Nicolas P. Rougier**[1,2,3†]**, Fabien B. Wagner**[1*†]

*For correspondence:
fabien.wagner@u-bordeaux.fr
(Fabien B. Wagner)

[†]These authors contributed equally to this work

[1]Univ. Bordeaux, CNRS, IMN, UMR 5293, F-33000 Bordeaux, France; [2]Univ. Bordeaux, INRIA, IMN, UMR 5293, F-33000 Bordeaux, France; [3]Univ. Bordeaux, CNRS, Bordeaux INP, LaBRI, UMR 5800, F-33400 Talence, France

Abstract

Neurostimulation of the hippocampal formation has shown promising results for modulating memory but the underlying mechanisms remain unclear. In particular, the effects on hippocampal theta-nested gamma oscillations and theta phase reset, which are both crucial for memory processes, are unknown. Moreover, these effects cannot be investigated using current computational models, which consider theta oscillations with a fixed amplitude and phase velocity. Here, we developed a novel computational model that includes the medial septum, represented as a set of abstract Kuramoto oscillators producing a dynamical theta rhythm with phase reset, and the hippocampal formation, composed of biophysically-realistic neurons and able to generate theta-nested gamma oscillations under theta drive. We showed that this system can exhibit bistability in a specific range of parameters and that a single stimulation pulse could switch the network behavior from non-oscillatory to a state producing theta-nested gamma oscillations. Next, we demonstrated that for a theta input too weak to generate theta-nested gamma oscillations, pulse train stimulation at the theta frequency could restore seemingly physiological oscillations. Importantly, the presence of phase reset influenced whether these two effects depended on the phase at which stimulation onset was delivered, which has practical implications for designing neurostimulation protocols that are triggered by the phase of ongoing theta oscillations. This novel model opens new avenues for studying the effects of neurostimulation on the hippocampal formation. Furthermore, our hybrid approach that combines different levels of abstraction could be extended in future work to other neural circuits that produce dynamical brain rhythms.

Introduction

Neurostimulation methods have emerged as promising therapeutic modalities to restore neurological functions in a broad range of motor and cognitive disorders (*Gupta et al., 2023*). In the context of learning and memory, deep brain stimulation (DBS) of the entorhinal area or hippocampus has been shown to either enhance (*Suthana et al., 2012*; *Suthana and Fried, 2014*; *Titiz et al., 2017*; *Jun et al., 2019*) or disrupt (*Jacobs et al., 2016*; *Goyal et al., 2018*; *Lozano et al., 2016*) memory encoding. These conflicting results may originate from differences in experimental protocols and

from a poor understanding of their biophysical underpinnings. Among such mechanisms, the involvement of hippocampal theta oscillations (4-12 Hz) and their interactions with higher-frequency gamma oscillations (30-120 Hz) in memory-related processes has been reported in multiple studies (*Lisman et al., 2005*; *de Almeida et al., 2007*; *Lega et al., 2012*; *Lin et al., 2017*; *Malkov et al., 2021*; *Abbaspoor et al., 2022*). Moreover, the modulation of gamma oscillations by the phase of theta oscillations in hippocampal circuits, a phenomenon termed theta-gamma phase-amplitude coupling (PAC), correlates with the efficacy of memory encoding and retrieval (*Jensen and Colgin, 2007*; *Tort et al., 2009*; *Canolty and Knight, 2010*; *Axmacher et al., 2010*; *Fell and Axmacher, 2011*; *Lisman and Jensen, 2013*; *Lega et al., 2016*). Neurostimulation protocols that affect these rhythms, such as theta burst stimulation (*Titiz et al., 2017*), have also been shown to optimally induce long-term potentiation (LTP) (*Larson and Munkácsy, 2015*). Another potential mechanism underlying the effect of hippocampal neurostimulation might be the reset of the phase of theta oscillations in response to exogenous inputs, such as a novel sensory input or a pulse of electrical stimulation applied to the fornix or perforant path (*Buño et al., 1978*; *Williams and Givens, 2003*). Theta phase reset is known to facilitate LTP (*McCartney et al., 2004*) and naturally occurs during both encoding and retrieval of associative memories (*Kota et al., 2020*). In this context, the design of a computational model that replicates memory-related theta-gamma oscillations and theta phase reset is of uttermost importance to investigate the effects of electrical stimulation on the hippocampal formation and possibly optimize neurostimulation protocols for memory improvement.

Models of memory-related theta-gamma oscillations in the hippocampal formation have been developed across different resolution levels, ranging from abstract mean-field approaches (*Traub et al., 1997*; *Onslow et al., 2014*; *Segneri et al., 2020*) to biophysically-realistic conductance-based models (*Lundqvist et al., 2006*; *Herman et al., 2013*; *Aussel et al., 2018*). Neural masses, which represent the mean activity of a neuronal population, can generate gamma oscillations through reciprocal interactions between an excitatory and inhibitory population (*Traub et al., 1997*; *Onslow et al., 2014*), or even using a self-projecting inhibitory population (*Segneri et al., 2020*). Under excitatory oscillatory input at theta frequencies, these models are capable of generating theta-nested gamma oscillations. Similarly, these oscillations can be observed in more complex models of the hippocampal formation composed of single-compartment excitatory and inhibitory neurons connected through conductance-based synapses following the Hodgkin-Huxley formalism (*Hodgkin and Huxley, 1952*), and driven by a fixed oscillatory theta input (*Aussel et al., 2018*). Theta-nested gamma oscillations also appear in multi-compartment models of prefrontal cortex activity during memory retrieval (*Lundqvist et al., 2006*; *Herman et al., 2013*). Finally, several models have investigated the functional link between the hippocampal theta rhythm and memory processes, showing that encoding and retrieval occur at different phases within each theta cycle through phasic changes in neuronal activity and LTP (e.g. (*Hasselmo et al., 2002*; *Cutsuridis et al., 2010*)).

In terms of neurostimulation, most computational work has focused on the mechanisms underlying DBS of the basal ganglia for motor disorders such as Parkinson's disease (*Rubin and Terman, 2004*; *Pirini et al., 2009*; *Mina et al., 2013*; *Ebert et al., 2014*), peripheral nerve stimulation (*Rattay et al., 2003*; *Kipping and Nogueira, 2022*), spinal cord stimulation (*Rattay et al., 2000*; *Capogrosso et al., 2013*), or has remained generic (*Basu et al., 2018*). However, models investigating neurostimulation of hippocampal circuits are scarce (*Hendrickson et al., 2016*; *Bingham et al., 2018*), and do not take into account the effects on theta-gamma oscillations. For example, a detailed multicompartment model of the rat dentate gyrus was able to replicate experimentally recorded local field potentials induced by different electrode placements and pulse amplitudes during stimulation of the perforant path (*Bingham et al., 2018*). To model the impact of neurostimulation on neuronal oscillations, a more abstract formalism based on Kuramoto phase oscillators (*Kuramoto, 1984*) has been introduced in the context of Parkinson's disease and essential tremor, which enabled the design of novel neurostimulation paradigms that enhance or disrupt neuronal synchrony in basal ganglia circuits (*Tass, 2003*; *Ebert et al., 2014*; *Asllani et al., 2018*; *Weerasinghe et al., 2019*).

To our knowledge, there is currently no model of the hippocampal formation that is able to

replicate both theta-gamma PAC and theta phase reset during neurostimulation. To investigate
the effects of neurostimulation on hippocampal circuits while taking into account these two mech-
anisms, we modified an existing biophysical model of the hippocampal formation (*Aussel et al.,*
2018). In this original model, however, the theta rhythm was considered as an oscillatory input of
fixed amplitude and phase velocity, which is inconsistent with theta phase reset. To circumvent this
limitation, we combined this biophysical model with abstract Kuramoto oscillators that acted as a
dynamical source of theta rhythm, thereby modeling medial septum inputs to the hippocampal for-
mation. This new hybrid dynamical model could generate both theta-nested gamma oscillations
and theta phase reset, following a particular phase response curve (PRC) inspired by experimental
literature (*Lengyel et al., 2005*; *Akam et al., 2012*; *Torben-Nielsen et al., 2010*).

We then leveraged this model to explore the effect of single-pulse and pulse-train stimulation
on theta-gamma oscillations. In the absence of theta input from the medial septum, single-pulse
stimulation produced a transient effect consisting of one or several bursts of activity, depending
on stimulation amplitude. The presence of multiple bursts depended on the single-cell calcium dy-
namics and M-type potassium adaptation current. In the presence of weak theta input, designed
to mimic a pathological state that impairs theta-gamma oscillations, single-pulse stimulation could
produce long-lasting or even persistent activity by switching the network to a highly synchronized
state characterized by theta-nested gamma oscillations. When phase reset was not included in the
model, this effect was more pronounced when the stimulation pulse was delivered at the peak of
the theta rhythm. However, when strong theta reset was considered, the phase at which stimula-
tion was delivered did not influence the outcome. In the presence of an even weaker theta input,
mimicking a pathological state that completely abolishes theta-gamma oscillations, only pulse train
stimulation could restore physiological theta-gamma oscillations during the stimulation period. As
for the previous results, this effect was phase-dependent only when theta reset was not included.

These results provide a new framework to interpret neurostimulation interventions that inter-
fere with hippocampal oscillations and aim at improving memory function. It can be further ex-
tended to investigate the effects of more complex neurostimulation protocols, and the impact of
stimulation location and amplitude on the observed network dynamics.

Results

A computational model of the hippocampal formation with dynamical theta input

We first developed a computational model of hippocampal circuits, able to generate both theta-
nested gamma oscillations and theta phase reset. To achieve this, we combined an existing con-
ductance-based model of the hippocampal formation (*Aussel et al., 2018*) with a set of Kuramoto
phase oscillators (*Kuramoto, 1984*) that were used to model the dynamical theta input originating
from pacemaker neurons in the medial septum (*Wang, 2002*) (*Figure 1*A). The model contained ex-
citatory and inhibitory neuronal populations in the entorhinal cortex (EC), dentate gyrus (DG), CA3
and CA1 fields within a coronal slice of the human hippocampus (*Figure 1*B-C). The output of the
hippocampal formation (i.e. the activity of CA1 pyramidal neurons) was provided as an input to the
Kuramoto oscillators, simulating the hippocampal-septal projections through the fornix (*Williams*
and Givens, 2003; *Nuñez and Buño, 2021*; *Takeuchi et al., 2021*). The oscillations produced by
the collective behavior of Kuramoto oscillators represented a population average of their activity:
highly synchronized and desynchronized states generated respectively high and low amplitude
theta oscillations (*Figure 1*D). The number of oscillators was chosen so that their synchronization
level (i.e. their order parameter) and their frequency distribution were sufficiently close to their
asymptotic behavior for a large number of oscillators (*Figure S1*).

As in the original model (*Aussel et al., 2018*), the input theta rhythm drove the network to pro-
duce spiking activity preferentially around the peak of theta in all excitatory and inhibitory popu-
lations (*Figure 2*A and *Figure S2*). Spectrograms of the CA1 population firing rates revealed that
these bursts of activity around each theta peak were characterized by oscillations around 60 Hz

(*Figure 1*A). This was confirmed by power spectral densities, which showed a clear peak at 4 Hz (corresponding to the theta drive) and increases in gamma band activity between 40 and 80 Hz (*Figure 2*B). To quantify PAC between theta and gamma oscillations, we used the modulation index (MI) (*Tort et al., 2008*, *2010*), which has been shown to outperform other similar measures (*Hülsemann et al., 2019*). However, we discovered that this metric would give erroneous results in our simulated datasets due to the absence of certain frequency components. To overcome this limitation and avoid artifacts, uniform noise was added to the firing rates prior to computing the MI (*Figure S3*). We first visualized the quantification of PAC using the comodulogram, which indicates the MI as a function of two frequencies corresponding to the modulating phase signal and the modulated amplitude signal. This analysis confirmed that gamma-band signals between 45 and 75 Hz were modulated by lower theta frequencies between 3 and 6 Hz, which we quantified by computing a global MI in frequency ranges encompassing theta (3-9 Hz) and gamma (40-80 Hz) (*Figure S2*B). To refine this analysis, we also computed a similar MI between the amplitude of various gamma frequencies and the phase of theta, which indicated that oscillations between 40 and 80 Hz occur preferentially around the peak of theta (*Figure 2*C and *Figure S2*B).

Next, we verified that our model was able to display theta phase reset during single-pulse stimulation (depolarizing pulse, 1-ms duration). This mechanism is tightly linked with the concept of PRC, which characterizes the phase delay or advancement that follows a single pulse delivered to an oscillatory system, as a function of the phase at which this input is delivered. Although there is no direct measurement of the PRC of septal neurons, such characterizations have been performed for individual pyramidal cells in the CA3 and CA1 fields of the hippocampus (*Lengyel et al., 2005*; *Kwag and Paulsen, 2009*; *Akam et al., 2012*). These PRCs appear biphasic and show a phase advancement (respectively delay) for stimuli delivered in the ascending (respectively descending) slope of theta. We modeled this behavior by a specific term (which we called the phase response function) in the general equation of the Kuramoto oscillators (see methods, *Equation 1*). Importantly, introducing a phase offset in the phase response function disrupted theta-nested gamma oscillations (*Figure S4*), which suggests that the septohippocampal circuitry must be critically tuned to be able to generate such oscillations. The strength of phase reset could also be adjusted by a gain that was manually tuned. In the presence of the physiological phase response function and of a sufficiently high reset gain, a single stimulation pulse delivered to all excitatory and inhibitory CA1 neurons could reset the phase of theta to a value close to its peaks (*Figure 2*A). We computed the PRC of our simulated data for different stimulation amplitudes and validated that our neuronal network behaved according to the phase response function set in our Kuramoto oscillators (*Figure 2*D). It should be noted that including this phase reset mechanism affected the generated theta rhythm even in the absence of stimulation, extending the duration of the theta peak and thereby slowing down the frequency of the generated theta rhythm.

Overall, we successfully developed a new model of the hippocampal formation able to exhibit theta-nested gamma oscillations and theta phase reset in response to stimulation. We then decided to explore further the effects of various stimulation protocols on its dynamics.

Effects of theta input and stimulation amplitudes on theta-gamma oscillations

We investigated the behavior of the model across multiple states of varying theta input amplitude in response to single-pulse stimulation delivered to CA1. In the absence of theta drive, a single stimulation pulse elicited either zero or two bursts of spiking activity (depending on stimulation amplitude) separated by about 200 ms (*Figure 3*A, C). We first sought to understand the origin of this second burst, as it showed that even a single pulse could induce transient periodic activity around 5 Hz in our model, a frequency within the theta range. A previous model has shown that the presence of Calcium-Activated Non-specific cationic (CAN) currents can lead to self-sustained theta oscillations in the hippocampus (*Giovannini et al., 2017*). Moreover, this study showed a direct link between the increased excitation provided by the CAN current and the spike-frequency adaptation properties of the M-current, directly affecting transitions from an asynchronous low-firing regime

to synchronous bursting. We tested the role of the CAN current in the response to single-pulse stimulation by completely removing it from our simulations, which abolished the second burst of activity (*Figure 3*B). Moreover, the time interval separating the two bursts likely resulted from the interplay between the depolarizing CAN current and hyperpolarizing M current (*Figure 3*A).

In the presence of theta input, the effects of single-pulse stimulation depended both on theta drive amplitude and stimulation amplitude, highlighting different regimes of network activity (*Figure 4* and *Figure S5*, *Figure S6*, *Figure S7*). For low theta input, theta-nested gamma oscillations were initially absent and could not be induced by stimulation (*Figure 4*A, B). At most, the stimulation could only elicit a few bursts of spiking activity that faded away after approximately 500 ms (*Figure 4*B). For increasing theta input, the network switched to a bistable regime: upon initialization at a state with no spiking activity, it could be kicked to a state with self-sustained theta-nested gamma oscillations by a single stimulation pulse of sufficiently high amplitude (*Figure 4*C). This regime existed for a range of theta inputs and stimulation amplitudes, as characterized by the post-stimulation theta power, gamma power, and theta-gamma PAC (*Figure 4*D). For the highest theta input, the network became monostable, spontaneously generating theta-nested gamma oscillations even when initialized at a state with no spiking activity (*Figure 4*D, last stimulation amplitude). Additionally, we investigated the effects of various synchronization levels of the Kuramoto oscillators on the production of theta-gamma oscillations and their response to single-pulse stimulation. The results showed that the production of theta-nested gamma oscillations depended mostly on the theta input amplitude and less on the synchronization level (*Figure S8*). Therefore, we decided to use the amplitude of the theta input as the main variable controlling whether the neuronal activity was within a physiological or pathological state.

Neurostimulation for restoring theta-gamma oscillations in pathological states

Based on the above analyses, we considered two pathological states: one with a moderate theta input associated with a bistable regime that allowed self-sustained oscillations (*Figure 4*, point C), and one with a weak theta input characterized by the complete absence of self-sustained oscillations (*Figure 4*, point B). In each case, we sought to assess whether single-pulse or pulse train stimulation could induce or restore theta-nested gamma oscillations and whether this effect depended on the phase at which stimulation was delivered (i.e. at the peak or trough of the theta cycle). We hypothesized that any possible phase relationship would also depend on the phase reset mechanism. To test this hypothesis, we ran a series of simulations using two different models: one without phase reset and one with strong phase reset (i.e. the reset gain was set at the value used in *Figure 2*).

In the case of a bistable network (moderate theta input) with phase reset, delivering a pulse at either the peak or trough of theta could induce theta-nested gamma oscillations (*Figure 5*A, C). By contrast, in the absence of phase reset, only stimulation delivered at the peak of theta was able to induce such oscillations (*Figure 5*B, D). Quantification of these results in terms of theta power, gamma power, and theta-gamma PAC showed strongly similar responses between stimulation delivered at the peak or trough of theta with phase reset enabled, similar but weaker responses with stimulation delivered at the peak of theta with phase reset disabled, and no response in the case of trough stimulation with phase reset disabled (*Figure 5*E).

In the case of a weak theta input that completely abolished neuronal oscillations, we delivered stimulation pulses continuously at a frequency matching that of the underlying theta rhythm for a duration of 2 s, in order to restore physiological oscillations (*Figure 6*). The stimulation onset was timed to either the peak or trough of the ongoing theta cycle. The continuous delivery of pulses produced similar results as single-pulse stimulation. With phase reset, pulse train stimulation restored theta-nested gamma oscillations within the whole network, irrespective of the phase of stimulation onset (*Figure 6*A, C). There was a slight difference between the responses caused by the first pulse, as trough stimulation missed one cycle before producing network-wide oscillations. Notably, stimulation delivered at the trough forced a reset of the phase of theta rhythm and led

²⁴⁰ to the subsequent delivery of pulses at the peak of theta. Interestingly, in the absence of phase
²⁴¹ reset, only peak-targeted stimulation induced theta-gamma oscillations in all network areas (*Figure 5*B, D). Comparing these simulations based on theta power, gamma power, and theta-gamma
²⁴² *ure 5*B, D). Comparing these simulations based on theta power, gamma power, and theta-gamma
²⁴³ PAC within CA1 (*Figure 6*E) showed similar differences as single-pulse stimulation, although less
²⁴⁴ striking because CA1 was driven directly by the stimulation. However, these differences were even
²⁴⁵ more pronounced in areas other than CA1, as can be seen by the complete absence of oscillations
²⁴⁶ for trough stimulation without phase reset (*Figure 6*D).

²⁴⁷ Discussion

²⁴⁸ Highlights

²⁴⁹ In summary, we have developed a novel computational model to investigate the effects of elec-
²⁵⁰ trical stimulation on a slice of the hippocampal formation, incorporating two important features
²⁵¹ related to memory: theta-nested gamma oscillations and theta phase reset. The key innovation
²⁵² compared to previous models (e.g. *Aussel et al.* (*2018*)) is the introduction of a set of abstract Ku-
²⁵³ ramoto oscillators, which represent pacemaker neurons in the medial septum and are interfaced
²⁵⁴ with biophysically realistic neuronal models in the hippocampus. From a methodological point of
²⁵⁵ view, this hybrid interfacing between two levels of abstraction represents an innovation in itself
²⁵⁶ and could be applied to other systems or brain structures that are driven by dynamical rhythms.
²⁵⁷ The main outcomes reported here relate to the importance of the theta reset mechanism when
²⁵⁸ examining the effects of neurostimulation on hippocampal oscillations.

²⁵⁹ A very interesting finding consists in the bistable nature of the model for certain values of the
²⁶⁰ theta amplitude (*Figure 4*). In the bistable regime, a single stimulation pulse could switch the net-
²⁶¹ work behavior from a state with no spiking activity to one with theta-nested gamma oscillations.
²⁶² Whether such bistability exists in vivo in the context of memory processes remains an open ques-
²⁶³ tion. Nevertheless, delivering a single stimulation pulse bilaterally to the human hippocampus dur-
²⁶⁴ ing a memory task is sufficient to impair memory encoding (*Lacruz et al., 2010*), suggesting that
²⁶⁵ even single-pulse stimulation can indeed have wide network effects that are behaviorally relevant.

²⁶⁶ The second main finding is that the timing of individual pulses with respect to the phase of
²⁶⁷ the ongoing theta rhythm matters differently depending on the presence or absence of phase re-
²⁶⁸ set (*Figure 5* and *Figure 6*). Human intracranial stimulation data indicate that the receptivity of
²⁶⁹ hippocampal circuits to single-pulse stimulation is modulated by the phase of theta (*Lurie et al.,*
²⁷⁰ *2022*). To our knowledge, however, the degree of phase reset that follows each stimulation pulse
²⁷¹ remains unknown and should be investigated in future experimental studies. From a technological
²⁷² point of view, the two regimes with and without phase reset have opposite predictions concerning
²⁷³ the need for closed-loop stimulation protocols that would trigger stimulation in real-time based
²⁷⁴ on the phase of ongoing theta oscillations. Such phase-triggered stimulation would be most use-
²⁷⁵ ful if the phase reset mechanism remains relatively limited. When this mechanism becomes too
²⁷⁶ strong, regular continuous stimulation appears sufficient to restore physiological theta-gamma
²⁷⁷ oscillations (*Figure 6*).

²⁷⁸ It should also be noted that the reset gain in our simulations (*Equation 1*, gain G_{reset}) was either
²⁷⁹ completely turned off or set at a high value, producing a strong effect that always reset the phase
²⁸⁰ of theta to its peak. In reality, the degree of theta phase reset is dynamic, depending on the en-
²⁸¹ vironment and associated task requirements (*Rizzuto et al., 2003*; *Mormann et al., 2005*; *Jackson*
²⁸² *et al., 2008*), and may be affected by neurodegenerative disorders that affect the connections be-
²⁸³ tween the hippocampal formation and the medial septum. Importantly, our proposed framework
²⁸⁴ can simulate intermediate values of the reset gain, which should ideally be fitted to experimental
²⁸⁵ data in future applications.

²⁸⁶ Finally, we modeled pathological states by reducing the maximum amplitude of the theta in-
²⁸⁷ put (*Equation 2*, gain G_{θ}) until theta-nested gamma oscillations were impaired or even abolished
²⁸⁸ (*Figure 4*). This choice was meant to simulate neurodegeneration in the medial septum, which is

known to be affected in Alzheimer's disease, leading to oscillatory disruptions (*Nelson et al., 2014*; *Hampel et al., 2018*; *Takeuchi et al., 2021*). Another possibility would be that neurodegeneration limits the ability of the septal pacemaker neurons to synchronize, thus producing a weaker collective theta rhythm without affecting the maximum amplitude of individual oscillators. Although we simulated this change in our model by reducing the synchronization parameter, the effects on hippocampal oscillations were more mitigated (*Figure S8*). Linking these different modeling parameters to experimental biomarkers will be important in future work.

Limitations

Even though we took great care in developing a precise representation of the hippocampal formation, the resulting model remains a simplification that could be further enriched. In particular, we deliberately modeled only a single theta generator, while multiple intra- and extra-hippocampal generators are known to co-exist (*Kocsis et al., 1999*). We decided to model septal pacemaker neurons projecting to the EC as the main source of hippocampal theta as reported in multiple experimental studies (*Buzsáki, 2002*; *Buzsáki et al., 2003*; *Hangya et al., 2009*; *Colgin, 2013*). However, previous models have also proposed that direct septal inputs are not essential for theta generation (*Wang, 2002*; *Mysin et al., 2019*), but play an important role in phase synchronization of hippocampal neurons. To account for these discrepancies, our model could be extended by considering more realistic connectivity patterns between the medial septum and the hippocampal formation, including glutamatergic, cholinergic, and GABAergic reciprocal connections (*Müller and Remy, 2018*), or by considering multiple sets of oscillators each representing one theta generator.

Another limitation of our model concerns synaptic transmission delays, which have been largely neglected and could affect the phase relationships between the medial septum and different hippocampal subfields. Experimental studies have indeed reported time delays in the population activities of connected anatomical structures (i.e. from EC to DG) (*Mizuseki et al., 2009*), with pyramidal cells in downstream areas like CA3 and CA1 preferentially firing at different phases of theta (*Dragoi and Buzsáki, 2006*). Propagation effects could also depend on the spatial scale of the model. We also decided to represent only a thin coronal slice of the hippocampal formation, and it remains unclear how an anatomically-accurate model of the whole structure would behave in terms of propagation of spontaneous and electrically-induced neuronal activity.

Importantly, we did not consider learning through synaptic plasticity, even though such mechanisms could drastically modify synaptic conduction for the whole network (*Borges et al., 2017*). Even more interestingly, the inclusion of spike-timing-dependent plasticity would enable the investigation of stimulation protocols aimed at promoting LTP, such as theta-burst stimulation (*Larson and Munkácsy, 2015*). This aspect would be of uttermost importance to make a link with memory encoding and retrieval processes (*Axmacher et al., 2006*; *Tsanov and Manahan-Vaughan, 2009*; *Jutras et al., 2013*) and with neurostimulation studies for memory improvement (*Titiz et al., 2017*; *Solomon et al., 2021*).

From the point of view of neurostimulation, future work is needed to extend the current model to a multi-compartment representation of neurites, since axons are known to be preferentially activated by extracellular electrical stimulation (*Rattay et al., 2003*). Specifically, multi-compartment cable models have been developed to investigate spike initiation and propagation by modeling the axon as a series of resistance-capacitance circuits following its trajectory (*Rattay et al., 2003*; *Joucla and Yvert, 2012*; *Ashida and Nogueira, 2018*). These models are particularly suited to study the effects of extracellular stimulation, which are non-intuitive and depend as a first approximation on the second spatial derivative of the electrical potential along the cell membrane (*Rattay, 1986*; *Rattay et al., 2003*; *McIntyre et al., 2004*; *Rattay et al., 2018*). Here, we have modeled electrical stimulation as an intracellular current applied equally across all neurons in the targeted area, which is extremely simplified but enables computational tractability. Future developments will focus on developing an equivalent multicompartment model with realistic axonal trajectories while making sufficient simplifications to allow for realistic computation times.

Outlook

Overall, this new model of the hippocampal formation represents a methodologically innovative basis to further explore multiple neurostimulation strategies that target hippocampal oscillations. Moreover, the limitations discussed above represent important avenues for future refinements, which will require significant work to overcome the costs in terms of computational tractability (e.g. modeling the whole hippocampus, or using multicompartment models of axonal trajectories and dendritic trees). Ultimately, such model refinements should allow the investigation of the effects of extracellular stimulation using charge-balanced biphasic pulses and multipolar electrode configurations with realistic electrode geometries. Most importantly, we believe that our current work may also serve as an inspiration for future computational models of oscillopathies (not necessarily limited to the hippocampus), which could benefit from interfacing abstract sets of synchronizing oscillators and investigating their interactions with biophysically-realistic neurons.

Materials and Methods

Computational Model

Overall Architecture

We aimed to develop a computational model of the hippocampal formation that is able to generate theta-nested gamma oscillations and takes into account the dynamic nature of the theta input from the medial septum, i.e. the reset of the theta phase following strong activity in the perforant path or fornix (*Buño et al., 1978*; *Williams and Givens, 2003*). To this end, we adapted a previous biophysical model of the human hippocampal formation under fixed sinusoidal input (*Aussel et al., 2018*) and interconnected it with an abstract representation of the medial septum, modeled as an assembly of Kuramoto oscillators (*Kuramoto, 1984*) (*Figure 1*).

More precisely, we modeled a 15-mm thick coronal slice of the hippocampal formation which comprises the DG, the CA3 and CA1 subfields of the hippocampus, and the EC, all populated with excitatory and inhibitory Hodgkin-Huxley neurons. Biologically, GABAergic neurons from the medial septum project to the EC, CA3, and CA1 fields of the hippocampus (*Toth et al., 1993*; *Hajós et al., 2004*; *Manseau et al., 2008*; *Hangya et al., 2009*; *Unal et al., 2015*; *Müller and Remy, 2018*). Although the respective roles of these different projections are not fully understood, previous computational studies have suggested that the direct projection from the medial septum to CA1 is not essential for the production of theta in CA1 microcircuits (*Mysin et al., 2019*). Since our modeling of the medial septum is only used to generate a dynamic theta rhythm, we opted for a simplified representation where the medial septum projects only to the EC, which in turn drives the different subfields of the hippocampus. In our model, Kuramoto oscillators are therefore connected to the EC neurons and they receive projections from CA1 neurons (see sections below for more details).

Medial septum: Kuramoto Oscillators

The medial septum contains pacemaker neurons (*Varga et al., 2008*; *Hangya et al., 2009*) that synchronize with one another and generate a global theta rhythm. In turn, these pacemaker neurons drive downstream areas that receive projections from the medial septum. Here, we modeled the entire medial septum neuronal assembly as a set of coupled phase oscillators following the Kuramoto model (*Kuramoto, 1984*), which generated the driving theta rhythm that was provided to the hippocampal formation through the EC (*Breakspear et al., 2010*).

Kuramoto oscillators have also been used to investigate the effects of neurostimulation on synchronized brain rhythms in the context of Parkinson's disease or essential tremor (*Tass, 2003*; *Weerasinghe et al., 2019*). Here, each Kuramoto oscillator was described by its phase θ_i, which evolves over time according to the following equation:

$$\frac{d\theta_i}{dt} = \omega_i + \frac{k}{N} \sum_{j=1}^{N} \sin(\theta_j - \theta_i) + G_{reset} X(t) Z(\theta_i) \tag{1}$$

The term ω_i denotes the natural frequency of oscillator i, and is normally distributed around the center frequency f_0 and with standard deviation σ. The synchronization parameter k represents the coupling strength of the group of oscillators, and N is the number of oscillators. Higher values for the synchronization parameter indicate stronger coupling between pairs of oscillators, thus affecting how fast or slow their phases tend to synchronize. The final product $G_{reset}X(t)Z(\theta_i)$ is used to describe the effects of external inputs on the phase of the oscillators. Here, the only external input originates in the projections from CA1 to the medial septum. It is described by the instantaneous firing rate $X(t)$ of the CA1 excitatory population. G_{reset} is an arbitrary gain that determines the strength of the CA1 input to the oscillators. Finally, the function $Z(\theta_i)$ describes how the effects of CA1 inputs on the oscillators depend on the phase of the ongoing theta rhythm.

The coherence and phase of the driving theta rhythm I_θ were computed using the order parameter r to extract the mean amplitude A and the mean phase ϕ of the ensemble of Kuramoto oscillators as follows:

$$\begin{cases} r(t) & = \frac{1}{N} \sum_{i=1}^{N} e^{j\,\theta_i(t)} \\ A(t) & = \mathrm{Re}(r) \\ \phi(t) & = \mathrm{Im}(r) \\ I_\theta(t) & = G_\theta\,A(t)\,\frac{\cos(\phi(t))+1}{2} \end{cases} \tag{2}$$

where the mean amplitude $A(t)$ and the mean phase $\phi(t)$ are derived by taking the real and imaginary part of the order parameter respectively, and the output theta rhythm $I_\theta(t)$ is a rectified cosine, multiplied by a gain G_θ that controls the maximum output amplitude (in nA).

To simulate the effect of the projections from CA1 to the medial septum on the phase of the oscillators, an approximation of the instantaneous firing rate of the CA1 excitatory population was used as the term $X(t)$. To obtain the instantaneous firing rate, we convolved the CA1 population spike train with an exponential kernel using the following equations:

$$\begin{cases} \frac{dX}{dt} & = -\frac{X}{\tau_{FR}} \\ X_{t+1} & = X_t + \frac{1}{N\,\tau_{FR}}, \forall\, t \in t^S \end{cases} \tag{3}$$

where t^S denotes the ordered set of the spike timings, and τ_{FR} determines the exponential decay of the kernel, which was set to 10 ms to compute population activity (for single neurons, a typical value is 100 ms) (*Gerstner et al., 2014*).

Hereafter, we call the term $Z(\theta)$ the phase response function, to distinguish it from the PRC obtained from experimental data or simulations (see section below *"Data Analysis"*, *"Phase Response Curve"*). Briefly, the PRC of an oscillatory system indicates the phase delay or advancement that follows a single pulse, as a function of the phase at which this input is delivered. The phase response function $Z(\theta)$ was chosen to mimic as well as possible experimental PRCs reported in the literature (*Lengyel et al., 2005*; *Kwag and Paulsen, 2009*; *Akam et al., 2012*). These PRCs appear biphasic and show a phase advancement (respectively delay) for stimuli delivered in the ascending (respectively descending) slope of theta. To accurately model this behavior, we used the following equation for the phase response function, where θ_{peak} represents the phase at which the theta rhythm reaches its maximum and the parameter ϕ_{offset} controls the desired phase offset from the peak:

$$Z(\theta_i) = -\sin(\theta_i - (\theta_{peak} + \phi_{offset})) \tag{4}$$

An overview of the default parameters for the Kuramoto oscillators and the bidirectional connections between medial septum-hippocampal formation can be found in *Table 1*.

Hippocampal formation: Hodgkin-Huxley Neurons

The following sections describe in detail how individual neurons and synapses were modeled, and are adapted from the original work by Aussel and colleagues (*Aussel et al., 2018*). Neurons were

Table 1. Full list of the default parameter values for the Kuramoto oscillators.

Parameter	Value	
Number of oscillators (N)	250	
Center frequency (f_0)	6	Hz
Standard deviation (σ)	0.5	Hz
Synchronization ratio ($\frac{k}{N}$)	15	
Phase reset gain (G_{reset})	4	
Peak phase (θ_{peak})	0	rad
Firing rate time constant (τ_{FR})	10	ms

modeled as conductance-based single compartments, following the Hodgkin-Huxley formalism (*Hodgkin and Huxley, 1952*), in line with previous work (*Aussel et al., 2022*, *2018*). The temporal evolution of the membrane potential of each neuron is described by a differential equation whose general form reads:

$$C_m \frac{dV_m}{dt} = -I_L - \sum_{channel} I_{channel} - \sum_{j \in [E,I]} I_{syn_j} + I_\theta + I_{stim} + \eta \tag{5}$$

I_L denotes the leakage current. $I_{channel}$ are currents associated with specific ion channels, namely potassium (I_K), fast sodium (I_{Na}), and low-threshold calcium (I_{Ca}) currents, the CAN current (I_{CAN}) (*Giovannini et al., 2017*), and the M-type potassium channel current (I_M) responsible for spike adaptation (*Kosenko et al., 2012*; *Sun and Kapur, 2012*; *Kwag et al., 2014*). I_{syn} represents the currents originating from synaptic inputs to the cell and can be either depolarizing (negative sign) or hyperpolarizing (positive sign). η is a Gaussian random noise term accounting for other external inputs and synaptic fluctuations. Theta input from the medial septum is modeled as a depolarizing current and is denoted by I_θ, while electrical stimulation is denoted by I_{stim}. Excitatory neurons represent pyramidal cells in EC, CA3, and CA1, and granule cells in DG. They were modeled with I_{Na}, I_K, and I_{Ca}, and I_M currents, with the addition of I_{CAN} for pyramidal cells. Fast-spiking interneurons in all areas were modeled with I_{Na} and I_K currents. The complete description for all of the above ionic channels and their corresponding currents can be found in *Giovannini et al.* (*2017*). Leakage currents followed the following equation:

$$I_L = (g_L \times A) \times (V_m - E_L) \tag{6}$$

where g_L is the maximum leakage conductance, A is the area of the single compartment corresponding to the membrane of a neuron, and E_L is the reversal potential of the leakage channel. Channel currents I_K, I_M, I_{CAN} obey the following set of equations:

$$I_K = g_K \times A \times n^4 \times (V_m - E_K)$$
$$I_M = g_M \times A \times p \times (V_m - E_M) \tag{7}$$
$$I_{CAN} = g_{CAN} \times A \times m_{CAN}^2 \times (V_m - E_{CAN})$$

where g_K, g_M, g_{CAN} are the maximum conductances for the respective channel, and n, p, m_{CAN} are the respective gating variables defined by the following differential equations:

$$\frac{dn}{dt} = \frac{n_\infty - n}{\tau_n}$$
$$\frac{dp}{dt} = \frac{p_\infty - p}{\tau_p} \tag{8}$$
$$\frac{dm_{CAN}}{dt} = \frac{m_{CAN,\infty} - m_{CAN}}{\tau_{m_{CAN}}}$$

For the potassium and CAN currents, the steady-state values for their corresponding gating variables n_∞ and $m_{CAN,\infty}$ and their corresponding time constants τ_K and τ_{CAN} depend on the following functions of the transition rate constants:

$$n_\infty = \frac{\alpha_n}{\alpha_n + \beta_n} \qquad\qquad m_{CAN,\infty} = \frac{\alpha_{m_{CAN}}}{\alpha_{m_{CAN}} + \beta_{m_{CAN}}}$$

$$\tau_n = \frac{0.2}{\alpha_n + \beta_n} \qquad\qquad \tau_{m_{CAN}} = \frac{0.2}{\alpha_{m_{CAN}} + \beta_{m_{CAN}}} \tag{9}$$

The sodium current (I_{Na}) and calcium current (I_{Ca}) follow a similar set of equations:

$$I_{Na} = g_{Na} \times A \times m^3 \times h \times (V_m - E_{Na})$$

$$I_{Ca} = g_{Ca} \times A \times m^2 \times h \times (V_m - E_{Ca}) \tag{10}$$

with two gating variables m and h defined by the following differential equations:

$$\frac{dm}{dt} = \frac{n_\infty - n}{\tau_n} \qquad\qquad \frac{dn}{dt} = \frac{n_\infty - n}{\tau_n}$$

$$m_\infty = \frac{\alpha_m}{\alpha_m + \beta_m} \qquad\qquad h_\infty = \frac{\alpha_h}{\alpha_h + \beta_h}$$

$$\tau_m = \frac{0.2}{\alpha_m + \beta_m} \qquad\qquad \tau_h = \frac{0.2}{\alpha_h + \beta_h} \tag{11}$$

The gating variable of I_{CAN} depends on the calcium concentration within the neuron ($[Ca]_i^{2+}$), given by:

$$\frac{d[Ca]_i^{2+}}{dt} = \gamma(I_{Ca}) + \frac{([Ca]_\infty^{2+} - [Ca]_i^{2+}}{\tau_{[Ca]^{2+}}}$$

$$\gamma(I_{Ca}) = \frac{-k_u \times I_{Ca}}{2 \times F \times d \times A} \tag{12}$$

where $\tau_{[Ca]^{2+}} = 1s$ represents the rate of calcium removal from the cell, $[Ca]_\infty^{2+} = 0.24\ mol/L$ is the calcium concentration if the calcium channel remains open for a duration of $\Delta T \to \infty$, $k_u = 10^4$ is a unit conversion constant, F is the Faraday constant, and $d = 1\ \mu m$ is the depth at which the calcium is stored inside the cell.

Noise (η), accounting for random inputs to the network, was simulated as intracellular current acting on the membrane voltage and following the properties of a Gaussian random variable with a mean of $0\ \mu V$ and a standard deviation of $10\ \mu V$ ($\eta_E \sim N(0, 10)\ \mu V$) for excitatory neurons and a mean of $0\ \mu V$ and standard deviation of $1\ \mu V$ ($\eta_I \sim N(0, 1)\ \mu V$) for inhibitory neurons. The ratio of 1:10 between the noise terms was adapted from the original work and it accounts for the higher excitability of the inhibitory neurons as well as the E-I population size ratios.

The original model introduced some parameters representing the vigilance state (i.e. active wakefulness vs slow-wave sleep). However, the present model only focused on the state of active wakefulness, since this is when memory-related theta-nested gamma oscillations occur. For all simulations, the parameters were set so that the network operated in the wakefulness regime in a healthy hippocampus (*Aussel et al., 2018*). The full expressions for all the parameters defined above can be found in *Table 2* for pyramidal cells and in *Table 3* for interneurons.

Synaptic Models

Inter-neuronal interactions were modeled as instantaneous AMPA and GABA-A synapses using the synaptic currents I_{syn_E} and I_{syn_I} respectively. Synaptic currents were described by the following bi-exponential differential equations:

$$I_{syn_{I,E}} = g_{I,E}(V_m - E_{I,E})$$

$$\frac{dg_{I,E}}{dt} = \frac{1}{\tau_{g_{I,E}}}(-g_{I,E} + h_{I,E})$$

$$\frac{dh_{I,E}}{dt} = -h_{I,E}\frac{1}{\tau_{h_{I,E}}} \tag{13}$$

Table 2. Full list of parameter values and expressions for pyramidal neurons

Parameter	Expression
A	$29.10^3\,cm^2$
C_m	$1.\,\mu F/cm^2$
g_L	$0.01\,mS/cm^2$
E_L	$-70\,mV$
g_K	$5\,mS/cm^2$
E_K	$-100\,mV$
$\alpha_{n,K}$	$-0.032\,\dfrac{V_m+40\,mV}{-1+e^{-0.02\,(V_m+40\,mV)}}$
$\beta_{n,K}$	$0.5\,e^{\frac{-(V_m+45\,mV)}{40\,mV}}$
g_{Na}	$50\,mS/cm^2$
E_{Na}	$50\,mV$
$\alpha_{m,Na}$	$-0.32\,\dfrac{V_m+42\,mV}{e^{-\frac{V_m+42\,mV}{4\,mV}}-1}$
$\beta_{m,Na}$	$0.28\,\dfrac{V_m+15\,mV}{e^{-\frac{V_m+15\,mV}{5\,mV}}-1}$
$\alpha_{h,Na}$	$0.128\,e^{-\frac{V_m+38\,mV}{18\,mV}}$
$\beta_{h,Na}$	$\dfrac{4}{1+e^{-\frac{V_m+15\,mV}{5\,mV}}}$
g_M	$90\,\mu S/cm^2$
E_M	$-100\,mV$
p_∞	$\dfrac{1}{1+e^{-0.01\,(V_m+35\,mV)}}$
τ_p	$\dfrac{1}{3.3\,e^{\frac{(V_m+35\,mV)}{20\,mV}}+e^{-\frac{(V_m+35\,mV)}{20\,mV}}}$
g_{Ca}	$0.1\,mS/cm^2$
E_{Ca}	$120\,mV$
$\alpha_{m,Ca}$	$-0.055\,\dfrac{V_m+27\,mV}{e^{-\frac{V_m+27\,mV}{17\,mV}}-1}$
$\beta_{m,Ca}$	$-0.94e^{\frac{V_m+75\,mV}{17\,mV}}$
$\alpha_{h,Ca}$	$-0.000457e^{\frac{V_m+13\,mV}{50\,mV}}$
$\beta_{h,Ca}$	$\dfrac{0.0065}{e^{-\frac{V_m+15\,mV}{28\,mV}}}$
g_{CAN}	$25\,\mu S/cm^2$
E_{CAN}	$-20\,mV$
$\alpha_{m,CAN}$	$0.0002\,e^{1.4\frac{[Ca]_i^{2+}}{0.5\,mol/L}}$
$\beta_{m,CAN}$	$0.0002\,e^{1.4}$

where $E_{I,E}$ are the synaptic resting potentials, and $\tau_{g_{I,E}}$ and $\tau_{h_{I,E}}$ are the synaptic time constants of rise and decay for inhibitory and excitatory neurons respectively. The occurrence of a pre-synaptic spike leads to an increase of the values h_I or h_E in the post-synaptic neuron by a fixed amount, which depends on the type of synapse and the region (due to the presence of cholinergic effects described in the initial model). Specific values for the intra-area and inter-area synaptic connections are given in *Table 4* and *Table 5* respectively.

Hippocampal formation: neuron types and numbers

Each area of the network is comprised of two populations, one excitatory and one inhibitory. Excitatory cells in the DG represent granule cells and pyramidal neurons in all other areas. Interneurons represent basket cells across all areas. The ratio between pyramidal neurons and interneurons was directly adapted from *Aussel et al.* (*2018*). The ratio between pyramidal neurons and interneurons was kept as a ratio of 10:1 for all areas except the dentate gyrus, where the ratio was 100:1. The number of neurons per subfield of the hippocampal formation is summarized in *Table 6*.

A two-dimensional simplified image depicting a coronal slice of the hippocampal formation

Table 3. Full list of parameter values and expressions for interneurons

Parameter	Expression
A	$14.10^3 cm^2$
C_m	$1.\mu F/cm^2$
g_L	$0.1 mS/cm^2$
E_L	$-65 mV$
g_K	$9 mS/cm^2$
E_K	$-90 mV$
$\alpha_{n,K}$	$0.01\frac{V_m+34\,mV}{1-e^{-0.1(V_m+34\,mV)}}$
$\beta_{n,K}$	$0.125 e^{-\frac{V_m+44\,mV}{80\,mV}}$
g_{Na}	$35 mS cm^2$
E_{Na}	$55 mV$
$\alpha_{m,Na}$	$0.1\frac{V_m+35\,mV}{1-e^{-0.1(V_m+35\,mV)}}$
$\beta_{m,Na}$	$4e^{-\frac{V_m+60\,mV}{18\,mV}}$
$\alpha_{h,Na}$	$0.07 e^{-\frac{V_m+58\,mV}{20\,mV}}$
$\beta_{h,Na}$	$\frac{1}{1+e^{-0.1(V_m+28\,mV)}}$

Table 4. A pre-synaptic spike causes an increase in the conductances h_e and h_i in the post-synaptic neuron. The values for the intra-area connections are given here. Empty cells indicate no connection between the populations.

	E → E	E → I	I → E	I → I
EC		20 pS	600 pS	
DG		180 pS	1800 pS	
CA3	20 pS	20 pS	600 pS	
CA1		60 pS	1800 pS	1800 pS

(**Aussel et al., 2018**) was used as a basis for a two-dimensional manifold that was uniformly populated by neurons following a density-driven approach (**Rougier, 2018**). Pyramidal neurons were uniformly distributed within the stratum pyramidale (or within the stratum granulosum for the dentate gyrus) and interneurons were uniformly distributed within the stratum oriens. Initial neuron positions were drawn from a blue noise distribution and a Voronoi diagram was computed. To adjust the positions of the neurons over a centroidal Voronoi diagram, the Lloyd relaxation algorithm was applied for 1000 iterations. Transitioning from a two-dimensional manifold to a 3D reconstruction of the hippocampal formation was achieved through the addition of the third coordinate with values uniformly distributed between 0 and 15 mm.

Hippocampal formation: inputs and connectivity

The hippocampal formation receives most of its external inputs from the EC. Activity from the EC is projected to all hippocampal subfields, starting with the DG. DG granule cells project onto the CA3 pyramidal cells via mossy cell fibers. CA3 projects in turn to CA1 via Schaffer collaterals. These connections form the tri-synaptic pathway. Direct connections from the EC towards the CA3 and CA1 subfields through the monosynaptic pathway are also considered. Pyramidal neurons from CA1 project to pyramidal neurons and interneurons in the EC, closing the hippocampal-entorhinal loop. CA1 pyramidal neurons also project to the medial septum through the fornix. An overview of the connections between areas is presented in panel B of *Figure 1*.

In the current model, EC pyramidal neurons and interneurons receive oscillatory theta input from the medial septum in the form of an excitatory intracellular current as described in *Equation 2* and *Equation 5*. Projections from CA1 towards the medial septum were modeled as a signal

Table 5. A pre-synaptic spike causes an increase in the conductances h_e and h_i in the post-synaptic neuron. The values for the inter-area connections are given here. Empty cells indicate no connection between the areas. Recurrent projections are not allowed and are marked with dashes.

Source \ Target	EC	DG	CA3	CA1
EC	-	20 pS	20 pS	20 pS
DG		-	180 pS	
CA3			-	20 pS
CA1	60 pS			-

Table 6. Number of neurons per subfield of the hippocampal formation, divided by neuron type.

Area	N_{Exc}	N_{Inh}
EC	10,000	1,000
DG	10,000	100
CA3	10,000	100
CA1	10,000	1,000

representing the collective firing rate of the CA1 pyramidal neurons. All connections towards and from the medial septum are summarized in panels A and B of *Figure 1*.

Synaptic connectivity between neurons within a region is characterized using a probability p and is distance-based, following a Gaussian-like distribution (*Equation 14*) with a width σ of $2500 \mu m$ (excitatory synapses) and $350 \mu m$ (inhibitory synapses). The value of A_{intra} defines the maximum probability of connection between two neurons separated by an infinitesimal distance (i.e. for $D \to 0$). The maximum values for A_{intra} are given in *Table 7*.

$$p = A_{intra} \ \exp(-\frac{D^2}{2\sigma^2}) \tag{14}$$

Inter-area connectivity followed a similar Gaussian-like distribution (*Equation 15*), however, only excitatory projections were considered and the distance D_z was computed only across the z-coordinate. Pyramidal neurons from the source area projected to pyramidal neurons and interneurons in the target area, with connectivity probabilities drawn from the said distribution with a width σ of $1000 \ \mu m$.

$$p = min(1, A_{inter} \exp(-\frac{D_z^2}{2\sigma^2})) \tag{15}$$

Table 7. Maximum probability of connection (A_{intra}, *Equation 14*) between neurons within each region.

	Py-Py	Py-Inh	Inh-Py	Inh-Inh
EC	0	0.37	0.54	0
DG	0	0.06	0.14	0
CA3	0.56	0.75	0.75	0
CA1	0	0.28	0.3	0.7

Tuning input gain and connection strengths: targeted firing rates

We adjusted the input gain (G_θ) and inter-area connection strengths (A_{inter}) to target an overall oscillatory rhythm $f_{osc,targ}$ at the driving frequency of the input, i.e. 6 Hz, and a mean firing rate of each excitatory population $f_{exc,targ}$ also at 6 Hz (meaning that each excitatory cell should spike on

average once per theta cycle). Because of the ratio between excitatory and inhibitory neurons in the model, this resulted in firing rates of about 60 Hz in inhibitory neurons. These targeted values were inspired by literature in behaving rodents showing that hippocampal pyramidal neurons typically fire at rates below 10 Hz, usually between 1 and 2 Hz, and that interneurons fire at rates between 20 and 80 Hz (*Hirase et al., 2001*). In practice, the obtained firing rates were constrained by the simplifications made in the model.

The mean population firing rate f_{exc} was computed by counting the number of spikes in the last second of a 3-s simulation run, to avoid edge effects due to the non-physiological initial conditions, and by averaging this number over time and across neurons. The oscillatory frequency f_{osc} was computed as the mean of the inverse of the timing between all pairs of consecutive peaks in the theta rhythm. Finally, the metric used to adjust parameters (input strength and connection strengths) was calculated as the Euclidean distance between the targeted and obtained firing rate and oscillatory rate:

$$J = \sqrt{(f_{exc} - f_{exc,targ})^2 + (f_{osc} - f_{osc,targ})^2} \tag{16}$$

Tuning input and connection strengths: detailed procedure

The input gain (G_θ, *Equation 2*) and inter-area connection strengths (A_{inter}, *Equation 15*) were sequentially adjusted using the following heuristics. All simulations were performed for a duration of 3 s, and the first 2 s were excluded from the analysis to avoid edge effects. The network was initialized with membrane voltages uniformly distributed in the range $[-70, -60]$ mV. Our initial point was a fully uncoupled model in which all the connection strengths were set to 0. The tuning procedure was performed in the absence of noise.

1. First, we adjusted the amplitude of the theta input to a value that would produce a mean firing rate of 6 Hz in the excitatory population of the EC. To this end, the input strength was progressively increased from 0 to 0.5 nA in steps of 0.01 nA, and the value that maximized our metric was selected.
2. We then adjusted similarly the connection strength from the EC to the DG, increasing it from 0 to 20 in steps of 0.1.
3. We repeated the procedure to jointly tune the connection strengths from EC and DG to CA3, assuming an equal contribution from these two areas. These two connection strengths were set to the same value and were increased from 0 to 5 in steps of 0.1.
4. We followed the same procedure to jointly tune the connection strengths from CA3 and EC to CA1, assuming again an equal contribution from these two areas.
5. The last connection from CA1 to EC was more difficult to tune because it closes the feedback loop and induces complex dynamics when too strong. To tune it, we temporarily uncoupled CA1 from all other structures and temporarily added a fixed sinusoidal input at the target frequency of 6 Hz to both excitatory and inhibitory CA1 neurons. We first adjusted this input amplitude to generate a mean firing rate of about 6 Hz in CA1 excitatory neurons. Next, the connection strength from CA1 to EC was adjusted to also achieve a mean firing rate of 6 Hz in EC excitatory neurons.
6. Because EC receives both an external input and feedback from CA1, we assumed that both these contributions should be decreased for the system to behave in a physiological range in the presence of the feedback loop. We, therefore, decided to divide by two the connection strength from CA1 to EC obtained in step 5. All other connection strengths were then set again to the values tuned in the previous steps, and the temporary external input to CA1 was removed. The external theta input to the EC was reinstated and tuned again as in step 1.

Following the above procedure, the inter-area synaptic connectivity parameters were set for all subsequent simulations. The values are summarized in *Table 8*. Empty cells denote no effective connectivity.

Table 8. Inter-area connection strengths (A_{inter}). The source is always the excitatory population of the subfield. The same values are used when targeting excitatory and inhibitory populations. Empty cells indicate no connections. EC: Entorhinal Cortex, DG: Dentate Gyrus.

Source \ Target	EC	DG	CA3	CA1
EC	-	13.0	0.14	1.1
DG		-	0.14	
CA3			-	1.1
CA1	0.2			-

Numerical Implementation

The model was implemented with the Brian2 libraries for Python (*Stimberg et al., 2019*) (see Data availability section for access to the code). Simulations were performed using a timestep of 0.1 ms and a total simulation duration ranging between 3-10 seconds (depending on the experiment, see results section). The average time for simulating 1 second for the complete model locally was 10 minutes.

Data Analysis

During each simulation, we monitored and exported the following data for subsequent analysis: (i) spike timings per neuron, (ii) time series of ionic currents, (iii) input theta rhythm and phase, and (iv) time series of electrical stimulation. Theta phase was wrapped between $[-\pi, \pi]$ with a phase of 0 radians corresponding to the peak of theta rhythm. All PAC analyses were performed using the *TensorPAC* toolbox for Python (*Combrisson et al., 2020*).

Firing rates

To obtain instantaneous firing rates, the corresponding spike trains were binned in 5-ms rectangular windows with a 90% overlap (i.e. consecutive windows were spaced by 0.5 ms). The number of spikes within each bin was normalized by the bin size and the number of neurons in the group, yielding instantaneous population firing rates. Where reported (i.e. values μ_I and μ_E in *Figure 2*A), the mean population firing rates within a given time window (typically lasting several seconds) were computed by binning all spikes in that time window and normalizing by the window width and the number of neurons in the population.

Spectral analyses

Power spectral density (PSD) estimates were calculated based on Welch's method. The periodogram was computed using 1-s windows with a 90% overlap, yielding a frequency resolution of 1 Hz. The average spectral power within a specific frequency band was calculated using Simpson's rule within the desired frequency band $[f_l, f_h]$. Spectrograms were computed using the short-time Fourier transform with a sliding Hann window of 100-ms width and 99% overlap, yielding a frequency resolution of 10 Hz.

Modulation Index

The MI (*Tort et al., 2008*) was used to estimate the degree of PAC between theta and gamma oscillations in the model. To compute the MI, the normalized firing rate traces were band-pass filtered in the frequency ranges of interest: 3-9 Hz for theta (referred to as the "phase signal") and 40-80 Hz for gamma (referred to as the "amplitude signal"). Then, the phase and amplitude time series were extracted from the filtered signals using the Hilbert transform. A histogram of the mean amplitude of gamma over the phase of theta was then extracted, using phase bins of 5 degrees. The MI was finally calculated as the Kullback-Leibler divergence between the mean amplitude distribution and the uniform distribution. A higher MI indicates stronger PAC. For a schematic representation

regarding the computation of the MI, refer to Figure 1 in *Tort et al.* (*2010*). These computations were performed using the provided *PreferredPhase* and *Pac* methods from the *TensorPAC* library for Python (*Combrisson et al., 2020*).

Comodulograms

Comodulograms represent the amount of PAC between two ranges of frequencies, used to extract respectively a phase and an amplitude signal. Specifically, we computed the MI between 80% overlapping 1-Hz frequency bands used to compute the phase of the signal (in the theta range), and 90% overlapping 10-Hz frequency bands used to compute the amplitude of the signal (in the gamma range). The resulting distribution of MI values was subsequently plotted as heat maps. An inherent problem with the computation of the MI using simulated data was the lack of frequency components in some frequency bands. Filtering the data within 1-Hz frequency bands thus created a flat signal, resulting in high values of the MI despite the absence of modulation. To overcome this limitation, we added uniform noise to the firing rate signals prior to computing the MI, with an amplitude of approximately 20% of the maximum instantaneous firing rates. The results with and without added noise are presented in *Figure S3*.

Phase dependency of PAC

For a given theta frequency, PAC depends on the phase of the underlying theta oscillation. To identify this relationship and the theta phase that maximizes coupling, we used the *PreferredPhase* function of the *TensorPAC* toolbox. More precisely, we applied a Hilbert transform to extract the phase of the theta signal (firing rate band-pass filtered between 3 and 9 Hz) and the amplitude of the gamma signal band-pass filtered within narrow (10-Hz wide) frequency ranges between 20 and 100 Hz with a 90% overlap. For each narrow gamma range, we binned the amplitude with respect to the phase in a similar way as in the calculation of the MI. We obtained a vector of the binned high-frequency amplitudes with respect to the phase of the low-frequency phase, represented as a polar plot as in *Figure 2*C.

Phase Response Curves

The PRC of an oscillatory system indicates the phase delay or advancement that follows a single pulse, as a function of the phase at which this input is delivered. To characterize the PRC of our computational model, we applied a single stimulation pulse to CA1 across different phases of the theta rhythm and calculated the resulting change in the theta phase. We split a single theta cycle into intervals of width $\pi/8$ radians and applied a single stimulation pulse of a given amplitude. For each case, we ran two simulations: one with a stimulation pulse and one without. Finally, we compared the theta phase 2.5 ms post-stimulation and at the same time but in the absence of stimulation. The resulting values for the phase difference $\Delta\phi$ were plotted against the stimulation phase and are presented in *Figure 2*D for varying stimulation amplitudes.

Acknowledgments

Simulations presented in this paper were carried out using the PlaFRIM experimental testbed, supported by Inria, CNRS (LABRI and IMB), Université de Bordeaux, Bordeaux INP and Conseil Régional d'Aquitaine (see https://www.plafrim.fr).

Additional information

Funding

This work was supported by the Bordeaux Neurocampus junior chair program co-funded by the Région Nouvelle-Aquitaine and the University of Bordeaux Initiative of Excellence (IdEx) (F.B.W., Neurocampus junior chair), and the European Union's Horizon Europe research and innovation program under European Research Council (ERC) Starting Grant agreement N° 101040391 (F.B.W., MEMOPROSTHETICS).

Author contributions

N.V., Conceptualization, Methodology, Software, Formal analysis, Investigation, Visualization (lead), Writing – original draft, Writing – review and editing; A.A., Methodology, Formal analysis, Investigation, Software, Visualization (supporting), Writing – original draft, Writing – review and editing, Supervision; N.P.R., Conceptualization, Methodology, Formal analysis, Investigation, Visualization (supporting), Writing – original draft, Writing – review and editing, Supervision; F.B.W., Conceptualization, Methodology, Formal analysis, Investigation, Visualization (supporting), Writing – original draft, Writing – review and editing, Supervision, Funding acquisition.

Author ORCIDs

Nikolaos Vardalakis https://orcid.org/0000-0002-5436-4091 https://orcid.org/0000-0002-5436-4091

Amélie Aussel https://orcid.org/0000-0003-0498-2905 https://orcid.org/0000-0003-0498-2905

Nicolas P. Rougier https://orcid.org/0000-0002-6972-589X https://orcid.org/0000-0002-6972-589X

Fabien B. Wagner https://orcid.org/0000-0002-9582-6109 https://orcid.org/0000-0002-9582-6109

Conflicts of interest

The authors declare no conflicts of interest.

Data availability

All model files and analysis scripts are available at the following repositories under the GNU General Public License v3.0 license:

ZENODO https://www.doi.org/10.5281/zenodo.7702041

GITHUB https://github.com/NikVard/memstim-hh

References

Abbaspoor S, Hussin AT, Hoffman KL. Theta- and gamma-band oscillatory uncoupling in the macaque hippocampus; 2022.

Akam T, Oren I, Mantoan L, Ferenczi E, Kullmann DM. Oscillatory dynamics in the hippocampus support dentate gyrus–CA3 coupling. Nat Neurosci. 2012 May; 15(5):763–768. http://www.nature.com/articles/nn.3081, doi: 10.1038/nn.3081.

de Almeida L, Idiart M, Lisman JE. Memory retrieval time and memory capacity of the CA3 network: role of gamma frequency oscillations. Learn Mem. 2007 Nov; 14(11):795–806. doi: 10.1101/lm.730207.

Ashida G, Nogueira W. Spike-Conducting Integrate-and-Fire Model. eNeuro. 2018 Jul; 5(4):ENEURO.0112–18.2018. https://www.eneuro.org/lookup/doi/10.1523/ENEURO.0112-18.2018, doi: 10.1523/ENEURO.0112-18.2018.

Asllani M, Expert P, Carletti T. A minimally invasive neurostimulation method for controlling abnormal synchronisation in the neuronal activity. PLOS Computational Biology. 2018 Jul; 14(7):e1006296. https://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1006296, doi: 10.1371/journal.pcbi.1006296, publisher: Public Library of Science.

Aussel A, Buhry L, Tyvaert L, Ranta R. A detailed anatomical and mathematical model of the hippocampal formation for the generation of sharp-wave ripples and theta-nested gamma oscillations. J Comput Neurosci. 2018 Dec; 45(3):207–221. http://link.springer.com/10.1007/s10827-018-0704-x, doi: 10.1007/s10827-018-0704-x.

Aussel A, Ranta R, Aron O, Colnat-Coulbois S, Maillard L, Buhry L. Cell to network computational model of the epileptic human hippocampus suggests specific roles of network and channel dysfunctions in the ictal and interictal oscillations. J Comput Neurosci. 2022 Nov; 50(4):519–535. doi: 10.1007/s10827-022-00829-5.

Axmacher N, Henseler MM, Jensen O, Weinreich I, Elger CE, Fell J. Cross-frequency coupling supports multi-item working memory in the human hippocampus. PNAS. 2010 Feb; 107(7):3228–3233. https://www.pnas.org/content/107/7/3228, doi: 10.1073/pnas.0911531107, publisher: National Academy of Sciences Section: Biological Sciences.

Axmacher N, Mormann F, Fernández G, Elger CE, Fell J. Memory formation by neuronal synchronization. Brain Research Reviews. 2006 Aug; 52(1):170–182. https://linkinghub.elsevier.com/retrieve/pii/S0165017306000099, doi: 10.1016/j.brainresrev.2006.01.007.

Basu I, Crocker B, Farnes K, Robertson MM, Paulk AC, Vallejo DI, Dougherty DD, Cash SS, Eskandar EN, Kramer MM, Widge AS. A neural mass model to predict electrical stimulation evoked responses in human and non-human primate brain. J Neural Eng. 2018 Dec; 15(6):066012. https://iopscience.iop.org/article/10.1088/1741-2552/aae136, doi: 10.1088/1741-2552/aae136.

Bingham CS, Loizos K, Yu GJ, Gilbert A, Bouteiller JMC, Song D, Lazzi G, Berger TW. Model-Based Analysis of Electrode Placement and Pulse Amplitude for Hippocampal Stimulation. IEEE Trans Biomed Eng. 2018 Oct; 65(10):2278–2289. https://ieeexplore.ieee.org/document/8269392/, doi: 10.1109/TBME.2018.2791860.

Borges RR, Borges FS, Lameu EL, Batista AM, Iarosz KC, Caldas IL, Antonopoulos CG, Baptista MS. Spike timing-dependent plasticity induces non-trivial topology in the brain. Neural Networks. 2017 Apr; 88:58–64. https://linkinghub.elsevier.com/retrieve/pii/S0893608017300102, doi: 10.1016/j.neunet.2017.01.010.

Breakspear M, Heitmann S, Daffertshofer A. Generative Models of Cortical Oscillations: Neurobiological Implications of the Kuramoto Model. Front Hum Neurosci. 2010; 4. https://www.frontiersin.org/articles/10.3389/fnhum.2010.00190/full, doi: 10.3389/fnhum.2010.00190, publisher: Frontiers.

Buzsáki G, Buhl DL, Harris KD, Csicsvari J, Czéh B, Morozov A. Hippocampal network patterns of activity in the mouse. Neuroscience. 2003 Jan; 116(1):201–211. http://www.sciencedirect.com/science/article/pii/S0306452202006693, doi: 10.1016/S0306-4522(02)00669-3.

Buzsáki G. Theta Oscillations in the Hippocampus. Neuron. 2002 Jan; 33(3):325–340. https://www.sciencedirect.com/science/article/pii/S089662730200586X, doi: 10.1016/S0896-6273(02)00586-X.

Buño W, Garcia-Sanchez JL, Garcia-Austt E. Reset of hippocampal rhythmical activities by afferent stimulation. Brain Res Bull. 1978; 3(1):21–28. doi: 10.1016/0361-9230(78)90057-6.

Canolty RT, Knight RT. The functional role of cross-frequency coupling. Trends in Cognitive Sciences. 2010 Nov; 14(11):506–515. https://linkinghub.elsevier.com/retrieve/pii/S1364661310002068, doi: 10.1016/j.tics.2010.09.001.

Capogrosso M, Wenger N, Raspopovic S, Musienko P, Beauparlant J, Bassi Luciani L, Courtine G, Micera S. A Computational Model for Epidural Electrical Stimulation of Spinal Sensorimotor Circuits. J Neurosci. 2013 Dec; 33(49):19326–19340. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6618777/, doi: 10.1523/JNEUROSCI.1688-13.2013.

Colgin LL. Mechanisms and Functions of Theta Rhythms. Annual Review of Neuroscience. 2013; 36(1):295–312. https://doi.org/10.1146/annurev-neuro-062012-170330, doi: 10.1146/annurev-neuro-062012-170330.

Combrisson E, Nest T, Brovelli A, Ince RAA, Soto JLP, Guillot A, Jerbi K. Tensorpac: An open-source Python toolbox for tensor-based phase-amplitude coupling measurement in electrophysiological brain signals. PLOS Computational Biology. 2020 Oct; 16(10):e1008302. https://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1008302, doi: 10.1371/journal.pcbi.1008302, publisher: Public Library of Science.

Cutsuridis V, Cobb S, Graham BP. Encoding and retrieval in a model of the hippocampal CA1 microcircuit. Hippocampus. 2010; 20(3):423–446. http://onlinelibrary.wiley.com/doi/abs/10.1002/hipo.20661, doi: 10.1002/hipo.20661, _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/hipo.20661.

Dragoi G, Buzsáki G. Temporal encoding of place sequences by hippocampal cell assemblies. Neuron. 2006 Apr; 50(1):145–157. doi: 10.1016/j.neuron.2006.02.023.

Ebert M, Hauptmann C, Tass PA. Coordinated reset stimulation in a large-scale model of the STN-GPe circuit. Frontiers in Computational Neuroscience. 2014; 8:154. https://www.frontiersin.org/article/10.3389/fncom.2014.00154, doi: 10.3389/fncom.2014.00154.

Fell J, Axmacher N. The role of phase synchronization in memory processes. Nat Rev Neurosci. 2011 Feb; 12(2):105–118. http://www.nature.com/articles/nrn2979, doi: 10.1038/nrn2979.

Gerstner W, Kistler WM, Naud R, Paninski L. Neuronal Dynamics: From Single Neurons to Networks and Models of Cognition. USA: Cambridge University Press; 2014.

Giovannini F, Knauer B, Yoshida M, Buhry L. The CAN-In network: A biologically inspired model for self-sustained theta oscillations and memory maintenance in the hippocampus. Hippocampus. 2017 Apr; 27(4):450–463. doi: 10.1002/hipo.22704.

Goyal A, Miller J, Watrous AJ, Lee SA, Coffey T, Sperling MR, Sharan A, Worrell G, Berry B, Lega B, Jobst BC, Davis KA, Inman C, Sheth SA, Wanda PA, Ezzyat Y, Das SR, Stein J, Gorniak R, Jacobs J. Electrical Stimulation in Hippocampus and Entorhinal Cortex Impairs Spatial and Temporal Memory. J Neurosci. 2018 May; 38(19):4471–4481. doi: 10.1523/JNEUROSCI.3049-17.2018.

Gupta A, Vardalakis N, Wagner FB. Neuroprosthetics: from sensorimotor to cognitive disorders. Commun Biol. 2023 Jan; 6(1):1–17. https://www.nature.com/articles/s42003-022-04390-w, doi: 10.1038/s42003-022-04390-w, number: 1 Publisher: Nature Publishing Group.

Hajós M, Hoffmann WE, Orbán G, Kiss T, Érdi P. Modulation of septo-hippocampal θ activity by GABAA receptors: an experimental and computational approach. Neuroscience. 2004 Jan; 126(3):599–610. https://linkinghub.elsevier.com/retrieve/pii/S0306452204002301, doi: 10.1016/j.neuroscience.2004.03.043.

Hampel H, Mesulam MM, Cuello AC, Farlow MR, Giacobini E, Grossberg GT, Khachaturian AS, Vergallo A, Cavedo E, Snyder PJ, Khachaturian ZS. The cholinergic system in the pathophysiology and treatment of Alzheimer's disease. Brain. 2018 Jul; 141(7):1917–1933. https://doi.org/10.1093/brain/awy132, doi: 10.1093/brain/awy132.

Hangya B, Borhegyi Z, Szilágyi N, Freund TF, Varga V. GABAergic neurons of the medial septum lead the hippocampal network during theta activity. J Neurosci. 2009 Jun; 29(25):8094–8102. doi: 10.1523/JNEUROSCI.5665-08.2009.

Hasselmo ME, Bodelón C, Wyble BP. A Proposed Function for Hippocampal Theta Rhythm: Separate Phases of Encoding and Retrieval Enhance Reversal of Prior Learning. Neural Computation. 2002 Apr; 14(4):793–817. https://www.mitpressjournals.org/doi/abs/10.1162/089976602317318965, doi: 10.1162/089976602317318965.

Hendrickson PJ, Yu GJ, Song D, Berger TW. A Million-Plus Neuron Model of the Hippocampal Dentate Gyrus: Critical Role for Topography in Determining Spatiotemporal Network Dynamics. IEEE Transactions on Biomedical Engineering. 2016 Jan; 63(1):199–209. doi: 10.1109/TBME.2015.2445771, conference Name: IEEE Transactions on Biomedical Engineering.

Herman PA, Lundqvist M, Lansner A. Nested theta to gamma oscillations and precise spatiotemporal firing during memory retrieval in a simulated attractor network. Brain Res. 2013 Nov; 1536:68–87. doi: 10.1016/j.brainres.2013.08.002.

Hirase H, Leinekugel X, Czurkó A, Csicsvari J, Buzsáki G. Firing rates of hippocampal neurons are preserved during subsequent sleep episodes and modified by novel awake experience. Proceedings of the National Academy of Sciences. 2001 Jul; 98(16):9386–9390. https://www.pnas.org/doi/10.1073/pnas.161274398, doi: 10.1073/pnas.161274398, publisher: Proceedings of the National Academy of Sciences.

Hodgkin AL, Huxley AF. A quantitative description of membrane current and its application to conduction and excitation in nerve. J Physiol. 1952 Aug; 117(4):500–544. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1392413/.

Hülsemann MJ, Naumann E, Rasch B. Quantification of Phase-Amplitude Coupling in Neuronal Oscillations: Comparison of Phase-Locking Value, Mean Vector Length, Modulation Index, and Generalized-Linear-Modeling-Cross-Frequency-Coupling. Frontiers in Neuroscience. 2019; 13. https://www.frontiersin.org/article/10.3389/fnins.2019.00573.

Jackson J, Dickson CT, Bland BH. Median Raphe Stimulation Disrupts Hippocampal Theta Via Rapid Inhibition and State-Dependent Phase Reset of Theta-Related Neural Circuitry. Journal of Neurophysiology. 2008 Jun; 99(6):3009–3026. https://www.physiology.org/doi/10.1152/jn.00065.2008, doi: 10.1152/jn.00065.2008.

Jacobs J, Miller J, Lee SA, Coffey T, Watrous AJ, Sperling MR, Sharan A, Worrell G, Berry B, Lega B, Jobst BC, Davis K, Gross RE, Sheth SA, Ezzyat Y, Das SR, Stein J, Gorniak R, Kahana MJ, Rizzuto DS. Direct Electrical Stimulation of the Human Entorhinal Region and Hippocampus Impairs Memory. Neuron. 2016 Dec; 92(5):983–990. https://www.sciencedirect.com/science/article/pii/S0896627316308364, doi: 10.1016/j.neuron.2016.10.062.

Jensen O, Colgin LL. Cross-frequency coupling between neuronal oscillations. Trends in Cognitive Sciences. 2007 Jul; 11(7):267–269. https://www.sciencedirect.com/science/article/pii/S1364661307001271, doi: 10.1016/j.tics.2007.05.003.

Joucla S, Yvert B. Modeling extracellular electrical neural stimulation: From basic understanding to MEA-based applications. Journal of Physiology-Paris. 2012 May; 106(3-4):146–158. https://linkinghub.elsevier.com/retrieve/pii/S0928425711000416, doi: 10.1016/j.jphysparis.2011.10.003.

Jun S, Kim JS, Chung CK. Direct Stimulation of Human Hippocampus During Verbal Associative Encoding Enhances Subsequent Memory Recollection. Front Hum Neurosci. 2019 Feb; 13:23. https://www.frontiersin.org/article/10.3389/fnhum.2019.00023/full, doi: 10.3389/fnhum.2019.00023.

Jutras MJ, Fries P, Buffalo EA. Oscillatory activity in the monkey hippocampus during visual exploration and memory formation. Proceedings of the National Academy of Sciences of the United States of America. 2013; 110(32):13144–13149. doi: 10.1073/pnas.1302351110.

Kipping D, Nogueira W. A Computational Model of a Single Auditory Nerve Fiber for Electric-Acoustic Stimulation. J Assoc Res Otolaryngol. 2022 Dec; 23(6):835–858. doi: 10.1007/s10162-022-00870-2.

Kocsis B, Bragin A, Buzsáki G. Interdependence of Multiple Theta Generators in the Hippocampus: a Partial Coherence Analysis. J Neurosci. 1999 Jul; 19(14):6200–6212. https://www.jneurosci.org/content/19/14/6200, doi: 10.1523/JNEUROSCI.19-14-06200.1999, publisher: Society for Neuroscience Section: ARTICLE.

Kosenko A, Kang S, Smith IM, Greene DL, Langeberg LK, Scott JD, Hoshi N. Coordinated signal integration at the M-type potassium channel upon muscarinic stimulation. EMBO J. 2012 May; 31(14):3147–3156. doi: 10.1038/emboj.2012.156.

Kota S, Rugg MD, Lega BC. Hippocampal Theta Oscillations Support Successful Associative Memory Formation. J Neurosci. 2020 Dec; 40(49):9507–9518. http://www.jneurosci.org/lookup/doi/10.1523/JNEUROSCI.0767-20.2020, doi: 10.1523/JNEUROSCI.0767-20.2020.

Kuramoto Y. Chemical Oscillations, Waves, and Turbulence, vol. 19 of Springer Series in Synergetics. Haken H, editor, Berlin, Heidelberg: Springer Berlin Heidelberg; 1984. http://link.springer.com/10.1007/978-3-642-69689-3, doi: 10.1007/978-3-642-69689-3.

Kwag J, Jang HJ, Kim M, Lee S. M-type potassium conductance controls the emergence of neural phase codes: a combined experimental and neuron modelling study. J R Soc Interface. 2014 Oct; 11(99):20140604. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4233740/, doi: 10.1098/rsif.2014.0604.

Kwag J, Paulsen O. The timing of external input controls the sign of plasticity at local synapses. Nat Neurosci. 2009 Oct; 12(10):1219–1221. http://www.nature.com/articles/nn.2388, doi: 10.1038/nn.2388.

Lacruz ME, Valentín A, Seoane JJG, Morris RG, Selway RP, Alarcón G. Single pulse electrical stimulation of the hippocampus is sufficient to impair human episodic memory. Neuroscience. 2010 Oct; 170(2):623–632. doi: 10.1016/j.neuroscience.2010.06.042.

Larson J, Munkácsy E. Theta-burst LTP. Brain Res. 2015 Sep; 1621:38–50. doi: 10.1016/j.brainres.2014.10.034.

Lega B, Burke J, Jacobs J, Kahana MJ. Slow-Theta-to-Gamma Phase-Amplitude Coupling in Human Hippocampus Supports the Formation of New Episodic Memories. Cereb Cortex. 2016 Jan; 26(1):268–278. doi: 10.1093/cercor/bhu232.

Lega BC, Jacobs J, Kahana M. Human hippocampal theta oscillations and the formation of episodic memories. Hippocampus. 2012 Apr; 22(4):748–761. https://onlinelibrary.wiley.com/doi/10.1002/hipo.20937, doi: 10.1002/hipo.20937.

Lengyel M, Kwag J, Paulsen O, Dayan P. Matching storage and recall: hippocampal spike timing–dependent plasticity and phase response curves. Nat Neurosci. 2005 Dec; 8(12):1677–1683. http://www.nature.com/articles/nn1561, doi: 10.1038/nn1561.

Lin JJ, Rugg M, Das S, Stein J, Rizzuto D, Kahana M, Lega B. Theta band power increases in the posterior hippocampus predict successful episodic memory encoding in humans. Hippocampus. 2017 Oct; 27(10):1040–1053. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6517838/, doi: 10.1002/hipo.22751.

Lisman JE, Talamini LM, Raffone A. Recall of memory sequences by interaction of the dentate and CA3: A revised model of the phase precession. Neural Networks. 2005 Nov; 18(9):1191–1201. https://linkinghub.elsevier.com/retrieve/pii/S0893608005002017, doi: 10.1016/j.neunet.2005.08.008.

Lisman J, Jensen O. The Theta-Gamma Neural Code. Neuron. 2013 Mar; 77(6):1002–1016. https://linkinghub.elsevier.com/retrieve/pii/S0896627313002316, doi: 10.1016/j.neuron.2013.03.007.

Lozano AM, Fosdick L, Chakravarty MM, Leoutsakos JM, Munro C, Oh E, Drake KE, Lyman CH, Rosenberg PB, Anderson WS, Tang-Wai DF, Pendergrass JC, Salloway S, Asaad WF, Ponce FA, Burke A, Sabbagh M, Wolk DA, Baltuch G, Okun MS, et al. A Phase II Study of Fornix Deep Brain Stimulation in Mild Alzheimer's Disease. Journal of Alzheimer's Disease. 2016 Jan; 54(2):777–787. https://content.iospress.com/articles/journal-of-alzheimers-disease/jad160017, doi: 10.3233/JAD-160017, publisher: IOS Press.

Lundqvist M, Rehn M, Djurfeldt M, Lansner A. Attractor dynamics in a modular network model of neocortex. Network: Computation in Neural Systems. 2006 Jan; 17(3):253–276. https://www.tandfonline.com/doi/full/10.1080/09548980600774619, doi: 10.1080/09548980600774619.

Lurie SM, Kragel JE, Schuele SU, Voss JL. Human hippocampal responses to network intracranial stimulation vary with theta phase. eLife. 2022 Dec; 11:e78395. https://elifesciences.org/articles/78395, doi: 10.7554/eLife.78395.

Malkov AE, Shevkova L, Latyshkova A, Kitchigina V. Theta and gamma hippocampal-neocortical oscillations during the episodic-like memory test: impairment in epileptogenic rats. Neuroscience; 2021.

Manseau F, Goutagny R, Danik M, Williams S. The Hippocamposeptal Pathway Generates Rhythmic Firing of GABAergic Neurons in the Medial Septum and Diagonal Bands: An Investigation Using a Complete Septohippocampal Preparation In Vitro. J Neurosci. 2008 Apr; 28(15):4096–4107. https://www.jneurosci.org/content/28/15/4096, doi: 10.1523/JNEUROSCI.0247-08.2008, publisher: Society for Neuroscience Section: Articles.

McCartney H, Johnson AD, Weil ZM, Givens B. Theta reset produces optimal conditions for long-term potentiation. Hippocampus. 2004; 14(6):684–687. http://onlinelibrary.wiley.com/doi/abs/10.1002/hipo.20019, doi: 10.1002/hipo.20019, _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/hipo.20019.

McIntyre CC, Grill WM, Sherman DL, Thakor NV. Cellular effects of deep brain stimulation: model-based analysis of activation and inhibition. J Neurophysiol. 2004 Apr; 91(4):1457–1469. doi: 10.1152/jn.00989.2003.

Mina F, Benquet P, Pasnicu A, Biraben A, Wendling F. Modulation of epileptic activity by deep brain stimulation: a model-based study of frequency-dependent effects. Front Comput Neurosci. 2013; 7. https://www.frontiersin.org/articles/10.3389/fncom.2013.00094/full, doi: 10.3389/fncom.2013.00094, publisher: Frontiers.

Mizuseki K, Sirota A, Pastalkova E, Buzsáki G. Theta Oscillations Provide Temporal Windows for Local Circuit Computation in the Entorhinal-Hippocampal Loop. Neuron. 2009 Oct; 64(2):267–280. https://www.sciencedirect.com/science/article/pii/S0896627309006734, doi: 10.1016/j.neuron.2009.08.037.

Mormann F, Fell J, Axmacher N, Weber B, Lehnertz K, Elger CE, Fernández G. Phase/amplitude reset and theta-gamma interaction in the human medial temporal lobe during a continuous word recognition memory task. Hippocampus. 2005; 15(7):890–900. https://onlinelibrary.wiley.com/doi/10.1002/hipo.20117, doi: 10.1002/hipo.20117.

Mysin IE, Kitchigina VF, Kazanovich YB. Phase relations of theta oscillations in a computer model of the hippocampal CA1 field: Key role of Schaffer collaterals. Neural Networks. 2019 Aug; 116:119–138. https://linkinghub.elsevier.com/retrieve/pii/S0893608019301029, doi: 10.1016/j.neunet.2019.04.004.

Müller C, Remy S. Septo–hippocampal interaction. Cell Tissue Res. 2018 Sep; 373(3):565–575. https://doi.org/10.1007/s00441-017-2745-2, doi: 10.1007/s00441-017-2745-2.

Nelson AR, Kolasa K, McMahon LL. Noradrenergic Sympathetic Sprouting and Cholinergic Reinnervation Maintains Non-Amyloidogenic Processing of AβPP. Journal of Alzheimer's Disease. 2014 Jan; 38(4):867–879. https://content.iospress.com/articles/journal-of-alzheimers-disease/jad130608, doi: 10.3233/JAD-130608, publisher: IOS Press.

Nuñez A, Buño W. The Theta Rhythm of the Hippocampus: From Neuronal and Circuit Mechanisms to Behavior. Front Cell Neurosci. 2021; 15. https://www.frontiersin.org/articles/10.3389/fncel.2021.649262/full, doi: 10.3389/fncel.2021.649262, publisher: Frontiers.

Onslow ACE, Jones MW, Bogacz R. A Canonical Circuit for Generating Phase-Amplitude Coupling. PLoS ONE. 2014 Aug; 9(8):e102591. https://dx.plos.org/10.1371/journal.pone.0102591, doi: 10.1371/journal.pone.0102591.

Pirini M, Rocchi L, Sensi M, Chiari L. A computational modelling approach to investigate different targets in deep brain stimulation for Parkinson's disease. J Comput Neurosci. 2009 Feb; 26(1):91–107. doi: 10.1007/s10827-008-0100-z.

Rattay F, Minassian K, Dimitrijevic MR. Epidural electrical stimulation of posterior structures of the human lumbosacral cord: 2. quantitative analysis by computer modeling. Spinal Cord. 2000 Aug; 38(8):473–489. doi: 10.1038/sj.sc.3101039.

Rattay F, Resatz S, Lutter P, Minassian K, Jilge B, Dimitrijevic MR. Mechanisms of Electrical Stimulation with Neural Prostheses: Mechanisms in Electrical Stimulation. Neuromodulation: Technology at the Neural Interface. 2003 Jan; 6(1):42–56. http://doi.wiley.com/10.1046/j.1525-1403.2003.03006.x, doi: 10.1046/j.1525-1403.2003.03006.x.

Rattay F. Analysis of Models for External Stimulation of Axons. IEEE Trans Biomed Eng. 1986 Oct; BME-33(10):974–977. http://ieeexplore.ieee.org/document/4122186/, doi: 10.1109/TBME.1986.325670.

Rattay F, Bassereh H, Stiennon I. Compartment models for the electrical stimulation of retinal bipolar cells. PLOS ONE. 2018 Dec; 13(12):e0209123. https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0209123, doi: 10.1371/journal.pone.0209123, publisher: Public Library of Science.

Rizzuto DS, Madsen JR, Bromfield EB, Schulze-Bonhage A, Seelig D, Aschenbrenner-Scheibe R, Kahana MJ. Reset of human neocortical oscillations during a working memory task. Proceedings of the National Academy of Sciences. 2003 Jun; 100(13):7931–7936. http://www.pnas.org/cgi/doi/10.1073/pnas.0732061100, doi: 10.1073/pnas.0732061100.

Rougier NP. A Density-Driven Method for the Placement of Biological Cells Over Two-Dimensional Manifolds. Front Neuroinform. 2018 Mar; 12:12. http://journal.frontiersin.org/article/10.3389/fninf.2018.00012/full, doi: 10.3389/fninf.2018.00012.

Rubin JE, Terman D. High frequency stimulation of the subthalamic nucleus eliminates pathological thalamic rhythmicity in a computational model. J Comput Neurosci. 2004 Jun; 16(3):211–235. doi: 10.1023/B:JCNS.0000025686.47117.67.

Segneri M, Bi H, Olmi S, Torcini A. Theta-Nested Gamma Oscillations in Next Generation Neural Mass Models. Front Comput Neurosci. 2020; 14. https://www.frontiersin.org/articles/10.3389/fncom.2020.00047/full, doi: 10.3389/fncom.2020.00047, publisher: Frontiers.

Solomon EA, Sperling MR, Sharan AD, Wanda PA, Levy DF, Lyalenko A, Pedisich I, Rizzuto DS, Kahana MJ. Theta-burst stimulation entrains frequency-specific oscillatory responses. Brain Stimulation. 2021 Sep; 14(5):1271–1284. https://linkinghub.elsevier.com/retrieve/pii/S1935861X21002163, doi: 10.1016/j.brs.2021.08.014.

Stimberg M, Brette R, Goodman DF. Brian 2, an intuitive and efficient neural simulator. eLife. 2019 Aug; 8:e47314. https://doi.org/10.7554/eLife.47314, doi: 10.7554/eLife.47314, publisher: eLife Sciences Publications, Ltd.

Sun J, Kapur J. M-type potassium channels modulate Schaffer collateral-CA1 glutamatergic synaptic transmission. J Physiol. 2012 Aug; 590(16):3953–3964. doi: 10.1113/jphysiol.2012.235820.

Suthana N, Fried I. Deep brain stimulation for enhancement of learning and memory. Neuroimage. 2014 Jan; 85 Pt 3:996–1002. doi: 10.1016/j.neuroimage.2013.07.066.

Suthana N, Haneef Z, Stern J, Mukamel R, Behnke E, Knowlton B, Fried I. Memory Enhancement and Deep-Brain Stimulation of the Entorhinal Area. New England Journal of Medicine. 2012 Feb; 366(6):502–510. https://doi.org/10.1056/NEJMoa1107212, doi: 10.1056/NEJMoa1107212, publisher: Massachusetts Medical Society _eprint: https://doi.org/10.1056/NEJMoa1107212.

Takeuchi Y, Nagy AJ, Barcsai L, Li Q, Ohsawa M, Mizuseki K, Berényi A. The Medial Septum as a Potential Target for Treating Brain Disorders Associated With Oscillopathies. Frontiers in Neural Circuits. 2021; 15:66. https://www.frontiersin.org/article/10.3389/fncir.2021.701080, doi: 10.3389/fncir.2021.701080.

Tass PA. A model of desynchronizing deep brain stimulation with a demand-controlled coordinated reset of neural subpopulations. Biological Cybernetics. 2003 Aug; 89(2):81–88. http://link.springer.com/10.1007/s00422-003-0425-7, doi: 10.1007/s00422-003-0425-7.

Titiz AS, Hill MRH, Mankin EA, M Aghajan Z, Eliashiv D, Tchemodanov N, Maoz U, Stern J, Tran ME, Schuette P, Behnke E, Suthana NA, Fried I. Theta-burst microstimulation in the human entorhinal area improves memory specificity. eLife. 2017 Oct; 6:e29515. https://doi.org/10.7554/eLife.29515, doi: 10.7554/eLife.29515, publisher: eLife Sciences Publications, Ltd.

Torben-Nielsen B, Uusisaari M, Stiefel K. A comparison of methods to determine neuronal phase-response curves. Frontiers in Neuroinformatics. 2010; 4:6. https://www.frontiersin.org/article/10.3389/fninf.2010.00006, doi: 10.3389/fninf.2010.00006.

Tort ABL, Komorowski R, Eichenbaum H, Kopell N. Measuring Phase-Amplitude Coupling Between Neuronal Oscillations of Different Frequencies. J Neurophysiol. 2010 Aug; 104(2):1195–1210. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2941206/, doi: 10.1152/jn.00106.2010.

Tort ABL, Komorowski RW, Manns JR, Kopell NJ, Eichenbaum H. Theta–gamma coupling increases during the learning of item–context associations. PNAS. 2009 Dec; 106(49):20942–20947. https://www.pnas.org/content/106/49/20942, doi: 10.1073/pnas.0911331106, publisher: National Academy of Sciences Section: Biological Sciences.

Tort ABL, Kramer MA, Thorn C, Gibson DJ, Kubota Y, Graybiel AM, Kopell NJ. Dynamic cross-frequency couplings of local field potential oscillations in rat striatum and hippocampus during performance of a T-maze task. PNAS. 2008 Dec; 105(51):20517–20522. https://www.pnas.org/content/105/51/20517, doi: 10.1073/pnas.0810524105, publisher: National Academy of Sciences Section: Biological Sciences.

Toth K, Borhegyi Z, Freund T. Postsynaptic targets of GABAergic hippocampal neurons in the medial septum-diagonal band of broca complex. J Neurosci. 1993 Sep; 13(9):3712–3724. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6576440/, doi: 10.1523/JNEUROSCI.13-09-03712.1993.

Traub RD, Jefferys JGR, Whittington MA. Simulation of Gamma Rhythms in Networks of Interneurons and Pyramidal Cells. Journal of Computational Neuroscience. 1997; p. 10.

Tsanov M, Manahan-Vaughan D. Long-Term Plasticity Is Proportional to Theta-Activity. PLoS One. 2009 Jun; 4(6):e5850. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2688745/, doi: 10.1371/journal.pone.0005850.

Unal G, Joshi A, Viney TJ, Kis V, Somogyi P. Synaptic Targets of Medial Septal Projections in the Hippocampus and Extrahippocampal Cortices of the Mouse. J Neurosci. 2015 Dec; 35(48):15812–15826. https://www.jneurosci.org/content/35/48/15812, doi: 10.1523/JNEUROSCI.2639-15.2015, publisher: Society for Neuroscience Section: Articles.

Varga V, Hangya B, Kránitz K, Ludányi A, Zemankovics R, Katona I, Shigemoto R, Freund TF, Borhegyi Z. The presence of pacemaker HCN channels identifies theta rhythmic GABAergic neurons in the medial septum. J Physiol. 2008 Aug; 586(16):3893–3915. doi: 10.1113/jphysiol.2008.155242.

Wang XJ. Pacemaker Neurons for the Theta Rhythm and Their Synchronization in the Septohippocampal Reciprocal Loop. Journal of Neurophysiology. 2002 Feb; 87(2):889–900. https://journals.physiology.org/doi/full/10.1152/jn.00135.2001, doi: 10.1152/jn.00135.2001, publisher: American Physiological Society.

Weerasinghe G, Duchet B, Cagnan H, Brown P, Bick C, Bogacz R. Predicting the effects of deep brain stimulation using a reduced coupled oscillator model. PLoS Comput Biol. 2019 Aug; 15(8):e1006575. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6701819/, doi: 10.1371/journal.pcbi.1006575.

Williams JM, Givens B. Stimulation-induced reset of hippocampal theta in the freely performing rat. Hippocampus. 2003 Jan; 13(1):109–116. https://onlinelibrary.wiley.com/doi/10.1002/hipo.10082, doi: 10.1002/hipo.10082, publisher: John Wiley & Sons, Ltd.

979 **Figures**

Figure 1. Dynamical computational model of the hippocampal formation and medial septum oscillatory drive. A. Anatomical representation of the neuronal types and interconnections within and between the medial septum and the hippocampal formation (EC: entorhinal cortex, DG: dentate gyrus, CA3 and CA1 fields of the hippocampus) **B.** Simplified anatomy of the hippocampal formation, modeled as a 15-mm thick cylindrical slice, with spatially segregated excitatory and inhibitory neurons (blue: excitatory neurons, consisting of granule cells in DG and pyramidal cells in other areas; red: inhibitory basket cells). **C.** Model architecture and connectivity. Each area is comprised of one excitatory and one inhibitory neuronal population. Theta drive is provided through input from the medial septum, which is modeled as a set of 250 Kuramoto oscillators and receives feedback connections from CA1. Electrical stimulation Is modeled as an intracellular current affecting both excitatory and inhibitory populations in the targeted area (shown here for CA1). **D.** Illustration of Kuramoto oscillators with two different levels of synchronization. Each dot represents one oscillator, its position on the circle indicates its phase and its color its angular velocity. Higher synchronization corresponds to a clustering of the dots around a similar phase. "r" indicates the order parameter, which is a measure of synchronization.

A. Theta Rhythm

f_θ = 4.2 Hz

Phase

$\pi/2$

CA1 Spiking Activity

μ_I: 50.2 Hz

μ_E: 12.8 Hz

CA1 Firing Rates

Inhibitory

Excitatory

100 Hz

Spectrograms

Inhibitory

Excitatory

a.u.

Time [s]

B. CA1 Spectral Analysis

Inhibitory

Excitatory

Max

PSD

Frequency [Hz]

C. Phase-Amplitude Coupling

θ_{trough}

80 Hz

60 Hz

40 Hz

θ_{peak}

MI: 0.19

min   max

D. Phase Response Curve

$\pi/4$

$\pi/8$

$\Delta\phi$ [rad]

$-\pi/8$

$-\pi/4$

Stim. Amp. [nA]

2 5 10 20 40

$-\pi$ $-\pi/2$ 0 $\pi/2$ π

Stim. Phase [rad]

Figure 2. Theta-nested gamma oscillations and theta phase reset in response to a single stimulation pulse. A. Representative example of the network behavior, spontaneously producing theta-nested gamma oscillations and characterized by a reset of the theta phase following a single stimulation pulse (vertical grey line, applied here in CA1 at a theta phase of $\pi/2$, i.e. in the middle of the descending slope). Top to bottom: theta rhythm originating from the medial septum and provided as an input to the EC (f_θ: mean oscillation frequency), instantaneous phase of the theta rhythm, raster plots indicating the spiking activity of CA1 excitatory (blue) and inhibitory (red) neurons (μ_E and μ_I: mean firing rates within the shaded area), average population firing rates in CA1 (computed as a windowed moving average with a sliding window of 100 ms with 99% overlap), spectrograms for each CA1 population (windowed short-time fast Fourier transform using a Hann sliding window: 100 ms with 99% overlap). Spectrograms show gamma oscillations (around 60 Hz) modulated by the underlying theta rhythm (~ 4 Hz|), indicating theta-gamma PAC. Theta phase reset after stimulation is associated with a rebound of spiking activity and theta-nested gamma oscillations. **B.** Power spectral densities of the CA1 firing rates. Theta peaks are found at 4 Hz for excitatory and inhibitory cells. Gamma activity is located between 40 and 80 Hz. **C.** PAC as a function of theta phase and gamma frequency. The polar plot represents the amplitude of gamma oscillations (averaged across all theta cycles, see methods) at each phase of theta (theta range: 3-9 Hz, phase indicated as angular coordinate) and for different gamma frequencies (radial coordinate, binned in 10-Hz ranges), indicating that gamma oscillations between 40 and 80 Hz occur preferentially around the peak of theta. The MI gives an overall quantification of how the phase of low-frequency oscillations (3-9 Hz) modulates the amplitude of higher-frequency oscillations (40-80 Hz) (see *Methods* and *Figure supplement 2*). **D.** PRC in response to a single stimulation pulse applied in CA1 at various phases of the ongoing theta rhythm and for various stimulation amplitudes (color-coded). The phase difference (left y-axis) shows the theta phase induced by the stimulation pulse (computed 2.5 ms after the pulse), compared to the phase computed at the same time in a scenario without stimulation. Positive and negative phase differences respectively indicate phase advances and delays. The grey trace shows the normalized amplitude of theta (right y axis) for different phases, used to indicate the peak and trough of the rhythm. Stimulation applied in the ascending slope of theta ($[-\pi, 0]$) produced a phase advance and accelerated the rhythm towards its peak (0 radians). Conversely, stimulation during the descending slope ($[0, \pi]$) produced a phase delay that slowed down the rhythm. Higher stimulation amplitudes yielded a stronger effect.

Figure 3. CAN and adaptation (M) currents modulate neuronal responses to single-pulse stimulation in the absence of theta input. A.
Network response to single-pulse stimulation in the absence of medial septum input. Stimulation (grey vertical line, 10 nA, applied in CA1) induced an instantaneous burst of activity lasting about 20 ms in both excitatory and inhibitory CA1 neurons, followed by a secondary burst approximately 200 ms later (raster plot and firing rate traces), associated with specific CAN- and M- currents dynamics (bottom traces, illustrated for a representative CA1 excitatory neuron). Positive (I_M) and negative (I_{CAN}) currents indicate respectively a hyperpolarizing and depolarizing effect on the cell membrane potential. **B.** Similar representation as in A., but in the absence of CAN channel. Stimulation induced only a single burst of activity, indicating that CAN channels are necessary to observe a rebound of activity. **C.** Number of bursts in CA1 spiking activity following a single stimulation pulse at various amplitudes (x-axis), shown both in the presence and absence of CAN channels in excitatory neurons. The absence of the CAN current leads to the abolition of the second burst, irrespective of stimulation amplitude.

Figure 4. Medial septum oscillatory drive and stimulation amplitude govern the steady-state response to single-pulse stimulation. A-C. Network responses to single-pulse stimulation (vertical line) under medial septum input (with phase reset), shown for different amplitudes of the medial septum oscillatory drive (A-C: increasing oscillator amplitudes). **A.** Low oscillatory input: stimulation induces only two bursts of spiking activity as in Figure 3. **B.** Medium oscillatory input: stimulation induces transient activity that fades away after about 500 ms. **C.** Higher oscillatory input: a single stimulation pulse switches network behavior from no activity to sustained oscillations driven by the medial septum. **D.** Steady-state response to single-pulse stimulation as a function of medial septum oscillatory input (x-axis) and stimulation amplitude (y-axis), characterized by three metrics: theta power (3-9 Hz), gamma power (40-80 Hz), and PAC (quantified using the MI). White dots: parameter combinations corresponding to panels A-C.

Figure 5. Single-pulse stimulation phase differentially affects network responses depending on the presence of theta phase reset. All results shown here were obtained for parameters from Figure 4C (theta oscillation amplitude: 0.18 nA, stimulation amplitude: 8.0 nA). A single stimulation pulse was delivered at the peak (**A, B**) or trough (**C, D**) of the underlying theta rhythm, either in the presence (**A, C**) or absence (**B, D**) of theta phase reset. With phase reset, both peak and trough stimulation switch network behavior from no activity to sustained oscillations. Without phase reset, only peak stimulation can induce sustained oscillations. **E.** Quantification of theta power, gamma power, and PAC (measured using the MI) in CA1 excitatory (blue) and inhibitory (red) populations in all 4 cases (metrics are computed in the shaded areas of panels A-D).

Figure 6. Pulse train stimulation restores theta-nested gamma oscillations depending on stimulation timing and theta phase reset. All results shown here were obtained for parameters from Figure 4B (theta oscillation amplitude: 0.07 nA; stimulation amplitude: 8 nA). Representations are similar to Figure 5, with the difference that stimulation consisted of a pulse train delivered at 6 Hz for a duration of 2 s (individual pulses indicated by grey dots, first pulse by a triangle). The pulse train was delivered at the peak (A, B) or trough (C, D) of the underlying theta rhythm, either in the presence (A, C) or absence (B, D) of theta phase reset. With phase reset, both peak and trough stimulation switch network behavior from no activity to sustained oscillations. Without phase reset, only peak stimulation can induce sustained oscillations. **E.** Quantification of theta power, gamma power, and PAC (measured using the MI) in CA1 excitatory (blue) and inhibitory (red) populations in all 4 cases (metrics are computed during the pulse train, within the shaded areas of panels A-D).

980 **Appendix**

981 **Supplementary Figures**

Figure S1. (Figure 1 – Figure Supplement 1) Influence of the number of Kuramoto oscillators on their collective dynamics. Left: influence of the number of Kuramoto oscillators (N) on the convergence over time of the order parameter (r), which is a measure of synchronization of the set of oscillators. The above results were obtained using a synchronization ratio k/N of 25. Upon initialization, the oscillators' natural frequencies (ω_i) were normally distributed with a center frequency ($f_0 = 6$ Hz). Initial phases were uniformly distributed around the unit circle ($U \sim [0, 2\pi]$). **Right**: histograms of the natural frequencies (ω_i), together with the distribution mean (μ) and standard error of the mean ($\sigma/\sqrt{N}$). As the number of oscillators increases, the mean frequency appears closer to the desired center frequency of 6 Hz, and the standard error of the mean is reduced. Overall, increasing the number of oscillators past 250 did not yield any substantial change in the convergence of the order parameter or the distribution of natural frequencies. This number of oscillators was therefore chosen for all subsequent analyses.

Figure S2. (Figure 2 – Figure Supplement 1) Whole-network behavior producing theta-nested gamma oscillations and theta phase reset in response to a single stimulation pulse. This figure shows the behavior of all areas of the hippocampal formation during the simulation illustrated in Figure 2A. All areas display bursts of gamma activity that occur almost synchronously at the peak of each theta cycle in both excitatory and inhibitory neuronal populations (μ_E and μ_I: mean firing rates within the shaded area). The CA1 excitatory population firing rate is supplied back to the medial septum. A single stimulation pulse (vertical line and grey triangle) delivered to both CA1 populations at a phase of $\pi/2$ (i.e. halfway through the descending slope) resets the phase of the theta rhythm according to the PRC shown in Figure 2D.

Figure S3. (Figure 2 – Figure Supplement 2) Quantification of PAC with and without noise. A. Quantifying PAC in the absence of noise produced inaccurate identification of the coupled frequency bands, due to the complete absence of oscillations at some frequencies. All analyses are based on the CA1 firing rates (top traces) during a representative simulation. Power spectral densities of these firing rates (**left**) indicate that some frequencies have a power of 0. PAC of the excitatory population was assessed using two graphical representations, the polar plot (**middle**) and comodulogram (**right**), and quantified using the MI. The comodulogram was calculated by computing the MI across 80% overlapping 1-Hz frequency bands in the theta range and across 90% overlapping 10-Hz frequency bands in the gamma range and subsequently plotted as a heat map. In the absence of noise, a slow theta frequency centered around 5 Hz is found to modulate a broad range of gamma frequencies between 40 and 100 Hz. The value indicated on the comodulogram indicates the average MI in the 3-9 Hz theta range and 40-80 Hz gamma range. As in Figure 2, the polar plot represents the amplitude of gamma oscillations (averaged across all theta cycles) at each phase of theta (theta range: 3-9 Hz, phase indicated as angular coordinate) and for different gamma frequencies (radial coordinate, binned in 1-Hz ranges). **B.** Adding uniform noise to the firing rate (with an amplitude ranging between 15 and 25% of the maximum firing rate) improved the identification of the coupled frequency bands. In this case, the slower theta frequency centered around 5 Hz modulates a gamma band located between 45 and 75 Hz.

Figure S4. (Figure 2 – Figure Supplement 3) Network behavior generated by Kuramoto oscillators with non-physiological phase response functions. Each panel is similar to Figure 2A, but with a different offset added to the phase response function of the Kuramoto oscillators (see methods, *Equation 4*) **A.** A phase offset of $-\pi/2$ leads to an overall theta oscillation of 4 Hz, with a second peak following the main theta peak. **B.** A phase offset of $+\pi/2$ reduces the peak of theta, resetting the rhythm to the middle of the ascending phase. **C.** A phase offset of $\pm\pi$ leads to the CA1 output resetting the theta rhythm to the trough of theta. A sudden desynchronization of the network of Kuramoto oscillators temporarily abolishes gamma oscillations.

Figure S5. (Figure 4 – Figure Supplement 1) Whole-network behavior in response to single-pulse stimulation under low theta oscillatory drive. This figure shows the behavior of all areas of the hippocampal formation during the simulation illustrated in Figure 4A. A single stimulation pulse delivered to both CA1 populations induces two bursts of spiking activity that propagate only from CA1 to EC.

Figure S6. (Figure 4 – Figure Supplement 2) Whole-network behavior in response to single-pulse stimulation under medium theta oscillatory drive. This figure shows the behavior of all areas of the hippocampal formation during the simulation illustrated in Figure 4B. A single stimulation pulse delivered to both CA1 populations induces transient oscillatory activity that propagates across all areas. All neurons briefly follow the theta rhythm driven by the medial septum, but this activity quickly fades away after a few theta cycles.

Figure S7. (Figure 4 – Figure Supplement 3) Whole-network behavior in response to single-pulse stimulation under higher theta oscillatory drive. This figure shows the behavior of all areas of the hippocampal formation during the simulation illustrated in Figure 4C. A single stimulation pulse switches network behavior from no activity to sustained oscillations driven by the medial septum

Figure S8. (Figure 4 – Figure Supplement 4) Synchronization level of the oscillators minimally affects the production of theta-nested gamma oscillations. A-C. Network responses to single-pulse stimulation (vertical line) under medial septum input (with phase reset), shown for different synchronization parameter ratios of the Kuramoto oscillators (A-C: $\frac{k}{N}$ of 5, 15, and 35 respectively; stimulation amplitude: 8 nA). **Panels A.** and **B.** show that the network retains its bistable characteristics as presented in Figure 4 across a large range of synchronization levels. **Panel C.** indicates that the model can produce oscillations without external stimulation when the synchronization level is higher. **D.** Steady-state response to single-pulse stimulation as a function of medial septum oscillatory input (x-axis) and synchronization parameter ratio (k/N, y-axis), characterized by three metrics: theta power (3-9 Hz), gamma power (40-80 Hz), and PAC (quantified using the MI). White dots: parameter combinations corresponding to panels A-C