The conserved protein CBA1 is required for vitamin B$_{12}$ uptake in

different algal lineages

Andrew P. Sayer[1*], Marcel Llavero-Pasquina[1¶], Katrin Geisler[1], Andre Holzer[1‡], Freddy Bunbury[1§],

Gonzalo I. Mendoza-Ochoa[1], Andrew D. Lawrence[2], Martin J. Warren[3,4], Payam Mehrshahi[1], Alison

G. Smith[1*]

[1]Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge, CB2 3EA, UK

[2]School of Biological Sciences, University of Southampton, Southampton SO17 1BJ, UK

[3]School of Biosciences, University of Kent, Canterbury, Kent, CT2 7NJ, UK

[4]Quadram Institute Bioscience, Norwich Research Park, Norwich, NR4 7UA, UK

Current addresses:

* CRUK-AZ Functional Genomics Centre, Milner Therapeutics Institute, University of Cambridge,

Puddicombe Way, Cambridge, CB2 0AW, UK

¶ Environmental Science and Technology Institute (ICTA-UAB), Universitat Autònoma de Barcelona

(UAB) 08193 Bellaterra (Cerdanyola del Vallès) - Spain

‡Center for Bioinformatics and Department of Computer Science, Saarland University, Saarbrücken,

Germany

§Department of Plant Biology, Carnegie Science, 260 Panama Street, Stanford CA 94305, USA

* Author for correspondence: Alison Smith

Email: as25@ cam.ac.uk; orcid.org/0000-0001-6511-5704;

Phone: +44-1223 333952; Fax: +44-1223 333953

Short title: Identification of a B_{12} uptake protein in algae

One sentence summary: Knockout mutants and physiological studies demonstrate that the CBA1 protein is essential for uptake of vitamin B_{12} in both *Chlamydomonas reinhardtii* and the unrelated *Phaeodactylum tricornutum.*

Keywords: cobalamin, *Chlamydomonas reinhardtii, Phaeodactylum tricornutum*, insertional mutagenesis, CLiP mutants, CRISPR-Cas9, riboswitch

Manuscript length: 7241 words

Abstract

Microalgae play an essential role in global net primary productivity and global biogeochemical cycling, but despite their phototrophic lifestyle, over half of algal species depend on a supply of the corrinoid vitamin B_{12} (cobalamin) for growth. This essential organic micronutrient is produced only by a subset of prokaryotic organisms, which implies that for algal species to use this compound, they must first acquire it from external sources. Previous studies have identified protein components involved in vitamin B_{12} uptake in bacterial species and humans. However, little is known about how it is taken up in algae. Here, we demonstrate the essential role of a protein, CBA1 (for cobalamin acquisition protein 1), in B_{12} uptake in *Phaeodactylum tricornutum*, using CRISPR-Cas9 to generate targeted knockouts, and in *Chlamydomonas reinhardtii*, by insertional mutagenesis. In both cases, CBA1 knockout lines are no longer able to take up exogenous vitamin B_{12}. Complementation of the *C. reinhardtii* mutants with the wildtype *CBA1* gene restores B_{12} uptake, and regulation of *CBA1* expression via a riboswitch element can be used to control the phenotype. When visualised by confocal microscopy, a YFP-fusion with *C. reinhardtii* CBA1 shows association with membranes. A bioinformatics analysis found that CBA1-like sequences are present in all the major eukaryotic phyla. Its presence is correlated with B_{12}-dependent enzymes in many, although not all, taxa, suggesting CBA1 has a conserved role. Our results thus provide insight into the molecular basis of algal B_{12} acquisition, a process that likely underpins many interactions in aquatic microbial communities.

INTRODUCTION

Microalgae are a diverse group of eukaryotic organisms that thrive in all aquatic environments. They form the basis of most aquatic food chains and are major contributors to global primary productivity, with marine microalgae responsible for an estimated 30% of total carbon fixation (Field et al., 1998). Understanding the drivers that support algal growth is thus of considerable ecological importance. Despite their photoautotrophic lifestyle, a widespread trait in algae is dependence on an external source of an organic micronutrient, vitamin B_{12} (cobalamin), a complex cobalt-containing corrinoid molecule. Approximately half of algal species surveyed across the eukaryotic tree of life require B_{12} for growth (Croft et al., 2005). However, the proportion of B_{12}-dependent species differs between algal groups, from 30% (n=148) of Chlorophytes to 96% (n=27) of algal species that participate in harmful algal blooms (Tang et al., 2010). Within algal lineages, there is no evidence that any can produce B_{12} *de novo*, so this auxotrophy is not due to loss of one or more biosynthetic genes. Rather, the requirement for B_{12} stems from the fact that it is an essential cofactor for methionine synthase (METH), and species that can grow without supplementation have an alternative, B_{12}-independent, isoform of this enzyme called METE (Croft et al., 2005; Helliwell et al., 2011). Many microalgae, including the green alga *Chlamydomonas reinhardtii* and the unrelated diatom *Phaeodactylum tricornutum*, encode both forms of methionine synthase and utilise METE in the absence of exogenous B_{12}, but take up and utilise the compound if it becomes available (Helliwell et al., 2011). Under those conditions, the expression of *METE*, which has been found to have a lower catalytic rate than METH (Gonzalez et al. 1992), is repressed, and cells rely on METH activity.

The biosynthetic pathway for B_{12} is confined to prokaryotes (Warren et al., 2002) and indeed only a subset of bacteria encode the entire set of 20 or so enzymes required to synthesise corrinoids from the common tetrapyrrole precursor (Shelton et al., 2019), with many eubacterial species also reliant on an external source. In some cases, this is due to the loss of one or a few enzymes of the biosynthetic pathway, but in many bacteria the pathway is absent altogether and auxotrophy is the consequence of relying on one or more B_{12}-dependent enzymes, such as METH. In microalgae, supplementation of cultures of *P. tricornutum* with B_{12} increases its growth rate subtly (Bertrand et al., 2012) and in *C.*

reinhardtii use of METH confers thermal tolerance (Xie et al., 2013). More direct evidence for a

selective advantage is demonstrated by the fact that an experimentally-evolved *metE* mutant of *C.*

reinhardtii predominates in mixed populations with wild-type cells over tens of cell generations, as

long as B_{12} is included in the medium (Helliwell et al., 2015). This is despite the fact that in the

absence of B_{12}, the *metE* mutant is non-viable within a few days (Bunbury et al., 2020).

The minimum levels of B_{12} in the medium needed to support growth of laboratory cultures of algal

B_{12}-auxotrophs are in the range of 10-50 pM (Croft et al., 2005), whereas B_{12} concentrations have

been reported to be just 5-13 pM in freshwater systems (Ohwada, 1973). A similar value of 6.2 pM is

the average value in most marine environments, although up to 87 pM could be detected in some

coastal waters (Sañudo-Wilhelmy et al., 2014), which may be linked to the higher cobalt

concentrations measured there (Panzeca et al., 2009). Given the limiting levels of B_{12} in the

environment, its relatively short half-life (in the order of days) in surface water (Carlucci et al., 2007;

Sañudo-Wilhelmy et al., 2014), and that as a large polar molecule it is unlikely to simply diffuse

across cellular membranes, it is clear that algae must have an efficient means to take up B_{12}. In

bacteria, the molecular mechanisms for B_{12} uptake have been extensively characterised. The B_{12}

transport and utilisation (*btu*) operon is perhaps the best known (Kadner, 1990), comprising BtuB, a

TonB-dependent transporter in the outer membrane, a B_{12}-binding protein, BtuF, located in the

periplasm, and BtuC and BtuD, components of an ATP-binding cassette (ABC) transporter that sits in

the inner membrane (Borths et al., 2002). In mammals, dietary B_{12} is bound to intrinsic factor in the

ileum and taken up from the gut via receptor-mediated endocytosis (Nielsen et al., 2012). It is then

transported between and within cells via multiple B_{12} transport proteins (Banerjee et al., 2021; Choi

and Ford, 2021). These include LMBD1/ABCD4, the latter being an integral membrane ABC

transporter in the lysosomal membrane of gut epithelial cells, which facilitates delivery of B_{12} into the

cytosol, and MRP1 (or ABCC1), another ABC transporter that has sequence similarity to BtuCD and

is involved in export of free B_{12} into the plasma where it binds to the main B_{12} transport protein,

transcobalamin (Beedholm-Ebsen et al., 2010). Mice *mrp1* mutants were still able to transport a small

amount of cobalamin out of cells, indicating redundant mechanisms for this function that have not yet

been identified. Cobalamin circulating in the plasma bound to transcobalamin can then be taken up by

other cells via receptor-mediated endocytosis (Nielsen et al., 2012).

In contrast to these well-studied processes in bacteria and mammals, the understanding of B_{12}

acquisition in microalgae is more limited. A survey of microalgal species, including marine and

freshwater taxa and those that require B_{12} (for example *Euglena gracilis*, *Thalassiosira pseudonana*)

and non-requirers (such as *P. tricornutum*, *Dunaliella primolecta*), found that many released a 'B_{12}-

binder' into the medium, likely a protein, that appeared to sequester B_{12} from solution and thereby

inhibited growth of B_{12}-dependent algae (Pintner and Altmeyer, 1979). Its role was unknown, but it

was postulated that it might be involved in competition for resources between microalgal species in

the environment. Subsequently, a protein was purified from the medium of cultures of *T. pseudonana*

with a high affinity binding constant of 2 pM for B_{12} (Sahni et al., 2001). In its native state it was an

oligomer of >400 kDa, with subunits of ~80 kDa and the amino acid profile was determined, but it

was not possible to obtain sufficient amounts to characterise further. A different approach was taken

by Bertrand et al. (2012), who conducted a transcriptomics and proteomics study of *P. tricornutum*

and *T. pseudonana* grown under low or sufficient B_{12} conditions. This led to the identification of a

gene highly upregulated at the transcript and protein level in the absence of B_{12}. Overexpression of

this protein in *P. tricornutum* resulted in an increase in the rate of B_{12} uptake, and the protein was

named CoBalamin Acquisition protein 1 (CBA1) although no direct role was established. In this study

we have taken a mutagenesis approach to try to identify genes responsible for B_{12} uptake in both *P.*

tricornutum and *C. reinhardtii*, including extending the work on CBA1. In addition, we have

determined the extent to which candidate proteins are conserved throughout the algal lineages,

making   use   of   recent   increases   in   algal   sequencing   data.

RESULTS

***P. tricornutum CBA1* knockout lines do not take up B$_{12}$**

Previous work showed that overexpression of CBA1 in *P. tricornutum* conferred enhanced B$_{12}$ uptake

rates (Bertrand et al., 2012) but the study did not demonstrate whether it was essential for this process.

To address this question, *CBA1* knockout lines were generated in *P. tricornutum* strain 1055/1 (Table

S1) by CRISPR-Cas9 editing, using a homologous recombination repair template that included a

nourseothricin resistance (*NAT*) cassette (Figure 1a). CRISPR-Cas9 lines were cultured on selective

media and screened for the absence of WT alleles at the *PtCBA1* locus (Phatr3_J48322) using PCR

(Figure 1b). When the *PtCBA1* gene was amplified (top panel, Figure 1b) from ΔCBA1-1 with

primers flanking the homologous recombination regions, two bands were detected; the larger of these

corresponded to the WT amplicon, whilst the smaller band corresponded to a replacement of *CBA1* by

NAT, suggesting that this strain is a mono-allelic knockout. For ΔCBA1-2, the *PtCBA1* gene primers

amplified a single smaller product, suggesting that this was a bi-allelic knockout, whereas the *PtCBA1*

ORF primers (bottom panel of Figure 1b) did not amplify anything, indicating a disruption

specifically in this region. Similarly, no band was detected with primers that amplify across the 5' end

of the NAT knock-in (HR primers), which might indicate further disruptions upstream of the 5'HR

region of ΔCBA1-2. Although a larger band than for WT was amplified in ΔCBA1-3 using the

PtCBA1 gene primers, those for the *PtCBA1* ORF amplified a smaller product; in both cases a single

band was observed indicating a bi-allelic deletion at the sgRNA target sites.

To test whether the ΔCBA1 lines were affected in their ability to take up vitamin B$_{12}$ we developed a

standardised B$_{12}$-uptake assay, detailed in Materials and Methods. In brief, algal cells were grown to

the same growth stage and adjusted to the same cell density, then incubated in media containing a

known amount of cyanocobalamin for one hour. Thereafter, cells were pelleted by centrifugation and

the amount of B$_{12}$ determined in the cell pellet and the media fraction using a *Salmonella typhimurium*

bioassay (Bunbury et al., 2020). For each sample, the B$_{12}$ measured in the cellular and media fractions

were added to provide an estimated 'Total' and compared to the amount of B$_{12}$ added initially (Figure

1c, dashed line), to determine the extent of recovery. For the WT strain, most of the added B_{12} was

found in the cellular fraction. The mono-allelic knockout line ΔCBA1-1 consistently showed ~20-

30% B_{12} uptake relative to the WT strain. This suggested that a single copy of *PtCBA1* is sufficient to

confer B_{12} uptake in *P. tricornutum*, but not to the same extent as the WT strain. In contrast, for the

two bi-allelic knockout lines (ΔCBA1-2 and ΔCBA1-3) no B_{12} was detected in the cellular fraction in

any experiment, indicating that vitamin B_{12} uptake was fully impaired in the absence of a functional

PtCBA1 copy, at least at the limit of detection of the B_{12} bioassay (of the order of 10 pg). These

results expand our understanding of *PtCBA1* by demonstrating that its presence is essential for B_{12}

uptake and indicates that there is no functional redundancy to *PtCBA1*.

Insertional mutagenesis identified the *C. reinhardtii* homologue of *CBA1*

Bertrand et al. (2012) reported that there were no detectable *CBA1* homologues in algal lineages

outside the Stramenopiles, so to investigate B_{12} uptake in *C. reinhardtii*, we decided to take an

insertional mutagenesis approach. We took advantage of the fact that B_{12} represses expression of the

METE gene at the transcriptional level via the promoter (P_{METE}), and that reporter genes driven by this

genetic element respond similarly (Helliwell et al., 2014), to develop a highly sensitive screen for

lines no longer able to respond to B_{12}. We hypothesised that, since P_{METE} is likely to respond

specifically to intracellular B_{12}, P_{METE} would not be repressed in strains unable to take up B_{12} from the

media, so the reporter would be expressed and functional. If the reporter were an antibiotic resistance

gene, this would allow identification of B_{12} uptake mutants in a more high-throughput manner than

the B_{12}-uptake assay. The background strain for insertional mutagenesis was made by transforming *C.*

reinhardtii strain UVM4 (Neupert et al., 2009) with plasmid pAS_R1 containing a paromomycin

resistance gene (*aphVIII*) under control of P_{METE} (Figure 2a, top construct). Lines of this strain were

tested for their responsiveness to B_{12} and paromomycin. One line, UVM4-T12, showed the

appropriate sensitivity with increasing repression of growth in paromomycin as B_{12} concentrations

were increased, the effect being more marked at 15-20 µg·ml^{-1} paromomycin than at 5-10 µg·ml^{-1}

(Figure 2b). This line thus allowed for an easily quantifiable growth phenotype that was

proportionally related to B_{12} concentration.

Insertional mutagenesis was carried out by transforming UVM4-T12 with a plasmid (pHyg3)

containing a hygromycin resistance gene (*aphVII)* under the control of the constitutively expressed

β2-tubulin promoter (Figure 2a, bottom construct), generating a population of UVM4-T12::pHyg3

lines with the cassette randomly inserted into the nuclear genome. By plating the products of the

transformation on solid TAP media supplemented with a range of paromomycin, hygromycin and

vitamin B_{12} concentrations (see Methods), 7 colonies were obtained. This was estimated to be from

approximately 5000 primary transformants, determined by plating the same volume on TAP plates

with the antibiotics but without B_{12}. These 7 putative insertional mutant (IM) lines were then assessed

for their ability to take up B_{12} using the B_{12} uptake assay. For UVM4, UVM4-T12 and insertional

lines from the plate without B_{12} (labelled Control 1-3), similar amounts of B_{12} were recovered from

the cellular and media fractions (Figure S1). This was also the case for 6 of the IM lines, suggesting

that they could still take up B_{12} and were likely false positives of the initial screen. However, no B_{12}

could be detected in the cellular fraction of UVM4-T12::pHyg3 #IM4 (hereafter referred to as IM4),

indicating that this mutant line did not take up B_{12}.

To obtain independent corroboration that IM4 was impaired in B_{12} uptake, cells of this mutagenized

line were incubated with a fluorescently-labelled B_{12} derivative, B_{12}-BODIPY (Lawrence et al., 2018),

and then imaged using confocal microscopy. *C. reinhardtii* cells were incubated in TAP medium

without B_{12}-BODIPY or with 1 µM B_{12}-BODIPY for 1 hour, washed with fresh media and

subsequently imaged. There was no signal detected in the channel used for B_{12}-BODIPY (589 nm

excitation; 607-620 nm detection) in samples without B_{12}-BODIPY added (Figure S2, top two rows),

indicating that the imaging protocol was specific to this compound. When B_{12}-BODIPY was added,

UVM4-T12 showed the B_{12}-BODIPY signal located within the algal cell (Figure S2, third row),

indicating that this signal could be effectively detected by the imaging protocol and that B_{12}-BODIPY

was being transported into the cells. In contrast, there was no B_{12}-BODIPY signal in IM4 cells,

supporting the hypothesis that B_{12} uptake was impaired in this mutant (Figure S2, bottom row). In

addition, the response of the *METE* gene to B_{12} in IM4 was assessed by RT-qPCR. UVM4 and IM4

cultures were grown in media with or without addition of B_{12} for 4 days in continuous light, after

which the cultures were harvested for RNA extraction and cDNA synthesis. As expected, *METE* was

repressed in UVM4 in the presence of B_{12} compared to no supplementation (Figure 3a), whereas IM4

showed similar *METE* expression in both conditions. This provided further support for disrupted B_{12}

uptake in this line.

To identify the genomic location of the causal mutation in IM4, short-read whole genome sequencing

was performed on DNA samples from UVM4, UVM4-T12 and IM4. The location of the pHyg3

cassette in IM4 was identified as described in Methods and found to have disrupted the

Cre12.g508644 locus (Figure S3a), an unannotated gene. To corroborate that disruption of the

Cre12.g508644 was responsible for the uptake-phenotype, two independent mutant lines of the gene

(LMJ-119922 and LMJ-042227) were ordered from the Chlamydomonas library project (CLiP)

collection (Li et al., 2016) and verified to be disrupted at this locus by PCR (Figure S3a). However,

when these knockout lines were tested for the ability to take up B_{12} using the B_{12} uptake assay, they

were both found to be able to do so to a similar extent as their parental strain, cw15 (Figure S3b). This

suggested that *Cre12.g508644* did not encode a protein essential for B_{12} uptake.

We therefore examined the genome sequence data more closely to determine the genetic cause for the

B_{12}-uptake phenotype of IM4. We had identified putative homologues of human proteins involved in

receptor-mediated endocytosis of B_{12}, such as ABCD4, LMBD1 (Rutsch et al., 2009; Coelho et al.,

2012) and MRP1 (Beedholm-Ebsen et al., 2010), in the *C. reinhardtii* genome by BLAST (data not

shown). However, given the widespread percentage of SNPs in the IM4 genome compared to UVM4,

it was not possible to identify any candidate causal mutations with confidence. Instead, manual

inspection of the DNA sequencing reads mapped to the reference strain revealed one locus,

Cre02.g081050, annotated as flagella-associated protein 24 (FAP24), where there was a unique

discontinuity, suggesting that there was an insertion at exon 2 in IM4 (Figure 3b; Figure S4a). The

sequence was bordered by a genome duplication of 8 bp (shown in blue in Figure S4a) and exhibited

imperfect inverted repeats at the terminal regions (TIRs), indicative of a transposable element. Reads

could not be assembled across the discontinuity to obtain the complete sequence of the insertion, but

using the left and right junction sequences as queries, three regions encoding two very similar genes

were identified (Figure S4b).

Remarkably, when the *Cre02.g081050* protein was used as a query in a BLAST search, one of the hits

recovered was the *PtCBA1* protein (22.9% sequence identity), even though the reciprocal sequence

search had not picked up the *C. reinhardtii* gene (Bertrand et al., 2012). The Phyre2 structural

prediction server (Kelley et al., 2015) was used to model the 3D structures of PtCBA1 and the *C.*

reinhardtii protein encoded by *Cre02.g081050* (Figure S5). The modelled proteins showed a high

degree of structural similarity to one another (root mean squared deviation (RMSD) = 2.333),

particularly with respect to the arrangement of alpha helices and lower cleft. Due to the sequence

similarity and predicted structural similarity, these proteins appeared to be homologous to one another

and Cre02.g081050 is hereafter referred to as CrCBA1.

To determine whether disruption of *CrCBA1* in IM4 was responsible for the impaired B_{12} uptake, we

investigated whether it was possible to restore its ability to take up B_{12} by transforming IM4 with the

wild-type *CrCBA1*. Construct pAS_C2 was designed with the *CrCBA1* promoter, *CrCBA1* open

reading frame (ORF) and terminator and included a 3' mVenus tag attached by a poly-glycine linker

(Figure 3c). IM4 was transformed with pAS_C2, and resulting lines were tested for the ability to take

up B_{12} using the B_{12} uptake assay. As observed previously, UVM4 was able to take up B_{12} whilst IM4

was unable to do so (Figure 3d). The CBA1 complementation line IM4::pAS_C2 showed B_{12} in the

cellular fraction at similar levels as in UVM4, thereby indicating that the mutant phenotype had been

complemented.

CrCBA1 CLiP mutant is unable to take up B_{12} and is complemented by the WT _CrCBA1_ gene

Given the many genetic changes in line IM4 compared to the parental UVM4-T12 strain caused by

the mutagenesis, it was essential to have independent corroboration that mutation of *CrCBA1* caused

the inability to take up B_{12}. Accordingly, we obtained two further CLiP mutants (LMJ-135929 and

LMJ-040682) with disruptions in intron 2 and introns 6/7 respectively of *CrCBA1* (Figure S6a) and

assessed them for their ability to take up B_{12} (Figure S6b). No B_{12} was detected in cells of LMJ-

040682, indicating complete inhibition of B_{12} uptake. Although LMJ-135929 cells accumulated some

B_{12}, this was less than half the amount of its parent strain cw15, suggesting partial impairment in

uptake, similar to the phenotype of the monoallelic *PtCBA1* knockout line (Figure 1c). However,

heterozygosity cannot be the explanation for *C. reinhardtii,* which is haploid, and instead indicates

that LMJ-135929 was likely to have just partial knockdown of the gene, probably because the

insertion is in an intron.

Nonetheless, to provide further confirmation that mutations in *CrCBA1* were responsible for the

observed impaired B_{12} uptake, we again tested whether the phenotype could be complemented with

the wild-type *CrCBA1* gene using both plasmid pAS_C2 (Figure 3b) and an additional construct

pAS_C3 (Figure 4a), in which expression of *CrCBA1* can be controlled by a thiamine pyrophosphate

(TPP) repressible riboswitch, RS_{THI4_4N} (Mehrshahi et al., 2020). In the absence of thiamine

supplementation of the cultures, the riboswitch is not active and the gene containing it is transcribed

and translated as normal; with thiamine addition, alternative splice sites are utilised, leading to

inclusion of an upstream ORF containing a stop codon in the mRNA, preventing translation from the

downstream start codon. LMJ-040682 was transformed with both pAS_C2 and pAS_C3, and

representative transformant lines selected via antibiotic resistance were obtained. These, together with

their parental strains were grown in the presence or absence of 10 μM thiamine for 5 days, and then

used in the B_{12} uptake assay. Transformants of both LMJ-040682::pAS_C2 and LMJ-

040682::pAS_C3 were found to take up B_{12} to a similar extent as their parental strain cw15 when

grown in the absence of thiamine (Figure 4b). However, when 10 μM thiamine was included in the

culture medium, LMJ-040682::pAS_C3 showed virtually no B_{12} uptake. This riboswitch-mediated

conditional complementation of the phenotype in LMJ-040682::pAS_C3 demonstrated conclusively

that B_{12} uptake in *C. reinhardtii* is dependent on the presence of CrCBA1.

CrCBA1 shows an association with membranes and is highly upregulated under B_{12}-deprivation

To investigate the subcellular location of CrCBA1, we used several bioinformatic targeting prediction

tools. CrCBA1 is annotated as a flagella-associated protein in the Phytozome v5.6 *C. reinhardtii*

annotation. However, both DeepLoc (Almagro Armenteros et al., 2017) and SignalP (Almagro

Armenteros et al., 2019) indicated a hydrophobic sequence with the characteristics of a signal peptide

at the N-terminus of CrCBA1 and predicted it would be targeted to the endoplasmic reticulum (ER).

Additionally, it was predicted to contain a transmembrane helix at its C-terminus by InterPro

(Mitchell et al., 2019).

We next investigated the subcellular location of CrCBA1 *in vivo* by imaging two lines of LMJ-

040682::pAS_C2, where the CBA1 is tagged with mVenus, with confocal microscopy. No mVenus

was detected in the parental LMJ-040682 cells, whereas a clear fluorescent signal was observed in

LMJ-040682::pAS_C2 #A10 and LMJ-040682::pAS_C2 #D10 (Figure 5). In these complemented

lines, the mVenus signal was absent from the chloroplast, nucleus and flagella, but instead could be

seen within the cell localising both to the plasma membrane and to regions that may be

endomembranes such as the ER. This is consistent with findings from *P. tricornutum* showing a

similar distribution (Bertrand et al., 2012). Together these data indicate that CBA1 is likely to be

associated with membranes, and therefore, may have a conserved role in the B_{12} uptake process.

Further evidence for the role of CBA1 in B_{12} uptake was obtained by taking advantage of a B_{12}-

dependent mutant of *C. reinhardtii*, metE7 (Helliwell et al., 2015; Bunbury et al., 2020). We tested

the effect of B_{12}-deprivation over time on the expression of the *CrCBA1* gene by RT-qPCR in the

mutant and determined the rate of B_{12} uptake over a similar period. Within 6h of B_{12} removal, there

was a ~250-fold induction of the *CrCBA1* transcript, followed by a slow decline over the next 60h

(Figure 6a). After resupply of B_{12} there was then a rapid ~100-fold decline within 8h. The B_{12} uptake

capacity of metE7 followed a similar profile, increasing 3-fold over the first 12 hours of B_{12} depletion,

from ~6.5 x 10^5 molecules B_{12}/cell/hour to 1.86 x 10^6 molecules B_{12}/cell/hour (Figure 6b), then

declining slowly. This induction profile is characteristic of a nutrient-starvation response shown by

many transporters, including in *C. reinhardtii* those for Fe (Allen et al., 2007), and for *CBA1* in the

B_{12}-dependent diatom, *Thalassiosira pseudonana* (Bertrand et al., 2012).

333   Widespread distribution of CBA1 in algae

Having shown the importance of *PtCBA1* and *CrCBA1* for B_{12} uptake in their respective species, we

re-examined how prevalent CBA1-like proteins are in Nature. Searches with BLASTP using *PtCBA1*

resulted in no significant homologues in species outside the Stramenopiles (Bertrand et al., 2012).

Instead, we created a hidden Markov model (HMM), using the *C. reinhardtii* CBA1 amino acid

sequence and CBA1 sequences from *P. tricornutum*, *T. pseudonana*, *Fragilariopsis cylindrus*,

Aureococcus anophagefferens and *Ectocarpus siliculosus* (Bertrand et al., 2012), to identify more

accurately CBA1-like proteins in other organisms. The EukProt database of curated eukaryotic

genomes (Richter et al. 2022) includes representatives from the Archaeplastida (designated by

EukProt as Chloroplastida), which encompass green algae, red algae, glaucophytes and all land plants,

as well as phyla that include algae with complex plastids, namely Stramenopiles (which include

diatoms), Alveolata (including dinoflagellates), Rhizaria and Haptophyta, and the animals (both

Metazoa and basal Choanoflagellates), the fungi and Amoebozoa. This database was queried with the

CBA1 HMM model, using a cutoff e-value of 1e-20, and 277 hits were obtained (Figure S7;

Supplementary Table S3). No candidates were found in the Metazoa, but CBA1 homologues were

identified in all other phyla, including all photosynthetic groups, fungi and amoebozoa and in

choanoflagellates, unicellular and colonial flagellated organisms considered to be the closest living

relatives of animals (King et al., 2008).

Given that higher plants have no B_{12}-dependent enzymes, the presence of a putative B_{12}-binding

protein in several angiosperms, both monocot and dicot, and the gymnosperm *Ginkgo biloba*, was

somewhat surprising. To address this conundrum, we investigated to what extent CBA1 was

associated with vitamin B_{12} dependence by determining the distribution of the different isoforms of

methionine synthase, METH and METE. Using the same HMM approach as before, the protein

sequences were searched against the EukProt database and the combination of presence and absence

of CBA1, METH and METE across eukaryotic species groups was compiled (Figure 7;

Supplementary Table S4). What is immediately apparent is that the combination of the three proteins

is quite different in the various lineages. In the major algal groups, the Chlorophyta and the SAR

clade (Stramenopiles, Alveolata and Rhizaria), METH sequences were found in the majority of

genomes analysed and their presence was correlated with CBA1. In the genomes of the Chlorophyta

and the SAR clade that encoded METE only (7 taxa in total), CBA1 was absent in all but one, the

diatom *Thalassionema nitzschiodes*. Equal numbers of Alveolata species encoded METH and CBA1,

or METH only; interestingly, the latter were all non-photosynthetic lineages. Grouping the data from

these 4 algal groups, a Chi Square test was significant for CBA1 and METH being more often both

present or both absent *($X2$ (1, $N = 86$) = 9.2, $p = 0.00240$)*. The association could be due to linkage,

although in neither *C. reinhardtii* nor *P. tricornutum* are the two genes on the same chromosome,

making this unlikely. Alternatively, there is a fitness advantage in both genes being acquired or lost

together.

Most fungal taxa lacked both METH and CBA1, but we found examples of 6 species that were

predicted to be B_{12} users (METH present) and 5 of these were also predicted to contain CBA1-like

sequences: *Allomyces macrogynus*, *Spizellomyces punctatus*, *Rhizophagus irregularis*, *Rhizopus*

delemar and *Phycomyces blakesleeanus*. CBA1-like sequences were identified in the Opisthokonta

and Amoebozoa, although were less prevalent, with ~23% of choanoflagellates and 8% of amoeboid

species being like algae in having both METH and CBA1. CBA1 was entirely absent from the

Metazoa. In contrast, in the Streptophyta, which include multicellular green algae and all land plants,

the majority lack METH, but almost 80% of species were found to contain CBA1-like sequences.

This implies that Streptophyta CBA1 sequences may have gained a different function, which would

be consistent with the lack of B_{12}-dependent metabolism in these organisms. In summary, these data

suggest that CBA1 is associated with vitamin B_{12} use to different degrees in different eukaryotic

groups, with there being a greater association in obligate and facultative B_{12} users than in those

organisms that do not utilise B_{12}.

The many putative CBA1 homologues in algal lineages and their strong association with B_{12} uptake

provided an opportunity to identify conserved, and thus likely functionally important, residues.

Accordingly, a multiple sequence alignment of proteins matching the CBA1 HMM query was

generated (Figure S7). Highlighted in green in the similarity matrix at the top are nine conserved

regions with several almost completely conserved residues; these are shown in more detail in Figure

8a for selected taxa representing different algal groups. Further insight came from inspection of the

model of the 3D structure of CrCBA1 generated by the Phyre2 structural prediction server. The

analysis showed that regions of CrCBA1 showed similarity to bacterial periplasmic binding proteins,

including the B_{12}-binding protein BtuF. A structure is available of *E. coli* BtuF in complex with B_{12}

(Borths et al., 2002), so we compared this to the modelled CrCBA1 structure. Although there is little

sequence similarity, alignment of the two structures resulted in an RMSD of 3.362 and enabled the

relative position of B_{12} to be placed in the lower cleft of CrCBA1, shown in red in Figure 8b.

Mapping of the highly conserved residues onto this structure found that many (P251, V253, W255,

W394, F395 and E396) were in a cluster around the relative position of B_{12}. Another cluster of highly

conserved residues were located at the end of the upper alpha helix (P118, L136, F214, F215, N216

and E218). Both clusters represent promising mutational targets to investigate CrCBA1 function.

DISCUSSION

In this study we have shown experimentally that a conserved protein, CBA1, is required for the

uptake of the micronutrient B_{12} in two taxonomically distant algae, the diatom *P. tricornutum* (Figure

1) and the chlorophyte *C. reinhardtii* (Figures 3 & 4). Strains with knockouts of the gene were unable

to take up $B_{12,}$ demonstrating that there is no functional redundancy of this protein in either organism.

This is also the first *in vivo* evidence that CBA1 is present outside the Stramenopiles. Moreover, we

found widespread occurrence of CBA1 homologues with considerable sequence conservation across

eukaryotic lineages (Figures 7 and S7). The strong correlation of CBA1 with the B_{12}-dependent

methionine synthase, METH, in algal lineages, provides evidence that CBA1 is a key component of

the B_{12} uptake process in evolutionarily distinct microalgae, and the structural similarities between

CBA1 and BtuF (Figure 8b), suggest it may operate as a B_{12}-binding protein. The highly conserved

residues identified in the algal homologues (Figure 8a) offer the means to establish which are

functionally important, facilitated by the uptake assay we established.

Nonetheless, the mechanistic role of CBA1 in the process of B_{12} acquisition in algae is not yet clear.

Previous physiological studies of B_{12} uptake by microalgae, such as the haptophyte *Diacronema*

lutheri (Droop, 1968), indicated a biphasic process: firstly rapid irreversible adsorption of B_{12} to the

cell exterior, followed by a slower second step of B_{12} uptake into the cell, consistent with endocytosis.

CBA1 is unlikely to be associated with the binding of B_{12} in the cell wall, however. This is because

the *C. reinhardtii* strains used in this study, UVM4 and CW15, were cell wall deficient, and therefore

likely also deficient in cell wall proteins that bind B_{12}; the lack of a B_{12}-BODIPY signal from the cell

surface in IM4 (Figure S2) supports this hypothesis. Further use of this fluorescent probe offers the

possibility to monitor the localisation of B_{12}-BODIPIY over time to gain insights into the stages of B_{12}

uptake, as has been done in other organisms (Lawrence et al., 2018). In addition, confocal microscopy

of CBA1-mVenus fusion protein in *C. reinhardtii* (Figure 5) showed an apparent association of

CrCBA1 with the plasma membrane and endomembranes, which is similar to that for ER-localised

proteins (Mackinder et al., 2017). Moreover, in a proteomics study of lipid droplets (which form by

budding from the ER) CBA1 was in the top 20 most abundant proteins (Goold et al., 2016). Bertrand

et al. (2012) found that PtCBA1 had a signal peptide and fluorescently tagged PtCBA1 was also

targeted to the ER. Nonetheless, based on its predicted 3D structure and the fact that it has at most one

transmembrane helix, CBA1 does not appear to be a transporter itself. Instead, given its structural

similarity to BtuF, a distinct possibility is that CBA1 is the soluble component of an ABC transporter,

either at the plasma membrane or an internal membrane, and likely will interact with one or more

other proteins to allow B_{12} uptake to occur, at least some of them being those involved in receptor-

mediated endocytosis, as is the case for B_{12} acquisition in humans (Rutsch et al., 2009; Beedholm-

Ebsen et al., 2010; Coelho et al., 2012). In this context, there are known similarities between

endocytosis in *C. reinhardtii* and humans (Denning and Fulton, 1989; Bykov et al., 2017), and several

putative homologues have been identified by sequence similarity in the alga. Testing the B_{12}-uptake

capacity of mutants of these proteins would be one approach to investigate whether their roles are also

conserved.

In contrast to the situation in algae, the Streptophyta live in a B_{12}-free world, neither synthesising nor

utilising this cofactor. This is exemplified by the fact that in our analysis only one species, the

charophyte alga *Cylindrocystis brebissonii*, encoded METH. Despite this, more than three-quarters of

this group encode a CBA1 homologue (Figures 7 & S7). Since the majority of the conserved residues

(Figure 8a) are also found in putative CBA1 sequences in the angiosperms such as *Arabidopsis*,

including those around the potential binding pocket, it is possible that the streptophyte protein has

acquired a new function that still binds a tetrapyrrole molecule. Intriguingly, the reverse is observed in

the Metazoa, where METH is almost universal, but CBA1 is entirely absent. However, some

Choanoflagellates and some species of fungi do appear to encode both METH and CBA1, suggesting

that they utilise B_{12}, a trait only recently recognised to occur in fungi (Orłowska et al., 2021). It will

be of interest therefore to test whether CBA1 is involved in B_{12} uptake in these organisms, for

example by gene knockout studies.

The importance of B_{12} availability for phytoplankton productivity has been demonstrated across

several marine ecosystems by amendment experiments (e.g. Bertrand et al., 2011; Koch et al., 2012;

Joglar et al., 2021), where addition of B_{12} led to algal blooms and affected the composition and

stability of microbial communities. The mode of acquisition of this micronutrient is thus likely to be

highly conserved and subject to significant ecological and evolutionary selection pressure to be

retained. Moreover, the role of B_{12} at the cellular level may well provide a direct connection between

environmental conditions and the epigenetic status of the genome: methionine synthase is the key

enzyme in C1 metabolism, linking the folate and methylation cycles and thus responsible for

maintaining levels of S-adenosylmethionine (SAM) the universal methyl donor (Hanson & Roje

2001; Mentch & Locasale, 2016). In this context, it is noteworthy that the knockout of *CBA1* in the

IM4 line was the result of insertion of a class II transposable element into the gene. This mobilization

is likely to reflect epigenetic alterations of the autonomous element, presumably as a result of cellular

stress either from the antibiotic selection, or the transformation procedure, or both. Recent

classification of the transposons in *C. reinhardtii* indicate that the transposon inserted into *CBA1* in

IM4 is a member of the KDZ superfamily of class II TIR elements named Kyakuja-3_cRei (Craig et

al. 2021). If the phenomenon of inactivation of a gene that is deleterious (in this case allowing B_{12} to

be taken up and repress the antibiotic resistance gene) via transposition is a general response in *C.*

reinhardtii, repeating the screen for CBA1 mutants might allow observation of further transposition

events, and enable characterisation of this group of elements at the functional level. Moreover, it

could be adopted as a more general methodology to identify candidate genes involved in other

physiological processes, by tying their expected effects to deleterious outcomes through synthetic

biology constructs and screening surviving mutants by sequencing.

MATERIALS AND METHODS
483
Organisms and growth conditions
Strains, media and growth conditions used in this study are listed in Table S1. If required, antibiotics,
vitamin B_{12} (cyanocobalamin) and thiamine were added to the medium at concentrations indicated.
Algal culture density was measured using a Z2 particle count analyser (Beckman Coulter Ltd.) and
optical density (OD) at 730 nm was measured using a FluoStar OPTIMA (BMG labtech) plate reader
or a CLARIOstar plate reader (BMG labtech). Bacterial growth was recorded by measuring OD_{595}.
490
Algal B_{12}-uptake assay
Algal cultures were grown to stationary phase and cyanocobalamin salt (Sigma) was added (*P.*
tricornutum: 600 pg; *C. reinhardtii*: 150 pg) to 5×10^6 cells in a final volume of 1 ml in f/2 or TAP
medium respectively. The samples were incubated at 25°C under continuous light with shaking for 1
hour and inverted every 30 minutes to aid mixing. Samples were centrifuged and the supernatant
(media fraction) transferred into a fresh microcentrifuge tube. The cell pellet was resuspended in 1 ml
water. Both samples were boiled for 10-20 minutes to release any cellular or bound B_{12} into solution,
and then centrifuged to pellet debris. The supernatant was used in the *S. typhimurium* B_{12} bioassay as
described in Bunbury et al. (2020). The amount of B_{12} in the sample was calculated by comparison to
a standard curve of known B_{12} concentrations fitted to a 4 parameter logistic equation $f(x) = c + (d-$
$c)(1+\exp(b(\log(x)-\log(e))))$ (Ritz et al., 2015). This standard curve was regenerated with every
bioassay experiment.
503
Generating *P. tricornutum* CBA1 knockout lines using CRISPR-Cas9
CRISPR/Cas9 genome editing applied the single guide RNA (sgRNA) design strategy described in
Hopes et al., (2017). Details are provided in the Supplementary methods. *P. tricornutum CCAP*
1055/1 cells were co-transformed with linearised plasmids pMLP2117 and pMLP2127 using a
NEPA21 Type II electroporator (Nepa Gene) as previously described (Yu et al., 2021). After plating
on 1% agar selection plates containing 75 mg·l^{-1} zeocin and incubation for 2-3 weeks, zeocin resistant
colonies were picked into 96 well plates containing 200 µl of f/2 media with 75 mg·l^{-1} zeocin. After
seven days strains were subcultured into fresh media either containing 75 mg·l^{-1} zeocin or 300 mg·l^{-1}
nourseothricin, and genotyped with a three-primer PCR using PHIRE polymerase (Thermo Fisher

Scientific) with primers gCBA1.fwd, gCBA1.rv and NAT.rv (Table S2). Five promising colonies
resistant to nourseothricin and with genotypes showing homologous recombination or indels were re-
streaked on 75 mg·l^{-1} zeocin f/2 plates to obtain secondary monoclonal colonies. Twelve secondary
colonies were picked for each primary colony after 2-3 weeks and again genotyped with a three-
primer PCR. Promising colonies were genotyped in further detail with primer pairs
gCBA1.fwd/gCBA1.rv, gCBA1.fwd/NAT.rv and gCBA1in.fwd/gCBA1in.rv (Table S2).
519

Construct assembly and *C. reinhardtii* transformation
Constructs were generated using Golden Gate cloning, using parts from the *Chlamydomonas* MoClo
toolkit (Crozet et al., 2018) and some that were created in this work. All parts relating to
Cre02.g081050 were domesticated from UVM4 genomic DNA, with BpiI and BsaI sites removed
from the promoter, ORF and terminator by PCR based mutagenesis. A list of plasmids used in this
study is shown in Table S2. Transformation of *C. reinhardtii* cultures with linearised DNA was
carried out by electroporation essentially as described by Mehrshahi et al. (2020) before plating on
TAP-agar plates with the appropriate antibiotics.
528

Insertional mutagenesis was performed as above, however, cultures were grown to a density of
approximately 1×10^7 cells/ml and were incubated with 500 ng transgene cassette. After allowing the
cells to recover overnight in TAP plus 60 mM sucrose at 25°C in low light (less than 10 μmol photon
$m^{-2}.s^{-1}$ at 100 rpm), between 200 - 250 μl of transformants were plated on solid TAP media (square
12x12 cm petri dishes) containing ranges of 15-20 μg/ml hygromycin, 20-50 μg/ml paromomycin and
48-1024 ng/l vitamin B12, and the plates were incubated in standing incubators.
535

Confocal laser scanning microscopy
C. reinhardtii transformants carrying the pAS_C2 construct were imaged in a confocal laser scanning
microscope (TCS SP8, Leica Microsystems, Germany) with an HC PL APO CS2 40x/1.30 aperture
oil-immersion lens. Images were taken using the sequential mode provided by the Leica LAS
software, with the channel used for mVenus and brightfield detection being taken first with excitation
from a white light source at 486 nm and emissions were detected between 520 - 567 nm, followed by
chlorophyll detection (excitation 514 nm, emission 687-724 nm). The overlay images were produced
automatically by the Leica LAS software. Inkscape was used to increase the lightness and decrease
the contrast of all the images in the same manner.
545

546 **Quantitative real-time PCR**
547 Quantification of steady state levels of transcripts was carried out according to Bunbury et al. (2020),
548 using random hexamer primers for cDNA synthesis. The qPCR data was analysed using the ΔΔCT
549 method with an assumed amplification efficiency of 2. Log2(2-ΔCT) values were plotted in the
550 resulting figures.
551

Whole genome sequencing
Genomic DNA was extracted from *C. reinhardtii* cells by phenol-chloroform extraction and
sequenced using the NovaSeq sequencing platform by Novogene (Cambridge, UK) to produce 150 bp
paired-end reads. This involved RNase treatment and library preparation with the NEBNext Ultra II
DNA Library Prep Kit (PCR-free), which generated 350 bp inserts. The raw sequencing data for this
study have been deposited in the European Nucleotide Archive (ENA) at EMBL-EBI under accession
number PRJEB58730 (https://www.ebi.ac.uk/ena/browser/view/PRJEB58730). Novogene performed
all quality filtering, summary statistics and bioinformatic analysis. The location of the Hyg3 cassette
was determined by identifying loci that comprised reads from IM4 that mapped between genomic
DNA and pHyg3, and cross-referencing these loci against the parental strains. The TE identification
was carried out similarly, full details are provided in Supplementary Methods.
563

Bioinformatics pipeline
The EukProt database was assessed for the presence of METE, METH and CBA1 (Richter et al.,
2022). The query used for CBA1 was a hidden Markov model (HMM) generated from the protein
fasta sequences: Phatr3_J48322, Thaps3 11697, Fracy1 241429, Fracy1 246327, Auran1 63075,

Ectocarpus siliculosus D8LMT1 and Cre02.g081050.t1.2 by first aligning using MAFFT (Katoh and
Standley, 2013) version 7.470 with the --auto option, and then building a HMM using hmmbuild
(hmmer 3.2.1). Additionally, protein fasta (Cre06.g250902, Cre03.g180750), PFAM (PF02310,
PF02965, PF00809, PF02574, PF01717, PF08267) and KO (K00548, K00549) queries were searched
against EukProt to identify sequences with similarity to METE and METH. The queries were
searched against EukProt using hmmsearch (HMMER 3.1b2). The default bitscore thresholds were
used for KO and PFAM queries. The threshold used for CBA1 HMM, and the CrMETE and CrMETH
protein fasta sequences, was a full-length e-value of 1e-20. For each protein, all individual queries
were required to be significant to classify the protein as present. The best hit in each species was
identified by taking the protein with the greatest geometric mean of full length bitscores for the
queries. The dataset was joined with taxonomic information from EukProt and completeness
information calculated using BUSCO version 4.1.4 and eukaryote_odb10 (Manni et al., 2021).
580
581

Acknowledgements

We thank Catherine Sutherland for help with maintaining and screening the CRISPR/Cas9 mutants of *P.*

tricornutum and Dr Lorraine Archer for lab management. We are grateful to Dr Amanda Hopes and Prof

Thomas Mock (University of East Anglia) for the parts used in gene editing of *PtCBA1*. The plasmid Hyg3

used in the insertional mutagenesis was obtained from the Chlamydomonas Resource Centre

(www.chlamycollection.org). This work was supported by the UK's Biotechnology and Biological Sciences

Research Council (BBSRC) Doctoral Training Partnership, grant no. BB/M011194/1 to A.P.S., M.L.P. and

A.G.S; grant no. BB/M018180/ 1 to P.M. and A.G.S.; grant no. BB/L002957/1 and BB/R021694/1 to K.G.

and AGS; grant no. BB/L014130/1 to G.I.M, K.G., P.M. and A.G.S; grant no. BB/S002197/1 to M.J.W;

University of Cambridge Broodbank Fellowship to G.I.M; Royal Society grant no. INF\R2\180062 to

M.J.W.; and Bill & Melinda Gates Foundation grant OPP1144 and Gates Cambridge Trust (Graduate

Student Fellowship) to A.H. For the purpose of open access, the authors have applied a Creative Commons

Attribution (CC BY) licence to any Author Accepted Manuscript version arising from this submission.

Author contributions

APS designed and performed research, analysed data and wrote the article with contributions from all

the authors. KG and MLP carried out the CRISPR/Cas9 editing of Phaeodactylum and contributed to

writing the article. AH carried out the bioinformatics analysis to identify the putative transposable

elements. MJW & ADL synthesised the BODIPY-labelled B$_{12}$ and contributed to writing the article.

KG, GMO and PM supervised aspects of the project and contributed to writing the article. AGS

conceived the project, obtained the funding, supervised the project and wrote the article with

contributions from all the authors. AGS agrees to serve as the author responsible for contact and

ensures communication.

References

Allen, MD, del Campo JA, Kropat J, Merchant SS (2007) FEA1, FEA2, and FRE1, encoding
two homologous secreted proteins and a candidate ferrireductase, are expressed coordinately with
FOX1 and FTR1 in iron-deficient *Chlamydomonas reinhardtii*. Eukaryotic Cell **6:** 1841–1852

Almagro Armenteros J, Sønderby CK, Sønderby SK, Nielsen H, Winther O (2017) DeepLoc:
Prediction of protein subcellular localization using deep learning. Bioinformatics **33**: 3387–3395

Almagro Armenteros J, Tsirigos KD, Sønderby CK, Petersen TN, Winther O, Brunak S,
Heijne G von, Nielsen H (2019) SignalP 5.0 improves signal peptide predictions using deep
neural networks. Nature Biotechnology **37**: 420–423

Banerjee R, Gouda H, Pillay S (2021) Redox-linked coordination chemistry directs vitamin B_{12}
trafficking. Accounts of Chemical Research **54**: 2003–2013

Beedholm-Ebsen R, Wetering KVD, Hardlei T, Nexø E, Borst P, Søren K, Moestrup SK
(2010) Identification of multidrug resistance protein 1 (MRP1/ABCC1) as a molecular gate for
cellular export of cobalamin. Blood **115**: 1632–1639

Bertrand EM, Saito MA, Rose JM, Riesselman CR, Lohan MC, Noble AE, Lee PA, DiTullio
GR (2011) Vitamin B_{12} and iron colimitation of phytoplankton growth in the Ross Sea. Limonol
Oceanogr **52**: 1079-1093

Bertrand EM, Allen AE, Dupont CL, Norden-Krichmar TM, Bai J, Valas RE, Saito MA
(2012) Influence of cobalamin scarcity on diatom molecular physiology and identification of a
cobalamin acquisition protein. Proc Natl Acad Sci U S A **109**: E1762–71

Borths EL, Locher KP, Lee AT, Rees DC (2002) The structure of *Escherichia coli* BtuF and
binding to its cognate ATP binding cassette transporter. Proc Natl Acad Sci U S A **99**: 16642–7

Bunbury F, Helliwell KE, Mehrshahi P, Davey MP, Salmon DL, Holzer A, Smirnoff N,
Smith AG (2020) Responses of a newly evolved auxotroph of Chlamydomonas to B_{12}
deprivation. Plant Physiology **183**: 167–178

Bykov YS, Schaffer M, Dodonova SO, Albert S, Plitzko JM, Baumeister W, Engel BD,
Briggs JA (2017) The structure of the COPI coat determined within the cell. eLife **6**: e32493

Carlucci FA, Silbernagel BS, McNally MP (2007) The influence of temperature and solar
radiation on persistence of vitamin B12, thiamine, and biotin in seawater. Journal of Phycology **5**:
302–305

Choi CC, Ford RC (2021) ATP binding cassette importers in eukaryotic organisms. Biological
Reviews **96**: 1318–1330

Coelho D, Kim JC, Miousse IR, Fung S, Moulin M du, Buers I, Suormala T, Burda P,
Frapolli M, Stucki M, et al (2012) Mutations in ABCD4 cause a new inborn error of vitamin
B12 metabolism. Nature Genetics **44**: 1152–1155

Craig RJ, Hasan AR, Ness RW, Keightley PD (2021) Comparative genomics of
Chlamydomonas. Plant Cell **33**:1016-1041

Croft MT, Lawrence AD, Raux-Deery E, Warren MJ, Smith AG (2005) Algae acquire
vitamin B_{12} through a symbiotic relationship with bacteria. Nature **438**: 90–3

Crozet P, Navarro FJ, Willmund F, Mehrshahi P, Bakowski K, Lauersen KJ, Pérez-Pérez
M-E, Auroy P, Gorchs Rovira A, Sauret-Gueto S, et al (2018) Birth of a photosynthetic
chassis: A MoClo toolkit enabling synthetic biology in the microalga *Chlamydomonas*
reinhardtii. ACS Synthetic Biology **7**: 2074–2086

Denning GM, Fulton AB (1989) Purification and characterization of clathrin-coated vesicles
from *Chlamydomonas*. The Journal of Protozoology **36**: 334–340

Droop MR (1968) Vitamin B12 and marine ecology. IV. The kinetics of uptake, growth and
inhibition in *Monochrysis lutheri*. Journal of the Marine Biological Association of the United
Kingdom **48**: 689–733

Field CB, Behrenfeld MJ, Randerson JT, Falkowski P (1998) Primary production of the
biosphere: Integrating terrestrial and oceanic components. Science **281**: 237–240

Gonzalez JC, Banerjee RV, Huang S, Sumner JS, Matthews RG (1992) Comparison of
cobalamin-independent and cobalamin-dependent methionine synthases from *Escherichia coli*:
Two solutions to the same chemical problem. Biochemistry **31**: 6045–6056

Goold HD, Cuiné S, Légeret B, Liang Y, Brugière S, Auroy P, Javot H, Tardif M, Jones B,
Beisson F, et al (2016) Saturating light induces sustained accumulation of oil in plastidal lipid
droplets in *Chlamydomonas reinhardtii*. Plant Physiology **171**: 2406–2417

Hanson AD, Roje S (2001) One-Carbon metabolism in higher plants. Ann Rev Plant Physiol **52**:
119-137

Helliwell KE, Collins S, Kazamia E, Purton S, Wheeler GL, Smith AG (2015) Fundamental
shift in vitamin B_{12} eco-physiology of a model alga demonstrated by experimental evolution.
ISME J **9**: 1446–1455

Helliwell KE, Scaife MA, Sasso S, Araujo APU, Purton S, Smith AG (2014) Unraveling
vitamin B_{12}-responsive gene regulation in algae. Plant Physiol **165**: 388–397

Helliwell KE, Wheeler GL, Leptos KC, Goldstein RE, Smith AG (2011) Insights into the
evolution of vitamin B_{12} auxotrophy from sequenced algal genomes. Molecular Biology and
Evolution **28**: 2921–2933

Hopes A, Nekrasov V, Belshaw N, Grouneva I, Kamoun S, Mock T (2017) Genome editing in
diatoms using CRISPR-cas to induce precise bi-allelic deletions. Bio Protoc **7**: e2625

Joglar V, Pontiller B, Martínez-García S, Fuentes-Lema A, Pérez-Lorenzo M, Lundin D,
Pinhassi J Fernández E, Teira E (2021) Microbial plankton community structure and function
responses to vitamin B_{12} and B_1 amendments in an upwelling system. Appl Environ Microbiol **87**:
e0152521.

Kadner RJ (1990) Vitamin B_{12} transport in *Escherichia coli*: Energy coupling between
membranes. Molecular Microbiology **4**: 2027–2033

Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7:
Improvements in performance and usability. Molecular Biology and Evolution **30**: 772–780

Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJE (2015) The Phyre2 web portal for
protein modeling, prediction and analysis. Nat Protoc **10**: 845–58

King N, Westbrook MJ, Young SL, Kuo A, Abedin M, Chapman J, et al. (2008). The
genome of the choanoflagellate *Monosiga brevicollis* and the origin of metazoans. *Nature*. **451**:
783–788.

Koch F, Hattenrath-Lehmann TK, Goleski JA, Sañudo-Wilhelmy S, Fisher NS, Gobler CJ.
2012. Vitamin B_1 and B_{12} uptake and cycling by plankton communities in coastal
ecosystems. *Front Microbiol* 3:363.

Lawrence AD, Nemoto-Smith E, Deery E, Baker JA, Schroeder S, Brown DG, Tullet JMA,
Howard MJ, Brown IR, Smith AG, et al (2018) Construction of fluorescent analogs to follow
the uptake and distribution of cobalamin (vitamin B_{12}) in bacteria, worms, and plants. Cell Chem
Biol **25**: 941–951

Li X, Zhang R, Patena W, Gang SS, Blum SR, Ivanova N, Yue R, Robertson JM, Lefebvre
PA, Fitz-Gibbon ST, et al (2016) An indexed, mapped mutant library enables reverse genetics
studies of biological processes in *Chlamydomonas reinhardtii*. Plant Cell **28**: 367–87

Mackinder LCM, Chen C, Leib RD, Patena W, Blum SR, Rodman M, Ramundo S, Adams
CM, Jonikas MC (2017) A spatial interactome reveals the protein organization of the algal CO_2
concentrating mechanism. Cell **171**: 133–147.e14

Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM (2021) BUSCO update: Novel
and streamlined workflows along with broader and deeper phylogenetic coverage for scoring of
eukaryotic, prokaryotic, and viral genomes. Molecular Biology and Evolution **38**: 4647–4654

Mehrshahi P, Nguyen GTDT, Gorchs Rovira A, Sayer A, Llavero-Pasquina M, Lim Huei
Sin M, Medcalf EJ, Mendoza-Ochoa GI, Scaife MA, Smith AG (2020) Development of novel
riboswitches for synthetic biology in the green alga *Chlamydomonas*. ACS Synthetic Biology **9**:
1406–1417

Mentch SJ, Locasale JW (2016) One-carbon metabolism and epigenetics: understanding the
specificity Annal NY Acad Sci **1363**: 91-8

Mitchell AL, Attwood TK, Babbitt PC, Blum M, Bork P, Bridge A, Brown SD, Chang H-Y,
El-Gebali S, Fraser MI et al. (2019) InterPro in 2019: improving coverage, classification and
access to protein sequence annotations. Nucl Acids Res **47:** D351–D360

Neupert J, Karcher D, Bock R (2009) Generation of *Chlamydomonas* strains that efficiently
express nuclear transgenes. The Plant Journal **57**: 1140–1150

Nielsen MJ, Rasmussen MR, Andersen CBF, Nexø E, Moestrup SK (2012) Vitamin B_{12}
transport from food to the body's cells—a sophisticated, multistep pathway. Nature Reviews
Gastroenterology & Hepatology **9**: 345–354

Ohwada K (1973) Seasonal cycles of vitamin B_{12}, thiamine and biotin in Lake Sagami. Patterns
of their distribution and ecological significance. Internationale Revue der gesamten
Hydrobiologie und Hydrographie **58**: 851–871

Orłowska M, Steczkiewicz K, Muszewska A (2021) Utilization of cobalamin is ubiquitous in
early-branching fungal phyla. Genome Biology and Evolution **13**: evab043

Panzeca C, Beck AJ, Tovar-Sanchez A, Segovia-Zavala J, Taylor GT, Gobler CJ, Sañudo-
Wilhelmy SA (2009) Distributions of dissolved vitamin B_{12} and Co in coastal and open-ocean
environments. Estuarine, Coastal and Shelf Science **85**: 223–230

Pintner IJ, Altmeyer VL (1979) Vitamin B_{12}-binder and other algal inhibitors Journal of
Phycology **15**: 391–398

Richter DJ, Berney C, Strassert JFH, Poh Y-P, Herman EK, Muñoz-Gómez SA, Wideman
JG, Burki F, Vargas C de (2022) EukProt: A database of genome-scale predicted proteins across
the diversity of eukaryotes. Peer Community Journal **2**: e56

Ritz C, Baty F, Streibig JC, Gerhard D (2015) Dose-response analysis using R. PLoS One **10**:
e0146021

Rutsch F, Gailus S, Miousse IR, Suormala T, Sagné C, Toliat MR, Nürnberg G, Wittkampf
T, Buers I, Sharifi A, et al (2009) Identification of a putative lysosomal cobalamin exporter
altered in the cblF defect of vitamin B_{12} metabolism. Nature Genetics **41**: 234–239

Sahni MK, Spanos S, Wahrman MZ, Sharma GM (2001) Marine corrinoid-binding proteins
for the direct determination of vitamin B_{12} by radioassay. Analytical Biochemistry **289**: 68–76

Sañudo-Wilhelmy SA, Gómez-Consarnau L, Suffridge C, Webb EA (2014) The role of B
vitamins in marine biogeochemistry. Annual Review of Marine Science **6**: 339–367

Shelton AN, Seth EC, Mok KC, Han AW, Jackson SN, Haft DR, Taga ME (2019) Uneven
distribution of cobamide biosynthesis and dependence in bacteria predicted by comparative
genomics. The ISME Journal **13**: 789–804

Tang YZ, Koch F, Gobler CJ (2010) Most harmful algal bloom species are vitamin B_1 and B_{12}
auxotrophs. Proc Natl Acad Sci U S A **107**: 20756–61

Warren MJ, Raux E, Schubert HL, Escalante-Semerena JC (2002) The biosynthesis of
adenosylcobalamin (vitamin B_{12}). Natural product reports **19**: 390–412

Xie B, Bishop S, Stessman D, Wright D, Spalding MH, Halverson LJ (2013) *Chlamydomonas*
reinhardtii thermal tolerance enhancement mediated by a mutualistic interaction with vitamin B_{12}-
producing bacteria. ISME J **7**: 1544–55

Yu Z, Geisler K, Leontidou T, Young REB, Vonlanthen SE, Purton S, Abell C, Smith AG
(2021) Droplet-based microfluidic screening and sorting of microalgal populations for strain
engineering applications. Algal Res **56**: 102293

754
755
756
757

 Figure Legends
759
Figure 1. Disruption of *Phaeodactylum tricornutum CBA1* (*PtCBA1*) using CRISPR-
Cas9 yielded lines with impaired B$_{12}$ uptake. a) Schematic showing CRISPR-Cas9 sgRNA
target sites and the homology repair template design used to generate mutant lines in *PtCBA1*
(Phatr3_J48322). The homology repair template schematic is annotated with the 5' homology
region (HR) and 3'HR, the *FCPB* promoter, nourseothricin resistance gene (*NAT*) and *FCPC*
terminator. The *PtCBA1* gene is annotated with the ORF, the 5'HR and 3'HR regions used in
the homology template and the regions of the ORF targeted by sgRNA (vertical bars). Primer
positions used for the analysis of putative mutant lines are shown with arrowheads. **b)** PCR of
regions across and within wild-type (WT) and mutant *PtCBA1* in 3 independent CRISPR-
Cas9 lines (ΔCBA1) showing indel mutations in the mutants. PCR products from different
sets of primers indicated in panel a are shown. M = marker, - Ctrl = no DNA template. **c)** A
B$_{12}$ uptake assay was performed as described in Materials and Methods, to determine the
amount of B$_{12}$ in the media and the cells after 1h incubation of *P. tricornutum* cells in 600 pg
B$_{12}$. The 'Total' was inferred by the addition of the cell and media fractions. The dashed line
indicates the amount of B$_{12}$ added to the experiment. Standard deviation error bars are shown,
n=4. Statistical analysis was performed on the media fraction, and Tukey's test identified the
following comparisons to be significantly different from one another: WT vs No Algae
($p<1e^{-12}$); WT vs ΔCBA1-1 ($p<1e^{-10}$); WT vs ΔCBA1-2 ($p<1e^{-12}$); WT vs ΔCBA1-3 ($p<1e^{-11}$
); No Algae vs ΔCBA1-1 ($p<1e^{-03}$); No Algae vs ΔCBA1-3 ($p<0.05$); and ΔCBA1-1 vs
ΔCBA1-2 ($p<1e^{-02}$).
780
781
Figure 2. Generation and use of *C. reinhardtii* reporter strain UVM4-T12 for insertional
mutagenesis. a) Schematic of the constructs used for insertional mutagenesis of *C.*
reinhardtii. The pAS_R1 construct was designed to control expression of the paromomycin
resistance gene (*aphVIII*) via B$_{12}$ mediated repression of the *METE* promoter (P_{METE}). The
pHyg3 construct encoded a constitutively expressed hygromycin resistance gene (*aphVII*), to
be used for insertional mutagenesis. **b)** Growth of *C. reinhardtii* B$_{12}$ reporter strain UVM4-
T12 bearing pAS_R1 plasmid, in response to vitamin B$_{12}$ and paromomycin concentration in
the media according to the algal dose-response assay. The predicted dose-response model is
shown in black, with 95% confidence intervals in grey.
791
792
Figure 3. *C. reinhardtii* insertional mutant IM4 is defective in B$_{12}$ response and uptake,
and can be functionally complemented with *CrCBA1*. a) Effect of vitamin B$_{12}$ on *METE*
gene expression in UVM4 and IM4, determined by RT-qPCR. UVM4 and IM4 were grown
in TAP media with or without 1000 ng·l^{-1} vitamin B$_{12}$ for 4 days at 25°C, 120 rpm and in
continuous light (90 µE·m^{-2}·s^{-1}). Boxplots of the log$_2$ transformed relative expression level of
METE to the *RACK1* housekeeping gene are shown, n=6. Significant comparisons were
identified using Tukey's test: UVM4 + 1000 ng·l^{-1} vitamin B$_{12}$ from UVM4 No Addition
($p<1e^{-08}$), IM4 No Addition ($p<1e^{-08}$) and IM4 + 1000 ng·l^{-1} vitamin B12 ($p<1e^{-07}$). **b)**
Schematic of the Cre02.g081050 gene showing the position of the insertion site (indicated
with a black triangle) determined by whole genome sequencing (Figure S4). **c)** Schematic of
the pAS_C2 construct designed to express *CrCBA1* fused to the fluorescent reporter mVenus.
CrCBA1-mVenus was under the control of the *CrCBA1* promoter and terminator. pAS_C2
also contained the spectinomycin resistance gene *aadA,* driven by the *PSAD* promoter and
PSAD terminator. **d)** B$_{12}$-uptake assay with UVM4, IM4 and IM4::pAS_C2 (n =4 separate
transformants with high mVenus expression). Dashed line indicates the amount of B$_{12}$ added

to the assay. Standard deviation error bars are shown. Statistical analysis was performed on
the media fraction, and Tukey's test identified the following comparisons to be significantly
different from one another: No Algae vs UVM4 ($p<1e^{-05}$); No Algae vs IM4 ($p<0.05$); No
Algae vs IM4::pAS_C2 ($p<1e^{-03}$); UVM4 vs 1.G2 ($p<1e^{-09}$); and 1.G2 vs 1.G2::pAS_C2
($p<1e^{-06}$).
813
814
Figure 4. CLiP mutants in CrCBA1 are impaired in their ability to take up B$_{12}$. a)
Schematic of the pAS_C3 construct designed to express *CrCBA1* in a controllable manner
using a thiamine repressible riboswitch (RS$_{THI4\_4N}$) to allow repression of *CrCBA1* through
the addition of thiamine (Mehrshahi et al., 2020). **b)** B$_{12}$-uptake assay with cw15, LMJ-
040682 and mean of 3 independent transformants of LMJ-040682::pAS_C2 and LMJ-
040682::pAS_C3. The growth conditions were modified compared to previous assays: lines
were grown with or without 10 µM thiamine supplementation for 5 days in a 16/8 light/dark
cycle, and 8 hours after the dark to light transition the cultures were used for the algal B$_{12}$-
uptake assay. The dashed line indicates the amount of B$_{12}$ added to the sample. Standard
deviation error bars are shown. Statistical analysis was performed on the media fraction.
Tukey's test identified the following algal strains to be significantly different from one
another in media without thiamine (not reporting comparisons against the No Algae control
condition): cw15 vs LMJ-040682 ($p<1e^{-10}$); LMJ-040682 vs LMJ-040682::pAS_C2 ($p<1e^{-09}$); and LMJ-040682 vs LMJ-040682::pAS_C3 ($p<1e^{-09}$). Additionally, Tukey's test found
828
829 the following strain to show a significant difference due to thiamine addition: LMJ-
830 040682::pAS_C3 ($p<1e^{-07}$).
831
832
833 **Figure 5. Confocal microscopy of complemented *C. reinhardtii CrCBA1* knockout lines**
834 **showing an association between CrCBA1 and membranes.** LMJ-040682 and LMJ-
835 040682::pAS_C2 A10 and D10 lines were imaged according to the protocol outlined in the
836 materials and methods. Channels shown (left to right) are brightfield, chlorophyll, mVenus
837 and an overlay. Microscope settings are described in Methods.
838
Figure 6. CBA1 expression and B$_{12}$ uptake capacity in a B$_{12}$-dependent mutant of C.
reinhardtii (metE7) during B$_{12}$ starvation and add-back. a) CBA1 expression measured
by RT-qPCR and expressed relative to the housekeeping gene RACK1 using the $2^{\wedge}(\Delta Ct)$
method. Vertical dashed lines denote when B$_{12}$ was removed and added. b) B$_{12}$ uptake
capacity of starved metE7 cells (expressed as 10^6 molecules of B$_{12}$ per cell over 1h) at the
same 6 time points during B$_{12}$ starvation; it was not possible to perform the uptake assay on
cells to which B$_{12}$ had already been added. Cell density measurements were performed by
counting plated cells in dilution series, and so included non-viable cells. For CBA1
expression and B$_{12}$ uptake, 3 and 6 biological replicates were used, respectively, with points
representing means, and error bars representing standard deviations.
849
850
Figure 7. Distribution of CBA1 and methionine synthase sequences across Eukaryotic
groups. The EukProt database (Richter et al., 2022) was searched for METE, METH and
CBA1 queries, as described in the materials and methods. Organisms were only considered if
they contained at least one valid methionine synthase hit (METE or METH) and their
genomes were >70% complete, as measured by BUSCO (Manni et al., 2021). Eukaryotic
classes were filtered for those with greater than 5 genomes and the numbers of taxa for each
class are indicated in brackets. The different combinations of CBA1, METE and METH were

calculated for each species (Supplementary Table 4) and summarised as a percentage of the
total number of taxa in each class, with gradual shading to show the variation in distribution
between the different classes.
861
Figure 8. Identification and predicted structural location of CrCBA1 conserved
residues. a) Sequences with similarity to CBA1 were identified from the EukProt database
(Richter et al., 2022) using a manually generated CBA1 Hidden Markov Model (HMM), as
described in the materials and methods. A selection of 18 taxa from several eukaryotic
supergroups were chosen and conserved regions from the protein are presented. Specific
residues indicated by * are: K78, P118, L136, E206, F214, F215, N216, E218, P251, V253,
W255, G289, W394, F395, E396 and D408. Protein sequences are coloured according to the
Clustal colour-scheme using Geneious Prime 2021.1.1 (www.geneious.com). For each highly
conserved region, the corresponding position and amino acid from the CrCBA1 sequence
(Cre02.g081050) is indicated. **b)** The predicted 3D structure of CrCBA1 was assessed using
the Phyre2 structural prediction server using the intensive mode settings (dark blue). Highly
conserved regions of CrCBA1 are indicated in light blue and labelled. CrCBA1 was aligned
to the crystal structure of *E. coli* BtuF in complex with B_{12} (pdb: 1n2z). This enabled the
relative position of B_{12} (shown in red) to be superimposed onto CrCBA1.
876
877
878

a)

B_{12} ⊣ P_{METE} → *aphVIII* — T_{CA1}

pAS_R1

$P_{\beta2TUBULIN}$ → *aphVII* — T_{RBCS2}

pHyg3

b)

UVM4::pAS_R1 (UVM4-T12)

OD_{730}

Vitamin B_{12} concentration (ng/l)

Paromomycin concentration (µg/ml)   ○ 5   □ 10   ◇ 15   ▲ 20

a)

Relative level of *METE* expression (log2)

UVM4 | IM4

Vitamin B$_{12}$ (ng/l) added to media

No Addition — 1000

b)

Cre02.g081050

1.085Mbp   1.087Mbp   1.089Mbp

5'   3'

ATG

*

IM4 insertion site
of putative mobile
genetic element

c)

Cre02.g081050 cassette

pAS_C2 *Promoter* *ORF* *mVenus* *Terminator*

d)

pg of B$_{12}$ in fraction

Cell

Media

Total (Cell + Media)

No Algae   UVM4   IM4   IM4::pAS_C2

Class (no. of taxa)		% with METH	% with METH+METE	% with METE
Chlorophyta (18)	CBA1	61	17	0
	No CBA1	11	6	6
Stramenopiles (48)	CBA1	48	17	2
	No CBA1	4	21	8
Alveolata (11)	CBA1	45	0	0
	No CBA1	45	0	9
Rhizaria (9)	CBA1	67	0	0
	No CBA1	11	22	0
Streptophyta (22)	CBA1	0	5	73
	No CBA1	0	0	23
Amoebozoa (13)	CBA1	0	8	0
	No CBA1	31	62	0
Choanoflagellata (22)	CBA1	23	0	0
	No CBA1	77	0	0
Metazoa (42)	CBA1	0	0	0
	No CBA1	83	12	5
Fungi (27)	CBA1	0	19	7
	No CBA1	0	4	70

Parsed Citations

Allen, MD, del Campo JA, Kropat J, Merchant SS (2007) FEA1, FEA2, and FRE1, encoding two homologous secreted proteins and a candidate ferrireductase, are expressed coordinately with FOX1 and FTR1 in iron-deficient Chlamydomonas reinhardtii. Eukaryotic Cell 6: 1841–1852
Google Scholar: Author Only Title Only Author and Title

Almagro Armenteros J, Sønderby CK, Sønderby SK, Nielsen H, Winther O (2017) DeepLoc: Prediction of protein subcellular localization using deep learning. Bioinformatics 33: 3387–3395
Google Scholar: Author Only Title Only Author and Title

Almagro Armenteros J, Tsirigos KD, Sønderby CK, Petersen TN, Winther O, Brunak S, Heijne G von, Nielsen H (2019) SignalP 5.0 improves signal peptide predictions using deep neural networks. Nature Biotechnology 37: 420–423
Google Scholar: Author Only Title Only Author and Title

Banerjee R, Gouda H, Pillay S (2021) Redox-linked coordination chemistry directs vitamin B12 trafficking. Accounts of Chemical Research 54: 2003–2013
Google Scholar: Author Only Title Only Author and Title

Beedholm-Ebsen R, Wetering KVD, Hardlei T, Nexø E, Borst P, Søren K, Moestrup SK (2010) Identification of multidrug resistance protein 1 (MRP1/ABCC1) as a molecular gate for cellular export of cobalamin. Blood 115: 1632–1639
Google Scholar: Author Only Title Only Author and Title

Bertrand EM, Saito MA, Rose JM, Riesselman CR, Lohan MC, Noble AE, Lee PA, DiTullio GR (2011) Vitamin B12 and iron colimitation of phytoplankton growth in the Ross Sea. Limonol Oceanogr 52: 1079-1093
Google Scholar: Author Only Title Only Author and Title

Bertrand EM, Allen AE, Dupont CL, Norden-Krichmar TM, Bai J, Valas RE, Saito MA (2012) Influence of cobalamin scarcity on diatom molecular physiology and identification of a cobalamin acquisition protein. Proc Natl Acad Sci U S A 109: E1762–71
Google Scholar: Author Only Title Only Author and Title

Borths EL, Locher KP, Lee AT, Rees DC (2002) The structure of Escherichia coli BtuF and binding to its cognate ATP binding cassette transporter. Proc Natl Acad Sci U S A 99: 16642–7
Google Scholar: Author Only Title Only Author and Title

Bunbury F, Helliwell KE, Mehrshahi P, Davey MP, Salmon DL, Holzer A, Smirnoff N, Smith AG (2020) Responses of a newly evolved auxotroph of Chlamydomonas to B12 deprivation. Plant Physiology 183: 167–178
Google Scholar: Author Only Title Only Author and Title

Bykov YS, Schaffer M, Dodonova SO, Albert S, Plitzko JM, Baumeister W, Engel BD, Briggs JA (2017) The structure of the COPI coat determined within the cell. eLife 6: e32493
Google Scholar: Author Only Title Only Author and Title

Carlucci FA, Silbernagel BS, McNally MP (2007) The influence of temperature and solar radiation on persistence of vitamin B12, thiamine, and biotin in seawater. Journal of Phycology 5: 302–305
Google Scholar: Author Only Title Only Author and Title

Choi CC, Ford RC (2021) ATP binding cassette importers in eukaryotic organisms. Biological Reviews 96: 1318–1330
Google Scholar: Author Only Title Only Author and Title

Coelho D, Kim JC, Miousse IR, Fung S, Moulin M du, Buers I, Suormala T, Burda P, Frapolli M, Stucki M, et al (2012) Mutations in ABCD4 cause a new inborn error of vitamin B12 metabolism. Nature Genetics 44: 1152–1155
Google Scholar: Author Only Title Only Author and Title

Craig RJ, Hasan AR, Ness RW, Keightley PD (2021) Comparative genomics of Chlamydomonas. Plant Cell 33:1016-1041
Google Scholar: Author Only Title Only Author and Title

Croft MT, Lawrence AD, Raux-Deery E, Warren MJ, Smith AG (2005) Algae acquire vitamin B12 through a symbiotic relationship with bacteria. Nature 438: 90–3
Google Scholar: Author Only Title Only Author and Title

Crozet P, Navarro FJ, Willmund F, Mehrshahi P, Bakowski K, Lauersen KJ, Pérez-Pérez M-E, Auroy P, Gorchs Rovira A, Sauret-Gueto S, et al (2018) Birth of a photosynthetic chassis: A MoClo toolkit enabling synthetic biology in the microalga Chlamydomonas reinhardtii. ACS Synthetic Biology 7: 2074–2086
Google Scholar: Author Only Title Only Author and Title

Denning GM, Fulton AB (1989) Purification and characterization of clathrin-coated vesicles from Chlamydomonas. The Journal of Protozoology 36: 334–340
Google Scholar: Author Only Title Only Author and Title

Droop MR (1968) Vitamin B12 and marine ecology. IV. The kinetics of uptake, growth and inhibition in Monochrysis lutheri. Journal of the Marine Biological Association of the United Kingdom 48: 689–733
Google Scholar: Author Only Title Only Author and Title

Field CB, Behrenfeld MJ, Randerson JT, Falkowski P (1998) Primary production of the biosphere: Integrating terrestrial and oceanic components. Science 281: 237–240
Google Scholar: Author Only Title Only Author and Title

Gonzalez JC, Banerjee RV, Huang S, Sumner JS, Matthews RG (1992) Comparison of cobalamin-independent and cobalamin-dependent methionine synthases from Escherichia coli: Two solutions to the same chemical problem. Biochemistry 31: 6045–6056
Google Scholar: Author Only Title Only Author and Title

Goold HD, Cuiné S, Légeret B, Liang Y, Brugière S, Auroy P, Javot H, Tardif M, Jones B, Beisson F, et al (2016) Saturating light induces sustained accumulation of oil in plastidal lipid droplets in Chlamydomonas reinhardtii. Plant Physiology 171: 2406–2417
Google Scholar: Author Only Title Only Author and Title

Hanson AD, Roje S (2001) One-Carbon metabolism in higher plants. Ann Rev Plant Physiol 52: 119-137
Google Scholar: Author Only Title Only Author and Title

Helliwell KE, Collins S, Kazamia E, Purton S, Wheeler GL, Smith AG (2015) Fundamental shift in vitamin B12 eco-physiology of a model alga demonstrated by experimental evolution. ISME J 9: 1446–1455
Google Scholar: Author Only Title Only Author and Title

Helliwell KE, Scaife MA, Sasso S, Araujo APU, Purton S, Smith AG (2014) Unraveling vitamin B12-responsive gene regulation in algae. Plant Physiol 165: 388–397
Google Scholar: Author Only Title Only Author and Title

Helliwell KE, Wheeler GL, Leptos KC, Goldstein RE, Smith AG (2011) Insights into the evolution of vitamin B12 auxotrophy from sequenced algal genomes. Molecular Biology and Evolution 28: 2921–2933
Google Scholar: Author Only Title Only Author and Title

Hopes A, Nekrasov V, Belshaw N, Grouneva I, Kamoun S, Mock T (2017) Genome editing in diatoms using CRISPR-cas to induce precise bi-allelic deletions. Bio Protoc 7: e2625
Google Scholar: Author Only Title Only Author and Title

Joglar V, Pontiller B, Martínez-García S, Fuentes-Lema A, Pérez-Lorenzo M, Lundin D, Pinhassi J Fernández E, Teira E (2021) Microbial plankton community structure and function responses to vitamin B12 and B1 amendments in an upwelling system. Appl Environ Microbiol 87: e0152521.
Google Scholar: Author Only Title Only Author and Title

Kadner RJ (1990) Vitamin B12 transport in Escherichia coli: Energy coupling between membranes. Molecular Microbiology 4: 2027–2033
Google Scholar: Author Only Title Only Author and Title

Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. Molecular Biology and Evolution 30: 772–780
Google Scholar: Author Only Title Only Author and Title

Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJE (2015) The Phyre2 web portal for protein modeling, prediction and analysis. Nat Protoc 10: 845–58
Google Scholar: Author Only Title Only Author and Title

King N, Westbrook MJ, Young SL, Kuo A, Abedin M, Chapman J, et al. (2008). The genome of the choanoflagellate Monosiga brevicollis and the origin of metazoans. Nature. 451: 783–788.
Google Scholar: Author Only Title Only Author and Title

Koch F, Hattenrath-Lehmann TK, Goleski JA, Sañudo-Wilhelmy S, Fisher NS, Gobler CJ. 2012. Vitamin B1 and B12 uptake and cycling by plankton communities in coastal ecosystems. Front Microbiol 3:363.
Google Scholar: Author Only Title Only Author and Title

Lawrence AD, Nemoto-Smith E, Deery E, Baker JA, Schroeder S, Brown DG, Tullet JMA, Howard MJ, Brown IR, Smith AG, et al (2018) Construction of fluorescent analogs to follow the uptake and distribution of cobalamin (vitamin B12) in bacteria, worms, and plants. Cell Chem Biol 25: 941–951
Google Scholar: Author Only Title Only Author and Title

Li X, Zhang R, Patena W, Gang SS, Blum SR, Ivanova N, Yue R, Robertson JM, Lefebvre PA, Fitz-Gibbon ST, et al (2016) An indexed, mapped mutant library enables reverse genetics studies of biological processes in Chlamydomonas reinhardtii. Plant Cell 28: 367–87
Google Scholar: Author Only Title Only Author and Title

Mackinder LCM, Chen C, Leib RD, Patena W, Blum SR, Rodman M, Ramundo S, Adams CM, Jonikas MC (2017) A spatial interactome reveals the protein organization of the algal CO2 concentrating mechanism. Cell 171: 133–147.e14
Google Scholar: Author Only Title Only Author and Title

Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM (2021) BUSCO update: Novel and streamlined workflows along with broader and deeper phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes. Molecular Biology and Evolution 38: 4647–4654
Google Scholar: Author Only Title Only Author and Title

Mehrshahi P, Nguyen GTDT, Gorchs Rovira A, Sayer A, Llavero-Pasquina M, Lim Huei Sin M, Medcalf EJ, Mendoza-Ochoa GI, Scaife MA, Smith AG (2020) Development of novel riboswitches for synthetic biology in the green alga Chlamydomonas. ACS Synthetic Biology 9: 1406–1417
Google Scholar: Author Only Title Only Author and Title

Mentch SJ, Locasale JW (2016) One-carbon metabolism and epigenetics: understanding the specificity Annal NY Acad Sci 1363: 91-8
Google Scholar: Author Only Title Only Author and Title

Mitchell AL, Attwood TK, Babbitt PC, Blum M, Bork P, Bridge A, Brown SD, Chang H-Y, El-Gebali S, Fraser MI et al. (2019) InterPro in 2019: improving coverage, classification and access to protein sequence annotations. Nucl Acids Res 47: D351–D360
Google Scholar: Author Only Title Only Author and Title

Neupert J, Karcher D, Bock R (2009) Generation of Chlamydomonas strains that efficiently express nuclear transgenes. The Plant Journal 57: 1140–1150
Google Scholar: Author Only Title Only Author and Title

Nielsen MJ, Rasmussen MR, Andersen CBF, Nexø E, Moestrup SK (2012) Vitamin B12 transport from food to the body's cells-a sophisticated, multistep pathway. Nature Reviews Gastroenterology & Hepatology 9: 345–354
Google Scholar: Author Only Title Only Author and Title

Ohwada K (1973) Seasonal cycles of vitamin B12, thiamine and biotin in Lake Sagami. Patterns of their distribution and ecological significance. Internationale Revue der gesamten Hydrobiologie und Hydrographie 58: 851–871
Google Scholar: Author Only Title Only Author and Title

Orłowska M, Steczkiewicz K, Muszewska A (2021) Utilization of cobalamin is ubiquitous in early-branching fungal phyla. Genome Biology and Evolution 13: evab043
Google Scholar: Author Only Title Only Author and Title

Panzeca C, Beck AJ, Tovar-Sanchez A, Segovia-Zavala J, Taylor GT, Gobler CJ, Sañudo-Wilhelmy SA (2009) Distributions of dissolved vitamin B12 and Co in coastal and open-ocean environments. Estuarine, Coastal and Shelf Science 85: 223–230
Google Scholar: Author Only Title Only Author and Title

Pintner IJ, Altmeyer VL (1979) Vitamin B12-binder and other algal inhibitors Journal of Phycology 15: 391–398

Richter DJ, Berney C, Strassert JFH, Poh Y-P, Herman EK, Muñoz-Gómez SA, Wideman JG, Burki F, Vargas C de (2022) EukProt: A database of genome-scale predicted proteins across the diversity of eukaryotes. Peer Community Journal 2: e56
Google Scholar: Author Only Title Only Author and Title

Ritz C, Baty F, Streibig JC, Gerhard D (2015) Dose-response analysis using R. PLoS One 10: e0146021
Google Scholar: Author Only Title Only Author and Title

Rutsch F, Gailus S, Miousse IR, Suormala T, Sagné C, Toliat MR, Nürnberg G, Wittkampf T, Buers I, Sharifi A, et al (2009) Identification of a putative lysosomal cobalamin exporter altered in the cblF defect of vitamin B12 metabolism. Nature Genetics 41: 234–239
Google Scholar: Author Only Title Only Author and Title

Sahni MK, Spanos S, Wahrman MZ, Sharma GM (2001) Marine corrinoid-binding proteins for the direct determination of vitamin B12 by radioassay. Analytical Biochemistry 289: 68–76
Google Scholar: Author Only Title Only Author and Title

Sañudo-Wilhelmy SA, Gómez-Consarnau L, Suffridge C, Webb EA (2014) The role of B vitamins in marine biogeochemistry. Annual Review of Marine Science 6: 339–367
Google Scholar: Author Only Title Only Author and Title

Shelton AN, Seth EC, Mok KC, Han AW, Jackson SN, Haft DR, Taga ME (2019) Uneven distribution of cobamide biosynthesis and dependence in bacteria predicted by comparative genomics. The ISME Journal 13: 789–804
Google Scholar: Author Only Title Only Author and Title

Tang YZ, Koch F, Gobler CJ (2010) Most harmful algal bloom species are vitamin B1 and B12 auxotrophs. Proc Natl Acad Sci U S A 107: 20756–61

Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Warren MJ, Raux E, Schubert HL, Escalante-Semerena JC (2002) The biosynthesis of adenosylcobalamin (vitamin B12). Natural product reports 19: 390–412

Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Xie B, Bishop S, Stessman D, Wright D, Spalding MH, Halverson LJ (2013) Chlamydomonas reinhardtii thermal tolerance enhancement mediated by a mutualistic interaction with vitamin B12-producing bacteria. ISME J 7: 1544–55

Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Yu Z, Geisler K, Leontidou T, Young REB, Vonlanthen SE, Purton S, Abell C, Smith AG (2021) Droplet-based microfluidic screening and sorting of microalgal populations for strain engineering applications. Algal Res 56: 102293

Google Scholar: <u>Author Only</u> <u>Title Only</u> <u>Author and Title</u>

Figure 1

Figure 1. Disruption of *Phaeodactylum tricornutum CBA1* (*PtCBA1*) using CRISPR-Cas9 yielded lines with impaired B$_{12}$ uptake. a) Schematic showing CRISPR-Cas9 sgRNA target sites and the homology repair template design used to generate mutant lines in *PtCBA1* (Phatr3_J48322). The homology repair template schematic is annotated with the 5' homology region (HR) and 3'HR, the *FCPB* promoter, nourseothricin resistance gene (*NAT*) and *FCPC* terminator. The *PtCBA1* gene is annotated with the ORF, the 5'HR and 3'HR regions used in the homology template and the regions of the ORF targeted by sgRNA (vertical bars). Primer positions used for the analysis of putative mutant lines are shown with arrowheads. **b)** PCR of regions across and within wild-type (WT) and mutant *PtCBA1* in 3 independent CRISPR-Cas9 lines (ΔCBA1) showing indel mutations in the mutants. PCR products from different sets of primers indicated in panel a are shown. M = marker, - Ctrl = no DNA template. **c)** A B$_{12}$ uptake assay was performed as described in Materials and Methods, to determine the amount of B$_{12}$ in the media and the cells after 1h incubation of *P. tricornutum* cells in 600 pg B$_{12}$. The 'Total' was inferred by the addition of the cell and media fractions. The dashed line indicates the amount of B$_{12}$ added to the experiment. Standard deviation error bars are shown, n=4. Statistical analysis was performed on the media fraction, and Tukey's test identified the following comparisons to be significantly different from one another: WT vs No Algae ($p<1e^{-12}$); WT vs ΔCBA1-1 ($p<1e^{-10}$); WT vs ΔCBA1-2 ($p<1e^{-12}$); WT vs ΔCBA1-3 ($p<1e^{-11}$); No Algae vs ΔCBA1-1 ($p<1e^{-03}$); No Algae vs ΔCBA1-3 ($p<0.05$); and ΔCBA1-1 vs ΔCBA1-2 ($p<1e^{-02}$).

Figure 2

Figure 2. Generation and use of *C. reinhardtii* reporter strain UVM4-T12 for insertional mutagenesis. a) Schematic of the constructs used for insertional mutagenesis of *C. reinhardtii*. The pAS_R1 construct was designed to control expression of the paromomycin resistance gene (*aphVIII*) via B_{12} mediated repression of the *METE* promoter (P_{METE}). The pHyg3 construct encoded a constitutively expressed hygromycin resistance gene (*aphVII*), to be used for insertional mutagenesis. **b)** Growth of *C. reinhardtii* B_{12} reporter strain UVM4-T12 bearing pAS_R1 plasmid, in response to vitamin B_{12} and paromomycin concentration in the media according to the algal dose-response assay. The predicted dose-response model is shown in black, with 95% confidence intervals in grey.

Figure 3

Figure 3. *C. reinhardtii* insertional mutant IM4 is defective in B$_{12}$ response and uptake, and can be functionally complemented with *CrCBA1*. a) Effect of vitamin B$_{12}$ on *METE* gene expression in UVM4 and IM4, determined by RT-qPCR. UVM4 and IM4 were grown in TAP media with or without 1000 ng·l^{-1} vitamin B$_{12}$ for 4 days at 25°C, 120 rpm and in continuous light (90 μE·m^{-2}·s^{-1}). Boxplots of the log$_2$ transformed relative expression level of *METE* to the *RACK1* housekeeping gene are shown, n=6. Significant comparisons were identified using Tukey's test: UVM4 + 1000 ng·l^{-1} vitamin B$_{12}$ from UVM4 No Addition (p<1e^{-08}), IM4 No Addition (p<1e^{-08}) and IM4 + 1000 ng·l^{-1} vitamin B12 (p<1e^{-07}). **b)** Schematic of the Cre02.g081050 gene showing the position of the insertion site (indicated with a black triangle) determined by whole genome sequencing (Figure S4). **c)** Schematic of the pAS_C2 construct designed to express *CrCBA1* fused to the fluorescent reporter mVenus. *CrCBA1-mVenus* was under the control of the *CrCBA1* promoter and terminator. pAS_C2 also contained the spectinomycin resistance gene *aadA,* driven by the *PSAD* promoter and *PSAD* terminator. **d)** B$_{12}$-uptake assay with UVM4, IM4 and IM4::pAS_C2 (n =4 separate transformants with high mVenus expression). Dashed line indicates the amount of B$_{12}$ added to the assay. Standard deviation error bars are shown. Statistical analysis was performed on the media fraction, and Tukey's test identified the following comparisons to be significantly different from one another: No Algae vs UVM4 (p<1e^{-05}); No Algae vs IM4 (p<0.05); No Algae vs IM4::pAS_C2 (p<1e^{-03}); UVM4 vs 1.G2 (p<1e^{-09}); and 1.G2 vs 1.G2::pAS_C2 (p<1e^{-06}).

Figure 4

Figure 4. CLiP mutants in CrCBA1 are impaired in their ability to take up B$_{12}$. a) Schematic of the pAS_C3 construct designed to express *CrCBA1* in a controllable manner using a thiamine repressible riboswitch (RS$_{THI4\_4N}$) to allow repression of *CrCBA1* through the addition of thiamine (Mehrshahi et al., 2020). **b)** B$_{12}$-uptake assay with cw15, LMJ-040682 and mean of 3 independent transformants of LMJ-040682::pAS_C2 and LMJ-040682::pAS_C3. The growth conditions were modified compared to previous assays: lines were grown with or without 10 µM thiamine supplementation for 5 days in a 16/8 light/dark cycle, and 8 hours after the dark to light transition the cultures were used for the algal B$_{12}$-uptake assay. The dashed line indicates the amount of B$_{12}$ added to the sample. Standard deviation error bars are shown. Statistical analysis was performed on the media fraction. Tukey's test identified the following algal strains to be significantly different from one another in media without thiamine (not reporting comparisons against the No Algae control condition): cw15 vs LMJ-040682 ($p<1e^{-10}$); LMJ-040682 vs LMJ-040682::pAS_C2 ($p<1e^{-09}$); and LMJ-040682 vs LMJ-040682::pAS_C3 ($p<1e^{-09}$). Additionally, Tukey's test found the following strain to show a significant difference due to thiamine addition: LMJ-040682::pAS_C3 ($p<1e^{-07}$).

Figure 5

Figure 5. Confocal microscopy of complemented *C. reinhardtii CrCBA1* knockout lines showing an association between CrCBA1 and membranes. LMJ-040682 and LMJ-040682::pAS_C2 A10 and D10 lines were imaged according to the protocol outlined in the materials and methods. Channels shown (left to right) are brightfield, chlorophyll, mVenus and an overlay. Microscope settings are described in Methods.

Figure 6

Figure 6. CBA1 expression and B$_{12}$ uptake capacity in a B$_{12}$-dependent mutant of C. reinhardtii (metE7) during B$_{12}$ starvation and add-back. a) CBA1 expression measured by RT-qPCR and expressed relative to the housekeeping gene RACK1 using the 2^(ΔCt) method. Vertical dashed lines denote when B$_{12}$ was removed and added. b) B$_{12}$ uptake capacity of starved metE7 cells (expressed as 10^6 molecules of B$_{12}$ per cell over 1h) at the same 6 time points during B$_{12}$ starvation; it was not possible to perform the uptake assay on cells to which B$_{12}$ had already been added. Cell density measurements were performed by counting plated cells in dilution series, and so included non-viable cells. For CBA1 expression and B$_{12}$ uptake, 3 and 6 biological replicates were used, respectively, with points representing means, and error bars representing standard deviations.

Figure 7

Class (no. of taxa)		% with METH	% with METH+METE	% with METE
Chlorophyta (18)	CBA1	61	17	0
	No CBA1	11	6	6
Stramenopiles (48)	CBA1	48	17	2
	No CBA1	4	21	8
Alveolata (11)	CBA1	45	0	0
	No CBA1	45	0	9
Rhizaria (9)	CBA1	67	0	0
	No CBA1	11	22	0
Streptophyta (22)	CBA1	0	5	73
	No CBA1	0	0	23
Amoebozoa (13)	CBA1	0	8	0
	No CBA1	31	62	0
Choanoflagellata (22)	CBA1	23	0	0
	No CBA1	77	0	0
Metazoa (42)	CBA1	0	0	0
	No CBA1	83	12	5
Fungi (27)	CBA1	0	19	7
	No CBA1	0	4	70

Figure 7. Distribution of CBA1 and methionine synthase sequences across Eukaryotic groups. The EukProt database (Richter et al., 2022) was searched for METE, METH and CBA1 queries, as described in the materials and methods. Organisms were only considered if they contained at least one valid methionine synthase hit (METE or METH) and their genomes were >70% complete, as measured by BUSCO (Manni et al., 2021). Eukaryotic classes were filtered for those with greater than 5 genomes and the numbers of taxa for each class are indicated in brackets. The different combinations of CBA1, METE and METH were calculated for each species (Supplementary Table 4) and summarised as a percentage of the total number of taxa in each class, with gradual shading to show the variation in distribution between the different classes.

Figure 8

Figure 8. Identification and predicted structural location of CrCBA1 conserved residues. a) Sequences with similarity to CBA1 were identified from the EukProt database (Richter et al., 2022) using a manually generated CBA1 Hidden Markov Model (HMM), as described in the materials and methods. A selection of 18 taxa from several eukaryotic supergroups were chosen and conserved regions from the protein are presented. Specific residues indicated by * are: K78, P118, L136, E206, F214, F215, N216, E218, P251, V253, W255, G289, W394, F395, E396 and D408. Protein sequences are coloured according to the Clustal colour-scheme using Geneious Prime 2021.1.1 (www.geneious.com). For each highly conserved region, the corresponding position and amino acid from the CrCBA1 sequence (Cre02.g081050) is indicated. **b)** The predicted 3D structure of CrCBA1 was assessed using the Phyre2 structural prediction server using the intensive mode settings (dark blue). Highly conserved regions of CrCBA1 are indicated in light blue and labelled. CrCBA1 was aligned to the crystal structure of *E. coli* BtuF in complex with B_{12} (pdb: 1n2z). This enabled the relative position of B_{12} (shown in red) to be superimposed onto CrCBA1.

Figure S1. Characterisation of B$_{12}$ uptake in *C. reinhardtii* insertional mutant lines. To determine whether any of the 7 insertional mutant lines (IM) isolated from mutagenesis of UVM4::T12 showed impaired B$_{12}$ uptake, the B$_{12}$-uptake assay was performed as described in the materials and methods. Control-1, Control-2 and Control-3 lines were picked from solid media with paromomycin and hygromycin but without vitamin B$_{12}$, whereas IM1-IM7 lines were picked from solid media containing paromomycin, hygromycin and vitamin B$_{12}$. The total was inferred by the addition of the cell and media fractions. The dashed line indicates the amount of B$_{12}$ added in the uptake assay. Standard deviation error bars are shown, n=4. Statistical analysis was performed on the media fraction, and Tukey's test identified the following comparisons to be significantly different from one another (only reporting IM strains different from UVM4, UVM4-T12 or Control-[1,2,3] strains): IM4 vs UVM4 ($p<1e^{-12}$), UVM4-T12 ($p<1e^{-12}$), Control-1 ($p<1e^{-12}$), Control-2 ($p<1e^{-12}$), Control-3 ($p<1e^{-12}$); and IM2 vs UVM4 ($p<0.05$).

Figure S2

Figure S2. Visualisation of B$_{12}$-BODIPY uptake in *C. reinhardtii* using confocal microscopy. To assess B$_{12}$ uptake more directly, the insertional mutant line IM4 (impaired B$_{12}$ uptake) and its parental line UVM4-T12 were incubated with the fluorescent B$_{12}$ analogue B$_{12}$-BODIPY and the samples were imaged using confocal microscopy, as described in the materials and methods. Channels shown are brightfield (greyscale), chlorophyll (red), B$_{12}$-BODIPY (yellow) and an overlay.

Figure S3

Figure S3. *C. reinhardtii* knockout lines of *Cre12.g508644* are able to take up B₁₂. a) Schematic showing the structure of *Cre12.g508644* with annotations for the 5'UTR (medium grey), start codon (ATG), exons (light grey), introns (black lines), stop codon (*) and 3'UTR (black). The location of the IM4 pHyg3 insertion site (identified by DNA sequencing and validated using PCR) is indicated with a black triangle. The predicted disruption sites in CLiP (Li et al., 2016) knockout strains LMJ-119922 and LMJ-042227 are indicated with grey triangles. These were also confirmed by PCR (insets) **b)** B₁₂-uptake assay of the CLiP mutants and their background strain cw15. The dashed line shows the amount of B₁₂ added to the experiment. Total = inferred by addition of amounts determined in the cellular and media fractions. Standard deviation error bars are shown, No Algae (n=10), cw15 (n=10), LMJ-119922 (n=10) and LMJ-042227 (n=6). Statistical analysis was performed on the media fraction, and Tukey's test identified the following comparisons to be significantly different from one another: No Algae vs cw15 ($p<1e^{-12}$); No Algae vs LMJ-119922 ($p<1e^{-12}$); No Algae vs LMJ-042227 ($p<1e^{-08}$); cw15 vs LMJ-042227 ($p<1e^{-03}$); and LMJ-119922 vs LMJ-042227 ($p<0.05$).

Figure S4

a) Insertion site and sequence found in IM4

1,084,548 · Left junction (+) 148 bp · ? bp · Right junction (+) 227 bp + 8 bp · 1,084,549

b) Chromosomal regions with similarity to the IM4 left and right junction sequences

Figure S4. Structure and sequence of a second insertion in the IM4 strain. a) Mapping of the WGS data to the reference strain revealed an extra sequence between positions 1,084,548 bp and 1,084,549 bp of chromosome 2 in the *C. reinhardtii* v5.0 Phytozome reference genome corresponding to the Cre02.g081050 locus. An 8 bp target site duplication 'TACTACAG' was identified flanking the insertion (blue). The sequences of the left (red) and right (green) insertion junctions were identified from DNA sequencing reads and confirmed with PCR and sequencing. The sequence between these left and right junctions has not been determined. **b)** Chromosomal regions with similarity to the left (148 bp) and right (227 bp) junction sequences were identified by a BLAST search. Shown are three regions with sequence similarity to the left and right junctions and where the left and right junctions are a similar distance apart (~10 Kb) and in the same orientation. The position of the matching left junction (red) and right junction (green) sequences are indicated. The matching strand is indicated in brackets. The number of matching base pairs from the BLAST search is listed in brackets (target/query).

Figure S5

Figure S5. The predicted structures of CrCBA1 and PtCBA1 show a high degree of structural similarity. Structural prediction of CrCBA1 (Cre02.g081050) (blue) and PtCBA1 (Phatr3_J48322) (gold) was performed using the Phyre2 structural prediction server (Kelley et al., 2015). CrCBA1 was modelled with 56% of residues predicted with greater than 90% accuracy; PtCBA1 was modelled with 69% of residues predicted with greater than 90% accuracy. Structures were aligned using the super command in PyMOL, root mean squared deviation = 2.333. Conserved alpha helices are seen in the centre of the image, as is a common cleft region at the bottom of the image.

a)

Figure S6. Independent mutant lines of *CrCBA1* show defective B$_{12}$ uptake. A) Schematic showing the structure of *CrCBA1* (Cre02.g081050) with annotations for the 5'UTR (medium grey), start codon (ATG), exons (light grey), introns (black lines), stop codon (*) and 3'UTR (black). The location of predicted disruption sites in *CrCBA1* CliP knockout strains LMJ-135929 and LMJ-040682 are indicated (triangles). Primer positions used for the analysis of mutant lines are shown with arrows. Insets show that primers amplifying between exons 2 and 4 produced a band of the expected size for control lines cw15, LMJ-040682 and LMJ-042227, whereas LMJ-135929 showed an increased amplicon size, indicating an insertion in this region. Primers between exons 4 and 8 produced a band of the expected size in UVM4 and the same-background control lines LMJ-042227 and LMJ-135929, whereas LMJ-040682 lacked this band, indicating a disruption in this region. **B)** B$_{12}$-uptake assay. The dashed line indicates the amount of B$_{12}$ added to the sample. Standard deviation error bars are shown, No Algae (n=14), cw15 (n=10), LMJ-135929 (n=6) and LMJ-040682 (n=18). Statistical analysis was performed on the media fraction, and Tukey's test identified the following comparisons to be significantly different from one another: No Algae vs cw15 ($p<1e^{-12}$); No Algae vs LMJ-135929 ($p<1e^{-02}$); No Algae vs LMJ-040682 ($p<1e^{-02}$); cw15 vs LMJ-135929 ($p<1e^{-11}$); cw15 vs LMJ-040682 ($p<1e^{-12}$); and LMJ-135929 vs LMJ-040682 ($p<1e^{-06}$).

Figure S7

a)

b)

Figure S7. Sequences with similarity to CBA1 are found throughout Eukaryota.
Sequences with similarity to CBA1 were identified from the EukProt database (Richter et al., 2022) using a manually generated hidden Markov model for CBA1, as described in the materials and methods. Positively classified CBA1 sequences were aligned with MAFFT (--auto option) (Katoh and Standley, 2013) and trimmed with trimai (-automated1 option) (Capella-Gutiérrez et al., 2009), and Iqtree version 1.6.10 (options: -bb 1000, -safe, -bnni, -alrt 1000, -st AA, -seed 1000, -msub nuclear, -t RANDOM, -m TEST) (Nguyen et al., 2015) was used to produce a gene tree. Visualisation was performed using iTOL (Letunic and Bork, 2019). A multiple sequence alignment of the full length protein sequences was generated and visualised using Geneious Prime 2021.1.1 (https://www.geneious.com), and is shown adjacent to the tree (options: Clustal colour scheme, agreements to consensus highlighted, sliding window size = 5 bp, and sites with over 50% gaps hidden). A conservation track and consensus track are shown above the alignment panel, with regions of conserved residues indicated, detailed more in Figure 8a.