1   **Antibiotics promote intestinal growth of carbapenem-resistant *Enterobacteriaceae***

2   **by enriching nutrients and depleting microbial metabolites**

4   Alexander Y. G. Yip[1], Olivia G. King[2], Oleksii Omelchenko[1], Sanjana Kurkimat[1], Victoria Horrocks[1],

5   Phoebe Mostyn[1], Nathan Danckert[3], Rohma Ghani[3,4], Giovanni Satta[5], Frances J. Davies[4], Thomas

6   B. Clarke[2], Benjamin H. Mullish[3,6], Julian R. Marchesi[3], Julie A. K. McDonald[1]*

Affiliations:

[1] Department of Life Sciences, Imperial College London, London SW7 2AZ, United Kingdom.

[2] Department of Infectious Disease, Imperial College London, London SW7 2AZ, United Kingdom.

[3] Division of Digestive Diseases, Department of Metabolism, Digestion and Reproduction, Faculty of

Medicine, St Mary's Hospital Campus, Imperial College London, London, United Kingdom.

[4] Department of Infectious Disease, Imperial College Healthcare NHS Trust, London, United

Kingdom.

[5] UCL Centre for Clinical Microbiology, University College London, United Kingdom.

[6] Departments of Gastroenterology and Hepatology, St Mary's Hospital, Imperial College Healthcare

NHS Trust, Paddington, London, UK

* Corresponding author

Correspondence:

Dr. Julie A. K. McDonald

Imperial College London

Room 1.44, Flowers Building

Exhibition Road

London SW7 2AZ

Email: julie.mcdonald@imperial.ac.uk

Telephone: +44 (0)20 7594 5247

Acknowledgements: This work was supported by funding awarded to JAKM: a Medical Research Council (MRC) New Investigator Research Grant (MR/W025655/1), a Wellcome Trust Institutional Strategic Support Fund Springboard Fellowship, and startup funds from the Department of Life Sciences at Imperial College London. FJD received funding from the Medical Research Council (MRC) Clinical Academic Research Partnership Scheme. TBC received funding from a UKRI Impact Accelerator Award (MR/X502959/1). BHM is the recipient of an NIHR Academic Clinical Lectureship (CL-2019-21-002). The Division of Digestive Diseases received financial and infrastructure support from the National Institute for Health Research (NIHR) Imperial Biomedical Research Centre (BRC) based at Imperial College Healthcare NHS Trust and Imperial College London.

Author contributions:

Alexander Y. G. Yip, MRes = formal analysis: equal; investigation: lead; methodology: equal; visualisation: equal; writing – review & editing: supporting

Olivia G. King, MSc = investigation: supporting; writing – review & editing: supporting

Oleksii Omelchenko, MRes = formal analysis: supporting; investigation: supporting

Sanjana Kurkimat, MSc = formal analysis: supporting; investigation: supporting

Victoria Horrocks, PhD = formal analysis: supporting; investigation: supporting; writing – review & editing: supporting

Phoebe Mostyn, MRes = formal analysis: supporting; investigation: supporting

Nathan Danckert, PhD = investigation: supporting

Rohma Ghani, MB BS = resources: supporting, writing – review & editing: supporting

Giovanni Satta, PhD = resources: supporting, writing – review & editing: supporting

Frances J. Davies, PhD = resources: supporting; writing – review & editing: supporting

Thomas B. Clarke, PhD = investigation: supporting; methodology: supporting; writing – review & editing: supporting

Benjamin H. Mullish, PhD = resources: supporting; writing – review & editing: supporting

Julian R. Marchesi, PhD = resources: supporting; writing – review & editing: supporting

Julie A. K. McDonald, PhD = conceptualisation: lead; data curation: lead; formal analysis: equal; funding acquisition: lead; methodology: equal; project administration: lead; resources: lead; supervision: lead; visualisation: equal; writing – original draft: lead; writing – review & editing: lead

Abbreviations: ^{1}H-NMR, proton nuclear magnetic resonance; AMiGA, Analysis of Microbial Growth Assays; ANOVA, analysis of variance; BCFA, branched chain fatty acid; CAZ, ceftazidime; CFU, colony forming units; CIP, ciprofloxacin; CRE, carbapenem-resistant *Enterobacteriaceae*; CRO, ceftriaxone; CTX, cefotaxime; ETP, ertapenem; FDR, false discovery rate; IPM, imipenem/cilastatin; LB, Luria broth; LOD, limit of detection; MEM, meropenem; NDM-1, New Delhi Metallo-β-lactamase 1; OD_{600}, optical density at 600 nm; PBS, phosphate-buffered saline; PCR, polymerase chain reaction; rCCA, regularised canonical correlation analysis; SCFA, short chain fatty acid; TSP, 3-(Trimethylsilyl)propionic-2,2,3,3-d4 acid sodium salt; TZP, piperacillin/tazobactam

Keywords: gut microbiota; nutrient competition; metabolite inhibition; colonisation resistance

ABSTRACT

The intestine is the primary colonisation site for carbapenem-resistant *Enterobacteriaceae* (CRE) and serves as a reservoir of CRE that cause invasive infections (e.g. bloodstream infections). Antibiotics disrupt colonisation resistance mediated by the gut microbiota, promoting the expansion of CRE within the intestine. We used *ex vivo* faecal cultures to measure the impact of antibiotics (that promote CRE intestinal colonisation) on the faecal microbiota from healthy human donors. We demonstrated that antibiotics decreased the abundance of gut commensals (including *Bifidobacteriaceae* and *Bacteroidales*) in human faecal microbiota, resulting in an enrichment of nutrients and a depletion of microbial metabolites. The nutrient utilisation abilities, nutrient preferences, and metabolite inhibition susceptibilities of carbapenem-resistant *Escherichia coli*, *Klebsiella pneumoniae*, and *Enterobacter cloacae* strains were measured. Nutrients (which were elevated with antibiotics) acted as carbon and nitrogen sources to support CRE growth, where CRE

showed an ordered preference for specific nutrients. These nutrients were also increased in faeces from antibiotic-treated mice but decreased following intestinal colonisation with carbapenem-resistant *E. coli*. Microbial metabolites (which decreased with antibiotics) were inhibitory towards CRE growth *in vitro*. Carbapenem-resistant *E. coli* growth was decreased in faecal samples from mice treated with a mixture of inhibitory metabolites compared with PBS-treated mice. These findings demonstrate that killing gut commensals with antibiotics disrupts colonisation resistance by enriching nutrients that support CRE growth and depleting metabolites that inhibit CRE growth. These results support the development of new microbiome therapeutics to prevent CRE intestinal colonisation, which would also prevent the subsequent development of invasive CRE infections.

INTRODUCTION

Antibiotic resistance presents a serious threat to human health, resulting in treatment failures, infection relapses, longer hospitalisations, poor clinical outcomes, and increased healthcare costs[1]. Treatment options are increasingly limited, less effective, and may involve administration of antibiotics that are more toxic to the patient[2]. There is an urgent need to develop new treatments for infections by antibiotic-resistant bacteria. In particular, alternative approaches are required to prevent the development of invasive infections by extremely drug-resistant pathogens such as carbapenem-resistant *Enterobacteriaceae* (CRE), which present an urgent public health threat[2]. However, there are many difficulties associated with the development of new antibiotics to treat these multidrug-resistant infections[2].

The intestine is the primary colonisation site for CRE and serves as a reservoir of CRE that seed difficult-to-treat invasive infections (such as bloodstream infections and recurrent urinary tract infections)[3]. Therefore, one approach to prevent the development of invasive CRE infections would be to prevent CRE from colonising the intestine, or to decolonise patients with pre-existing CRE intestinal colonisation. However, to do so we must first understand what drives CRE colonisation in the intestines of susceptible patients.

Healthy gut microbiota exhibit colonisation resistance, where gut commensals prevent pathogens from colonising the intestine. However, it is well established that broad spectrum antibiotics disrupt colonisation resistance and significantly promote the expansion of pathogens within the intestine[4]. In particular, carbapenems, piperacillin/tazobactam, ciprofloxacin, and cephalosporins are known to promote CRE intestinal colonisation[5]. The mechanisms of colonisation resistance that healthy gut microbiota use to protect against CRE intestinal colonisation are not fully understood. However, based on findings from other pathogens, nutrient competition and metabolite inhibition are expected to play an important role[6-10].

Bacteria must have access to nutrients that support their growth to successfully colonise the intestine. However, nutrients are limited within the intestine as bacteria with similar or "overlapping" nutrient utilisation abilities will compete for these nutrients and occupy similar niches[11]. Different bacteria have different nutrient utilisation abilities, and the diversity and concentration of the available nutrients will impact their growth[12]. Although bacteria may be able to utilise a variety of nutrients, they generally have a prioritised utilisation of nutrients that varies between different bacterial species[13]. The nutrient-niche hypothesis predicts that antibiotic-mediated disruption of the gut microbiota leads to reduced competition for nutrients and an increase in nutrient availability in the gut, which could promote *Enterobacteriaceae* growth[14].

Nutrient metabolism by the gut microbiota results in the production of metabolites, some of which may be inhibitory towards pathogen growth[7,8]. Polysaccharide fermentation results in the production of metabolites such as short chain fatty acids (SCFAs; e.g. formate, acetate, propionate, butyrate, valerate) and organic acids (e.g. succinate, lactate, ethanol)[15]. Protein fermentation results in the production of SCFAs and branched chain fatty acids (BCFAs; e.g. isobutyrate and isovalerate)[15]. However, killing gut commensals with antibiotics can decrease the production of microbial metabolites and enable pathogen growth[7,16].

Broad-spectrum antibiotics kill a wide range of gut commensals and promote the intestinal colonisation with CRE[5]. However, to date previous studies have only identified a very small number

of nutrients or metabolites that are disrupted with antibiotics and involved in CRE intestinal colonisation[16,17]. Given the severe impact of broad-spectrum antibiotics on the gut microbiota, we expected that these antibiotics would have a much more significant impact on nutrient metabolism in the intestine. Understanding how antibiotics disrupt colonisation resistance to promote CRE growth will facilitate the rational design of microbiome therapeutics to prevent or treat CRE intestinal colonisation, and therefore prevent the subsequent development of difficult-to-treat invasive CRE infections. However, microbiome therapeutics that only deplete a small subset of the available nutrients that CRE can utilise will have limited clinical impact to prevent or restrict CRE growth, as CRE will switch nutrient metabolism to utilise alternative available nutrients.

We hypothesised that antibiotics significantly enrich for wide range and diversity of nutrients and deplete a wide range and diversity of metabolites, resulting in a significant change in the nutrient and metabolite landscape CRE encounter in antibiotic-treated gut microbiota and creating a niche that promotes CRE growth. We demonstrated that broad-spectrum antibiotics that promote CRE growth enriched for a wide range of nutrients in the intestine, generating a "nutrient-enriched niche" that supports CRE growth. We also demonstrated that these broad-spectrum antibiotics also depleted a wide range of metabolites and generated a "metabolite-depleted niche" that no longer inhibited CRE growth. We showed that these nutrients could act as carbon and nitrogen sources to support CRE growth, where CRE showed an ordered preference for specific nutrients. Oxygen availability is also increased in an antibiotic-treated gut, and we showed that CRE growth was higher on these nutrients in the presence of oxygen. Finally, we showed that microbial metabolites (that were decreased with antibiotics) were inhibitory towards CRE growth. Together, these results demonstrated that killing gut commensals with antibiotics reduces colonisation resistance by disrupting nutrient metabolism. This results in an intestinal niche that supports the expansion of CRE by enriching nutrients that support CRE growth and depleting metabolites that inhibit CRE growth. We demonstrated that bacterial consortia that compose microbiome therapeutics must fully occupy the same nutrient-defined intestinal niche that CRE occupy in the gut to effectively prevent or restrict CRE intestinal growth.

RESULTS

Antibiotics decreased the abundance of important bacterial families in faecal microbiota

We performed *ex vivo* faecal culture experiments to measure the effects of eight broad-spectrum antibiotics on the faecal microbiota from 11 healthy human donors. Antibiotics tested included meropenem (MEM), imipenem/cilastatin (IPM), ertapenem (ETP), piperacillin/tazobactam (TZP), ciprofloxacin (CIP), ceftriaxone (CRO), ceftazidime (CAZ), and cefotaxime (CTX). These antibiotics are frequently used clinically and known to promote the intestinal colonisation with CRE[5]. Faecal cultures are highly reproducible, allowing for the parallel testing of multiple groups that are inoculated from a single faecal sample[18]. For each experiment a faecal sample was inoculated into a distal gut growth medium supplemented with one of the eight antibiotics or not supplemented with antibiotics ("antibiotic-naïve" control). First, we measured how these different antibiotics (that promote susceptibility to CRE intestinal colonisation) impacted the abundance of gut commensal taxa from faecal microbiota.

We found that antibiotics decreased the absolute abundance of several bacterial families, in particular *Bifidobacteriaceae* and families from the order *Bacteroidales* (**Fig. 1a**). Carbapenems decreased the abundance of *Bifidobacteriaceae*, *Coriobacteriaceae*, *Bacteroidaceae*, *Barnesiellaceae*, *Butyricicoccaceae,* and other bacterial families. TZP decreased the abundance of *Bifidobacteriaceae* and *Bacteroidaceae*. CIP decreased the abundance of *Bifidobacteriaceae*, *Barnesiellaceae* and *Sutterellaceae*. Cephalosporins decreased the abundance of *Bifidobacteriaceae* only.

Overall, antibiotic treatment resulted in a decrease in the abundance of several bacterial families in faecal microbiota, in particular *Bifidobacteriaceae* and families from the *Bacteroidales* order.

Figure 1: CRE growth is promoted in antibiotic-treated human faecal microbiota that had a significant decrease in the abundance of several bacterial families, a significant increase in the concentration of nutrients, and a significant decrease in the concentration of microbial metabolites. a) Log2 fold change in bacterial families significantly decreased with antibiotics (blue) or significantly increased with antibiotics (red), relative to antibiotic-naïve faecal cultures. Wilcoxon signed rank test of log-transformed abundances with Benjamini Hochberg false discovery rate (FDR) correction, p<0.05. Bacterial families, nutrients, and metabolites that were not significantly changed with antibiotics were not plotted (white). n = 11 healthy human faecal donors. **b)** Log2 fold change in nutrients and metabolites significantly increased with antibiotics (red) and nutrients and metabolites significantly decreased with antibiotics (blue), relative to antibiotic-naïve faecal cultures. Wilcoxon signed rank test with Benjamini Hochberg FDR, p<0.05. Bacterial families, nutrients, and metabolites that were not significantly changed with antibiotics were not plotted (white). n = 11 healthy human faecal donors. **c)** TZP treatment significantly promoted the growth of CRE in human faecal microbiota. Each CRE strain was spiked into TZP-treated or water-treated faecal cultures at a concentration of 10^3 CFU/ml at the start of the experiment (represented by the black horizontal dashed line). Limit of detection represented by the grey horizontal dotted line. n = 5 healthy human faecal donors, Mann-Whitney U test of log transformed CFU/ml, ** = P ≤ 0.01.

Antibiotics enriched nutrients and depleted microbial metabolites in faecal microbiota

Antibiotic-mediated killing of gut commensals can reduce competition for nutrients, resulting in an increased availability of nutrients that could support pathogen growth. Antibiotics also decrease the concentration of microbial metabolites which may be inhibitory to pathogen growth. However, we do not know which nutrients and metabolites are impacted by treatment with different antibiotics that promote the intestinal colonisation with CRE. Therefore, we also measured the effects of the different antibiotics on nutrient availability and metabolite production in antibiotic-treated and antibiotic-naïve faecal cultures using ^{1}H-NMR spectroscopy.

Antibiotic treatment of faecal microbiota increased the concentration of several nutrients, including monosaccharides (in particular glucose), disaccharides, amino acids, uracil, and succinate (**Fig. 1b**). Antibiotics also decreased the concentration of microbial metabolites, including SCFAs, BCFAs, lactate, 5-aminovalerate, and ethanol (**Fig. 1b**). Butyrate decreased in ETP, TZP, CIP, and CRO-treated faecal microbiota but increased in MEM, IPM, CAZ, and CTX-treated faecal microbiota.

Next, we spiked CRE isolates into antibiotic-treated and antibiotic-naïve faecal cultures to determine whether the increase in nutrient availability and decrease in the concentration of microbial metabolites led to environmental conditions that promoted CRE growth. We chose to test TZP in this experiment because TZP has previously been demonstrated to promote the intestinal colonisation of CRE in humans and mice.[5,19,20] Moreover, changes in bacterial families, nutrient availability, and microbial metabolites in TZP-treated faecal microbiota were representative of changes in faecal microbiota treated with the other antibiotics that were tested in this study (**Fig. 1a,b**). We demonstrated that carbapenem-resistant *E. coli* and *K. pneumoniae* growth were significantly promoted in TZP-treated faecal cultures and suppressed in antibiotic-naïve faecal cultures (**Fig. 1c**).

Overall, antibiotic treatment of a faecal microbiota resulted in an increase in monosaccharides, disaccharides, amino acids, uracil, and succinate and a decrease in the concentration of SCFAs, BCFAs, lactate, 5-aminovalerate, and ethanol. This created a nutrient-enriched niche that supported

CRE growth and a metabolite-depleted niche that no longer inhibited CRE growth.

Bacterial taxa that were decreased with antibiotics were positively correlated with microbial

metabolites and negatively correlated with nutrients

Regularised canonical correlation analysis (rCCA) is an unsupervised approach used to analyse the

correlation structure between two multivariate datasets that have been acquired from the same

samples[21]. We used rCCA modelling to measure correlations between 16S rRNA gene sequencing

data and [1]H-NMR spectroscopy data from antibiotic-treated and antibiotic-naïve faecal cultures. The

aim of this analysis was to link changes in bacterial families to changes in nutrients and metabolites

following antibiotic treatment, to provide targets for the future development of microbiome

therapeutics to restore colonisation resistance against CRE.

Unit representation plots show sample clustering projected into the XY-variate space (in this study,

X and Y represent the 16S rRNA sequencing dataset and [1]H-NMR spectroscopy dataset)[21]. The unit

representation plots showed a separation between antibiotic-treated and antibiotic-naïve faecal

cultures along the first canonical variate, indicating antibiotic treatment caused a separation in these

groups based on their bacterial taxa and nutrient/metabolite profiles (**Fig. S1**).

Correlation circle plots show the correlation structure between features and the canonical variates[21].

In these plots, features projected in the same direction from the origin have a positive correlation,

and features projected in opposite directions from the origin have a negative correlation. Features

sitting at farther distances from the origin have stronger correlations than features sitting closer to

the origin (features with weak correlations are not plotted). In this study, correlation circle plots

showed moderate and strong negative correlations between bacterial families that were decreased

with antibiotics and monosaccharides, disaccharides, amino acids, uracil, and succinate (**Fig. 2**).

They also showed moderate and strong positive correlations between bacterial families that were

decreased with antibiotics and SCFAs, BCFAs, 5-aminovalerate, and ethanol (**Fig. 2**).

Therefore, antibiotic treatment (using antibiotics that promote CRE intestinal colonisation) resulted

in a reduction of bacterial families that are positively correlated with metabolites (indicating that these

bacterial families may produce these metabolites) and negatively correlated with nutrients (indicating

that these bacterial families may consume these nutrients). This highlights the potential role of these

gut commensals for providing colonisation resistance against CRE in antibiotic-naïve faecal

microbiota.

Figure 2. Bacterial families that were decreased in antibiotic-treated human faecal microbiota were positively correlated with microbial metabolites and negatively correlated with nutrients. rCCA model correlating 16S rRNA gene sequencing data (family level) and ^{1}H-NMR spectroscopy data (n = 11 healthy faecal donors). Correlation circle plot showing correlations between features from antibiotic-treated and antibiotic-naïve samples. Nutrients and metabolites are shown in blue and bacterial families are shown in orange. The following abbreviations are used for nutrients and metabolites: arabinose, Ara; fructose, Fru; fucose, Fuc; galactose, Gal; glucose, Glc; mannose, Man; ribose, Rib; xylose, Xyl; N-acetylglucosamine, GlcNAc; maltose, Mal; sucrose, Suc; trehalose, Tre; alanine, Ala; arginine, Arg; aspartate, Asp; glutamate, Glu; glycine, Gly; isoleucine, Ile; leucine, Leu; lysine, Lys; methionine, Met; phenylalanine, Phe; proline, Pro; threonine, Thr; tryptophan, Trp; tyrosine, Tyr; valine, Val; uracil, Ura; succinate, SA; lactate, LA; 5-aminovalerate, 5-AVA; formate, FA; acetate, AA; propionate, PA; butyrate, BA; valerate, VA; isobutyrate, iso-BA; isovalerate, iso-VA; ethanol, EtOH.

CRE can grow using nutrients that were elevated in antibiotic-treated faecal microbiota

We demonstrated that monosaccharides, disaccharides, amino acids, uracil, and succinate were

elevated with antibiotic treatment of faecal microbiota (**Fig. 1b**). Our next aim was to demonstrate

that these nutrients could support CRE growth. The growth of carbapenem-resistant *Escherichia coli*

(NDM-1), *Klebsiella pneumoniae* (NDM-1), and *Enterobacter cloacae* (NDM-1) were measured in a

mixture of these nutrients supplemented into M9 minimal medium (lacking any other carbon or

nitrogen sources) and changes in OD_{600} were measured over time. Antibiotic-mediated disruption of

the gut microbiota significantly increases oxygen availability in the gut, which may allow

Enterobacteriaceae to grow better on available carbon sources[14]. Therefore, we investigated CRE

growth in both the presence and absence of oxygen.

We showed that CRE growth was supported by the nutrient mixture, where these nutrients were

acting as the sole carbon and nitrogen sources (**Fig. 3**). For each CRE isolate growth was

significantly higher under aerobic conditions compared to anaerobic conditions.

Therefore, CRE can grow on nutrients that were elevated with antibiotics, and increased oxygen

availability promotes higher levels of CRE growth on these nutrients. This led us to conclude that the

significant increase of these nutrients following antibiotics contributes to the loss of colonisation

resistance against CRE.

Figure 3: CRE growth was supported by a mixture of nutrients that were elevated in antibiotic-treated faecal microbiota. AMiGA-predicted growth curves for NDM-1 *E. coli*, NDM-1 *K. pneumoniae*, and NDM-1 *E. cloacae* grown on M9 minimal medium supplemented with a mixture of nutrients under anaerobic or aerobic conditions. Growth of each isolate was measured with 6 replicates in 2 independent experiments. The predicted mean of growth is shown with bold lines and the predicted 95% credible intervals are shown with the shaded bands.

CRE can use monosaccharides, disaccharides, and amino acids as carbon sources to support their growth

Above we demonstrated that a mixture of monosaccharides, disaccharides, amino acids, uracil, and succinate were able to support CRE growth in a minimal medium (**Fig. 3**). However, it was not clear which specific nutrients were utilised by each CRE isolate. Therefore, we next tested which individual nutrients could act as sole carbon sources to support CRE growth. The carbon utilisation abilities of *E. coli* (NDM-1), *K. pneumoniae* (NDM-1), and *E. cloacae* (NDM-1) were demonstrated by adding a single carbon source into M9 minimal medium and monitoring changes in OD_{600} over time.

When grown anaerobically, *E. coli*, *K. pneumoniae*, and *E. cloacae* could use most monosaccharides and disaccharides as carbon sources to support their growth (**Fig. 4**), with some carbon sources supporting higher levels of growth than others (**Fig. S2**). *E. coli* could grow on arabinose, fructose, galactose, glucose, mannose, xylose, N-acetylglucosamine, and trehalose. *K. pneumoniae* could grow on arabinose, fructose, galactose, glucose, mannose, xylose, N-acetylglucosamine, maltose, sucrose, and trehalose. *E. cloacae* could grow on arabinose, fructose, galactose, glucose, mannose,

ribose, xylose, N-acetylglucosamine, maltose, sucrose, and trehalose. Carbon sources that did not support *E. coli*, *K. pneumoniae*, and *E. cloacae* growth under anaerobic conditions are shown in **Fig. S3**.

When grown aerobically, *E. coli*, *K. pneumoniae*, and *E. cloacae* could use the same carbon sources that were used anaerobically plus additional carbon sources to support their growth (**Fig. 4**). Again, some carbon sources supported higher levels of growth than others (**Fig. S2**). *E. coli* could grow on arabinose, fructose, galactose, glucose, mannose, ribose, xylose, N-acetylglucosamine, maltose, sucrose, trehalose, and alanine. *K. pneumoniae* could grow on arabinose, fructose, galactose, glucose, mannose, ribose, xylose, N-acetylglucosamine, maltose, sucrose, trehalose, alanine, glutamate, and proline. *E. cloacae* could grow on arabinose, fructose, fucose, galactose, glucose, mannose, ribose, xylose, N-acetylglucosamine, maltose, sucrose, trehalose, alanine, and glutamate. Carbon sources that did not support *E. coli*, *K. pneumoniae*, and *E. cloacae* growth under aerobic conditions are shown in **Fig. S3**.

Overall, *E. coli*, *K. pneumoniae*, and *E. cloacae* were able to utilise individual nutrients as carbon sources to support their growth. However, each isolate was able to grow to higher levels when grown aerobically versus anaerobically. Moreover, each isolate was also able to grow on a larger number of carbon sources aerobically versus anaerobically.

Figure 4: CRE growth was supported by many individual carbon sources that were elevated in antibiotic-treated faecal microbiota. AMiGA-predicted growth curves for NDM-1 *E. coli*, NDM-1 *K. pneumoniae*, and NDM-1 *E. cloacae* grown on M9 minimal medium supplemented with a single carbon source (or water as the no carbon control) under anaerobic or aerobic conditions. Growth of each isolate was measured with 6 replicates in 2-3 independent experiments. The predicted mean of growth is shown with bold lines and the predicted 95% credible intervals are shown with the shaded bands.

CRE can use amino acids, N-acetylglucosamine, and uracil as nitrogen sources to support their growth

Above we showed that most amino acids and uracil were not used as carbon sources by CRE. However, amino acids, N-acetylglucosamine, and uracil may instead act as nitrogen sources to support CRE growth. Therefore, we next tested the nitrogen utilisation abilities of *E. coli* (NDM-1), *K. pneumoniae* (NDM-1), and *E. cloacae* (NDM-1) by adding a single nitrogen source into M9 minimal medium and monitoring changes in OD_{600} over time.

When grown anaerobically, *E. coli*, *K. pneumoniae*, and *E. cloacae* could use some amino acids and N-acetylglucosamine as nitrogen sources to support their growth (**Fig. 5**), with some nitrogen sources supporting higher levels of growth than others (**Fig. S4**). *E. coli* could grow on arginine, aspartate, valine, and N-acetylglucosamine. *K. pneumoniae* and *E. cloacae* could grow on arginine, aspartate, and N-acetylglucosamine.

When grown aerobically, *E. coli*, *K. pneumoniae*, and *E. cloacae* could use the same nitrogen sources that were used anaerobically plus additional nitrogen sources to support their growth (**Fig. 5**). Again, some nitrogen sources supported higher levels of growth than others (**Fig. S4**). *E. coli* could grow on alanine, arginine, aspartate, glutamate, glycine, isoleucine, leucine, lysine, methionine, proline, threonine, tryptophan, tyrosine, valine, N-acetylglucosamine, and uracil. *K. pneumoniae* could grow on alanine, arginine, aspartate, glutamate, glycine, isoleucine, leucine, lysine, methionine, proline, and uracil. *E. cloacae* could grow on alanine, arginine, aspartate, glutamate, glycine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, threonine, N-acetylglucosamine, and uracil.

Overall, *E. coli*, *K. pneumoniae*, and *E. cloacae* were able to utilise individual nutrients as nitrogen sources to support their growth. Each isolate was able to grow to higher levels when grown aerobically versus anaerobically. Moreover, each isolate was also able to grow on a larger number of nitrogen sources aerobically versus anaerobically.

Figure 5: CRE growth was supported by individual nitrogen sources that were elevated in antibiotic-treated faecal microbiota. AMiGA-predicted growth curves for NDM-1 *E. coli*, NDM-1 *K. pneumoniae*, and NDM-1 *E. cloacae* grown on M9 minimal medium supplemented with a single nitrogen source (or water as the no nitrogen control) under anaerobic or aerobic conditions. Growth of each isolate was measured with 6 replicates in 2-3 independent experiments. The predicted mean of growth is shown with bold lines and the predicted 95% credible intervals are shown with the shaded bands. NH$_4$Cl was used as a positive control for growth as a sole nitrogen source.

CRE had an ordered preference of nutrients when provided with a mixture of nutrients

Above we demonstrated that CRE can use monosaccharides, disaccharides, and some amino acids as carbon sources and amino acids, N-acetylglucosamine, and uracil as nitrogen sources to support their growth. However, carbon utilisation assays were carried out using a single nitrogen source and nitrogen utilisation assays were carried out using a single carbon source. As we demonstrated in **Fig. 1b**, there are several carbon and nitrogen sources that are available to support CRE growth following antibiotic treatment, and it is not clear whether CRE have preferences for some carbon or nitrogen sources over others. It is also not clear whether the presence of some carbon or nitrogen sources will influence the utilisation of other carbon or nitrogen sources, as has been previously demonstrated for commensal *E. coli*[22]. Therefore, the nutrient utilisation abilities of *E. coli* (NDM-1), *K. pneumoniae* (NDM-1), and *E. cloacae* (NDM-1) were measured by adding a mixture of these nutrients into M9 minimal medium (lacking any other carbon or nitrogen sources) and monitoring changes in nutrient concentration over time by ^{1}H-NMR spectroscopy.

E. coli, *K. pneumoniae*, and *E. cloacae* were able to utilise nearly all the monosaccharides and disaccharides tested (**Fig. 6**). When grown anaerobically, *E. coli* and *E. cloacae* had high utilisation levels of all monosaccharides and disaccharides. *K. pneumoniae* had high utilisation of all monosaccharides and disaccharides except for fucose, which was not utilised. When grown aerobically, *E. coli* had high utilisation of all monosaccharides and disaccharides except for sucrose, which had moderate utilisation. *K. pneumoniae* had high utilisation of all monosaccharides and disaccharides except for fucose, which was not utilised. *E. cloacae* had high utilisation of all monosaccharides and disaccharides.

E. coli, *K. pneumoniae*, and *E. cloacae* showed variable amino acid utilisation patterns. Under anaerobic conditions, *E. coli* had high utilisation of aspartate and lysine, moderate utilisation of tryptophan, and low utilisation of the other amino acids (except for alanine or proline, which were not utilised). *K. pneumoniae* had high utilisation of aspartate and lysine, and low utilisation of the other amino acids (except for alanine, proline, and tryptophan, which were not utilised). *E. cloacae* had

high utilisation of aspartate, threonine, and arginine, and low utilisation of the other amino acids (except for alanine, proline, tryptophan, and tyrosine, which were not utilised). Under aerobic conditions, *E. coli* had high utilisation of aspartate and tryptophan, moderate utilisation of glutamate, lysine, and threonine, and low utilisation of the other amino acids (except for proline, which was not utilised). *K. pneumoniae* had high utilisation of aspartate, lysine, and threonine, moderate utilisation of glutamate, and low utilisation of the other amino acids (except for tryptophan, which was not utilised). *E. cloacae* had high utilisation of aspartate, threonine, and arginine, moderate utilisation of glutamate, and low utilisation of the other amino acids (except for proline, tryptophan, and tyrosine, which were not utilised). *E. cloacae* produced low levels of lysine both anaerobically and aerobically.

Uracil utilisation also varied by isolate under anaerobic and aerobic conditions. *E. coli* had moderate utilisation of uracil under anaerobic and aerobic conditions. *K. pneumoniae* did not use uracil under anaerobic conditions but utilised uracil at low levels under aerobic conditions. *E. cloacae* did not use uracil under anaerobic and aerobic conditions.

The rate at which nutrients were utilised was different for *E. coli*, *K. pneumoniae*, and *E. cloacae* (**Fig. S5**). Some nutrients were used more quickly than others, indicating that each CRE isolate had its own ordered preference of nutrients. Moreover, aerobically incubated CRE cultures utilised most nutrients more rapidly than anaerobically incubated CRE cultures.

Succinate was produced by all CRE isolates under both anaerobic and aerobic conditions. Other metabolites produced by the CRE isolates included acetate, ethanol, and formate (**Fig. S6**).

In summary, these results led us to conclude that *E. coli*, *K. pneumoniae*, and *E. cloacae* had an ordered preference for nutrients and that nutrient utilisation was different in the presence and absence of oxygen.

Figure 6: CRE isolates had an ordered preference of nutrients when cultured in a mixture of nutrients that were elevated in antibiotic-treated faecal microbiota. Percent of nutrients remaining in M9 minimal medium supplemented with a mixture of nutrients, inoculated with NDM-1 *E. coli*, NDM-1 *K. pneumoniae*, or NDM-1 *E. cloacae*, and incubated under anaerobic or aerobic conditions. Nutrient concentrations were measured by ^{1}H-NMR spectroscopy. Two-way mixed ANOVA followed by pairwise comparisons with Bonferroni correction. * = P ≤ 0.05, ** = P ≤ 0.01, *** = P ≤ 0.001, **** = P ≤ 0.0001, n=3 replicates for each CRE isolate and n=4 replicates for the sterile media controls.

Carbapenem-resistant* E. coli *utilised nutrients that were elevated in the faecal microbiota from TZP-treated mice

Next, we used a mouse model of carbapenem-resistant *E. coli* intestinal colonisation to measure changes in nutrient availability in mouse faeces following TZP treatment and NDM-1 *E. coli* colonisation. Mice were administered TZP or saline for 5 days, and NDM-1 *E. coli* was orally administered to the mice on the third day of TZP/saline administration (**Fig. 7a**). TZP-treated mice had significantly higher *E. coli* counts in their faeces compared to saline-treated mice, demonstrating that antibiotics disrupted colonisation resistance against carbapenem-resistant *E. coli* (**Fig. 7b**).

Faecal samples were collected from mice before TZP administration, after TZP administration but before *E. coli* administration, and after TZP and *E. coli* administration. Changes in nutrients and metabolites were measured by ^{1}H-NMR spectroscopy. The faeces of TZP-treated mice had significantly increased concentrations of nutrients compared to faeces collected prior to TZP treatment, including increases in monosaccharides (arabinose, fructose, galactose, glucose, xylose, N-acetylglucosamine), disaccharides (sucrose, trehalose), and amino acids (arginine, aspartate, glutamate, isoleucine, leucine, lysine, phenylalanine, proline, threonine, tryptophan, tyrosine, valine) (**Fig. 7c**, heatmap on the left). TZP treatment also decreased the concentration of microbial metabolites, including acetate, propionate, isobutyrate, ethanol, and lactate. Following colonisation with NDM-1 *E. coli* there was a significant decrease in nutrients that we demonstrated can be utilised by *E. coli* to support its growth, including monosaccharides (arabinose, fructose, galactose, glucose, xylose, N-acetylglucosamine), disaccharides (sucrose, trehalose), amino acids (alanine, arginine, aspartate, glutamate, lysine, tryptophan, tyrosine), and uracil (**Fig. 7c**, heatmap on the right).

Overall, antibiotic treatment of faecal microbiota from mice resulted in an increase in the concentration of nutrients and a decrease in the concentration of microbial metabolites, supporting observations from our *ex vivo* human faecal culture experiments (**Fig. 1**). Colonisation of the mouse intestine with NDM-1 *E. coli* resulted in a decrease in nutrients that we demonstrated can be utilised by *E. coli in vitro* (**Figs. 4, 5, 6**), supporting our hypothesis that *E. coli* consumes these nutrients to

support its growth.

Figure 7: Nutrients that were elevated with TZP treatment decreased with *E. coli* colonisation in faeces from a mouse model of intestinal CRE colonisation. a) Experimental design. Mice were administered TZP or saline for 5 days and were fed by oral gavage with NDM-1 *E. coli* on the third day of TZP/saline administration. Faeces were collected before TZP administration, after TZP administration but before *E. coli* administration, and after TZP and *E. coli* administration. **b)** NDM-1 *E. coli* counts from mouse faecal pellets following *E. coli* administration (at the end of the TZP dosing period). The box excludes the upper and lower 25% (quartiles) of data, and the lines go to maximum and minimum values. Unpaired t-test of log10-transformed counts. n=5 mice per group, **** = P ≤ 0.0001. **c)** Log2 fold change in nutrients and metabolites significantly changed with interventions. For comparison of post-TZP treated faeces relative to pre-TZP treated faeces (heatmap on the left), nutrients and metabolites that were significantly increased with TZP are shown in red, and nutrients and metabolites that were significantly decreased with TZP are shown in blue (n=9 mice per group). For comparison of TZP-treated faeces post-*E. coli* colonisation relative to TZP-treated faeces pre-*E. coli* colonisation (heatmap on the right), nutrients and metabolites that were significantly decreased with *E. coli* colonisation are shown in blue (n=4 mice per group). Paired t-test with Benjamini Hochberg FDR, p<0.05. Nutrients and metabolites that were not significantly different were not plotted (white). LOD = limit of detection.

CRE growth was inhibited by microbial metabolites that were depleted in antibiotic-treated

faecal microbiota

Microbial metabolites have previously been demonstrated to inhibit the growth of intestinal

pathogens[7,16]. Nutrients are metabolised to produce metabolites, and therefore increased nutrient availability is linked to reduced metabolite production. We demonstrated that microbial metabolites were decreased in antibiotic-treated faecal microbiota (**Fig. 1b**). Therefore, we next tested whether microbial metabolites (that were decreased with antibiotics) were capable of inhibiting CRE growth.

First, we measured the concentration of microbial metabolites in faeces from 12 healthy human donors to determine the concentration of these metabolites in antibiotic-naïve faecal microbiota (**Table S1**). We used these measurements to define a minimum, average, and maximum concentration to test in the metabolite inhibition assays.

First, we tested whether a mixture of metabolites that were decreased with antibiotics (formate, acetate, propionate, butyrate, valerate, isobutyrate, isovalerate, lactate, 5-aminovalerate, and ethanol) could directly inhibit CRE growth in an inhibition assay. LB was supplemented with the metabolite mixture, where each metabolite was added to mimic the average concentration that was measured in healthy human faeces. As pH has been demonstrated to influence the ability of SCFAs to inhibit pathogen growth, the metabolite mixture was tested at three pH conditions mimicking the pH found in the healthy large intestine: 6, 6.5, and 7[16,23]. We showed that the metabolite mixture inhibited the growth of *E. coli*, *K. pneumoniae*, and *E. cloacae* at pH 6, 6.5, and 7 (**Fig. 8a**). However, inhibition was higher at pH 6 and 6.5 compared to pH 7.

Next, we measured CRE growth in the presence of each individual metabolite at three different concentrations: the minimum, average, and maximum concentrations measured in healthy faeces. These experiments were performed at pH 6.5 as this pH was representative of the large intestine and showed strong inhibition in the mixed metabolite experiment (as shown in **Fig. 8a**). *E. coli* was inhibited by acetate at low, average, and high concentrations, by propionate, valerate, and ethanol at average and high concentrations, and by butyrate at high concentrations (**Fig. 8b**). *K. pneumoniae* was inhibited by butyrate at low, average, and high concentrations, and by acetate, propionate, valerate, isobutyrate, and isovalerate at average and high concentrations (**Fig. 8c**). *E. cloacae* was inhibited by butyrate and valerate at low, average, and high concentrations, by acetate, propionate,

isobutyrate, and isovalerate at average and high concentrations, and by ethanol at the high

concentration (**Fig. 8d**).

In summary, we demonstrated that specific microbial metabolites (that were decreased following

antibiotic treatment of faecal microbiota) were inhibitory towards CRE isolates. This led us to

conclude that the significant decrease of these metabolites following antibiotics contributes to the

loss of colonisation resistance against CRE.

Figure 8: CRE growth was inhibited by microbial metabolites that were decreased in antibiotic-treated faecal microbiota. a) CRE growth was inhibited by a mixture of metabolites that were decreased in antibiotic-treated faecal microbiota. NDM-1 *E. coli*, NDM-1 *K. pneumoniae*, or NDM-1 *E. cloacae* were grown in LB supplemented with a metabolite mixture (mimicking average human faecal concentrations) or unsupplemented ("no metabolite" control) at pH 6, 6.5, or 7. Growth of each isolate was measured with 6 replicates in 2 independent experiments. Unpaired t-test comparing metabolite mixture to the no metabolite control at the corresponding pH. **b-d)** CRE growth was inhibited by individual metabolites in a dose-dependent manner. Growth of NDM-1 *E. coli*, NDM-1 *K. pneumoniae*, or NDM-1 *E. cloacae* in LB supplemented with an individual metabolite at low, average, or high concentrations (mimicking human faecal concentrations) or unsupplemented ("no metabolite" control) at pH 6.5. Growth of each isolate was measured with 6 replicates in 2 independent experiments. One-way ANOVA followed by Dunn's multiple comparison test (each metabolite was compared to the no metabolite control). * = P ≤ 0.05, ** = P ≤ 0.01, *** = P ≤ 0.001, **** = P ≤ 0.0001.

Carbapenem-resistant E. coli *growth was inhibited by a mixture of microbial metabolites in a mouse model of CRE intestinal colonisation*

Next, we administered a mixture of inhibitory metabolites as an intervention in a mouse model of intestinal CRE colonisation to determine whether these metabolites could inhibit the growth of NDM-1 *E. coli in vivo*. TZP-treated mice colonised with NDM-1 *E. coli* were administered a mixture of metabolites (acetate, propionate, butyrate, and valerate) or PBS (as a vehicle control; **Fig. 9a**). We chose to test these metabolites in mice as they were inhibitory against all three CRE isolates when tested individually (**Fig. 8**) and when tested together as a mixture (**Fig. S7**). Each metabolite was administered to the mice in their triglyceride form (i.e. as a mixture of glycerol triacetate, glycerol tripropionate, glycerol tributyrate, and glycerol trivalerate) to avoid rapid absorption of SCFAs following oral gavage and to obtain sufficient concentrations of these SCFAs in the gastrointestinal tract[7]. The triglyceride forms of these SCFAs are hydrolysed by lipases to release acetate, propionate, butyrate, and valerate, respectively, in the intestine[24].

Prior to the administration of the metabolite mixture or PBS to mice, there was no significant difference in NDM-1 *E. coli* counts in the faeces from each group (p>0.9999). However, there was a significant decrease in NDM-1 *E. coli* counts in the faeces of metabolite-treated mice compared to PBS-treated mice (**Fig. 9b**, p=0.0397). After 3 doses of the metabolite mixture or PBS, metabolite-treated mice had an average of 87% less *E. coli* CFU per gram of faeces compared to PBS-treated mice.

Overall, we demonstrated that metabolites that were decreased in antibiotic-treated human faecal microbiota (acetate, propionate, butyrate, and valerate) were able to inhibit NDM-1 *E. coli* in a mouse model of intestinal CRE colonisation, highlighting their important role in providing colonisation resistance against CRE.

Figure 9: Microbial metabolites significantly decreased carbapenem-resistant *E. coli* growth in faeces from a mouse model of intestinal CRE colonisation. a) Experimental design. Mice were administered TZP for 5 days and were orally administered NDM-1 *E. coli* on the third day of TZP administration. Mice were then fed by oral gavage with a mixture of metabolites (glycerol triacetate, glycerol tripropionate, glycerol tributyrate, and glycerol trivalerate) or PBS for 3 days. **b)** NDM-1 *E. coli* counts were quantified from mouse faecal pellets just before metabolite mixture or PBS administration (pre-intervention) and at the end of metabolite mixture or PBS administration (post-intervention). The box excludes the upper and lower 25% (quartiles) of data, and the lines go to maximum and minimum values. Mann-Whitney U test of log10-transformed counts. n=5 mice per group, * = P ≤ 0.05.

DISCUSSION

In this study we demonstrated that antibiotic-mediated disruption of the gut microbiota (using antibiotics that promote CRE intestinal colonisation) led to the development of a nutrient-enriched and metabolite-depleted niche that was favourable for CRE growth. We showed that CRE can use these elevated nutrients as carbon and nitrogen sources to support their growth, where CRE show an order of preference for specific nutrients. CRE growth was higher on these nutrients in an oxygenated environment, which has been associated with antibiotic treatment. We also showed that specific microbial metabolites that were decreased with antibiotics were able to inhibit CRE growth.

We showed that healthy faecal microbiota treated with antibiotics that promote CRE intestinal colonisation resulted in a decrease of several bacterial families, in particular *Bifidobacteriaceae* (decreased with all antibiotics tested), families belonging to *Bacteroidales* (decreased with carbapenems, TZP, and CIP), and *Coriobacteriaceae* (decreased with carbapenems). Our results are supported by previous studies in healthy humans which also demonstrated a decrease in *Bifidobacterium* and *Bacteroides* with these antibiotics[25-30]. Our results are also supported by a recent study which demonstrated that CRE-positive patients had lower amounts of *Bacteroides dorei*, *Bifidobacterium bifidum*, *Bifidobacterium pseudocatenulatum*, and *Collinsella aerofaciens* in their faeces compared to CRE-negative patients[31]. Moreover, a recent study demonstrated that *Bacteroidota* and *Actinobacteria* (including *Bifidobacterium*) engraft very highly within the intestine following faecal microbiota transplantation[32]. This highlights the possible role of *Bifidobacterium* and *Bacteroides* in decolonising CRE-positive patients following faecal microbiota transplantation.

Polysaccharides, proteins, and mucins are major food sources for the gut microbiota in the large intestine. Primary degraders break down these complex substrates, resulting in the extracellular release of breakdown products (oligosaccharides, disaccharides, monosaccharides, peptides, and amino acids)[15]. These simple compounds can support the growth of secondary degraders. *Bacteroides* are important primary degraders (but can also degrade simple compounds) and *Bifidobacterium* are important secondary degraders in the gut[33]. Therefore, this supports the observation that decreases in the abundance of these bacteria following antibiotics results in an enrichment of simple nutrients.

The nutrient-niche hypothesis predicts that expansion of *Enterobacteriaceae* in an antibiotic-disrupted gut microbiota is due to an increase in the availability of nutrients, where competition for these nutrients would normally restrict *Enterobacteriaceae* growth[14]. We demonstrated that antibiotics (that promote susceptibility to CRE) increased the availability of nutrients that can support CRE expansion within the gut. The monosaccharides, disaccharides, and amino acids of interest in this study were derived from the breakdown of complex polysaccharides, proteins, and mucin, including starch (glucose, maltose, trehalose)[34-37], inulin (fructose, glucose, sucrose)[38], pectin

(glucose, arabinose, galactose, xylose, mannose, fucose)[39], xylan (xylose, arabinose)[40], arabinogalactan (arabinose, galactose)[41], mucin (N-acetylglucosamine, galactose, fucose, amino acids)[42], peptone (amino acids)[43,44], and yeast extract (amino acids, ribose, uracil)[45]. Our results led us to conclude that CRE have a growth advantage in antibiotic-perturbed gut microbiota by acting as secondary fermenters of monosaccharides, disaccharides, and amino acids, highlighting the important role of microbial cross-feeding in antibiotic-disrupted gut microbiota.

Previous studies demonstrated that antibiotics could enrich nutrients to support the growth of intestinal pathogens. Cefoperazone (which promoted *Clostridioides difficile* intestinal colonisation) increased nutrients (mannitol, fructose, sorbitol, raffinose, stachyose, and some amino acids) and decreased metabolites (valerate and isovalerate) in the caecal microbiota of mice[46]. *In vitro* experiments demonstrated that these nutrients supported *C. difficile* growth. Another study demonstrated that sialic acid was increased in the caecal microbiota of mice treated with antibiotics that promote susceptibility to *Salmonella enterica* serovar Typhimurium or *C. difficile*[10]. *S.* Typhimurium and *C. difficile* mutants that were unable to utilise sialic acid showed impaired growth within the intestine. In this study we found an enrichment of nutrients associated with polysaccharides, mucin, and proteins, indicating that antibiotics cause a disruption of secondary fermentation in the gut, where enrichment of these simple nutrients can support CRE growth.

Oxygen is a limiting resource for *Enterobacteriaceae* in the gut, and oxygenation of the gut that occurs following antibiotics has been hypothesised to drive the expansion of *Enterobacteriaceae*[14]. Our results demonstrate that an increase in oxygen availability results in an increase in CRE growth on nutrients that were enriched with antibiotics (compared to growth under anaerobic conditions). This observation is supported by a recent longitudinal shotgun metagenomics sequencing analysis of stool from CRE-positive patients which showed that aerobic respiration and oxidative phosphorylation pathways (e.g. the pentose phosphate pathway) were enriched during CRE colonisation (where antibiotic usage was common)[31]. Rivera-Chávez and colleagues demonstrated that antibiotic treatment led to increased gut oxygenation that resulted in the aerobic expansion of *S.* Typhimurium, demonstrating the advantage that oxygen can provide to facultatively anaerobic

pathogens[47]. Overall, our work supports the hypothesis that CRE growth is enhanced on carbon and

nitrogen sources (that become available due to antibiotic treatment) in the presence of oxygen.

The metabolism of monosaccharides and disaccharides produces SCFAs and the metabolism of

amino acids produces SCFAs and BCFAs[15,48]. We demonstrated that antibiotic treatment of the

faecal microbiota results in the disruption of nutrient metabolism, leading to the development of a

metabolite depleted environment. We showed that formate, acetate, propionate, butyrate, valerate,

isobutyrate, isovalerate, lactate, 5-aminovalerate, and ethanol were decreased by antibiotics.

Bifidobacterium, *Bacteroides*, and *Collinsella* (taxa that were decreased with antibiotics) are known

to produce many of these metabolites. *Bifidobacterium* produces acetate, lactate, ethanol, and

formate[15,49]. *Bacteroides* produces acetate, propionate, lactate, formate, ethanol, isobutyrate, and

isovalerate[15,50]. *Collinsella* produces ethanol, formate, and lactate[51].

We showed that a mixture of these 10 metabolites was inhibitory against CRE at pH values

representative of the human large intestine (pH 6, 6.5, and 7). However, the metabolite mixture was

more inhibitory at pH 6 and 6.5 compared to pH 7. Our work is supported by a previous study by

Sorbara and colleagues that demonstrated that acetate, propionate, and butyrate were decreased

in caecal content from mice treated with ampicillin[16]. They showed that a mixture of acetate,

propionate, and butyrate inhibited CRE growth at pH 5.75 (the pH of the mouse cecum) but not pH

7. It is possible that we observed higher levels of inhibition in the metabolite mixture at pH 7 due to

the inclusion of additional inhibitory metabolites in the mixture. This is supported by previous

observations that valerate can inhibit the growth of *C. difficile* at pH 7[7].

We demonstrated that 7 microbial metabolites (acetate, propionate, butyrate, valerate, isobutyrate,

isovalerate, and ethanol) inhibit CRE growth. Future research is required to solve the mechanisms

that these inhibitory metabolites use to inhibit intestinal CRE colonisation. It is unclear whether these

metabolites may act using one mechanism of inhibition or whether they may mediate inhibition by

multiple different mechanisms. We saw differences in the metabolites that inhibited *E. coli*, *K.*

pneumoniae, and *E. cloacae* growth, and it is not clear whether there are differences in the

mechanisms in these three different species. It is also unclear whether these metabolites may have synergistic inhibitory effects, or whether these mechanisms may be impacted by different gut environmental conditions. Future studies are needed to investigate how these metabolites impact pathogens, other members of the gut microbiota, and the host to provide colonisation resistance against CRE.

In this study we investigated the impact of microbial metabolites on the growth of CRE isolates. However, it is important to further investigate the impact of microbial metabolites on the host, as this may also impact the development of invasive CRE infections. For example, both butyrate and valerate have been demonstrated to improve gut barrier integrity[52,53]. Therefore, a depletion of these metabolites following antibiotic treatment may promote the translocation of CRE from the gut to the bloodstream.

In summary, we provide evidence that antibiotics create an intestinal niche that supports CRE expansion by enriching for nutrients that support CRE growth and by depleting metabolites that are inhibitory towards CRE growth. Understanding how antibiotics disrupt colonisation resistance against CRE is critical for the rational design of microbiome therapeutics that prevent or treat CRE intestinal colonisation. These microbiome therapeutics could reduce the risk of patients developing invasive CRE infections, reduce the recurrence of invasive CRE infections in chronically colonised patients, and reduce the spread of CRE to susceptible patients. A second-generation microbiome therapeutic may be composed of gut commensals that can outcompete CRE for nutrients and convert them into inhibitory metabolites. A third-generation microbiome therapeutic may be composed of a metabolite or mixture of metabolites that inhibit CRE growth, which could be administered to CRE colonised patients while they are receiving antibiotic treatment.

In the short term, our results could be used to help reduce the risk of patients becoming colonised with CRE. Our data could guide the prescription of antibiotics whereby clinicians avoid prescribing antibiotics that elevate specific nutrients that support CRE growth and deplete specific metabolites that inhibit CRE growth. Our data could also be used to identify patients that are at particular risk for

CRE intestinal colonisation by screening faecal samples for these nutrients and metabolites.

MATERIALS AND METHODS

Carbapenem-resistant **Enterobacteriaceae** *isolates:*

New Delhi Metallo-β-lactamase 1 (NDM-1) *E. coli*, NDM-1 *K. pneumoniae*, and NDM-1 *E. cloacae*

were isolated from patients with intestinal CRE colonisation from the Imperial College Healthcare

NHS Trust (London, UK) via rectal swab (with approval from the London - Queen Square Research

Ethics Committee, 19/LO/0112). The species were classified by matrix assisted laser desorption

ionisation-time of flight mass spectrometry and later confirmed with 16S rRNA gene sequencing and

whole genome sequencing. CRE status was determined using X-pert Carba-R PCR detection

(Cepheid, Sunnyvale, CA)[54]. Genome sequencing data for these isolates have been deposited into

the European Nucleotide Archive under the project accession number PRJEB60914 and strains are

available upon request.

We also tested NDM-4 *E. coli* ST167 (NCTC 14333), OXA-48 *E. coli* ST410 (NCTC 14324), KPC-3

K. pneumoniae ST258 (NCTC 13438), and KPC-3 *K. pneumoniae* ST11 (NCTC 14327) as these are

prevalent clonal groups that are responsible for the rapid emergence of CRE clinical isolates

worldwide.[55]

Ethical approval:

Ethical approval was received by the London - Queen Square Research Ethics Committee

(19/LO/0112) and the South Central - Oxford C Research Ethics Committee (16/SC/0021 and

20/SC/0389) to collect faecal samples from human donors. Faecal donors were healthy human

donors aged between 18-65 years old and had not received antibiotic treatment in the 6+ months

prior to donation. Informed consent was obtained from human participants.

Faecal culture experiments:

We performed *ex vivo* faecal culture experiments to measure the effects of eight broad-spectrum antibiotics on the faecal microbiota collected from 11 healthy human donors (6 males, 5 females). Female faecal donors were 33.2 ± 16.0 years old and male faecal donors were 29.7 ± 6.4 years old. Faecal culture experiments were performed under anaerobic conditions using a standardised distal gut growth medium mimicking nutrients found in the distal gut, including polysaccharides, proteins, and mucin[56]. For each donor, fresh faeces were inoculated into the distal gut growth medium at a 2% (w/v) concentration. Faecal cultures were supplemented with water (antibiotic-naïve control) or one of the following antibiotics (at concentrations measured in faeces): 2 µg/ml MEM, 2 µg/ml IPM, 37 µg/ml ETP, 139 µg/ml TZP, 152 µg/ml CRO, 22 µg/ml CAZ, 2 µg/ml CTX, 139 µg/ml CIP[18,26-28,57-60]. These antibiotics are known to promote the intestinal colonisation with CRE and are used frequently in routine clinical practice[5]. Faecal cultures were run in triplicate and incubated anaerobically at 37°C for 24 hours. Samples were analysed by proton nuclear magnetic resonance (^{1}H-NMR) spectroscopy (for nutrient and metabolite profiling) and 16S rRNA gene sequencing with 16S rRNA gene qPCR (for bacterial composition and biomass).

A second set of *ex vivo* faecal culture experiments was performed to measure the CRE growth in TZP-treated and antibiotic-naïve faecal microbiota. Faecal samples were donated from 5 healthy human donors (2 males, 3 females), and these faecal samples were separate from the faecal samples collected above. Female faecal donors were 31.3 ± 5.5 years old and male faecal donors were 31.0 ± 9.9 years old. For each donor, fresh faeces were inoculated into the distal gut growth medium supplemented with TZP (139 µg/ml) or water and spiked with 10^3 CFU/ml one of the following CRE isolates: NDM-1 *E. coli*, NDM-4 *E. coli* (ST167), OXA-48 *E. coli* (ST410), KPC-3 *K. pneumoniae* (ST258), or KPC-3 *K. pneumoniae* (ST11). Faecal cultures were run in duplicate and incubated anaerobically at 37°C for 24 hours. Samples were plated on Brilliance CRE agar to quantify CRE growth.

16S rRNA gene sequencing and 16S rRNA gene qPCR:

DNA was extracted from 250 µl of faecal culture sample using the DNeasy PowerLyzer PowerSoil

Kit (Qiagen, Germany) according to manufacturer's instructions, with the addition of a bead beating

step for 3 minutes at speed 8 in a Fast-Prep 24 bead beater (MP Biomedicals, USA). DNA was

stored at -80°C until it was ready to be used.

Illumina's 16S metagenomic sequencing library preparation protocol was used to generate sample

libraries amplifying the V1-V2 regions of the 16S rRNA gene, as previously described[61]. The

SequalPrep Normalisation Plate Kit (Life Technologies, UK) was used to clean up and normalise the

index PCR reactions. The NEBNext Library Quant Kit for Illumina (New England Biolabs, UK) was

used to quantify sample libraries. Sample libraries were sequenced on an Illumina MiSeq platform

(Illumina Inc., UK) using the MiSeq Reagent Kit v3 (Illumina) and paired-end 300bp chemistry. 16S

rRNA gene qPCR was performed to quantify the bacterial biomass as previously described[7].

16S rRNA gene sequencing data were imported into R and processed using the standard DADA2

pipeline (version 1.18.0)[62]. The SILVA bacterial database version 138.1 was used to classify the

sequence variants. 16S rRNA gene sequencing data was expressed as absolute abundances using

the 16S rRNA gene qPCR data, as previously described[7].

¹H-NMR spectroscopy:

Supernatants were prepared from human faeces by vortexing 300 mg of faeces in 900 µl of high-

performance liquid chromatography-grade water for 10 minutes at 3000 rpm followed by

centrifugation at 17,000 × g for 10 minutes at 4°C. Supernatants were prepared from faecal culture

samples by centrifugation at 17,000 × g for 10 minutes at 4°C. Supernatants were prepared from

mouse faeces by vortexing 20 mg of faeces in 560 µl of high-performance liquid chromatography-

grade water for 10 minutes at 3000 rpm followed by centrifugation at 17,000 × g for 10 minutes at

4°C.

[1]H-NMR spectroscopy was performed as previously described[7]. Briefly, sample supernatants were

randomised and defrosted at room temperature for 1 hour. Sample supernatants were centrifuged

for 10 min at 17,000 × g and 4°C and the pellet was discarded. For human faeces and faecal batch

culture samples, 400 µl of supernatant was added to 250 µl NMR buffer (28.85 g Na_2HPO_4, 5.25 g

NaH_2PO_4, 1 mM 3-(Trimethylsilyl)propionic-2,2,3,3-d_4 acid sodium salt (TSP), 3 mM NaN_3, deuterium

oxide to 1 L, pH 7.4). For mouse faecal samples 540 µl of supernatant was added to 60 µl NMR

buffer (20.4 g KH_2PO_4, 5.8 mM TSP, 2 mM NaN_3, deuterium oxide to 100 mL, pH 7.4). Each 5 mm

NMR tube was filled with 600 µl of sample and loaded into the SampleJet system.

[1]H-NMR spectroscopy was performed using the Bruker AVANCE III HD 800 MHz spectrometer

(Bruker Bio-Spin, Rheinstetten, Germany) using a 5mm CPTCI 1H-13C/15N/D Z-gradient cryoprobe.

1D spectra (a standard NOESYGPPR1D pulse sequence (RD-90°-t1-90°-tm-90°-ACQ)) as well as

2D spectra (standard 2D JRESGPPRQF pulse sequence) were acquired. A recycle delay of 4

seconds and mixing time of 100 ms was used. The 90° pulse length was approximately 10 µs. For

the human faecal samples and faecal batch culture samples 32 scans were recorded. For the mouse

faecal samples 128 scans were recorded.

NMR spectra underwent auto processing, which includes exponential window function (line

broadened by 0.3 Hz), auto phasing (zero-order), auto baseline correction, and auto referencing.

NMR spectra were imported into MATLAB r2019b (The Mathworks, USA) using a custom script.

Spectra were aligned and the water (4.6 – 5 ppm) and TSP (-0.2 – 0.2 ppm) peaks were cut.

Compounds were identified by comparison to spectral databases in Chenomx NMR Suite (Chenomx,

Canada). Representative peaks were integrated for quantification of compounds. A signal-to-noise

ratio cut-off value of 3:1 was used to define the limit of detection for peak detection.[63] The minimal

signal-to-noise ratio was 6.1:1 for the 32 scan human faecal dataset, 3.8:1 for the 32 scan faecal

batch culture dataset, and 4.0:1 for the 123 scan mouse faecal dataset.

mixOmics analysis:

rCCA modelling was used to correlate 16S rRNA gene sequencing data (family level) with ^{1}H-NMR

data for faecal cultures samples using the *mixOmics* library within R, as previously described[7,21]. In

unit representation plots each point represents a single faecal culture sample that was projected into

the XY-variate space. In the correlation circle plots features projected in the same direction from the

origin have a positive correlation, and features projected in opposite directions from the origin have

a negative correlation. Features sitting at farther distances from the origin have stronger correlations

than features sitting closer to the origin.

Carbon and nitrogen utilisation assays:

Compounds that were enriched in antibiotic-treated faecal microbiota were tested as nutrients in

growth assays to determine whether they could act as carbon or nitrogen sources to support CRE

growth. Growth of CRE isolates on sole carbon or nitrogen sources were tested by inoculating each

CRE isolate into a minimal medium supplemented with the nutrient of interest. First, CRE isolates

were passaged from frozen glycerol stocks onto R2A plates. Cells were suspended in M9 minimal

media to a turbidity of 42% using a Turbidimeter (Biolog Inc., Hayward, CA), then further diluted 1 in

6 to inoculate the carbon or nitrogen utilisation assays. M9 minimal media contained the following

(at final concentrations): 22 mM KH_2PO_4, 42 mM Na_2HPO_4, 9 mM NaCl, 0.49 mM $MgSO_4 \cdot 7H_2O$,

0.09 mM $CaCl_2$, 0.011 mM $FeSO_4 \cdot 7H_2O$. For the carbon assays the M9 minimal medium was also

supplemented with 0.1% (w/v) NH_4Cl (as the nitrogen source) and the carbon sources were

individually tested at 0.5% (w/v) concentration. For the nitrogen assays the M9 minimal medium was

also supplemented with 0.5% (w/v) glucose (as the carbon source) and the nitrogen sources were

individually tested at 0.1% (w/v) concentration. Cultures were incubated anaerobically and

aerobically in a plate reader and optical density (OD_{600}) readings were collected every 15 minutes

for 24 hours.

Mixed nutrient preference assays:

CRE isolates were passaged from frozen glycerol stocks on R2A plates. Cells were suspended in M9 minimal media to a turbidity of 42% using a Turbidimeter, and then further diluted 1 in 6 to inoculate the nutrient preference assays. M9 minimal media was supplemented with 0.015% (w/v) of each of the following nutrients (as the sole carbon and nitrogen sources): L-arabinose, D-fructose, L-fucose, D-galactose, D-glucose, D-mannose, D-ribose, D-xylose, N-acetylglucosamine, D-maltose, sucrose, D-trehalose, L-alanine, L-arginine, L-aspartate, L-glutamate, L-glycine, L-isoleucine, L-leucine, L-lysine, L-methionine, L-phenylalanine, L-proline, L-threonine, L-tryptophan, L-tyrosine, L-valine, uracil, and succinate. Cultures were incubated anaerobically and aerobically in a plate reader and OD_{600} readings were collected every 15 minutes for 24 hours. Cultures were also incubated anaerobically and aerobically for 24 h and samples were collected at 0, 4, 8, and 24 hours post-inoculation for ^{1}H-NMR spectroscopy. The starting nutrient concentration was measured by the integration of a representative peak in the NMR spectrum from samples collected at 0 hours and this concentration was set to 100% to enable measurements of percent nutrient remaining at 4, 8, and 24 hours.

Analysis of Microbial Growth Assays (AMiGA):

Growth curves were analysed in Python (v3.6.5) using the AMiGA software[64]. Growth of each isolate was measured with 6 replicates from 2-3 independent experiments. Growth curves were normalised to the no substrate control using the subtraction method. AMiGA performed Gaussian Process regression to test differential growth between the substrate and the no substrate control. The functional difference in the OD_{600} (and its credible interval) was also computed between the substrate and the no substrate control.

Metabolite measurements from healthy human faeces:

The concentration of 10 microbial metabolites were measured from the faeces of 12 healthy human donors (5 males, 7 females) by ^{1}H-NMR spectroscopy. This was a separate set of faecal samples than the faecal samples that were used to inoculate the faecal culture experiments. Female faecal

donors were 30.9 ± 12.1 years old and male donors were 27.8 ± 5.3 years old. Metabolite

concentrations were measured from NMR spectra using the Chenomx NMR Suite (v8.6). **Table S2**

shows the approximate minimum, average, and maximum concentrations of each metabolite that

was tested in the metabolite inhibition assays, which were based on the measurements of the

microbial metabolites from the healthy donor faeces.

Mixed metabolite inhibition assay:

Microbial metabolites that were present in antibiotic-naïve faecal microbiota but depleted in

antibiotic-treated faecal microbiota were tested as potential inhibitory compounds in growth assays.

We tested whether a mixture of microbial metabolites (that were decreased in antibiotic-treated

faecal microbiota) would be able to inhibit the growth of each CRE isolate. The metabolite mixture

was composed of formate, acetate, propionate, butyrate, valerate, isobutyrate, isovalerate, lactate,

5-aminovalerate, and ethanol. Each metabolite was added to the mixture at a concentration that

mimicked the average concentration measured in healthy human faecal samples (**Table S2**). Luria

broth (LB) was supplemented with the metabolite mixture and compared to a no metabolite control.

Overnight cultures of each CRE isolate were inoculated into the supplemented LB at 10^3 CFU/ml.

The assay was run at pH 6, 6.5, and 7 to mimic the pH range found in healthy faeces[23,65]. Cultures

were incubated anaerobically for 16 hours at 37°C. Growth was measured at OD_{600} after 16 hours

of incubation.

We also measured the growth of CRE isolates in LB supplemented with a mixture of acetate,

propionate, butyrate, and valerate. Each metabolite was added to the mixture at a concentration that

mimicked the average concentration measured in healthy human faecal samples (**Table S2**). LB was

supplemented with the metabolite mixture and compared to a no metabolite control. The assay was

run at pH 6.5. Overnight cultures of each CRE isolate were inoculated into the supplemented LB at

10^3 CFU/ml. Cultures were incubated anaerobically for 16 hours at 37°C. Growth was measured at

OD_{600} after 16 hours of incubation.

Individual metabolite inhibition assay:

We next tested whether individual microbial metabolites (that were decreased in antibiotic-treated faecal microbiota) would be able to inhibit the growth of each CRE isolate. The individual metabolites tested included formate, acetate, propionate, butyrate, valerate, isobutyrate, isovalerate, lactate, 5-aminovalerate, and ethanol. Each metabolite was tested at three concentrations that mimicked the low, average, and high concentrations measured in healthy human faecal samples (**Table S2**). LB was supplemented with each metabolite and compared to a no metabolite control. Overnight cultures of each CRE isolate were inoculated into the supplemented LB at 10^3 CFU/ml. The assay was run at pH 6.5 as this was representative of the distal gut and showed inhibition in the mixed metabolite assay. Cultures were incubated anaerobically for 16 hours at 37°C. Growth was measured at OD_{600} after 16 hours of incubation.

Mouse model of intestinal colonisation with carbapenem-resistant E. coli*:*

Ethical approval for mouse experiments was received from the Imperial College London Animal Welfare and Ethical Review Body (PF93C158E) and performed under the authority of the UK Home Office, as described in the Animals (Scientific Procedures) Act 1986. We abided by standards outlined in the Animal Research: Reporting of In Vivo Experiments (ARRIVE) guidelines.

Eight- to ten-week-old female wild-type C57BL/6 mice (purchased from Envigo, Huntingdon, UK) were acclimatised for 1 week prior to the start of the experiment. Each test group contained 5 mice that were housed 5 per cage (in individually ventilated cages). Mice were provided autoclaved food (RM1, Special Diet Services, Essex, UK), water (provided ad libitum), and bedding (Aspen chip 2 bedding, NEPCO, Warrensburg, New York). Mice were maintained at 20–22°C in a 12-hour light and 12-hour dark cycle (all interventions were performed during the light cycle).

We performed a mouse experiment to measure changes in nutrient availability in mouse faeces following TZP treatment and NDM-1 *E. coli* colonisation, and to measure NDM-1 *E. coli* growth in

mouse faeces following the administration of a mixture of inhibitory metabolites. We used a

previously published mouse model of intestinal CRE colonisation as described by Perez and

colleagues (**Figs. 7a and 9a**)[19]. The dosing schedule of the metabolite mixture was designed based

on a previous study from our group, where we investigated the effects of the microbial metabolite

valerate on the intestinal colonisation with *Clostridioides difficile* in a mouse model.[7]

In this study, mice were subcutaneously administered 8 mg of TZP (in 0.2 ml saline) or 0.2 ml of

saline once per day for 5 days to mimic the daily TZP dose recommended for human adults[19]. On

the third day of TZP or saline administration, 10^6 CFU of carbapenem-resistant (NDM-1) *E. coli*

(diluted in 0.2 ml saline) were orally administered to the mice. Next, TZP-treated mice that were

colonised with *E. coli* were fed by oral gavage with 0.2 ml of a mixture of metabolites (glycerol

triacetate, 65 mM; glycerol tripropionate, 15 mM; glycerol tributyrate, 15 mM; and glycerol trivalerate,

3.5 mM; diluted in PBS) or 0.2 ml of PBS once per day for 3 days. Faecal samples were collected

for NDM-1 *E. coli* plate counts (on Brilliance CRE agar, Oxoid, Basingstoke, UK) and for ^{1}H-NMR

spectroscopy.

Statistical analysis:

Statistical tests were performed using IBM SPSS Statistics Software 27 (IBM Corp, Armonk, New

York), GraphPad Prism 9.4.1 (La Jolla, California), or in R (v4.2.2). A p value less than 0.05 was

considered significant. 16S rRNA gene sequencing data (weighted as absolute abundances) were

analysed using Wilcoxon signed rank test with Benjamini & Hochberg FDR correction using the

DA.wil function within in the DAtest R package version 2.7.18. Peak integration values from the ^{1}H-

NMR data were analysed using Wilcoxon signed rank test with Benjamini & Hochberg FDR

correction using the DA.wil function within in the DAtest R package version 2.7.18 or a paired t-test

(two-sided) with Benjamini & Hochberg FDR correction using the pairwise_t_test function within the

rstatix R package version 0.7.1. A Mann-Whitney U test was used to compare log10-transformed

CRE counts from TZP-treated faecal cultures compared to water-treated faecal cultures.

Regularised canonical correlation analysis (rCCA) was used to correlate the 16S rRNA gene

sequencing data and ^{1}H-NMR data using the mixOmics library within R (as described above)[7,21]. For the carbon and nitrogen utilisation assays the growth curves were analysed in Python (v3.6.5) using the AMiGA software (as described above)[64]. For the nutrient preference experiment changes in the concentration of nutrients and metabolites for each CRE isolate were compared to the sterile media control over time using a two-way mixed ANOVA followed by pairwise comparisons with Bonferroni correction. For the *in vitro* mixed metabolite inhibition test, growth of the CRE isolate in the metabolite mixture was compared to growth in the no metabolite control at each pH using an unpaired t-test (two-sided). For the *in vitro* individual metabolite inhibition test, growth of the CRE isolate in the presence of each metabolite was compared to the no metabolite control using a one-way ANOVA followed by Dunn's multiple comparison test. For the mouse experiments, a paired t-test (two-sided) or a Mann-Whitney U test was used to compare log10-transformed NDM-1 *E. coli* counts from the faecal pellets of metabolite-treated mice compared to PBS-treated mice.

REFERENCES:

1. Dadgostar, P. Antimicrobial resistance: implications and costs. *Infect Drug Resist* **12**, 3903-3910 (2019).

2. Ventola, C. L. The antibiotic resistance crisis: part 1: causes and threats. *P T* **40**, 277-83 (2015).

3. Manges, A. R., Steiner, T. S. & Wright, A. J. Fecal microbiota transplantation for the intestinal decolonization of extensively antimicrobial-resistant opportunistic pathogens: a review. *Infect. Dis. (Lond)* **48**, 587-592 (2016).

4. Kim, S., Covington, A. & Pamer, E. G. The intestinal microbiota: Antibiotics, colonization resistance, and enteric pathogens. *Immunol Rev* **279**, 90-105 (2017).

5. UK Health Security Agency. Framework of actions to contain carbapenemase-producing Enterobacterales. *UK Health Security Agency* https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/1107705/Framework_of_actions_to_contain_CPE.pdf (2022).

6. Reed, A. D. & Theriot, C. M. Contribution of inhibitory metabolites and competition for nutrients to colonization resistance against *Clostridioides difficile* by commensal *Clostridium*. *Microorganisms* **9**, 371 (2021).

7. McDonald, J. A. K. *et al*. Inhibiting growth of *Clostridioides difficile* by restoring valerate, produced
by the intestinal microbiota. *Gastroenterology* **155**, 1495-1507 (2018).

8. Jacobson, A. *et al*. A gut commensal-produced metabolite mediates colonization resistance to
Salmonella infection. *Cell. Host Microbe* **24**, 296-307.e7 (2018).

9. Ferreyra, J. A. *et al*. Gut microbiota-produced succinate promotes *C. difficile* infection after
antibiotic treatment or motility disturbance. *Cell. Host Microbe* **16**, 770-777 (2014).

10. Ng, K. M. *et al*. Microbiota-liberated host sugars facilitate post-antibiotic expansion of enteric
pathogens. *Nature* **502**, 96-99 (2013).

11. Pudlo, N. A. *et al*. Phenotypic and genomic diversification in complex carbohydrate-degrading
human gut bacteria. *mSystems* **7**, e0094721 (2022).

12. Chang, D. E. *et al*. Carbon nutrition of *Escherichia coli* in the mouse intestine. *PNAS USA* **101**,
7427-32 (2004).

13. Tuncil, Y. E. *et al*. Reciprocal prioritization to dietary glycans by gut bacteria in a competitive
environment promotes stable coexistence. *mBio* **8**, 1068 (2017).

14. Rivera-Chávez, F., Lopez, C. A. & Bäumler, A. J. Oxygen as a driver of gut dysbiosis. *Free Radic*
Biol Med **105**, 93-101 (2016).

15. Oliphant, K. & Allen-Vercoe, E. Macronutrient metabolism by the human gut microbiome: major
fermentation by-products and their impact on host health. *Microbiome* **7**, 91 (2019).

16. Sorbara, M. T. *et al*. Inhibiting antibiotic-resistant *Enterobacteriaceae* by microbiota-mediated
intracellular acidification. *J. Exp. Med.* **216**, 84-98 (2019).

17. Djukovic, A. *et al*. *Lactobacillus* supports *Clostridiales* to restrict gut colonization by multidrug-
resistant *Enterobacteriaceae*. *Nat Commun* **13**, 5617 (2022).

18. Ladirat, S. E. *et al*. High-throughput analysis of the impact of antibiotics on the human intestinal
microbiota composition. *J Microbiol Methods* **92**, 387-97 (2013).

19. Perez, F., Pultz, M. J., Endimiani, A., Bonomo, R. A. & Donskey, C. J. Effect of antibiotic
treatment on establishment and elimination of intestinal colonization by KPC-producing
Klebsiella pneumoniae in mice. *Antimicrob Agents Chemother* **55**, 2585–2589 (2011).

20. Venturini, C., Bowring, B., Fajardo-Lubian, A., Devine, C. & Iredell, J. Effects of antibiotic

treatment with piperacillin/tazobactam versus ceftriaxone on the composition of the murine gut

microbiota. *Antimicrob Agents Chemother* **65**, 1504 (2021).

21. K. Le Cao, F. Rohart, I. Gonzalez and S. Dejean. mixOmics: Omics Data Integration Project. R

package version 6.1.2. https://bioconductor.org/packages/mixOmics/ (2017).

22. Autieri, S. M. *et al*. L-fucose stimulates utilization of D-ribose by *Escherichia coli* MG1655

DeltafucAO and *E. coli* Nissle 1917 DeltafucAO mutants in the mouse intestine and in M9

minimal medium. *Infect Immun* **75**, 5465-75 (2007).

23. Shelton, C. D. *et al*. *Salmonella enterica* serovar Typhimurium uses anaerobic respiration to

overcome propionate-mediated colonization resistance. *Cell Rep* **38**, 110180 (2022).

24. Schwartz, B. The effect of temperature on the rate of hydrolysis of triglycerides by pancreatic

lipase. *J. Gen. Physiol.* **27**, 113-118 (1943).

25. Bergan, T., Nord, C. E. & Thorsteinsson, S. B. Effect of meropenem on the intestinal microflora.

Eur J Clin Microbiol Infect Dis **10**, 524-7 (1991).

26. Kager, L., Brismar, B., Malmborg, A. S. & Nord, C. E. Imipenem concentrations in colorectal

surgery and impact on the colonic microflora. *Antimicrob Agents Chemother* **33**, 204-8 (1989).

27. Pletz, M. *et al*. Ertapenem pharmacokinetics and impact on intestinal microflora, in comparison

to those of ceftriaxone, after multiple dosing in male and female volunteers. *Antimicrob Agents*

Chemother **48**, 3765-72 (2004).

28. Nord, C. E., Brismar, B., Kasholm-Tengve, B. & Tunevall, G. Effect of piperacillin/tazobactam

treatment on human bowel microflora. *J Antimicrob Chemother* **31**, 61-5 (1993).

29. Stewardson, A. J. *et al*. Collateral damage from oral ciprofloxacin versus nitrofurantoin in

outpatients with urinary tract infections: a culture-free analysis of gut microbiota. *Clin Microbiol*

Infect **21**, 344.e1-11 (2015).

30. Guk, J. *et al*. Modeling the bacterial dynamics in the gut microbiota following an antibiotic-induced

perturbation. *CPT Pharmacometrics Syst Pharmacol* **11**, 906-918 (2022).

31. Kang, J. *et al*. Long-term ecological and evolutionary dynamics in the gut microbiomes of

carbapenemase-producing *Enterobacteriaceae* colonized subjects. *Nat Microbiol* **7**, 1516-1524

(2022).

32. Ianiro, G. *et al*. Variability of strain engraftment and predictability of microbiome composition after

fecal microbiota transplantation across different diseases. *Nat Med* **28**, 1913-1923 (2022).

33. Fernandez-Julia, P., Commane, D. M., van Sinderen, D. & Munoz-Munoz, J. Cross-feeding

interactions between human gut commensals belonging to the *Bacteroides* and *Bifidobacterium*

genera when grown on dietary glycans. *Microbiome Res Rep* **1**, 1-12 (2022).

34. DeMartino, P. & Cockburn, D. W. Resistant starch: impact on the gut microbiome and health.

Curr Opin Biotechnol **61**, 66-71 (2020).

35. Cerqueira, F. M., Photenhauer, A. L., Pollet, R. M., Brown, H. A. & Koropatkin, N. M. Starch

digestion by gut bacteria: crowdsourcing for carbs. *Trends Microbiol* **28**, 95-108 (2020).

36. Vital, M. *et al*. Metagenomic Insights into the degradation of resistant starch by human gut

microbiota. *Appl Environ Microbiol* **84**, 1562 (2018).

37. Chi, Z., Liu, J. & Zhang, W. Trehalose accumulation from soluble starch by *Saccharomycopsis*

fibuligera sdu. *Enzyme Microb Technol* **28**, 240-245 (2001).

38. Niness, K. R. Inulin and oligofructose: what are they? *J Nutr* **129**, 1402S-6S (1999).

39. Kaya, M., Sousa, A. G., Crépeau, M. J., Sørensen, S. O. & Ralet, M. C. Characterization of citrus

pectin samples extracted under different conditions: influence of acid type and pH of extraction.

Ann Bot **114**, 1319-26 (2014).

40. Bastawde, K. B. Xylan structure, microbial xylanases, and their mode of action. *World J Microbiol*

Biotechnol **8**, 353-68 (1992).

41. Kelly, G. S. Larch arabinogalactan: clinical relevance of a novel immune-enhancing

polysaccharide. *Altern Med Rev* **4**, 96-103 (1999).

42. Pultz, N. J., Hoskins, L. C. & Donskey, C. J. Vancomycin-resistant enterococci may obtain

nutritional support by scavenging carbohydrate fragments generated during mucin degradation

by the anaerobic microbiota of the colon. *Microb Drug Resist* **12**, 63-7 (2006).

43. Smid, E. J., Poolman, B. & Konings, W. N. Casein utilization by lactococci. *Appl Environ Microbiol*

57, 2447-52 (1991).

44. Assaf, A. *et al*. Evaluation of the impact of buffered peptone water composition on the

discrimination between *Salmonella enterica* and *Escherichia coli* by Raman spectroscopy. *Anal*

Bioanal Chem **412**, 3595-3604 (2020).

45. Raja Makendran Nandan Raja Rajendran. Enzymatic Conversion of RNA from Yeast Extract to

Guanosine monophosphate (a flavoring agent). Master's thesis, Chalmers University of

Technology https://odr.chalmers.se/handle/20.500.12380/166692 (2012).

46. Theriot, C. M. *et al*. Antibiotic-induced shifts in the mouse gut microbiome and metabolome

increase susceptibility to *Clostridium difficile* infection. *Nat. Commun.* **5**, 3114 (2014).

47. Rivera-Chávez, F. *et al*. Depletion of butyrate-producing clostridia from the gut microbiota drives

an aerobic luminal expansion of *Salmonella*. *Cell Host Microbe* **19**, 443-54 (2016).

48. Mortensen, P. B., Holtug, K. & Rasmussen, H. S. Short-chain fatty acid production from mono-

and disaccharides in a fecal incubation system: implications for colonic fermentation of dietary

fiber in humans. *J Nutr* **118**, 321-5 (1988).

49. Van der Meulen, R., Adriany, T., Verbrugghe, K. & De Vuyst, L. Kinetic analysis of bifidobacterial

metabolism reveals a minor role for succinic acid in the regeneration of NAD+ through its growth-

associated production. *Appl Environ Microbiol* **72**, 5204-10 (2006).

50. Rios-Covian, D., Salazar, N., Gueimonde, M. & de Los Reyes-Gavilan, C G. Shaping the

metabolism of intestinal *Bacteroides* population through diet to improve human health. *Front

Microbiol* **8**, 376 (2017).

51. Kageyama, A., Benno, Y. & Nakase, T. Phylogenetic and phenotypic evidence for the transfer of

Eubacterium aerofaciens to the genus *Collinsella* as *Collinsella aerofaciens* gen. nov., comb.

nov. *Int J Syst Bacteriol* **49**, 557-65 (1999).

52. Ghosh, S., Whitley, C. S., Haribabu, B. & Jala, V. R. Regulation of intestinal barrier function by

microbial metabolites. *Cell Mol Gastroenterol Hepatol* **11**, 1463-1482 (2021).

53. Gao, G. *et al*. Effects of valerate on intestinal barrier function in cultured Caco-2 epithelial cell

monolayers. *Mol Biol Rep* **49**, 1817-1825 (2022).

54. Otter, J. A. *et al*. Detecting carbapenemase-producing *Enterobacterales* (CPE): an evaluation of

an enhanced CPE infection control and screening programme in acute care. *J Antimicrob

Chemother* **75**, 2670-2676 (2020).

55. Kantele, A. *et al*. Dynamics of intestinal multidrug-resistant bacteria colonisation contracted by

visitors to a high-endemic setting: a prospective, daily, real-time sampling study. *Lancet Microbe*

2, e151-e158 (2021).

56. McDonald, J. A. *et al*. Evaluation of microbial community reproducibility, stability and composition

in a human distal gut chemostat model. *J. Microbiol. Methods* **95**, 167-174 (2013).

57. Moon, Y. S., Chung, K. C. & Gill, M. A. Pharmacokinetics of meropenem in animals, healthy

volunteers, and patients. *Clin Infect Dis* **24**, S249–S255 (1997).

58. Burdet, C. *et al*. Ceftriaxone and cefotaxime have similar effects on the intestinal microbiota in

human volunteers treated by standard-dose regimens. *Antimicrob Agents Chemother* **63**, 2244

(2019).

59. Pessôa de Menezes e Silva, C H. Elaboration and evaluation of a new screening medium for

detection and presumptive identification of extended-spectrum beta-lactamase-producing

organisms (ESBL). *Braz J Microbiol* **31**, 271-274 (2000).

60. Brismar, B., Edlund, C., Malmborg, A. S. & Nord, C. E. Ciprofloxacin concentrations and impact

of the colon microflora in patients undergoing colorectal surgery. *Antimicrob Agents Chemother*

34, 481-3 (1990).

61. Mullish, B. H. *et al*. Functional microbiomics: evaluation of gut microbiota-bile acid metabolism

interactions in health and disease. *Methods* **149**, 49-58 (2018).

62. Callahan, B. J. *et al*. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat*

Methods **13**, 581-3 (2016).

63. Webb, A. Increasing the sensitivity of magnetic resonance spectroscopy and imaging. *Anal*

Chem **84**, 9-16 (2012).

64. Midani, F. S., Collins, J. & Britton, R. A. AMiGA: software for automated analysis of microbial

growth assays. *mSystems* **6**, e0050821 (2021).

65. Payne, A. N., Zihler, A., Chassard, C. & Lacroix, C. Advances and perspectives in *in vitro* human

gut fermentation modeling. *Trends Biotechnol.* **30**, 17-25 (2012).