Integrated Drivers of Basal and Acute Immunity in Diverse Human Populations

Aisha Souquette[1,2], E. Kaitlynn Allen[1], Christine M. Oshansky[1], Li Tang[3], Sook-san Wong[4], Trushar Jeevan[4], Lei Shi[3], Stanley Pounds[3], George Elias[5,6,7,8], Guillermina Kuan[9,10], Angel Balmaseda[10,11], Raul Zapata[10,11], Kathryn Shaw-Saliba[12], Pierre Van Damme[13], Viggo Van Tendeloo[5], Juan Carlos Dib[14], Benson Ogunjimi[6,7,8,15], Richard Webby[4], Stacey Schultz-Cherry[4], Andrew Pekosz[12,16], Richard Rothman[12], Aubree Gordon[17], Paul G. Thomas[1]*

[1] Department of Immunology, St. Jude Children's Research Hospital, Memphis, Tennessee, 38105, USA; [2] University of Tennessee Health Science Center, Memphis, Tennessee, 38163, USA; [3] Department of Biostatistics, St. Jude Children's Research Hospital, Memphis, Tennessee, 38105, USA; [4] Department of Infectious Disease, St. Jude Children's Research Hospital, Memphis, Tennessee, 38105, USA; [5] Laboratory of Experimental Hematology (LEH), Vaccine and Infectious Disease Institute, University of Antwerp, Antwerp, 2610, Belgium; [6] Center for Health Economics Research and Modeling Infectious Diseases (CHERMID), Vaccine and Infectious Disease Institute, University of Antwerp, Antwerp, 2610, Belgium; [7] Antwerp Center for Translational Immunology and Virology (ACTIV), Vaccine and Infectious Disease Institute, University of Antwerp, Antwerp, 2610, Belgium; [8] Antwerp Unit for Data Analysis and Computation in Immunology and Sequencing (AUDACIS), University of Antwerp, Antwerp, 2610, Belgium; [9] Centro de Salud Sócrates Flores Vivas, Ministry of Health, Managua, 12014, Nicaragua; [10] Centro Nacional de Diagnóstico y Referencia, Ministry of Health, Managua, 16064, Nicaragua; [11] Sustainable Sciences Institute (Site - Managua, Nicaragua), San Francisco, California, 94102, USA; [12] Department of Emergency Medicine, Johns Hopkins University School of Medicine, Baltimore, MD, 21209, USA; [13] Center for the Evaluation of Vaccination (CEV), Vaccine and Infectious Disease Institute, University of Antwerp, Antwerp, 2610, Belgium; [14] Department of Medicine, Universidad del Norte, Barranquilla, 081001, Colombia; [15] Department of Paediatrics, Antwerp University Hospital, Antwerp, 2610, Belgium; [16] W. Harry Feinstone Department of Molecular Microbiology and Immunology, The Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland 21205, USA; [17] Department of Epidemiology, School of Public Health, University of Michigan, Ann Arbor, Michigan, 48109, USA

*Corresponding author: paul.thomas@stjude.org

Keywords: immune variation, influenza, herpesvirus, co-infection, systems immunology, basal immunity, acute immunity, immunogenetics, immunophenotype, correlate of severity

Abstract

Prior studies have identified genetic, infectious, and biological associations with immune competence and disease severity; however, there have been few integrative analyses of these factors and study populations are often limited in demographic diversity. Utilizing samples from 1,705 individuals in 5 countries, we examined putative determinants of immunity, including: single nucleotide polymorphisms, ancestry informative markers, herpesvirus status, age, and sex. In healthy subjects, we found significant differences in cytokine levels, leukocyte phenotypes, and gene expression. Transcriptional responses also varied by cohort, and the most significant determinant was ancestry. In influenza infected subjects, we found two disease severity immunophenotypes, largely driven by age. Additionally, cytokine regression models show each determinant differentially contributes to acute immune variation, with unique and interactive, location-specific herpesvirus effects. These results provide novel insight into the scope of immune heterogeneity across diverse populations, the integrative effects of factors which drive it, and the consequences for illness outcomes.

Introduction

Functional consequences of immune heterogeneity are demonstrated by variation in vaccine efficacy, responses to therapeutic treatment, infectious disease severity, and susceptibility to autoimmunity. Understanding the landscape and drivers of immune variation would provide valuable insight into immune competence and points of intervention to improve disease outcome. Indeed, baseline immune predictors of vaccine responses and neoplastic disease prognosis have already been identified in a variety of studies.[1,2]

Immune variation, at baseline and in response to immune challenge, is dynamically influenced by unique and interactive effects of intrinsic and extrinsic host factors. Intrinsic factors are internal to the host and include variables such as age, sex, genetics, and microbiome. Extrinsic factors are external to the host and include variables such as diet, environment, and acquired infections. Each of these determinants can influence multiple facets of immunity, including, but not limited to, susceptibility to infection or immune mediated disease, the basal activation state of the immune system, pathogen detection, downstream signaling, priming of adaptive immunity, leukocyte phenotype and function, and/or response to therapeutic treatment.[3–22] For example, in a study of three independent influenza cohorts, we found that a single nucleotide polymorphism rs3448144-A in *IFITM3* is enriched in severe patients, governs transcription factor binding and promoter activity, and alters methylation of the *IFITM3* promoter in CD8 T cells, which correlates with cell frequency in nasal washes.[12] Coinciding with these results, recent studies have reported associations between rs3448144 and COVID-19 disease severity.[23,24] Importantly, the collective impact of immune modulation induced by intrinsic and extrinsic host factors can alter vaccination responses and illness outcome. However, further study is needed to clarify the quantitative effects of these variables, as conflicting conclusions have occurred across studies, possibly due to skews in age distribution, geographical location, or ancestry impacting observed results. Ultimately, this limits both our understanding of the extent of immune heterogeneity and the general applicability of conclusions drawn from single cohorts.

Here, we quantify the contribution of intrinsic and extrinsic host factors on the setpoint of
baseline immunity, the transcriptional response to *in vitro* immune challenge, and the magnitude
and quality of the immune response to *in vivo* influenza infection in diverse populations around
the world. Immune measures include cytokines (from plasma and nasal wash), cellular
phenotypes, and/or gene expression (both at baseline and post PBMC stimulation with PRR
ligands). Of these, cytokines were our primary and most extensively used measurement, with
data collected at the RNA and protein level. While most prior work has focused on cell
phenotypes (by flow cytometry), cytokines have several advantages from both a scientific and
clinical standpoint. Samples are easily obtained and stored, and analytes can quickly and
precisely be measured with minimal equipment, sample input, and processing. In addition, they
have a substantial dynamic range and offer broad profiling of innate and adaptive immunity. For
influenza infected subjects, outcome measures include presence/absence of symptoms,
symptom severity scores, viral load, and duration of influenza virus shedding. To explain
variation in these measures, linear regression modeling was utilized, which allows for an easily
interpretable and transparent integrative analysis of all independent variables and a
quantification of their unique, interactive, and collective effects on the immune measure of
interest. Collectively, the data presented here, representing 123,178,117 data points from 1,705
individuals, show age, sex, herpesviruses, and host genetics can independently and
interactively influence baseline immunity and the acute immune response to influenza virus
infection. Of these determinants, ancestry is the most significant factor. Importantly, these
factors differentially contribute to variation in a given immune measure, and the cumulative
impact of immune modulation by these factors results in unique immune profiles associated with
disease severity. Our results demonstrate that simultaneous assessment of diverse populations
is necessary to better understand the scope of immune variation and the factors that contribute
to it.

Results

Explanatory factor variation within and across diverse populations

This study includes samples from 1,705 healthy or influenza infected subjects from 5 countries
and 8 distinct populations (**Fig 1**). North American populations include the Memphis cohort and
the Baltimore cohort. Latin American populations include the Nicaragua cohort and the
Colombia cohorts, which include subjects from the local city of Calabazo and two indigenous
populations, Seywiaka and Umandita. Samples also include a European cohort from Belgium
and an Asian cohort from Taiwan. Select assays and corresponding analyses focused on our
two most diverse cohorts, Memphis and Nicaragua, which have the largest sample size,
availability, and variety of sample types collected. These populations also capture
environmental diversity, and, collectively, represent three distinct racial/ethnic groups
(White/Caucasian, Hispanic/Latino, Black/African American), two of which are highly admixed,
thereby representing greater genetic diversity. Enrichment of diverse, understudied racial/ethnic
groups is a key feature of our study, as recent work has shown that increasing genetic diversity,
as opposed to increasing the sample size of European ancestry groups, provides substantial

improvement to genetic analyses, such as fine-mapping functional variants and portability of
polygenic risk scores.[25,26]
123
124 Before examining baseline and acute immunity, we assessed the degree of variation in
125 explanatory variables across cohorts. Here, intrinsic host factors include biological and genetic
126 variables. Biological factors include age and sex, and the distribution of each varied by cohort
127 (**Fig 2A-B**). Overall, the study was 58% female and age ranged from <1 year up to 92 years of
128 age.
129
Heritable contribution to immune variation was determined for a subset of healthy and infected
subjects with available DNA or PBMCs (n = 355, 234 Memphians and 121 Nicaraguans), using
genome-wide ancestry informative markers and candidate functional, immune-related SNPs.
Genetic based ancestry, as determined by ancestry informative markers (AIMs), accounts for
inherent genetic differences across diverse populations, typically from distinct geographical
regions. This allows for a genetic and quantitative representation of admixed populations, as
opposed to discrete classifications of race, which do not adequately represent populations with
mixed ancestral backgrounds. AIMs were analyzed utilizing a principal component analysis
(PCA), and the top two principal component values were included in downstream regression
analyses. Graphical representation of the PCA results shows distinct clusters that were similar
to the self-reported racial and ethnic makeup of the Memphis (75% African/Black American and
25% Caucasian/White American) and Nicaragua (100% Latino) cohorts, with a subset of
ancestrally mixed individuals falling between clusters (**Fig 2C**). Importantly, only AIMs were able
to distinguish between ancestral population clusters (**Supp Fig 1A-B**) and account for genetic
variation within a given population (**Supp Fig 1C**). This highlights the limited applicability of
current "Race" and "Ethnicity" categories collected in the clinical setting and applied in
biomedical research, and supports current arguments for the need for inclusion of more
appropriate factors in human studies aiming to distinguish between biological, epidemiological,
and social determinants of health and disease.[27,28] For the candidate SNP list, we examined
common SNPs in functional regions of genes encoding the 41 cytokines in our assay and in 3
transcription factors important in initiating antiviral immunity, namely IRF3, IRF7, and all 3
subunits of NFkB (detailed description of prioritization is in the Methods). While some SNPs
have similar allele frequencies across the Memphis and Nicaragua cohorts, several, such as
rs13900 in MCP1 and rs3024496 in IL10, show significant variation and opposing major alleles
in the two populations (**Fig 2D**, **Supp Table 1**), indicating genetic differences in immune-related
genes with potential functional consequences due to ancestry.
156
Herpesviruses were utilized as an extrinsic factor of infectious exposure because they are
globally prevalent, cause chronic infections that are generally well tolerated by
immunocompetent hosts, and have previously been associated with modulation of the immune
system.[29–32] The serostatus of cytomegalovirus (CMV), Epstein-Barr virus (EBV), herpes
simplex virus (HSV, 1 and 2 combined), human herpesvirus 6 (HHV6), and varicella-zoster virus
(VZV) were determined for healthy and influenza infected subjects with available plasma from
all eight cohorts (n = 1,532). Herpesvirus seroprevalence ranged from as low as 47.5% up to
97.5% (**Fig 2E**). Although the median age varies across cohorts, comparisons can be made

between cohorts of similar age ranges and/or with age adjustment and these results
demonstrate how herpesvirus prevalence varies across distinct groups. For example, although
the Baltimore, Taiwan, and Belgium cohorts are similar in age, Belgium has 31-35% lower
prevalence of HSV but a 9-24% higher prevalence of EBV. Additionally, despite having a similar
age group, the Colombian cohort has 22% higher prevalence of CMV, 31% higher prevalence of
EBV, and a 32% lower prevalence of HHV6 compared to Nicaragua. Importantly, with the
exception of the Belgium cohort, which has the most limited age range in participants, each
cohort exhibits a positive correlation between age and the total number of herpesviruses an
individual has (**Supp Fig 1D**). This is particularly important to account for in downstream
analyses because influenza disease severity is age associated, very young individuals (infants
and children under 5) tend to have more severe infection and tend to be herpesvirus
seronegative. Highlighting the ubiquity of herpesviruses, of the 1,532 individuals in this analysis,
only 10 (0.587%) were negative for CMV, HSV, EBV, and HHV6, and only 118 (6.92%) tested
positive for a single herpesvirus (**Fig 2F**). These data underscore the importance of accounting
for multiple herpesvirus infections when trying to assess their contributions to a given disease
outcome, as most individuals have more than one. However, due to the high prevalence of
HHV6 and VZV, the herpesvirus serostatuses in downstream analyses are focused on CMV,
EBV, and HSV.

Distinct basal immune profiles mold response to immune challenge

To begin assessing baseline immune status, the levels of 38 plasma/serum cytokines were
measured in 631 healthy subjects.[33–38] Cytokine levels were natural log (ln) transformed and
visualized using a t-Distributed Stochastic Neighbor Embedding (t-SNE) plot which aids in
dimensionality reduction while utilizing data from all cytokines measured. These results show
marked variation in cytokine levels, with distinct clustering by cohort. Some populations exhibit
tight clusters (Baltimore), whereas others exhibit dispersed clusters (Colombia) or even two
distinct clusters (Memphis), suggesting the variance of cytokine profiles across individuals
varies by population and, consequently, focusing on a population with minimal variation across
subjects may underestimate the extent of immune variation and bias results (**Fig 3A**). Split
clustering of the Memphis cohort was not due to age, sex, herpesvirus serostatus, race,
season/year, assay plate, or time between assay runs (**Supp Fig 2A-G**). Moreover, in a
subsequent analysis, cohort clusters and their positions relative to each other remain after
removal of Memphis cohort data (**Supp Fig 2H**). Variation was also observed at the individual
cytokine level for immunomodulatory cytokines, pro-inflammatory cytokines, chemokines, and
growth factors (**Fig 3B**).
199
To examine the effects of intrinsic and extrinsic host factors on basal cytokine levels, a linear
regression model (Model #1, **Fig 1**) was used to take into account the effects of: age, sex,
cohort (accounting for genetics and unknown environmental factors specific to the
geographically distinct populations), each herpesvirus alone, any interaction between a
herpesvirus and age (due to the age association with serostatus), and any interaction between
herpesviruses (since most individuals are infected with more than one). These results show
herpesviruses have largely unique effects on basal cytokine levels, and the herpesvirus which
caused the most effects was CMV (**Fig 3C** and **Supp Fig 3**). Each additional standard deviation

unit increase in age (years) is cumulatively associated with altered GRO, IFNg, Eotaxin, and
IL12p70, thus the larger the difference in age, the greater the magnitude of age effects (**Supp**
Fig 3). Male sex was associated with decreased TGFa and cohort differentially affected all
cytokines (**Supp Fig 3**). The adjusted R^2 result from a linear regression model is an estimate of
how much variability in the dependent variable the model can account for, and the "adjustment"
refers to a penalty to the R^2 value due to the incorporation of additional variables. This value
was highest for the IL4 model at 0.4709 (47.1%) (**Supp Fig 3**). Another output from the
regression models is an estimate of a coefficient for each factor which represents the change in
the dependent variable per unit change in the factor. Comparing coefficients across factors
provides insight into the magnitude of the effect of that factor. In these healthy cytokine models,
cohort was associated with the strongest effects on circulating cytokine levels (**Supp Fig 3**).
Collectively, these data suggest distinct populations have unique cytokine profiles, the effect of
which likely represents divergent baseline immune states, significant impacts on downstream
effector functions, including immune cell differentiation and functional responses to stimuli.
222
To further explore the potential for distinct immunophenotypes, we focused on two cohorts with
PBMC samples, Memphis and Nicaragua, which allowed for more comprehensive investigation
and statistical analysis. Flow cytometry was utilized to characterize the activation and/or
differentiation state of circulating lymphocytes and antigen presenting cells at baseline. Samples
included 28 Memphians (average age = 22, median age = 18) and 19 Nicaraguans (average
age = 20, median age = 15). Nicaraguans exhibited a trend towards higher regulatory T cell
(Treg)-like CD25+ CD4 T cells (padj = 0.071, **Supp Table 2**). To validate these findings, the
Treg transcription factor FOXP3 (forkhead box protein P3) was included in a follow up
experiment and results confirmed Nicaraguans have an increased frequency of FOXP3+ CD4
Tregs (p = 0.028, **Fig 3D**). Nicaraguans also showed significantly lower frequency and number
of activated CD4 and CD8 T cells (padj = 5.5e-4 for both subsets), but higher plasmablasts
(padj = 3.04e-3) and CD8 effector memory T cells (padj = 0.014), the latter being consistent with
increased prevalence of chronic infections (**Fig 3D**). Moreover, Memphians showed a higher
frequency and number of pro-inflammatory plasmacytoid dendritic cells (pDCs) (padj = 2.37e-3)
(**Fig 3D**).
238
Substantial variation was also observed between the two cohorts at the RNA level after PBMC
stimulation with plain media (an unstimulated control), lipopolysaccharide (a bacterial TLR4
ligand), Pam3CSK4 (a bacterial TLR1/2 ligand), transfected poly(I:C) (representative of RNA
virus infection), or transfected poly(dA:dT) (representative of DNA virus infection). For each
stimulation condition, samples included 94 Memphians and 94 Nicaraguans. Examples of genes
which varied by cohort include, but are not limited to, *IFI16* (a PRR important for herpesvirus
detection), *IRF1* and *IRF9* (upstream immune response transcription factors), *TBK1* (a
downstream immune signaling protein), *CCL4* and *CXCL2* (chemokines), *CD86* (an antigen
presenting cell activation marker and ligand for a co-stimulatory molecule expressed on T cells),
KLRC1 (inhibitory receptor NKG2A, expressed on NK and T cells), and *IFNG* (an antiviral
cytokine) (**Fig 4A, Supp Table 3**). These genes represent each step of the immune response
and highlight the numerous ways and potential for unique immune profiles to arise in diverse
populations. Notably, differential gene expression was stimulation condition dependent,

highlighting the need for future studies to investigate immune variation under a variety of
immune pressures.
254
Coinciding with the basal cytokine results (**Fig 3A**), PCA analysis of the media (unstimulated
control) gene expression data shows substantial variation in the basal immune transcriptional
profile (**Fig 4B**). To determine which factors were driving this variation, linear regression
modeling was utilized to account for age, sex, herpesvirus status, and ancestry (Model #2, **Fig
1**). These results show basal immune transcriptional heterogeneity is largely driven by genetic
ancestry (affected 62.2% of 90 genes), followed by herpesvirus infection (22.2% of genes), and
demographic variables, age and sex (7.8% of genes) (**Fig 4C, Supp Fig 4A, Supp Table 3**). A
similar model was utilized to assess determinants of stimulated gene expression, and
accounted for stimulant, age, sex, herpesvirus status, ancestry, and the basal immune profile
(Model #3, **Fig 1**). Similar to the basal response, ancestry was the most prominent determinant,
contributing to statistically significant alterations in 83% of genes examined, the basal
transcriptional profile affected 74% of genes (on top of the effects already captured by
ancestry), and herpesvirus status affected 66%, while demographic variables (age, sex)
affected 31% (**Fig 4C, Supp Fig 4B, Supp Table 3**).
269
Exploring these transcriptional results further, expression quantitative trait loci (eQTL) were
determined to define SNPs that were associated with altered gene expression and demonstrate
a means by which genetic effects could contribute to the observed transcriptional heterogeneity.
These data were analyzed with linear regression modeling in order to account for: genotypes for
immune-related SNPs of interest, genetic based ancestry, age, sex, herpesvirus status, and the
stimulation condition (Model #4, **Fig 1**). Of the 94 prioritized candidate SNPs, 36 SNPs in 26
genes (24 cytokines and 2 transcription factors) were significant factors in the expression of
their respective genes (**Supp Fig 4C, Supp Table 4**). Notably, 13 of these genes also showed a
measurable heritable influence in Brodin et al. at the protein level (GCSF, IFNg, IL10, IL12p70,
IL13, IL15, IL6, TNFb, GROa, IL8, MCP1, TGFa, and VEGF).[7] While further study is required to
determine the mechanism behind the eQTLs, these data demonstrate how genetic variation can
significantly affect gene expression both at baseline and during immune responses. With
respect to immune variation, this would be of most interest when eQTL allele frequencies vary
across diverse populations, such as that shown for rs13900 in *MCP1*, which had a minor allele
frequency (MAF) of 4% in Memphis and 38% in Nicaragua (**Fig 2D, Supp Table 1**).

Divergent age-associated influenza cytokine correlates of severity

Results from unstimulated, baseline analyses demonstrate significant immune heterogeneity
across diverse populations at the RNA, protein, and cellular level. Further, these data establish
that basal immune variation is differentially affected by intrinsic and extrinsic host factors. The
cumulative effect of modulation in the basal immune state may lead to unique immune profiles
in response to infection. In support of this, the *in vitro* TLR stimulation results show distinct
transcriptional responses between the Memphis and Nicaragua cohorts. To further assess the
extent to which unique immunophenotypes may occur, we examined how robust immune
signatures are across distinct cohorts during acute viral infection *in vivo*.
294

We determined cytokine correlates of severity in three independent human cohorts of naturally
acquired influenza virus infection. Cytokine levels were measured in plasma and/or the nasal
wash from influenza positive subjects in the Memphis (n = 165), Baltimore (n = 222), and
Taiwan (n = 226) cohorts. Each cohort was independently analyzed and results were compared
to identify unique and shared correlates. Utilizing a subset of samples, a previous study in the
Memphis cohort found increased levels of nasal wash IFNa2, MCP3, IL6, and VEGF correlated
with total, upper respiratory (URT), and/or lower respiratory (LRT) symptom scores. [33]
Additionally, increased plasma MCP3, IL10, and IL6 predicted hospitalization and were
positively associated with symptom scores, and IP10 positively correlated with influenza viral
load. [33] The current Memphis analysis expands upon this dataset and includes three additional
seasons of cytokine data.
306
There are key similarities and differences amongst the cohorts included in this analysis that are
important to consider in interpretation of the results. First, the Memphis cohort is centered
around pediatric recruitment, and thus is enriched in children and young adults, though they
also enrolled adult contacts. In contrast, the Baltimore and Taiwan cohorts are both adult
studies. Additionally, the Baltimore and Memphis cohorts are similar in genetic and geographical
background, whereas Taiwan is distinct from the rest. Further, the Baltimore, and Taiwan
cohorts each have binary (presence/absence) disease outcome data whereas the Memphis
cohort includes both binary and continuous outcomes, the latter of which requires a different
statistical approach. Importantly, binary disease outcomes with a high prevalence dilute the
specificity of our severity measures, and thus are removed from downstream analyses. Results
below are focused on continuous variables or binary variables with a prevalence <40% (range
14-39%, additional details in Study Subjects section of Methods).
319
All cohorts exhibit increased levels of plasma IL6, TNFa, GCSF, and IL8 in hospitalized,
pneumonia, and/or supplemental oxygen patients. This suggests these cytokines are associated
with severe influenza disease in general, irrespective of age and geographical location (**Fig 5,**
Supp Fig 5, Supp Table 5). However, these analytes may reflect severe (respiratory) viral
infection more broadly, as they have also been associated with severe COVID-19.[39–42] Cytokine
profiles as a whole, in both the plasma and nasal wash, are distinct by age groups, possibly
reflecting unique pathophysiologies. The severe pediatric response is characterized by global
hyper-inflammatory cytokine levels, whereas the severe adult response shows largely
decreased cytokine levels with a selective inflammatory response of the aforementioned
analytes (**Fig 5, Supp Fig 5, Supp Table 5)**. Notably, although the directionality is similar
among significant results in the adult cohorts, only decreased IFNg was shared, and thus
represents the most robust correlate for this age group. All additional plasma cytokine correlates
of severity in these adult populations were also associated with lower cytokine levels, albeit
specific to one of the adult cohorts (**Fig 5**, **Supp Fig 5**, **Supp Table 5**).
334
These data demonstrate correlates of severity can differ across distinct populations and
highlight the need for future work to simultaneously examine immune signatures of health and
disease in diverse populations for a more comprehensive analysis and identification of robust
therapeutic targets. Additionally, studies which aim to identify cytokine correlates should utilize a

broad panel, as focusing on a few select analytes is insufficient to characterize the overall
cytokine profile. These results also suggest that a "one size fits all" therapeutic approach may
not be efficacious, as distinct immune profiles may exist by extremes of age and likely require
different treatments. Multiple severity associated immunotypes were also recently observed in a
study of SARS-CoV-2 cellular responses, wherein 3 distinct cellular immunophenotypes were
identified for severe COVID-19.[43]

Age specific herpesvirus associations with influenza severity

In addition to immune correlates of disease severity, independent studies have identified
associations between age, sex, and CMV and the outcome of vaccination or infection with
influenza in humans.[33,44,45] To expand upon these findings, we examined disease outcome
utilizing a linear regression model which accounts for age, sex, CMV, EBV, and HSV serostatus,
as very few individuals are only infected with CMV (**Fig 2E-F**, **Supp Fig 1**).
351
In Memphis, the pediatric cohort, CMV serostatus was the only significant variable and it was
associated with improved outcome. Patients with a positive serostatus are enriched in the non-
severe subjects, and show a decrease of 29.1 ± 10.93 ($p = 0.0103$) in peak lower respiratory
symptom severity score (range = 0 to 176) (**Fig 6A, Supp Table 6**). Conversely, in the adult
Baltimore cohort, a positive CMV serostatus was associated with increased risk of pneumonia
(OR = 2.904, $p = 0.0457$) or hospitalization (OR = 4.966, $p = 5.72e-5$) (**Fig 6A, Supp Table 6**).
358
Aside from symptom severity, the Memphis cohort included two additional measures of the
quality of the immune response to influenza infection - influenza viral load was determined using
qRT-PCR (quantitative reverse transcription polymerase chain reaction) and functional antibody
titers were measured using microneutralization assays. To analyze these data, linear regression
modeling was utilized to assess the effect of biological (age, sex) and infectious (herpesvirus
serostatus) factors, while accounting for study day (when applicable). Neither CMV, EBV, nor
HSV had any effect on influenza viral load ($p = 0.66, 0.39$, and 0.69, respectively) (**Fig 6B,
Supp Table 6**). However, a positive HSV serostatus is associated with a decrease in duration of
influenza virus shedding by approximately 3.14 ± 1.42 days ($p = 0.0304$, **Fig 6B, Supp Table
6**). These results suggest that HSV co-infection may alter viral clearance mechanisms, such that
it decreases the days an individual sheds influenza virus. Although, there was no detected effect
on influenza viral load, one challenge with observational human studies is unknown true day of
infection and the viral load upon initial infection.
372
Analysis of microneutralization titers show that a positive CMV serostatus is associated with an
increase in functional antibody titers by approximately 144 ± 54.50 ($p = 0.0086$) (**Supp Table 6**).
Additionally, male sex is associated with an increase in antibody titer of 199 ± 60.70 ($p =
0.00112$) and titer decreases by 105.04 ± 30.60 per standard deviation unit change in age ($p =
0.00067$). These results are consistent with previous observations by Furman et al. which show
that CMV seropositive young adults exhibit augmented antibody responses to Fluzone
vaccination compared to CMV seronegative.[44]
380

Collectively, these data suggest a dichotomous, age-specific role of CMV on disease severity.
This is consistent with prior work which has shown CMV improves vaccine responses in young
populations, but increases frailty in the elderly.[44,46–48] Initial benefits may be mediated by trained
innate immunity, in which constant surveillance of chronic herpesvirus infection (in order to
mitigate reactivation) necessitates altered epigenetic and transcriptional profiles which confer
augmented response to restimulation.[49] However, the cumulative effects of CMV on immune
aging over time, such as through inflammaging and memory inflation, may eventually exceed
beneficial effects and lead to impaired, immunosenescent responses and, subsequently, poorer
outcome.[50–52] Future study is required to delineate the extent to which trained innate immunity,
CMV associated immunosenescence, or an alternative mechanism is the cause of these
differential influences on heterologous immunity.

Differential contributions of host genetics and herpesviruses to acute immune variation

The correlates of severity analysis demonstrates immune heterogeneity in acute immune
responses, and thus variation in putative therapeutic targets. Moreover, it identified biological
and infectious associations with influenza disease outcome. Having observed biological and
infectious contributions to basal cytokine levels, we sought to determine whether the association
between disease severity and these factors may be due, at least in part, to their effects on
cytokines, particularly those that were identified as a correlate of severity. These results will also
provide insight into whether such analyses would be useful in the identification of optimum
therapeutic targets by determining the extent to which there are differential contributions of
intrinsic and extrinsic host factors on cytokines of interest.
402
To achieve this, plasma and nasal wash samples were used from 83 subjects with naturally
acquired influenza infection in the Memphis cohort from time points throughout the duration of
infection. A total of 38 and 41 cytokines were measured in plasma (n = 237) and nasal wash (n
= 325), respectively. Since eQTLs were identified for a subset of analytes, this analysis begins
with an assessment of the effects of age, sex, and herpesviruses on cytokine levels, while
accounting for influenza infection related variables that can contribute to the immune state
(Models #5-6, **Fig 1**). This is followed by a second regression analysis, focused on cytokines
with an eQTL identified in the baseline analyses, which expands upon the previous model and
incorporates genetics by adding AIMs PCA values and SNP genotypes (Models #7-8, **Fig 1**).
412
For all analytes, a linear regression model was used to take into account the effects of: age,
sex, study day, influenza viral load, each herpesvirus alone, any interaction between a
herpesvirus and age, and any interaction between herpesviruses (Models #5-6, **Fig 1**). These
data show each herpesvirus has its own unique signature of immune modulation that is location
specific. In plasma (**Fig 6C, Supp Fig 6**), CMV alters the most cytokine levels independently,
which coincides with its primary site of latency in cells of the myeloid lineage. Furthermore,
herpesvirus interactions are most prevalent in the plasma, versus in the nasal wash, where
independent herpesvirus effects are more common. In the nasal wash (**Fig 6F, Supp Fig 7**), the
herpesvirus with the most independent effects is EBV, which predominantly infects the
nasopharynx region. In the plasma, the adjusted R^2 value reached 0.358 (or 35.8%) for IL10,
and herpesvirus interactions are generally associated with the strongest effects (i.e.

magnitudes) on cytokine levels as compared to the effects of independent herpesviruses or
other variables (i.e. age, influenza viral load) (**Supp Fig 6**). In the nasal wash, the adjusted R^2
value reached 0.332 (or 33.2%) for RANTES, and three factors accounted for most effects on
cytokine levels - viral load, age, and EBV, with the strongest effects from the latter two (**Supp**
Fig 7).
429
For the 23 cytokines which had eQTLs and protein level data, a second regression model was
performed for plasma and nasal wash, which included each factor previously described (age,
sex, herpesviruses, study day, viral load), in addition to the AIMs principal components 1 and 2
(genetic based ancestry) and the genotype for each SNP that significantly correlated with the
respective cytokine's gene expression (Models #7-8, **Fig 1**). GCSF and its SNPs which trended
toward a correlation were also included because fewer samples had detectable gene
expression (**Supp Table 7**). In the plasma, when there was an effect from incorporating
genetics into the cytokine regression models, it generally increased the amount of variability
accounted for by the model; however, the nasal wash models generally decreased. This is likely
due, at least in part, to the fact that eQTL status was determined from stimulated PBMCs as
opposed to nasal wash cells, which are at the site of infection, have a distinct milieu, and are
exposed to the external environment. Eight out of 24 cytokine models had a significant genetic
component and increased variability accounted for, two of which solely included a significant
AIMs PC without a significant association with a curated SNP (**Supp Table 7**). These cytokines
include chemokines (Eotaxin, MDC, MIP1a), immunomodulatory IL10, NK and memory T cell
proliferation inducing IL15, and growth factors (FGF2, GCSF, TGFa). The biggest change in
adjusted R^2, upon inclusion of genetic factors, occurred in Eotaxin with a 43.35% increase in the
variability accounted for by the model, for a final adjusted R^2 value of 0.5568 (**Fig 7A**).
Significant factors in the Eotaxin model include: sex, influenza viral load, SNP rs5027581, CMV
and its age interaction, the CMV and HSV interaction, the EBV and CMV interaction, and the
interaction of EBV, CMV, and HSV. The next highest adjusted R^2 was 0.3915 (39.15%) for
plasma IL10 and significant factors include age, sex, viral load, SNP rs3024496, the HSV and
age interaction, and the EBV and CMV interaction (**Fig 7B**).

Validation of the model: rs3024496 genotype modulates IL10 expression

To follow up on genetic associations from our analyses, we selected two associated SNPs with
potentially functional consequences for *in vitro* assays to assess effects on promoter activity.
SNP rs3024496 in *IL10* was chosen because IL10 is a correlate of severe influenza infection,
and its genotype prevalence varied across diverse populations (Memphis: 39% TT, 47% CT,
14% CC; Nicaragua: 67% TT, 32% CT, 1% CC). Of note, two SNPs within the gene encoding
IL10 were identified as eQTLs (rs3024496, rs3024498). However, including both SNPs in the
IL10 regression model served as a conditional SNP analysis and suggested there was only one
independent signal resulting from rs3024496. SNP rs2304204 in *IRF3* was also selected
because it is a transcription factor upstream of antiviral responses, with varying allele
prevalences across cohorts (Memphis: 40% GG, 40% AG, 20% AA; Nicaragua: 11% GG, 45%
AG, 44% AA). Additionally, an independent analysis using the GTEx Portal confirmed our
transcriptional results and showed rs2304204 is an eQTL in tissues such as the thyroid (p =
1.1e-19) and lung (p = 0.000027).

For each target, two luciferase reporter assay plasmids were created, which only differed in

sequence at the SNP site (**Fig 7C**, detailed in Methods). Unstimulated A549s exhibited higher

luciferase expression when the C allele was present for IL10-rs3024496 and when the G allele

was present for IRF3-rs2304204 (**Fig 7D**). Results were consistent 24 hours post infection with

influenza A virus and showed a higher magnitude of differential activity (**Fig 7E**). Coupled with

the eQTL findings, these results confirm rs3024496 is a functional SNP directly affecting IL10

promoter activity and coincide with the association predicted by the regression model - the CC

genotype was associated with higher plasma IL10 than the CT and TT genotypes which had

coefficient estimates of -0.3315 and -0.6830, respectively, when compared to the reference CC

genotype (**Fig 7B**). SNP rs2304204 in *IRF3* was also verified as a functional SNP with

coinciding directionality. This finding, coupled with the varying allele prevalence across diverse

populations, demonstrates a mechanism by which broad transcriptional changes could occur,

both at baseline and in response to immune challenge, as was observed in this study (**Fig 4**).

Together, these results demonstrate the integrative manner in which biological, infectious, and

genetic factors can influence variation in cytokines at the protein level during naturally acquired

influenza infection. The observation that these effects occur in proteins identified as correlates

of severity, highlights how variation in these factors can lead to distinct immunophenotypes

during acute infection and the potential impact on infectious disease outcome. Collectively, this

underscores the need for future studies to search for correlates of protection or severity across

diverse populations and to consider the factors which contribute to their variation in order to

identify the number of severe immune profiles, the most broadly applicable treatment, and

practical therapeutic options (i.e. those more affected by factors we can control or target).

Discussion

Studies of human immune variation thus far suggest genetics and age contribute to 20-40% and

5% of immune variance, respectively, leaving 55-75% possibly due to sex or environmental

influences.[53] However, studies to date often focus on a single ancestral group, geographical

location, immunological measure, and/or age range, and are sometimes limited in inclusion of

both sexes.[4,6–8,20,21,54–56] Often, this is in an attempt to limit additional variance attributed to

determinant categories that are not the focus of a given study. While statistically beneficial and

still informative, it is impossible to ascertain the scope of immune variation when studies are

focused on specific subsets of the human population and this limits the applicability of the

conclusions that are drawn. Moreover, this can obscure phenotypes, leading to false positive or

negative associations, and inconsistent results across studies that ultimately present a barrier to

translating results from the laboratory to effective, broadly applicable therapeutic treatments.

Lastly, comparisons of relative importance can be better understood with simultaneous

assessment of multiple determinant categories, and with increasing technologies for

multiparameter assessment, this approach is becoming more tractable, as demonstrated by this

study.

The demographics, immune measures, and analytical approach make this study unique from

prior work. First, the populations included are diverse in ancestral background, geographical

location, age, and sex. Second, immune measures were determined at baseline and during

immune challenge (both *in vitro* and *in vivo*), and include cell phenotypes, cytokines (both
circulatory and at the site of infection), and gene expression in response to PRR ligands
representative of both bacterial and viral infection. Third, the analytical approach utilized linear
regression modeling which allowed us to simultaneously account for and quantify the unique
and interactive effects of biological, infectious, and genetic factors. Fourth, our genetic approach
was based on ancestry informative markers and a targeted list of candidate immune-related
SNPs. This allowed for increased power compared to GWAS studies by limiting the impact of
multiple comparisons and provided clearly defined genetic factors that can be further explored in
mechanistic studies that aim to identify causal factors compared to heritability scores which are
unable to provide direct functional insight.
520
Investigation of baseline immunity shows differences in cellular phenotypes, cytokine levels, and
the basal transcriptional profile. Heterogeneity in baseline gene expression was primarily
influenced by genetic ancestry, followed by (in order of importance) herpesvirus infection, age,
and sex. Activation and acquisition of effector function is dependent on both the
microenvironment and differentiation state of leukocytes, and variation in these facets of basal
immunity were reflected in response to pathogen-like challenge. Stimulation with bacterial or
viral ligands showed differential transcriptional responses across the Nicaragua and Memphis
cohorts in genes important for each aspect of acute immunity. Differences between cohorts
were often dependent on the stimulatory conditions, suggesting that disparate
immunophenotypes can be context/infection dependent and further investigation of immune
variation under a wide spectrum of immune pressures is needed. Furthermore, the
transcriptional response to stimulation is largely affected by genetic ancestry. Remaining
determinants, in order of impact, include the basal transcriptional profile, herpesvirus status, and
demographic variables, age and sex. Together, these results highlight the substantial impact of
genetics on immunity, underscore the need for more studies of diverse ancestral backgrounds,
and provide a greater understanding of the relationship between the baseline immune state and
the magnitude and quality of acute immunity.
538
In further support of unique immune signatures across distinct populations, analysis of plasma
cytokines in three independent human cohorts of influenza infection showed two influenza
disease severity profiles, largely driven by age, wherein the severe pediatric response exhibits
global hyper-inflammatory cytokine levels and the severe adult response exhibits generally
decreased cytokine levels, with increases of select cytokines (IL6, TNFa, GCSF, IL8). This
indicates different therapeutic approaches may be needed and underscores the need for future
work to search for therapeutic targets across diverse populations in order to identify the most
robust and broadly applicable options. Such studies would advance the scientific and medical
communities closer to achieving personalized medicine by providing valuable insight into the
number of distinct severity profiles, the appropriate method to treat severe populations/profiles,
and the minimum number of therapeutic approaches required to effectively treat patients of all
demographics.
551
Results from individual cytokine regression models, both at baseline and during infection, show
each category of independent variables contributes to immune variation, these effects are not

mutually exclusive, the magnitudes of modulation vary by the cytokine of interest, and there are
interactive effects between and within categories. For example, herpesviruses have unique and
interactive effects on cytokine levels, suggesting independent and collective mechanisms of
immune modulation and emphasizing the need for future studies to include additional
herpesviruses in their experimental design. Interestingly, the effects of herpesviruses at
baseline are much smaller compared to the effects during acute infection with influenza. This
may be due, at least in part, to the limited transcriptional profile of herpesviruses during latency,
whereas increased transcriptional activity, perhaps in response to immune cell differentiation or
signaling, would allow for greater interference due to increased viral protein production. Indeed,
established methods for reactivating herpesviruses include stimulation with PRR ligands and
differentiation of myeloid cells.[57,58]

Of the 23 cytokine models with acute infection (study day, viral load), genetic (SNP genotypes,
AIMs), chronic infectious exposure (herpesvirus serostatus) and biological (age, sex) factors, 8
had a significant genetic component and improved model performance, each of which were
identified as a correlate of influenza disease severity. Functionally relevant SNPs, identified in
eQTL and protein cytokine level regression analyses, included SNPs with different allele
frequencies across populations and SNPs with similar allele frequencies across populations but
variations within populations, highlighting genetic contributions to both inter- and intra-
population immune variation, respectively. Importantly, variation in genotype prevalence across
or within populations may lead to different immune responses and subsequent immune
outcomes if the altered protein is a correlate of severity. For example, SNP rs3024496 was
identified as an eQTL for IL10, its distribution of genotypes varied between Memphis and
Nicaragua, and prior work in infected subjects from the Memphis cohort found IL10 to be a
correlate of hospitalization and symptom severity during influenza infection.[33] Moreover, IL10
had the second highest adjusted R^2 value in the infected regression model, and factors which
contributed to its variation, at the protein level, included SNP rs3024496, age, sex, viral load,
and herpesviruses. Luciferase assays further support the modeling results and show the
functional impact of rs3204496, such that the C allele is associated with higher promoter activity
when unstimulated or infected with influenza A virus, with the latter condition indicating a higher
magnitude of differential gene expression. This demonstrates the powerful implication of
genetics on the overall immunophenotype across ancestrally distinct populations, and supports
recent work which highlights the value in and need for increased studies of diverse, particularly
multiethnic, populations to improve our understanding of human immunology.[25,59,60]

It's postulated that immune variation can be modeled as an "immunological landscape", with
individuals lying at different coordinates according to their immune profile, this provides a
framework for the interaction of genetic, environmental, and biological factors in shaping the
human immune response.[53] In the initial description, it was suggested that the starting position
of an individual within the landscape is largely due to genetics, whereas age changes this
position in a consistent and predictable manner across subjects, and local environment may
drive convergence between two individuals if shared (such as cohabitation) or diverge two
individuals if they are from distinct environmental backgrounds.[53] The results presented here
suggest an even more complex and multidimensional immune space with dynamic landscapes

depending on the immune pressure and interactions between variables which change the
position of an individual, in addition to single variable influences. Our results show there are
clusters of distinct immunophenotypes across this landscape, both at baseline and during
infection. Importantly, the observation that gene expression differences between two diverse
populations depended on the stimulatory condition suggests that the immunological landscape
is context dependent, thus unique immunophenotype profiles and clusters may exist during
infection that were not observed at baseline. Additionally, unique correlates of severity across
diverse populations and shared correlates in cohorts with common features (e.g. age range)
suggest distinct clusters of immune profiles are associated with differential illness outcomes and
more than one severity-immunotype cluster may exist for a given disease. These observations,
and others presented here, likely would have been obscured if the study solely focused on a
single population, especially if biased in a given demographic. This underscores the need for
future studies to include diverse subjects when examining immune variation, whether at
baseline or during infection. While it is challenging to conduct studies that are diverse in all
desired categories, this study shows we can leverage multiple cohorts enriched in specific target
populations and conduct comparative analyses to gain insight into overlaps or unique signatures
of immunity as it relates to immune competence and disease severity.

Limitations of the Study

Although simultaneous investigation of multiple cohorts provided novel insight, there are several
limitations in this study. The independent designs of each cohort and differences in study
endpoints affected sampling and data collection. One of the baseline cohorts collected serum,
as opposed to plasma; however, we believe the impact was minimal, as it only applies to basal
analyses, and a study which compared plasma versus serum cytokine levels, using Luminex
multiplex assays, shows results are well correlated.[61] Differences in sampling types, timing, and
quantities limited baseline and acute infection PBMC availability for several cohorts. Thus, we
were unable to include acute cell phenotypes and connect baseline immune profiles with acute
immunophenotypes. Additionally, multiple approaches were utilized for symptom reporting and
clinical data collection, which required different statistical approaches for correlates of severity
analyses and limited our ability to include other factors, such as body mass index. Lastly, there
may be unknown differences in handling and/or storage across the multiple study sites, which
we attempted to account for by including "Cohort" as a factor in our models, when appropriate.

Main Figure Legends

Figure 1. *Study populations and model summary.* Table of cohort characteristics (top) and overview of regression model analyses (bottom, labeled A-D in order of appearance).

Figure 2. *Variation of explanatory factors across diverse populations.* Age (A) and sex (B) distribution for each cohort (n = 1,705). (C) Results from the principal component analysis of AIMs (n = 285). (D) Donut plots show the proportion of individuals in each cohort with a given genotype for a SNP of interest (n = 352), plot labels are encoded as follows: Protein-Gene-SNP. (E) Table of herpesvirus serostatus (%) by cohort (n = 1,532). (F) Venn-diagram of herpesvirus infection by cohort. Numbers in the bottom right represent the number of individuals negative for all indicated herpesviruses. For easier visualization, the herpesvirus with the highest overall seroprevalence (VZV) was excluded.

Figure 3. *Basal immune profiles in ancestrally and environmentally distinct populations.* (A) A t-SNE plot based on plasma/serum cytokines and colored by cohort (n = 631). (B) Violin plot demonstrates individual cytokine variation for EGF (n = 623). Black points in the center of the violin plot represent the mean cytokine level and line segments represent +/- the standard deviation. (C) Schematic of herpesvirus effects on basal cytokine levels based on linear regression modeling results (model displayed above diagram, n ≈ 500). Cytokines listed have an adjusted R^2 value > 0.08 and FDR < 0.05. (D) Boxplots depict differences in basal cell phenotypes in PBMC samples from healthy subjects in the Memphis (n = 28) and Nicaragua (n = 19) cohorts. Presented p-values are from Mann-Whitney U tests, post adjustment for multiple comparisons (when applicable).

Figure 4. *Distinct basal immune profiles mold response to immune challenge.* (A) Gene expression by cohort under different stim conditions for IFI16 ($n_{AllStim}$ = 903), CD86 ($n_{AllStim}$ = 906), and IFNG ($n_{AllStim}$ = 774). Displayed p-values were adjusted for multiple comparisons and are from Mann Whitney U tests. (B) Principal component analysis of baseline (unstim) gene expression colored by Cohort (n = 102). (C) Summary of linear regression Models #2 (n ≈ 188) and #3 ($n_{AllStim}$ ≈ 350) that were utilized to determine factors which contribute to baseline and post stim gene expression variation, respectively (see Summary of Models section for details). Displayed values are the number of genes that a factor significantly affected out of a total 90 genes.

Figure 5. *Acute cytokine profiles delineate two age-associated signatures of influenza disease severity.* (A) Heatmap of highlighted results from plasma (left) and nasal wash (right) cytokine correlates of severity analysis, with significant adjusted p-values displayed, for the Memphis (n = 165), Baltimore (n = 222), and Taiwan (n = 226) cohorts. (B) Boxplots show example associations between plasma or nasal wash cytokine levels and symptom severity during influenza infection for each cohort. Presented p-values are from Mann Whitney U tests, and results were adjusted for multiple comparisons by controlling the FDR (< 0.1).

Figure 6. *Herpesviruses have unique and age specific effects on influenza disease severity.* (A-B) Significant herpesvirus associations with influenza disease severity. Severe Memphis was

categorized based on lower respiratory symptom severity scores. (C-D) Schematic of
herpesvirus effects on acute cytokine levels in plasma (C, n ≈ 200) and nasal wash (D, n ≈ 300)
based on linear regression modeling results (models displayed above diagrams). Cytokines
listed have an adjusted R^2 value > 0.08 and FDR < 0.05.

Figure 7. *Transcription and protein level of IL10 is modulated by rs3024496.* (A-B) For
cytokines with an eQTL, a linear regression model was performed including: sex, age,
herpesvirus (HV) serostatus, interactions between HV serostatus and age, interactions between
HVs, study day, viral load, AIMs PC1 and PC2, and eQTL SNP genotype (n ≈ 200). The top two
results, based on adjusted R^2 value, are shown and include the coefficient estimate of the
change in the dependent variable per unit change in the independent variable (Coeff Estimate),
the corresponding standard error (Std Error), t score (Statistic), and p-value (highlighted when
significant). (C) Schematic of luciferase assays. Boxplots compare Firefly:Renilla ratios from
unstimulated (D) or influenza virus infected (E) luciferase assays for A and G alleles of IRF3
rs2304204 (left), and T and C alleles of IL10 rs3024496 (right). Presented p-values are from
Mann-Whitney U tests.

Supplemental Item Legends

Supplemental Figure 1. *Explanatory variables by Race/Ethnicity and age.* (A-C) Principal
component analysis of AIMs colored by Race and/or Ethnicity. (D) Total number of
herpesviruses by age, separated by cohort.

Supplemental Figure 2. *Cohort tSNE clusters are not associated with demographics or chronic
virus infection.* Basal cytokine tSNE plot colored by age (A), sex (B), herpesvirus serostatus (C-
F), and Race (G, for Memphis only, all other cohorts in gray). (H) Basal cytokine tSNE with
Memphis cohort data removed.

Supplemental Figure 3. *Heatmap summary of baseline cytokine regression models.* (A)
Summary of coefficient estimates for each factor included in the baseline plasma/serum linear
regression model. Cytokine models are listed in decreasing order of the adjusted R^2 value
(shown on the far right) and coefficients are colored if FDR < 0.05. Red is associated with an
increase in the cytokine level per unit change in the factor, and green is associated with a
decrease in the cytokine level. (B) Basal plasma/serum cytokine level distributions.

Supplemental Figure 4. *Highlights of gene expression associations with independent
variables.* (A-B) Example of associations based on results from Model #2 (A, determinants of
baseline gene expression) and Model #3 (B, determinants of stimulated gene expression). The
y-axis is gene expression and the x-axis is the independent variable (age in years, herpesvirus
serostatus, ancestry informative marker principal component, baseline gene expression
principal component). (C) Examples of eQTLs. *MCP1* (n = 833) and *FLT3LG* (n = 868) gene
expression by genotype for rs13900 and rs2946871, respectively.

Supplemental Figure 5. *Plasma and nasal wash cytokine correlates of severity.* Heatmap of
plasma (A) and nasal wash (B) cytokine correlates of severity analysis, with significant adjusted

p-values displayed. Colors are based on correlation coefficients or a Cohen's D effect size, depending on whether a symptom is continuous or binary, respectively. Presented p-values are from correlations (Spearman and Pearson) or Mann Whitney U tests, and results were adjusted for multiple comparisons by controlling the FDR (< 0.1).

Supplemental Figure 6. *Heatmap summary of acute plasma cytokine regression models.* Samples were from influenza infected subjects. (A) Summary of coefficient estimates for each factor included in the infected plasma linear regression model. Cytokine models are listed in decreasing order of the adjusted R^2 value (shown on the far right) and coefficients are colored if FDR < 0.05. Red is associated with an increase in the cytokine level per unit change in the factor, and green is associated with a decrease in the cytokine level. (B) Acute plasma cytokine level distributions.

Supplemental Figure 7. *Heatmap summary of acute nasal wash cytokine regression models.* Samples were from influenza infected subjects. (A) Summary of coefficient estimates for each factor included in the infected nasal wash linear regression model. Cytokine models are listed in decreasing order of the adjusted R^2 value (shown on the far right) and coefficients are colored if FDR < 0.05. Red is associated with an increase in the cytokine level per unit change in the factor, and green is associated with a decrease in the cytokine level. (B) Acute nasal wash cytokine level distributions.

Supplemental Table 1. *Immune-related SNP genotype frequency in Memphis and Nicaragua.*

Supplemental Table 2. *Results from basal cell phenotype comparisons in PBMCs from healthy Nicaraguans and Memphians.*

Supplemental Table 3. *Gene expression Mann Whitney U and linear regression model results.*

Supplemental Table 4. *Regression results for eQTL analysis.*

Supplemental Table 5. *Cytokine correlates of severity analysis results.*

Supplemental Table 6. *Biological and herpesvirus severity associations.*

Supplemental Table 7. *Results from integrative analysis of biological, genetic, and herpesvirus effects on cytokine levels during influenza infection.*

Acknowledgements

We thank the participants and clinical study support staff in each of the cohorts; Emily Walker and the Hartwell Center for processing Infinium Global Screening Array-24 v3.0 chips; Jessica Gaevert for technical assistance with RNA extractions; and ALSAC (American Lebanese Syrian Associated Charities). Select figures (Fig 1, Fig 3C, Fig 6E-F) were created using an academic license of BioRender software. This work was funded by: the Hercules Foundation – Belgium and the University of Antwerp (Methusalem funding; BOF Concerted Research Action), and the

Research Foundation Flanders, project grant G.0409.12N and 1861219N (B.O.); the National
Institute of Allergy and Infectious Diseases, US National Institutes of Health, under HHS
contract HHSN272201400007C for Johns Hopkins Center of Excellence in Influenza Research
and Surveillance (JH CEIRS) (A.P. and R.R.), HHS contract HHSN272201400006C for the St.
Jude Center of Excellence for Influenza Research and Surveillance (SJ CEIRS) (R.W., S.S.C.,
P.G.T.), HHS contract 75N93021C00016 for the St. Jude Centers of Excellence for Influenza
Research and Response (SJ CEIRR) (R.W., S.S.C., P.G.T.), HHS contract 75N93019C00052
for the Center for Influenza Vaccine Research for High Risk Populations (CIVR-HRP) of the
Collaborative Influenza Vaccine Innovation Centers (R.W., S.S.C., P.G.T.), RO1 5R01AI120997
(A.G.), and RO1 AI107625 (P.G.T.).

The content is solely the responsibility of the authors and does not necessarily represent the
official views of the National Institutes of Health.

Author Contributions

773 Conceptualization, A.S. and P.G.T.; Methodology, A.S., E.K.A., L.T., and P.G.T.; Investigation,
774 A.S., E.K.A., C.O., S.W., and T.J.; Formal Analysis, A.S., L.S., and S.P.; Visualization, A.S.;
775 Writing – Original Draft, A.S.; Writing – Review & Editing, A.S., E.K.A., and P.G.T.; Resources,
776 G.K., A.B., R.Z., K.S.S. J.C.D., S.H., G.E., P.V.D., V.V.T., B.O., R.W., S.S.C., A.P., R.R., A.G.,
777 P.G.T.; Supervision, P.G.T.; Funding Acquisition, B.O., R.W., S.S.C., A.P., R.R., A.G., P.G.T.

Declaration of Interests

779 P.G.T. has consulted or received honorarium and/or travel support from Illumina, JNJ, Pfizer,
780 and 10X. P.G.T. serves on the Scientific Advisory Board of ImmunoScape and CytoAgents. All
781 remaining authors declare no competing interests.

Data Availability

783 Supporting data can be obtained from the authors upon reasonable request.

Code Availability

785 Analyses were conducted using R, a publicly available software, as detailed in the Methods. All
786 code is available upon request.

Methods

Study Subjects

Memphis Cohort

790 Plasma, nasal wash, and PBMCs were collected from healthy and influenza infected (n = 383)
791 subjects in Memphis, Tennessee.[33] Recruited subjects were 75% Black/African American and
792 25% White/Caucasian. To be included in the study, a flu positive patient must meet the clinical

definition of influenza at the time of enrollment and symptoms must have begun within the previous 96 hours or less. Enrollment included index patients and household contacts (both flu positive and negative). Exclusion criteria included refusal to consent to nasal lavage samples, a history of receiving immunoglobulin or other blood products within the three months prior to enrollment, or any condition that in the opinion of the clinical site investigator place the subject at any risk or injury or render the subject unable to meet the study requirements. For index patients, nasal lavages and blood were collected at the day of enrollment (day 0) and days 3, 7, 10, and 28. Household contacts were also enrolled and provided nasal swabs on days 0. 3. 7, and 14, and 28, with blood on days 0 and 28. If at any point a household contact tested influenza positive, they were re-enrolled as a converted contact case and followed the same course of sample collection as the initial index patient. Symptom severity scores were collected throughout the duration of infection and were based on participant rankings using a visual analog scale and data consisted of five categories- total, upper respiratory, lower respiratory, gastrointestinal, and systemic. The peak total and lower respiratory symptom scores were utilized for severity analyses. Hospitalization status was also utilized and occurred in 25.9% of influenza positive participants. Written informed consent was obtained for participants or their guardians and written assent was acquired from age-appropriate subjects. This study was approved by the Institutional Review Boards of St. Jude Children's Research Hospital and the University of Tennessee Health Science Center/Le Bonheur Children's Hospital.

Colombia Cohorts

Healthy plasma samples were collected at a single time point from subjects in the local city of Calabazo (n = 100) and two indigenous populations, Seywiaka (n = 50) and Umandita (n = 50), which are 1.5 and 9.5 hours walk away from Calabazo, respectively.[35] Recruited subjects were 100% Hispanic/Latino. All individuals willing to participate by completing the study questionnaires and permitting the withdrawal of blood and nasal swabs were included in the study. Exclusion criteria included children who are less than two years old when baseline enrollment is performed and anyone with a known immunosuppressive condition or ongoing receipt of immunosuppressive therapy. Signed consent or assent was obtained as appropriate and this study was approved by the Indigenous Health Council, the Tropical Health Foundation ethics committee, and the St. Jude Children's Research Hospital Institutional Review Board.

Belgium Cohort

Healthy serum samples were collected at a single time point from 40 neonatal intensive care unit or pediatric unit nurses.[36] Recruited subjects were 100% White/Caucasian. Inclusion criteria included a minimum of two years working in the respective hospital unit, a maximum of one year between blood sampling and last VZV exposure for pediatric unit nurses, and a minimum of one year between blood sampling and last VZV exposure for neonatal intensive care unit nurses. Exclusion criteria included immunosuppressed state (due to disease or medication), any malignancy in the last five years, receipt of blood products within three months of blood sampling, coagulation disorders, and any condition the investigator believed might interfere with the study. Written informed consent was obtained from all study participants and this study was approved by the ethics board of the Antwerp University Hospital, Antwerp, Belgium.

Baltimore and Taiwan Cohorts

Plasma and/or nasal wash samples were collected from healthy or influenza-infected subjects recruited from the East Baltimore and Bayview campuses in Baltimore, Maryland, USA (n = 225) and the Chang Gung Memorial Hospitals in the Taipei, Taiwan metropolitan area (n = 266).[37,38,62,63] Recruited subjects from Taiwan were 100% Asian, and those from Baltimore were 67% Black/African American, 23% White/Caucasian, 1% American Indian or Alaskan Native, 9% Other. Possible index cases were recruited if, within the past 7 days of enrollment, they reported or documented fever and new or increased cough, headache, or sore throat. Cases were confirmed influenza positive if they received a positive influenza virus test result from the hospital visit (GeneXpert). Exclusion criteria included children younger than the age of five, currently incarcerated, unable to write or speak English, unable to provide informed consent, or unable to provide a telephone number for follow-up. Clinical coordinators obtained written informed consent, demographic, and clinical data. Samples were collected at the initial visit and at a 3-5 week follow-up visit. Additionally, symptom information included the presence or absence of wheezing and/or shortness of breath (WSOB), pneumonia (based on radiological or microscopic findings), hospitalization (for influenza associated illness), and administration of supplemental oxygen. In the Baltimore cohort, these symptoms occurred in 75%, 26%, 31%, and 17% of subjects, respectively. In the Taiwan cohort, the prevalence was 67%, 39%, 24%, and 19%, respectively. Due to the high prevalence of WSOB, it was excluded from downstream analyses for both cohorts. These studies were reviewed and approved by the Institutional Review Boards of Johns Hopkins School of Medicine and the Chang Gung Memorial Hospital (IRB00091667).

Nicaragua Cohort

Plasma samples were collected from healthy and influenza-infected subjects (n = 591) recruited at the Health Center Sócrates Flores Vivas (HCSFV) in Managua, Nicaragua.[34] Recruited subjects were 100% Hispanic/Latino. Inclusion criteria include that an index case must live in the HCSFV catchment area, be positive for influenza by QuickVue Influenza A+B rapid test, experience onset of symptoms within the previous 48 hours, live with at least one additional person who agrees to participate in the study, and have no household contacts who have had influenza-like illness in the previous two weeks. There are no exclusion criteria for participation in this study. Household members were also recruited and monitored for influenza infection through 5 home visits, 2-3 days apart, over a 10-14 day period. Blood samples were collected at the initial visit and a 30 day follow-up visit, and participants submitted questionnaires regarding signs and symptoms of acute respiratory illness. Participants provided written informed consent and parental permission was obtained from parents or legal guardians of children, in addition to verbal assent from children aged six years and older. This study was approved by the Institutional Review Boards at the University of Michigan, Centro Nacional de Diagnóstico y Referencia (Ministry of Health, Nicaragua), and University of California, Berkeley.

Herpesvirus Serostatus

Cytomegalovirus (CMV), Epstein-Barr virus (EBV), herpes simplex virus 1 (HSV1), herpes simplex virus 2 (HSV2), and human herpesvirus 6 (HHV6) serostatus was determined

using plasma and the following ELISA kits, respectively: GWB-892399, GWB- DA817E,
GWB-ECB3C0, GWB-C2349A, and Abnova KA1457.

Cytokine Measurement

Plasma and nasal wash cytokine, chemokine, and growth factor levels were measured
using the Luminex MAP system with the Milliplex HCYTOMAG-60K assay, and samples
were processed according to the kit manual. Prior to statistical analyses, analytes were
removed from the dataset if less than 15-40% of their values were above or below the lower
and upper limits of detection, respectively. This value was selected based on total sample
size and the prevalence of independent and dependent variables of interest.

Flow Cytometry

PBMCs were stained for the following surface markers: Panel 1) CCR7-FITC (Biolegend
Cat. 353216), CD25-APC (Biolegend Cat. 302610), CD3-PerCP eFluor710 (Thermo Cat. 46-
0037-42), CD38-BV785 (Biolegend Cat. 303530), CD4-BV650 (BD Biosciences Cat. 563875),
CD45RA-PE Dazzle/594 (Biolegend Cat. 304146), CD45RO-BV605 (Biolegend Cat. 304238),
CD69-AF700 (BD Biosciences Cat. 310922), CD8-APC Fire750 (Biolegend Cat. 344746), gd
TCR-PE/Cy7 (Biolegend Cat. 331222), PD1-BV711 (Biolegend Cat. 329928), TIGIT-PE
(Biolegend Cat. 372704), and TIM3-BV421 (BD Biosciences Cat. 565562). Panel 2) CD11b-
BV785 (Biolegend Cat. 301346), CD11c-PE/Cy7 (Biolegend Cat. 301608), CD123-BV421
(Biolegend Cat. 306018), CD14-APC eFluor780 (Thermo Cat. 47-0149-42), CD16-FITC
(Biolegend Cat. 360715), CD19-PerCP eFluor710 (Thermo Cat. 46-0198-42), CD1c-BV711
(Biolegend Cat. 331536), CD3-PerCP eFluor710 (Thermo Cat. 46-0037-42), CD56-BV605
(Biolegend Cat. 318334), CD68-APC (Biolegend Cat. 333810), CD80-PE (Biolegend Cat.
305208), and HLA DR-BV650 (Biolegend Cat. 307650). Panel 3) CD3-AF700 (Biolegend Cat.
300323), CD4-APC Fire750 (Biolegend Cat. 300559), FOXP3-PE (Biolegend Cat. 320107).
Briefly, after thawing from cryopreservation and plating in a 96-well round bottom plate, cells
were spun down and resuspended in 50µL of human Fc block (BD Biosciences Cat. 564220) in
DPBS and incubated for 10 minutes at room temperature. Afterwards, 50µL of a Live/Dead
Aqua (Tonbo Cat. 13-0870-T100) and pre-titrated surface antibody cocktail in DPBS were
added to each well and cells were incubated for 30 minutes on ice and in the dark. Additionally,
the BD Cytofix/Cytoperm kit (BD Biosciences Cat. 554714) and FOXP3 Transcription Factor
Staining Buffer set (eBioscience Cat. 00-5523-00) were used for intracellular staining of CD68-
APC (Panel 2) and FOXP3-PE (Panel 3), respectively and in accordance with the kit manual.
Cells were washed, resuspended in FACS (fluorescence-activated cell sorting) buffer and
analyzed on a BD Fortessa cell analyzer and analyzed using FlowJo software.

Microneutralization Assays

Microneutralization titers were performed according to guidelines outlined by the World
Health Organization (WHO) (Global Influenza Surveillance Network, 2011). Reported results
are the reciprocal dilution which inhibited virus growth as determined by an ELISA readout
or hemagglutination assay.

 Viral Titers

Influenza viral load was measured with quantitative reverse-transcription real-time polymerase
chain reaction (qRT-PCR) assays and RNA from nasal swabs. Nucleic acid was
extracted from samples using the MagMAXtm- 96 AI/ND Viral RNA Isolation kit on the
KingFisher magnetic processor platform. Afterwards, a 25mL qRT-PCR reaction was set up
utilizing the Qiagen One-Step RT-PCR kit, influenza A matrix or influenza B non-structural
protein primers and the protocol outlined by the WHO(Global Influenza Surveillance Network,
2011).

Human Genetic Studies

Prioritization of SNPs

The UCSC Human Genome Browser build GRCh37/19 was used to curate a list of 308 common
SNPs in genes encoding 41 cytokines measured by the Milliplex assay HCYTOMAG-60K and
for 3 transcription factors important in initiating antiviral immunity, namely IRF3, IRF7, and all 3
subunits of NFkB. Briefly, putative functional SNPs were prioritized and included the following
classes of SNPs: missense, splice, non-coding, and those which are located in the 3' or 5'
untranslated region (UTR) of the promoter region. Utilizing 1000 Genomes Project data, the
curated list of SNPs was reduced to 209 by filtering for SNPs with minor allele frequencies
(MAF) which significantly vary across ancestrally distinct populations. For a given SNP, this was
calculated using the MAFs of the African (AFR), American (AMR), Eastern Asian (EAS),
European (EUR), and South Asian (SAS) populations. A SNP was considered to significantly
vary if the standard deviation of the MAFs across these populations was greater than 5. Lastly,
the prioritized list of SNPs was reduced to 94 following a tagSNP analysis to identify the most
informative SNPs in a region based on linkage disequilibrium. The tagSNP analysis was
conducted by utilizing the SNPclip tool from the National Cancer Institute
(https://ldlink.nci.nih.gov/?tab=snpclip) and data from African (YRI), African American (ASW),
Mexican (MXL), and European (CEU) populations, as these most reflect the populations in this
study.

DNA Extraction

For the Memphis cohort, 90% of samples were extracted in a clinical HLA core at St. Jude, the
remaining 10% were extracted from peripheral blood mononuclear cells using the Zymo Quick-
DNA Microprep Plus Kit (Cat. D4074). All DNA samples for the Nicaragua cohort were extracted
from 200mL of neutrophil pellets using the Qiagen QIAamp DNA Mini Kit (Cat. 51306).
947
All samples were quantified using a Nanodrop, the average 260/280 ratio was 1.86
(range 1.16 - 2.04). For Fluidigm SNP Type assays, all samples met the minimum DNA
input requirement of 10ng/mL, and all ancestry informative marker samples had an input
of 250ng.

SNP Genotyping

Genotypes for the curated list of 94 SNPs were determined using the Fluidigm SNP Type

Assay on the BioMark HD System and 96.96 Dynamic Array integrated fluidic circuits.

SNP Type assays were designed using Fluidigm's D3 Assay Design Tool. The

assay was performed in accordance with the Fluidigm protocol, using PCR grade water

as a negative control, and including the specific target amplification step with 14 cycles of

amplification.

Ancestry Informative Markers

Genetic based ancestry was determined using the Infinium Global Screening Array-24 v3.0

BeadChip which measures 654,027 markers including multiethnic genome-wide content,

curated clinical research variants, and quality control markers. High quality DNA was submitted

to the St. Jude Hartwell Center for preparation and processing.

Gene Expression Assays

PBMC Stimulation

PBMCs from 188 subjects (94 Memphians and 94 Nicaraguans) were plated at a max of

850,000 cells per well and resuspended in 1µg/mL of lipopolysaccharide, Pam3CSK4,

transfected poly(I:C), transfected poly(dA:dT), or plain media as an unstimulated control.

Plates were incubated at 37°C for 24 hours. Following stimulation, plates were spun at

500g for 5 minutes. Supernatants were transferred to clean 96-well plates and stored at

-80°C. Pellets were resuspended in 300µL of lysis buffer and stored at -80°C in a labeled

epitube until ready for extraction.

Gene Expression

RNA was extracted using the Zymo Quick-RNA extraction kit (Cat. R1050), and stored in a 96-

well plate at -80°C until ready for PCR. Prior to running gene expression assays, stored RNA

was thawed and normalized to the highest amount of RNA possible, with respect to stimulation

condition. The input RNA was 14ng for unstimulated control, 18ng for Pam3CSK4, 40ng for

lipopolysaccharide, 50ng for transfected poly(I:C), and 50ng for transfected poly(dA:dT). The

expression of 96 immune genes was measured using the Fluidigm Fast Gene Expression

Analysis Using EvaGreen on the BioMark HD System and 96.96 Dynamic Array integrated

fluidic circuits. Primers were designed using Fluidigm's D3 Assay Design Tool. The assay was

performed in accordance with the Fluidigm protocol, including the specific target amplification

step with 16 cycles of amplification, and 2 water samples per chip as negative controls. Quality

control of gene expression results included removing samples/genes with no measurable gene

expression or samples with melting curve peaks that were more than one degree away from the

consensus peak melting temperature. Following QC steps, 4 genes were removed altogether

(*IFNL2, IL9, SFTPD, TMEM173*) and 92 remained for further analysis (2 housekeeping - *ACTB,*

GAPDH; 90 immune - *AIM2, AREG, BIK, CCL11, CCL2, CCL22, CCL3, CCL4, CCL5, CCL7,*

CD40LG, CD69, CD80, CD86, CGAS, CSF3, CX3CL1, CX3CR1, CXCL1, CXCL10, CXCL2,

DDX58, EGF, EOMES, FGF2, FLT3LG, GATA3, GZMB, IFI16, IFIT1, IFITM3, IFNA2, IFNB1,

IFNG, IFNL1, IL10, IL12A, IL12B, IL13, IL15, IL17A, IL18, IL1A, IL1B, IL1RN, IL2, IL4, IL5, IL6,

IL8, IRF1, IRF3, IRF7, IRF9, ISG15, JUN, KLRC1, KLRK1, LTA, MAVS, MX1, MX2, NFATC1,

NFKB1, NFKB2, NFKBIA, NLRP3, OAS1, PDGFA, PDGFB, PTGS2, REL, RELA, RELB,

RORC, *SELL*, *SOCS1*, *SOCS3*, *STAT1*, *STAT3*, *STAT4*, *TBK1*, *TBX21*, *TGFA*, *TGFB1*, *TNF*,

TNFAIP3, *TREX1*, *VEGFA*, *VEGFC*). Lastly, prior to conducting comparative analyses, results

were normalized again using housekeeping gene expression (*ACTB*).

Luciferase Assays

Determination of IL10-rs3024496 and IRF3-rs2304204 as functional SNPs was achieved using

luciferase assay constructs ordered from GenScript. For IL10-rs3024496, a 1,1114bp region

(chr1:206,945,768-206,946,881, GRCh37/19), encompassing the IL10 promoter, was inserted

upstream of the *luc2* gene in the pGL4.10[*luc2*] plasmid (Promega Cat. E6651). Following this

step, two final constructs were created, one with the C allele and one with the T allele for

rs3024496, by cloning a 2,364bp region (chr. 1:206,939,683-206,942,046, GRCh37/19) that

encompassed the IL10 3'UTR and the SNP of interest. For IRF3-rs2304204, a 1,394bp region

(chr19:50,168,220-50,169,613, GRCh37/19), encompassing the IRF3 promoter, was inserted

upstream of the *luc2* gene in the pGL4.10[*luc2*] plasmid (Promega Cat. E6651). As the SNP

was located within the promoter region, two versions of this construct were created and cloned,

one with the G allele and one with the A allele for rs2304204.

1009

A549s (ATCC Cat. CCL-185) were transfected with each plasmid using the TransIT LT

Transfection Reagent (Mirius Bio Cat. MIR 2300). Briefly, $1.5x10^5$ cells/well were seeded into

12-well plates, and incubated overnight at 37°C and 5% CO_2. The following day, 1µg of

pGL4.10 test plasmid and 0.01µg of pRL-TK control plasmid (Promega Cat. E2241) were added

per well, according to the TransIT LT protocol. All transfections were conducted in triplicate. At

24 hours post transfection, cells were collected and luciferase expression was measured using

the Dual-Luciferase Assay kit (Promega Cat. E1910) on the GloMax platform in accordance with

the kit protocol. For influenza stimulated results, at 24 hours post transfection, cells were

washed with PBS, 1mL of $6.0x10^5$ pfu/mL Influenza A virus (A/Singappore/INFIMH-16-

019/2016, multiplicity of infection = 4) was added to each well, and plates were incubated at

37°C and 5% CO_2 for 2 hours. Untreated control plates received infection media instead of

inoculum. Following the 2 hour incubation, inoculum was removed, cells were washed with PBS,

1mL of infection media was added to each well, and plates were incubated for an additional 24

hours at 37°C and 5% CO_2, after which luciferase expression was measured.

Statistical Analysis

Detailed descriptions of specific applied statistical approaches can be found in figure legends

and/or the main text. All analyses were performed with R using base functions (lm, p.adjust) and

generally the following packages: Rtsne, GGally, pheatmap, ggpubr, broom, formattable, and

tidyverse.[64–70] Dimension reduction techniques included principal component analysis and t-

Distributed Stochastic Neighbor Embedding. For the cytokine tSNE analysis, any missing values

were replaced with the median of that cytokine level, with respect to the cohort. Pearson and

Spearman correlations were used to determine cytokine correlates of severity for continuous

symptom outcomes. The unstimulated gene expression PCA required complete observations

(i.e. no missing values), and thus was generated from 102 subjects and 67 genes. Multivariate

analyses were conducted utilizing logistic or linear regression modeling, depending on whether

the outcome variable was binary or continuous, respectively. For all models, continuous

variables were standardized using the scale() function in R. All remaining analyses were Mann-
Whitney U or Kruskal Wallis tests, depending on whether there were two or more comparison
groups, respectively. All herpesvirus association analyses only included CMV, EBV, and HSV,
due to the overall high seroprevalence of HHV6 (%) and VZV (%). When appropriate and
indicated, p-values were adjusted for multiple comparisons using the Benjamini-Hochberg
method for controlling the false discovery rate.[71]

References

1. Tsang, J.S., Schwartzberg, P.L., Kotliarov, Y., Biancotto, A., Xie, Z., Germain, R.N., Wang, E., Olnes, M.J., Narayanan, M., Golding, H., et al. (2014). Global Analyses of Human Immune Variation Reveal Baseline Predictors of Postvaccination Responses. Cell *157*, 499–513. 10.1016/j.cell.2014.03.031.

2. Hwang, S., Kwon, A.-Y., Jeong, J.-Y., Kim, S., Kang, H., Park, J., Kim, J.-H., Han, O.J., Lim, S.M., and An, H.J. (2020). Immune gene signatures for predicting durable clinical benefit of anti-PD-1 immunotherapy in patients with non-small cell lung cancer. Sci. Rep. *10*, 643. 10.1038/s41598-019-57218-9.

3. Vujkovic-Cvijin, I., Dunham, R.M., Iwai, S., Maher, M.C., Albright, R.G., Broadhurst, M.J., Hernandez, R.D., Lederman, M.M., Huang, Y., Somsouk, M., et al. (2013). Dysbiosis of the Gut Microbiota Is Associated with HIV Disease Progression and Tryptophan Catabolism. Sci. Transl. Med. *5*, 193ra91-193ra91. 10.1126/scitranslmed.3006438.

4. Orrù, V., Steri, M., Sole, G., Sidore, C., Virdis, F., Dei, M., Lai, S., Zoledziewska, M., Busonero, F., Mulas, A., et al. (2013). Genetic Variants Regulating Immune Cell Levels in Health and Disease. Cell *155*, 242–256. 10.1016/j.cell.2013.08.041.

5. Drake, K.A., Torgerson, D.G., Gignoux, C.R., Galanter, J.M., Roth, L.A., Huntsman, S., Eng, C., Oh, S.S., Yee, S.W., Lin, L., et al. (2014). A genome-wide association study of bronchodilator response in Latinos implicates rare variants. J. Allergy Clin. Immunol. *133*, 370-378.e15. 10.1016/j.jaci.2013.06.043.

6. Ye, C.J., Feng, T., Kwon, H.-K., Raj, T., Wilson, M.T., Asinovski, N., McCabe, C., Lee, M.H., Frohlich, I., Paik, H., et al. (2014). Intersection of population variation and autoimmunity genetics in human T cell activation. Science *345*. 10.1126/science.1254665.

7. Brodin, P., Jojic, V., Gao, T., Bhattacharya, S., Angel, C.J.L., Furman, D., Shen-Orr, S., Dekker, C.L., Swan, G.E., Butte, A.J., et al. (2015). Variation in the Human Immune System Is Largely Driven by Non-Heritable Influences. Cell *160*, 37–47. 10.1016/j.cell.2014.12.020.

8. Carr, E.J., Dooley, J., Garcia-Perez, J.E., Lagou, V., Lee, J.C., Wouters, C., Meyts, I., Goris, A., Boeckxstaens, G., Linterman, M.A., et al. (2016). The cellular composition of the human immune system is shaped by age and cohabitation. Nat. Immunol. *17*, 461–468. 10.1038/ni.3371.

9. Lu, Y., Biancotto, A., Cheung, F., Remmers, E., Shah, N., McCoy, J.P., and Tsang, J.S. (2016). Systematic Analysis of Cell-to-Cell Expression Variation of T Lymphocytes in a Human Cohort Identifies Aging and Genetic Associations. Immunity *45*, 1162–1175. 10.1016/j.immuni.2016.10.025.

10. Li, Y., Oosting, M., Smeekens, S.P., Jaeger, M., Aguirre-Gamboa, R., Le, K.T.T., Deelen, P., Ricaño-Ponce, I., Schoffelen, T., Jansen, A.F.M., et al. (2016). A Functional Genomics Approach to Understand Variation in Cytokine Production in Humans. Cell *167*, 1099-1110.e14. 10.1016/j.cell.2016.10.017.

11. Molony, R.D., Nguyen, J.T., Kong, Y., Montgomery, R.R., Shaw, A.C., and Iwasaki, A. (2017). Aging impairs both primary and secondary RIG-I signaling for interferon induction in human monocytes. Sci. Signal. *10*, eaan2392. 10.1126/scisignal.aan2392.

12. Allen, E.K., Randolph, A.G., Bhangale, T., Dogra, P., Ohlson, M., Oshansky, C.M., Zamora, A.E., Shannon, J.P., Finkelstein, D., Dressen, A., et al. (2017). SNP-mediated disruption of CTCF binding at the IFITM3 promoter is associated with risk of severe influenza in humans. Nat. Med. *23*, 975–983. 10.1038/nm.4370.

13. Henrickson, S.E., Manne, S., Dolfi, D.V., Mansfield, K.D., Parkhouse, K., Mistry, R.D., Alpern, E.R., Hensley, S.E., Sullivan, K.E., Coffin, S.E., et al. (2018). Genomic Circuitry Underlying Immunological Response to Pediatric Acute Respiratory Infection. Cell Rep. *22*, 411–426. 10.1016/j.celrep.2017.12.043.

14. Gignoux, C.R., Torgerson, D.G., Pino-Yanes, M., Uricchio, L.H., Galanter, J., Roth, L.A.,

Eng, C., Hu, D., Nguyen, E.A., Huntsman, S., et al. (2019). An admixture mapping meta-analysis implicates genetic variation at 18q21 with asthma susceptibility in Latinos. J. Allergy Clin. Immunol. *143*, 957–969. 10.1016/j.jaci.2016.08.057.

15. Lim, H.K., Huang, S.X.L., Chen, J., Kerner, G., Gilliaux, O., Bastard, P., Dobbs, K., Hernandez, N., Goudin, N., Hasek, M.L., et al. (2019). Severe influenza pneumonitis in children with inherited TLR3 deficiency. J. Exp. Med. *216*, 2038–2056. 10.1084/jem.20181621.

16. Kotliarov, Y., Sparks, R., Martins, A.J., Mulè, M.P., Lu, Y., Goswami, M., Kardava, L., Banchereau, R., Pascual, V., Biancotto, A., et al. (2020). Broad immune activation underlies shared set point signatures for vaccine responsiveness in healthy individuals and disease activity in patients with lupus. Nat. Med. *26*, 618–629. 10.1038/s41591-020-0769-8.

17. Herati, R.S., Silva, L.V., Vella, L.A., Muselman, A., Alanio, C., Bengsch, B., Kurupati, R.K., Kannan, S., Manne, S., Kossenkov, A.V., et al. (2021). Vaccine-induced ICOS+CD38+ circulating Tfh are sensitive biosensors of age-related changes in inflammatory pathways. Cell Rep. Med. *2*. 10.1016/j.xcrm.2021.100262.

18. Brenes-Chacon, H., Garcia-Mauriño, C., Moore-Clingenpeel, M., Mertz, S., Ye, F., Cohen, D.M., Ramilo, O., and Mejias, A. (2021). Age-dependent Interactions Among Clinical Characteristics, Viral Loads and Disease Severity in Young Children With Respiratory Syncytial Virus Infection. Pediatr. Infect. Dis. J. *40*, 116–122. 10.1097/INF.0000000000002914.

19. Erwin, E.A., Jaramillo, L.M., Smith, B., Kruszewski, P.G., Kahwash, B., Grayson, M.H., Mejias, A., and Ramilo, O. (2021). Sex Differences in Blood Transcriptional Profiles and Clinical Phenotypes in Pediatric Patients with Eosinophilic Esophagitis. J. Allergy Clin. Immunol. Pract. *9*, 3350-3358.e8. 10.1016/j.jaip.2021.06.043.

20. O'Neill, M.B., Quach, H., Pothlichet, J., Aquino, Y., Bisiaux, A., Zidane, N., Deschamps, M., Libri, V., Hasan, M., Zhang, S.-Y., et al. (2021). Single-Cell and Bulk RNA-Sequencing Reveal Differences in Monocyte Susceptibility to Influenza A Virus Infection Between Africans and Europeans. Front. Immunol. *12*.

21. Randolph, H.E., Fiege, J.K., Thielen, B.K., Mickelson, C.K., Shiratori, M., Barroso-Batista, J., Langlois, R.A., and Barreiro, L.B. (2021). Genetic ancestry effects on the response to viral infection are pervasive but cell type specific. Science. 10.1126/science.abg0928.

22. Alanio, C., Verma, A., Mathew, D., Gouma, S., Liang, G., Dunn, T., Oldridge, D.A., Weaver, J.E., Kuri-Cervantes, L., Pampena, M.B., et al. (2022). Cytomegalovirus latent infection is associated with an increased risk of COVID-19-related hospitalization. J. Infect. Dis., jiac020. 10.1093/infdis/jiac020.

23. Schönfelder, K., Breuckmann, K., Elsner, C., Dittmer, U., Fistera, D., Herbstreit, F., Risse, J., Schmidt, K., Sutharsan, S., Taube, C., et al. (2021). The influence of IFITM3 polymorphisms on susceptibility to SARS-CoV-2 infection and severity of COVID-19. Cytokine *142*, 155492. 10.1016/j.cyto.2021.155492.

24. Cuesta-Llavona, E., Albaiceta, G.M., García-Clemente, M., Duarte-Herrera, I.D., Amado-Rodríguez, L., Hermida-Valverde, T., Enríquez-Rodriguez, A.I., Hernández-González, C., Melón, S., Alvarez-Argüelles, M.E., et al. (2021). Association between the interferon-induced transmembrane protein 3 gene (IFITM3) rs34481144 / rs12252 haplotypes and COVID-19. Curr. Res. Virol. Sci. *2*, 100016. 10.1016/j.crviro.2021.100016.

25. Graham, S.E., Clarke, S.L., Wu, K.-H.H., Kanoni, S., Zajac, G.J.M., Ramdas, S., Surakka, I., Ntalla, I., Vedantam, S., Winkler, T.W., et al. (2021). The power of genetic diversity in genome-wide association studies of lipids. Nature *600*, 675–679. 10.1038/s41586-021-04064-3.

26. Ishigaki, K., Sakaue, S., Terao, C., Luo, Y., Sonehara, K., Yamaguchi, K., Amariuta, T., Too, C.L., Laufer, V.A., Scott, I.C., et al. (2022). Multi-ancestry genome-wide association analyses identify novel genetic mechanisms in rheumatoid arthritis. Nat. Genet. *54*, 1640–

1651. 10.1038/s41588-022-01213-w.
27. Mersha, T.B., and Abebe, T. (2015). Self-reported race/ethnicity in the age of genomic
research: its potential impact on understanding health disparities. Hum. Genomics *9*, 1.
10.1186/s40246-014-0023-x.
28. Borrell, L.N., Elhawary, J.R., Fuentes-Afflick, E., Witonsky, J., Bhakta, N., Wu, A.H.B.,
Bibbins-Domingo, K., Rodríguez-Santana, J.R., Lenoir, M.A., Gavin, J.R., et al. (2021).
Race and Genetic Ancestry in Medicine — A Time for Reckoning with Racism. N. Engl. J.
Med. *384*, 474–480. 10.1056/NEJMms2029562.
29. Virgin, H.W., Wherry, E.J., and Ahmed, R. (2009). Redefining Chronic Viral Infection. Cell
138, 30–50. 10.1016/j.cell.2009.06.036.
30. Kahan, S.M., Wherry, E.J., and Zajac, A.J. (2015). T cell exhaustion during persistent viral
infections. Virology *479–480*, 180–193. 10.1016/j.virol.2014.12.033.
31. Souquette, A., and Thomas, P.G. (2018). Past Life and Future Effects—How Heterologous
Infections Alter Immunity to Influenza Viruses. Front. Immunol. *9*, 1071.
10.3389/fimmu.2018.01071.
32. Sehrawat, S., Kumar, D., and Rouse, B.T. (2018). Herpesviruses: Harmonious Pathogens
but Relevant Cofactors in Other Diseases? Front. Cell. Infect. Microbiol. *8*.
33. Oshansky, C.M., Gartland, A.J., Wong, S.-S., Jeevan, T., Wang, D., Roddam, P.L., Caniza,
M.A., Hertz, T., Devincenzo, J.P., Webby, R.J., et al. (2014). Mucosal immune responses
predict clinical outcomes during influenza infection independently of age and viral load. Am.
J. Respir. Crit. Care Med. *189*, 449–462. 10.1164/rccm.201309-1616OC.
34. Ng, S., Lopez, R., Kuan, G., Gresh, L., Balmaseda, A., Harris, E., and Gordon, A. (2016).
The Timeline of Influenza Virus Shedding in Children and Adults in a Household
Transmission Study of Influenza in Managua, Nicaragua. Pediatr. Infect. Dis. J. *35*, 583–
586. 10.1097/INF.0000000000001083.
35. Altan, E., Dib, J.C., Gulloso, A.R., Juandigua, D.E., Deng, X., Bruhn, R., Hildebrand, K.,
Freiden, P., Yamamoto, J., Schultz-Cherry, S., et al. (2019). Effect of Geographic Isolation
on the Nasal Virome of Indigenous Children. J. Virol. *93*. 10.1128/JVI.00681-19.
36. Elias, G., Souquette, A., Heynderickx, S., Meester, I.D., Jansens, H., Beutels, P., Damme,
P.V., Smits, E., Thomas, P.G., Tendeloo, V.V., et al. (2018). Altered CD4+ T cell immunity
in nurses occupationally exposed to viral pathogens. Clin. Exp. Immunol. *194*, 192–204.
10.1111/cei.13193.
37. Liu, H., Gong, Y.-N., Shaw-Saliba, K., Mehoke, T., Evans, J., Liu, Z.-Y., Lewis, M., Sauer,
L., Thielen, P., Rothman, R., et al. (2021). Differential disease severity and whole-genome
sequence analysis for human influenza A/H1N1pdm virus in 2015–2016 influenza season.
Virus Evol. *7*, veab044. 10.1093/ve/veab044.
38. Blumenkrantz, D.R., Mehoke, T., Shaw-Saliba, K., Powell, H., Wohlgemuth, N., Liu, H.,
Macias, E., Evans, J., Lewis, M., Medina, R., et al. (2021). Identification of H3N2 NA and
PB1-F2 genetic variants and their association with disease symptoms during the 2014–15
influenza season. Virus Evol. *7*, veab047. 10.1093/ve/veab047.
39. Mudd, P.A., Crawford, J.C., Turner, J.S., Souquette, A., Reynolds, D., Bender, D.,
Bosanquet, J.P., Anand, N.J., Striker, D.A., Martin, R.S., et al. (2020). Distinct inflammatory
profiles distinguish COVID-19 from influenza with limited contributions from cytokine storm.
Sci. Adv., eabe3024. 10.1126/sciadv.abe3024.
40. Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., Zhang, L., Fan, G., Xu, J., Gu, X., et
al. (2020). Clinical features of patients infected with 2019 novel coronavirus in Wuhan,
China. The Lancet *395*, 497–506. 10.1016/S0140-6736(20)30183-5.
41. Hadjadj, J., Yatim, N., Barnabei, L., Corneau, A., Boussier, J., Smith, N., Péré, H., Charbit,
B., Bondet, V., Chenevier-Gobeaux, C., et al. (2020). Impaired type I interferon activity and
inflammatory responses in severe COVID-19 patients. Science *369*, 718–724.
10.1126/science.abc6027.

42. Bonnet, B., Cosme, J., Dupuis, C., Coupez, E., Adda, M., Calvet, L., Fabre, L., Saint-
Sardos, P., Bereiziat, M., Vidal, M., et al. (2021). Severe COVID-19 is characterized by the
co-occurrence of moderate cytokine inflammation and severe monocyte dysregulation.
eBioMedicine *73*. 10.1016/j.ebiom.2021.103622.
43. Mathew, D., Giles, J.R., Baxter, A.E., Oldridge, D.A., Greenplate, A.R., Wu, J.E., Alanio, C.,
Kuri-Cervantes, L., Pampena, M.B., D'Andrea, K., et al. (2020). Deep immune profiling of
COVID-19 patients reveals distinct immunotypes with therapeutic implications. Science *369*,
eabc8511. 10.1126/science.abc8511.
44. Furman, D., Jojic, V., Sharma, S., Shen-Orr, S.S., Angel, C.J.L., Onengut-Gumuscu, S.,
Kidd, B.A., Maecker, H.T., Concannon, P., Dekker, C.L., et al. (2015). Cytomegalovirus
infection enhances the immune response to influenza. Sci. Transl. Med. *7*, 281ra43-
281ra43. 10.1126/scitranslmed.aaa2293.
45. Mettelman, R.C., and Thomas, P.G. (2020). Human Susceptibility to Influenza Infection and
Severe Disease. Cold Spring Harb. Perspect. Med., a038711.
10.1101/cshperspect.a038711.
46. Wikby, A., Johansson, B., Olsson, J., Löfgren, S., Nilsson, B.-O., and Ferguson, F. (2002).
Expansions of peripheral blood CD8 T-lymphocyte subpopulations and an association with
cytomegalovirus seropositivity in the elderly: the Swedish NONA immune study. Exp.
Gerontol. *37*, 445–453. 10.1016/S0531-5565(01)00212-1.
47. Savva, G.M., Pachnio, A., Kaul, B., Morgan, K., Huppert, F.A., Brayne, C., and Moss, P.A.H.
(2013). Cytomegalovirus infection is associated with increased mortality in the older
population. Aging Cell *12*, 381–387. 10.1111/acel.12059.
48. Wang, G.C., Kao, W.H.L., Murakami, P., Xue, Q.-L., Chiou, R.B., Detrick, B., McDyer, J.F.,
Semba, R.D., Casolaro, V., Walston, J.D., et al. (2010). Cytomegalovirus Infection and the
Risk of Mortality and Frailty in Older Women: A Prospective Observational Cohort Study.
Am. J. Epidemiol. *171*, 1144–1152. 10.1093/aje/kwq062.
49. Netea, M.G., Joosten, L.A.B., Latz, E., Mills, K.H.G., Natoli, G., Stunnenberg, H.G., O'Neill,
L.A.J., and Xavier, R.J. (2016). Trained immunity: A program of innate immune memory in
health and disease. Science *352*. 10.1126/science.aaf1098.
50. Tu, W., and Rao, S. (2016). Mechanisms Underlying T Cell Immunosenescence: Aging and
Cytomegalovirus Infection. Front. Microbiol. *7*.
51. Souquette, A., Frere, J., Smithey, M., Sauce, D., and Thomas, P.G. (2017). A constant
companion: immune recognition and response to cytomegalovirus with aging and
implications for immune fitness. GeroScience *39*, 293–303. 10.1007/s11357-017-9982-x.
52. Jergović, M., Contreras, N.A., and Nikolich-Žugich, J. (2019). Impact of CMV upon immune
aging: facts and fiction. Med. Microbiol. Immunol. (Berl.) *208*, 263–269. 10.1007/s00430-
019-00605-w.
53. Liston, A., Carr, E.J., and Linterman, M.A. (2016). Shaping Variation in the Human Immune
System. Trends Immunol. *37*, 637–646. 10.1016/j.it.2016.08.002.
54. Evans, D.M., Frazer, I.H., and Martin, N.G. (1999). Genetic and environmental causes of
variation in basal levels of blood cells. Twin Res. Hum. Genet. *2*, 250–257.
10.1375/twin.2.4.250.
55. Roederer, M., Quaye, L., Mangino, M., Beddall, M.H., Mahnke, Y., Chattopadhyay, P., Tosi,
I., Napolitano, L., Terranova Barberio, M., Menni, C., et al. (2015). The Genetic Architecture
of the Human Immune System: A Bioresource for Autoimmunity and Disease Pathogenesis.
Cell *161*, 387–403. 10.1016/j.cell.2015.02.046.
56. Lakshmikanth, T., Muhammad, S.A., Olin, A., Chen, Y., Mikes, J., Fagerberg, L.,
Gummesson, A., Bergström, G., Uhlen, M., and Brodin, P. (2020). Human Immune System
Variation during 1 Year. Cell Rep. *32*, 107923. 10.1016/j.celrep.2020.107923.
57. Cook, C.H., Trgovcich, J., Zimmerman, P.D., Zhang, Y., and Sedmak, D.D. (2006).
Lipopolysaccharide, Tumor Necrosis Factor Alpha, or Interleukin-1β Triggers Reactivation of

Latent Cytomegalovirus in Immunocompetent Mice. J. Virol. *80*, 9151–9158.
10.1128/JVI.00216-06.
58. Sinclair, J. (2008). Human cytomegalovirus: Latency and reactivation in the myeloid lineage.
J. Clin. Virol. *41*, 180–185. 10.1016/j.jcv.2007.11.014.
59. Pino-Yanes, M., Gignoux, C.R., Galanter, J.M., Levin, A.M., Campbell, C.D., Eng, C.,
Huntsman, S., Nishimura, K.K., Gourraud, P.-A., Mohajeri, K., et al. (2015). Genome-wide
association study and admixture mapping reveal new loci associated with total IgE levels in
Latinos. J. Allergy Clin. Immunol. *135*, 1502–1510. 10.1016/j.jaci.2014.10.033.
60. Peng, K., Safonova, Y., Shugay, M., Popejoy, A.B., Rodriguez, O.L., Breden, F., Brodin, P.,
Burkhardt, A.M., Bustamante, C., Cao-Lormeau, V.-M., et al. (2021). Diversity in
immunogenomics: the value and the challenge. Nat. Methods *18*, 588–591.
10.1038/s41592-021-01169-5.
61. Rosenberg-Hasson, Y., Hansmann, L., Liedtke, M., Herschmann, I., and Maecker, H.T.
(2014). Effects of serum and plasma matrices on multiplex immunoassays. Immunol. Res.
58, 224–233. 10.1007/s12026-014-8491-6.
62. Fink, A.L., Liu, H., Shaw-Saliba, K., Mehoke, T., Evans, J., Liu, Z.-Y., Lewis, M., Sauer, L.,
Thielen, P., Chen, K.-F., et al. (2021). Regional differences in vaccine uptake and
serological responses to vaccine and circulating strains of H1N1 viruses among patients
with confirmed influenza. J. Clin. Virol. Plus *1*, 100034. 10.1016/j.jcvp.2021.100034.
63. Zapf, A.J., Hardick, J., McBryde, B., Sauer, L.M., Fenstermacher, K.Z.J., Ricketts, E.P., Lin,
Y.-C., Chen, K.-F., Hsieh, Y.-H., Dugas, A., et al. (2022). Impact of coinfection status and
comorbidity on disease severity in adult emergency department patients with influenza B.
Influenza Other Respir. Viruses *16*, 236–246. 10.1111/irv.12907.
64. R Core Team (2021). R: A Language and Environment for Statistical Computing.
65. Krijthe, J.H. (2015). Rtsne: T-Distributed Stochastic Neighbor Embedding using Barnes-Hut.
66. Kolde, R. (2019). pheatmap: Pretty Heatmaps.
67. Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L.D., François, R., Grolemund,
G., Hayes, A., Henry, L., Hester, J., et al. (2019). Welcome to the Tidyverse. J. Open
Source Softw. *4*, 1686. 10.21105/joss.01686.
68. David Robinson, Alex Hayes, and Simon Couch (2021). broom: Convert Statistical Objects
into Tidy Tibbles.
69. Kun Ren and Kenton Russell (2021). formattable: Create "Formattable" Data Structures.
70. Schloerke, B., Cook, D., Larmarange, J., Briatte, F., Marbach, M., Thoen, E., Elberg, A.,
Toomet, O., Crowley, J., Hofmann, H., et al. (2021). GGally: Extension to "ggplot2."
71. Benjamini, Y., and Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical
and Powerful Approach to Multiple Testing. J. R. Stat. Soc. Ser. B Methodol. *57*, 289–300.

Study n = 1,705	Unit/Key	Memphis n = 383	Nicaragua n = 591	Baltimore n = 225	Taiwan n = 266	Colombia n = 200	Belgium n = 40
Age	Median (Range)	17 (0-71)	11 (0-84)	43 (18-89)	41 (20-92)	13 (2-77)	42 (22-53)
Sex	% Female ♀♂	61.4	55.7	56.0	57.9	56.5	85
Subjects Included	Healthy / Influenza Infected	Healthy, Infected	Healthy, Infected	Healthy, Infected	Healthy, Infected	Healthy	Healthy
Sample Types	Nasal Wash, Plasma, PBMC	Nasal Wash, Plasma, PBMC	Plasma, PBMC	Plasma	Plasma	Plasma	Plasma
Timepoints	Single or Multiple	Multiple	Single	Single	Single	Single	Single
Symptoms	Continuous (Scale) or Binary (presence/absence)	Continuous	Binary	Binary	Binary	-	-
Analysis	Cytokine Severity Correlates / Baseline Immunity / Acute Immunity	Cytokine, Baseline, Acute	Cytokine, Baseline	Cytokine, Baseline	Cytokine, Baseline	Baseline	Baseline

Figure 1. Study populations and model summary. Table of cohort characteristics (top) and overview of regression model analyses (bottom, labeled A-D in order of appearance).

Figure 2. Variation of explanatory factors across diverse populations. Age (A) and sex (B) distribution for each cohort (n = 1,705). (C) Results from the principal component analysis of AIMs (n = 285). (D) Donut plots show the proportion of individuals in each cohort with a given genotype for a SNP of interest (n = 352), plot labels are encoded as follows: Protein-Gene-SNP. (E) Table of herpesvirus serostatus (%) by cohort (n = 1,532). (F) Venn-diagram of herpesvirus infection by cohort. Numbers in the bottom right represent the number of individuals negative for all indicated herpesviruses. For e sier visualization, the herpesvirus with the highest overall seroprevalence (VZV) was excluded.

Figure 3. Basal immune profiles in ancestrally and environmentally distinct populations.
(A) A t-SNE plot based on plasma/serum cytokines and colored by cohort (n = 631). (B) Violin plot demonstrates individual cytokine variation for EGF (n = 623). Black points in the center of the violin plot represent the mean cytokine level and line segments represent +/- the standard deviation.
(C) Schematic of herpesvirus effects on basal cytokine levels based on linear regression modeling results (model displayed above diagram, n ≈ 500). Cytokines listed have an adjusted R² value > 0.08 and FDR < 0.05. (D) Boxplots depict differences in basal cell phenotypes in PBMC samples from healthy subjects in the Memphis (n = 28) and Nicaragua (n = 19) cohorts. Presented p-values are from Mann-Whitney U tests, post adjustment for multiple comparisons (when applicable).

Figure 4. Distinct basal immune profiles mold response to immune challenge. (A) Gene expression by cohort under different stim conditions for *IFI16* ($n_{AllStim}$ = 903), *CD86* ($n_{AllStim}$ = 906), and *IFNG* ($n_{AllStim}$ = 774). Displayed p-values were adjusted for multiple comparisons and are from Mann Whitney U tests. (B) Principal component analysis of baseline (unstim) gene expression colored by Cohort (n = 102). (C) Summary of linear regression Models #2 (n ≈ 188) and #3 ($n_{AllStim}$ ≈ 350) that were utilized to determine factors which contribute to baseline and post stim gene expression variation, respectively (see Summary of Models section for details). Displayed values are the number of genes that a factor significantly affected out of a total 90 genes.

Figure 5. Acute cytokine profiles delineate two age-associated signatures of influenza disease severity. (A) Heatmap of highlighted results from plasma (left) and nasal wash (right) cytokine correlates of severity analysis, with significant adjusted p-values displayed, for the Memphis (n = 165), Baltimore (n = 222), and Taiwan (n = 226) cohorts. (B) Boxplots show example associations between plasma or nasal wash cytokine levels and symptom severity during influenza infection for each cohort. Presented p-values are from Mann Whitney U tests, and results were adjusted for multiple comparisons by controlling the FDR (< 0.1).

Figure 6. Herpesviruses have unique and age specific effects on influenza disease severity. (A-B) Significant herpesvirus associations with influenza disease severity. Severe Memphis was categorized based on lower respiratory symptom severity scores. (C-D) Schematic of herpesvirus effects on acute cytokine levels in plasma (C, n ≈ 200) and nasal wash (D, n ≈ 300) based on linear regression modeling results (models displayed above diagrams). Cytokines listed have an adjusted R^2 value > 0.08 and FDR < 0.05.

Figure 7. Transcription and protein level of IL10 is modulated by rs3024496. (A-B) For cytokines with an eQTL, a linear regression model was performed including: sex, age, herpesvirus (HV) serostatus, interactions between HV serostatus and age, interactions between HVs, study day, viral load, AIMs PC1 and PC2, and eQTL SNP genotype (n ≈ 200). The top two results, based on adjusted R² value, are shown and include the coefficient estimate of the change in the dependent variable per unit change in the independent variable (Coeff Estimate), the corresponding standard error (Std Error), t score (Statistic), and p-value (highlighted when significant). (C) Schematic of luciferase assays. Boxplots compare Firefly:Renilla ratios from unstimulated (D) or influenza virus infected (E) luciferase assays for A and G alleles of IRF3 rs2304204 (left), and T and C alleles of IL10 rs3024496 (right). Presented p-values are from Mann-Whitney U tests.

Supplemental Figure 1. Explanatory variables by Race/Ethnicity and age. (A-C) Principal component analysis of AIMs colored by Race and/or Ethnicity. (D) Total number of herpesviruses by age, separated by cohort.

Supplemental Figure 2. Cohort tSNE clusters are not associated with demographics or chronic virus infection. Basal cytokine tSNE plot colored by age (A), sex (B), herpesvirus serostatus (C-F), and Race (G, for Memphis only, all other cohorts in gray). (H) Basal cytokine tSNE with Memphis cohort data removed.

A

Cytokine	Age	Sex.Male	CMVPos	Age:CMVPos	HSVPos	Age:HSVPos	EBVPos	Age:EBVPos	EBVPos:CMVPos	CMVPos:HSVPos	EBVPos:HSVPos	EBVPos:CMVPos:HSVPos	Cohort Belgium	Cohort Calabazo	Cohort Memphis	Cohort Nicaragua	Cohort Seywiaka	Cohort Taiwan	Cohort Umandita	Adj.R2
IL4	-0.1088	-0.1187	0.0732	-0.0120	0.0176	0.0470	0.2812	0.0541	-0.4330	0.1111	-0.2029	0.1721	-1.8587	-1.8428	-1.7906	-1.8317	-1.9067	-1.2541	-1.9483	0.4709
GRO	-0.6083	-0.1828	0.5188	0.0176	0.6678	0.1884	0.9633	0.3739	-0.6153	-0.1712	-0.8587	0.4887	-1.2974	-1.4589	-3.3088	-1.4349	-2.4034	-1.2187	-2.6628	0.4569
sCD40L	-0.1498	0.0320	0.2733	-0.0063	0.4675	-0.0766	0.3363	0.1580	-0.0530	-0.2256	-0.4437	0.0684	0.0267	-1.8331	-1.3420	-0.1152	-2.1052	-0.5331	-1.8609	0.3721
EGF	-0.3680	-0.0638	0.1286	-0.1198	-0.0218	-0.0842	0.1586	0.4614	0.1728	-0.2553	-0.1412	0.1270	0.5745	-0.8113	-1.8451	0.3987	-1.0405	-0.6201	-1.3360	0.3222
TNFa	-0.1924	0.1087	-0.3801	-0.1434	-0.2171	0.2375	-0.0024	0.0227	0.3725	0.4636	-0.0266	-0.3256	-0.3021	-0.6190	-0.8691	0.3070	0.0880	0.4516	0.0620	0.3145
MIP1b	0.0118	-0.0054	0.1301	-0.2592	-0.1558	0.0151	0.2618	0.1504	-0.0750	0.0339	-0.1831	-0.0257	-0.7073	-1.3946	-0.3720	0.2491	-1.0017	0.2679	-1.1129	0.2908
GCSF	0.2368	-0.1938	-0.3033	-0.1047	-0.3581	0.0370	-0.4705	-0.1489	0.3321	0.3941	0.4278	-0.4806	-1.3700	-1.2568	-0.6437	0.3056	-1.2812	0.1689	-1.0055	0.2707
FGF2	0.0477	-0.1376	-0.1519	-0.1824	-0.4421	0.0613	-0.3831	0.0135	0.1077	0.0934	0.3625	-0.1602	-0.4969	-0.5252	-0.5932	0.7341	-0.1671	0.2722	-0.4174	0.2456
IL15	0.1659	0.0308	0.1109	-0.1188	-0.2467	0.0471	-0.2469	-0.0874	-0.3082	-0.1951	0.1223	0.4140	-0.5866	-0.6610	-0.4542	-0.0102	-0.6615	0.3304	-0.6074	0.2446
IFNg	0.5748	-0.0621	-0.0782	-0.3273	-0.5823	-0.2260	-0.5147	-0.1701	-0.0694	-0.2742	0.4063	0.3563	-0.0580	-0.9925	0.4446	0.3864	-1.0031	0.7206	-1.0720	0.2373
Eotaxin	0.4972	0.0587	0.0545	-0.1183	0.0555	-0.0861	0.2356	-0.2032	0.1275	-0.2862	-0.1655	0.2181	0.0882	0.0130	-0.7203	0.1396	0.2171	0.0646	0.3574	0.2262
MCP1	0.0113	0.0475	-0.4893	-0.1279	-0.2861	0.0624	-0.2374	0.1047	0.6307	0.5500	0.1808	-0.6520	0.1039	-0.3207	-0.6930	0.1921	-0.0696	0.0964	0.0760	0.2251
IL10	-0.0720	0.1462	-0.1937	-0.0798	-0.0833	0.4251	-0.3078	-0.2187	0.1091	0.4206	0.0041	-0.2021	0.1391	0.0583	0.0943	0.5325	0.3709	0.9496	0.2144	0.2187
MIP1a	-0.1897	0.0489	-0.2085	-0.1010	0.2399	0.0405	0.3956	0.2503	0.2535	0.0191	-0.4526	0.0301	0.0186	-0.8508	0.4048	0.2604	-0.8784	-0.6070	-0.8068	0.2172
GMCSF	-0.0420	-0.1299	-0.3589	-0.2353	-0.2667	0.1421	-0.1504	0.0603	0.6074	0.2353	0.2579	-0.4960	-0.5482	-0.7287	0.5406	0.6355	-0.5638	0.2539	-0.6541	0.1934
TGFa	0.1096	0.0475	-0.2944	-0.0777	-0.2855	-0.0444	-0.1424	-0.0366	0.2444	0.3439	0.2182	-0.3986	0.4111	-0.2401	-0.1366	0.8170	0.2590	-0.0253	-0.1227	0.1914
MDC	-0.1701	0.0475	-0.0030	-0.0357	0.0114	0.0750	0.0614	0.0291	0.0310	0.0788	0.0274	-0.0529	-0.5556	-0.6780	-0.3266	-0.3886	-0.8918	-0.4388	-0.8713	0.1854
IL1RA	-0.1225	-0.2492	-0.3479	-0.3561	-0.1754	0.0903	-0.2355	0.1782	0.1426	0.6326	-0.0311	-0.3671	-0.8538	0.7046	-0.5987	1.1648	0.0248	0.7523	-0.4907	0.1841
IFNa2	-0.1405	-0.0134	-0.6085	-0.2437	-0.7392	0.2623	-0.6239	0.0823	0.6951	0.7066	0.6490	-0.7791	-0.5000	-0.5509	0.0846	0.9336	-0.5658	0.2821	-0.2326	0.1797
Fractalkine	-0.1954	-0.0797	0.3100	-0.1096	0.0769	0.2075	-0.0076	-0.0092	-0.1755	-0.3041	-0.0156	0.0794	-0.3072	-0.6019	-0.1779	0.7816	-0.7219	-0.1262	-0.4398	0.1509
Flt3.Ligand	0.0986	0.0026	0.4639	-0.1211	-0.4175	-0.0003	-0.5485	0.0870	-0.2417	-0.4601	0.5555	0.2208	0.0102	-0.5042	-0.2653	0.7390	-0.2370	-0.2853	-0.3680	0.1497
IL17A	0.4172	-0.0808	0.2671	-0.1709	-0.3182	-0.0586	-0.4994	-0.1961	-0.3258	-0.8089	0.2834	0.7680	0.2855	-0.6726	-0.2004	0.6091	-0.7006	0.2974	-0.6203	0.1465
VEGF	0.0869	-0.0136	0.4686	-0.1109	-0.0462	0.0133	0.0006	0.0578	-0.4046	-0.4549	0.0484	0.3181	-0.0270	-0.0227	0.1478	0.6481	0.0207	0.0315	-0.0022	0.1461
IL1a	0.2248	0.1829	-1.1481	-0.0985	-0.7966	-0.0128	-0.8889	-0.1142	0.9937	1.1015	0.6754	-0.9216	0.2377	-0.0466	0.1490	0.9348	-0.1420	0.9872	0.1367	0.1233
MCP3	-0.0118	-0.0920	-0.5290	-0.2323	-0.6159	0.1024	-0.4191	0.0784	0.7628	0.6483	0.4008	-0.7783	-0.3575	-0.1492	0.3900	0.7740	0.1501	0.1652	-0.1290	0.1125
IL12p70	0.5703	0.0794	0.2996	-0.1585	-0.1648	-0.5545	-0.2743	-0.3378	-0.6801	0.2929	0.7445	-0.1870	-0.4830	0.3366	0.3190	-0.5343	-0.0240	-0.4817		0.1036
IL8	-0.1511	-0.0145	-0.0129	-0.1972	0.6285	0.0617	0.5794	0.3017	0.1426	-0.1806	-1.0008	0.1956	0.7903	-0.6442	0.6688	0.5908	0.1193	0.0702	-0.2125	0.1002
IP10	-0.1878	0.0471	-0.0404	-0.1303	0.0631	0.2009	-0.1414	0.1662	0.0457	0.1263	-0.1158	0.1388	-0.1655	0.4218	0.3975	0.2474	0.8169	0.6865	0.4257	0.0840
IL6	0.1951	0.0601	0.2211	-0.0953	-0.2674	0.0235	-0.0431	-0.1372	-0.2875	-0.0399	0.1320	0.2065	-0.2199	-0.2561	0.3124	-0.0198	-0.2808	0.2884	-0.2456	0.0777
IL1b	0.3180	-0.0189	-0.7883	-0.1694	-0.7561	-0.0158	-0.7097	-0.1642	0.8976	0.5514	0.7211	-0.6407	-0.0731	-0.0862	0.4355	0.3602	-0.0538	0.0319	-0.0863	0.0771
IL12p40	0.1574	0.0094	-0.7549	-0.2386	-0.4214	0.1237	-0.5394	-0.0696	0.8085	0.5738	0.3174	-0.5956	-0.3054	-0.2273	0.0002	0.3845	-0.0974	-0.1160	-0.0919	0.0693
TNFb	0.0286	-0.1087	-0.5919	-0.1881	-0.9711	0.0448	-0.7963	0.0793	0.5967	0.8155	0.8129	-0.7705	-0.4168	-0.2177	-0.1584	0.5117	-0.0477	0.1885	-0.3609	0.0621
IL13	0.1430	-0.0892	-0.3477	-0.1975	-0.6860	-0.0172	-0.6252	0.0084	0.4115	0.3556	0.6543	-0.4551	-0.3710	-0.2655	0.0542	0.2374	-0.1504	0.0387	-0.3375	0.0460
IL2	0.2218	-0.0281	-0.5328	-0.1709	-0.4215	0.0323	-0.6362	-0.0919	0.5919	0.3098	0.4380	-0.4106	-0.1207	-0.1569	-0.0214	0.4073	-0.1495	0.0150	-0.0307	0.0449

Coefficient Scale: -3.5 | -2.5 | -1.5 | -0.5 | 0 | 0.5 | 1.5 | 2.5 | 3.5

B

Supplemental Figure 3. Heatmap summary of baseline cytokine regression models. (A) Summary of coefficient estimates for each factor included in the baseline plasma/serum linear regression model. Cytokine models are listed in decreasing order of the adjusted R² value (shown on the far right) and coefficients are colored if FDR < 0.05. Red is associated with an increase in the cytokine level per unit change in the factor, and green is associated with a decrease in the cytokine level. (B) Basal plasma/serum cytokine level distributions.

Supplemental Figure 4. Highlights of gene expression associations with independent variables. (A-B) Example of associations based on results from Model #2 (A, determinants of baseline gene expression) and Model #3 (B, determinants of stimulated gene expression). The y-axis is gene expression and the x-axis is the independent variable (age in years, herpesvirus serostatus, ancestry informative marker principal component, baseline gene expression principal component). (C) Examples of eQTLs. *MCP1* (n = 833) and *FLT3LG* (n = 868) gene expression by genotype for rs13900 and rs2946871, respectively.

Supplemental Figure 5. Plasma and nasal wash cytokine correlates of severity.
Heatmap of plasma (A) and nasal wash (B) cytokine correlates of severity analysis, with significant adjusted p-values displayed. Colors are based on correlation coefficients or a Cohen's D effect size, depending on whether a symptom is continuous or binary, respectively. Presented p-values are from correlations (Spearman and Pearson) or Mann Whitney U tests, and results were adjusted for multiple comparisons by controlling the FDR (< 0.1).

A

Cytokine	Age	Study.Day	Sex.Male	Log.VCN	CMVPos	Age:CMVPos	HSVPos	Age:HSVPos	EBVPos	Age:EBVPos	EBVPos:CMVPos	CMVPos:HSVPos	EBVPos:HSVPos	EBVPos:CMVPos:HSVPos	Adj.R2
IL10	-0.5874	-0.0587	0.2509	0.4600	-0.6830	-0.1276	0.0425	0.2695	-0.1583	0.2630	1.5257	0.9956	0.4518	-1.7824	0.3584
IP10	-0.6166	0.0080	0.2041	0.6615	-1.2799	-0.5005	-0.1757	0.0431	-0.1985	1.2459	1.3050	2.2717	-0.1422	-1.9641	0.3489
IL15	0.0194	-0.0056	0.3004	0.2022	-0.6360	-0.3822	-0.0968	0.4781	-0.3892	-0.2145	1.8998	2.9631	0.6586	-4.1683	0.3010
TNFb	-0.0570	-0.0041	0.3022	0.1136	0.0288	0.0495	0.6813	0.8302	-0.3057	-0.7850	1.9288	0.7604	0.1641	-2.7216	0.2942
MIP1b	-2.1096	0.1970	0.1437	0.1408	-1.5901	-0.9589	2.9331	1.3542	1.6877	1.8147	1.5511	0.7308	-3.1939	-0.6401	0.2521
IL4	-0.2507	-0.0301	0.1829	0.0568	0.0586	0.0144	0.5657	0.5337	-0.0145	-0.3234	1.4769	0.8452	0.1422	-2.5686	0.2490
IL1a	-1.1253	0.0378	0.2607	0.5205	-1.8751	-0.6607	0.0804	1.0447	0.1628	0.8981	3.3974	2.7584	-0.0032	-3.5819	0.2464
MCP1	-0.1291	0.0940	-0.1171	0.3070	-0.8931	-0.5235	-0.9928	0.0853	-0.4618	0.5924	0.3722	2.2769	0.2700	-1.2507	0.2420
TNFa	-1.2841	0.1046	0.2800	0.1284	-0.9306	-0.2655	1.9287	0.8857	0.7262	0.8280	1.3744	0.3214	-1.5506	-0.7523	0.2249
IFNa2	-1.0016	0.0474	0.1560	0.3892	-0.0615	-0.1000	1.0870	1.1556	0.6803	0.3403	0.8192	3.0638	-1.2218	-3.5546	0.2214
Fractalkine	-0.7381	0.0531	0.0261	0.2013	-1.1847	-0.6522	1.0444	0.6399	0.3609	0.8132	2.0190	1.6061	-0.9654	-2.2066	0.2048
IL13	0.1338	0.0105	0.3999	0.1737	-0.0269	-0.2466	0.2157	0.3634	-0.4284	-0.6090	1.9360	1.4272	0.6039	-2.9886	0.2021
FGF2	-0.4958	0.0618	0.3218	0.1446	-0.9162	-0.2508	0.8524	0.6301	0.2189	0.3021	1.6238	1.7633	-0.6028	-2.2256	0.1947
IL12p70	-0.1672	0.0352	0.4088	0.1452	-1.0714	-0.5844	0.6820	0.1821	0.3160	0.6094	2.1132	1.7136	-0.0988	-2.7607	0.1890
EGF	-0.5943	0.2254	0.2382	0.3282	-1.7527	-0.8944	-1.1247	0.9160	-0.5112	0.8791	1.3319	4.1539	0.0541	-3.0222	0.1834
MCP3	-0.9384	0.1528	0.0983	0.2180	-0.6707	-0.1748	0.6836	1.0980	-0.1973	0.0844	1.9107	2.6431	0.0211	-3.7237	0.1740
IL2	0.0183	-0.0137	0.4153	0.1262	-0.8029	-0.7567	0.1492	0.2193	0.1643	0.3543	1.4518	1.7735	-0.0757	-2.0331	0.1732
GMCSF	-0.7331	0.0554	0.4098	0.3321	-1.5228	-1.4543	-0.9168	0.6720	0.0500	1.3538	1.3310	3.2242	-0.3184	-2.0213	0.1672
sCD40L	-0.4422	0.2783	-0.0512	0.3645	-0.5653	-0.3441	-0.9859	0.7662	-0.6361	0.5531	0.2032	2.3928	0.3830	-1.9433	0.1585
IL1Ra	-0.3795	0.1372	-0.0451	0.2043	-0.9516	0.0224	0.2344	0.8292	0.0720	-0.0254	1.9640	3.0805	-0.0448	-4.3557	0.1549
IL6	-0.7663	0.1117	0.4695	0.1763	-1.0349	-0.5904	1.7590	-0.3755	1.2898	1.4923	1.6158	0.5870	-1.9787	-0.9350	0.1542
Flt3.Ligand	0.6184	0.0532	0.5497	0.2149	-0.3946	-0.4729	-0.2288	-0.4844	-0.1184	0.1700	0.8500	1.1729	0.2747	-1.2686	0.1488
IL8	-1.7305	0.5313	-0.0053	0.2662	-0.7178	-0.2163	2.4720	0.2646	1.1112	1.8013	1.0996	2.4508	-1.7499	-3.4827	0.1461
IL12p40	-0.1598	-0.0652	0.2525	0.0584	-0.8477	-0.6326	-0.3722	0.2447	-0.3926	0.3431	1.6514	2.0133	0.7380	-2.5471	0.1460
GRO	-1.2059	0.1903	0.1118	0.3476	-1.1951	-0.8754	-0.1583	1.1881	-0.0013	1.1770	1.1350	2.4497	-0.2608	-1.8318	0.1415
TGFa	0.3623	0.1109	0.2830	0.0699	-0.7113	-0.5358	-0.0178	0.0092	-0.1350	0.0021	1.0660	1.6230	-0.1107	-1.6913	0.1273
MDC	-0.0337	0.0884	0.0143	0.1374	-0.4573	-0.3960	-0.3726	-0.3434	-0.2581	0.6558	0.2235	1.0355	-0.0414	-0.3764	0.1255
Eotaxin	-0.1278	0.1375	0.1731	0.2429	-0.3089	-0.6248	-0.2619	0.2205	-0.0332	0.7042	-0.0434	1.2704	-0.4706	-0.6325	0.1233
IL17A	0.9563	0.0394	0.4188	0.1510	-1.7088	-0.6783	-1.0232	-0.8812	0.0553	0.4966	1.6534	3.3257	0.1436	-3.0768	0.1203
GCSF	-1.1270	-0.0436	-0.2817	0.2141	-0.8567	-0.4152	1.7337	0.5869	0.3970	0.9875	1.5321	0.4635	-1.4084	-0.7805	0.1186
IL1b	-0.9188	0.1101	-0.2564	-0.0428	-0.3924	-0.2932	1.2011	-0.0083	2.1916	1.0974	-0.4535	-0.0092	-2.4446	0.6532	0.1149
MIP1a	-1.1026	0.1199	-0.1066	0.1179	-1.6884	-1.2280	1.0940	-0.1170	0.5728	1.9323	1.9453	1.6499	-1.5113	-1.3446	0.1148
VEGF	0.1944	0.0473	0.6644	0.2818	-1.2003	-0.3599	-0.5785	-0.3831	0.3459	0.5275	1.2994	2.9979	-0.2648	-2.6507	0.1040
IL3	-0.1420	0.0019	0.0837	0.0227	-0.2677	-0.3249	0.2363	0.4353	0.2337	0.0320	0.5315	0.7438	-0.3464	-0.8183	0.0981
IFNg	0.3701	-0.0429	0.5934	0.2926	-1.3498	-0.7027	-0.7589	-0.3456	0.0905	0.4439	1.8268	3.0936	0.1470	-3.1874	0.0851
IL9	0.1107	-0.0550	0.3552	-0.0201	-0.3268	0.0646	-0.3496	0.1152	-0.3608	-0.4376	0.7928	1.9024	0.6888	-2.1145	0.0685
IL7	-0.0377	0.0675	0.1323	0.0283	0.0167	0.2053	0.2239	0.1528	-0.0270	-0.2860	0.5495	1.1632	0.0828	-1.5705	0.0208

Coefficient Scale: -5 -4 -3 -2 -1 0 1 2 3 4 5

B

Supplemental Figure 6. Heatmap summary of acute plasma cytokine regression models. Samples were from influenza infected subjects. (A) Summary of coefficient estimates for each factor included in the infected plasma linear regression model. Cytokine models are listed in decreasing order of the adjusted R^2 value (shown on the far right) and coefficients are colored if FDR < 0.05. Red is associated with an increase in the cytokine level per unit change in the factor, and green is associated with a decrease in the cytokine level. (B) Acute plasma cytokine level distributions.

A

Cytokine	Age	Study.Day	Sex.Male	Log.VCN	CMVPos	Age:CMVPos	HSVPos	Age:HSVPos	EBVPos	Age:EBVPos	EBVPos:CMVPos	CMVPos:HSVPos	EBVPos:HSVPos	EBVPos:CMVPos:HSVPos	Adj.R2
RANTES	-1.4776	-0.0022	-0.1340	0.1326	-0.6570	0.0316	-0.1059	0.4904	-0.4976	1.0175	0.6927	0.4233	0.0585	-0.4770	0.332
IFNa2	-0.6786	-0.1032	-0.1711	0.3624	0.4912	-0.1265	-0.4204	-0.1818	-0.6756	0.7716	-0.4320	-0.6776	0.3987	1.0873	0.321
MIP1b	-2.0749	-0.0859	0.0685	0.5850	-0.4289	-0.0608	0.4615	0.4124	-0.0169	1.5636	0.4344	0.1856	-0.6486	-0.1606	0.314
FGF2	-0.9453	0.0054	-0.0363	0.1898	-0.4048	-0.0185	0.0024	0.2167	-0.2611	0.7276	0.5318	0.4335	-0.0917	-0.5024	0.306
PDGFABBB	-1.0112	-0.0144	-0.1526	0.1787	-0.4939	-0.0454	-0.1640	0.2566	-0.4063	0.8132	0.4856	0.5301	0.0294	-0.3881	0.301
IL10	-1.2074	-0.0908	-0.0346	0.4968	-0.1795	-0.0075	0.1769	0.1550	-0.2585	0.9289	0.1132	-0.2709	-0.1841	0.5233	0.300
IL6	-1.3389	-0.0603	-0.0887	0.5361	-0.4547	-0.0856	0.3458	0.2499	-0.3158	1.0644	0.5040	-0.2156	-0.3461	0.3720	0.295
IFNg	-0.5473	-0.0120	-0.0721	0.1977	-0.1668	-0.0704	-0.0058	0.1069	-0.2155	0.4809	0.1567	0.1250	-0.0383	0.0356	0.289
VEGF	-1.2603	-0.0204	-0.1035	0.2228	-0.4571	-0.1230	-0.1465	0.2048	-0.2344	1.0962	0.4564	0.8210	-0.2707	-0.5466	0.289
Fractalkine	-0.9536	-0.0627	-0.0777	0.3599	-0.0663	-0.2316	0.1727	-0.0968	-0.3078	1.1194	-0.0745	0.0414	-0.4442	0.6122	0.286
IL1Ra	-1.5814	-0.0808	-0.1758	0.2606	-0.1519	-0.1531	-0.0443	0.1136	-0.5321	1.4833	0.1563	0.6069	-0.1814	-0.3167	0.285
sCD40L	-1.0431	-0.0310	0.0013	0.2368	-0.7654	-0.0721	-0.3201	0.2271	-0.4050	0.8180	0.8582	0.8641	0.0351	-0.7212	0.280
TNFa	-0.9269	-0.0169	-0.0985	0.3963	-0.2459	-0.0121	0.1210	0.2813	-0.1593	0.6229	0.2086	-0.2959	-0.2360	0.3603	0.257
MIP1a	-1.5491	-0.0418	-0.0631	0.4249	-0.3489	-0.0444	0.3185	0.3686	-0.0028	1.1586	0.3076	0.1130	-0.5590	-0.0429	0.256
GMCSF	-0.7425	-0.0036	-0.0716	0.1493	-0.2507	-0.0562	-0.0154	0.1673	-0.0763	0.6065	0.1228	0.3017	-0.2294	-0.0552	0.255
MDC	-0.8485	-0.0478	-0.0280	0.0818	-0.4146	-0.0355	-0.3037	0.1879	-0.3376	0.6294	0.3510	0.5851	0.1050	-0.3136	0.237
IL12p70	-0.7346	-0.0168	-0.0352	0.1454	-0.4191	-0.0267	-0.0433	0.1728	-0.2587	0.5753	0.5139	0.3598	-0.0595	-0.3769	0.236
Eotaxin	-0.6210	-0.0609	-0.0609	0.0896	-0.3334	-0.0274	-0.1205	0.1168	-0.1387	0.5131	0.2653	0.4822	-0.0637	-0.3557	0.235
GCSF	-2.8086	-0.3114	-0.0862	0.0777	-0.9625	-0.1586	-0.1860	0.5492	-0.7696	1.8876	1.4126	0.4714	0.2993	-0.5759	0.232
IL1b	-1.5241	-0.0248	-0.1652	0.1845	-0.4864	-0.0436	0.1096	0.4152	-0.1683	1.1221	0.2826	0.1023	-0.3644	0.1624	0.221
Flt3.Ligand	-0.3865	-0.0110	-0.0404	0.1006	-0.2368	-0.0312	-0.2022	0.1058	-0.2252	0.3026	0.1796	0.2419	0.0896	-0.1123	0.214
IL1a	-1.1099	-0.0584	-0.1163	0.2400	-0.5514	-0.3175	0.0336	0.2933	-0.1003	1.0428	0.6520	0.6426	-0.5574	-0.2620	0.211
IP10	-0.1511	-0.5635	-0.0908	0.6460	0.1458	-0.0607	0.3088	-0.6153	-1.0982	-0.0377	0.0748	-1.0626	0.0199	1.8618	0.203
IL12p40	-0.6352	-0.0118	-0.0883	0.1569	-0.5532	-0.0622	-0.2406	0.1319	-0.2871	0.5500	0.4877	0.6876	0.0324	-0.4913	0.199
MCP3	-0.4562	-0.0073	-0.0187	0.1593	-0.3465	-0.0278	-0.0533	0.1007	-0.1650	0.3659	0.3396	0.2897	-0.0884	-0.1794	0.199
MCP1	-0.7382	-0.1627	-0.2032	0.4581	-0.1866	-0.1374	0.6990	-0.0148	-0.2400	0.7581	0.2491	-0.4924	-0.9743	0.8910	0.178
TGFa	-0.3740	-0.0011	-0.0473	0.0157	-0.2120	-0.0136	-0.0499	0.0983	-0.0870	0.2870	0.1674	0.1740	-0.0402	-0.0986	0.163
IL8	-2.1434	-0.2907	-0.1995	0.1478	-0.1616	-0.0700	0.7039	0.1712	-0.2434	1.4735	0.0179	-0.7922	-0.5714	1.2125	0.160
GRO	-1.2874	-0.2565	-0.1324	0.0403	-0.1200	-0.1532	0.3730	0.1332	-0.6436	0.6075	0.2947	-0.5608	-0.0695	0.7886	0.159
IL7	-0.2078	-0.0147	0.0189	0.1046	-0.3871	-0.0271	-0.2584	0.0368	-0.3235	0.1907	0.4509	0.3308	0.2146	-0.3213	0.154
EGF	0.3233	-0.1739	-0.1145	-0.1100	-1.2474	-0.0413	-0.2612	-0.6705	-0.5924	0.1431	1.0814	1.1319	0.0997	-0.7124	0.118
PDGFAA	-0.5016	-0.1477	0.0180	-0.0420	-0.1029	-0.1962	0.0005	-0.0014	-0.4792	0.3703	0.0085	0.1481	-0.2555	0.4710	0.105

Coefficient Scale: -3 -2 -1 0 1 2 3

B

Supplemental Figure 7. Heatmap summary of acute nasal wash cytokine regression models. Samples were from influenza infected subjects. (A) Summary of coefficient estimates for each factor included in the infected nasal wash linear regression model. Cytokine models are listed in decreasing order of the adjusted R² value (shown on the far right) and coefficients are colored if FDR < 0.05. Red is associated with an increase in the cytokine level per unit change in the factor, and green is associated with a decrease in the cytokine level. (B) Acute nasal wash cytokine level distributions.