# FIGNL1 AAA+ ATPase remodels RAD51 and DMC1 filaments in meiotic DNA replication and recombination

**Masaru Ito[1]\*, Asako Furukohri[1], Kenichiro Matsuzaki[1, 2], Yurika Fujita[1], Atsushi Toyoda[3], and Akira Shinohara[1]\***

[1]Institute for Protein Research, Osaka University, Suita, Osaka 565-0871, Japan
[2]Department of Advanced Bioscience, Graduate School of Agriculture, Kindai University, Nara city, Nara 631-8505, Japan
[3]Advanced Genomics Center, National Institute of Genetics, Mishima, Shizuoka 411-8540, Japan

Running title: FIGNL1 in mouse meiosis

Keywords: meiosis; homologous recombination; FIGNL1; RAD51; DMC1

\*Corresponding author:
Masaru Ito; msrito2@protein.osaka-u.ac.jp
Akira Shinohara; ashino@protein.osaka-u.ac.jp

Lead contact:
Akira Shinohara
Institute for Protein Research, Osaka University,
3-2 Yamadaoka, Suita, Osaka 565-0871 JAPAN
Phone: 81-6-6879-8624
FAX: 81-6-6879-8626
E-mail: ashino@protein.osaka-u.ac.jp
ORCID: 0000-0003-4207-8247

**Summary**

31
The formation of RAD51/DMC1 filaments on single-stranded (ss)DNAs, which is
essential for homology search and strand exchange in DNA double-strand break
(DSB) repair and for the protection of stalled DNA replication forks, is tightly
regulated in time and space. FIGNL1 AAA+++ ATPase plays positive and
negative roles in the RAD51-mediated recombination in human cells. However,
the role of FIGNL1 in gametogenesis remains unsolved. Here, we characterized
a germ-line-specific conditional knockout (cKO) mouse of *FIGNL1*. The *Fignl1*
cKO male mice showed defective chromosome synapsis and impaired meiotic
DSB repair with the accumulation of RAD51/DMC1 on chromosomes in mid-
meiotic prophase I, supporting a role of FIGNL1 in a post-assembly stage of
RAD51/DMC1 filaments. *Fignl1* cKO spermatocytes accumulate RAD51 and
DMC1 ensembles on chromosomes not only in early meiotic prophase I but also
in meiotic S-phase. These RAD51/DMC1 assemblies are independent of meiotic
DSB formation. Finally, we showed that purified FIGNL1 dismantles RAD51
filament on double-stranded (ds)DNA as well as ssDNA. These results suggest
a critical role of FIGNL1 to limit the uncontrolled assembly of RAD51 and DMC1
on native dsDNAs during the meiotic S-phase and meiotic prophase I.
49

**Introduction**

RAD51 is a central player in homologous recombination in eukaryotes by catalyzing homology-directed repair such as the repair of DNA double-strand breaks (DSBs) in both somatic and meiotic cells. RAD51 also plays a role in S-phase for the protection of stalled DNA replication forks [1, 2], which is independent of RAD51-mediated strand exchange [3, 4]. Moreover, RAD51 is important for the protection of chromosome ends such as telomeres by promoting the incorporation of Telomeric repeat-containing RNA (TERRA) into telomeres for the formation of a unique telomere DNA structure [5]. RAD51-mediated DNA transactions are critical in the maintenance of genome stability, thus suppressing tumorigenesis.

As an active form for homology search and strand exchange, multiple protomers of RAD51 bind to single-stranded (ss)DNAs in the presence of ATP to form a right-handed helical filament termed RAD51 nucleofilament, in which ssDNAs are extended 1.5-2.0-fold relative to the B-form DNA [6-9]. RAD51 nucleofilament is highly dynamic; its assembly and disassembly are controlled by multiple proteins/complexes. The RAD51 mediator promotes the assembly of RAD51 nucleofilament on the ssDNAs coated with an ssDNA-binding protein, RPA (Replication protein A), which hampers RAD51 binding to the ssDNA. The RAD51 mediator in humans includes BRCA2(-PALB2-DSS1) [10], BRCA1-BARD1 [11], RAD52 [12-14], SWI5-SFR1/MEI5 [15, 16], and five RAD51 paralogs (RAD51B, -C, -D, XRCC2, and -3) in vertebrates including humans. The RAD51 paralogs form two distinct complexes: RAD51C-XRCC3 and RAD51B-RAD51C-RAD51D-XRCC2 [17]. SWSAP1 is a distinct member of a RAD51 paralog [18-20], which forms a complex with SWS1 and SPIDR. In addition to its role in RAD51 assembly, the SWI5-SFR1/MEI5 stabilizes RAD51 filament by suppressing ATP hydrolysis by RAD51 [16, 19, 21]. At the post-assembly stage, RAD51 nucleofilament works together with the SWI2/SNF2 family, RAD54 and RAD54B [22, 23], and RAD51AP1 [24-26] for efficient homology search and strand exchange with a target double-stranded DNA (dsDNA) for the formation of displacement (D)-loop. RAD51-mediated strand invasion provides a template for DNA synthesis for further processing of the intermediates for the completion of the repair.

The dynamics of the RAD51 filament are also negatively regulated by proteins that disassemble the filament. Various DNA helicases are involved in the disassembly of RAD51 filament, which include BLM [27-29], FANCJ [30], RECQ5 [31, 32], and FBH1 [33, 34] as well as a degenerate DNA helicase, PARI [35]. RADX is an

ssDNA binding protein that modulates RAD51 filament assembly [36, 37]. RAD51-
mediated invasion intermediates are also remodeled by other DNA helicases
such as WRN, RTEL1, FANCM, and RECQ1 [38]. The stabilization and
destabilization of RAD51 filaments are associated with the choice of a
homologous recombination pathway. The disruption of RAD51 filaments and/or
the D-loop leads to the synthesis-dependent strand annealing pathway (SDSA)
for non-crossover formation [39]. On the other hand, stabilization of DNA
intermediates such as D-loop facilitates the formation of an intermediate with
Holliday structure, whose resolution leads to both crossover (CO) and non-
crossover (NCO) recombinants.

Meiosis is a specialized cell division that produces haploid gametes.
Homologous recombination, specifically crossing-over, is essential for
chromosome segregation of homologous chromosomes during meiosis I. To
ensure the crossover formation in meiosis, highly programmed regulation is
implanted on the recombination pathway [40-42]. A meiosis-specific RAD51 cousin,
DMC1, confers the uniqueness on homology search in meiotic recombination [43].
The cooperation of RAD51 and DMC1 promotes homology search between
homologous chromosomes rather than sister chromatids whose mechanism is
one big enigma in the field [44, 45]. RAD51/DMC1-mediated recombination requires
meiosis-specific proteins as well as proteins playing a role in somatic cells.
Knockout (KO) of the most of genes necessary for homologous recombination,
*Rad51* and RAD51 mediators (*Brca2*, *Rad51B*, *-C*, *-D*, *Xrcc2*, and *-3*), in mice
leads to embryonic lethality [46-52]. *Swsap1* (and *Sws1* and *Spidr*) KO mice are
viable but are sterile [18, 53]. Indeed, SWSAP1-SWS1-SPIDR is necessary for the
efficient formation of RAD51/DMC1 focus, which is a cytologically detectable
immuno-stained structure, in male meiosis [18, 53, 54]. Moreover, a meiosis-specific
HSF2BP(MEILB2)-BRME1(MEIOK21) complex promotes the assembly of
RAD51/DMC1 working together with BRCA2 mainly in male meiotic
recombination [55-58]. Like RAD51 mediators, KO mice of some negative regulators
such as BLM are also embryonically lethal [59] while *Rad54* KO mice are viable
and fertile [60].

A recently identified recombination regulator, FIGNL1, encodes an AAA+
ATPase which belongs to microtubule severing proteins including Katanin,
Spastin, and Fidgetin with an N-terminal microtubule-binding domain [61]. FIGNL1
was identified as a RAD51-interacting protein and is required for homologous
recombination in human somatic cells [62]. FIGNL1 knockdown does not affect

DSB-induced RAD51 focus formation [53, 62], suggesting its role in the post-RAD51
assembly step. In addition to its positive role in RAD51 assembly, FIGNL1 also
functions as a negative regulator for homologous recombination. FIGNL1
depletion rescues defective RAD51-focus formation in human cells depleted for
SWSAP1 or SWS1, but not for RAD51C [53]. Moreover, purified FIGNL1 promotes
the dissociation of RAD51 protein from ssDNAs, which is suppressed by
SWSAP1 [53]. In *Arabidopsis thaliana* meiosis, a mutant of *FIGL1*, FIGNL1 ortholog,
as well as its partner, *FLIP*, shows increased frequencies of COs during meiosis,
indicating a role as an anti-CO factor [63, 64]. The role of FIGL1-FLIP as the anti-CO
factor is also seen in rice meiosis [65, 66]. Importantly, the *figl1* mutation can
suppress defective RAD51/DMC1 focus formation in *brca2* mutant meiocytes,
indicating antagonism of FIGL with the RAD51 mediator [67]. These suggest that
RAD51 mediators might play a distinct function in different recombination
pathways. Although FIGNL1 plays a role in homologous recombination by
regulating the SWSAP1 complex in somatic cells [53], it remains unknown how
FIGNL1 regulates recombination and works together with SWSAP1 during
meiosis.
In this study, we analyzed the role of FIGNL1 in mice, particularly in male
meiosis. *Fignl1* KO mice show embryonic lethality, indicating that FIGNL1 plays
an essential role in early embryonic development. To explore the role of FIGNL1
in meiosis, we constructed a germ-cell specific conditional knockout (cKO) mouse
and found that *Fignl1* depletion in the testis led to defective spermatogenesis.
Meiotic chromosome spreads from *Fignl1* cKO spermatocytes showed a two-fold
increase in the numbers of RAD51 and DMC1 foci relative to controls, suggesting
the role of FIGNL1 in the post-assembly stage of not only RAD51 filament but
also DMC1 filament. This defective disassembly results in the defective loading
of proteins necessary for crossover formation. We also found that FIGNL1
suppresses RAD51 and DMC1 assembly in meiotic S-phase and early leptonema,
which are independent of meiotic DSB formation. This is supported by the
formation of RAD51 and DMC1 foci in spermatocytes from *Spo11* null mice
defective in meiotic DSB formation. Moreover, *Fignl1* cKO ameliorates defective
RAD51- and DMC1-focus formation in KO spermatocytes of *Swasp1,* a
RAD51/DMC1 mediator, supporting the antagonistic role of FIGNL1 to SWSAP1
in the assembly of not only RAD51 but also DMC1. Together, our study
establishes a dual role of FIGNL1 in mammalian male meiosis. The same
conclusion was obtained by Baudat's group by characterizing the *Fignl1* cKO as

well as *Flip* cKO (FLIP is a FIGNL1-interacting protein, accompanying paper).

**Results**

***Fignl1* is essential for early mouse embryogenesis**

To determine the functions of FIGNL1 in mice, we created a conditional allele of

*Fignl1* gene, *Fignl1$^{flox}$*, in which the exon 3 encoding an open reading frame of

the full-length protein is flanked by the *loxP* sites (Fig. 1a) in C57BL/6 background.

A null allele of the gene, *Fignl1$^{\Delta}$*, was obtained by breeding with *CAG-Cre+* mice

that constitutively express Cre recombinase. *Fignl1$^{\Delta/+}$* heterozygous mice grew

normally and did not show any apparent defects. Breeding between male and

female of *Fignl1$^{\Delta/+}$* produced *Fignl1$^{+/+}$* and *Fignl1$^{\Delta/+}$* pups, but no *Fignl1$^{\Delta/\Delta}$*

homozygous pups (0/186 from 38 litters), showing that *Fignl1$^{\Delta/\Delta}$* mice are not

viable. This is consistent with the reported lethality of *Fignl1* knockout (KO) mice

in the International Mouse Phenotyping Consortium (IMPC)

(https://www.mousephenotype.org). In a detailed analysis of different embryonic

stages by the breeding of heterozygous parents, we could not obtain *Fignl1$^{\Delta/\Delta}$*

homozygous embryo at both E7.5 and E13.5 stages (0/29 and 0/37, respectively),

showing very early lethality of mice homozygous for *Fignl1* deletion. Early

embryonic lethality in mice homozygous for the homologous recombination

genes has been reported; *Rad51* [52], *Brca2*, [68] and Rad51 paralogs (*Rad51B*, *-C*,

*-D*, *Xrcc2*, and *-3*) [46-51].

***Fignl1* conditional knockout mice are defective in spermatogenesis**

To know the role of FIGNL1 in male meiosis, we generated *Fignl1$^{flox/\Delta}$* mice

harboring a germ-line specific Cre recombinase gene, *Stra8-Cre*. The *Stra8-Cre*

gene ensured the expression of Cre not only in spermatocytes (meiotic cells) but

also in spermatogonia (mitotic stem cells) [69]. *Fignl1$^{flox/\Delta}$ Stra8-Cre$^{+}$*, hereafter

*Fignl1* conditional knockout *(*cKO), male mice were characterized for

spermatogenesis. *Fignl1* cKO adult male mice showed normal body weight

without any morphological defects but reduced sizes of the testis by ~50%

relative to controls (*Fignl1$^{+/+}$, Fignl1$^{flox/\Delta}$,* and *Fignl1$^{+/+}$ Stra8-Cre$^{+}$*), which is a

characteristic of mutants with defective spermatogenesis (Fig. 1b, c). The

western blotting analysis confirmed decreased amounts of FIGNL1 protein in the

whole testis extracts (Fig. 1d). The level of residual FIGNL1 protein is consistent

with the efficiency of Stra8-Cre-mediated excision of *Fignl1$^{flox}$* allele estimated by

the transmission of *Fignl1$^{\Delta}$* allele from *Fignl1$^{flox/+}$ Stra8-Cre$^{+}$* male mice (91.3%;

116/127). Sperm count showed a ~6-fold reduction in its number; residual sperms
might come from $Fignl1^{flox/\Delta}$ cells where Stra8-Cre recombinase failed excision of
the $Fignl1^{\Delta}$ allele (Fig. 1e). Histological analysis showed that seminiferous
tubules in *Fignl1* cKO testes contained decreased numbers of late spermatocytes
and spermatids (Fig. 1f). These indicate a critical role of FIGNL1 in
spermatogenesis.

To analyze meiotic defects in *Fignl1* cKO male mice, we prepared surface-
spread chromosomes of spermatocytes and immuno-stained for meiosis-specific
chromosomal proteins, SYCP3(Scp3) and SYCP1(Scp1), which are components
of lateral/axial elements and central region of the synaptonemal complex (SC),
respectively [70-72]. In leptonema of control spermatocytes, SYCP3 shows thin-line
staining structures that indicate elongating chromosome axes (from short to long
lines). In zygonema, regions of two thin chromosome axes are paired locally to
form a thick line of SYCP3, where SYCP1 proteins are deposited for the assembly
of the SC central region. From early zygonema to early pachynema, SC
elongates along entire chromosomes, resulting in pachytene chromosome
configuration with 19 fully paired thick SYCP3 lines colocalized with SYCP1 and
partially paired X-Y bivalent. In *Fignl1* cKO spermatocytes, like in controls,
SYCP3-positive lines were formed in leptonema and short lines of SYCP1 in
zygonema were also observed. However, there were fewer pachytene and
diplotene cells in *Fignl1* cKO relative to controls (22.6±2.4% and 11.4±6.2%
compared to 44.7±11.0% and 24.3±7.7%, respectively; Fig. 1g). We also
evaluated the expression of FIGNL1 protein in testicular cell squashes by
categorizing SYCP3-positive meiotic cells into two: cells before pachynema (i.e.
leptonema and zygonema) with a pan-nuclear $\gamma$H2AX staining and those after
pachynema (i.e. pachynema and diplonema) with a large $\gamma$H2AX staining on the
X-Y pair (the sex body) [73] (Extended Data Fig. 1). In controls, cells both before
and after pachynema were 100% positive for FIGNL1 staining. In contrast, while
only ~25% of cells prior to pachynema were positive for FIGNL1, cells after
pachynema were nearly 100% positive in *Fignl1* cKO (Fig. 1h). This indicates that
$Fignl1^{\Delta/\Delta}$ cells are depleted at either late zygonema or early pachynema from
*Fignl1* cKO testes. Further analysis of surface-spread chromosomes revealed
abnormal SC formation such as synapsis between non-homologous
chromosomes (67.7±2.0% of cells compared to 13.6±4.1% in controls) and
multivalent synapsis (26.8±10.9 % of cells compared to 3.1±2.7% in controls) in
*Fignl1* cKO (Fig. 1i, j), which could be defined as "pachytene-like" stage as seen

in other meiotic recombination defective mutants such as *Dmc1* and *Meiob* KO,
etc. [74-76]. These results show that FIGNL1 is necessary for full synapsis of meiotic
chromosomes in late zygonema and/or early pachynema and that most of the
FIGNL1-depleted cells seem to die at these stages or do not progress further.
234

**FIGNL1 promotes the disassembly of DMC1 in male meiosis**

To know the effect of *Fignl1* cKO on meiotic recombination, we examined the
localization of DMC1 on surface-spread chromosomes of spermatocytes,
together with SYCP3 as a marker for the progression of meiotic prophase I. As
shown previously [77-79], in control spermatocytes, DMC1 foci appear in leptonema,
peak in their number at early zygonema, and gradually decrease throughout
zygonema (Fig. 2a, b). In *Fignl1* cKO, DMC1 foci followed similar kinetics as seen
in controls. However, at any stage, the number of DMC1 foci in *Fignl1* cKO was
more than that in controls. In early zygonema, ~500 DMC1 foci (449±89 foci) are
observed in *Fignl1* cKO, which is 2.3-fold more than in controls (194±32 foci).
DMC1-focus number in *Fignl1* cKO clearly increased through meiotic prophase I
progression. In late zygonema, the number of DMC1 foci in *Fignl1* cKO was
reduced by half of that in early zygonema (220±70 foci compared to 449±89 foci),
indicating a turnover of the foci. These results suggest that FIGNL1 suppresses
excess DMC1 focus formation and/or timely progression of DMC1-mediated
recombination.
In *Fignl1* cKO spermatocytes, we always observed two populations: cells
with a high number of DMC1 foci and cells with a similar number of foci to controls.
We assumed that the latter is *Fignl1^{flox/Δ}* cells where Stra8-Cre recombinase failed
excision of the *Fignl1^{Δ}* allele. Note that we included these foci for our
quantification and that this population becomes an internal control for our analysis.
256

**Compromised meiotic recombination in *Fignl1* cKO**

ssDNA-binding proteins such as RPA and MEIOB-SPATA22 play a role in the
early/middle stage of meiotic recombination both before and after the DMC1
assembly. We examined the localization of RPA2, a subunit of RPA, on meiotic
chromosomes. In control spermatocytes, RPA2 foci appear from leptonema with
200-300 foci on average in mid/late leptonema (223±73 foci; Fig. 3a, b). The
number of RPA2 foci increased up to ~400 foci in zygonema (376±45 foci in mid-
zygonema) and persisted throughout zygonema (332±59 foci in late zygonema).
The RPA2-focus number in *Fignl1* cKO mid/late leptonema increased up to 300-

400 foci (368±100 foci) and a similar number persisted in the early zygotene
stage (409±59 foci). Different from control mid/late leptonema where the RPA2-
focus number is ~2-fold more than the DMC1 focus number (223±73 RPA2 foci
and 96±57 DMC1 foci), the number of RPA2 foci in *Fignl1* cKO is comparable to
that of DMC1 foci (368±100 RPA2 foci and 369 ±156 DMC1 foci). The number of
RPA2 foci in *Fignl1* cKO gradually decreased from mid-zygonema and was
significantly lower than controls in late zygonema (281±48 foci compared to
332±59 in controls; $p<0.0001$, Mann-Whitney *U*-test; Fig. 3b), suggesting a role
of FIGNL1 in the dynamic behavior of an RPA-associated recombination event.
We also analyzed the histone $\gamma$H2AX staining as a marker of DSB formation and
its repair. Interestingly, *Fignl1* cKO spermatocytes with increased RPA2 foci
showed a similar or slightly reduced intensity of $\gamma$H2AX from leptonema to the late
zygonema stage compared to controls (Extended Data Fig. 2a, b).
In meiotic recombination, cytological analyses showed that RPA2 foci
persist longer than DMC1/RAD51 foci from mid/late zygonema to early/mid
pachynema, suggesting a role of RPA in a late stage of meiotic recombination [80].
In zygonema (from early zygonema), foci containing the MutS$\gamma$ complex with
MSH4 and MSH5 appear specifically on synapsed regions and show similar
behavior to RPA on meiotic chromosomes to regulate late recombination events
[81]. We studied MSH4 localization and observed the reduced MSH4 foci in *Fignl1*
cKO in mid/late zygonema (50±22 foci versus 107±47 foci in controls; Fig. 3c, d).
Given a possibility that the reduction in MSH4-focus number is an indirect
consequence of synapsis defects in *Fignl1* cKO, we also analyzed the density of
MSH4 foci per unit length of SC. MSH4-focus number and SC length (total
SYCP1 length) showed a linear relationship in control mid-zygonema ($r=0.89$; Fig.
3e). In *Fignl1* cKO, a correlation between MSH4-focus number and SC length
was also positive, but lower than in controls ($r=0.58$). Importantly, the density of
MSH4 foci per unit length of SC is reduced in *Fignl1* cKO relative to controls
(0.73±0.22 foci per $\mu$m of SC versus 0.98±0.18 foci per $\mu$m of SC in controls; Fig.
3f). These suggest that FIGNL1 is required for efficient loading of MSH4 for
meiotic recombination and downstream events such as SC elongation.
Crossover formation is a key outcome of meiotic recombination. The
crossover sites on meiotic chromosomes are marked with MutL$\gamma$ complex such
as MLH1 foci [82]. Control late pachytene cells show ~25 MLH1 foci per nucleus.
On the other hand, we could not detect any MLH1 foci on late
zygotene/pachytene-like chromosomes of *Fignl1* cKO spermatocytes (Extended

Data Fig. 2c). This is consistent with the observation that FIGNL1 is essential for
the progression of spermatocytes into a pachytene stage where MLH1 foci form.
304

**FIGNL1 promotes the disassembly of RAD51 in male meiosis**
In human cultured cells, FIGNL1 regulates RAD51 assembly [53, 62]. We examined
the localization of RAD51 on meiotic chromosomes. As reported previously [77-79],
in control spermatocytes, RAD51 foci are observed from mid/late leptonema and
peak in early zygonema, and then decrease toward pachynema (Fig. 2c, d). The
average number of RAD51 foci is 107±56 and 181±27 foci per nucleus in control
mid/late leptotene and early zygotene stages, respectively. As seen for DMC1,
we detected an increased number of RAD51 foci in *Fignl1* cKO cells. The average
number of RAD51 foci in early zygonema is 380±98 foci, which is more than twice
compared to that of controls (181±27 foci). The number of RAD51 foci in
zygonema is similar to that of DMC1 foci, though ~700 RAD51 foci were observed
in some nuclei. When the zygotene stage progressed from early to late, the
RAD51-focus number in *Fignl1* cKO gradually decreased as in controls. However,
even at late zygonema, we detected a significant increase in RAD51-focus
number in *Fignl1* cKO relative to controls (182±86 foci compared to 128±32 foci
in controls; $p$=0.0163, Mann-Whitney $U$-test; Fig. 2d). These results show that
FIGNL1 promotes the timely resolution of RAD51 ensembles during the zygotene
stage.
323

**Increased RAD51/DMC1 foci in early leptonema in *Fignl1* cKO**
In contrast to the zygotene stage where the numbers of DMC1 and RAD51 foci
are comparable, focus numbers of DMC1 and RAD51 were clearly different at
early leptonema only in *Fignl1* cKO (Fig. 2a-d). In mid/late leptonema when
DMC1 and RAD51 foci appear on elongating SYCP3 stretches in controls, the
number of DMC1 foci in *Fignl1* cKO is 369±156 foci per cell, which is about 3-4
times more of that in controls (96±57 foci). The number of RAD51 foci in mid/late
leptonema in *Fignl1* cKO is 387±134 foci, 3-4 times more than in controls (107±56
foci). In early leptonema, where SYCP3 short stretches are yet elongated, DMC1
and RAD51 foci rarely appear in control spermatocytes. However, *Fignl1* cKO
showed 355±178 RAD51 foci, which exceeded the number of DMC1 foci
(194±148 foci). These suggest that FIGNL1 suppresses the formation of excess
numbers of RAD51 and DMC1 foci in early meiosis I, and that, in the absence of
FIGNL1, RAD51 and DMC1 loading is uncoupled.

**Meiotic S-phase in *Fignl1* cKO accumulates RAD51/DMC1 foci**

In wild-type meiosis, the formation of both RAD51 and DMC1 foci is mediated by

SPO11-induced DSBs in leptonema [83, 84]. Increased number of RAD51 and

DMC1 foci on early leptotene chromosome of *Fignl1* cKO spermatocytes led us

to analyze preleptotene (meiotic S-phase) spermatocytes in order to characterize

the origin of excess RAD51/DMC1 foci in the absence of FIGNL1. For *in vitro*

labeling of DNA replication, we incubated testis cell suspension with EdU for 1

hour in a culture medium. The preleptotene spermatocytes, where a limited

number of SYCP3 short stretches forms along with a numerous number of

SYCP3 dots, were categorized into two classes: pan dotty staining of EdU in early

preleptonema and clustered dotty staining of EdU in mid/late preleptonema. In

controls, both early and mid/late preleptotene EdU-positive spermatocytes

showed very few foci of RAD51 (Fig. 4a, b). On the other hand, we could detect

RAD51 foci in EdU-positive spermatocytes of *Fignl1* cKO, indicating RAD51-

focus formation during the meiotic S phase. The number of RAD51 foci in the

meiotic S-phase is around 300 in focus-positive nuclei (298±110 foci and

337±108 foci in early and mid/late preleptonema, respectively), which is ~3/4 of

the RAD51 focus number in meiotic prophase I of *Fignl1* cKO mice, but ~1.5-fold

more of that in early zygonema of control mice (compare Figs. 2d and 4b). Some

RAD51 foci in the meiotic S-phase are colocalized with EdU dots/lines,

suggesting that RAD51 foci are associated with DNA replication. However, when

DNA replication labeling with another thymidine analog IdU was shortened for 10

min, most of the RAD51 foci did not colocalize with IdU foci (Extended Data Fig.

3a-c), though the degree of colocalization was significantly higher than random

colocalization (3.7±1.9% of IdU foci colocalized with RAD51 compared to

2.5±1.3%, $p$=0.0261, Mann-Whitney $U$-test; 9.1±4.5% of RAD51 foci colocalized

with IdU compared to 6.2±3.3%, $p$=0.0439, Mann-Whitney $U$-test; Extended Data

Fig. 3c). It is unlikely that RAD51 is frequently associated with ongoing DNA

replication in meiotic S-phase.

Further evaluation of DMC1 localization in *Fignl1* cKO revealed the

presence of preleptotene spermatocytes containing DMC1 foci (Fig. 4c, d). The

number of DMC1 foci in focus-positive nuclei is around 150 (136±71 foci and

164±70 foci in early and mid/late preleptonema, respectively), about half of

RAD51 foci in *Fignl1* cKO. Of note is that very few DMC1 foci were detected in

control preleptotene spermatocytes. This suggests that FIGNL1 limits the

assembly of mitotic recombination machinery with RAD51 on chromatin during
the meiotic S-phase as well as meiotic recombination machinery containing
DMC1.

**Uncoupled loading of RAD51 and DMC1 in *Fignl1* cKO spermatocytes**
Despite the overall increase in RAD51 and DMC1 focus numbers on any
analyzed stages of *Fignl1* cKO spermatocytes, ~2-fold more RAD51 foci
compared to DMC1 foci in preleptotene and early leptotene stages suggest
uncoupling of RAD51 and DMC1 loading onto chromosomes in the absence of
FIGNL1. Co-staining of RAD51 and DMC1 confirmed different kinetics of RAD51
and DMC1 in *Fignl1* cKO (Fig 2e, f, Extended Data Fig. 4); DMC1 focus numbers
gradually increase from preleptonema to late leptonema/early zygonema
(177±74, 325±136, and 407±69 foci in preleptonema, early leptonema, and late
leptonema/early zygonema, respectively; Fig, 2f), whereas RAD51 focus
numbers are nearly constant in those stages (399±120, 476±109, and 480±95
foci in preleptonema, early leptonema, and late leptonema/early zygonema,
respectively). Colocalization analysis revealed a gradual increase in the number
of RAD51-DMC1 co-foci as DMC1 foci increased. In preleptotene and early
leptotene spermatocytes, 50-60% of DMC1 foci colocalized with RAD51 foci
(57.0±10.6% and 53.1±11.0% in preleptonema and early leptonema,
respectively; Fig, 2g) and less than 40% of RAD51 foci colocalized with DMC1
foci (25.7±9.9% and 35.3±12.0% in preleptonema and early leptonema,
respectively). The degree of colocalization at these stages is significantly lower
than that in the late leptotene/early zygotene stage where comparable levels of
colocalization with controls were observed (65.8±10.1% and 67.2±7.6% of DMC1
foci colocalized with RAD51 in *Fignl1* cKO and controls, respectively; 56.2±9.6%
and 59.3±8.9% of RAD51 foci colocalized with DMC1 foci in *Fignl1* cKO and
controls, respectively; Fig. 2g). These suggest that coupled loading of RAD51
and DMC1 in meiotic S-phase and early meiotic prophase I is compromised in
*Fignl1* cKO spermatocytes. On the other hand, the fact that more than half of
DMC1 foci are colocalized with RAD51 in preleptonema suggests that DMC1
preferentially binds to meiotic chromatin where RAD51 is already bound.

**RAD51/DMC1-focus formation in *Spo11* KO *Fignl1* cKO spermatocytes**
Both RAD51 and DMC1 foci appear in the meiotic S-phase in *Fignl1* cKO
spermatocytes, which is much earlier than meiotic prophase I with programmed

DSB formation by SPO11. To examine whether RAD51/DMC1 focus formation in
*Fignl1* cKO mice is independent of meiotic DSB formation or not, we constructed
*Spo11* KO *Fignl1* cKO mice (*Spo11*$^{-/-}$ *Fignl1*$^{flox/\Delta}$ *Stra8-Cre*$^{+}$), which is defective in
SPO11-mediated DSB formation [83, 84]. As shown previously, spermatocytes from
control *Spo11* KO mice (*Spo11*$^{-/-}$ *Fignl1*$^{flox/\Delta}$) did not show any RAD51 foci on
meiotic chromosomes (Fig. 5a, b, Extended Data Fig. 5a). On the other hand,
spermatocytes from *Spo11* KO *Fignl1* cKO show on average 300-400 RAD51 foci
from leptotene to zygotene(-like) stages (407±77, 390±64, and 292±34 foci for
focus-positive nuclei in leptonema, early zygonema, and late zygonema,
respectively; Fig. 5b). Moreover, *Spo11* KO *Fignl1* cKO also accumulates DMC1
foci with similar numbers to RAD51 foci (374±108, 391±57, and 342±54 foci for
focus-positive nuclei in leptonema, early zygonema, and mid-zygonema,
respectively; Fig. 5c, d, Extended Data Fig. 5b). Comparable numbers of RAD51
and DMC1 foci between *Fignl1* cKO and *Spo11* KO *Fignl1* cKO indicate that a
large fraction of RAD51 and DMC1 foci on meiotic chromatin in the absence of
FIGNL1 is independent of SPO11, thus meiotic DSBs. Like *Spo11* KO, *Spo11*
KO *Fignl1* cKO spermatocytes lack pun-nuclear $\gamma$H2AX staining (Extended Data
Fig. 5c, d), supporting the assembly of RAD51/DMC1 complexes in the absence
of endogenous DSBs. *Spo11* KO *Fignl1* cKO spermatocytes died in the
pachytene-like stage with the formation of the pseud-sex body (Extended Data
Fig. 5d) as seen in *Spo11* KO spermatocytes [83, 84]. We also detected
RAD51/DMC1 foci in preleptotene spermatocytes of *Spo11* KO *Fignl1* cKO mice
(Fig. 5f). This confirms that RAD51/DMC1 focus formation during meiotic S-phase
is independent of the SPO11 function. Importantly, the degree of RAD51-DMC1
colocalization from preleptonema to early zygonema in *Spo11* KO *Fignl1* cKO is
comparable to that in *Fignl1* cKO and more than half of DMC1 foci are colocalized
with RAD51 at those stages in *Spo11* KO *Fignl1* cKO (Fig. 5e-g). This suggests
that DMC1 preferentially binds to RAD51-bound meiotic chromatin, which is most
likely dsDNA.
439

**RAD51 binds to and persists on meiotic recombination sites in *Fignl1* cKO**
**spermatocytes**
As described above, the absence of FIGNL1 increases the SPO11-independent
formation of RAD51 and DMC1 foci in meiotic prophase I, some of which would
come from the meiotic S-phase. Comparable numbers of RAD51 and DMC1 foci
in the presence and absence of SPO11 (i.e., meiotic DSBs) sought us to check

whether RAD51/DMC1 can bind to meiotic recombination hotspots, especially

ssDNA. Chromatin immunoprecipitation followed by single-stranded DNA

sequencing (ChIP-SSDS) for RAD51 [85-87] revealed that in *Fignl1* cKO testes, like

in controls, RAD51 binds to recombination hotspots where DMC1 strongly binds

in wild-type testes (Fig. 6a, Extended Data Fig. 6a). Importantly, when RAD51

binding around hotspots is ranked based on the number of SPO11-oligos in wild-

type, a similar binding profile of RAD51 is seen between *Fignl1* cKO and controls,

with a small increase in RAD51 ChIP-SSDS signals in two out of three *Fignl1*

cKO mice (Fig. 6b, Extended Data Fig. 6b-d).

To address whether RAD51 persists on meiotic recombination sites in the

absence of FIGNL1, we also measured the colocalization frequency of RAD51

and RPA2 in mid/late zygonema, given that RPA2 localizes to recombination sites

even after RAD51/DMC1 disassembly in wild-type meiosis and can be a marker

of recombination sites on mid/late zygotene chromosomes. In control mid/late

zygotene spermatocytes, out of 234±44 RPA2 foci, 41±15 foci were colocalized

with RAD51 foci, which corresponds to 17.2±4.3% of RPA2 foci (Fig. 6c-e). *Fignl1*

cKO exhibited two classes of populations; one is similar to controls (~30%) and

the other shows 40-80% of colocalization, with 45.4±21.9% of RPA2 foci

colocalized with RAD51 for all populations. This is consistent with the idea that

RAD51 binds to ssDNA on meiotic recombination sites in the absence of FIGNL1

and suggests that RAD51 persists on RPA2-associated recombination sites.

***Fignl1* deficiency rescues RAD51/DMC1 assembly defects in *Swsap1***

Our previous study in human cells showed that FIGNL1 antagonizes the function

of the RAD51 mediator, SWSAP1 [53]. In mouse male meiosis, SWSAP1 is

necessary for efficient RAD51/DMC1 focus formation [18, 53]. We tested the effect

of *Fignl1* cKO on RAD51/DMC1 assembly in *Swsap1* KO cells by generating

*Swsap1* KO *Fignl1* cKO male mice (*Swasp1⁻/⁻ Fignl1flox/Δ Stra8-Cre⁺*). As reported

[18, 53], surface-spread chromosomes from *Swasp1* KO spermatocytes (*Swsap1⁻/⁻*

*Fignl1flox/Δ*) showed a large reduction of RAD51 foci with 10-30 foci (~1/10 of

control *Fignl1flox/Δ*) in leptonema and zygonema (29±26, 22±15, and 20±14 foci in

leptonema, early zygonema, and mid zygonema, respectively; Fig. 7a, b). Like

*Fignl1* cKO, *Swsap1* KO *Fignl1* cKO spermatocytes showed an accumulation of

RAD51 foci on meiotic chromosomes. The number of RAD51 foci from leptonema

to mid-zygonema in *Swsap1* KO *Fignl1* cKO is very similar to that in *Fignl1* cKO

(320±217, 308±221, and 194±179 foci in *Swsap1* KO *Fignl1* cKO and 399±145,

377±98, and 282±91 foci in *Fignl1* cKO in leptonema, early zygonema, and mid

zygonema, respectively). This indicates that in mouse spermatogenesis without

FIGNL1, RAD51 assembly does not depend on SWSAP1. This is consistent with

the results in human cultured cells. On the other hand, in late zygonema, the

number of RAD51 foci in *Swsap1* KO *Fignl1* cKO was comparable to *Swsap1* KO

and much lower than that in *Fignl1* cKO (10±12, 7±8, and 189±79 foci in *Swsap1*

KO *Fignl1* cKO, *Swsap1* KO, and *Fignl1* cKO, respectively; Fig. 7b). This

suggests that only *Swsap1$^{-/-}$ Fignl1$^{flox/\Delta}$* cells where Stra8-Cre recombinase failed

excision of *Fignl1$^{flox}$* allele can progress into late zygotene stage in *Swsap1* KO

*Fignl1* cKO testes. Moreover, similar results were observed for DMC1; the

number of DMC1 foci in *Swsap1* KO *Fignl1* cKO was comparable to *Fignl1* cKO

(Fig. 7c, d). Thus, reduced DMC1-focus formation in *Swsap1* KO is also

suppressed by *Fignl1* cKO. It is likely that the activity of FIGNL1 to suppress

RAD51/DMC1 assembly is separable from its activity regulated by SWSAP1.

**FIGNL1 disrupts RAD51-dsDNA complex *in vitro***

We previously showed that FIGNL1 promotes the dissociation of RAD51 from

ssDNAs *in vitro* [53]. The above results in mouse spermatocytes suggest that

FIGNL1 can disrupt RAD51 filaments on chromosomal DNAs, both ssDNAs and

dsDNAs. To address whether FIGNL1 promotes the dissociation of RAD51

filaments on dsDNAs *in vitro*, we performed the electrophoretic mobility shift

assay (EMSA). First, we confirmed that a purified human FIGNL1ΔN, which lacks

N-terminal 284 amino acids [53], disrupted the pre-formed RAD51-ssDNA

complexes in the presence of ATP (Fig. 8a, b). Under this condition, FIGNL1ΔN

could disrupt ~30% of RAD51-ssDNA complex (24.6±6.9% of unbound DNAs

were increased to 42.9±2.1% by the addition of 1.2 μM of FIGNL1ΔN). When

FIGNL1ΔN was incubated with a pre-formed RAD51-dsDNA complex, the most

of dsDNA was recovered as an unbound form (25.8±11.1% of unbound DNAs

were increased to 80.0±11.6%). This suggests that FIGNL1ΔN could disrupt the

RAD51-dsDNA complex more efficiently than the RAD51-ssDNA complex *in vitro*,

supporting the *in vivo* results.

**Discussion**

In this study, we analyzed the role of FIGNL1 AAA+ ATPase, which plays a critical

role in homologous recombination in somatic cells, in mouse male meiosis. *Fignl1*

cKO mice accumulated RAD51/DMC1 on meiotic chromosomes with defective

chromosome synapsis. Our analysis revealed the dual function of FIGNL1 during spermatogenesis (Fig. 8c). One is to promote meiotic recombination by regulating the dissociation of RAD51/DMC1 ensembles on the ssDNA. The other is to protect the genome by disrupting the non-functional form of RAD51/DMC1 ensembles on the dsDNA. Consistent with the idea, purified FIGNL1 can disrupt the RAD51-dsDNA complex as well as the RAD51-ssDNA complex *in vitro*. The same conclusion was obtained by Baudat's group by characterizing the *Fignl1* cKO as well as *Flip* cKO (accompanying paper).

**FIGNL1 promotes the post-assembly stage of RAD51/DMC1 in meiotic recombination**

In meiotic recombination, RAD51/DMC1 filaments on ssDNAs play a critical role in homology search and strand exchange with homologous dsDNA, leading to the formation of the D-loop, in which the 3'-OH end of invading strand becomes a substrate for recombination-associated DNA synthesis. Biochemical analyses showed that RAD51/DMC1-mediated D-loop is a poor substrate for *in vitro* DNA synthesis by DNA polymerase $\delta$ and its accessory factors [88-90]. To promote efficient DNA synthesis on the recombination intermediate, RAD51/DMC1 bound to the intermediate should be removed by factors such as RAD54, a RAD51 remodeler [88-90]. Although the male KO mice for Rad54 and for both Rad54 and Rad54B, a Rad54 paralog, are fertile, the cKO spermatocytes show abnormal loading of RAD51 on chromosomes in pachynema and diplonema [60]. This is consistent with the role of RAD54 in the post-assembly stage of RAD51 (DMC1 had not been analyzed in detail). Our *in vivo* and *in vitro* studies suggest the role of FIGNL1 in the post-assembly stage of the RAD51/DMC1 filaments (this study). It is likely that FIGNL1 disrupts not only RAD51 but also DMC1 filaments on recombination intermediates for subsequent DNA synthesis [89, 91] (Fig. 8c). Post-assembly role of FIGNL1 in RAD51-mediated homologous recombination was reported previously in human cells with DNA damage [62]. Consistent with the post-assembly role of FINGL1, *Fignl1* cKO spermatocytes reduce the formation of MSH4 foci, which appears after the loading and dissociation of RAD51/DMC1 [81]. *Fignl1* cKO spermatocytes seem not to enter the pachytene stage for the loading of the MutLγ complex such as MLH1 and most *Fignl1* KO cells die around late zygonema. The meiotic defects seen in *Fignl1* cKO are similar to those in the mutants where RAD51/DMC1 foci are accumulated such as *Msh4*, *Msh5*, *Mcm8*, *Trip13,* and *Mcmdc2* [92-98]. In mouse spermatocytes, FIGNL1 seems to promote

ZMM-dependent CO formation. This is different from the role of FIGNL1 ortholog,
FIGL1, in *A. thaliana*. *Arabidopsis FIGL1* mutant meiocytes showed normal or
slightly elevated CO formation [63, 64]. Indeed, the depletion of FIGL1 or its partner
FLIP suppresses reduced CO formation in the absence of pro-CO proteins, ZMM
[63, 64]. This suggests the role of plant FIGL1-FLIP as an anti-CO factor, which
disrupts early recombination intermediates such as RAD51/DMC1 filaments on
the ssDNA prior to or during the strand invasion.
561
**FIGNL1 disrupts non-productive RAD51 and DMC1 ensembles on duplex**
**DNAs**
RAD51(DMC1)-focus formation depends on exogenous genotoxic treatment or
SPO11-catalyzed DSB formation associated with the resection, thus the
formation of ssDNAs. The most striking phenotypes seen in *Fignl1* cKO
spermatocytes are the accumulation of RAD51 and DMC1 foci in unchallenged
meiotic S-phase cells and spermatocytes in the absence of SPO11. This
accumulation of RAD51/DMC1 foci in the absence of DSBs, therefore the
ssDNAs, could be simply explained by the non-specific assembly of
RAD51/DMC1 filaments on chromatin DNAs, thus the binding to dsDNA (Fig. 8c).
Interestingly, human RAD51 and DMC1 bind to both ssDNAs and dsDNA with
similar affinity *in vitro* [17, 99] and RAD51 bears an activity to remodel the
nucleosome upon the dsDNA binding [100-102]. On the other hand, *in vivo* DNA
binding of RAD51 and DMC1 is restricted to ssDNAs. How cells discriminate
ssDNAs among lots of competitors, dsDNAs, in the RAD51/DMC1 binding
remains unsolved. Based on the results described here, we propose that FIGNL1
actively dismantles RAD51 and DMC1 filaments on dsDNAs, which is otherwise
non-productive for the recombination and/or might become obstacles to
chromosomal events on duplex DNAs such as DNA replication, transcription, and
others. Indeed, Bishop and his colleagues reported activities of the RAD54
translocase family to remove RAD51 and DMC1 from dsDNAs. In yeast meiosis,
Spo11-independent focus formation of Dmc1, but not of Rad51, depends on
Tid1/Rdh54, a Rad54 homolog [103]. And mitotic spontaneous Rad51 focus
formation on yeast chromosomes is increased in the absence of three
translocases, Rad54, Tid1/Rdh54, and Uls1 [104]. Moreover, in unchallenged
human cells, the absence of two RAD54 family proteins RAD54 (or RAD54L) and
RAD54B leads to the accumulation of spontaneous RAD51 foci particularly when
RAD51 is over-expressed [105]. Further studies are necessary to reveal the

relationship between FIGNL1 and RAD54 translocases in the remodeling of
RAD51/DMC1 assembles on duplex DNAs.
One of the striking features of RAD51/DMC1 assembles on duplex DNAs is
that these ensembles are quite stable. The number of RAD51 in both meiotic S-
phase and early meiotic prophase I from early leptonema to early zygonema is
constant. Little increase in RAD51-focus number during early meiotic prophase I
suggest that spermatocytes show very limited additional assembly of RAD51 foci
during this stage. On the other hand, DMC1 shows a 2-fold increase in focus
number from early leptonema (and meiotic S-phase) to early zygonema. This
increase could be induced by SPO11-mediated DSBs. Interestingly, the
colocalization frequency of RAD51 with DMC1 in *Fignl1* cKO increased from
~30% to ~60%, which is similar to the colocalization frequency seen in control
early zygonema, while the colocalization frequency of DMC1 with RAD51 is
nearly constant (with 60-70% frequency). This suggests that RAD51 ensembles
even on the dsDNA tend to recruit DMC1 to form co-ensembles.
605
**FIGNL1 antagonizes SWSAP1 for RAD51/DMC1 assembly on meiotic**
**chromosomes**
FIGNL1 shows antagonism to the RAD51 mediator. In humans, *FIGNL1*
depletion suppresses RAD51-assembly defect induced by the *SWSAP1*
depletion [53]. In meiosis of the *A. thaliana*, *FIGL1* (*A. thaliana FIGNL1*) and/or *FLIP*
deletion restores reduced RAD51/DMC1 focus formation in the mutants of the
*BRCA2* [67] as well as defective DMC1-focus formation in the mutant of the *SDS1*,
which promotes DMC1 assembly [63]. Given that *Swsap1* KO spermatocytes
reduced the RAD51 and DMC1 foci [18, 53], we asked whether *Fignl1* cKO could
also suppress these meiotic defects in *Swsap1* KO spermatocytes and found that
the number of both RAD51/DMC1 foci in *Swsap1* KO *Fignl1* cKO is similar to
those in *Fignl1* cKO. This is simply interpreted as, like in human cells, *Fignl1* cKO
suppresses RAD51/DMC1 assembly defect in *Swsap1* KO. However, at this point,
we do not know whether RAD51/DMC1 foci are associated with ssDNAs in
meiotic prophase I cells of *Swsap1* KO *Fignl1* cKO as shown in *Fignl1* cKO,
although it is likely. In addition, different from RAD51/DMC1 assembly on ssDNAs,
these suggest that RAD51/DMC1 assembly on dsDNAs in the absence of
FIGNL1 does not require the RAD51/DMC1 mediator, SWSAP1.
624

## Methods

### Mice

The care and use of mice in this study were performed under the guideline for the proper conduct of animal experiments (Society Council of Japan). These procedures were approved by the Institutional Animal Care Committee at Institute for Protein Research, Osaka University (approval ID; 25-03-0).

All mice used in this study were congenic to the C57BL/6 background. Genotyping of mice was performed by PCR on genomic DNA isolated from mice. Primers used for genotyping are listed in Supplementary table 1. Mature adult mice (8-20 weeks old) were used for experiments otherwise mentioned.

*Fignl1$^{flox}$* allele was constructed by Cyagen Bioscience Co. Ltd. ES cells with the *Fignl1$^{flox}$* allele were injected into C57BL/6 albino blastocytes, which were transplanted into CD-1 pseudo-pregnant female mice. The founder mice with the correct *Fignl1$^{flox}$* allele were backcrossed to C57BL/6 mice. *Fignl1$^{+/\Delta}$* allele was constructed by breeding *Fignl1$^{flox/+}$* with *CAG-Cre* mice. *Stra8-Cre* transgenic mice were previously described [106] and provided *by* the RIKEN Animal Resource Center. *Spo11$^{-/-}$* mouse was previously described [83] and *Spo11$^{+/-}$* mice with a mixed background of C57BL/6 and 129/SvJ were backcrossed to C57BL/6 mice prior to experiments. *Swsap1$^{-/-}$* mice were described in our previous study [53].

### Histology

Testes from adult mice were fixed in 10% buffered formalin overnight at room temperature, washed twice with PBS, and stored in 70% EtOH at 4°C prior to embedding in paraffin and sectioning. The sections of 5 $\mu$m thickness were deparaffinized and stained with hematoxylin and eosin (H&E).

### Spermatocyte chromosome spreads

Surface-spread chromosomes of spermatocytes were prepared as described previously [107, 108] with slight modification. Testes were dissected, the tunica albuginea was removed, and adherent extra-tubular tissues were removed by rinsing and dissociating seminiferous tubules in PBS using a pair of 25-gauge needles in a 35 mm petri dish (Corning, 430165). Dissociated seminiferous tubules were incubated in hypotonic extraction buffer (30 mM Tris-HCl pH 8.0, 50 mM sucrose, 17 mM trisodium citrate dihydrate, 5 mM EDTA, 0.5 mM Dithiothreitol (DTT) and 0.5 mM phenylmethylsulphonyl fluoride (PMSF), pH 8.2-

8.3; HEB) for 15-45 min at room temperature. After quickly rinsing seminiferous

tubules in 100 mM sucrose, some tubules were placed in 40 $\mu$l of 100 mM sucrose

on a glass depression slide. Tubules were torn to pieces using two fine forceps,

and pieces of tubular remnants were removed. The volume was increased to 40

$\mu$l by adding 100 mM sucrose and the cell suspension was made by pipetting. A

clean glass slide (Matsunami, MAS-01) was dipped into freshly made 1%

paraformaldehyde (PFA) solution pH 9.2 (pH set by 1.25 M sodium borate)

containing 0.15% Triton X-100 in a 50 ml Falcon tube, and then extra 1% PFA /

0.15% Triton X-100 solution was removed. 20 $\mu$l of cell suspension was placed

at the upper right corner of the slide and was slowly dispersed in horizontal and

vertical directions several times to make cells homogenous on the slide. Slides

were placed and dried in a closed humid chamber with hot tap water overnight at

room temperature, followed by drying with lids ajar for 3 h and then with lids

removed for 1-2 hr. Slides were washed once for 5 min in deionized water and

twice for 5 min in 0.4% DryWell (Fujifilm) in a coplin jar and air-dried at room

temperature. Slides were either directly processed for immunostaining as below

or stored in aluminum foil at -80°C prior to immunostaining.

For the preparation of testicular cell suspension, testes were dissected, the

tunica albuginea was removed, and seminiferous tubules were transferred into 1

ml of testis cell isolation medium (104 mM NaCl, 45 mM KCl, 1.2 mM $MgSO_4$, 0.6

mM $NaHPO_4$, 0.15 (w/v) glucose; TIM) containing 2 mg of collagenase

(Worthington, LS004186) for incubation at 32°C for 55 min with shaking at 53 rpm.

Tubules were pelleted by centrifugation at 1,220 x $g$ for 2 min at room temperature,

washed three times with 3 ml of TIM, and suspended in 1 ml of TIM containing

0.7 mg of Trypsin (Worthington, LS003740) and 4 $\mu$g of DNase I (Roche, 104159)

for incubation at 32°C for 15 min with shaking at 53 rpm. Trypsinization was

stopped by adding 5 mg of Trypsin inhibitor (Sigma, T9003) and 10 $\mu$g of DNase

I, and cells were separated by pipetting and filtered through a 70 $\mu$m cell strainer

(Corning, 431751). TIM was added up to 7.5 ml and cells were pelleted by

centrifugation at 1,220 x $g$ for 3 min at room temperature. After the removal of

supernatants, 0.75 $\mu$g of DNase I was added, cell pellets were dislodged and

suspended in 7.5 ml of TIM for cell counting. Pelleting was repeated, and the cell

pellet was suspended in 0.75 µg of DNase I and PBS (Nakarai, 14249-24) to 1

$x10^6$ cell/ml. 1 ml of cell suspension was then transferred to a 1.5 ml tube and

centrifuged at 200 x $g$ for 5 min at room temperature. After carefully removing the

supernatant, cells were suspended in 30 $\mu$l of HEB and incubated for 15-45 min

at room temperature. 30 µl of 100 mM sucrose was added, and 20 µl of cell

suspension was placed on a clean glass slide (Matsunami, MAS-01) coated with

100 µl of freshly made 1% paraformaldehyde (PFA) solution pH 9.2 (pH set by

1.25 M sodium borate) containing 0.15% Triton X-100. Cells were slowly

dispersed in horizontal and vertical directions several times to get homogenous

on the slide. Slides were placed and dried in a closed humid chamber with hot

tap water overnight at room temperature, followed by drying with lids ajar for 3 h

and then with lids removed for 1-2 h. Slides were dried, washed, and stored as

described above.

**Immunofluorescence staining**

Slides were rehydrated with Tris-buffered saline (TBS) pH 8.0 containing 0.05%

of Triton X-100 (TBST) for 3 min, blocked twice with blocking buffer (1% normal

goat or donkey serum, 3% bovine serum albumin (BSA), 1x TBS pH 8.0, 0.05%

of Triton X-100, 0.05% sodium azide) for 15 min at room temperature and

incubated with primary antibodies in antibody dilution buffer (10% normal goat or

donkey serum, 3% BSA, 1x TBS pH 8.0, 0.05% of Triton X-100, 0.05% sodium

azide) in a humid chamber overnight at room temperature. Slides were quickly

rinsed with TBST, washed twice with TBST for 5 min, blocked twice with blocking

buffer for 15 min at room temperature, and incubated with secondary antibodies

in antibody dilution buffer in a humid chamber for 1 h at 37°C. Slides were quickly

rinsed with TBST, washed three times with TBST for 5 min, washed once with

Milli-Q water for 2 min, and air-dried before mounting with ProLong Diamond

Antifade Mountant (Thermo Fisher Scientific, P36970). For detection of IdU,

rehydrated slides were denatured with 4N HCl for 10 min at room temperature

before blocking. EdU was detected using Click-iT EdU Imaging Kit with Alexa

Fluor 647 azides according to the manufacturer's instruction (Invitrogen). All

primary and secondary antibodies used are listed in Supplementary table 2.

**Testicular cell squashes**

Testes were dissected, the tunica albuginea was removed, and adherent extra-

tubular tissues were removed by rinsing and dissociating seminiferous tubules in

PBS using a pair of 25-gauge needles in a 35 mm petri dish (Corning, 430165).

Dissociated seminiferous tubules were fixed in freshly prepared 2%

formaldehyde containing 0.1% Triton X-100 for 10 min at room temperature.

Small pieces of tubules were placed on a clean glass slide (Matsunami, MAS-01),

torn using two fine forceps, and squashed under a coverslip with pressure from
the blunt end of pencil followed by a thumb. Slides were frozen briefly in liquid
nitrogen and either directly processed for immunostaining or stored at -80°C prior
to immunostaining. After the removal of coverslips, slides were washed three
times with PBS for 5 min and immunostaining was performed as described above.
738

**Image acquisition**
Images of mouse spermatocyte spreads testicular cell squashes were acquired
using a computer-assisted fluorescence microscope system (DeltaVision;
Applied Precision) with a 60× NA 1.4 oil immersion objective. Image
deconvolution was performed using an image workstation (SoftWorks; Applied
Precision) and afterward processed using Imaris (Oxford Instruments, UK) and
Photoshop (Adobe, USA) software tools. Images in each figure are matched
exposure.
747

**Image analysis**
Comparisons were made between animals that were either littermate or matched
by age. Single cells were manually cropped, and numbers of foci were counted
by the auto-thresholding signal intensity in Imaris software (Oxford Instrument)
and clear foci that were missed by auto-thresholding, especially adjacent foci,
were manually counted. Signal intensities were measured using Image J software
by subtracting signal intensities of single cells from those of regions of interest
(ROIs) manually drawn next to cells as backgrounds.
Meiotic prophase I stages on spermatocyte spreads were defined by
SYCP3 and SYCP1 staining using standard criteria. Leptonema was defined by
short SYCP3 stretches without evidence of synapsis determined by SYCP1
staining. Zygonema was defined by longer stretches of SYCP3 with various
degrees of synapsis: early, mid- and late zygonema was defined by having <25%,
25-75%, and >75% of synapsis, respectively. Pachynema was defined by the full
synapsis of all autosomes. Diplonema was defined by de-synapsis beginning
from mostly the middle of chromosomes with various degrees of residual
synapsis. Preleptonema was defined by very short SYCP3 stretches with EdU or
IdU-positive staining.
766

**EdU incorporation**
1 ml of 1 x$10^6$ cell/ml testicular cell suspension in PBS was centrifuged at 200 x

*g* for 5 min at room temperature, and after removal of the supernatant, cells were
suspended in 500 μl of D-MEM (Wako, 044-29765) / 10% FBS containing 10 μM
EdU. After incubation at 37°C for 1 h in a well of a 12-well dish (Violamo), cells
were pelleted by centrifugation at 200 x *g* for 5 min at room temperature, washed
in 1 ml of PBS once, and suspended in 30 μl HEB for incubation for 15-30 min at
room temperature. After adding 30 μl of 100 mM sucrose, cells were fixed, and
the slides were dried, washed, and stored as described above.
776

**IdU pulse incorporation**
778   1 ml of 1 x10$^6$ cell/ml testicular cell suspension in PBS was centrifuged at 200 x
779   *g* for 5 min at room temperature, and after removal of the supernatant, cells were
780   suspended in 500 μl of D-MEM (Wako, 044-29765) / 10% FBS containing 20 μM
781   IdU (Sigma, I7125). After incubation at 37°C for 10 min in a well of a 12-well dish
782   (Violamo), cells were subjected to chromosome spreads as described above.
783

**Rad51 Chromatin immuno-precipitation – single-stranded DNA sequencing**
**(ChIP-SSDS)**
ChIP was performed as described previously [85]. One testis from the control and
a pair of testes from *Fignl1* cKO mice (14-18 weeks) were dissected, the tunica
albuginea was removed, and seminiferous tubules were transferred into 10 ml of
1% formaldehyde (Pierce, 28908) in PBS (Nakarai, 14249-24) and immediately
homogenized in a Dounce homogenizer with 20 strokes of the tight-fit pestle. The
homogenate was passed through a 70 μm cell strainer (Falcon, 352350), and
after a total of 10 min of incubation at room temperature, crosslinking was
quenched by adding 2.5 M glycine to a final concentration of 125 mM. After 5 min
of incubation at room temperature, cells were pelleted by centrifugation at 900 x
*g* for 5 min at 4°C, washed once with 10 ml of cold PBS, and suspended in 1 ml
of cold PBS. The cell suspension was transferred to a 1.5 ml tube, centrifuged at
900 x *g* for 5 min at 4°C, and after removal of the supernatant, the cell pellet was
snap-frozen on liquid nitrogen and stored at -80°C. The frozen cell pellet was
suspended in 500 μl of cell lysis buffer (0.25% Triton X-100, 10 mM EDTA, 0.5
mM EGTA, 10 mM Tris-HCl pH 8.0) supplemented with protease inhibitor (Roche,
04693159001), incubated for 10 min on ice, centrifuged at 300 x *g* for 3 min at
4°C, and the nuclei pellet was resuspended in 500 μl of lysis wash buffer (200
mM NaCl, 1 mM EDTA, 0.5 mM EGTA, 10 mM Tris-HCl pH 8.0) supplemented
with protease inhibitor. After centrifugation at 300 x *g* for 3 min at 4°C, the nuclei

pellet was resuspended in 200 μl of RIPA buffer (10 mM Tris-HCl pH 8.0, 1 mM
EDTA, 0.5 mM EGTA, 1% Triton X-100, 0.1% sodium deoxycholate, 0.1% SDS,
140 mM NaCl) supplemented with protease inhibitor and split into two ~130 μl
aliquots and sonicated on the Covaris M220 for 5 min 30 sec at 5-9°C with a
setting of Peak Power 50.0, Duty Factor 20.0, and Cycles/Burst 200. Sheared
nuclei were combined, and after adding 700 μl of RIPA buffer, centrifuged at
15,000 rpm for 15 min at 4°C to remove insoluble debris. 800 μl of the chromatin-
containing supernatant was mixed with 50 μl of Dynabeads protein G bound to
9.7 μg of RAD51 antibody (Novus, NB100-148) and incubated overnight on a
rotating wheel at 4°C for immunoprecipitation. Beads were washed once with 1
ml of cold ChIP wash buffer 1 (0.1% SDS, 1% Triton X-100, 2 mM EDTA, 20 mM
Tris-HCl pH 8.0, 150 mM NaCl), 1 ml of cold ChIP wash buffer 2 (0.1% SDS, 1%
Triton X-100, 2 mM EDTA, 20 mM Tris-HCl pH 8.0, 500 mM NaCl), 1 ml of cold
ChIP wash buffer 3 (0.25 M LiCl, 1% NP-40, 1% sodium deoxycholate, 1 mM
EDTA, 10 mM Tris-HCl pH 8.0), and twice with 1 ml of cold TE (1 mM EDTA and
10 mM Tris-HCl pH 8.0), followed by elution with 100 μl of elution buffer (0.1 M
NaHCO$_3$ and 1% SDS) for 30 min at 65°C with vortexing every 10 min. The elutes
were incubated overnight at 65°C for de-crosslinking, neutralized by adding 4 μl
of 1M Tris-HCl (pH 6.5) and 2 μl of 0.5 M EDTA, and RNA was digested by adding
2 μl of 10 mg/ml RNase-A and incubation for 15 min 37°C. After adding 5 μl of 10
mg/ml Proteinase-K (Wako, 165-21043) and incubation for 1 h at 55°C, DNA was
purified using QIAquick PCR Purification Kit (QIAGEN).

SSDS was performed as described previously (Brick et al., 2018). 7-10 ng
of ChIP DNA was further sonicated on the Covaris M220 for 8 min 30 sec at room
temperature with a setting of Peak Power 75.0, Duty Factor 10.0, and
Cycles/Burst 200, followed by end-repair and dA-tailing using ChIP-Seq Library
Prep Reagent Set for Illumina (NEB, E6200L). DNA was incubated for 3 min at
95°C and returned to room temperature for kinetic enrichment for single-stranded
DNA. After adapter ligation using IDT for Illumina TruSeq DNA UD Indexes
(Illumina, 20022370), SSDS libraries were amplified during 15 cycles of PCR
using TruSeq DNA Nano LP (Illumina, 20016328), purified with AMPureXP beads
(Beckman Coulter, A63880), and sequenced on HiSeq2500 in the rapid mode in
a 50 bp PE run. Around 70 million paired reads were generated per sample.

A published bioinformatic pipeline (https://nf-co.re/ssds/dev) was run for the
resulting reads for the identification of single-stranded sequences and mapping
to the mm10 genome [109]. Heatmaps were generated using Deeptools [110].

**Western blot**

Testes were dissected, the tunica albuginea was removed, and seminiferous
tubules were snap frozen on liquid nitrogen and stored at -80°C before cell lysis.
Tubules from either one testis (*Fignl1*$^{+/+}$ *Stra8-Cre*$^+$ and *Fignl1*$^{flox/\Delta}$) or two testes
(*Fignl1*$^{flox/\Delta}$ *Stra8-Cre*$^+$) were homogenized in 1 ml of RIPA buffer (50 mM Tris-HCl
pH 7.5, 150 mM NaCl, 1 mM EDTA, 1% NP-40, 0.5% sodium deoxycholate, 0.1%
sodium dodecyl sulfate (SDS)) supplemented with protease inhibitor (Roche,
04693159001)) on ice, incubated for 15 min on ice, sonicated with a Bioruptor for
30 min (30 cycles of 30 sec ON / 30 sec OFF) at 4°C with a setting of High
intensity, and incubated for 15 min on ice. The samples were centrifuged at
14,000 rpm for 15 min at 4°C and the supernatants were collected as whole-testis
extracts. Protein concentrations were measured by Brad-ford assay and the
concentrations between samples were made constant by adding RIPA buffer
supplemented with protease inhibitor. Whole-testis extract was mixed with 4x
SDS-PAGE loading buffer (250 mM Tris-HCl pH 6.8, 8% SDS, 40% glycerol, 572
mM β-mercaptoethanol, 0.05% bromophenol blue) and boiled for 5 min at 100°C
and subjected to immunoblotting.

Protein samples were separated on 10% gels by SDS-PAGE and
transferred to PVDF membranes by wet transfer method in Towbin buffer (25 mM
Tris, 192 mM glycine, 20% ethanol) at 60 V for 150 min at 4°C. Membranes were
blocked with PBS containing 2.5% non-fat milk for 1 h at room temperature and
incubated with primary antibodies in 2.5% non-fat milk in PBS overnight at 4°C.
Membranes were washed three times with PBS containing 0.05% Tween-20
(PBST) for 10 min and incubated with HRP-conjugated secondary antibodies in
PBST for 1 h at room temperature. Membranes were washed three times with
PBST for 10 min and the signal was developed by ECL Prime Western Blotting
Detection Reagent (Amersham) and detected by ImageQuant LAS4000 (GE
Healthcare).

**Antibody generation**

His-tagged mouse SYCP3 was overexpressed from pET28c-SYCP3 plasmid by
the addition of 0.2 mM IPTG in BL21(DE3) at 37°C for 4 h. Cell extracts were
prepared in Lysis buffer (50 mM Tris-HCl pH 7.5, 500 mM NaCl) by sonication
and the supernatants were collected after centrifugation at 20,400 x *g* for 30 min
at 4°C. After adding Tween-20 to a final concentration of 0.2%, the cell lysates

were incubated with Ni-NTA Agarose (QIAGEN) for 2 h at 4°C and the lysate-
resin slurry was washed in Lysis buffer containing 50 mM Imidazole, 100 mM
Imidazole, 300 mM Imidazole. Ni-NTA-bound His-SYCP3 was eluted in Lysis
buffer containing 3M Imidazole and 3M Urea (elution 1) by incubation for 1 h at
4°C and in Lysis buffer containing 0.1% SDS (elution 2) by incubation for 30 min
at room temperature. Elution 2 was concentrated using MICROCON Centrifugal
Filter Device YM-10 (Millipore) and subjected to immunization to a rat.
884

**Purification of RAD51 and FINGL1ΔN**
GST-tagged human FIGNL1ΔN (N-terminal 284 aa deletion) was overexpressed
from pGEX-6P-1-*Fignl1ΔN* plasmid by the addition of 0.25 mM IPTG in
BL21(DE3) pLysS at 30°C for overnight. Cell extracts were prepared in Lysis
buffer (25 mM Tris-HCl pH 7.6, 150 mM NaCl, 10% Glycerol, 1 mM DTT) and
GST-FIGNL1ΔN was recovered by GSTrap HP column (Cytiva). FIGNL1ΔN was
eluted from the GSTrap column by the cleavage of GST tag by Prescission
protease (#27-0843-01, Cytiva) in SP buffer (20 mM HEPES-NaOH pH 7.5, 10%
Glycerol, 1 mM DTT) supplemented with 100 mM NaCl. FIGNL1ΔN was further
purified by HiTrap SP column (Cytiva) with a linear gradient of 100 mM-1M in SP
buffer. The peak fractions are dialyzed against SP buffer supplemented with 100
mM NaCl, concentrated with Amicon Ultra 30K (Millipore), and stored at -80°C.
Human RAD51 was purified as described previously using pET-HsRad51 plasmid
[111].
899

**Electrophoresis mobility shift assay (EMSA)**
Double-stranded DNA substrate was prepared by annealing 5'-Cy5-labeled, 40-
mer oligo DNA (Oligo 1, Cy5-
TAATACAAAATAAGTAAATGAATAAACAGAGAAAATAAAG, IDT) and its
complementary DNA (Oligo 2,
CTTTATTTTCTCTGTTTATTCATTTACTTATTTTGTATTA, Fasmac). Oligo 1
was used as an ssDNA substrate. For the assay, DNA (0.8 μM in bp/nt) was pre-
incubated with 1 μM of RAD51 in 10 μl of the buffer (20 mM HEPES-NaOH pH7.5,
100 ug/ml BSA, 1 mM DTT, 10 mM MgCl$_2$, 2 mM ATP) at 37°C for 5 min to form
RAD51-DNA complex. Then the indicated concentration of FIGNL1ΔN was added
to the reaction and further incubated at 37°C for 15 min. In the control reaction
without FIGNL1ΔN, the same volume of dialysis buffer was added instead of the
protein. DNA-protein complexes were analyzed using 8% native polyacrylamide

gel electrophoresis with 1×TBE. Cy5-labelled DNA was visualized by LAS4000

(GE healthcare) and unbound DNA was quantified using ImageQuant TL (GE

healthcare). Images were further processed to adjust brightness using Photoshop

(Adobe, USA) software.

**Statistical analysis and reproducibility**

Statistical analyses were performed using Graphpad Prism software v.9 and R

version 4.2.2 (https://www.r-project.org). Bars in figures and values in

manuscripts indicate means ± standard deviations (SDs) otherwise mentioned.

Statistic parameters and tests and sample sizes are described in the figures

and/or figure legends. Sample sizes were not predetermined using any statistical

tests.

At least two animals of each genotype were analyzed and similar results

were obtained otherwise mentioned. Three independent experiments were

performed for biochemical analysis.

**Data availability**

The raw and processed sequencing data have been deposited in the Gene

Expression Omnibus (GEO) repository under accession number GSE227944.

**Acknowledgments**

We thank Drs. F. Baudat and B. de Massy for sharing unpublished results. We

acknowledge Drs. F. Pratto and V. Jayakumar for SSDS analysis. We are grateful

to Ms. S. Kondo for the generation of anti-SYCP3. We acknowledge Drs. S.

Keeney, Y. Fujiwara, and K. Ishiguro for *spo11* mice and Dr. H. Kurumizaka for

the RAD51 expression plasmid. We are also indebted to members of the

Shinohara lab, particularly Ms. S. Aoyama, A. Maeda, C. Watanabe, and S.

Hashimoto for mouse care. This work was supported by JSPS KAKENHI Grant

Numbers; 19H00981 to A.S., 20K15716 and 16H06279 (PAGS) to M.I., and

19H03157 to A.F.

**Author contributions**

M.I. and A.S. conceived and designed the experiments. M.I. performed all

experiments on mouse meiosis. M.I., Y.F., and A.T. analyzed SSDS data. A.F.

performed biochemical experiments. K.M. set up Fignl1 mice. M.I. and A.S.

analyzed the data. A.S. prepared the original draft and wrote the manuscript with

help from M.I.

**Competing interest**

The authors declare no competing interests.

**References**

**References**

1.    Sonoda, E. *et al.* Rad51-deficient vertebrate cells accumulate
chromosomal breaks prior to cell death. *EMBO J* **17**, 598-608 (1998).

2.    Tashiro, S., Walter, J., Shinohara, A., Kamada, N. & Cremer, T. Rad51
accumulation at sites of DNA damage and in postreplicative chromatin. *J*
*Cell Biol* **150**, 283-291 (2000).

3.    Mason, J.M., Chan, Y.L., Weichselbaum, R.W. & Bishop, D.K. Non-
enzymatic roles of human RAD51 at stalled replication forks. *Nat Commun*
**10**, 4410 (2019).

4.    Schlacher, K. *et al.* Double-strand break repair-independent role for
BRCA2 in blocking stalled replication fork degradation by MRE11. *Cell* **145**,
529-542 (2011).

5.    Feretzaki, M. *et al.* RAD51-dependent recruitment of TERRA lncRNA to
telomeres through R-loops. *Nature* **587**, 303-308 (2020).

6.    Ogawa, T., Yu, X., Shinohara, A. & Egelman, E.H. Similarity of the yeast
RAD51 filament to the bacterial RecA filament. *Science* **259**, 1896-1899
(1993).

7.    Shinohara, A., Ogawa, H. & Ogawa, T. Rad51 protein involved in repair
and recombination in S. cerevisiae is a RecA-like protein. *Cell* **69**, 457-470
(1992).

8.    Sung, P. Catalysis of ATP-dependent homologous DNA pairing and strand
exchange by yeast RAD51 protein. *Science* **265**, 1241-1243 (1994).

9.    Sung, P. & Robberson, D.L. DNA strand exchange mediated by a RAD51-
ssDNA nucleoprotein filament with polarity opposite to that of RecA. *Cell*
**82**, 453-461 (1995).

10.    Jensen, R.B., Carreira, A. & Kowalczykowski, S.C. Purified human BRCA2
stimulates RAD51-mediated recombination. *Nature* **467**, 678-683 (2010).

11.    Zhao, W. *et al.* BRCA1-BARD1 promotes RAD51-mediated homologous
DNA pairing. *Nature* **550**, 360-365 (2017).

12.    New, J.H., Sugiyama, T., Zaitseva, E. & Kowalczykowski, S.C. Rad52
protein stimulates DNA strand exchange by Rad51 and replication protein
A. *Nature* **391**, 407-410 (1998).

13.    Shinohara, A. & Ogawa, T. Stimulation by Rad52 of yeast Rad51-mediated
recombination. *Nature* **391**, 404-407 (1998).

14.    Van Dyck, E., Stasiak, A.Z., Stasiak, A. & West, S.C. Binding of double-

strand breaks in DNA by human Rad52 protein. *Nature* **398**, 728-731
(1999).

15.   Akamatsu, Y., Dziadkowiec, D., Ikeguchi, M., Shinagawa, H. & Iwasaki, H.
Two different Swi5-containing protein complexes are involved in mating-
type switching and recombination repair in fission yeast. *Proc Natl Acad*
*Sci U S A* **100**, 15770-15775 (2003).

16.   Tsai, S.P. *et al.* Rad51 presynaptic filament stabilization function of the
mouse Swi5-Sfr1 heterodimeric complex. *Nucleic Acids Res* **40**, 6558-
6569 (2012).

17.   Masson, J.Y. *et al.* Identification and purification of two distinct complexes
containing the five RAD51 paralogs. *Genes Dev* **15**, 3296-3307 (2001).

18.   Abreu, C.M. *et al.* Shu complex SWS1-SWSAP1 promotes early steps in
mouse meiotic recombination. *Nat Commun* **9**, 3961 (2018).

19.   Liu, T., Wan, L., Wu, Y., Chen, J. & Huang, J. hSWS1.SWSAP1 is an
evolutionarily conserved complex required for efficient homologous
recombination repair. *J Biol Chem* **286**, 41758-41766 (2011).

20.   Martino, J. *et al.* The human Shu complex functions with PDS5B and
SPIDR to promote homologous recombination. *Nucleic Acids Res* **47**,
10151-10165 (2019).

21.   Lu, C.H. *et al.* Swi5-Sfr1 stimulates Rad51 recombinase filament assembly
by modulating Rad51 dissociation. *Proc Natl Acad Sci U S A* **115**, E10059-
E10068 (2018).

22.   Tanaka, K., Kagawa, W., Kinebuchi, T., Kurumizaka, H. & Miyagawa, K.
Human Rad54B is a double-stranded DNA-dependent ATPase and has
biochemical properties different from its structural homolog in yeast,
Tid1/Rdh54. *Nucleic Acids Res* **30**, 1346-1353 (2002).

23.   Zhang, Z., Fan, H.Y., Goldman, J.A. & Kingston, R.E. Homology-driven
chromatin remodeling by human RAD54. *Nat Struct Mol Biol* **14**, 397-405
(2007).

24.   Liang, F. *et al.* Promotion of RAD51-Mediated Homologous DNA Pairing
by the RAD51AP1-UAF1 Complex. *Cell Rep* **15**, 2118-2126 (2016).

25.   Parplys, A.C. *et al.* NUCKS1 is a novel RAD51AP1 paralog important for
homologous recombination and genome stability. *Nucleic Acids Res* **43**,
9817-9834 (2015).

26.   Wiese, C. *et al.* Promotion of homologous recombination and genomic
stability by RAD51AP1 via RAD51 recombinase enhancement. *Mol Cell*

**28**, 482-490 (2007).

27.   Bugreev, D.V., Mazina, O.M. & Mazin, A.V. Bloom syndrome helicase
stimulates RAD51 DNA strand exchange activity through a novel
mechanism. *J Biol Chem* **284**, 26349-26359 (2009).

28.   Bugreev, D.V., Yu, X., Egelman, E.H. & Mazin, A.V. Novel pro- and anti-
recombination activities of the Bloom's syndrome helicase. *Genes Dev* **21**,
3085-3094 (2007).

29.   Xue, C. *et al.* Single-molecule visualization of human BLM helicase as it
acts upon double- and single-stranded DNA substrates. *Nucleic Acids Res*
**47**, 11225-11237 (2019).

30.   Sommers, J.A. *et al.* FANCJ uses its motor ATPase to destabilize protein-
DNA complexes, unwind triplexes, and inhibit RAD51 strand exchange. *J*
*Biol Chem* **284**, 7505-7517 (2009).

31.   Schwendener, S. *et al.* Physical interaction of RECQ5 helicase with
RAD51 facilitates its anti-recombinase activity. *J Biol Chem* **285**, 15739-
15745 (2010).

32.   Xue, C. *et al.* Single-molecule visualization of human RECQ5 interactions
with single-stranded DNA recombination intermediates. *Nucleic Acids Res*
**49**, 285-305 (2021).

33.   Chu, W.K. *et al.* FBH1 influences DNA replication fork stability and
homologous recombination through ubiquitylation of RAD51. *Nat Commun*
**6**, 5931 (2015).

34.   Simandlova, J. *et al.* FBH1 helicase disrupts RAD51 filaments in vitro and
modulates homologous recombination in mammalian cells. *J Biol Chem*
**288**, 34168-34180 (2013).

35.   Moldovan, G.L. *et al.* Inhibition of homologous recombination by the
PCNA-interacting protein PARI. *Mol Cell* **45**, 75-86 (2012).

36.   Adolph, M.B. *et al.* RADX controls RAD51 filament dynamics to regulate
replication fork stability. *Mol Cell* **81**, 1074-1083 e1075 (2021).

37.   Dungrawala, H. *et al.* RADX Promotes Genome Stability and Modulates
Chemosensitivity by Regulating RAD51 at Replication Forks. *Mol Cell* **67**,
374-386 e375 (2017).

38.   Croteau, D.L., Popuri, V., Opresko, P.L. & Bohr, V.A. Human RecQ
helicases in DNA repair, recombination, and replication. *Annu Rev*
*Biochem* **83**, 519-552 (2014).

39.   Scully, R., Panday, A., Elango, R. & Willis, N.A. DNA double-strand break

repair-pathway choice in somatic mammalian cells. *Nat Rev Mol Cell Biol*
**20**, 698-714 (2019).

40.  Grey, C. & de Massy, B. Chromosome Organization in Early Meiotic
Prophase. *Front Cell Dev Biol* **9**, 688878 (2021).

41.  Ito, M. & Shinohara, A. Chromosome architecture and homologous
recombination in meiosis. *Front Cell Dev Biol* **10**, 1097446 (2022).

42.  Yadav, V.K. & Claeys Bouuaert, C. Mechanism and Control of Meiotic DNA
Double-Strand Break Formation in S. cerevisiae. *Front Cell Dev Biol* **9**,
642737 (2021).

43.  Bishop, D.K., Park, D., Xu, L. & Kleckner, N. DMC1: a meiosis-specific
yeast homolog of E. coli recA required for recombination, synaptonemal
complex formation, and cell cycle progression. *Cell* **69**, 439-456 (1992).

44.  Brown, M.S. & Bishop, D.K. DNA strand exchange and RecA homologs in
meiosis. *Cold Spring Harb Perspect Biol* **7**, a016659 (2014).

45.  Crickard, J.B. & Greene, E.C. The biochemistry of early meiotic
recombination intermediates. *Cell Cycle* **17**, 2520-2530 (2018).

46.  Deans, B., Griffin, C.S., Maconochie, M. & Thacker, J. Xrcc2 is required
for genetic stability, embryonic neurogenesis and viability in mice. *EMBO*
*J* **19**, 6675-6685 (2000).

47.  Kuznetsov, S.G., Haines, D.C., Martin, B.K. & Sharan, S.K. Loss of
Rad51c leads to embryonic lethality and modulation of Trp53-dependent
tumorigenesis in mice. *Cancer Res* **69**, 863-872 (2009).

48.  Lim, D.S. & Hasty, P. A mutation in mouse rad51 results in an early
embryonic lethal that is suppressed by a mutation in p53. *Mol Cell Biol* **16**,
7133-7143 (1996).

49.  Pittman, D.L. & Schimenti, J.C. Midgestation lethality in mice deficient for
the RecA-related gene, Rad51d/Rad51l3. *Genesis* **26**, 167-173 (2000).

50.  Prakash, R. *et al.* XRCC3 loss leads to midgestational embryonic lethality
in mice. *DNA Repair (Amst)* **108**, 103227 (2021).

51.  Shu, Z., Smith, S., Wang, L., Rice, M.C. & Kmiec, E.B. Disruption of
muREC2/RAD51L1 in mice results in early embryonic lethality which can
Be partially rescued in a p53(-/-) background. *Mol Cell Biol* **19**, 8686-8693
(1999).

52.  Tsuzuki, T. *et al.* Targeted disruption of the Rad51 gene leads to lethality
in embryonic mice. *Proc Natl Acad Sci U S A* **93**, 6236-6240 (1996).

53.  Matsuzaki, K., Kondo, S., Ishikawa, T. & Shinohara, A. Human RAD51

paralogue SWSAP1 fosters RAD51 filament by regulating the anti-
recombinase FIGNL1 AAA+ ATPase. *Nat Commun* **10**, 1407 (2019).

54. Prakash, R. *et al.* Distinct pathways of homologous recombination
controlled by the SWS1-SWSAP1-SPIDR complex. *Nat Commun* **12**, 4255
(2021).

55. Shang, Y. *et al.* MEIOK21: a new component of meiotic recombination
bridges required for spermatogenesis. *Nucleic Acids Res* **48**, 6624-6639
(2020).

56. Takemoto, K. *et al.* Meiosis-Specific C19orf57/4930432K21Rik/BRME1
Modulates Localization of RAD51 and DMC1 to DSBs in Mouse Meiotic
Recombination. *Cell Rep* **31**, 107686 (2020).

57. Zhang, J., Fujiwara, Y., Yamamoto, S. & Shibuya, H. A meiosis-specific
BRCA2 binding protein recruits recombinases to DNA double-strand
breaks to ensure homologous recombination. *Nat Commun* **10**, 722 (2019).

58. Zhang, J. *et al.* The BRCA2-MEILB2-BRME1 complex governs meiotic
recombination and impairs the mitotic BRCA2-RAD51 function in cancer
cells. *Nat Commun* **11**, 2055 (2020).

59. Chester, N., Kuo, F., Kozak, C., O'Hara, C.D. & Leder, P. Stage-specific
apoptosis, developmental delay, and embryonic lethality in mice
homozygous for a targeted disruption in the murine Bloom's syndrome
gene. *Genes Dev* **12**, 3382-3393 (1998).

60. Wesoly, J. *et al.* Differential contributions of mammalian Rad54 paralogs
to recombination, DNA damage repair, and meiosis. *Mol Cell Biol* **26**, 976-
989 (2006).

61. McNally, F.J. & Roll-Mecak, A. Microtubule-severing enzymes: From
cellular functions to molecular mechanism. *J Cell Biol* **217**, 4057-4069
(2018).

62. Yuan, J. & Chen, J. FIGNL1-containing protein complex is required for
efficient homologous recombination repair. *Proc Natl Acad Sci U S A* **110**,
10640-10645 (2013).

63. Fernandes, J.B. *et al.* FIGL1 and its novel partner FLIP form a conserved
complex that regulates homologous recombination. *PLoS Genet* **14**,
e1007317 (2018).

64. Girard, C. *et al.* AAA-ATPase FIDGETIN-LIKE 1 and Helicase FANCM
Antagonize Meiotic Crossovers by Distinct Mechanisms. *PLoS Genet* **11**,
e1005369 (2015).

65.  Hu, Q. *et al.* Meiotic Chromosome Association 1 Interacts with TOP3alpha
and Regulates Meiotic Recombination in Rice. *Plant Cell* **29**, 1697-1708
(2017).
66.  Yang, S. *et al.* FIGNL1 Inhibits Non-homologous Chromosome
Association and Crossover Formation. *Front Plant Sci* **13**, 945893 (2022).
67.  Kumar, R., Duhamel, M., Coutant, E., Ben-Nahia, E. & Mercier, R.
Antagonism between BRCA2 and FIGL1 regulates homologous
recombination. *Nucleic Acids Res* **47**, 5170-5180 (2019).
68.  Sharan, S.K. *et al.* Embryonic lethality and radiation hypersensitivity
mediated by Rad51 in mice lacking Brca2. *Nature* **386**, 804-810 (1997).
69.  Sadate-Ngatchou, P.I., Payne, C.J., Dearth, A.T. & Braun, R.E. Cre
recombinase activity specific to postnatal, premeiotic male germ cells in
transgenic mice. *Genesis* **46**, 738-742 (2008).
70.  Yuan, L., Brundell, E. & Hoog, C. Expression of the meiosis-specific
synaptonemal complex protein 1 in a heterologous system results in the
formation of large protein structures. *Exp Cell Res* **229**, 272-275 (1996).
71.  Yuan, L. *et al.* The murine SCP3 gene is required for synaptonemal
complex assembly, chromosome synapsis, and male fertility. *Mol Cell* **5**,
73-83 (2000).
72.  Yuan, L. *et al.* The synaptonemal complex protein SCP3 can form
multistranded, cross-striated fibers in vivo. *J Cell Biol* **142**, 331-339 (1998).
73.  Mahadevaiah, S.K. *et al.* Recombinational DNA double-strand breaks in
mice precede synapsis. *Nat Genet* **27**, 271-276 (2001).
74.  Luo, M. *et al.* MEIOB exhibits single-stranded DNA-binding and
exonuclease activities and is essential for meiotic recombination. *Nat
Commun* **4**, 2788 (2013).
75.  Souquet, B. *et al.* MEIOB targets single-strand DNA and is necessary for
meiotic recombination. *PLoS Genet* **9**, e1003784 (2013).
76.  Yoshida, K. *et al.* The mouse RecA-like gene Dmc1 is required for
homologous chromosome synapsis during meiosis. *Mol Cell* **1**, 707-718
(1998).
77.  Ashley, T. *et al.* Dynamic changes in Rad51 distribution on chromatin
during meiosis in male and female vertebrates. *Chromosoma* **104**, 19-28
(1995).
78.  Ikeya, T., Shinohara, A., Sato, S., Tabata, S. & Ogawa, T. Localization of
mouse Rad51 and Lim15 proteins on meiotic chromosomes at late stages

of prophase 1. *Genes Cells* **1**, 379-389 (1996).

79.  Tarsounas, M., Morita, T., Pearlman, R.E. & Moens, P.B. RAD51 and
DMC1 form mixed complexes associated with mouse meiotic
chromosome cores and synaptonemal complexes. *J Cell Biol* **147**, 207-
220 (1999).

80.  Shi, B. *et al.* Dual functions for the ssDNA-binding protein RPA in meiotic
recombination. *PLoS Genet* **15**, e1007952 (2019).

81.  Moens, P.B. *et al.* The time course and chromosomal localization of
recombination-related proteins at meiosis in the mouse are compatible
with models that can resolve the early DNA-DNA interactions without
reciprocal recombination. *J Cell Sci* **115**, 1611-1622 (2002).

82.  Baker, S.M. *et al.* Involvement of mouse Mlh1 in DNA mismatch repair and
meiotic crossing over. *Nat Genet* **13**, 336-342 (1996).

83.  Baudat, F., Manova, K., Yuen, J.P., Jasin, M. & Keeney, S. Chromosome
synapsis defects and sexually dimorphic meiotic progression in mice
lacking Spo11. *Mol Cell* **6**, 989-998 (2000).

84.  Romanienko, P.J. & Camerini-Otero, R.D. The mouse Spo11 gene is
required for meiotic chromosome synapsis. *Mol Cell* **6**, 975-987 (2000).

85.  Hinch, A.G. *et al.* The Configuration of RPA, RAD51, and DMC1 Binding
in Meiosis Reveals the Nature of Critical Recombination Intermediates.
*Mol Cell* **79**, 689-701 e610 (2020).

86.  Khil, P.P., Smagulova, F., Brick, K.M., Camerini-Otero, R.D. & Petukhova,
G.V. Sensitive mapping of recombination hotspots using sequencing-
based detection of ssDNA. *Genome Res* **22**, 957-965 (2012).

87.  Smagulova, F. *et al.* Genome-wide analysis reveals novel molecular
features of mouse recombination hotspots. *Nature* **472**, 375-378 (2011).

88.  Altmannova, V. *et al.* The role of bivalent ions in the regulation of D-loop
extension mediated by DMC1 during meiotic recombination. *iScience* **25**,
105439 (2022).

89.  Li, X. & Heyer, W.D. RAD54 controls access to the invading 3'-OH end
after RAD51-mediated DNA strand invasion in homologous recombination
in Saccharomyces cerevisiae. *Nucleic Acids Res* **37**, 638-646 (2009).

90.  Li, X., Stith, C.M., Burgers, P.M. & Heyer, W.D. PCNA is required for
initiation of recombination-associated DNA synthesis by DNA polymerase
delta. *Mol Cell* **36**, 704-713 (2009).

91.  Bugreev, D.V. *et al.* The resistance of DMC1 D-loops to dissociation may

account for the DMC1 requirement in meiosis. *Nat Struct Mol Biol* **18**, 56-
60 (2011).

92.   Finsterbusch, F. *et al.* Alignment of Homologous Chromosomes and
Effective Repair of Programmed DNA Double-Strand Breaks during
Mouse Meiosis Require the Minichromosome Maintenance Domain
Containing 2 (MCMDC2) Protein. *PLoS Genet* **12**, e1006393 (2016).

93.   Her, C., Wu, X., Bailey, S.M. & Doggett, N.A. Mouse MutS homolog 4 is
predominantly expressed in testis and interacts with MutS homolog 5.
*Mamm Genome* **12**, 73-76 (2001).

94.   Kneitz, B. *et al.* MutS homolog 4 localization to meiotic chromosomes is
required for chromosome pairing during meiosis in male and female mice.
*Genes Dev* **14**, 1085-1097 (2000).

95.   Li, X.C. & Schimenti, J.C. Mouse pachytene checkpoint 2 (trip13) is
required for completing meiotic recombination but not synapsis. *PLoS*
*Genet* **3**, e130 (2007).

96.   Lutzmann, M. *et al.* MCM8- and MCM9-deficient mice reveal
gametogenesis defects and genome instability due to impaired
homologous recombination. *Mol Cell* **47**, 523-534 (2012).

97.   McNairn, A.J., Rinaldi, V.D. & Schimenti, J.C. Repair of Meiotic DNA
Breaks and Homolog Pairing in Mouse Meiosis Requires a
Minichromosome Maintenance (MCM) Paralog. *Genetics* **205**, 529-537
(2017).

98.   Roig, I. *et al.* Mouse TRIP13/PCH2 is required for recombination and
normal higher-order chromosome structure during meiosis. *PLoS Genet* **6**
(2010).

99.   Benson, F.E., Stasiak, A. & West, S.C. Purification and characterization of
the human Rad51 protein, an analogue of E. coli RecA. *EMBO J* **13**, 5764-
5771 (1994).

100.   Dupaigne, P. *et al.* Rad51 polymerization reveals a new chromatin
remodeling mechanism. *PLoS One* **3**, e3643 (2008).

101.   North, J.A. *et al.* ATP-dependent nucleosome unwrapping catalyzed by
human RAD51. *Nucleic Acids Res* **41**, 7302-7312 (2013).

102.   Senavirathne, G., Mahto, S.K., Hanne, J., O'Brian, D. & Fishel, R. Dynamic
unwrapping of nucleosomes by HsRAD51 that includes sliding and
rotational motion of histone octamers. *Nucleic Acids Res* **45**, 685-698
(2017).

103. Holzen, T.M., Shah, P.P., Olivares, H.A. & Bishop, D.K. Tid1/Rdh54
promotes dissociation of Dmc1 from nonrecombinogenic sites on meiotic
chromatin. *Genes Dev* **20**, 2593-2604 (2006).

104. Shah, P.P. *et al.* Swi2/Snf2-related translocases prevent accumulation of
toxic Rad51 complexes during mitotic growth. *Mol Cell* **39**, 862-872 (2010).

105. Mason, J.M. *et al.* RAD54 family translocases counter genotoxic effects of
RAD51 in human tumor cells. *Nucleic Acids Res* **43**, 3180-3196 (2015).

106. Tokuhiro, K. *et al.* Calreticulin is required for development of the cumulus
oocyte complex and female fertility. *Sci Rep* **5**, 14254 (2015).

107. de Boer, E., Lhuissier, F.G. & Heyting, C. Cytological analysis of
interference in mouse meiosis. *Methods Mol Biol* **558**, 355-382 (2009).

108. Peters, A.H., Plug, A.W., van Vugt, M.J. & de Boer, P. A drying-down
technique for the spreading of mammalian meiocytes from the male and
female germline. *Chromosome Res* **5**, 66-68 (1997).

109. Brick, K., Pratto, F., Sun, C.Y., Camerini-Otero, R.D. & Petukhova, G.
Analysis of Meiotic Double-Strand Break Initiation in Mammals. *Methods*
*Enzymol* **601**, 391-418 (2018).

110. Ramirez, F. *et al.* deepTools2: a next generation web server for deep-
sequencing data analysis. *Nucleic Acids Res* **44**, W160-165 (2016).

111. Matsuo, Y., Sakane, I., Takizawa, Y., Takahashi, M. & Kurumizaka, H.
Roles of the human Rad51 L1 and L2 loops in DNA binding. *FEBS J* **273**,
3148-3159 (2006).

112. Lange, J. *et al.* The Landscape of Mouse Meiotic Double-Strand Break
Formation, Processing, and Repair. *Cell* **167**, 695-708 e616 (2016).

113. Boekhout, M. *et al.* REC114 Partner ANKRD31 Controls Number, Timing,
and Location of Meiotic DNA Breaks. *Mol Cell* **74**, 1053-1068 e1058 (2019).

1268
1269

**Figure legends**

**Fig. 1. Defective spermatogenesis in *Fignl1* cKO mice.**

(a) Scheme of mouse *Fignl1 flox* and *Δ* alleles. Yellow and blue boxes and black triangles represent untranslated regions (UTRs), coding sequence (CDS), and *loxP* sequences, respectively.

(b) Representative image of testes in $Fignl1^{+/+}$ $Stra8\text{-}Cre^+$ (left) and $Fignl1^{flox/\Delta}$ $Stra8\text{-}Cre^+$ (right) mice.

(c) Testis weights of mice with different combinations of genotypes. Adult testes (9-20 weeks) were analyzed. The red bars are means.

(d) Immunoblotting of FIGNL1 from whole-testis extracts of mice with different genotypes. $\alpha$-tubulin is a loading control.

(e) Sperm numbers in $Fignl1^{+/+}$ $Stra8\text{-}Cre^+$ (black circles) and $Fignl1^{flox/\Delta}$ $Stra8\text{-}Cre^+$ (blue open circles) mice. Sperms from adult Cauda epididymis (11-17 weeks) were counted. The red bars are means.

(f) Representative images of seminiferous tubule sections from the testis of $Fignl1^{+/+}$ $Stra8\text{-}Cre^+$ (Ctrl) and $Fignl1^{flox/\Delta}$ $Stra8\text{-}Cre^+$ (*Fignl1* cKO) mice. The sections were stained with hematoxylin and eosin (H&E). Sg, spermatogonia; Sc (L), leptotene spermatocytes; Sc (P), pachytene spermatocytes; St, spermatids.

(g) Distribution of different meiotic prophase I stages in Ctrl (gray bars) and *Fignl1* cKO (blue open bars). >100 cells were counted from each animal. The red bars are SDs from three animals of each genotype.

(h) Distribution of FIGNL1-positive and -negative cells at different meiotic prophase I stages in Ctrl (gray bars) and *Fignl1* cKO (blue open bars). Squashed testicular cells were immunostained for FIGNL1, γH2AX, and SYCP3. >100 leptotene/zygotene cells (with pan-nuclear γH2AX staining) and pachytene/diplotene cells (with the sex body-specific γH2AX staining) were analyzed from each animal. The means of two mice of each genotype are indicated. Lept., leptonema; Zyg., zygonema; Pach., pachynema; dipl., diplonema. Representative images of testicular cell squashes are shown in Extended Data Fig. 1.

(i) Representative images of spermatocyte chromosome spreads immunostained for SYCP1 (green) and SYCP3 (magenta) at zygonema in Ctrl and *Fignl1* cKO. Bottom panels are magnified images of regions with yellow dotted rectangles.

(j) Quantification of the frequency of synapsis defects in Ctrl (gray bars) and *Fignl1* cKO (blue open bars). >90 cells were analyzed from each animal. The red

bars are SDs from three mice of each genotype.

In (f)-(i), genotypes of indicated animals are Ctrl, *Fignl1$^{+/+}$ Stra8-Cre$^+$*; *Fignl1* cKO,

*Fignl1$^{flox/\Delta}$ Stra8-Cre$^+$*. The results of unpaired *t*-tests are indicated in the graphs:

****$p \leq 0.0001$.

Scale bars, 50 $\mu$m in (f) and 10 $\mu$m for whole-nucleus images and 2 $\mu$m for

magnified images in (i).

1312

**Fig. 2. Accumulation of RAD51 and DMC1 in meiotic prophase I in *Fignl1***

**cKO spermatocytes.**

(a) Representative images of spermatocyte chromosome spreads

immunostained for DMC1 (white in the top panels and green in the bottom panels)

and SYCP3 (magenta) at indicated meiotic prophase I stages in Ctrl and *Fignl1*

cKO.

(b) Quantification of DMC1 focus numbers at different meiotic prophase I stages

in Ctrl (black circles) and *Fignl1* cKO (blue open circles). The red bars are means.

(c) Representative images of spermatocyte chromosome spreads

immunostained for RAD51 (white in the top panels and green in the bottom

panels) and SYCP3 (magenta) at indicated meiotic prophase I stages in Ctrl and

*Fignl1* cKO.

(d) Quantification of RAD51 focus numbers at different meiotic prophase I stages

in Ctrl (black circles) and *Fignl1* cKO (blue open circles). The red bars are means.

(e) Quantification of focus numbers of RAD51 (green), DMC1 (magenta) and

RAD51-DMC1 colocalization (brown) at different meiotic prophase I stages in Ctrl

(filled circles) and *Fignl1* cKO (open circles). The black bars are means.

Representative images of spermatocyte chromosome spreads are shown in

Extended Data Fig. 4.

(f) Quantification of the frequency of RAD51-DMC1 colocalization at different

meiotic prophase I stages in Ctrl (filled circles) and *Fignl1* cKO (open circles). The

degrees of RAD51 foci colocalizing with DMC1 (left, green) and DMC1 foci

colocalizing with RAD51 (right, magenta) are shown. The black bars are means.

Genotypes of indicated animals are Ctrl, *Fignl1$^{+/+}$ Stra8-Cre$^+$*; *Fignl1* cKO,

*Fignl1$^{flox/\Delta}$ Stra8-Cre$^+$*. The results of the Mann-Whitney *U*-test are indicated in

the graphs: ****$p \leq 0.0001$. The total number of cells analyzed is indicated below

the graphs.

Scale bars in (a) and (c), 10 $\mu$m. Prelept., preleptonema; lept., leptonema; zyg.,

zygonema.

1342

**Fig. 3. Deregulated meiotic recombination in *Fignl1* cKO spermatocytes.**

(a) Representative images of spermatocyte chromosome spreads
immunostained for RPA2 (white in the top panels and green in the bottom panels)
and SYCP3 (magenta) at indicated meiotic prophase I stages in Ctrl and *Fignl1*
cKO.

(b) Quantification of RPA2 focus numbers at different meiotic prophase I stages
in Ctrl (black circles) and *Fignl1* cKO (blue open circles). The red bars are means.

(c) Representative images of spermatocyte chromosome spreads
immunostained for MSH4 (white in the left panels and green in the right panels),
SYCP1 (magenta) and SYCP3 (blue) at mid zygonema in Ctrl and *Fignl1* cKO.
Bottom panels are magnified images of regions with dotted yellow.

(d) Quantification of MSH4 focus numbers at mid zygonema in Ctrl (black circles)
and *Fignl1* cKO (blue open circles). The red bars are means.

(e) The correlation between MSH4 focus number and total SC length at mid
zygonema in Ctrl (black circles) and *Fignl1* cKO (blue open circles). Pearson's *r*
is shown in parentheses. A black solid line and a blue dashed line indicate linear
regression for Ctrl and *Fignl1* cKO, respectively.

(f) Quantification of MSH4 focus density on the SCs at mid-zygonema in Ctrl
(black circles) and *Fignl1* cKO (blue open circles). The red bars are means. In (e)
and (f), mid zygotene nuclei with 40-120 $\mu$m of total SC length per nucleus were
analyzed.

Genotypes of indicated animals are: Ctrl, *Fignl1$^{+/+}$ Stra8-Cre$^+$*; *Fignl1* cKO,
*Fignl1$^{flox/\Delta}$ Stra8-Cre$^+$*. The results of the Mann-Whitney *U*-test are indicated in
the graphs: ****$p \leq 0.0001$. The total number of cells analyzed is indicated below
the graphs.

Scale bars in (a) and (c), 10 $\mu$m and 2 $\mu$m for magnified images in (c).

1369

**Fig. 4. Accumulation of RAD51 and DMC1 in pre-leptotene *Fignl1* cKO**
**spermatocytes.**

(a) Representative images of spermatocyte chromosome spreads
immunostained for RAD51 (left), SYCP3 (middle) and EdU (right) at
preleptonema in Ctrl and *Fignl1* cKO.

(b) Quantification of RAD51 focus numbers at preleptonema in Ctrl (black circles)
and *Fignl1* cKO (blue open circles). The red solid bars and dashed bars are
means for total cells and RAD51-positive cells (with >10 RAD51 foci; 26/38 and

25/48 cells in early and mid/late preleptonema, respectively), respectively.

(c) Representative images of spermatocyte chromosome spreads
immunostained for DMC1 (left), SYCP3 (middle), and EdU (right) at
preleptonema in Ctrl and *Fignl1* cKO.

(d) Quantification of DMC1 focus numbers at preleptonema in Ctrl (black circles)
and *Fignl1* cKO (blue open circles). The red solid bars and dashed bars are
means for total cells and DMC1-positive cells (with >30 DMC1 foci; 28/39 and
26/45 cells in early and mid/late preleptonema, respectively), respectively.

Genotypes of indicated animals are Ctrl, *Fignl1$^{+/+}$ Stra8-Cre$^+$*; *Fignl1* cKO,
*Fignl1$^{flox/\Delta}$ Stra8-Cre$^+$*. The results of the Mann-Whitney *U*-test are indicated in
the graphs: ****$p \leq 0.0001$. The total numbers of cells analyzed are indicated
below the graphs.

Scale bars in (a) and (c), 10 $\mu$m.

1391

**Fig. 5. SPO11-independent accumulation of RAD51 and DMC1 in *Fignl1* cKO**
**spermatocytes.**

(a) Representative images of spermatocyte chromosome spreads
immunostained for RAD51 (white in the top panels and green in the bottom
panels) and SYCP3 (magenta) at leptonema in *Spo11* KO, *Fignl1* cKO and *Spo11*
KO *Fignl1* cKO.

(b) Quantification of RAD51 focus numbers at different meiotic prophase I stages
in *Spo11* KO (orange circles), *Fignl1* cKO (blue open circles), and *Spo11* KO
*Fignl1* cKO (purple open circles). The red bars are means. Cells in which Stra8-
Cre recombinase seemed to fail excision of the *Fignl1$^\Delta$* allele in *Spo11* KO *Fignl1*
cKO (<30 RAD51 foci) are excluded. Quantification including these cells is shown
in Extended Data Fig. 5a.

(c) Representative images of spermatocyte chromosome spreads
immunostained for DMC1 (white in the top panels and green in the bottom panels)
and SYCP3 (magenta) at early zygonema in *Spo11* KO, *Fignl1* cKO and *Spo11*
KO *Fignl1* cKO.

(d) Quantification of DMC1 focus numbers at different meiotic prophase I stages
in *Spo11* KO (orange circles), *Fignl1* cKO (blue open circles) and *Spo11* KO
*Fignl1* cKO (purple open circles). The red bars are means. Cells in which Stra8-
Cre recombinase seemed to fail excision of the *Fignl1$^\Delta$* allele in *Spo11* KO *Fignl1*
cKO (<60 DMC1 foci) are excluded. Quantification including these cells is shown
in Extended Data Fig. 5b.

(e) Representative images of spermatocyte chromosome spreads
immunostained for RAD51 (green), DMC1 (magenta) and SYCP3 (blue) at late
zygonema/early leptonema in *Spo11* KO *Fignl1* cKO. The bottom panels are
magnified images of regions with yellow dotted rectangles.

(f) Quantification of focus numbers of RAD51 (green open circles), DMC1
(magenta open circles), and RAD51-DMC1 colocalization (brawn open circles) at
different meiotic prophase I stages in *Spo11* KO *Fignl1* cKO. The black bars are
means.

(g) Quantification of the frequency of RAD51-DMC1 colocalization at different
meiotic prophase I stages in *Spo11* KO *Fignl1* cKO. The degrees of RAD51 foci
colocalizing with DMC1 (left, green open circles) and DMC1 foci colocalizing with
RAD51 (right, magenta open circles) are shown. The black bars are means.

Genotypes of indicated animals are: *Spo11* KO, *Spo11$^{-/-}$ Fignl1$^{flox/\Delta}$*; *Fignl1* cKO,
*Fignl1$^{flox/\Delta}$ Stra8-Cre$^+$*; *Spo11* KO *Fignl1* cKO, *Spo11$^{-/-}$ Fignl1$^{flox/\Delta}$ Stra8-Cre$^+$*. The
results of the Mann-Whitney *U*-test are indicated in the graphs: \*\*\*\**p* ≤ 0.0001.
The total number of cells analyzed is indicated below the graphs.

Scale bars in (a), (c) and (e), 10 $\mu$m and 2 $\mu$m for magnified images in (e). Prelept.,
preleptonema; lept., leptonema; zyg., zygonema.

1432

**Fig. 6. Persistent RAD51 binding at recombination sites in *Fignl1* cKO**
**spermatocytes.**

(a) RAD51 ChIP-SSDS signals (green) in Ctrl and *Fignl1* cKO with Spo11-oligo
counts (blue) and DMC1 ChIP-SSDS signals (black) in wild-type at an indicated
chromosomal region with a hotspot on mouse chromosome 2.

(b) The heatmap of RAD51 ChIP-SSDS signals (green) in Ctrl and *Fignl1* cKO
and Spo11-oligo counts (blue) and DMC1 ChIP-SSDS signals (black) in wild-type.
The 4,000 most active meiotic recombination hotspots on autosomes identified
by Spo11-oligo in wild-type are ordered by SPO11-oligos counts [112]. ChIP-SSDS
signals are centered relative to hotspot centers. In (a) and (b), Spo11-oligo and
DMC1 ChIP-SSDS data are from previous studies [112, 113].

(c) Representative images of spermatocyte chromosome spreads
immunostained for RAD51 (green), RPA2 (magenta) and SYCP3 (blue) at mid
zygonema in Ctrl and *Fignl1* cKO. Bottom panels are magnified images of regions
with yellow dotted rectangles. Scale bars, 10 $\mu$m for whole-nucleus images and
2 $\mu$m for magnified images.

(d) Quantification of focus numbers of RAD51 (green), RPA2 (magenta) and

RAD51-RPA2 colocalization (brawn) at mid zygonema in Ctrl (filled circles) and
*Fignl1* cKO (open circles). The black bars are means.
(e) Quantification of the frequency of RAD51-RPA2 colocalization at mid-
zygonema in Ctrl (black circles) and *Fignl1* cKO (blue open circles). The degree
of RPA2 foci colocalizing with RAD51 is shown. The red bars are means.
Genotypes of indicated animals are: Ctrl, *Fignl1$^{+/+}$ Stra8-Cre$^+$*; *Fignl1* cKO,
*Fignl1$^{flox/\Delta}$ Stra8-Cre$^+$*. The results of the Mann-Whitney *U*-test are indicated: ****$p$
$\leq$ 0.0001. The total numbers of cells analyzed are indicated below the graphs.
1458
**Fig. 7. *Fignl1* deficiency rescues defects in RAD51 and DMC1 assembly**
**caused by *Swsap1* deficiency.**
(a) Representative images of spermatocyte chromosome spreads
immunostained for RAD51 (white in the top panels and green in the bottom
panels) and SYCP3 (magenta) at early zygonema in Ctrl, *Swsap1* KO*, Fignl1*
cKO and *Swsap1* KO *Fignl1* cKO.
(b) Quantification of RAD51 focus numbers at different meiotic prophase I stages
in Ctrl (black circles), *Swsap1* KO (orange circles), *Fignl1* cKO (blue open circles)
and *Swsap1* KO *Fignl1* cKO (purple open circles). The red bars are means.
(c) Representative images of spermatocyte chromosome spreads
immunostained for DMC1 (white in the top panels and green in the bottom panels)
and SYCP3 (magenta) at early zygonema in Ctrl, *Swsap1* KO*, Fignl1* cKO, and
*Swsap1* KO *Fignl1* cKO.
(d) Quantification of DMC1 focus numbers at different meiotic prophase I stages
in Ctrl (black circles), *Swsap1* KO (orange circles), *Fignl1* cKO (blue open circles),
and *Swsap1* KO *Fignl1* cKO (purple open circles). The red bars are means.
Genotypes of indicated animals are *Swsap1* KO, *Swsap1$^{-/-}$ Fignl1$^{flox/\Delta}$*; *Fignl1*
cKO, *Fignl1$^{flox/\Delta}$ Stra8-Cre$^+$*; *Swsap1* KO *Fignl1* cKO, *Swsap1$^{-/-}$ Fignl1$^{flox/\Delta}$ Stra8-*
*Cre$^+$*. The results of the Mann-Whitney *U*-test are indicated in the graphs: ****$p \leq$
0.0001. The total numbers of cells analyzed are indicated below the graphs.
Scale bars in (a) and (c), 10 $\mu$m.
1480
**Fig. 8. FIGNL1 disrupts RAD51-DNA complexes.**
(a) Electrophoresis mobility shift assay showing *in vitro* disruption of RAD51-DNA
complexes by FIGNL1. A representative gel image among triplicate is shown.
Human RAD51 protein was incubated with either Cy5-labelled ssDNA (40nt) or
dsDNA (40bp) in the presence of ATP. After the formation of the protein-DNA

complex, purified human FINGL1ΔN was added at the indicated concentration
and further incubated. The complexes were analyzed by polyacrylamide gel
electrophoresis and the fluorescence from the DNAs was captured.
(b) Quantification of free ssDNA or dsDNA in (A). The red bars are means and
SDs from three independent experiments.
(c) Model of FIGNL1 function in spermatocytes. In meiotic S-phase and early
meiotic prophase I, FIGNL1 dissociates RAD51 and DMC1 from dsDNAs. At
meiotic DSB sites, FIGNL1 removes RAD51 and DMC1 from ssDNAs, which
facilitates efficient loading of MutSγ (MSH4-MSH5) and further processing of
homologous recombination to crossover outcome.
1496

**Extended Data Figures:**
**Extended Data Fig. 1. Depletion of FIGNL1 from early meiotic prophase I**
**spermatocytes, related to Fig. 1.**
Representative images of testicular cell squashes immunostained for FIGNL1
(green), γH2AX (blue) and SYCP3 (magenta) in Ctrl (*Fignl1*[+/+] *Stra8-Cre*[+]) and
*Fignl1* cKO (*Fignl1*[flox/Δ] *Stra8-Cre*[+]). SYCP3-positive prophase I spermatocytes
were categorized into two by γH2AX staining: leptotene/zygotene cells (L/Z) and
pachytene/diplotene cells (P/D) show pan-nuclear γH2AX staining and the sex
body-specific γH2AX staining, respectively. Scale bars, 10 μm.
1506

**Extended Data Fig. 2. Deregulated meiotic recombination in *Fignl1* cKO**
**spermatocytes, related to Fig. 3.**
(a) Representative images of spermatocyte chromosome spreads
immunostained for γH2AX (white in the top panels and green in the bottom
panels) and SYCP3 (magenta) at indicated meiotic prophase I stages in Ctrl and
*Fignl1* cKO.
(b) Quantification of γH2AX intensity at different meiotic prophase I stages in Ctrl
(black circles) and *Fignl1* cKO (blue open circles). Two independent experiments
showed reduced (left; pooled results from two animals of each genotype) and one
experiment showed similar (right) levels of γH2AX intensity in *Fignl1* cKO
spermatocytes compared to Ctrl. The red bars are means. The results of the
Mann-Whitney *U*-tests are indicated: ****$p \leq 0.0001$. The total numbers of cells
analyzed are indicated below the graphs.
(c) Representative images of spermatocyte chromosome spreads
immunostained for MLH1 (white in the top panels and green in the bottom panels),

SYCP1 (magenta), and SYCP3 (blue) at late pachynema in Ctrl and late
zygonema (pachynema-like) in *Fignl1* cKO. The right panels are magnified
images of regions with yellow dotted rectangles.
Genotypes of indicated animals are Ctrl, *Fignl1$^{+/+}$ Stra8-Cre$^+$*; *Fignl1* cKO,
*Fignl1$^{flox/\Delta}$ Stra8-Cre$^+$*.
Scale bars in (a) and (c), 10 $\mu$m and 2 $\mu$m for magnified images in (c).
1528
**Extended Data Fig. 3. Infrequent localization of RAD51 at ongoing**
**replication fork in preleptotene *Fignl1* spermatocytes, related to Fig. 4.**
(a) Representative images of spermatocyte chromosome spreads
immunostained for IdU (green), RAD51 (magenta), and SYCP3 at early
preleptonema in *Fignl1* cKO. The bottom panels are magnified images of regions
with yellow dotted rectangles. Scale bars, 10 µm for whole-nucleus images and
2 $\mu$m for magnified images.
(b) Quantification of focus numbers of IdU (green open circles), RAD51 (magenta
open circles), and IdU-RAD51 colocalization (brawn open circles) at early
preleptonema in *Fignl1* cKO. The black bars are means.
(c) Quantification of the frequency of IdU-RAD51 colocalization at early
preleptonema in *Fignl1* cKO. The degrees of IdU foci colocalizing with RAD51
(left, green open circles) and RAD51 foci colocalizing with IdU (right, magenta
open circles) are shown. The black bars are means. The colocalization frequency
was measured with and without rotating the IdU image 90º clockwise relative to
the RAD51 image. The results of the Mann-Whitney *U*-tests are indicated.
The Genotype of *Fignl1* cKO animals is *Fignl1$^{flox/\Delta}$ Stra8-Cre$^+$*.
1546
**Extended Data Fig. 4. Uncoupled loading of RAD51 and DMC1 in *Fignl1* cKO**
**spermatocytes, related to Fig. 2 and 4.**
Representative images of spermatocyte chromosome spreads immunostained
for RAD51 (green), DMC1 (magenta), and SYCP3 (blue) at indicated meiotic
prophase I stages in Ctrl (*Fignl1$^{+/+}$ Stra8-Cre$^+$*) and *Fignl1* cKO (*Fignl1$^{flox/\Delta}$ Stra8-*
*Cre$^+$*). The bottom panels are magnified images of regions with yellow dotted
rectangles. Scale bars, 10 µm for whole-nucleus images and 2 $\mu$m for magnified
images.
1555
**Extended Data Fig. 5. Accumulation of RAD51 and DMC1 with low levels of**
**γH2AX in *Spo11* KO *Fignl1* cKO spermatocytes, related to Fig. 5.**

(a) Quantification of RAD51 focus numbers at different meiotic prophase I stages
in *Spo11* KO (orange circles), *Fignl1* cKO (blue open circles), and *Spo11* KO
*Fignl1* cKO (purple open circles). The red bars are means.
(b) Quantification of DMC1 focus numbers at different meiotic prophase I stages
in *Spo11* KO (orange circles), *Fignl1* cKO (blue open circles), and *Spo11* KO
*Fignl1* cKO (purple open circles). The red bars are means.
(c) Representative images of spermatocyte chromosome spreads
immunostained for γH2AX (green), SYCP3 (magenta), and RAD51 (white in
insets) at indicated meiotic prophase I stages in *Fignl1* cKO, *Spo11* KO and
*Spo11* KO *Fignl1* cKO. Scale bars, 10 μm.
(d) Quantification of γH2AX intensity at different meiotic prophase I stages in
*Fignl1* cKO (blue open circles), *Spo11* KO (orange circles), and *Spo11* KO *Fignl1*
cKO (purple open circles). The red bars are means.
Genotypes of indicated animals are: *Spo11* KO, *Spo11$^{-/-}$ Fignl1$^{flox/\Delta}$*; *Fignl1* cKO,
*Fignl1$^{flox/\Delta}$ Stra8-Cre$^+$*; *Spo11* KO *Fignl1* cKO, *Spo11$^{-/-}$ Fignl1$^{flox/\Delta}$ Stra8-Cre$^+$*. The
results of the Mann-Whitney *U*-test are indicated in the graphs: ****$p \leq 0.0001$.
The total number of cells analyzed is indicated below the graphs.
1575
**Extended Data Fig. 6. RAD51 binding at recombination sites in *Fignl1* cKO**
**spermatocytes, related to Fig. 6.**
(a) RAD51 ChIP-SSDS signals (green) in Ctrl and *Fignl1* cKO with Spo11-oligo
counts (blue) and DMC1 ChIP-SSDS signals (black) in wild-type at an indicated
chromosomal region with multiple hotspots on mouse chromosome 3.
(b) The heatmap of RAD51 ChIP-SSDS signals (green) in Ctrl and *Fignl1* cKO
and Spo11-oligo counts (blue) and DMC1 ChIP-SSDS signals (black) in wild-type,
as shown in Fig. 6b. Three mice of each genotype were analyzed and two RAD51
antibodies with different lot numbers were used for ChIP. The results of Ctrl #1
and *Fignl1* cKO #1 are shown in Fig. 6a, b, Extended Data Fig. 6a.
(c) Comparison of RAD51 ChIP-SSDS signals at meiotic recombination hotspots.
Each spot represents the sum of RAD51 ChIP-SSDS signals around ±2,000 bp
of one of the 4,000 most active meiotic recombination hotspots on autosomes
(black) used for heatmap representation in Fig. 6b, Extended Data Fig. 6a and
163 active hotspots on sex chromosomes (magenta) with similar Spo11-oligo
counts as the 4,000 active autosomal hotspots in wild-type [112]. Pearson's *r* for
autosomal hotspots is also shown.
(d) Metaplots of RAD51 ChIP-SSDS signals around meiotic recombination

hotspots. Smoothed RAD51 ChIP-SSDS signals in Ctrl (top) and *Fignl1* cKO

(bottom) around ±2,000 bp of the 4,000 most active meiotic recombination

hotspots on autosomes used for heatmap representation are shown. Three mice

of each genotype were analyzed. Black, orange and blue lines represent the

results of #1, #2 and #3, respectively.

Genotypes of indicated animals are: Ctrl, *Fignl1$^{+/+}$ Stra8-Cre$^+$*; *Fignl1* cKO,

*Fignl1$^{flox/\Delta}$ Stra8-Cre$^+$*.

1601

1602     **Source Data**

1603     Uncropped images of immunoblotting in Fig. 1d and an unprocessed image of

1604     electrophoresis mobility shift assay in Fig. 8a are provided.

1605

1606    **Supplementary information**

1607

1608    **Supplementary Table 1. Primers used for genotyping in this study**

| Primer | Sequence (5'->3') |
|--------|-------------------|
| *Fignl1* wild-type and flox allele, forward | TATCCATATTGGGTGTCTCCTGTTAC |
| *Fignl1* wild-type and flox allele, reverse | TAGCACACACACACTGGATATAACT |
| *Fignl1* deletion allele, forward | CCGAGTATCCATATTGGGTG |
| *Fignl1* deletion allele, reverse | TGTCTGCTGTAGTCATGAGG |
| *Stra8-Cre* transgenic allele, forward | CTCCAAGGGGGTAAGGTGTAGC |
| *Stra8-Cre* transgenic allele, reverse | TTCCAGGGCGCGAGTTGATAG |
| *Swsap1* wild-type allele, forward | GAGATATCGCTTTTGATTGGTC |
| *Swsap1* wild-type allele, reverse | GAATTGAGGATGGGTACTGG |
| *Swsap1* deletion allele, forward | CGGTTTCCATATGGGGATTGGTGGCGACGA |
| *Swsap1* deletion allele, reverse | TGCCTTATTTCCTGATCCAGGCTAGCTGTC |
| *Spo11* wild-type allele, forward | TGAGATACATGGAGGAAGATGG |
| *Spo11* deletion allele, forward | CTGAGCCCAGAAAGCGAAGGAA |
| *Spo11,* common reverse | ATGTTAGTCGGCACAGCAGTAG |

1623

1624    **Supplementary Table 2. Antibodies used in this study**

| Antibody | Source | Identifier | Application | Dilution |
|----------|--------|-----------|-------------|----------|
| Rat anti-SYCP3 (Serum) | this study | N/A | IF | 1:5,000 |
| Mouse anti-SYCP3 | Abcam | ab 97672 | IF | 1:750 |
| Rabbit anti-SYCP1 | Novus Biologicals | NB300-229 | IF | 1:500 |
| Mouse anti-MLH1 | BD Pharmingen | 51-1327QR | IF | 1:50 |
| Rabbit anti-MSH4 | Abcam | ab58666 | IF | 1:200 |
| Rabbit anti-RPA2 | Abcam | ab76420 | IF | 1:200 |
| Rat anti-RPA2 | Cell Signaling | 2208 | IF | 1:100 |
| Rabbit anti-DMC1 | Santa Cruz | sc-22768 | IF | 1:200 |
| Rabbit anti-RAD51 | Santa Cruz | sc-8349 | IF | 1:100 |
| Mouse anti-RAD51 | Novus Biologicals | NB100-148 | IF, ChIP | 1:100 |
| Mouse anti-BrdU | BD Biosciences | BD347580 | IF | 1:200 |
| Rabbit anti-FIGNL1 | Proteintech | 17604-1-AP | IF | 1:100 |
| | | | IB | 1:1,000 |
| Mouse anti-α tubulin | SIGMA | T6074 | IB | 1:3,000 |
| Goat anti-mouse 488 | Molecular Probes | A-11029 | IF | 1:1,000 |
| Goat anti-rabbit 488 | Molecular Probes | A-11034 | IF | 1:1,000 |

| 1642 | Goat anti-rat 488 | Molecular Probes A-11006 | IF | 1:1,000 |
|---|---|---|---|---|
| 1643 | Goat anti-mouse 594 | Molecular Probes A-11032 | IF | 1:1,000 |
| 1644 | Goat anti-rat 594 | Molecular Probes A-11007 | IF | 1:1,000 |
| 1645 | Goat anti-rabbit 647 | Molecular Probes A-21245 | IF | 1:500 |
| 1646 | Goat anti-rabbit IgG-HRP | Southern Biotech 4030-05 | IB | 1:10,000 |
| 1647 | Goat anti-mouse IgG-HRP | Thermo Scientific 31430 | IB | 1:10.000 |

1648    IF, immunofluorescence; IB, immunoblotting.; ChIP, chromatin immunoprecipitation

1649

Ito et al., Figure 1

**a** *Fignl1*
flox — Exon 1 — Exon 2 — Exon 3
Δ

**b** *Fignl1* +/+ flox/Δ
*Stra8-Cre* + +
5 mm

**c**
Testis weight (as % body weight)
n= 6 6 7 7
*Fignl1* +/+ flox/Δ +/+ flox/Δ
*Stra8-Cre* - +

**d** *Fignl1* +/+ flox/Δ flox/Δ
*Stra8-Cre* + - +
75 — FIGNL1
50 — α-tubulin

**e**
Sperm per mouse (x 10⁶)
p= 0.0008
n= 3 3
*Fignl1* +/+ flox/Δ
*Stra8-Cre* +

**f** *Stra8-Cre⁺ Fignl1⁺/⁺* (Ctrl)   *Stra8-Cre⁺ Fignl1^flox/Δ* (*Fignl1* cKO)
Sg
St
Sc (L)
Sc (P)
Sg
Sc (L)
St

**g**
Percent of prophase cells
■ Ctrl □ *Fignl1* cKO
Leptonema Zygonema Pachynema Diplonema

**h**
Percent of FIGNL1-positive cells
■ Ctrl □ *Fignl1* cKO
Lept./zyg. Pach./dipl.

**i** Ctrl   *Fignl1* cKO
Zygonema
SYCP1
SYCP3
ii i
i Non-homologous synapsis
ii Multi-chromosomal synapsis

**j**
Percent of zygotene cells with defective synapsis
■ Ctrl □ *Fignl1* cKO
**** p= 0.0214
Non-homologous Multi-chromosomal

**a**

**c** Eary preleptonema

**b**

**d**

**c** Mid zygonema

**a**

**b**

**a**

**b**

**c**

**a** Early preleptonema

*Fignl1* cKO

SYCP3 | IdU | RAD51 | IdU RAD51

**b**

Foci per nucleus

n= 38  38  38

IdU  RAD51  Colocalization

Early preleptonema

**c**

Percent of IdU foci colocalization with RAD51

*p*= 0.0261

n= 21  21

Early prelept.  Early prelept. 90° turn

Percent of RAD51 foci colocalization with IdU

*p*= 0.0439

n= 21  21

Early prelept.  Early prelept. 90° turn

Ito et al., Extended Data Figure 4

**a**

**b**

**c**

(a) An uncropped immunoblotting image for FIGNL1 shown in Fig. 1d.
(b) An uncropped immunoblotting image for α-tubulin shown in Fig. 1d.
(c) An unprocessed image of electrophoresis mobility shift assay shown in Fig. 8a.