FIGNL1 AAA+ ATPase remodels RAD51 and DMC1 filaments in pre-meiotic DNA replication and meiotic recombination

Masaru Ito[1]*, Asako Furukohri[1], Kenichiro Matsuzaki[1, 2], Yurika Fujita[1], Atsushi Toyoda[3], and Akira Shinohara[1]*

[1]Institute for Protein Research, Osaka University, Suita, Osaka 565-0871, Japan

[2]Department of Advanced Bioscience, Graduate School of Agriculture, Kindai University, Nara, Nara 631-8505, Japan

[3]Advanced Genomics Center, National Institute of Genetics, Mishima, Shizuoka 411-8540, Japan

Running title: Suppression of RAD51/DMC1 assembly by FIGNL1

Keywords: meiosis; homologous recombination; FIGNL1; RAD51; DMC1

*Corresponding author:

Masaru Ito; msrito2@protein.osaka-u.ac.jp

Akira Shinohara; ashino@protein.osaka-u.ac.jp

Lead contact:

Akira Shinohara

Institute for Protein Research, Osaka University,

3-2 Yamadaoka, Suita, Osaka 565-0871 JAPAN

Phone: 81-6-6879-8624

FAX: 81-6-6879-8626

E-mail: ashino@protein.osaka-u.ac.jp

ORCID: 0000-0003-4207-8247

Abstract

The formation of RAD51/DMC1 filaments on single-stranded (ss)DNAs essential

for homology search and strand exchange in DNA double-strand break (DSB)

repair is tightly regulated. FIGNL1 AAA+++ ATPase controls RAD51-mediated

recombination in human cells. However, its role in gametogenesis remains

unsolved. Here, we characterized a germ-line-specific conditional knockout

(cKO) mouse of *FIGNL1*. *Fignl1* cKO male mice showed defective chromosome

synapsis and impaired meiotic DSB repair with the accumulation of

RAD51/DMC1 on meiotic chromosomes, supporting a positive role of FIGNL1 in

homologous recombination at a post-assembly stage of RAD51/DMC1 filaments.

Fignl1 cKO spermatocytes also accumulate RAD51/DMC1 on chromosomes in

pre-meiotic S-phase. These RAD51/DMC1 assemblies are independent of

meiotic DSB formation. We also showed that purified FIGNL1 dismantles RAD51

filament on double-stranded (ds)DNA as well as ssDNA. These results suggest

an additional role of FIGNL1 in limiting the non-productive assembly of

RAD51/DMC1 on native dsDNAs during pre-meiotic S-phase and meiotic

prophase I. (149 words)

Introduction

RAD51 is a central player in homologous recombination in eukaryotes by catalyzing homology-directed repair such as the repair of DNA double-strand breaks (DSBs) in both somatic and meiotic cells. RAD51 also plays a role in S-phase for the protection of stalled DNA replication forks [1, 2], which is independent of RAD51-mediated strand exchange [3, 4]. As an active form for homology search and strand exchange, multiple protomers of RAD51 bind to single-stranded (ss)DNAs in the presence of ATP to form a right-handed helical filament termed RAD51 nucleofilament, in which ssDNAs are extended 1.5-2.0-fold relative to the B-form DNA [5-8]. RAD51 nucleofilament is highly dynamic; its assembly and disassembly are controlled by multiple proteins/complexes. The RAD51 mediator promotes the assembly of RAD51 nucleofilament on the ssDNAs coated with an ssDNA-binding protein, RPA (Replication protein A), which hampers RAD51 binding to the ssDNA. The RAD51 mediator in humans includes BRCA2(-PALB2-DSS1) [9], BRCA1-BARD1 [10], RAD52 [11-13], SWI5-SFR1/MEI5 [14, 15], and five RAD51 paralogs (RAD51B, -C, -D, XRCC2, and -3) in vertebrates including humans. The RAD51 paralogs form two distinct complexes: RAD51C-XRCC3 and RAD51B-RAD51C-RAD51D-XRCC2 [16]. SWSAP1 is a distinct member of a RAD51 paralog [17-19], which forms a complex with SWS1 and SPIDR. At the post-assembly stage, RAD51 nucleofilament works together with the SWI2/SNF2 family, RAD54 and RAD54B [20, 21], and RAD51AP1 [22-24] for efficient homology search and strand exchange with a target double-stranded DNA (dsDNA) for the formation of displacement (D)-loop. RAD51-mediated strand invasion provides a template for DNA synthesis for further processing of the intermediates for the completion of the repair.

The dynamics of the RAD51 filament are also negatively regulated by proteins that disassemble the filament. Various DNA helicases are involved in the disassembly of RAD51 filament, which include BLM [25-27], FANCJ [28], RECQ5 [29, 30], and FBH1 [31, 32] as well as a degenerate DNA helicase, PARI [33]. RAD51-mediated invasion intermediates are also remodeled by other DNA helicases such as WRN, RTEL1, FANCM, and RECQ1 [34]. RAD54 translocase also disassembles RAD51 from dsDNA in recombination intermediates [35]. The stabilization and destabilization of RAD51 filaments are associated with the choice of a homologous recombination pathway [36].

Meiosis is a specialized cell division that produces haploid gametes. Homologous recombination, specifically crossing-over, is essential for

chromosome segregation of homologous chromosomes during meiosis I. To

ensure the crossover (CO) formation in meiosis, highly programmed regulation is

implanted on the recombination pathway [37-39]. A meiosis-specific RAD51 cousin,

DMC1, is critical for homology search in meiotic recombination in many species

[40]. The cooperation of RAD51 and DMC1 promotes homology search between

homologous chromosomes rather than sister chromatids whose mechanism is

one big enigma in the field [41, 42]. RAD51/DMC1-mediated recombination requires

meiosis-specific proteins as well as proteins playing a role in somatic cells.

Knockout (KO) of the most of genes necessary for homologous recombination,

Rad51 and RAD51 mediators (*Brca2*, *Rad51B*, *-C*, *-D*, *Xrcc2*, and *-3*), in mice

leads to embryonic lethality [43-49]. *Swsap1* (and *Sws1* and *Spidr*) KO mice are

viable but are sterile, with an exception of *Spidr* KO female mice being sub-fertile

[17, 50, 51]. Indeed, SWSAP1-SWS1-SPIDR is necessary for the efficient formation

of RAD51/DMC1 focus, which is a cytologically detectable immuno-stained

structure, in male meiosis [17, 50-52]. Like RAD51 mediators, KO mice of some

negative regulators such as BLM are also embryonically lethal [53] while *Rad54* KO

mice are viable and fertile [54].

A recently identified recombination regulator, FIGNL1, encodes an AAA+

ATPase which belongs to microtubule severing proteins including Katanin,

Spastin, and Fidgetin with an N-terminal microtubule-binding domain [55]. FIGNL1

was identified as a RAD51-interacting protein and is required for homologous

recombination in human somatic cells [56]. FIGNL1 knockdown does not affect

DSB-induced RAD51 focus formation [51, 56], suggesting its role in the post-RAD51

assembly step. Despite its positive role in homologous recombination, FIGNL1

functions as a negative regulator for RAD51 assembly. FIGNL1 depletion rescues

defective RAD51-focus formation in human cells depleted for SWSAP1 or SWS1,

but not for RAD51C [51]. Moreover, purified FIGNL1 promotes the dissociation of

RAD51 protein from ssDNAs, which is suppressed by SWSAP1 [51]. In *Arabidopsis*

thaliana meiosis, a mutant of *FIGL1*, FIGNL1 ortholog, as well as its partner, *FLIP*,

shows increased frequencies of COs during meiosis, indicating a role as an anti-

CO factor [57, 58]. The role of FIGL1-FLIP as the anti-CO factor is also seen in rice

meiosis [59, 60]. Importantly, the *figl1* mutation can suppress defective

RAD51/DMC1 focus formation in *brca2* mutant meiocytes, indicating antagonism

of FIGL with the RAD51 mediator [61]. These suggest that RAD51 mediators might

play a distinct function in different recombination pathways. Although FIGNL1

plays a role in homologous recombination by regulating the SWSAP1 complex in

somatic cells [51], it remains unknown how FIGNL1 regulates recombination and
works together with SWSAP1 during meiosis.

In this study, we analyzed the role of FIGNL1 in mice in meiosis, by
constructing a germ-cell specific conditional knockout (cKO) mouse and found
that *Fignl1* depletion in the testis led to defective spermatogenesis. Meiotic
chromosome spreads from *Fignl1* cKO spermatocytes showed a two-fold
increase in the numbers of RAD51 and DMC1 foci relative to controls, suggesting
the role of FIGNL1 in the post-assembly stage of not only RAD51 filament but
also DMC1 filament. This defective disassembly results in the impaired loading
of proteins necessary for CO formation. We also found that FIGNL1 suppresses
RAD51 and DMC1 assembly in pre-meiotic S-phase and early leptonema, which
are independent of meiotic DSB formation. Moreover, *Fignl1* cKO ameliorates
defective RAD51- and DMC1-focus formation in KO spermatocytes of *Swasp1,* a
RAD51/DMC1 mediator, supporting the antagonistic role of FIGNL1 to SWSAP1
in the assembly of both RAD51 and DMC1. Together, our study establishes a
dual role of FIGNL1 in mammalian male meiosis. The same conclusion was
obtained by Baudat's group by characterizing the *Fignl1* cKO as well as *Firrm*
cKO (FIRRM is a FIGNL1-interacting protein, accompanying paper).

Results

***Fignl1* is essential for early mouse embryogenesis**

To determine the functions of FIGNL1 in mice, we created a conditional allele of
Fignl1 gene, *Fignl1^{flox}*, in which the exon 3 encoding an open reading frame of
the full-length protein is flanked by the *loxP* sites (Fig. 1a) in C57BL/6 background.
A null allele of the gene, *Fignl1^Δ*, was obtained by breeding with *CAG-Cre+* mice
that constitutively express Cre recombinase. *Fignl1^{Δ/+}* heterozygous mice grew
normally and did not show any apparent defects. Breeding between male and
female of *Fignl1^{Δ/+}* produced *Fignl1^{+/+}* and *Fignl1^{Δ/+}* pups, but no *Fignl1^{Δ/Δ}*
homozygous pups (0/186 from 38 litters), showing that *Fignl1^{Δ/Δ}* mice are not
viable. This is consistent with the reported lethality of *Fignl1* knockout (KO) mice
in the International Mouse Phenotyping Consortium (IMPC)
(https://www.mousephenotype.org). In a detailed analysis of different embryonic
stages by the breeding of heterozygous parents, we could not obtain *Fignl1^{Δ/Δ}*
homozygous embryo at both E7.5 and E13.5 stages (0/29 and 0/37, respectively),
showing very early lethality of mice homozygous for *Fignl1* deletion. Early
embryonic lethality in mice homozygous for the homologous recombination

genes has been reported; *Rad51* [49], *Brca2*, [62] and Rad51 paralogs (*Rad51B*, *-C*,
-D, *Xrcc2*, and *-3*) [43-48].
159
Fignl1 conditional knockout mice are defective in spermatogenesis

To know the role of FIGNL1 in male meiosis, we generated *Fignl1$^{flox/\Delta}$* mice
harboring a germ-line specific Cre recombinase gene, *Stra8-Cre*. The *Stra8-Cre*
gene ensured the expression of Cre not only in spermatocytes (meiotic cells) but
also in spermatogonia [63]. *Fignl1$^{flox/\Delta}$ Stra8-Cre$^+$,* hereafter *Fignl1* conditional
knockout *(*cKO), male mice were characterized for spermatogenesis. *Fignl1* cKO
adult male mice showed normal body weight without any morphological defects
but reduced sizes of the testis by ~50% relative to controls (*Fignl1$^{+/+}$, Fignl1$^{flox/\Delta}$,*
and *Fignl1$^{+/+}$ Stra8-Cre$^+$*), which is a characteristic of mutants with defective
spermatogenesis (Fig. 1b, c). Western blotting analysis confirmed decreased
amounts of FIGNL1 protein in whole testis extracts (Fig. 1d). The level of residual
FIGNL1 protein is consistent with the efficiency of Stra8-Cre-mediated excision
of *Fignl1^{flox}* allele estimated by the transmission of *Fignl1$^{\Delta}$* allele from *Fignl1$^{flox/+}$*
Stra8-Cre$^+$ male mice (91.3%; 116/127). Sperm counts were ~6-fold lower in
Fignl1 cKO relative to controls (Fig. 1e). Residual sperms in *Fignl1* cKO might
come from *Fignl1$^{flox/\Delta}$* cells where Stra8-Cre recombinase failed excision of the
Fignl1^{flox} allele. Histological analysis showed that seminiferous tubules in *Fignl1*
cKO testes contained decreased numbers of late spermatocytes and spermatids
(Fig. 1f). These indicate a critical role of FIGNL1 in spermatogenesis. Consistent
with this, FIGNL1 expression was detected in PLZF-positive spermatogonia and
SYCP3-positive spermatocytes, but not in Sertoli cells and round spermatids
(Supplementary Fig. 1a).
To analyze meiotic defects in *Fignl1* cKO male mice, we prepared surface-
spread chromosomes of spermatocytes and immuno-stained for meiosis-specific
chromosomal proteins, SYCP3(Scp3) and SYCP1(Scp1), which are components
of lateral/axial elements and central region of the synaptonemal complex (SC),
respectively [64-66]. In leptonema of control spermatocytes, SYCP3 shows thin-line
staining structures that correspond to elongating chromosome axes (from short
to long lines). In zygonema, regions of two thin chromosome axes are paired
locally to form a thick line of SYCP3, where SYCP1 proteins are deposited for the
assembly of the SC central region. From early zygonema to early pachynema,
SC elongates along entire chromosomes, resulting in pachytene chromosome
configuration with 19 fully-paired thick SYCP3 lines colocalized with SYCP1 and

partially-paired X-Y bivalent. In *Fignl1* cKO spermatocytes, like in controls,
SYCP3-positive short and long lines were formed in leptonema, and short lines
of SYCP1 in zygonema were also observed. However, there were fewer
pachytene and diplotene cells in *Fignl1* cKO relative to controls (22.6±2.4% and
11.4±6.2% compared to 44.7±11.0% and 24.3±7.7%, respectively; Fig. 1g). We
also evaluated the expression of FIGNL1 protein in testicular cell squashes by
categorizing SYCP3-positive meiotic cells into two: cells before pachynema (i.e.
leptonema and zygonema) with a pan-nuclear γH2AX staining and those after
pachynema (i.e. pachynema and diplonema) with a dense γH2AX staining on the
X-Y pair (the sex body) [67] (Supplementary Fig. 1b). In controls, cells both before
and after pachynema were 100% positive for FIGNL1 staining. In contrast, while
only ~25% of cells prior to pachynema were positive for FIGNL1, cells after
pachynema were nearly 100% positive in *Fignl1* cKO (Fig. 1h). This indicates that
Fignl1$^{Δ/Δ}$ cells are fully lost before pachynema from *Fignl1* cKO testes and only
Fignl1$^{flox/Δ}$ cells where Stra8-Cre recombinase failed excision of the *Fignl1*^{flox}
allele progress further to complete spermatogenesis. Further analysis of surface-
spread chromosomes revealed abnormal SC formation such as synapsis
between non-homologous chromosomes (67.7±2.0% of cells compared to
13.6±4.1% in controls) and multivalent synapsis (26.8±10.9 % of cells compared
to 3.1±2.7% in controls) in *Fignl1* cKO (Fig. 1i, j), which could be defined as
"pachytene-like" stage as seen in other meiotic recombination defective mutants
such as *Dmc1* and *Meiob* KO, etc. [68-70]. These results show that FIGNL1 is
necessary for full synapsis of homologous chromosomes in late zygonema and/or
early pachynema and that most of the FIGNL1-depleted cells seem to die at these
stages or do not progress further.
218
Compromised meiotic recombination in *Fignl1* cKO
ssDNA-binding proteins such as RPA and MEIOB-SPATA22 play a role in the
early/middle stage of meiotic recombination both before and after the RAD51 and
DMC1 assembly. To know the effect of *Fignl1* cKO on meiotic recombination, we
examined the localization of RPA2, a middle-subunit of RPA, on chromosome
spreads of spermatocytes, together with SYCP3 as a marker for the progression
of meiotic prophase I. In control spermatocytes, RPA2 foci appear from
leptonema with 200-300 foci on average in mid/late leptonema (223±73 foci; Fig.
2a, b). The number of RPA2 foci increased up to ~400 foci in zygonema (376±45
foci in mid-zygonema) and persisted throughout zygonema (332±59 foci in late

zygonema). The RPA2-focus number in *Fignl1* cKO mid/late leptonema was
increased up to 300-400 foci (368±100 foci) and a similar number persisted in the
early zygotene stage (409±59 foci). The number of RPA2 foci in *Fignl1* cKO
gradually decreased from mid-zygonema and was significantly lower than
controls in late zygonema (281±48 foci compared to 332±59 in controls; *p*<0.0001,
Mann-Whitney *U*-test; Fig. 2b), suggesting a role of FIGNL1 in the dynamic
behavior of an RPA-associated recombination event. We also analyzed the
histone γH2AX staining as a marker of DSB formation and its repair. Interestingly,
Fignl1 cKO spermatocytes with increased RPA2 foci showed a similar or slightly
reduced intensity of γH2AX from leptonema to the late zygonema stage
compared to controls (Supplementary Fig. 2a, b).

In meiotic recombination, previous cytological analyses showed that RPA2
foci persist longer than DMC1/RAD51 foci from mid/late zygonema to early/mid
pachynema, suggesting a role of RPA in a late stage of meiotic recombination [71].
In zygonema (from early zygonema), foci containing the MutSγ complex with
MSH4 and MSH5, which regulate recombination events for CO formation, appear
specifically on synapsed regions and show similar behavior to RPA on meiotic
chromosomes [72]. We analyzed MSH4 focus formation and found that the number
of MSH4 foci in *Fignl1* cKO was lower than in controls in mid/late zygonema
(50±22 foci versus 107±47 foci in controls; Fig. 2c, d). Given the possibility that
the reduction in MSH4-focus number is an indirect consequence of synapsis
defects in *Fignl1* cKO, we also analyzed the density of MSH4 foci per unit length
of SC. MSH4-focus number and SC length (total SYCP1 length) showed a linear
relationship in control mid-zygonema (*r*=0.89; Fig. 2e). In *Fignl1* cKO, a
correlation between MSH4-focus number and SC length was also positive, but
lower than in controls (*r*=0.58). Importantly, the density of MSH4 foci per unit
length of SC is reduced in *Fignl1* cKO relative to controls (0.73±0.22 foci per μm
of SC versus 0.98±0.18 foci per μm of SC in controls; Fig. 2f). These suggest that
FIGNL1 is required for efficient loading of MSH4 for meiotic recombination.

CO formation is a key outcome of meiotic recombination. The CO sites on
meiotic chromosomes are marked with the foci of MutLγ complex such as MLH1
[73]. Control late pachytene cells show ~25 MLH1 foci per nucleus. On the other
hand, we could not detect any MLH1 foci on late zygotene/pachytene-like
chromosomes of *Fignl1* cKO spermatocytes (Supplementary Fig. 2c). This is
consistent with the observation that FIGNL1 is essential for the progression of
spermatocytes into a pachytene stage with MLH1 foci.

FIGNL1 promotes the disassembly of DMC1 in male meiosis

We next examined the localization of DMC1 on meiotic chromosomes. As shown

previously [74-76], in control spermatocytes, DMC1 foci appear in leptonema, peak

in their number at early zygonema, and gradually decrease throughout zygonema

(Fig. 3a, b). In *Fignl1* cKO, DMC1 foci followed similar kinetics as seen in controls.

However, at any stages, the number of DMC1 foci in *Fignl1* cKO was more than

that in controls. In early zygonema, ~500 DMC1 foci (449±89 foci) are observed

in *Fignl1* cKO, which is 2.3-fold more than in controls (194±32 foci). DMC1-focus

number in *Fignl1* cKO clearly increased through meiotic prophase I progression.

In late zygonema, the number of DMC1 foci in *Fignl1* cKO was reduced by half

of that in early zygonema (220±70 foci compared to 449±89 foci), indicating a

turnover of the foci. These results suggest that FIGNL1 suppresses excess

DMC1-focus formation and/or timely progression of DMC1-mediated

recombination.

In *Fignl1* cKO spermatocytes, we always observed two populations: cells

with a high number of DMC1 foci and cells with a similar number of foci to controls.

We assumed that the latter is *Fignl1$^{flox/\Delta}$* cells where Cre recombinase failed

excision of the *Fignl1^{flox}* allele into *Fignl1$^{\Delta}$*. Note that we included these foci for

our quantification and that this population becomes an internal control for our

analysis.

FIGNL1 promotes the disassembly of RAD51 in spermatocytes

In human cultured cells, FIGNL1 regulates RAD51 assembly [51, 56]. We examined

the localization of RAD51 on meiotic chromosomes. As reported previously [74-76],

in control spermatocytes, RAD51 foci are observed from mid/late leptonema and

peak in early zygonema, and then decrease toward pachynema (Fig. 3c, d). The

average number of RAD51 foci is 107±56 and 181±27 foci per nucleus in control

mid/late leptotene and early zygotene stages, respectively. As seen for DMC1,

we detected an increased number of RAD51 foci in *Fignl1* cKO cells. The average

number of RAD51 foci in early zygonema is 380±98 foci, which is more than twice

compared to that of controls (181±27 foci). The number of RAD51 foci in

zygonema is similar to that of DMC1 foci, with ~600 RAD51 foci in some early

zygotene nuclei. When the zygotene stage progressed from early to late, the

RAD51-focus number in *Fignl1* cKO gradually decreased as in controls. However,

even at late zygonema, we detected a significant increase in RAD51-focus

number in *Fignl1* cKO relative to controls (182±86 foci compared to 128±32 foci
in controls; *p*=0.0163, Mann-Whitney *U*-test; Fig. 2d). These results show that
FIGNL1 promotes the timely resolution of RAD51 ensembles during the zygotene
stage.
305   We note that the focus intensities of RAD51 and DMC1 in *Fignl1* cKO
306   spermatocytes were slightly decreased compared to controls (Supplementary Fig.
307   3), suggesting that in *Fignl1* cKO spermatocytes, more sites for the accumulation
308   of RAD51/DMC1 molecules are present, fewer molecules of RAD51/DMC1
309   protein are on each accumulation site because of a limited number of
310   RAD51/DMC1 protein in a cell.
311

Increased RAD51/DMC1 foci in early leptonema in *Fignl1* cKO
In contrast to the zygotene stage where the numbers of DMC1 and RAD51 foci
are comparable in both controls and *Fignl1* cKO, focus numbers of DMC1 and
RAD51 were clearly different at early leptonema only in *Fignl1* cKO (Fig. 3a-d).
In mid/late leptonema when DMC1 and RAD51 foci appear on elongating SYCP3
stretches in controls, the number of DMC1 foci in *Fignl1* cKO is 369±156 foci per
cell, which is about 3-4 times more of that in controls (96±57 foci). The number
of RAD51 foci in mid/late leptonema in *Fignl1* cKO is 387±134 foci, 3-4 times
more than in controls (107±56 foci). In early leptonema, where SYCP3 short
stretches are yet to be elongated, DMC1 and RAD51 foci rarely appear in control
spermatocytes. However, *Fignl1* cKO formed a relatively larger number of RAD51
foci than DMC1: the number of RAD51 foci was 355±178, which exceeds the
number of DMC1 foci (194±148 foci) in early leptonema. These suggest that
FIGNL1 suppresses the formation of excess numbers of RAD51 and DMC1 foci
in early meiosis I, and that, in the absence of FIGNL1, RAD51 and DMC1 loading
is uncoupled (see below).
328

Pre-meiotic S-phase in *Fignl1* cKO accumulates RAD51/DMC1 foci
In wild-type meiosis, the focus formation of both RAD51 and DMC1 is mediated
by SPO11-induced DSBs in leptonema [77, 78]. The increased number of RAD51
and DMC1 foci on the early leptotene chromosome of *Fignl1* cKO spermatocytes
led us to analyze preleptotene (pre-meiotic S-phase) spermatocytes in order to
characterize the origin of excess RAD51/DMC1 foci that form in the absence of
FIGNL1. For *in vitro* labeling of DNA replication, we incubated testis cell
suspension with EdU for 1 hour in a culture medium. The preleptotene

spermatocytes, where a limited number of SYCP3 short stretches forms along
with a numerous number of "tiny" SYCP3 dots, were categorized into two classes:
pan dotty staining of EdU in early preleptonema and clustered dotty staining of
EdU in mid/late preleptonema. In controls, both early and mid/late preleptotene
EdU-positive spermatocytes showed very few foci of RAD51 (Fig. 4a, b). On the
other hand, we could detect RAD51 foci in EdU-positive spermatocytes of *Fignl1*
cKO, indicating RAD51-focus formation during pre-meiotic S phase. The number
of RAD51 foci during pre-meiotic S-phase is around 300 in focus-positive nuclei
(298±110 foci and 337±108 foci in early and mid/late preleptonema, respectively),
which is ~3/4 of the RAD51 focus number in meiotic prophase I of *Fignl1* cKO
mice, but ~1.5-fold more of that in early zygonema of control mice (compare Figs.
3d and 4b). Some RAD51 foci in pre-meiotic S-phase are colocalized with EdU
dots/lines (38.8±12.6% of RAD51 foci colocalized with EdU; Supplementary Fig.
4d), suggesting that RAD51 foci are associated with DNA replication. However,
when DNA replication labeling with another thymidine analog IdU was shortened
for 10 min, most of RAD51 foci did not colocalize with IdU foci (Supplementary
Fig. 4a-c), though the degree of colocalization was significantly higher than
random colocalization (3.7±1.9% of IdU foci colocalized with RAD51 compared
to 2.5±1.3%, p=0.0261, Mann-Whitney U-test; 9.1±4.5% of RAD51 foci
colocalized with IdU compared to 6.2±3.3%, p=0.0439, Mann-Whitney U-test;
Supplementary Fig. 4c). It is unlikely that RAD51 is associated with ongoing DNA
replication forks in *Fignl1* cKO spermatocytes.

Further evaluation of DMC1 localization in *Fignl1* cKO revealed the
presence of preleptotene spermatocytes containing DMC1 foci (Fig. 4c, d). The
number of DMC1 foci in focus-positive nuclei is around 150 (136±71 foci and
164±70 foci in early and mid/late preleptonema, respectively), about half of
RAD51 foci in *Fignl1* cKO. As seen for RAD51, most of DMC1 foci did not
colocalize with ongoing DNA replication forks when DNA replication was labeled
with EdU for 10 min (23.0±7.9% of DMC1 foci colocalized with EdU;
Supplementary Fig. 5a-c). Of note is that very few DMC1 foci were detected in
control preleptotene spermatocytes. This suggests that FIGNL1 limits the
assembly of mitotic recombination machinery with RAD51 on chromatin during
pre-meiotic S-phase as well as meiotic recombination machinery containing
DMC1.

In contrast to compromised meiotic recombination and homolog synapsis
as described above, pre-meiotic DNA replication in *Fignl1 c*KO seemed largely

normal, as the number of RPA2 foci, which are expected to correspond to
replication fork-associated ssDNAs, and IdU foci with 10 min labeling, which
indicates ongoing DNA replication forks, in early preleptotene spermatocytes
were comparable between *Fignl1* cKO and controls (624±241 RPA2 foci in *Fignl1*
cKO compared to 607±217 foci in controls; *p*=0.9414, Mann-Whitney *U*-test;
500±131 IdU foci in *Fignl1* cKO compared to 476±140 foci in controls; *p*=0.4652,
Mann-Whitney *U*-test; Supplementary Fig, 6a-c). However, we cannot exclude
the possibility that FIGNL1 plays a role in pre-meiotic DNA replication as recently
shown in *FIGNL1* KO human cell lines [79].
382
Uncoupled loading of RAD51 and DMC1 in *Fignl1* cKO spermatocytes
Despite the overall increase in RAD51 and DMC1 focus numbers at each
analyzed stage of *Fignl1* cKO spermatocytes, we observed ~2-fold more RAD51
foci compared to DMC1 foci in preleptotene and early leptotene stages,
suggesting uncoupling of RAD51 and DMC1 loading onto chromosomes in the
absence of FIGNL1. Co-staining of RAD51 and DMC1 confirmed different
kinetics of RAD51 and DMC1 in *Fignl1* cKO (Fig. 3e, f, Supplementary Fig. 7);
DMC1 focus numbers gradually increase from preleptonema to late
leptonema/early zygonema (177±74, 325±136, and 407±69 foci in preleptonema,
early leptonema, and late leptonema/early zygonema, respectively; Fig. 3f),
whereas RAD51-focus numbers are nearly constant in those stages (399±120,
476±109, and 480±95 foci in preleptonema, early leptonema, and late
leptonema/early zygonema, respectively). Colocalization analysis revealed a
gradual increase in the number of RAD51-DMC1 co-foci as DMC1 foci increased.
In preleptotene and early leptotene spermatocytes, 50-60% of DMC1 foci
colocalized with RAD51 foci (57.0±10.6% and 53.1±11.0% in preleptonema and
early leptonema, respectively; Fig, 3f) and less than 40% of RAD51 foci
colocalized with DMC1 foci (25.7±9.9% and 35.3±12.0% in preleptonema and
early leptonema, respectively). The degree of colocalization at these stages is
significantly lower than that in the late leptotene/early zygotene stage where
comparable levels of colocalization with controls were observed (65.8±10.1% and
67.2±7.6% of DMC1 foci colocalized with RAD51 in *Fignl1* cKO and controls,
respectively; 56.2±9.6% and 59.3±8.9% of RAD51 foci colocalized with DMC1
foci in *Fignl1* cKO and controls, respectively; Fig. 3f). These suggest that coupled
loading of RAD51 and DMC1 in pre-meiotic S-phase and early meiotic prophase
I is compromised in *Fignl1* cKO spermatocytes. On the other hand, the fact that

more than half of DMC1 foci are colocalized with RAD51 in preleptonema
suggests that DMC1 preferentially binds to meiotic chromatin where RAD51 is
already bound.

RAD51/DMC1-focus formation in *Spo11* KO *Fignl1* cKO spermatocytes
Both RAD51 and DMC1 foci appear in pre-meiotic S-phase in *Fignl1* cKO
spermatocytes, which is much earlier than meiotic prophase I with programmed
DSB formation by SPO11. To examine whether RAD51/DMC1-focus formation in
Fignl1 cKO mice is independent of meiotic DSB formation or not, we constructed
Spo11 KO *Fignl1* cKO mice (*Spo11*[-/-] *Fignl1*[flox/Δ] *Stra8-Cre*[+]), which are expected
to be defective in SPO11-mediated DSB formation [77, 78]. As shown previously,
spermatocytes from control *Spo11* KO mice (*Spo11*[-/-] *Fignl1*[flox/Δ]) did not show
any RAD51 foci on meiotic chromosomes (Fig. 5a, b, Supplementary Fig. 8a). On
the other hand, spermatocytes from *Spo11* KO *Fignl1* cKO show on average 300-
400 RAD51 foci from leptotene to zygotene(-like) stages (407±77, 390±64, and
292±34 foci for focus-positive nuclei in leptonema, early zygonema, and late
zygonema, respectively; Fig. 5b). Moreover, *Spo11* KO *Fignl1* cKO also
accumulates DMC1 foci with similar numbers to RAD51 foci (374±108, 391±57,
and 342±54 foci for focus-positive nuclei in leptonema, early zygonema, and mid-
zygonema, respectively; Fig. 5c, d, Supplementary Fig. 8b). Comparable
numbers of RAD51 and DMC1 foci between *Fignl1* cKO and *Spo11* KO *Fignl1*
cKO indicate that a large fraction of RAD51 and DMC1 foci on meiotic chromatin
in the absence of FIGNL1 is independent of SPO11, thus meiotic DSBs. Like
Spo11 KO, *Spo11* KO *Fignl1* cKO spermatocytes lack pan-nuclear γH2AX
staining and form few RPA2 foci (Supplementary Fig. 8c-f), supporting the
assembly of RAD51/DMC1 complexes in the absence of endogenous DSBs and
ssDNAs. *Spo11* KO *Fignl1* cKO spermatocytes died in the pachytene-like stage
with the formation of the pseudo-sex body (Supplementary Fig. 8c) as seen in
Spo11 KO spermatocytes [77, 78]. We also detected RAD51/DMC1 foci in
preleptotene spermatocytes of *Spo11* KO *Fignl1* cKO mice (Fig. 5f). This confirms
that the formation of RAD51 and DMC1 foci during pre-meiotic S-phase is
independent of the SPO11 function. Importantly, the degree of RAD51-DMC1
colocalization from preleptonema to early zygonema in *Spo11* KO *Fignl1* cKO is
comparable to that in *Fignl1* cKO and more than half of DMC1 foci are colocalized
with RAD51 at those stages in *Spo11* KO *Fignl1* cKO (Fig. 5e-g). This supports
the idea that DMC1 preferentially binds to RAD51-bound meiotic chromatin,

which is most likely dsDNA in the absence of meiotic DSB formation.

RAD51 binds to and persists on meiotic recombination sites in *Fignl1* cKO
spermatocytes

As described above, the absence of FIGNL1 increases the SPO11-independent
formation of RAD51 and DMC1 foci in meiotic prophase I, some of which would
come from pre-meiotic S-phase. Comparable numbers of RAD51 and DMC1 foci
in the presence and absence of SPO11 (i.e., meiotic DSBs) prompted us to check
whether RAD51/DMC1 can bind to meiotic recombination hotspots, especially
ssDNA. Chromatin immunoprecipitation followed by single-stranded DNA
sequencing (ChIP-SSDS) for RAD51 [80-82] revealed that in *Fignl1* cKO testes, like
in controls, RAD51 binds to recombination hotspots where DMC1 strongly binds
in wild-type testes (Fig. 6a, Supplementary Fig. 9a). Importantly, when RAD51
binding around hotspots is ranked based on the number of SPO11-oligos in wild-
type, a similar binding profile of RAD51 is seen between *Fignl1* cKO and controls,
with a small increase in RAD51 ChIP-SSDS signals in two out of three *Fignl1*
cKO mice (Fig. 6b, Supplementary Fig. 9b-d). We also ran RAD51 ChIP-SSDS
in *Spo11* KO and *Spo11* KO *Fignl1* cKO testes, confirming no RAD51 enrichment
at recombination hotspots and lower RAD51 enrichment in the entire genome
(Supplementary Fig. 10a-c). Most of non-overlapping 2-kb bins with high RAD51
enrichment in *Spo11* KO *Fignl1* cKO testes were common to *Spo11* KO testes
(94.1%; 241/256; Supplementary Fig. 10d), indicating a similar binding profile of
RAD51 on ssDNAs between *Spo11* KO and *Spo11* KO *Fignl1* cKO mice. These
suggest that, in the absence of FIGNL1, RAD51 does not bind to any specific
regions with ssDNAs in *Spo11* KO spermatocytes.

To address whether RAD51 persists on meiotic recombination sites in the
absence of FIGNL1, we also measured the colocalization frequency of RAD51
and RPA2 in mid/late zygonema, given that RPA2 localizes to recombination sites
even after RAD51/DMC1 disassembly in wild-type meiosis and can be a marker
of recombination sites on mid/late zygotene chromosomes. In control mid/late
zygotene spermatocytes, out of 234±44 RPA2 foci, 41±15 foci were colocalized
with RAD51 foci, which corresponds to 17.2±4.3% of RPA2 foci (Fig. 6c-e). *Fignl1*
cKO exhibited two classes of populations; one is similar to controls (~30%) and
the other shows 40-80% of colocalization, with 45.4±21.9% of RPA2 foci
colocalized with RAD51 for all populations. This is consistent with the idea that
RAD51 binds to ssDNA on meiotic recombination sites in the absence of FIGNL1,

suggesting that RAD51 persists on RPA2-associated recombination sites.

Fignl1 deficiency rescues RAD51/DMC1 assembly defects in *Swsap1*

Our previous study in human cells showed that FIGNL1 antagonizes the function
of the RAD51 mediator, SWSAP1 [51]. In mouse male meiosis, SWSAP1 is
necessary for efficient RAD51/DMC1 focus formation [17, 51]. We tested the effect
of *Fignl1* cKO on RAD51/DMC1 assembly in *Swsap1* KO cells by generating
Swsap1 KO *Fignl1* cKO male mice (*Swasp1$^{-/-}$ Fignl1$^{flox/\Delta}$ Stra8-Cre^{+}*). As reported
[17, 51], surface-spread chromosomes from *Swasp1* KO spermatocytes (*Swsap1$^{-/-}$*
Fignl1$^{flox/\Delta}$) showed a large reduction of RAD51 foci with 10-30 foci (~1/10 of
control *Fignl1$^{flox/\Delta}$*) in leptonema and zygonema (29±26, 22±15, and 20±14 foci in
leptonema, early zygonema, and mid zygonema, respectively; Fig. 7a, b). Like
Fignl1 cKO, *Swsap1* KO *Fignl1* cKO spermatocytes showed an accumulation of
RAD51 foci on meiotic chromosomes. The number of RAD51 foci from leptonema
to mid-zygonema in *Swsap1* KO *Fignl1* cKO is very similar to that in *Fignl1* cKO
(320±217, 308±221, and 194±179 foci in *Swsap1* KO *Fignl1* cKO and 399±145,
377±98, and 282±91 foci in *Fignl1* cKO in leptonema, early zygonema, and mid
zygonema, respectively). This indicates that in mouse spermatogenesis without
FIGNL1, RAD51 assembly does not depend on SWSAP1. This is consistent with
the results in human cultured cells. On the other hand, in late zygonema, the
number of RAD51 foci in *Swsap1* KO *Fignl1* cKO was comparable to *Swsap1* KO
and much lower than that in *Fignl1* cKO (10±12, 7±8, and 189±79 foci in *Swsap1*
KO *Fignl1* cKO, *Swsap1* KO, and *Fignl1* cKO, respectively; Fig. 7b). This
suggests that only *Swsap1$^{-/-}$ Fignl1$^{flox/\Delta}$* cells where Cre recombinase failed
excision of *Fignl1^{flox}* allele can progress into late zygotene stage in *Swsap1* KO
Fignl1 cKO testes. Moreover, similar results were observed for DMC1; the
number of DMC1 foci in *Swsap1* KO *Fignl1* cKO was comparable to *Fignl1* cKO
(Fig. 7c, d). Thus, reduced DMC1-focus formation in *Swsap1* KO is also
suppressed by *Fignl1* cKO. It is likely that the activity of FIGNL1 to suppress
RAD51/DMC1 assembly on intact dsDNAs is independent of SWSAP1-mediated
stable assembly of RAD51/DMC1 on DSB-induced ssDNAs.

FIGNL1 disrupts RAD51-dsDNA complex *in vitro*

We previously showed that FIGNL1 promotes the dissociation of RAD51 from
ssDNAs *in vitro* [51]. The above results in mouse spermatocytes suggest that
FIGNL1 can disrupt RAD51 filaments on chromosomal DNAs, both ssDNAs and

dsDNAs. To address whether FIGNL1 promotes the dissociation of RAD51
filaments on dsDNAs *in vitro*, we purified human FIGNL1ΔN, which lacks N-
terminal 284 amino acids [51], since a full-length human FIGNL1 protein was
insoluble when expressed in *E. coli*, and performed the electrophoretic mobility
shift assay (EMSA). First, we confirmed that a purified human FIGNL1ΔN
disrupted the pre-formed RAD51-ssDNA complexes in the presence of ATP (Fig.
8a, b). Under this condition, FIGNL1ΔN could disrupt ~30% of RAD51-ssDNA
complex (24.6±6.9% of unbound DNAs were increased to 42.9±2.1% by the
addition of 1.2 μM of FIGNL1ΔN). When FIGNL1ΔN was incubated with a pre-
formed RAD51-dsDNA complex, the most of dsDNA was recovered as an
unbound form (25.8±11.1% of unbound DNAs were increased to 80.0±11.6%).
This suggests that FIGNL1ΔN could disrupt the RAD51-dsDNA complex more
efficiently than the RAD51-ssDNA complex *in vitro*, supporting the *in vivo* results.
Alternatively, RAD51-ssDNA *in vitro* might form an aggregate more than RAD51-
dsDNA, which may affect the accessibility by FIGNL1 protein to disrupt the
complexes.
533
Discussion
In this study, we analyzed the role of FIGNL1 AAA+ ATPase, which plays a critical
role in homologous recombination in somatic cells, in mouse male meiosis. *Fignl1*
cKO mice displayed higher levels of RAD51 and DMC1 foci than controls, as well
as defective chromosome synapsis. Our analysis revealed the dual function of
FIGNL1 during spermatogenesis (Fig. 8c). One is to promote meiotic
recombination by regulating the dissociation of RAD51/DMC1 ensembles on the
ssDNA. The other is to protect the genome by disrupting the non-functional form
of RAD51/DMC1 ensembles on the dsDNA. Consistent with the idea, purified
FIGNL1 can disrupt the RAD51-dsDNA complex as well as the RAD51-ssDNA
complex *in vitro*. The same conclusion was obtained by Baudat's group by
characterizing the *Fignl1* cKO as well as *Firrm* cKO (accompanying paper).
546
FIGNL1 promotes the post-assembly step of RAD51/DMC1 in meiotic
recombination
In meiotic recombination, RAD51/DMC1 filaments on ssDNAs play a critical role
in homology search and strand exchange with homologous dsDNA, leading to
the formation of the D-loop, in which the 3'-OH end of the invading strand
becomes a substrate for recombination-associated DNA synthesis. Biochemical

analyses showed that RAD51/DMC1-mediated D-loop is a poor substrate for *in*
vitro DNA synthesis by DNA polymerase δ and its accessory factors [83-85]. To
promote efficient DNA synthesis on the recombination intermediate,
RAD51/DMC1 bound to the intermediate should be removed by factors such as
RAD54, a RAD51 remodeler [83-85]. Although the male KO mice for *Rad54* and
Rad54b, a *Rad54* paralog, and the double KO mice for *Rad54* and *Rad54b* are
fertile, the *Rad54 Rad54b* double KO spermatocytes show abnormal loading of
RAD51 on chromosomes in pachynema and diplonema [54]. This is consistent with
the role of RAD54 in the post-assembly step of RAD51 (DMC1 had not been
analyzed in detail). Our *in vivo* and *in vitro* studies suggest the role of FIGNL1 in
the post-assembly step of the RAD51/DMC1 filaments. It is likely that FIGNL1
disrupts not only RAD51 but also DMC1 filaments on recombination
intermediates for subsequent DNA synthesis [84, 86] (Fig. 8c), though other
possibilities including that proper disassembly of RAD51 and DMC1 filaments
promotes strand invasion won't be excluded. Post-assembly role of FIGNL1 in
RAD51-mediated homologous recombination was reported previously in human
cells with DNA damage [56]. Consistent with the post-assembly role of FINGL1,
Fignl1 cKO spermatocytes reduce the formation of MSH4 foci, which appears
after the loading and dissociation of RAD51/DMC1 [72]. *Fignl1* cKO spermatocytes
seem not to enter the pachytene stage for the loading of the MutLγ complex such
as MLH1 and most *Fignl1* KO cells die around late zygonema. The meiotic
defects seen in *Fignl1* cKO are similar to those in the mutants where
RAD51/DMC1 foci accumulate in a higher number than normal such as *Msh4*,
Msh5, *Mcm8*, *Trip13,* and *Mcmdc2* mutant spermatocytes [87-93]. In mouse
spermatocytes, FIGNL1 seems to promote ZMM-dependent CO formation. This
is different from the role of FIGNL1 ortholog, FIGL1, in *A. thaliana*. *Arabidopsis*
FIGL1 mutant meiocytes showed normal or slightly elevated CO formation [57, 58].
Indeed, the depletion of FIGL1 or its partner FLIP suppresses reduced CO
formation in the absence of pro-CO proteins, ZMM [57, 58]. This suggests the role
of plant FIGL1-FLIP as an anti-CO factor, which disrupts early recombination
intermediates such as RAD51/DMC1 filaments on the ssDNA prior to or during
the strand invasion. Although our attempt to analyze the role of FIGNL1 in mouse
oocytes failed due to insufficient excision of the *Fignl1^flox* allele in oocytes by the
Stra8-Cre transgene used in this study, a recent study by Yu and colleagues
demonstrated the accumulation of RAD51 in not only *Firrm/Flip* cKO
spermatocytes but also in oocytes with infertility in both male and female mice [79],

suggesting that the essential role of the FIGNL1-FIRRM complex in meiotic
recombination is shared between males and females.
591
FIGNL1 disrupts non-productive RAD51 and DMC1 ensembles on duplex
DNAs
RAD51/DMC1 focus formation normally occurs at sites of ssDNA tracks that are
associated with mitotic or meiotic DSBs. The most striking phenotypes seen in
Fignl1 cKO spermatocytes are the accumulation of RAD51 and DMC1 foci in
unchallenged pre-meiotic S-phase cells and spermatocytes in the absence of
SPO11-dependent meiotic DSBs. This accumulation of RAD51/DMC1 foci in the
absence of DSBs, therefore the ssDNAs, could be simply explained by the non-
specific assembly of RAD51/DMC1 filaments on chromatin DNAs, thus the
binding to dsDNA (Fig. 8c). Interestingly, human RAD51 and DMC1 bind to both
ssDNAs and dsDNA with similar affinity *in vitro* [16, 94] and RAD51 bears an activity
to remodel the nucleosome upon the dsDNA binding [95-97]. On the other hand,
detection of RAD51/DMC1 foci *in vivo* normally requires the presence of tracts of
ssDNAs. How cells discriminate ssDNAs among lots of competitors, dsDNAs, in
the RAD51/DMC1 binding remains unsolved. Based on the results described
here, we propose that FIGNL1 actively dismantles RAD51 and DMC1 filaments
on dsDNAs, which is otherwise non-productive for the recombination and/or
might become obstacles to chromosomal events on duplex DNAs such as DNA
replication, transcription, and others. Indeed, Bishop, Heyer and their colleagues
reported activities of the RAD54 translocase family to remove RAD51 and DMC1
from dsDNAs. In yeast meiosis, Spo11-independent focus formation of Dmc1, but
not of Rad51, depends on Tid1/Rdh54, a Rad54 homolog [98]. And mitotic
spontaneous Rad51-focus formation on yeast chromosomes is increased in the
absence of three translocases, Rad54, Tid1/Rdh54, and Uls1 [99]. Moreover, in
unchallenged human cells, the absence of two RAD54 family proteins RAD54 (or
RAD54L) and RAD54B leads to the accumulation of spontaneous RAD51 foci
particularly when RAD51 is over-expressed [100]. These *in vivo* studies were further
supported by biochemical analysis showing that Rad54 dissociates RAD51 from
dsDNAs [101-103]. Further studies are necessary to reveal the relationship between
FIGNL1 and RAD54 translocases in the disassembling of RAD51/DMC1 on
duplex DNAs. We stress that, distinct from RAD54, which dismantles RAD51
filaments only on dsDNAs (not on ssDNAs), FIGNL1 can disrupt RAD51
ensembles on both ss- and dsDNAs.

One of the striking features of RAD51/DMC1 assembles on duplex DNAs is
that these ensembles are quite stable. The number of RAD51 in both pre-meiotic
S-phase and early meiotic prophase I from early leptonema to early zygonema is
constant. Little increase in RAD51-focus number during early meiotic prophase I
suggest that spermatocytes show very limited additional assembly of RAD51 foci
during this stage. On the other hand, DMC1 shows a 2-fold increase in focus
number from early leptonema (and pre-meiotic S-phase) to early zygonema. This
increase could be induced by SPO11-mediated DSBs. Interestingly, the
colocalization frequency of RAD51 with DMC1 in *Fignl1* cKO increased from
~30% to ~60%, which is similar to the colocalization frequency seen in control
early zygonema, while the colocalization frequency of DMC1 with RAD51 is
nearly constant (with 60-70% frequency). This suggests that RAD51 ensembles
even on the dsDNA tend to recruit DMC1 to form co-ensembles.
638
FIGNL1 antagonizes SWSAP1 for RAD51/DMC1 assembly on meiotic
chromosomes
FIGNL1 shows antagonism to the RAD51 mediator. In human cells, *FIGNL1*
depletion suppresses RAD51-assembly defect induced by the *SWSAP1*
depletion [51]. In meiosis of the *A. thaliana*, *FIGL1* (*A. thaliana FIGNL1*) and/or *FLIP*
deletion restores reduced RAD51/DMC1-focus formation in the mutants of the
BRCA2 [61] as well as defective DMC1-focus formation in the mutant of the *SDS1*,
which promotes DMC1 assembly [57]. Since *Swsap1* KO spermatocytes reduced
the RAD51 and DMC1 foci [17, 51], we asked whether *Fignl1* cKO could also
suppress these meiotic defects in *Swsap1* KO spermatocytes and found that the
number of both RAD51/DMC1 foci in *Swsap1* KO *Fignl1* cKO is similar to those
in *Fignl1* cKO. This is simply interpreted as, like in human cells, *Fignl1* cKO
suppresses RAD51/DMC1-assembly defect in *Swsap1* KO. However, at this
point, we do not know whether RAD51/DMC1 foci are associated with ssDNAs in
meiotic prophase I cells of *Swsap1* KO *Fignl1* cKO as shown in *Fignl1* cKO,
although it is likely. In addition, different from RAD51/DMC1 assembly on ssDNAs,
these suggest that RAD51/DMC1 assembly on the dsDNA in the absence of
FIGNL1 does not require the RAD51/DMC1 mediator, SWSAP1.
657

Methods

Mice

The care and use of mice in this study were performed under the guideline for the proper conduct of animal experiments (Society Council of Japan). These procedures were approved by the Institutional Animal Care Committee at Institute for Protein Research, Osaka University (approval ID; 25-03-0).

All mice used in this study were congenic to the C57BL/6 background. Genotyping of mice was performed by PCR on genomic DNA isolated from mice. Primers used for genotyping are listed in Supplementary table 1. Mature adult mice (8-20 weeks old) were used for experiments otherwise mentioned.

Fignl1^{flox} allele was constructed by Cyagen Bioscience Co. Ltd. ES cells with the *Fignl1^{flox}* allele were injected into C57BL/6 albino blastocytes, which were transplanted into CD-1 pseudo-pregnant female mice. The founder mice with the correct *Fignl1^{flox}* allele were backcrossed to C57BL/6 mice. *Fignl1$^{+/\Delta}$* allele was constructed by breeding *Fignl1$^{flox/+}$* with *CAG-Cre* mice. *Stra8-Cre* transgenic mice were previously described [104] and provided *by* the RIKEN Animal Resource Center. *Spo11$^{-/-}$* mouse was previously described [77] and *Spo11$^{+/-}$* mice with a mixed background of C57BL/6 and 129/SvJ were backcrossed to C57BL/6 mice prior to experiments. *Swsap1$^{-/-}$* mice were described in our previous study [51].

Histology

Testes from adult mice were fixed in 10% buffered formalin overnight at room temperature, washed twice with PBS, and stored in 70% EtOH at 4°C prior to embedding in paraffin and sectioning. The sections of 5 μm thickness were deparaffinized and stained with periodic acid-Schiff (PAS).

Spermatocyte-chromosome spreads

Surface-spread chromosomes of spermatocytes were prepared as described previously [105, 106] with slight modification. Testes were dissected, the tunica albuginea was removed, and adherent extra-tubular tissues were removed by rinsing and dissociating seminiferous tubules in PBS using a pair of 25-gauge needles in a 35 mm petri dish (Corning, 430165). Dissociated seminiferous tubules were incubated in hypotonic extraction buffer (30 mM Tris-HCl pH 8.0, 50 mM sucrose, 17 mM trisodium citrate dihydrate, 5 mM EDTA, 0.5 mM Dithiothreitol (DTT) and 0.5 mM phenylmethylsulphonyl fluoride (PMSF), pH 8.2-

8.3; HEB) for 15-45 min at room temperature. After quickly rinsing seminiferous
tubules in 100 mM sucrose, some tubules were placed in 40 μl of 100 mM sucrose
on a glass depression slide. Tubules were torn to pieces using two fine forceps,
and pieces of tubular remnants were removed. The cell suspension was made by
pipetting, and 20 μl of cell suspension was placed on a clean glass slide
(Matsunami, MAS-01) coated with freshly made 1% paraformaldehyde (PFA)
solution pH 9.2 (pH set by 1.25 M sodium borate) containing 0.15% Triton X-100.
Slides were placed and dried in a closed humid chamber with hot tap water
overnight at room temperature, followed by drying with lids ajar for 3 h and then
with lids removed for 1-2 hr. Slides were washed once for 5 min in deionized
water and twice for 5 min in 0.4% DryWell (Fujifilm) in a coplin jar and air-dried at
room temperature. Slides were either directly processed for immunostaining as
below or stored in aluminum foil at -80^{o}C prior to immunostaining.
For the preparation of testicular cell suspension, testes were dissected, the
tunica albuginea was removed, and seminiferous tubules were transferred into 1
ml of testis cell isolation medium (104 mM NaCl, 45 mM KCl, 1.2 mM MgSO$_4$, 0.6
mM NaHPO$_4$, 0.15 (w/v) glucose; TIM) containing 2 mg of collagenase
(Worthington, LS004186) for incubation at 32^{o}C for 55 min with shaking at 53 rpm.
Tubules were washed three times with 3 ml of TIM, and suspended in 1 ml of TIM
containing 0.7 mg of Trypsin (Worthington, LS003740) and 4 μg of DNase I
(Roche, 104159) for incubation at 32^{o}C for 15 min with shaking at 53 rpm.
Trypsinization was stopped by adding 5 mg of Trypsin inhibitor (Sigma, T9003)
and 10 μg of DNase I, and cells were separated by pipetting and filtered through
a 70 μm cell strainer (Corning, 431751). Cells were washed once with 0.75 μg of
DNase I and 7.5 ml of TIM, and suspended in 0.75 μg of DNase I and PBS
(Nakarai, 14249-24) to 1 x10^6 cell/ml. 1 ml of cell suspension was centrifuged at
200 x g for 5 min at room temperature. After carefully removing the supernatant,
cells were suspended in 30 μl of HEB and incubated for 15-45 min at room
temperature. 30 μl of 100 mM sucrose was added, and 20 μl of cell suspension
was placed on a clean glass slide (Matsunami, MAS-01) coated with 100 μl of
freshly made 1% paraformaldehyde (PFA) solution pH 9.2 containing 0.15%
Triton X-100. Slides were placed and dried in a closed humid chamber, washed,
and stored as described above.
727
Immunofluorescence staining
Slides were rehydrated with Tris-buffered saline (TBS) pH 8.0 containing 0.05%

of Triton X-100 (TBST) for 3 min, blocked twice with blocking buffer (1% normal
goat or donkey serum, 3% bovine serum albumin (BSA), 1x TBS pH 8.0, 0.05%
of Triton X-100, 0.05% sodium azide) for 15 min at room temperature and
incubated with primary antibodies in antibody dilution buffer (10% normal goat or
donkey serum, 3% BSA, 1x TBS pH 8.0, 0.05% of Triton X-100, 0.05% sodium
azide) in a humid chamber overnight at room temperature. Slides were quickly
rinsed with TBST, washed twice with TBST for 5 min, blocked twice with blocking
buffer for 15 min at room temperature, and incubated with secondary antibodies
in antibody dilution buffer in a humid chamber for 1 h at 37°C. Slides were quickly
rinsed with TBST, washed three times with TBST for 5 min, washed once with
Milli-Q water for 2 min, and air-dried before mounting with ProLong Diamond
Antifade Mountant (Thermo Fisher Scientific, P36970). For detection of IdU,
rehydrated slides were denatured with 4N HCl for 10 min at room temperature
before blocking. EdU was detected using Click-iT EdU Imaging Kit with Alexa
Fluor 647 azides according to the manufacturer's instruction (Invitrogen). All
primary and secondary antibodies used are listed in Supplementary table 2.
746

Testicular cell squashes
Testes were dissected, the tunica albuginea was removed, and adherent extra-
tubular tissues were removed by rinsing and dissociating seminiferous tubules in
PBS using a pair of 25-gauge needles in a 35 mm petri dish (Corning, 430165).
Dissociated seminiferous tubules were fixed in freshly prepared 2%
formaldehyde containing 0.1% Triton X-100 for 10 min at room temperature.
Small pieces of tubules were placed on a clean glass slide (Matsunami, MAS-01),
torn using two fine forceps, and squashed under a coverslip with pressure from
the blunt end of pencil followed by a thumb. Slides were frozen briefly in liquid
nitrogen and either directly processed for immunostaining or stored at -80°C prior
to immunostaining. After the removal of coverslips, slides were washed three
times with PBS for 5 min and immunostaining was performed as described above.
759

Image acquisition
Images of mouse spermatocyte spreads testicular cell squashes were acquired
using a computer-assisted fluorescence microscope system (DeltaVision;
Applied Precision) with a $60\times$ NA 1.4 oil immersion objective. Image
deconvolution was performed using an image workstation (SoftWorks; Applied
Precision) and afterward processed using Imaris (Oxford Instruments, UK) and

Photoshop (Adobe, USA) software tools. Images in each figure are matched
exposure.
768
Image analysis
Comparisons were made between animals that were either littermate or matched
by age. Single cells were manually cropped, and numbers of foci were counted
by the auto-thresholding signal intensity in Imaris software (Oxford Instrument)
and clear foci that were missed by auto-thresholding, especially adjacent foci,
were manually counted. Signal intensities were measured using Image J software
by subtracting signal intensities of single cells from those of regions of interest
(ROIs) manually drawn next to cells as backgrounds.
Meiotic prophase I stages on spermatocyte spreads were defined by
SYCP3 and SYCP1 staining using standard criteria. Leptonema was defined by
short SYCP3 stretches without evidence of synapsis determined by SYCP1
staining. Zygonema was defined by longer stretches of SYCP3 with various
degrees of synapsis: early, mid- and late zygonema was defined by having <25%,
25-75%, and >75% of synapsis, respectively. Pachynema was defined by the full
synapsis of all autosomes. Diplonema was defined by de-synapsis beginning
from mostly the middle of chromosomes with various degrees of residual
synapsis. Preleptonema was defined by very short SYCP3 stretches with EdU or
IdU-positive staining.
787
EdU incorporation
1 ml of 1 x10^6 cell/ml testicular cell suspension in PBS was centrifuged at 200 x
g for 5 min at room temperature, and after removal of the supernatant, cells were
suspended in 500 μl of D-MEM (Wako, 044-29765) / 10% FBS containing 10 μM
EdU. After incubation at 37^{o}C for 10 min or 1 h in a well of a 12-well dish (Violamo),
cells were pelleted by centrifugation at 200 x g for 5 min at room temperature,
washed in 1 ml of PBS once, and suspended in 30 μl HEB for incubation for 15-
30 min at room temperature. After adding 30 μl of 100 mM sucrose, cells were
fixed, and the slides were dried, washed, and stored as described above.
797
IdU pulse incorporation
1 ml of 1 x10^6 cell/ml testicular cell suspension in PBS was centrifuged at 200 x
g for 5 min at room temperature, and after removal of the supernatant, cells were
suspended in 500 μl of D-MEM (Wako, 044-29765) / 10% FBS containing 20 μM

IdU (Sigma, I7125). After incubation at 37°C for 10 min in a well of a 12-well dish

(Violamo), cells were subjected to chromosome spreads as described above.

Rad51 Chromatin immuno-precipitation – single-stranded DNA sequencing

(ChIP-SSDS)

ChIP was performed as described previously [80]. One testis from the control and

a pair of testes from *Fignl1* cKO, *Spo11* KO, and *Spo11* KO *Fignl1* cKO mice (9-

20 weeks, except a *Spo11* KO mouse at 31 weeks) were dissected, the tunica

albuginea was removed, and seminiferous tubules were crosslinked in 10 ml of

1% formaldehyde (Pierce, 28908) in PBS (Nakarai, 14249-24) for 10 min at room

temperature. Crosslinking was quenched by adding 2.5 M glycine to a final

concentration of 125 mM. After 5 min of incubation at room temperature, cells

were washed once with 10 ml of cold PBS, and the cell pellet was snap-frozen

on liquid nitrogen and stored at -80°C. The frozen cell pellet was suspended in

500 µl of cell lysis buffer (0.25% Triton X-100, 10 mM EDTA, 0.5 mM EGTA, 10

mM Tris-HCl pH 8.0) supplemented with protease inhibitor (Roche,

04693159001), incubated for 10 min on ice, and centrifuged at 300 x *g* for 3 min

at 4°C. The nuclei pellet was washed in 500 µl of lysis wash buffer (200 mM NaCl,

1 mM EDTA, 0.5 mM EGTA, 10 mM Tris-HCl pH 8.0) supplemented with protease

inhibitor, resuspended in 200 µl of RIPA buffer (10 mM Tris-HCl pH 8.0, 1 mM

EDTA, 0.5 mM EGTA, 1% Triton X-100, 0.1% sodium deoxycholate, 0.1% SDS,

140 mM NaCl) supplemented with protease inhibitor, split into two ~130 µl

aliquots and sonicated on the Covaris M220 for 5 min 30 sec at 5-9°C with a

setting of Peak Power 50.0, Duty Factor 20.0, and Cycles/Burst 200. Sheared

nuclei were combined, and after adding 700 µl of RIPA buffer, centrifuged at

15,000 rpm for 15 min at 4°C to remove insoluble debris. 800 µl of the chromatin-

containing supernatant was mixed with 50 µl of Dynabeads protein G bound to

9.7 µg of RAD51 antibody (Novus, NB100-148) and incubated overnight on a

rotating wheel at 4°C for immunoprecipitation. Beads were washed once with 1

ml of cold ChIP wash buffer 1 (0.1% SDS, 1% Triton X-100, 2 mM EDTA, 20 mM

Tris-HCl pH 8.0, 150 mM NaCl), 1 ml of cold ChIP wash buffer 2 (0.1% SDS, 1%

Triton X-100, 2 mM EDTA, 20 mM Tris-HCl pH 8.0, 500 mM NaCl), 1 ml of cold

ChIP wash buffer 3 (0.25 M LiCl, 1% NP-40, 1% sodium deoxycholate, 1 mM

EDTA, 10 mM Tris-HCl pH 8.0), and twice with 1 ml of cold TE (1 mM EDTA and

10 mM Tris-HCl pH 8.0), followed by elution with 100 µl of elution buffer (0.1 M

NaHCO$_3$ and 1% SDS) for 30 min at 65°C with vortexing every 10 min. The elutes

were incubated overnight at 65°C for de-crosslinking, neutralized by adding 4 μl

of 1M Tris-HCl (pH 6.5) and 2 μl of 0.5 M EDTA, and RNA was digested by adding

2 μl of 10 mg/ml RNase-A and incubation for 15 min 37°C. After adding 5 μl of 10

mg/ml Proteinase-K (Wako, 165-21043) and incubation for 1 h at 55°C, DNA was

purified using QIAquick PCR Purification Kit (QIAGEN).

SSDS was performed as described previously (Brick et al., 2018). 7-10 ng

of ChIP DNA was further sonicated on the Covaris M220 for 8 min 30 sec at room

temperature with a setting of Peak Power 75.0, Duty Factor 10.0, and

Cycles/Burst 200, followed by end-repair and dA-tailing using ChIP-Seq Library

Prep Reagent Set for Illumina (NEB, E6200L). DNA was incubated for 3 min at

95°C and returned to room temperature for kinetic enrichment for single-stranded

DNA. After adapter ligation using IDT for Illumina TruSeq DNA UD Indexes

(Illumina, 20022370), SSDS libraries were amplified during 15 cycles of PCR

using TruSeq DNA Nano LP (Illumina, 20016328), purified with AMPureXP beads

(Beckman Coulter, A63880), and sequenced on HiSeq2500 in the rapid mode in

a 50 bp PE run. Around 70 million paired reads were generated per sample.

A published bioinformatic pipeline (https://nf-co.re/ssds/dev) was run for the

resulting reads for the identification of single-stranded sequences and mapping

to the mm10 genome [107]. Heatmaps were generated using Deeptools [108].

Western blot

Testes were dissected, the tunica albuginea was removed, and seminiferous

tubules were snap frozen on liquid nitrogen and stored at -80°C before cell lysis.

Tubules from either one testis (*Fignl1$^{+/+}$ Stra8-Cre^{+}* and *Fignl1$^{flox/Δ}$*) or two testes

(*Fignl1$^{flox/Δ}$ Stra8-Cre^{+}*) were homogenized in 1 ml of RIPA buffer (50 mM Tris-HCl

pH 7.5, 150 mM NaCl, 1 mM EDTA, 1% NP-40, 0.5% sodium deoxycholate, 0.1%

sodium dodecyl sulfate (SDS)) supplemented with protease inhibitor (Roche,

04693159001)) on ice, incubated for 15 min on ice, sonicated with a Bioruptor for

30 min (30 cycles of 30 sec ON / 30 sec OFF) at 4°C with a setting of High

intensity, and incubated for 15 min on ice. The samples were centrifuged at

14,000 rpm for 15 min at 4°C and the supernatants were collected as whole-testis

extracts. Protein concentrations were measured by Bradford assay and the

concentrations between samples were made constant by adding RIPA buffer

supplemented with protease inhibitor. Whole-testis extract was mixed with 4x

SDS-PAGE loading buffer (250 mM Tris-HCl pH 6.8, 8% SDS, 40% glycerol, 572

mM β-mercaptoethanol, 0.05% bromophenol blue) and boiled for 5 min at 100°C

and subjected to immunoblotting.

Protein samples were separated on 10% gels by SDS-PAGE and
transferred to PVDF membranes by wet transfer method in Towbin buffer (25 mM
Tris, 192 mM glycine, 20% ethanol) at 60 V for 150 min at 4°C. Membranes were
blocked with PBS containing 2.5% non-fat milk for 1 h at room temperature and
incubated with primary antibodies in 2.5% non-fat milk in PBS overnight at 4°C.
Membranes were washed three times with PBS containing 0.05% Tween-20
(PBST) for 10 min and incubated with HRP-conjugated secondary antibodies in
PBST for 1 h at room temperature. Membranes were washed three times with
PBST for 10 min and the signal was developed by ECL Prime Western Blotting
Detection Reagent (Amersham) and detected by ImageQuant LAS4000 (GE
Healthcare).

Antibody generation
His-tagged mouse SYCP3 was overexpressed from pET28c-SYCP3 plasmid by
the addition of 0.2 mM IPTG in BL21(DE3) at 37°C for 4 h. Cell extracts were
prepared in Lysis buffer (50 mM Tris-HCl pH 7.5, 500 mM NaCl) by sonication
and the supernatants were collected after centrifugation at 20,400 x *g* for 30 min
at 4°C. After adding Tween-20 to a final concentration of 0.2%, the cell lysates
were incubated with Ni-NTA Agarose (QIAGEN) for 2 h at 4°C and the lysate-
resin slurry was washed in Lysis buffer containing 50 mM Imidazole, 100 mM
Imidazole, 300 mM Imidazole. Ni-NTA-bound His-SYCP3 was eluted in Lysis
buffer containing 3M Imidazole and 3M Urea (elution 1) by incubation for 1 h at
4°C and in Lysis buffer containing 0.1% SDS (elution 2) by incubation for 30 min
at room temperature. Elution 2 was concentrated using MICROCON Centrifugal
Filter Device YM-10 (Millipore) and subjected to immunization to a rat.

Purification of RAD51 and FINGL1∆N
GST-tagged human FIGNL1∆N (N-terminal 284 aa deletion) was overexpressed
from pGEX-6P-1-*Fignl1∆N* plasmid by the addition of 0.25 mM IPTG in
BL21(DE3) pLysS at 30°C for overnight. Cell extracts were prepared in Lysis
buffer (25 mM Tris-HCl pH 7.6, 150 mM NaCl, 10% Glycerol, 1 mM DTT) and
GST-FIGNL1∆N was recovered by GSTrap HP column (Cytiva). FIGNL1∆N was
eluted from the GSTrap column by the cleavage of GST tag by Prescission
protease (#27-0843-01, Cytiva) in SP buffer (20 mM HEPES-NaOH pH 7.5, 10%
Glycerol, 1 mM DTT) supplemented with 100 mM NaCl. FIGNL1∆N was further

purified by HiTrap SP column (Cytiva) with a linear gradient of 100 mM-1M in SP
buffer. The peak fractions are dialyzed against SP buffer supplemented with 100
mM NaCl, concentrated with Amicon Ultra 30K (Millipore), and stored at -80°C.
Human RAD51 was purified as described previously using pET-HsRad51 plasmid
[109].

Electrophoresis mobility shift assay (EMSA)
Double-stranded DNA substrate was prepared by annealing 5'-Cy5-labeled, 40-
mer   oligo   DNA   (Oligo   1,   Cy5-
TAATACAAAATAAGTAAATGAATAAACAGAGAAAATAAAG,   IDT)   and   its
complementary   DNA   (Oligo   2,
CTTTATTTTCTCTGTTTATTCATTTACTTATTTTGTATTA,   Fasmac).   Oligo   1
was used as an ssDNA substrate. For the assay, DNA (0.8 μM in bp/nt) was pre-
incubated with 1 μM of RAD51 in 10 μl of the buffer (20 mM HEPES-NaOH pH7.5,
100 ug/ml BSA, 1 mM DTT, 10 mM MgCl$_2$, 2 mM ATP) at 37°C for 5 min to form
RAD51-DNA complex. Then the indicated concentration of FIGNL1ΔN was added
to the reaction and further incubated at 37°C for 15 min. In the control reaction
without FIGNL1ΔN, the same volume of dialysis buffer was added instead of the
protein. DNA-protein complexes were analyzed using 8% native polyacrylamide
gel electrophoresis with 1×TBE. Cy5-labelled DNA was visualized by LAS4000
(GE healthcare) and unbound DNA was quantified using ImageQuant TL (GE
healthcare). Images were further processed to adjust brightness using Photoshop
(Adobe, USA) software.

Statistical analysis and reproducibility
Statistical analyses were performed using Graphpad Prism software v.9 and R
version 4.2.2 (https://www.r-project.org). Bars in figures and values in
manuscripts indicate means ± standard deviations (SDs) otherwise mentioned.
Statistic parameters and tests and sample sizes are described in the figures
and/or figure legends. Sample sizes were not predetermined using any statistical
tests.
At least two animals of each genotype were analyzed and similar results
were obtained otherwise mentioned. Three independent experiments were
performed for biochemical analysis.

Data availability

The raw and processed sequencing data have been deposited in the Gene
Expression Omnibus (GEO) repository under accession number GSE227944.
Uncropped immunoblotting images and an unprocessed image of electrophoresis
mobility shift assay are provided in Source Data. The raw data used in plots and
graphs are also provide in Source Data. The raw microscopy images used for
figures and quantification are available from corresponding authors upon request.
952

953 **Code availability**
954 A published bioinformatic pipeline for single-stranded DNA sequencing analysis
955 is available at https://nf-co.re/ssds/dev.
956

Acknowledgments
We thank Drs. F. Baudat and B. de Massy for sharing unpublished results. We
acknowledge Drs. F. Pratto and V. Jayakumar for SSDS analysis. We are grateful
to Ms. S. Kondo for the generation of anti-SYCP3. We acknowledge Drs. S.
Keeney, Y. Fujiwara, and K. Ishiguro for *spo11* mice and Dr. H. Kurumizaka for
the RAD51 expression plasmid. We are also indebted to members of the
Shinohara lab, particularly Ms. S. Aoyama, A. Maeda, C. Watanabe, and S.
Hashimoto for mouse care. This work was supported by JSPS KAKENHI Grant
Numbers; 19H00981 to A.S., 20K15716 and 16H06279 (PAGS) to M.I., and
19H03157 to A.F.
967

Author contributions
M.I. and A.S. conceived and designed the experiments. M.I. performed all
experiments on mouse meiosis. M.I., Y.F., and A.T. analyzed SSDS data. A.F.
performed biochemical experiments. K.M. set up Fignl1 mice. M.I. and A.S.
analyzed the data. A.S. prepared the original draft and wrote the manuscript with
help from M.I.
974

975 **Competing interest**
976 The authors declare no competing interests.
977

References

1. Sonoda, E. *et al.* Rad51-deficient vertebrate cells accumulate chromosomal breaks prior to cell death. *EMBO J* **17**, 598-608 (1998).

2. Tashiro, S., Walter, J., Shinohara, A., Kamada, N. & Cremer, T. Rad51 accumulation at sites of DNA damage and in postreplicative chromatin. *J Cell Biol* **150**, 283-291 (2000).

3. Mason, J.M., Chan, Y.L., Weichselbaum, R.W. & Bishop, D.K. Non-enzymatic roles of human RAD51 at stalled replication forks. *Nat Commun* **10**, 4410 (2019).

4. Schlacher, K. *et al.* Double-strand break repair-independent role for BRCA2 in blocking stalled replication fork degradation by MRE11. *Cell* **145**, 529-542 (2011).

5. Ogawa, T., Yu, X., Shinohara, A. & Egelman, E.H. Similarity of the yeast RAD51 filament to the bacterial RecA filament. *Science* **259**, 1896-1899 (1993).

6. Shinohara, A., Ogawa, H. & Ogawa, T. Rad51 protein involved in repair and recombination in S. cerevisiae is a RecA-like protein. *Cell* **69**, 457-470 (1992).

7. Sung, P. Catalysis of ATP-dependent homologous DNA pairing and strand exchange by yeast RAD51 protein. *Science* **265**, 1241-1243 (1994).

8. Sung, P. & Robberson, D.L. DNA strand exchange mediated by a RAD51-ssDNA nucleoprotein filament with polarity opposite to that of RecA. *Cell* **82**, 453-461 (1995).

9. Jensen, R.B., Carreira, A. & Kowalczykowski, S.C. Purified human BRCA2 stimulates RAD51-mediated recombination. *Nature* **467**, 678-683 (2010).

10. Zhao, W. *et al.* BRCA1-BARD1 promotes RAD51-mediated homologous DNA pairing. *Nature* **550**, 360-365 (2017).

11. New, J.H., Sugiyama, T., Zaitseva, E. & Kowalczykowski, S.C. Rad52 protein stimulates DNA strand exchange by Rad51 and replication protein A. *Nature* **391**, 407-410 (1998).

12. Shinohara, A. & Ogawa, T. Stimulation by Rad52 of yeast Rad51-mediated recombination. *Nature* **391**, 404-407 (1998).

13. Van Dyck, E., Stasiak, A.Z., Stasiak, A. & West, S.C. Binding of double-strand breaks in DNA by human Rad52 protein. *Nature* **398**, 728-731 (1999).

14.   Akamatsu, Y., Dziadkowiec, D., Ikeguchi, M., Shinagawa, H. & Iwasaki, H.
Two different Swi5-containing protein complexes are involved in mating-
type switching and recombination repair in fission yeast. *Proc Natl Acad*
Sci U S A **100**, 15770-15775 (2003).

15.   Tsai, S.P. *et al.* Rad51 presynaptic filament stabilization function of the
mouse Swi5-Sfr1 heterodimeric complex. *Nucleic Acids Res* **40**, 6558-
6569 (2012).

16.   Masson, J.Y. *et al.* Identification and purification of two distinct complexes
containing the five RAD51 paralogs. *Genes Dev* **15**, 3296-3307 (2001).

17.   Abreu, C.M. *et al.* Shu complex SWS1-SWSAP1 promotes early steps in
mouse meiotic recombination. *Nat Commun* **9**, 3961 (2018).

18.   Liu, T., Wan, L., Wu, Y., Chen, J. & Huang, J. hSWS1.SWSAP1 is an
evolutionarily conserved complex required for efficient homologous
recombination repair. *J Biol Chem* **286**, 41758-41766 (2011).

19.   Martino, J. *et al.* The human Shu complex functions with PDS5B and
SPIDR to promote homologous recombination. *Nucleic Acids Res* **47**,
10151-10165 (2019).

20.   Tanaka, K., Kagawa, W., Kinebuchi, T., Kurumizaka, H. & Miyagawa, K.
Human Rad54B is a double-stranded DNA-dependent ATPase and has
biochemical properties different from its structural homolog in yeast,
Tid1/Rdh54. *Nucleic Acids Res* **30**, 1346-1353 (2002).

21.   Zhang, Z., Fan, H.Y., Goldman, J.A. & Kingston, R.E. Homology-driven
chromatin remodeling by human RAD54. *Nat Struct Mol Biol* **14**, 397-405
(2007).

22.   Liang, F. *et al.* Promotion of RAD51-Mediated Homologous DNA Pairing
by the RAD51AP1-UAF1 Complex. *Cell Rep* **15**, 2118-2126 (2016).

23.   Parplys, A.C. *et al.* NUCKS1 is a novel RAD51AP1 paralog important for
homologous recombination and genome stability. *Nucleic Acids Res* **43**,
9817-9834 (2015).

24.   Wiese, C. *et al.* Promotion of homologous recombination and genomic
stability by RAD51AP1 via RAD51 recombinase enhancement. *Mol Cell*
28, 482-490 (2007).

25.   Bugreev, D.V., Mazina, O.M. & Mazin, A.V. Bloom syndrome helicase
stimulates RAD51 DNA strand exchange activity through a novel
mechanism. *J Biol Chem* **284**, 26349-26359 (2009).

26.   Bugreev, D.V., Yu, X., Egelman, E.H. & Mazin, A.V. Novel pro- and anti-

recombination activities of the Bloom's syndrome helicase. *Genes Dev* **21**,
3085-3094 (2007).

27.   Xue, C. *et al.* Single-molecule visualization of human BLM helicase as it
acts upon double- and single-stranded DNA substrates. *Nucleic Acids Res*
47, 11225-11237 (2019).

28.   Sommers, J.A. *et al.* FANCJ uses its motor ATPase to destabilize protein-
DNA complexes, unwind triplexes, and inhibit RAD51 strand exchange. *J*
Biol Chem **284**, 7505-7517 (2009).

29.   Schwendener, S. *et al.* Physical interaction of RECQ5 helicase with
RAD51 facilitates its anti-recombinase activity. *J Biol Chem* **285**, 15739-
15745 (2010).

30.   Xue, C. *et al.* Single-molecule visualization of human RECQ5 interactions
with single-stranded DNA recombination intermediates. *Nucleic Acids Res*
49, 285-305 (2021).

31.   Chu, W.K. *et al.* FBH1 influences DNA replication fork stability and
homologous recombination through ubiquitylation of RAD51. *Nat Commun*
6, 5931 (2015).

32.   Simandlova, J. *et al.* FBH1 helicase disrupts RAD51 filaments in vitro and
modulates homologous recombination in mammalian cells. *J Biol Chem*
288, 34168-34180 (2013).

33.   Moldovan, G.L. *et al.* Inhibition of homologous recombination by the
PCNA-interacting protein PARI. *Mol Cell* **45**, 75-86 (2012).

34.   Croteau, D.L., Popuri, V., Opresko, P.L. & Bohr, V.A. Human RecQ
helicases in DNA repair, recombination, and replication. *Annu Rev*
Biochem **83**, 519-552 (2014).

35.   Heyer, W.D., Li, X., Rolfsmeier, M. & Zhang, X.P. Rad54: the Swiss Army
knife of homologous recombination? *Nucleic Acids Res* **34**, 4115-4125
(2006).

36.   Scully, R., Panday, A., Elango, R. & Willis, N.A. DNA double-strand break
repair-pathway choice in somatic mammalian cells. *Nat Rev Mol Cell Biol*
20, 698-714 (2019).

37.   Grey, C. & de Massy, B. Chromosome Organization in Early Meiotic
Prophase. *Front Cell Dev Biol* **9**, 688878 (2021).

38.   Ito, M. & Shinohara, A. Chromosome architecture and homologous
recombination in meiosis. *Front Cell Dev Biol* **10**, 1097446 (2022).

39.   Yadav, V.K. & Claeys Bouuaert, C. Mechanism and Control of Meiotic DNA

Double-Strand Break Formation in S. cerevisiae. *Front Cell Dev Biol* **9**,
642737 (2021).

40.   Bishop, D.K., Park, D., Xu, L. & Kleckner, N. DMC1: a meiosis-specific
yeast homolog of E. coli recA required for recombination, synaptonemal
complex formation, and cell cycle progression. *Cell* **69**, 439-456 (1992).

41.   Brown, M.S. & Bishop, D.K. DNA strand exchange and RecA homologs in
meiosis. *Cold Spring Harb Perspect Biol* **7**, a016659 (2014).

42.   Crickard, J.B. & Greene, E.C. The biochemistry of early meiotic
recombination intermediates. *Cell Cycle* **17**, 2520-2530 (2018).

43.   Deans, B., Griffin, C.S., Maconochie, M. & Thacker, J. Xrcc2 is required
for genetic stability, embryonic neurogenesis and viability in mice. *EMBO*
J **19**, 6675-6685 (2000).

44.   Kuznetsov, S.G., Haines, D.C., Martin, B.K. & Sharan, S.K. Loss of
Rad51c leads to embryonic lethality and modulation of Trp53-dependent
tumorigenesis in mice. *Cancer Res* **69**, 863-872 (2009).

45.   Lim, D.S. & Hasty, P. A mutation in mouse rad51 results in an early
embryonic lethal that is suppressed by a mutation in p53. *Mol Cell Biol* **16**,
7133-7143 (1996).

46.   Pittman, D.L. & Schimenti, J.C. Midgestation lethality in mice deficient for
the RecA-related gene, Rad51d/Rad51l3. *Genesis* **26**, 167-173 (2000).

47.   Prakash, R. *et al.* XRCC3 loss leads to midgestational embryonic lethality
in mice. *DNA Repair (Amst)* **108**, 103227 (2021).

48.   Shu, Z., Smith, S., Wang, L., Rice, M.C. & Kmiec, E.B. Disruption of
muREC2/RAD51L1 in mice results in early embryonic lethality which can
Be partially rescued in a p53(-/-) background. *Mol Cell Biol* **19**, 8686-8693
(1999).

49.   Tsuzuki, T. *et al.* Targeted disruption of the Rad51 gene leads to lethality
in embryonic mice. *Proc Natl Acad Sci U S A* **93**, 6236-6240 (1996).

50.   Huang, T. *et al.* SPIDR is required for homologous recombination during
mammalian meiosis. *Nucleic Acids Res* **51**, 3855-3868 (2023).

51.   Matsuzaki, K., Kondo, S., Ishikawa, T. & Shinohara, A. Human RAD51
paralogue SWSAP1 fosters RAD51 filament by regulating the anti-
recombinase FIGNL1 AAA+ ATPase. *Nat Commun* **10**, 1407 (2019).

52.   Prakash, R. *et al.* Distinct pathways of homologous recombination
controlled by the SWS1-SWSAP1-SPIDR complex. *Nat Commun* **12**, 4255
(2021).

53. Chester, N., Kuo, F., Kozak, C., O'Hara, C.D. & Leder, P. Stage-specific
apoptosis, developmental delay, and embryonic lethality in mice
homozygous for a targeted disruption in the murine Bloom's syndrome
gene. *Genes Dev* **12**, 3382-3393 (1998).

54. Wesoly, J. *et al.* Differential contributions of mammalian Rad54 paralogs
to recombination, DNA damage repair, and meiosis. *Mol Cell Biol* **26**, 976-
989 (2006).

55. McNally, F.J. & Roll-Mecak, A. Microtubule-severing enzymes: From
cellular functions to molecular mechanism. *J Cell Biol* **217**, 4057-4069
(2018).

56. Yuan, J. & Chen, J. FIGNL1-containing protein complex is required for
efficient homologous recombination repair. *Proc Natl Acad Sci U S A* **110**,
10640-10645 (2013).

57. Fernandes, J.B. *et al.* FIGL1 and its novel partner FLIP form a conserved
complex that regulates homologous recombination. *PLoS Genet* **14**,
e1007317 (2018).

58. Girard, C. *et al.* AAA-ATPase FIDGETIN-LIKE 1 and Helicase FANCM
Antagonize Meiotic Crossovers by Distinct Mechanisms. *PLoS Genet* **11**,
e1005369 (2015).

59. Hu, Q. *et al.* Meiotic Chromosome Association 1 Interacts with TOP3alpha
and Regulates Meiotic Recombination in Rice. *Plant Cell* **29**, 1697-1708
(2017).

60. Yang, S. *et al.* FIGNL1 Inhibits Non-homologous Chromosome
Association and Crossover Formation. *Front Plant Sci* **13**, 945893 (2022).

61. Kumar, R., Duhamel, M., Coutant, E., Ben-Nahia, E. & Mercier, R.
Antagonism between BRCA2 and FIGL1 regulates homologous
recombination. *Nucleic Acids Res* **47**, 5170-5180 (2019).

62. Sharan, S.K. *et al.* Embryonic lethality and radiation hypersensitivity
mediated by Rad51 in mice lacking Brca2. *Nature* **386**, 804-810 (1997).

63. Sadate-Ngatchou, P.I., Payne, C.J., Dearth, A.T. & Braun, R.E. Cre
recombinase activity specific to postnatal, premeiotic male germ cells in
transgenic mice. *Genesis* **46**, 738-742 (2008).

64. Yuan, L., Brundell, E. & Hoog, C. Expression of the meiosis-specific
synaptonemal complex protein 1 in a heterologous system results in the
formation of large protein structures. *Exp Cell Res* **229**, 272-275 (1996).

65. Yuan, L. *et al.* The murine SCP3 gene is required for synaptonemal

complex assembly, chromosome synapsis, and male fertility. *Mol Cell* **5**,
73-83 (2000).

66. Yuan, L. *et al.* The synaptonemal complex protein SCP3 can form
multistranded, cross-striated fibers in vivo. *J Cell Biol* **142**, 331-339 (1998).

67. Mahadevaiah, S.K. *et al.* Recombinational DNA double-strand breaks in
mice precede synapsis. *Nat Genet* **27**, 271-276 (2001).

68. Luo, M. *et al.* MEIOB exhibits single-stranded DNA-binding and
exonuclease activities and is essential for meiotic recombination. *Nat*
Commun **4**, 2788 (2013).

69. Souquet, B. *et al.* MEIOB targets single-strand DNA and is necessary for
meiotic recombination. *PLoS Genet* **9**, e1003784 (2013).

70. Yoshida, K. *et al.* The mouse RecA-like gene Dmc1 is required for
homologous chromosome synapsis during meiosis. *Mol Cell* **1**, 707-718
(1998).

71. Shi, B. *et al.* Dual functions for the ssDNA-binding protein RPA in meiotic
recombination. *PLoS Genet* **15**, e1007952 (2019).

72. Moens, P.B. *et al.* The time course and chromosomal localization of
recombination-related proteins at meiosis in the mouse are compatible
with models that can resolve the early DNA-DNA interactions without
reciprocal recombination. *J Cell Sci* **115**, 1611-1622 (2002).

73. Baker, S.M. *et al.* Involvement of mouse Mlh1 in DNA mismatch repair and
meiotic crossing over. *Nat Genet* **13**, 336-342 (1996).

74. Ashley, T. *et al.* Dynamic changes in Rad51 distribution on chromatin
during meiosis in male and female vertebrates. *Chromosoma* **104**, 19-28
(1995).

75. Ikeya, T., Shinohara, A., Sato, S., Tabata, S. & Ogawa, T. Localization of
mouse Rad51 and Lim15 proteins on meiotic chromosomes at late stages
of prophase 1. *Genes Cells* **1**, 379-389 (1996).

76. Tarsounas, M., Morita, T., Pearlman, R.E. & Moens, P.B. RAD51 and
DMC1 form mixed complexes associated with mouse meiotic
chromosome cores and synaptonemal complexes. *J Cell Biol* **147**, 207-
220 (1999).

77. Baudat, F., Manova, K., Yuen, J.P., Jasin, M. & Keeney, S. Chromosome
synapsis defects and sexually dimorphic meiotic progression in mice
lacking Spo11. *Mol Cell* **6**, 989-998 (2000).

78. Romanienko, P.J. & Camerini-Otero, R.D. The mouse Spo11 gene is

required for meiotic chromosome synapsis. *Mol Cell* **6**, 975-987 (2000).

79.   Zhang, Q. *et al.* The FLIP-FIGNL1 complex regulates the dissociation of
RAD51/DMC1 in homologous recombination and replication fork restart.
Nucleic Acids Res (2023).

80.   Hinch, A.G. *et al.* The Configuration of RPA, RAD51, and DMC1 Binding
in Meiosis Reveals the Nature of Critical Recombination Intermediates.
Mol Cell **79**, 689-701 e610 (2020).

81.   Khil, P.P., Smagulova, F., Brick, K.M., Camerini-Otero, R.D. & Petukhova,
G.V. Sensitive mapping of recombination hotspots using sequencing-
based detection of ssDNA. *Genome Res* **22**, 957-965 (2012).

82.   Smagulova, F. *et al.* Genome-wide analysis reveals novel molecular
features of mouse recombination hotspots. *Nature* **472**, 375-378 (2011).

83.   Altmannova, V. *et al.* The role of bivalent ions in the regulation of D-loop
extension mediated by DMC1 during meiotic recombination. *iScience* **25**,
105439 (2022).

84.   Li, X. & Heyer, W.D. RAD54 controls access to the invading 3'-OH end
after RAD51-mediated DNA strand invasion in homologous recombination
in Saccharomyces cerevisiae. *Nucleic Acids Res* **37**, 638-646 (2009).

85.   Li, X., Stith, C.M., Burgers, P.M. & Heyer, W.D. PCNA is required for
initiation of recombination-associated DNA synthesis by DNA polymerase
delta. *Mol Cell* **36**, 704-713 (2009).

86.   Bugreev, D.V. *et al.* The resistance of DMC1 D-loops to dissociation may
account for the DMC1 requirement in meiosis. *Nat Struct Mol Biol* **18**, 56-
60 (2011).

87.   Finsterbusch, F. *et al.* Alignment of Homologous Chromosomes and
Effective Repair of Programmed DNA Double-Strand Breaks during
Mouse Meiosis Require the Minichromosome Maintenance Domain
Containing 2 (MCMDC2) Protein. *PLoS Genet* **12**, e1006393 (2016).

88.   Her, C., Wu, X., Bailey, S.M. & Doggett, N.A. Mouse MutS homolog 4 is
predominantly expressed in testis and interacts with MutS homolog 5.
Mamm Genome **12**, 73-76 (2001).

89.   Kneitz, B. *et al.* MutS homolog 4 localization to meiotic chromosomes is
required for chromosome pairing during meiosis in male and female mice.
Genes Dev **14**, 1085-1097 (2000).

90.   Li, X.C. & Schimenti, J.C. Mouse pachytene checkpoint 2 (trip13) is
required for completing meiotic recombination but not synapsis. *PLoS*

Genet **3**, e130 (2007).

91. Lutzmann, M. *et al.* MCM8- and MCM9-deficient mice reveal
gametogenesis defects and genome instability due to impaired
homologous recombination. *Mol Cell* **47**, 523-534 (2012).

92. McNairn, A.J., Rinaldi, V.D. & Schimenti, J.C. Repair of Meiotic DNA
Breaks and Homolog Pairing in Mouse Meiosis Requires a
Minichromosome Maintenance (MCM) Paralog. *Genetics* **205**, 529-537
(2017).

93. Roig, I. *et al.* Mouse TRIP13/PCH2 is required for recombination and
normal higher-order chromosome structure during meiosis. *PLoS Genet* **6**
(2010).

94. Benson, F.E., Stasiak, A. & West, S.C. Purification and characterization of
the human Rad51 protein, an analogue of E. coli RecA. *EMBO J* **13**, 5764-
5771 (1994).

95. Dupaigne, P. *et al.* Rad51 polymerization reveals a new chromatin
remodeling mechanism. *PLoS One* **3**, e3643 (2008).

96. North, J.A. *et al.* ATP-dependent nucleosome unwrapping catalyzed by
human RAD51. *Nucleic Acids Res* **41**, 7302-7312 (2013).

97. Senavirathne, G., Mahto, S.K., Hanne, J., O'Brian, D. & Fishel, R. Dynamic
unwrapping of nucleosomes by HsRAD51 that includes sliding and
rotational motion of histone octamers. *Nucleic Acids Res* **45**, 685-698
(2017).

98. Holzen, T.M., Shah, P.P., Olivares, H.A. & Bishop, D.K. Tid1/Rdh54
promotes dissociation of Dmc1 from nonrecombinogenic sites on meiotic
chromatin. *Genes Dev* **20**, 2593-2604 (2006).

99. Shah, P.P. *et al.* Swi2/Snf2-related translocases prevent accumulation of
toxic Rad51 complexes during mitotic growth. *Mol Cell* **39**, 862-872 (2010).

100. Mason, J.M. *et al.* RAD54 family translocases counter genotoxic effects of
RAD51 in human tumor cells. *Nucleic Acids Res* **43**, 3180-3196 (2015).

101. Kiianitsa, K., Solinger, J.A. & Heyer, W.D. Rad54 protein exerts diverse
modes of ATPase activity on duplex DNA partially and fully covered with
Rad51 protein. *J Biol Chem* **277**, 46205-46215 (2002).

102. Kiianitsa, K., Solinger, J.A. & Heyer, W.D. Terminal association of Rad54
protein with the Rad51-dsDNA filament. *Proc Natl Acad Sci U S A* **103**,
9767-9772 (2006).

103. Solinger, J.A., Kiianitsa, K. & Heyer, W.D. Rad54, a Swi2/Snf2-like

recombinational repair protein, disassembles Rad51:dsDNA filaments.
Mol Cell **10**, 1175-1188 (2002).

104.   Tokuhiro, K. *et al.* Calreticulin is required for development of the cumulus
oocyte complex and female fertility. *Sci Rep* **5**, 14254 (2015).

105.   de Boer, E., Lhuissier, F.G. & Heyting, C. Cytological analysis of
interference in mouse meiosis. *Methods Mol Biol* **558**, 355-382 (2009).

106.   Peters, A.H., Plug, A.W., van Vugt, M.J. & de Boer, P. A drying-down
technique for the spreading of mammalian meiocytes from the male and
female germline. *Chromosome Res* **5**, 66-68 (1997).

107.   Brick, K., Pratto, F., Sun, C.Y., Camerini-Otero, R.D. & Petukhova, G.
Analysis of Meiotic Double-Strand Break Initiation in Mammals. *Methods*
Enzymol **601**, 391-418 (2018).

108.   Ramirez, F. *et al.* deepTools2: a next generation web server for deep-
sequencing data analysis. *Nucleic Acids Res* **44**, W160-165 (2016).

109.   Matsuo, Y., Sakane, I., Takizawa, Y., Takahashi, M. & Kurumizaka, H.
Roles of the human Rad51 L1 and L2 loops in DNA binding. *FEBS J* **273**,
3148-3159 (2006).

110.   Lange, J. *et al.* The Landscape of Mouse Meiotic Double-Strand Break
Formation, Processing, and Repair. *Cell* **167**, 695-708 e616 (2016).

111.   Boekhout, M. *et al.* REC114 Partner ANKRD31 Controls Number, Timing,
and Location of Meiotic DNA Breaks. *Mol Cell* **74**, 1053-1068 e1058 (2019).

1287

1288

 Figure legends

 Fig. 1. Defective spermatogenesis in *Fignl1* cKO mice.

 (a) Scheme of mouse *Fignl1 flox* and *Δ* alleles. Yellow and blue boxes and black
 triangles represent untranslated regions (UTRs), coding sequence (CDS), and
 loxP sequences, respectively.

 (b) Representative image of testes in *Fignl1$^{+/+}$ Stra8-Cre$^+$* (left) and *Fignl1$^{flox/Δ}$*
 Stra8-Cre$^+$ (right) mice.

 (c) Testis weights of mice with different combinations of genotypes. Adult testes
 (9-20 weeks) were analyzed. The red bars are means.

 (d) Immunoblotting of FIGNL1 from whole-testis extracts of mice with different
 genotypes. α-tubulin is a loading control.

 (e) Sperm numbers in *Fignl1$^{+/+}$ Stra8-Cre$^+$* (black circles) and *Fignl1$^{flox/Δ}$ Stra8-*
 Cre$^+$ (blue open circles) mice. Sperms from adult Cauda epididymis (11-17
 weeks) were counted. The red bars are means.

 (f) Representative images of seminiferous tubule sections from the testis of
 Fignl1$^{+/+}$ Stra8-Cre$^+$ (Ctrl) and *Fignl1$^{flox/Δ}$ Stra8-Cre$^+$* (*Fignl1* cKO) mice. The
 sections were stained with periodic acid-Schiff (PAS). Sg, spermatogonia; Sc (L),
 leptotene spermatocytes; Sc (P), pachytene spermatocytes; St, spermatids.

 (g) Distribution of different meiotic prophase I stages in Ctrl (gray bars) and *Fignl1*
 cKO (blue open bars). >100 cells were counted from each animal. The red bars
 are SDs from three animals of each genotype.

 (h) Distribution of FIGNL1-positive and -negative cells at different meiotic
 prophase I stages in Ctrl (gray bars) and *Fignl1* cKO (blue open bars). Squashed
 testicular cells were immunostained for FIGNL1, γH2AX, and SYCP3. >100
 leptotene/zygotene cells (with pan-nuclear γH2AX staining) and
 pachytene/diplotene cells (with the sex body-specific γH2AX staining) were
 analyzed from each animal. The means of two mice of each genotype are
 indicated. Lept., leptonema; Zyg., zygonema; Pach., pachynema; dipl.,
 diplonema. Representative images of testicular cell squashes are shown in
 Supplementary Fig. 1b.

 (i) Representative images of spermatocyte-chromosome spreads immunostained
 for SYCP1 (green) and SYCP3 (magenta) at zygonema in Ctrl and *Fignl1* cKO.
 The bottom panels are magnified images of regions with yellow dotted rectangles.

 (j) Quantification of the frequency of synapsis defects in Ctrl (gray bars) and
 Fignl1 cKO (blue open bars). >90 cells were analyzed from each animal. The red

bars are SDs from three mice of each genotype.

In (f)-(i), genotypes of indicated animals are Ctrl, *Fignl1$^{+/+}$ Stra8-Cre$^+$*; *Fignl1* cKO,

Fignl1$^{flox/\Delta}$ Stra8-Cre$^+$. The results of unpaired *t*-tests are indicated in the graphs:

****$p \leq 0.0001$.

Scale bars, 50 μm in (f) and 10 μm for whole-nucleus images and 2 μm for

magnified images in (i).

1331

Fig. 2. Deregulated meiotic recombination in *Fignl1* cKO spermatocytes.

(a) Representative images of spermatocyte-chromosome spreads

immunostained for RPA2 (white in the top panels and green in the bottom panels)

and SYCP3 (magenta) at indicated meiotic prophase I stages in Ctrl and *Fignl1*

cKO.

(b) Quantification of RPA2 focus numbers at different meiotic prophase I stages

in Ctrl (black circles) and *Fignl1* cKO (blue open circles). The red bars are means.

(c) Representative images of spermatocyte-chromosome spreads

immunostained for MSH4 (white in the left panels and green in the right panels),

SYCP1 (magenta) and SYCP3 (blue) at mid-zygonema in Ctrl and *Fignl1* cKO.

The bottom panels are magnified images of regions with dotted yellow.

(d) Quantification of MSH4 focus numbers at mid-zygonema in Ctrl (black circles)

and *Fignl1* cKO (blue open circles). The red bars are means.

(e) The correlation between MSH4 focus number and total SC length at mid-

zygonema in Ctrl (black circles) and *Fignl1* cKO (blue open circles). Pearson's *r*

is shown in parentheses. A black solid line and a blue dashed line indicate linear

regression for Ctrl and *Fignl1* cKO, respectively.

(f) Quantification of MSH4 focus density on the SCs at mid-zygonema in Ctrl

(black circles) and *Fignl1* cKO (blue open circles). The red bars are means. In (e)

and (f), mid-zygotene nuclei with 40-120 μm of total SC length per nucleus were

analyzed.

Genotypes of indicated animals are: Ctrl, *Fignl1$^{+/+}$ Stra8-Cre$^+$*; *Fignl1* cKO,

Fignl1$^{flox/\Delta}$ Stra8-Cre$^+$. The results of the Mann-Whitney *U*-test are indicated in

the graphs: ****$p \leq 0.0001$. The total number of cells analyzed is indicated below

the graphs.

Scale bars in (a) and (c), 10 μm and 2 μm for magnified images in (c).

1358

Fig. 3. Accumulation of RAD51 and DMC1 in meiotic prophase I in *Fignl1*

cKO spermatocytes.

(a) Representative images of spermatocyte-chromosome spreads
immunostained for DMC1 (white in the top panels and green in the bottom panels)
and SYCP3 (magenta) at indicated meiotic prophase I stages in Ctrl and *Fignl1*
cKO.
(b) Quantification of DMC1 focus numbers at different meiotic prophase I stages
in Ctrl (black circles) and *Fignl1* cKO (blue open circles). The red bars are means.
(c) Representative images of spermatocyte-chromosome spreads
immunostained for RAD51 (white in the top panels and green in the bottom
panels) and SYCP3 (magenta) at indicated meiotic prophase I stages in Ctrl and
Fignl1 cKO.
(d) Quantification of RAD51 focus numbers at different meiotic prophase I stages
in Ctrl (black circles) and *Fignl1* cKO (blue open circles). The red bars are means.
(e) Quantification of focus numbers of RAD51 (green), DMC1 (magenta), and
RAD51-DMC1 colocalization (brown) at different meiotic prophase I stages in Ctrl
(filled circles) and *Fignl1* cKO (open circles). The black bars are means.
Representative images of spermatocyte-chromosome spreads are shown in
Supplementary Fig. 7.
(f) Quantification of the frequency of RAD51-DMC1 colocalization at different
meiotic prophase I stages in Ctrl (filled circles) and *Fignl1* cKO (open circles). The
degrees of RAD51 foci colocalizing with DMC1 (left, green) and DMC1 foci
colocalizing with RAD51 (right, magenta) are shown. The black bars are means.
Genotypes of indicated animals are Ctrl, *Fignl1$^{+/+}$ Stra8-Cre^{+}*; *Fignl1* cKO,
Fignl1$^{flox/\Delta}$ Stra8-Cre^{+}. The results of the Mann-Whitney *U*-test are indicated in
the graphs: ****$p \le 0.0001$. The total number of cells analyzed is indicated below
the graphs.
Scale bars in (a) and (c), 10 μm. Prelept., preleptonema; lept., leptonema; zyg.,
zygonema.
1388
Fig. 4. Accumulation of RAD51 and DMC1 in preleptotene *Fignl1* cKO
spermatocytes.
(a) Representative images of spermatocyte-chromosome spreads
immunostained for RAD51 (left), SYCP3 (middle) and EdU (right) at
preleptonema in Ctrl and *Fignl1* cKO.
(b) Quantification of RAD51 focus numbers at preleptonema in Ctrl (black circles)
and *Fignl1* cKO (blue open circles). The red solid bars and dashed bars are
means for total cells and RAD51-positive cells (with >10 RAD51 foci; 26/38 and

25/48 cells in early and mid/late preleptonema, respectively), respectively.

(c)   Representative   images   of   spermatocyte-chromosome   spreads
immunostained for DMC1 (left), SYCP3 (middle), and EdU (right) at
preleptonema in Ctrl and *Fignl1* cKO.

(d) Quantification of DMC1 focus numbers at preleptonema in Ctrl (black circles)
and *Fignl1* cKO (blue open circles). The red solid bars and dashed bars are
means for total cells and DMC1-positive cells (with >30 DMC1 foci; 28/39 and
26/45 cells in early and mid/late preleptonema, respectively), respectively.

Genotypes of indicated animals are Ctrl, *Fignl1*$^{+/+}$ *Stra8-Cre*$^+$; *Fignl1* cKO,
Fignl1$^{flox/\Delta}$ *Stra8-Cre*$^+$. The results of the Mann-Whitney *U*-test are indicated in
the graphs: ****$p \leq 0.0001$. The total number of cells analyzed is indicated below
the graphs.

Scale bars in (a) and (c), 10 μm.

1410

Fig. 5. SPO11-independent accumulation of RAD51 and DMC1 in *Fignl1* cKO
spermatocytes.

(a)   Representative   images   of   spermatocyte-chromosome   spreads
immunostained for RAD51 (white in the top panels and green in the bottom
panels) and SYCP3 (magenta) at leptonema in *Spo11* KO, *Fignl1* cKO and *Spo11*
KO *Fignl1* cKO.

(b) Quantification of RAD51 focus numbers at different meiotic prophase I stages
in *Spo11* KO (orange circles), *Fignl1* cKO (blue open circles), and *Spo11* KO
Fignl1 cKO (purple open circles). The red bars are means. Cells in which Stra8-
Cre recombinase seemed to fail excision of the *Fignl1*$^\Delta$ allele in *Spo11* KO *Fignl1*
cKO (<30 RAD51 foci) are excluded. Quantification including these cells is shown
in Supplementary Fig. 8a.

(c)   Representative   images   of   spermatocyte-chromosome   spreads
immunostained for DMC1 (white in the top panels and green in the bottom panels)
and SYCP3 (magenta) at early zygonema in *Spo11* KO, *Fignl1* cKO and *Spo11*
KO *Fignl1* cKO.

(d) Quantification of DMC1 focus numbers at different meiotic prophase I stages
in *Spo11* KO (orange circles), *Fignl1* cKO (blue open circles), and *Spo11* KO
Fignl1 cKO (purple open circles). The red bars are means. Cells in which Stra8-
Cre recombinase seemed to fail excision of the *Fignl1*$^\Delta$ allele in *Spo11* KO *Fignl1*
cKO (<60 DMC1 foci) are excluded. Quantification including these cells is shown
in Supplementary Fig. 8b.

(e) Representative images of spermatocyte-chromosome spreads
immunostained for RAD51 (green), DMC1 (magenta), and SYCP3 (blue) at late
zygonema/early leptonema in *Spo11* KO *Fignl1* cKO. The bottom panels are
magnified images of regions with yellow dotted rectangles.
(f) Quantification of focus numbers of RAD51 (green open circles), DMC1
(magenta open circles), and RAD51-DMC1 colocalization (brawn open circles) at
different meiotic prophase I stages in *Spo11* KO *Fignl1* cKO. The black bars are
means.
(g) Quantification of the frequency of RAD51-DMC1 colocalization at different
meiotic prophase I stages in *Spo11* KO *Fignl1* cKO. The degrees of RAD51 foci
colocalizing with DMC1 (left, green open circles) and DMC1 foci colocalizing with
RAD51 (right, magenta open circles) are shown. The black bars are means.
Genotypes of indicated animals are: *Spo11* KO, *Spo11$^{-/-}$ Fignl1$^{flox/\Delta}$*; *Fignl1* cKO,
Fignl1$^{flox/\Delta}$ Stra8-Cre$^+$; *Spo11* KO *Fignl1* cKO, *Spo11$^{-/-}$ Fignl1$^{flox/\Delta}$ Stra8-Cre$^+$*. The
results of the Mann-Whitney *U*-test are indicated in the graphs: ****$p \leq 0.0001$.
The total number of cells analyzed is indicated below the graphs.
Scale bars in (a), (c) and (e), 10 μm and 2 μm for magnified images in (e). Prelept.,
preleptonema; lept., leptonema; zyg., zygonema.
1451
Fig. 6. Persistent RAD51 binding at recombination sites in *Fignl1* cKO
spermatocytes.
(a) RAD51 ChIP-SSDS signals (green) in Ctrl and *Fignl1* cKO with Spo11-oligo
counts (blue) and DMC1 ChIP-SSDS signals (black) in wild-type at an indicated
chromosomal region with a hotspot on mouse chromosome 2.
(b) The heatmap of RAD51 ChIP-SSDS signals (green) in Ctrl and *Fignl1* cKO
and Spo11-oligo counts (blue) and DMC1 ChIP-SSDS signals (black) in wild-type.
The 4,000 most active meiotic recombination hotspots on autosomes identified
by Spo11-oligo in wild-type are ordered by SPO11-oligos counts [110]. ChIP-SSDS
signals are centered relative to hotspot centers. In (a) and (b), Spo11-oligo and
DMC1 ChIP-SSDS data are from previous studies [110, 111].
(c) Representative images of spermatocyte-chromosome spreads
immunostained for RAD51 (green), RPA2 (magenta), and SYCP3 (blue) at mid-
zygonema in Ctrl and *Fignl1* cKO. The bottom panels are magnified images of
regions with yellow dotted rectangles. Scale bars, 10 μm for whole-nucleus
images and 2 μm for magnified images.
(d) Quantification of focus numbers of RAD51 (green), RPA2 (magenta) and

RAD51-RPA2 colocalization (brawn) at mid zygonema in Ctrl (filled circles) and
Fignl1 cKO (open circles). The black bars are means.
(e) Quantification of the frequency of RAD51-RPA2 colocalization at mid-
zygonema in Ctrl (black circles) and *Fignl1* cKO (blue open circles). The degree
of RPA2 foci colocalizing with RAD51 is shown. The red bars are means.
Genotypes of indicated animals are: Ctrl, *Fignl1$^{+/+}$ Stra8-Cre$^+$*; *Fignl1* cKO,
Fignl1$^{flox/Δ}$ Stra8-Cre$^+$. The results of the Mann-Whitney *U*-test are indicated: ****p
≤ 0.0001. The total number of cells analyzed is indicated below the graphs.
1477
Fig. 7. *Fignl1* deficiency rescues defects in RAD51 and DMC1 assembly
caused by *Swsap1* deficiency.
(a) Representative images of spermatocyte-chromosome spreads
immunostained for RAD51 (white in the top panels and green in the bottom
panels) and SYCP3 (magenta) at early zygonema in Ctrl, *Swsap1* KO*, Fignl1*
cKO and *Swsap1* KO *Fignl1* cKO.
(b) Quantification of RAD51 focus numbers at different meiotic prophase I stages
in Ctrl (black circles), *Swsap1* KO (orange circles), *Fignl1* cKO (blue open circles),
and *Swsap1* KO *Fignl1* cKO (purple open circles). The red bars are means.
(c) Representative images of spermatocyte-chromosome spreads
immunostained for DMC1 (white in the top panels and green in the bottom panels)
and SYCP3 (magenta) at early zygonema in Ctrl, *Swsap1* KO*, Fignl1* cKO, and
Swsap1 KO *Fignl1* cKO.
(d) Quantification of DMC1 focus numbers at different meiotic prophase I stages
in Ctrl (black circles), *Swsap1* KO (orange circles), *Fignl1* cKO (blue open circles),
and *Swsap1* KO *Fignl1* cKO (purple open circles). The red bars are means.
Genotypes of indicated animals are *Swsap1* KO, *Swsap1$^{-/-}$ Fignl1$^{flox/Δ}$*; *Fignl1*
cKO, *Fignl1$^{flox/Δ}$ Stra8-Cre$^+$*; *Swsap1* KO *Fignl1* cKO, *Swsap1$^{-/-}$ Fignl1$^{flox/Δ}$ Stra8-*
Cre$^+$. The results of the Mann-Whitney *U*-test are indicated in the graphs: ****p ≤
0.0001. The total number of cells analyzed is indicated below the graphs.
Scale bars in (a) and (c), 10 μm.
1499
Fig. 8. FIGNL1 disrupts RAD51-DNA complexes.
(a) Electrophoresis mobility shift assay showing *in vitro* disruption of RAD51-DNA
complexes by FIGNL1. A representative gel image among triplicate is shown.
Human RAD51 protein was incubated with either Cy5-labelled ssDNA (40nt) or
dsDNA (40bp) in the presence of ATP. After the formation of the protein-DNA

complex, purified human FINGL1ΔN was added at the indicated concentration
and further incubated. The complexes were analyzed by polyacrylamide gel
electrophoresis and the fluorescence from the DNAs was captured.
(b) Quantification of free ssDNA or dsDNA in (A). The red bars are means and
SDs from three independent experiments.
(c) Model of FIGNL1 function in spermatocytes. In pre-meiotic S-phase and early
meiotic prophase I, FIGNL1 dissociates RAD51 and DMC1 from dsDNAs. At
meiotic DSB sites, FIGNL1 removes RAD51 and DMC1 from ssDNAs, which
facilitates efficient loading of MutSγ (MSH4-MSH5) and further processing of
homologous recombination to crossover outcome.
1515
1516 **Supplementary information**
1517
1518 **Supplementary Figures legends**
1519
Supplementary Fig. 1. Depletion of FIGNL1 from early meiotic prophase I
spermatocytes, related to Fig. 1.
(a) Representative images of testicular cell squashes immunostained for FIGNL1
(green), PLZF (top, magenta) and SYCP3 (bottom, magenta), counterstained for
DNA with DAPI (blue). Cells in yellow dotted circles with condensed DAPI bodies
are Sertoli cells or round spermatids. Scale bars, 10 μm.
(b) Representative images of testicular cell squashes immunostained for FIGNL1
(green), γH2AX (blue), and SYCP3 (magenta) in Ctrl (*Fignl1*$^{+/+}$ *Stra8-Cre*$^+$) and
Fignl1 cKO (*Fignl1*$^{flox/Δ}$ *Stra8-Cre*$^+$). SYCP3-positive prophase I spermatocytes
were categorized into two by γH2AX staining: leptotene/zygotene cells (L/Z) and
pachytene/diplotene cells (P/D) show pan-nuclear γH2AX staining and the sex
body-specific γH2AX staining, respectively. Scale bars, 10 μm.
1532
Supplementary Fig. 2. Deregulated meiotic recombination in *Fignl1* cKO
spermatocytes, related to Fig. 2.
(a) Representative images of spermatocyte-chromosome spreads
immunostained for γH2AX (white in the top panels and green in the bottom
panels) and SYCP3 (magenta) at indicated meiotic prophase I stages in Ctrl and
Fignl1 cKO.
(b) Quantification of γH2AX intensity at different meiotic prophase I stages in Ctrl
(black circles) and *Fignl1* cKO (blue open circles). Two independent experiments

showed reduced (left; pooled results from two animals of each genotype) and one

experiment showed similar (right) levels of γH2AX intensity in *Fignl1* cKO

spermatocytes compared to Ctrl. The red bars are means. The results of the

Mann-Whitney *U*-tests are indicated: ****$p \leq 0.0001$. The total number of cells

analyzed is indicated below the graphs.

(c)   Representative   images   of   spermatocyte-chromosome   spreads

immunostained for MLH1 (white in the top panels and green in the bottom panels),

SYCP1 (magenta), and SYCP3 (blue) at late pachynema in Ctrl and late

zygonema (pachynema-like) in *Fignl1* cKO. The right panels are magnified

images of regions with yellow dotted rectangles.

Genotypes of indicated animals are Ctrl, *Fignl1*[+/+] *Stra8-Cre*[+]; *Fignl1* cKO,

Fignl1[flox/Δ] *Stra8-Cre*[+].

Scale bars in (a) and (c), 10 μm and 2 μm for magnified images in (c).

1554

Supplementary Fig. 3. Reduced focus intensity of RAD51 and DMC1 in

***Fignl1* cKO spermatocytes, related to Fig. 3.**

(a) Quantification of focus intensities of RAD51 at different meiotic prophase I

stages in Ctrl (black circles) and *Fignl1* cKO (blue open circles). The red bars are

means ± SDs. The total number of cells analyzed is 35 and 26 at leptonema and

early zygonema in Ctrl, respectively; 54 and 30 at leptonema and early zygonema

in *Fignl1* cKO, respectively.

(b) Quantification of focus intensities of DMC1 at different meiotic prophase I

stages in Ctrl (black circles) and *Fignl1* cKO (blue open circles). The red bars are

means ± SDs. The total number of cells analyzed is 64 and 22 at leptonema and

early zygonema in Ctrl, respectively; 61 and 33 at leptonema and early zygonema

in *Fignl1* cKO, respectively.

Genotypes of indicated animals are Ctrl, *Fignl1*[+/+] *Stra8-Cre*[+]; *Fignl1* cKO,

Fignl1[flox/Δ] *Stra8-Cre*[+]. The results of the Mann-Whitney *U*-test are indicated in

the graphs: ****$p \leq 0.0001$. The total number of foci analyzed is indicated below

the graphs.

1571

Supplementary Fig. 4. Infrequent localization of RAD51 at ongoing

replication fork in preleptotene *Fignl1* cKO spermatocytes, related to Fig. 4.

(a)   Representative   images   of   spermatocyte-chromosome   spreads

immunostained for IdU (green), RAD51 (magenta), and SYCP3 at early

preleptonema in Ctrl and *Fignl1* cKO. The bottom panels are magnified images

of regions with yellow dotted rectangles. Scale bars, 10 μm for whole-nucleus
images and 2 μm for magnified images.
(b) Quantification of focus numbers of IdU (green open circles), RAD51 (magenta
open circles), and IdU-RAD51 colocalization (brawn open circles) at early
preleptonema in *Fignl1* cKO. The black bars are means.
(c) Quantification of the frequency of IdU-RAD51 colocalization at early
preleptonema in *Fignl1* cKO. The degrees of IdU foci colocalizing with RAD51
(left, green open circles) and RAD51 foci colocalizing with IdU (right, magenta
open circles) are shown. The black bars are means. The colocalization frequency
was measured with and without rotating the IdU image 90° clockwise relative to
the RAD51 image.
(d) Comparison of the frequency of IdU-RAD51 (10' IdU labeling) and EdU-
RAD51 (60' EdU labeling) colocalization at early preleptonema in *Fignl1* cKO.
The degrees of RAD51 foci colocalizing with IdU (left, magenta open circles) and
RAD51 foci colocalizing with EdU (right, magenta open triangles) are shown. The
black bars are means.
Genotypes of indicated animals are Ctrl, *Fignl1$^{+/+}$ Stra8-Cre$^+$*; *Fignl1* cKO,
Fignl1$^{flox/Δ}$ Stra8-Cre$^+$. The results of the Mann-Whitney *U*-test are indicated in
the graphs: ****$p \leq 0.0001$. The total number of cells analyzed is indicated below
the graphs.
1597
Supplementary Fig. 5. Infrequent localization of DMC1 at ongoing
replication fork in preleptotene *Fignl1* cKO spermatocytes, related to Fig. 4.
(a)   Representative   images   of   spermatocyte-chromosome   spreads
immunostained for EdU (green), DMC1 (magenta), and SYCP3 at early
preleptonema in Ctrl and *Fignl1* cKO. The bottom panels are magnified images
of regions with yellow dotted rectangles. Scale bars, 10 μm for whole-nucleus
images and 2 μm for magnified images.
(b) Quantification of focus numbers of EdU (green open circles), DMC1 (magenta
open circles), and EdU-DMC1 colocalization (brawn open circles) at early
preleptonema in *Fignl1* cKO. The black bars are means.
(c) Quantification of the frequency of EdU-DMC1 colocalization at early
preleptonema in *Fignl1* cKO. The degrees of EdU foci colocalizing with DMC1
(left, green open circles) and DMC1 foci colocalizing with EdU (right, magenta
open circles) are shown. The black bars are means. The colocalization frequency
was measured with and without rotating the EdU image 90° clockwise relative to

the DMC1 image.

Genotypes of indicated animals are Ctrl, *Fignl1$^{+/+}$ Stra8-Cre^{+}*; *Fignl1* cKO,

Fignl1$^{flox/Δ}$ Stra8-Cre^{+}. The results of the Mann-Whitney *U*-test are indicated in

the graphs. The total number of cells analyzed is indicated below the graphs.

1617

Supplementary Fig. 6. Largely normal pre-meiotic DNA replication in *Fignl1*

cKO spermatocytes, related to Fig. 4.

(a) Representative images of spermatocyte-chromosome spreads

immunostained for RPA2 (left), SYCP3 (middle) and EdU (right) at preleptonema

in Ctrl and *Fignl1* cKO. Scale bars, 10 μm.

(b) Quantification of RPA2 focus numbers at early preleptonema in Ctrl (black

circles) and *Fignl1* cKO (blue open circles). The red bars are means.

(c) Quantification of IdU focus numbers (10' IdU labeling) at early preleptonema

in Ctrl (black circles) and *Fignl1* cKO (blue open circles). The red bars are means.

Genotypes of indicated animals are Ctrl, *Fignl1$^{+/+}$ Stra8-Cre^{+}*; *Fignl1* cKO,

Fignl1$^{flox/Δ}$ Stra8-Cre^{+}. The results of the Mann-Whitney *U*-test are indicated in

the graphs. The total number of cells analyzed is indicated below the graphs.

1630

Supplementary Fig. 7. Uncoupled loading of RAD51 and DMC1 in *Fignl1*

cKO spermatocytes, related to Fig. 3 and 4.

Representative images of spermatocyte-chromosome spreads immunostained

for RAD51 (green), DMC1 (magenta), and SYCP3 (blue) at indicated meiotic

prophase I stages in Ctrl (*Fignl1$^{+/+}$ Stra8-Cre^{+}*) and *Fignl1* cKO (*Fignl1$^{flox/Δ}$ Stra8-*

Cre^{+}). The bottom panels are magnified images of regions with yellow dotted

rectangles. Scale bars, 10 μm for whole-nucleus images and 2 μm for magnified

images.

1639

Supplementary Fig. 8. Accumulation of RAD51 and DMC1 with low levels of

γH2AX and few RPA2 foci in *Spo11* KO *Fignl1* cKO spermatocytes, related

to Fig. 5.

(a) Quantification of RAD51 focus numbers at different meiotic prophase I stages

in *Spo11* KO (orange circles), *Fignl1* cKO (blue open circles), and *Spo11* KO

Fignl1 cKO (purple open circles). The red bars are means.

(b) Quantification of DMC1 focus numbers at different meiotic prophase I stages

in *Spo11* KO (orange circles), *Fignl1* cKO (blue open circles), and *Spo11* KO

Fignl1 cKO (purple open circles). The red bars are means.

(c) Representative images of spermatocyte-chromosome spreads
immunostained for γH2AX (green), SYCP3 (magenta), and RAD51 (white in
insets) at indicated meiotic prophase I stages in *Fignl1* cKO, *Spo11* KO and
Spo11 KO *Fignl1* cKO. Scale bars, 10 μm.
(d) Quantification of γH2AX intensity at different meiotic prophase I stages in
Fignl1 cKO (blue open circles), *Spo11* KO (orange circles), and *Spo11* KO *Fignl1*
cKO (purple open circles). The red bars are means.
(e) Representative images of spermatocyte-chromosome spreads
immunostained for RPA2 (green) and SYCP3 (magenta) at early leptonema in
Spo11 KO and *Spo11* KO *Fignl1* cKO. Scale bars, 10 μm.
(f) Quantification of RPA2 focus numbers at early leptonema in *Spo11* KO
(orange circles) and *Spo11* KO *Fignl1* cKO (purple open circles). The red bars
are means.
Genotypes of indicated animals are: *Spo11* KO, *Spo11$^{-/-}$ Fignl1$^{flox/\Delta}$*; *Fignl1* cKO,
Fignl1$^{flox/\Delta}$ Stra8-Cre$^+$; *Spo11* KO *Fignl1* cKO, *Spo11$^{-/-}$ Fignl1$^{flox/\Delta}$ Stra8-Cre$^+$*. The
results of the Mann-Whitney *U*-test are indicated in the graphs: ****$p \leq 0.0001$.
The total number of cells analyzed is indicated below the graphs.
1666

Supplementary Fig. 9. RAD51 binding at recombination sites in *Fignl1* cKO
spermatocytes, related to Fig. 6.
(a) RAD51 ChIP-SSDS signals (green) in Ctrl and *Fignl1* cKO with Spo11-oligo
counts (blue) and DMC1 ChIP-SSDS signals (black) in wild-type at an indicated
chromosomal region with multiple hotspots on mouse chromosome 3.
(b) The heatmap of RAD51 ChIP-SSDS signals (green) in Ctrl and *Fignl1* cKO
and Spo11-oligo counts (blue) and DMC1 ChIP-SSDS signals (black) in wild-type,
as shown in Fig. 6b. Three mice of each genotype were analyzed and two RAD51
antibodies with different lot numbers were used for ChIP. The results of Ctrl #1
and *Fignl1* cKO #1 are shown in Fig. 6a, b, Supplementary Fig. 9a.
(c) Comparison of RAD51 ChIP-SSDS signals at meiotic recombination hotspots.
Each spot represents the sum of RAD51 ChIP-SSDS signals around ±2,000 bp
of one of the 4,000 most active meiotic recombination hotspots on autosomes
(black) used for heatmap representation in Fig. 6b and Supplementary Fig. 9b
and 163 active hotspots on sex chromosomes (magenta) with similar Spo11-oligo
counts as the 4,000 active autosomal hotspots in wild-type [110]. Pearson's *r* for
autosomal hotspots is also shown.
(d) Metaplots of RAD51 ChIP-SSDS signals around meiotic recombination

hotspots. Smoothed RAD51 ChIP-SSDS signals in Ctrl (top) and *Fignl1* cKO
(bottom) around ±2,000 bp of the 4,000 most active meiotic recombination
hotspots on autosomes used for heatmap representation are shown. Three mice
of each genotype were analyzed. Black, orange and blue lines represent the
results of #1, #2 and #3, respectively.
Genotypes of indicated animals are: Ctrl, *Fignl1$^{+/+}$ Stra8-Cre$^+$*; *Fignl1* cKO,
Fignl1$^{flox/Δ}$ Stra8-Cre$^+$.
1692

Supplementary Fig. 10. RAD51 binding in *Spo11* KO and *Spo11* KO *Fignl1*
cKO spermatocytes, related to Fig. 6.
(a) The heatmap of RAD51 ChIP-SSDS signals (green) in *Spo11* KO and *Spo11*
KO *Fignl1* cKO and Spo11-oligo counts (blue) and DMC1 ChIP-SSDS signals
(black) in wild-type, as shown in Fig. 6b. Two mice of each genotype were
analyzed.
(b) Metaplots of RAD51 ChIP-SSDS signals around meiotic recombination
hotspots. Smoothed RAD51 ChIP-SSDS signals in *Spo11* KO (left) and *Spo11*
KO *Fignl1* cKO (right) around ±2,000 bp of the 4,000 most active meiotic
recombination hotspots on autosomes used for heatmap representation are
shown. Two mice of each genotype were analyzed. Black and blue lines
represent the results of #1 and #2, respectively.
(c) Comparison of RAD51 ChIP-SSDS profiles in the entire genome. Mouse
genome was divided into non-overlapping 2-kb bins and top 1,000 bins with the
highest RAD51 ChIP-SSDS signals were analyzed. The 1,000 bins were either
categorized into three: high (>50), middle (50-20) and low (<20), based on sum
RAD51 ChIP-SSDS signals in each bin (left) or two: overlapping with previously
identified DMC1 ChIP-SSDS hotspots [1] and non-overlapping (right). The
frequency of each category was plotted for each animal of Ctrl (black open
circles), *Fignl1* cKO (blue open circles), *Spo11* KO (orange circles), and *Spo11*
KO *Fignl1* cKO (purple open circles). The means of three (Ctrl and *Fignl1* cKO)
or two (*Spo11* KO and *Spo11* KO *Fignl1* cKO) mice of each genotype are
indicated.
(d) Comparison of RAD51 ChIP-SSDS profiles in *Spo11* KO and *Spo11* KO
Fignl1 cKO. 262 bins with high or middle RAD51 ChIP-SSDS signals in both two
of *Spo11* KO *Fignl1* cKO mice in (c) were categorized into high, middle, and low
RAD51 ChIP SSDS signals in either of two *Spo11* KO mice by the same criteria
as (c).

Genotypes of indicated animals are: Ctrl, *Fignl1*$^{+/+}$ *Stra8-Cre*$^+$; *Fignl1* cKO,
Fignl1$^{flox/\Delta}$ *Stra8-Cre*$^+$; *Spo11* KO, *Spo11*$^{-/-}$ *Stra8-Cre*$^+$; *Spo11* KO *Fignl1* cKO,
Spo11$^{-/-}$ *Fignl1*$^{flox/\Delta}$ *Stra8-Cre*$^+$.
1724

1725 **Supplementary Table 1. Primers used for genotyping in this study**

Primer	Sequence (5'->3')
Fignl1 wild-type and flox allele, forward	TATCCATATTGGGTGTCTCCTGTTAC
Fignl1 wild-type and flox allele, reverse	TAGCACACACACACTGGATATAACT
Fignl1 deletion allele, forward	CCGAGTATCCATATTGGGTG
Fignl1 deletion allele, reverse	TGTCTGCTGTAGTCATGAGG
Stra8-Cre transgenic allele, forward	CTCCAAGGGGGTAAGGTGTAGC
Stra8-Cre transgenic allele, reverse	TTCCAGGGCGCGAGTTGATAG
Swsap1 wild-type allele, forward	GAGATATCGCTTTTGATTGGTC
Swsap1 wild-type allele, reverse	GAATTGAGGATGGGTACTGG
Swsap1 deletion allele, forward	CGGTTTCCATATGGGGATTGGTGGCGACGA
Swsap1 deletion allele, reverse	TGCCTTATTTCCTGATCCAGGCTAGCTGTC
Spo11 wild-type allele, forward	TGAGATACATGGAGGAAGATGG
Spo11 deletion allele, forward	CTGAGCCCAGAAAGCGAAGGAA
Spo11, common reverse	ATGTTAGTCGGCACAGCAGTAG

1740

1741 **Supplementary Table 2. Antibodies used in this study**

Antibody	Source	Identifier	Application	Dilution
Rat anti-SYCP3 (Serum)	this study	N/A	IF	1:5,000
Mouse anti-SYCP3	Abcam	ab97672	IF	1:750
Rabbit anti-SYCP1	Novus Biologicals	NB300-229	IF	1:500
Mouse anti-MLH1	BD Pharmingen	51-1327QR	IF	1:50
Rabbit anti-MSH4	Abcam	ab58666	IF	1:200
Rabbit anti-RPA2	Abcam	ab76420	IF	1:200
Rat anti-RPA2	Cell Signaling	2208	IF	1:100
Mouse anti-γH2AX	Millipore	05-636	IF	1:500
Rabbit anti-DMC1	Santa Cruz	sc-22768	IF	1:200
Rabbit anti-RAD51	Santa Cruz	sc-8349	IF	1:100
Mouse anti-RAD51	Novus Biologicals	NB100-148	IF, ChIP	1:100
Mouse anti-BrdU	BD Biosciences	BD347580	IF	1:200
Mouse anti-PLZF	Santa Cruz	sc-28319	IF	1:100
Rabbit anti-FIGNL1	Proteintech	17604-1-AP	IF	1:100

1757				IB	1:1,000
1758	Mouse anti-α tubulin	SIGMA	T6074	IB	1:3,000
1759	Goat anti-mouse 488	Molecular Probes	A-11029	IF	1:1,000
1760	Goat anti-rabbit 488	Molecular Probes	A-11034	IF	1:1,000
1761	Goat anti-rat 488	Molecular Probes	A-11006	IF	1:1,000
1762	Goat anti-mouse 594	Molecular Probes	A-11032	IF	1:1,000
1763	Goat anti-rat 594	Molecular Probes	A-11007	IF	1:1,000
1764	Goat anti-rabbit 647	Molecular Probes	A-21245	IF	1:500
1765	Goat anti-rabbit IgG-HRP	Southern Biotech	4030-05	IB	1:10,000
1766	Goat anti-mouse IgG-HRP	Thermo Scientific	31430	IB	1:10.000

1767 IF, immunofluorescence; IB, immunoblotting.; ChIP, chromatin immunoprecipitation

1768

Supplementary References

1770

1771 1. Brick, K., Smagulova, F., Khil, P., Camerini-Otero, R.D. & Petukhova, G.V.
1772 Genetic recombination is directed away from functional genomic elements
1773 in mice. *Nature* **485**, 642-645 (2012).

1774

Ito et al., Figure 1

a *Fignl1*
flox — Exon 1, Exon 2, Exon 3
Δ

b *Fignl1* +/+ flox/Δ
Stra8-Cre + +
5 mm

c
Testis weight (as % body weight)
****
****
****
n= 6 6 7 7
Fignl1 +/+ flox/Δ +/+ flox/Δ
Stra8-Cre − +

d *Fignl1* +/+ flox/Δ flox/Δ
Stra8-Cre + − +
75 — FIGNL1
50 — α-tubulin

e
Sperm per mouse (× 10^6)
p= 0.0008
n= 3 3
Fignl1 +/+ flox/Δ
Stra8-Cre +

f
Stra8-Cre⁺ Fignl1^{+/+} (Ctrl)
Stra8-Cre⁺ Fignl1^{flox/Δ} (*Fignl1* cKO)
Sg
St
Sc (L)
Sc (P)
Sg
Sc (L)
St

g
Percent of prophase cells
■ Ctrl □ *Fignl1* cKO
Leptonema, Zygonema, Pachynema, Diplonema

h
Percent of FIGNL1-positive cells
■ Ctrl □ *Fignl1* cKO
Lept./zyg., Pach./dipl.

i
Ctrl / *Fignl1* cKO
SYCP1 SYCP3
Zygonema
ii i
Non-homologous synapsis
Multi-chromosomal synapsis

j
Percent of zygotene cells with defective synapsis
■ Ctrl □ *Fignl1* cKO
****
p= 0.0214
Non-homologous, Multi-chromosomal

a

c Eary preleptonema

b

RAD51 foci per nucleus

- Ctrl
- *Fignl1* cKO

p= 0.0007 *p*= 0.0085

n= 25 38 45 48
Early Mid/late
Preleptonema

d

DMC1 foci per nucleus

- Ctrl
- *Fignl1* cKO

**** *p*=0.0003

n= 30 39 27 45
Early Mid/late
Preleptonema

Ito et al., Figure 6

a

dsDNA (40 bp)   ssDNA (40 nt)

RAD51   − +   +   +   − +   +   +
FIGNL1ΔN   − 0 0.6 1.2   − 0 0.6 1.2 (μM)

well

RAD51-DNA

dsDNA
ssDNA

b

% of unbound DNAs

FIGNL1ΔN   − 0 0.6 1.2   − 0 0.6 1.2
RAD51 (μM) − +   +   +   − +   +   +

dsDNA (40 bp)   ssDNA (40 nt)

c

Fignl1+/+

Fignl1Δ/Δ

Fignl1+/+

Fignl1Δ/Δ

Crossover

● RAD51
● DMC1
● RPA

⚡ Meiotic DSB

⬭ MutSγ

● MutLγ

a

b

a

b

c

a Early preleptonema

SYCP3 | IdU | RAD51 | IdU RAD51

Ctrl

Fignl1 cKO

b
Foci per nucleus

IdU RAD51 Colocalization
n= 38 38 38

Early preleptonema

c
Percent of IdU foci colocalizing with RAD51

p= 0.0261

n= 21 21

Early prelept.
Early prelept. 90° turn

Percent of RAD51 foci colocalizing with IdU

p= 0.0439

n= 21 21

Early prelept.
Early prelept. 90° turn

d
Percent of RAD51 foci colocalizing with EdU/IdU

****

n= 38 26

10' IdU
60' EdU

Early preleptonema

a

Early preleptonema

Fignl1 cKO

SYCP3 | EdU | DMC1 | EdU DMC1

b

Foci per nucleus

EdU DMC1 Colocalization

n= 16 16 16

Early preleptonema

c

Percent of EdU foci colocalizing with DMC1

p= 0.0087

n= 16 16

Early prelept.
Early prelept. 90° turn

Percent of DMC1 foci colocalizing with EdU

p= 0.0002

n= 16 16

Early prelept.
Early prelept. 90° turn

a Early preleptonema

Ctrl — RPA2 | SYCP3 | EdU

Fignl1 cKO — RPA2 | SYCP3 | EdU

b
● Ctrl
○ *Fignl1* cKO

p= 0.9414

RPA2 foci per nucleus

n= 21   19

Early preleptonema

c
● Ctrl
○ *Fignl1* cKO

p= 0.4652

IdU foci per nucleus

n= 26   38

Early preleptonema

a

Fignl1	+/+	flox/Δ	flox/Δ		+/+	flox/Δ	flox/Δ
Stra8-Cre	+	-	+		+	-	+

b

Fignl1	+/+	flox/Δ	flox/Δ		+/+	flox/Δ	flox/Δ
Stra8-Cre	+	-	+		+	-	+

c

	dsDNA (40 bp)				ssDNA (40 nt)			
RAD51	-	+	+	+	-	+	+	+
FIGNL1ΔN	-	0	0.6	1.2	-	0	0.6	1.2	(μM)

well

RAD51-DNA

dsDNA
ssDNA

(a) An uncropped immunoblotting image for FIGNL1 shown in Fig. 1d.
(b) An uncropped immunoblotting image for α-tubulin shown in Fig. 1d.
(c) An unprocessed image of electrophoresis mobility shift assay shown in Fig. 8a.