The genomics and evolution of inter-sexual mimicry and female-limited polymorphisms in damselflies

B. Willink[1,2*], K. Tunström[1], S. Nilén[3], R. Chikhi[4], T. Lemane[5], M. Takahashi[6†], Y. Takahashi[7],

E. I. Svensson[3], C. W. Wheat[1]

[1] Department of Zoology, Stockholm University, Stockholm 106-91, Sweden

[2] Department of Biological Sciences, National University of Singapore, Singapore 117558,

Singapore

[*] Corresponding author email: bwillink@gmail.com

[3] Department of Biology, Evolutionary Ecology Unit, Lund University, Lund 223-62, Sweden

[4] Institut Pasteur, Université Paris Cité, Sequence Bioinformatics, F-75015 Paris, France

[5] Univ. Rennes, Inria, CNRS, IRISA, Rennes F-35000, France

[6] Graduate School of Life Sciences, Tohoku University, 6-3 Aramaki, Aoba, Sendai, 980-8578,

Japan

[7] Graduate School of Science, Chiba University, 1-33 Yayoi, Inage, Chiba, 263-8522, Japan

[†] Current address: Graduate School of Agriculture, Kyoto University, Oiwakecho,

Kitashirakawa, Sakyoku, Kyoto, 606-8502, Japan

Keywords: balancing selection, ectopic recombination, insertion, *k*-mer GWAS, structural

variant, transposable elements, trans-species polymorphism

Abstract

Sex-limited morphs can provide profound insights into the evolution and genomic architecture of

complex phenotypes. Inter-sexual mimicry is one particular type of sex-limited polymorphism in

which a novel morph resembles the opposite sex. While inter-sexual mimics are known in both

sexes and a diverse range of animals, their evolutionary origin is poorly understood. Here, we

investigated the genomic basis of female-limited morphs and male mimicry in the Common

Bluetail damselfly. Differential gene expression between morphs has been documented in

damselflies, but no causal locus has been previously identified. We found that male-mimicry

originated in an ancestrally sexually-dimorphic lineage in association with multiple structural

changes, probably driven by transposable element activity. These changes resulted in ~900 kb of

novel genomic content that is partly shared by male mimics in a close relative, indicating that

male mimicry is a trans-species polymorphism. More recently, a third morph originated

following the translocation of part of the male-mimicry sequence into a genomic position ~3.5

mb apart. We provide evidence of balancing selection maintaining male-mimicry, in line with

previous field population studies. Our results underscore how structural variants affecting a

handful of potentially regulatory genes and morph-specific genes, can give rise to novel and

complex phenotypic polymorphisms.

MAIN

Sexual dimorphism is one of the most fascinating forms of intra-specific phenotypic variation in

animals. Sexes often differ in size and colour, as well as the presence of elaborated ornaments

and weaponry. Theoretical and empirical studies over many decades have developed a detailed

framework of sexual selection and sexual conflict, explaining why these differences arise and

how they become encoded in sex differentiation systems[1–3]. However, a growing number of

examples of inter-sexual mimicry[4–7] suggest that sexual dimorphism can be evolutionarily fragile

and quite dynamic. Inter-sexual mimicry has evolved in several lineages, when individuals of

one sex gain a fitness advantage, usually frequency- or density-dependent, due to their

resemblance to the opposite sex. For example, males who mimic females, as seen in the Ruff

(*Calidris pugnax*) and the Melanzona Guppy (*Poecilia parae*), forgo courtship and 'sneak'

copulations from dominant males[4,5], while females who mimic males, in damselflies and

hummingbirds, avoid excessive male-mating harassment[6,8]. Inter-sexual mimicry thus requires

the evolution of a novel sex-mimicking morph in a sexually-dimorphic ancestor. The occurrence

of inter-sexual mimicry may be a intermediate step in the evolution of sexual monomorphism, it

may be an ephemeral state, or it may be maintained as a stable polymorphism. In any case,

sexual mimics harbour genetic changes that attenuate or prevent the development of sex-specific

phenotypes, and can therefore provide insights into the essential building blocks of sexual

dimorphism[9].

Considerable research effort has been devoted to uncover the genetic basis of discrete phenotypic

polymorphisms, such as those associated with alternative reproductive or life-history strategies[10–

14]. Together, these studies highlight a vast diversity of mechanisms used by evolution to package

complex phenotypic differences into a single locus that is protected from the eroding effects of

recombination. On one extreme, phenotypic morphs may evolve via massive insertions,

deletions, or inversions that lock together dozens to hundreds of genes into supergenes[15–17]. On

the other end, much smaller structural variants (SVs), confined to a few thousand base pairs, can

modulate the expression of one or a few regulators of pleiotropic networks, resulting in markedly

different morphs[11,12,18]. We are clearly only starting to get a glimpse of the major themes among

these genetic mechanisms. For example, it is not known whether genomic architecture

determines the type and breadth of co-varying traits or the likelihood of polymorphisms evolving

in specific lineages[19].

A few of these studies have focused on sex-limited polymorphisms, where one of the morphs

shares the overall appearance, such as the colour pattern, of the opposite sex[10,14,20]. Such sex-

limited morphs may illustrate novel origins of sexual dimorphism, driven by either sexual

selection in males[14] or natural selection in females[18,21]. Alternatively, sex-limited polymorphisms

may arise with the evolution of inter-sexual mimicry. Crucially, empirical support for the

evolution of inter-sexual mimicry demands both a macroevolutionary context for the

polymorphism, showing that sexually dimorphism is ancestral, and a documented advantage of

sexual mimics in at least some social contexts. There is therefore a need to integrate genomic,

microevolutionary and phylogenetic evidence into our understanding of the evolutionary

dynamics of sexual dimorphism and inter-sexual mimicry. This integrative approach has been

overall rare, and applied mostly to the study of alternative male reproductive strategies[18,22]. Yet,

female mimicry of males may be more common than historically appreciated[23], and the genetic

basis of such mimicry remains largely unexplored[24–26].

The Common Bluetail damselfly *Ischnura elegans* (Odonata) has three female-limited morphs

(namely *O*, *A* and *I*) that differ in colouration, whereas males are always monomorphic[27]. *O*

females display the colour pattern and developmental colour changes inferred as ancestral in a

comparative analysis of the genus *Ischnura*[28] (Fig. 1). Male-like (*A*) females are considered

male mimics, who experience a frequency-dependent advantage of reduced male-mating and

premating harassment due to their resemblance to males[6]. Finally, the *I* morph shares its stripe

pattern and immature colouration with the *A* morph[27] (Fig. 1), but develops a yellow-brown

background colouration with age, eventually resembling the *O* morph upon sexual maturation[29]. *I*

females are only known in *I. elegans* and a few close relatives[28] (Fig. 1), and their evolutionary

relationship to *A* and *O* females remains unresolved. The behaviour, ecology, and population

biology of *I. elegans* have been intensely investigated for over two decades, making it one of the

best understood female-limited polymorphisms, in terms of how morphs differ in fitness-related

traits and how alternative morphs are maintained sympatrically over long periods[30–33].

Nonetheless, the molecular basis of this polymorphism remains unknown.

To advance our understanding of the evolution of complex phenotypes, such as sexual

dimorphism and sex-specific morphs, we identify the genomic region responsible for the female-

limited colour polymorphism in *I. elegans*. Using a combination of reference-based and

reference-free genome wide association studies (GWAS), upon morph-specific genome

assemblies, we revealed two novel regions adding up to ~900 kb, that are associated with the

evolutionary origin of the male-mimicking *A* morph. These structural variants, probably

generated and expanded by transposable element (TE) activity, are partly shared by male-

mimicking females of the Tropical Bluetail damselfly (*Ischnura senegalensis*), indicating that

male mimicry is a trans-species polymorphism. We also show that the novel *I* morph evolved via

an ectopic recombination event, where part of the *A*-unique genomic content was translocated

into an *O* genomic background. Finally, we examined the evolutionary dynamics of the colour

morph locus and explored expression patterns of genes located in this region. Together, our

results indicate that structural variation affecting a handful of genes and maintained by balancing

selection provides the raw material for the evolution of a male-mimicking phenotype in pond

damselflies.

RESULTS

Male mimicry is encoded by a locus with a signature of balancing selection

We started by conducting three reference-based GWAS, comparing all morphs against each

other in a pairwise fashion (Extended Data Fig. 1). We used an *A* morph genome assembly

(Supporting Text 1) as mapping reference because structural variant analyses revealed that *A*

females harbour genomic content that is absent in the other two morphs (see *Female morphs*

differ in genomic content below). The draft assembly was scaffolded against the Darwin Tree of

Life (DToL) reference genome to place the contigs in a chromosome level framework[34]. The

DToL reference genome contains the *O* allele (see Supporting Text 2) and is assembled with

chromosome resolution, except for chromosome 13, which is fragmented and consists of one

main and several unlocalized scaffolds.

All pairwise GWAS between morphs pointed to one and the same unlocalized scaffold of

chromosome 13 as the causal morph locus (Fig. 2a). Closer examination of this scaffold revealed

two windows of elevated divergence between morphs (Fig. 2b). First, a narrow region near the

start of the scaffold (~50 kb - 0.2 mb) captures highly significant SNPs in both *A* vs *O* and *I* vs *O*

comparisons (Fig. 2b). Thereafter and up to ~1.5 mb, an abundance of SNPs differentiates *A*

females from both *O* and *I* females, especially between ~0.6 and ~1.0 mb (Fig. 2b). These results

are mirrored by Fst values across both regions (Fig. 2c).

Next, we investigated whether the morph locus carries a signature of balancing selection, as

suggested by previous field studies of morph-frequency dynamics[31]. The larger genomic window

that uniquely distinguishes *A* females from both *I* and *O* females displays a signature of

balancing selection, indicated by highly positive values of Tajima's D, exceeding the 95

percentile of genome-wide estimates (Fig. 2d). Conversely, values of both Tajimas's D and π in

the narrower window that differentiates *O* females from both *A* and *I* females (~50 kb - 0.2 mb)

fall within the 95 percentile of genome-wide estimates (Fig. 2d-e).

Female morphs differ in genomic content

Previous studies have found that complex phenotypic polymorphisms are often underpinned by

structural variants (SVs), arising from genomic rearrangements such as insertions, deletions and

inversions[10,13,15,20]. As these variants can be difficult to detect in a reference-based analysis, we

employed a *k*-mer based GWAS approach[35] (Extended Data Fig. 1), which enables reference-free

identification of genomic divergence between morphs. Significant *k*-mers in these analyses could

represent regions that are present in one morph and absent in the other (i.e. insertions or

deletions), or regions that are highly divergent in their sequence (as in a traditional GWAS).

First, we investigated the divergence associated with the male-mimicking *A* morph. Pairwise

analyses revealed 568,039 and 508,031 *k*-mers (length = 31 bp) significantly associated with the

A vs *O* and *A* vs *I* comparisons, respectively. To determine whether the associated *k*-mers

represent differences in genomic content or sequence between the morphs, we mapped these *k*-

mers to morph-specific reference genomes. If the associated *k*-mers are due to novel sequences

found in one morph but not the other, we would expect a vast majority of the significant *k*-mers

to be found in only one of the two morphs in a pairwise comparison. If the significant *k*-mers are

instead owed to point mutations in high-identity sequences, there should be morph-specific k-

mers in both morphs.

Most ($> 98\%$) of the mapped k-mers in the A vs O and A vs I comparisons, aligned perfectly to a

single ~1.5 mb region of the unlocalized scaffold 2 of chromosome 13, in the A-morph assembly

(Fig. 3a; Extended Data Table 1). This is the same region of the A-morph assembly that was

previously identified in the standard GWAS (Fig. 2). In contrast, only ~0.3% of the associated k-

mers in the A vs O comparison were found anywhere in the O-assembly, and similarly, only

~0.2% of the significant k-mers in the A vs I analysis mapped to the I assembly (Extended Data

Table 1). These results thus suggested that a large region of genomic content is unique to the A

haplotype.

Given that A and I females share their immature colour pattern[29,36], we then tested for k-mer

associations that would distinguish both A and I females from O females and found 85,134 such

k-mers (Extended Data Table 1). When mapped to the A assembly, a majority of these k-mers

were found near the start of the unlocalized scaffold 2 of chromosome 13 (Fig. 3a), where we

previously reported pronounced divergence of O females (Fig. 2b-c). However, when mapped to

the I assembly, most of the significant k-mers were found in a different region of the same

scaffold, separated by approximately 3.5 mb (Fig. 3b). These results thus suggested that A and I

females share genomic content that is absent in O. However, in the I haplotype this content

occupies a different chromosomal location.

To further investigate the distribution of genomic content among morphs, we plotted the

standardized number of mapped reads (read depths) along the ~1.5 mb region of the A assembly

that included most of the significant k-mers (Extended Data Fig. 1). Here, we expected read

depth values around 0.5 (heterozygous) or 1.0 (homozygous) for all A samples, whereas I and O

samples should have read depths of 0, if genomic content is uniquely present in the *A* allele

(because *I* and *O* individuals lack the *A* allele, Fig. 1). Read depths confirmed that male-

mimicking *A* females are differentiated by genomic content. Specifically, there are two windows

of the *A* assembly (of ~400 kb and ~500 kb) where no *I* or *O* data maps to the assembly after

filtering repetitive sequences (Fig. 3c), and which are therefore uniquely present in *A* females.

These two windows of *A*-specific content are separated by a region between ~0.6 and ~1.0 mb

that is shared among all morphs (Fig. 3c), and highly divergent in SNP-based comparisons

involving the *A* morph (Fig. 2b). Finally, the region including most significant *k*-mers in the *A*

and *I* vs. *O* comparison is present in all *A* and *I* samples but absent in all *O* samples, except for

one individual (Fig. 3c; Supporting Text 3). As noted in the *k*-mer GWAS, this region of

genomic content shared by *A* and *I* individuals is located in different regions, separated by ~3.5

mb, in the two assemblies (Fig. 3d).

By combining reference-based GWAS, reference-free GWAS and read-depth approaches, we

have identified three haplotypes controlling morph development in the Common Bluetail. The *A*

and *I* haplotypes share ~150 kb that are absent in *O*. The *A* haplotype has two additional

windows of unique genomic content, adding up to ~900 kb. In the *A* haplotype, a single ~1.5 mb

window (hereafter the morph locus) thus contains the regions of unique genomic content, the

region exclusively shared between *A* and *I*, and the SNP-rich region present in all morphs. In the

I haplotype the region exclusively shared with *A* occupies a single and different locus separated

by about 3.5 mb (Fig. 4a). These large and compounded differences in genomic content between

haplotypes suggest that multiple structural changes on a multi-million base-pair region were

responsible for the evolution of novel female morphs in *Ischnura* damselflies.

TE propagation and recombination likely explain the origins of novel female morphs

Based on previous inferences of the historical order in which female morphs evolved (Fig. 1), we

hypothesized that genomic divergence first occurred between *O* and *A* females, with some

genomic content being then translocated from *A* into an *O* background, leading to the

evolutionary origin of *I* females. We analyzed structural variants between morphs to test this

hypothesis (Extended Data Fig. 1; Supporting Text 4), and uncovered evidence of a ~20 kb

sequence in the *O* haplotype that is duplicated and inverted in tandem in derived morphs (*A* and

I; Fig. 4b; Extended Data Fig. 2). An investigation of the reads mapping to the inversion

breakpoints suggested that additional duplications in the *A* genome, presumably via TE

proliferation, may be related to the evolution of inter-sexual mimicry (Fig. 4b; Extended Data

Fig. 3). Interestingly, TE content is enriched and recombination is reduced not just in the vicinity

of the morph locus, but across the entire chromosome 13 (Extended Data Fig. 4-5; Supporting

Text 4). Finally, evidence of a translocation of an *A*-derived genomic region back into an *O*

background (Extended Data Fig. 6; Supporting Text 4) implied that the *I* morph evolved from an

ectopic recombination event between between *A* and *O* morphs (Fig. 4b). This scenario is also

consistent with our previous *k*-mer GWAS and read-depth results, where we found that the only

region differentiating both *A* and *I* females from *O* females is located ~3.5 mb in the *I* haplotype.

Male mimicry is a trans-species polymorphism

Ancestral state reconstruction of female colour states had previously pointed to an ancient origin

of male mimicry in the clade that includes *I. elegans* and several other widely-distributed

Ischnura damselflies[28] (Fig. 1). We investigated whether male mimicry is in fact a trans-species

polymorphism using *de novo* genome assemblies from the closely related Tropical Bluetail

(*Ischnura senegalensis*) (Extended Data Fig. 1). *I. senegalensis* shares a common ancestor with *I.*

elegans about 5 Ma[28], and has both a male-mimicking *A* morph and a non-mimicking morph,

which resembles the *O* females of *I. elegans*[28,37] (Fig. 5a).

We reasoned that if morph divergence is ancestral, the genomic content that is uniquely present

in *A* females or shared by *A* and *I* females in *I. elegans* should be at least partly present in *A*

females of *I. senegalensis*, but absent in the alternative *O*-like female morph (see Supporting

Text 5). This prediction was supported by differences in standardized read depths between the *A*

and *O*-like pool of *I. senegalensis*, specifically at the morph locus of *I. elegans* (Fig. 5b;

Supporting Text 5). A shared genomic basis of inter-sexual mimicry for the two species was also

supported by the same ~20 kb inversion signature in the *A* pool against an *O* assembly, as

detected in *A* and *I* females of *I. elegans* (Extended Data Fig. 7). Finally, assembly alignments

between *O*-like and *A* haplotypes of *I. senegalensis* showed that the *A*-specific genomic region of

I. elegans is partly present in the *A* but not the *O*-like assembly of *I. senegalensis* (Fig. 5c).

Shared and morph-specific genes reside in the morph locus

Finally, we examined gene content and expression patterns in the morph locus. As female

morphs differ in genomic content as well as sequence, the phenotypic effects of the morph locus

could come about in at least three non-exclusive ways. First, entire gene models may be present

in some morphs and absent in others. Second, genes present in all morphs may differ in

expression patterns. Third, genes may encode different amino acid sequences in different female

morphs. We used newly generated and previously published[38] RNAseq data to investigate these

questions (Extended Data Fig. 1), and capitalized on the annotations of the reference genome of

I. elegans[34], as well as transcripts assembled *de novo* in our *A*-morph genome assembly.

Because the genetic basis of inter-sexual mimicry is shared between *I. elegans* and *I.*

senegalensis (Fig. 5), we focus on genes that are expressed in both species in at least one

individual (Fig. 6a).

Three transcripts (from two predicted genes) in the morph locus are expressed in *A* females of *I.*

senegalensis, and in *A* and *I* females of *I. elegans*, but never in *O* or *O*-like females (Fig. 6b).

Only one of these gene models (Afem.4094) could be functionally annotated, and appears to

encode a Long Interspaced Nuclear Element (LINE) retrotransposon in the clade Jockey

(Supporting Text 6). This gene also exhibited expression changes in *I* females that reflect their

colour development trajectory of initial resemblance to *A* females, followed by an overall

appearance similar to *O* females upon sexual maturation (Supporting Text 6). Notably,

RepeatModeler and *RepeatMasker* detected signatures of the Jockey family at the same locus as

the mapping locations of the *A* reads that had suggested a propagation of TEs in our SV analyses

(Fig. 6a; Extended Data Fig. 3). Thus, these results further support that TEs are responsible for

the evolution and expansion of the male-mimicry allele.

We also identified three gene models that are shared by all haplotypes and expressed in both

species. The three predicted genes encode zinc-finger domain proteins (Fig. 6b; Supporting Text

6), which are known to participate in transcriptional regulation[39]. However, we found no

conclusive evidence of differential expression, nor evidence of non-synonymous substitutions

between morphs shared by both *I. elegans* and *I. senegalensis* (Supporting Text 6). While we see

genes of a potentially regulatory function reside in the morph locus, understanding their role in

morph development will likely require higher temporal and spatial resolution of gene expression

data.

DISCUSSION

Sexual dimorphism, where males and females have markedly distinct colour patterns, has led to

multiple evolutionary origins of female-limited polymorphisms and potential male-mimicry in

Ischnura damselflies[28]. Here, we present a first genomic glance into how these morphs evolve,

setting the stage for future functional work to unravel the reversal of sexual phenotypes in

damselfly sexual mimicry. Male mimicry in the Common Bluetail is controlled by a single

genomic region in chromosome 13 (Fig. 2; 3). Our data suggests that this morph locus likely

evolved with the accumulation of novel and potentially TE-derived sequences in the male

mimicry haplotype (Fig. 4), which is shared by male-mimicking females of species diverging

more than 5 Ma (Fig. 5). More recently, a rare recombination event involving part of the novel *A*

genomic content has triggered the origin of a third female morph (Fig. 4), which shares its

sexually immature colouration and patterning with *A* females, and shares its sexually mature

overall appearance with *O* females[27]. The morph locus contains a handful of genes, some of

which may have evolved with TE propagation in the *A* haplotype, and are therefore absent from

O individuals (Fig. 6). However, existing annotations provide only a hint on how these genes

may influence morph development. Our results thus echo recent calls for a broader application of

functional validation tools, in order to understand how lineage-specific genes contribute to

phenotypic variation in natural populations[40].

This study underscores two increasingly recognized insights in evolutionary genomics. First,

there is mounting evidence that structural variants abound in natural populations and often

underpin complex and ecologically relevant phenotypic variation[41], such as discrete phenotypic

polymorphisms[10,13,15,20]. Nonetheless, traditional GWAS approaches based on SNPs can easily

miss structural variants, as these approaches are contingent on the genomic content of the

reference assembly[42]. Among other novel approaches to tackle this problem[42], a reference-free *k*-

mer based GWAS, as implemented here, is a powerful method to identify variation in genomic

content and sequence, especially when the genomic architecture of the trait of interest is initially

unknown[35]. In this study, we did not know *a priori* which of the three morphs, if any, would

harbour unique genomic content. Had we ignored differences in genomic content between

morphs and based our GWAS analysis solely upon the DToL (*O*) reference assembly, we would

have failed to identify SNPs between *I* and *O* morphs (Extended Data Fig. 8), and the origin of *I*

females via a translocation of *A* content would have been obscured.

Second, a role for TEs in creating novel and even adaptive phenotypic variation is increasingly

being recognized[43,44]. Here, we found that a ~400 kb region of unique genomic content, possibly

driven by LINE transposition is associated with the male-mimicry phenotype in at least two

species of *Ischnura* damselflies. TE activity can contribute to phenotypic evolution by multiple

mechanisms. For instance, TEs may modify the regulatory environment of genes in their vicinity,

by altering methylation[45] and chromatin conformation patterns[46], or by providing novel cis-

regulatory elements[47]. The male-mimicry region in *I. elegans* is located between two coding

genes with putative DNA-binding domains, and which may thus act as transcription factors.

However, our expression data does not provide unequivocal support for differential regulation of

either of these genes between female morphs. Importantly, currently available expression data

comes from adult specimens, as female morphs are not visually discernible in aquatic nymphs.

Yet, the key developmental differences that produce the adult morphs are likely directed by

regulatory variation during earlier developmental stages. Now that the morph locus has been

identified, future work can address differential gene expression at more relevant developmental

stages, before colour differences between morphs become apparent.

TEs can also contribute to phenotypic evolution if they become domesticated, for example, when

TE-encoded proteins are remodeled through evolutionary change to perform adaptive host

functions[48]. We found two transcripts located in *A* specific or *A/I* specific regions that are likely

derived from LINE retrotransposons and are actively expressed in the genomes that harbour

them (Fig. 6b). It is therefore possible that these transcripts participate in the development of

adult colour patterns, which are initially more similar between *A* and *I* females than between

either of these morphs and *O* females[27,29]. Yet, functional work on these transcripts is required to

ascertain their role in morph determination. Finally, TEs can become sources of novel small

regulatory RNAs which play important regulatory roles[49], including in insect sex

determination[50]. Thus, future work should also address non-coding RNA expression and function

in the morph locus.

Our results also provide molecular evidence for previous insights, gained by alternative research

approaches, on the micro- and macroevolution of female-limited colour polymorphisms. A

wealth of population data in Southern Sweden has shown that female-morph frequencies are

maintained by balancing selection, as they fluctuate less than expected due to genetic drift[31].

Behavioural and field experimental studies indicate that such balancing selection on female

morphs is mediated by negative frequency-dependent male harassment[51,52]. We add to these

earlier results, by showing a molecular signature consistent with balancing selection in the

genomic region where *A* females differ from both of the non-mimicking morphs. Sexual conflict

is expected to have profound effects on genome evolution, but there are few examples of

sexually-antagonistic traits with a known genetic basis, in which predictions about these genomic

effects can be tested[53,54]. Here, the signature of balancing selection on the morph locus matches

the expectation of inter-sexual conflict resulting in negative-frequency dependent selection and

maintaining alternative morph alleles over long periods.

Similarly, a recent comparative study based on phenotypic and phylogenetic data inferred a

single evolutionary origin of the male-mimicking morph shared by *I. elegans* and *I.*

senegalensis[28]. Our present results strongly support this common origin. This is because *A*

females in both species share unique genomic content, including associated transcripts, and an

inversion signature against the ancestral *O* morph (Fig. 5; Extended Data Fig. 7). Nonetheless,

these data are consistent with both an ancestral polymorphism and ancestral introgression being

responsible for the spread of male mimicry across the clade. A potential role for introgression in

the evolution of male mimicry is also suggested by rampant hybridization between *I. elegans* and

its closest relatives[55], and by the fact that *I. elegans* and *I. senegalensis* can hybridize millions of

years after their divergence, at least in laboratory settings[56]. The identification of the morph locus

in *I. elegans*, enables future comparative genomics studies to disentangle the relative roles of

long-term balancing selection and introgression in shaping the widespread phylogenetic

distribution of female-limited polymorphisms in *Ischnura* damselflies.

Finally, our results open up new lines of inquiry on how the genomic architecture and

chromosomal context of the female polymorphism may influence its evolutionary dynamics. Our

data is consistent with the evolution of a third morph due to an ectopic recombination event that

translocated genomic content from the *A* haplotype back into an *O* background. Ectopic

recombination can occur when TE propagation generates homologous regions in different

genomic locations[57,58], and may be facilitated by the excess of TE content in chromosome 13

(Exteded Data Fig. 4). The male reproductive morphs in the Ruff (*Calidris pugnax*) are one of

few previous examples of a novel phenotypic morph arising via recombination between two pre-

existing morph haplotypes[10]. In pond damselflies, female polymorphisms with three or more

female morphs are not uncommon, and in some cases female morphs exhibit graded resemblance

to males[59]. It is therefore possible that recombination, even if rare, has repeatedly generated

diversity in damselfly female morphs.

While recombination might have had a role in generating the the novel *I* morph, we observe

reduced recombination over the morph locus in comparison to the rest of the genome of *I.*

elegans (Extended Data Fig. 5). However, this reduction in recombination is not limited to the

morph locus and its vicinity, but rather pervasive across chromosome 13 (Extended Data Fig. 5).

This unexpected finding suggests two alternative causal scenarios. First, selection for reduced

recombination at the morph locus, following the origin of sexual mimicry, could have spilled

over chromosome 13, facilitating a subsequent accumulation of TEs and further reducing

recombination[60]. Second, TE enrichment and reduced recombination may have preceded the

evolution of female morphs, and facilitated the establishment and maintenance of the female

polymorphism by balancing selection.

Both historical scenarios are compatible with a morph locus reminiscent of a supergene, which is

defined by tight genetic linkage of multiple functional loci[61]. However, an alternative and

parsimonious explanation is that the novel sequences in *A* and *I* females and their flanking genes

may not code for anything important for the male-mimicking phenotype as such, but simply

disrupt a region of chromosome 13 that is required for the development of ancestral sexual

differentiation. The observation that *I* females carry part of the sequence that originated in *A* in a

different location of the scaffold (Fig. 4b), and still develop some male-like characters (e.g. black

thoracic stripes), could come about if insertions anywhere on a larger chromosomal region

disrupt female suppression of the male phenotype, although with variable efficacy depending on

the exact location or insertion size.

Concluding Remarks

Recent years have witnessed an explosion of studies uncovering the loci behind complex

phenotypic polymorphisms in various species. An emerging outlook is that not all

polymorphisms are created equal, with some governed by massive chromosomal

rearrangements[15–17], and others by a handful of regulatory sites[11,12,18]. Our results contribute to

this growing field by uncovering a single causal locus, that features structural variation and

morph-specific transcripts, in the female-limited morphs of *Ischnura* damselflies. These morphs

not only differ in numerous morphological and life-history traits[32,62,63] and gene expression

profiles[24,25], but they include a male mimic that is maintained by balancing frequency-dependent

selection. Our findings enable future studies on the developmental basis of such male mimicry,

with consequences for a broader understanding of the evolutionary dynamics of sexual

dimorphism and the cross-sexual transfer of trait expression[64].

METHODS

Ischnura elegans samples

Samples for morph-specific genome assemblies of *I. elegans* were obtained from F1 individuals

with genotypes *Ao*, *Io* and *oo* (one adult female of each genotype). In June 2019, recently-mated

O females were captured in field populations in Southern Sweden. These females oviposited in

the lab within 48 h, and their eggs were then released into outdoor cattle tanks seeded with

Daphnia and covered with synthetic mesh. Larvae thus developed under normal field conditions

and emerged as adults during the Summer of 2020. Emerging females were kept in outdoor

enclosures until completion of adult colour development[25,65]. Fully mature females were

phenotyped, collected in liquid nitrogen and kept at -80 ºC. Because all of these females carry a

copy of the most recessive allele *o*, individuals of the *A* and *I* morph are heterozygous, with

genotypes *Ao* and *Io*, respectively.

A total of 19 resequencing samples of each female morph of *I. elegans* were also collected from

local populations in Southern Sweden, within a 40 x 40 km area (Table S1). Samples were

submerged in 95% ethanol and stored in a -20 ºC freezer until extraction. Additionally, 24

individuals (six adult females of each morph and six males) were collected for RNAseq analysis

in a natural field population (Bunkeflostrand) in Southern Sweden, in early July 2019. These

samples were transported on carbonated ice and stored in -80 ºC until extraction.

Ischnura senegalensis samples

Adults of *I. senegalensis* (30 adult females of each morph) were collected for pool sequencing

from a population on Okinawa Island in Japan (26.148N, 127.795E) in May 2016. Samples were

visually determined to sex and morph and stored in 99% ethanol until extraction. Samples for

morph-specific genome assemblies of *I. senegalensis* were obtained from a population in

Clementi Forest, Singapore (1.33N, 103.78E). Because the *A* allele is recessive in *I.*

senegalensis, all females with the *A* phentoype are homozygous. To obtain a homozygous *O*-like

sample, we developed primers (forward: CGCGGTATGATATGGTCCGA, reverse:

GGCTGCTTACACCAATGCAA) for an *A*-specific sequence that is shared by *A* females of the

two species (318,131 - 318,213 bp on the *A* haplotype of *I. elegans*). We used the mapped pool-

seq data to identify fixed SNPs between species and tailor the primer sequences accordingly. We

then tested the primers in 20 *A* females of *I. senegalensis* using a 328 bp fragment of the Histone

H3 gene (forward: ATGGCTCGTACCAAGCAGACGGC, reverse:

ATATCCTTGGGCATGATGGTGAC)[66] as a positive control for the PCR reaction. Once

validated, we utilize these primers to identify *O*-like females lacking the *A* allele and selected

one of these samples for whole genome sequencing.

DNA extraction, library preparation and sequencing

High molecular weight (HMW) DNA was extracted from one *I. elegans* female of each genotype

(*Ao*, *Io*, *oo*), using the Nanobind® Tissue Big Extraction Kit (Cat. No. NB-900-701-01,

Circulomics Inc. (PacBio), MD, USA). HMW DNA was isolated from homozygous females of

each morph of *I. senegalensis*, using the Monarch® HMW DNA Extraction Kit for Tissue (Cat.

No. T3060S, New England BioLabs Inc., MA, USA). DNA from resequencing samples was

isolated using either a modified protocol for the DNeasy Blood and Tissue Kit (Cat. No. 19053,

Qiagen, Germany) or the KingFisher Cell and Tissue DNA Kit (Cat no. N11997, ThermoFisher

Scientific). *I. senegalensis* DNA was extracted from muscle tissues in thoraxes using Maxwell®

16 LEV Plant DNA Kit (Cat. No. AS1420, Promega, WI, USA). Details on extraction and

library preparation protocols are provided in the Supporting Text 1.

Sequencing libraries were constructed from each HMW DNA sample for the Nanopore LSK-110

ligation kit (Oxford Nanopore Technologies, UK). Adapter ligation and sequencing of *I. elegans*

samples were carried out at the Uppsala Genome Centre (NGI), hosted by SciLife Lab. Each

sample was sequenced on a PromethION R10.4 with 1 nuclease wash and two library loadings.

Library preparation and sequencing of *I. senegalensis* samples were carried out by the Integrated

Genomics Platform, Genome Institute of Singapore (GIS), A-STAR, Singapore. Each sample

was sequenced on a PromethION R9.4 flow cell, with 2 nuclease washes and three library

loadings.

RNA extraction and sequencing

Whole-thorax samples were grounded into a fine powder using a TissueLyser and used as input

for the Spectrum^{TM} Plant Total RNA Kit (Cat. No. STRN50, Sigma Aldrich, MO, USA),

including DNase I treatment (Cat. No. DNASE10, Sigma Aldrich, MO, USA). Library

preparation and sequencing were performed by SciLife Lab at the Uppsala Genome Centre

(NGI). Sequencing libraries were prepared from 300 ng of RNA, using the TrueSeq stranded

mRNA library preparation kit (Cat. No. 20020595, Illumina Inc., CA, USA) including polyA

selection and unique dual indexing (Cat. No. 20022371, Illumina Inc., CA, USA), according to

the manufacturer's protocol. Sequencing was performed on the Illumina NovaSeq 6000 SP

flowcell with paired-end reads of 150 bp.

De novo genome assembly

Bases in raw ONT reads from *I. elegans* were called using *Guppy* v 4.0.11 (*Ao* and *Io* data) and

Guppy v 5.0.11 (*oo* data) (https://nanoporetech.com/). Low quality reads (qscore < 7 for v 4.0.11

and < 10 for v 5.0.11) were subsequently discarded. High quality reads were assembled using the

Shasta long-read assembler v 0.7.0[67]. Each assembly was conducted under four different

configuration schemes, which modified the June 2020 Nanopore configuration file

(https://github.com/chanzuckerberg/shasta/blob/master/conf/Nanopore-Jun2020.conf) in

alternative ways (Table S3). Assembly metrics were compared among Shasta configurations for

each morph using *AsmQC*[68] (https://sourceforge.net/projects/amos/) and the *stats.sh* script in the

BBTools suite (https://sourceforge.net/projects/bbmap). The assembly with greater contiguity

(i.e. highest contig N50, highest average contig length and highest percentage of the main

genome in scaffolds > 50 kb) was selected for polishing and downstream analyses.

Bases in raw ONT reads from *I. senegalensis* samples were called using *Guppy* v 6.1.5. Reads

with quality score < 7 were subsequently discarded. High quality reads were assembled using the

Shasta long-read assembler v 0.7.0[67] and the configuration file T2 (Table S3), which was also

selected for the *Io* and *oo* assemblies of *I. elegans*.

Morph specific assemblies of *I. elegans* were first polished using the ONT reads mapped back to

their respective assembly with *minimap2* v 2.22-r1110[69], and the *PEPPER-Margin-DeepVariant*

pipeline r0.4[70]. Alternative haplotypes were subsequently filtered using *purge_dups* v 0.0.3[71], to

produce a single haploid genome assembly for each sample. The *I. elegans* draft assemblies were

then polished with short read data from one resequencing sample (TE-2564-SwD172_S37, Table

S1), using the *POLCA* tool in *MaSuRCA* v 4.0.4[72]. For every draft and final assembly of *I.*

elegans, we computed quality metrics as mentioned above and assessed the completeness of

conserved insect genes using *BUSCO* v 5.0.0[73] and the "insecta_odb10" database (Fig. S1). For

I. senegalensis, we report quality metrics of the final assemblies (Fig. S2).

Scaffolding with the Darwin Tree of Life super assembly

During the course of this study, a chromosome-level genome of *I. elegans* was assembled by the

Darwin Tree of Life (DToL) Project[34], based on long-read (PacBio) and short-read (Illumina)

data, as well as Hi-C (Illumina) chromatin interaction data. 99.5% of the total length of this

assembly is distributed across 14 chromosomes, one of which (no. 13) is fragmented and divided

into a main assembly and five unlocalized scaffolds.

We used RagTag v 2.10[74] to scaffold each our morph-specific assemblies based on the DToL

reference (Supporting Text 2). Scaffolding was conducted using the *nucmer* v 4.0.0[75] aligner and

default RagTag options. Morph-specific scaffolded genomes were also aligned to each other

using *nucmer* and a minimum cluster length of 100 bp. Alignments were then filtered to preserve

only the longest alignments in both reference and query sequences, and alignments of at least 5

kb. These assembly alignments were then used to visualize synteny patterns across morphs, in

the region uncovered in our association analyses (Extended Data Fig. 1), using the package

RIdeogram v 0.2.2[76] in *R* v 4.2.2[77].

Reference-based (SNP) GWAS

We first investigated genomic divergence between morphs using a standard GWAS approach

based on SNPs (Extended Data Fig. 1). Initially, we conducted preliminary analyses using

different morph assemblies as mapping reference. Once the *A*-specific genomic region was

confirmed, we designated the *A* assembly as the mapping reference for the main analyses. Short-

read data were mapped using *bwa-mem* v 0.7.17[78]. Optical and PCR duplicates were then flagged

in the unfiltered *bam* files using *GATK* v 4.2.0.0[79]. Variant calling, filtering and sorting were

conducted using bcftools v 1.12[80], excluding the flagged reads. We retained only variant sites

with mapping quality > 20, genotype quality > 30 and minor allele frequency > 0.02 (i.e. the

variant is present in more than one sample). To avoid highly repetitive content, we filtered

variants that had a combined depth across samples > 1360 (equivalent to all samples having ~

50% higher than average coverage), and variants located in sites annotated as repetitive in either

RepeatMasker v 1.0.93[81] or *Red* v 0.0.1[82]. The final variant calling file was analysed in pairwise

comparisons (*A* vs *O*, *A* vs *I*, *I* vs *O*) using *PLINK* v 1.9[83]

(http://pngu.mgh.harvard.edu/purcell/plink/). We report the -Log_{10} of P-values for SNP

associations in these pairwise comparisons.

Reference free (k-mer) GWAS

We created a list of all *k*-mers of length 31 in the short-read data (19 females per morph,

Extended Data Fig. 1) following Voichek and Weigel[35], and counting *k*-mers in each sample

using *KMC* v 3.1.0[84]. The *k*-mer list was filtered by the minor allele count; *k*-mers that appeared

in less than 5 individuals were excluded. *k*-mers were also filtered by percent canonized (i.e. the

percent of samples for which the reverse complement of the *k*-mer was also present). If at least

20% of the samples including a given *k*-mer contained its canonized form, the *k*-mer was kept in

the list. The *k*-mer list was then used to create a table recording the presence or absence of each

k-mer in each sample. A kinship matrix for all samples was calculated from this *k*-mer table, and

was converted to a *PLINK*[83] binary file, where the presence or absence of each *k*-mer is coded as

two homozygous variants. In this step, we further filtered the *k*-mers with a minor allele

frequency below 5%.

Because a single variant, be it a SNP or SV, will likely be captured by multiple *k*-mers,

significance testing of *k*-mer associations requires a method to control for the non-independence

of overlapping *k*-mers. We followed the approach developed by Voichek and Weigel[35], which

uses a linear mixed model (LMM) genome-wide association analysis implemented in *GEMMA* v

0.98.5[85], and computes P-value thresholds for associated *k*-mers based on phenotype

permutations. We thus report *k*-mers below the 5% false-positive threshold as *k*-mers

significantly associated with the female-polymorphism in *I. elegans*. We conducted three *k*-mers

based GWAS: 1) comparing male mimics to the putatively ancestral female morph (*A* vs *O*), 2)

comparing male mimics to the most derive female morph (*A* vs *I*), and 3) comparing both

derived female morphs (*A* and *I*) to the ancestral *O* females. For every analysis, we then mapped

the significant *k*-mers to all reference genomes using *Blast* v 2.22.28[86] for short sequences, and

removed alignments that were below 100% identity and below full-length. The mapped *k*-mers

thus indicate the proportion of relevant genomic content present in each morph and how this

content is distributed across each genome (Extended Data Table 1).

Read depth analysis

To validate the *k*-mer GWAS results of unique genomic content in *A*-females relative to both *I*-

and *O*-females, we plotted read-depth across our region of interest (the unlocalized scaffold 2 of

chromosome 13, see Results) in the *A* assembly (Extended Data Fig. 1). Short-read data (19

samples per morph) were mapped to the assembly with *bwa-mem* v 0.7.17[78] and reads with

mapping score < 20 were filtered, using *Samtools* v 1.14[87]. Long-read data (one sample per

morph) were also mapped to the assembly using *minimap2* v 2.22-r1110[69], and quality filtering

was conducted as above. Read depth was then averaged for each sample across 500 bp, non-

overlapping windows using *mosdepth* v 0.2.8[88]. We also annotated repetitive content in the

reference genome using *RepeatMasker* v 1.0.93[81] and *Red* v 0.0.1[82], and filtered windows with

more than 10% repetitive content under either method.

To account for differences in overall coverage between samples, we conducted the same

procedure on a large (~15 mb) non-candidate region in chromosome 11 and calculated a

"background read depth" as the mean read depth across the non-repetitive windows of this

region. We then expressed read-depth in the candidate region as a proportion of the background

read depth. Values around 1 thus indicate that a sample is homozygous for the presence of the

sequence in a window. Values around 0.5 suggest that the sample only has one copy of this

sequence in its diploid genome (i.e. it is heterozygous). Finally, values of 0 imply that the 500 bp

reference sequence is not present in the sample (i.e. the window is part of an insertion or

deletion).

We also investigated read-depth coverage on the *I* assembly, specifically across the region that

was identified in the *k*-mer based GWAS as capturing content that differentiated both *A* and *I*

females from *O* females (Fig. 3b, Extended Data Fig. 1). To do so, we followed the same

strategy as a above, except here we used a 15 mb region from chromosome 1 to estimate

background read depth.

Population genetics

We investigated the evolutionary consequences of morph divergence by estimating between-

morph Fst and population-wide Tajima's D and nucleotide diversity (π). For these analyses, we

used the *A* assembly as mapping reference and the same variant calling approach as described for

the SNP based GWAS, but applied different filtering criteria (Extended Data Fig. 1).

Specifically, invariant sites were retained and we only filtered sites with mapping quality score <

20 and combined depth across samples > 1360 (equivalent to ~50% excess coverage in all

samples). Fst and (π) were estimated in *pixy* v 1.2.5[89] across 30 kb windows. Fst was computed

using the *hudson* estimator[90]. Negative Fst values were converted to zero for plotting. Tajima's D

was estimated across 30 kb using *vcftools* v 0.1.17[91]. In all analyses, windows with > 10%

repetitive content according to either *RepeatMasker* v 1.0.93[81] or *Red* v 0.0.1[82] annotation were

excluded.

Structural variants

We used two complimentary approaches to identify SVs overlapping the genomic region

uncovered by both *k*-mer-based and SNP-based GWAS. First, we mapped the raw data from

each long-read sample to the assemblies of alternative morphs (e.g. *Ao* data mapped to *Io* and *oo*

assemblies), and called SVs using *Sniffles* v 1.0.10[92] (Extended Data Fig. 1). These SV calls may

represent fixed differences between morphs, within-morph polymorphisms, or products of

assembly error. We therefore used *SamPlot* v 1.3.0[93] and our short-read samples (n = 19 per

morph) to validate morph-specific SV calls (Extended Data Fig. 1). *Samplot* identifies and plots

reads with discordant alignments, which can result from specific types of SVs. For example, if

Sniffles called a 10 kb deletion in the *Ao* and *Io* long-read samples relative to the *oo* assembly,

we then constructed a Samplot for this region using short-read data, and expected to find support

for such deletion in *I* and *A* samples, but not in *O* samples. We complemented this validation

approach with a scan of the region of interest in each assembly, in windows of 250 and 500 kb,

again using *Samplot* and the short-read data. If a SV appeared to be supported by the majority of

short-read samples from an alternative morph, we zoomed in this SV and recorded the number of

samples supporting the call in each morph.

Linkage disequilibrium and transposable elements

To estimate linkage disequilibrium (LD), we used the same variant calling file as for the SNP

based GWAS, which included only variant sites and was filtered by mapping quality, genotyping

quality, minimum allele frequency, and read depth, as described above (Extended Data Fig. 1).

The file was downsampled to one variant every 100th using vcftools v 0.1.17[91], prior to LD

estimation. We estimated LD using *PLINK* v 1.9[83], and recorded R^2 values > 0.05 for pairs up to

15 mb apart or with 10,000 or fewer variants between them. We estimated LD for the

unlocalized scaffold 2 of chromosome 13, which contains the morph loci and is about ~ 15 mb in

the *A* assembly. For comparison, we also estimated LD across the first 15 mb of the fully

assembled chromosomes (1-12 and X), the main scaffold of chromosome 13, and the unlocalized

scaffolds 1, 3 and 4 of chromosome 13.

We used the TE annotations from *RepeatModeler* v 2.0.1 *RepeatMasker* v 1.0.93[81] and "One

code to find them all"[94] to quantify TE coverage in chromosome 13 in comparison to the rest of

the genome. We divided each chromosome into 1.5 mb windows, and computed the proportion

of each window covered by each TE family.

Evidence of a trans-species polymorphism

We used pool-seq data from the closely related Tropical Bluetail damselfly (*Ischnura*

senegalensis) to determine whether male-mimicry has a shared genetic basis in the two species

(Extended Data Fig. 1). First, we aligned the short-read data from the the two *I. senegalensis*

pools (*A* and *O*-like) to the *A* morph assembly of *I. elegans* using *bwa-mem* v 0.7.17[78], and

filtered reads with mapping score < 20, using *Samtools* v 1.14[87]. We then quantified read depth

as for the *I. elegans* resequencing data (see *Read depth analysis* above). To confirm that the

higher read-depth coverage of the *A* pool is specific to the putative morph locus, we also plotted

the distribution of read-depth differences between *O*-like and *A* pools across the rest of the

genome and compared it to the morph locus (Supporting Text 5). Next, we determined if the ~20

kb SV that characterizes *A* and *I* females of *I. elegans* is also present in *A* females of *I.*

senegalensis. To do this, we mapped the pool-seq data to the *O* assembly of *I. elegans* as above,

and scanned the region at the start of the scaffold 2 of chromosome 13 for SVs using Samplot v

1.3.0[93]. Finally, we aligned the morph-specific assemblies of *I. senegalensis* to the *A* assembly of

I. elegans, using *nucmer* v 4.0.0[75] and preserving alignments > 500 bp and with identity > 70%

(Extended Data Fig. 1). We visualized synteny patterns across the morph locus using the package

RIdeogram v 0.2.2[76] in *R* v 4.2.2[77].

Gene content and expression in the morph locus

We assembled transcripts in the *A* morph genome (Extended Data Fig. 1) to identify potential

gene models unique to the *A* or *A* and *I* morphs and which would therefore be absent from the *I.*

elegans reference annotation (based on the *O* haplotype). First, all raw RNAseq data from *I.*

elegans samples were mapped to the *A* assembly using HISAT2 v 2.2.1[95] and reads with

mapping quality < 60 were filtered using *Samtools* v 1.19[87]. Transcripts were then assembled in

StringTie v 2.1.4[96] under default options, and merged into a single gtf file. Transcript abundances

were quantified using this global set of transcripts as targets, and a transcript count matrix was

produced using the *prepDE.py3* script provided with *StringTie*. Mapped RNAseq data from *I.*

senegalensis was also used to assemble transcripts (Extended Data Fig. 1), but this time the

HISAT2 assembly was guided by the annotation based on *I. elegans* data, while allowing the

identification of novel transcripts. Transcript abundances were quantified as for *I. elegans*.

We analyzed differential gene expression using the package *edgeR* v 3.36[97] in *R* v 4.2.2[77].

Transcripts with fewer than one count per million in more than three samples were filtered.

Library sizes were normalized across samples using the trimmed mean of M-values method[98],

and empirical Bayes tagwise dispersion[99] was estimated prior to pairwise expression analyses.

Differential expression of genes in the morph loci was tested using two-tailed exact tests[100],

assuming negative-binomially distributed transcript counts and applying the Benjamini and

Hochberg's algorithm to control the false discovery rate (FDR)[101].

Nucleotide sequences of all transcripts mapped to the 1.5 mb morph locus in the *A* assembly

were selected. Coding sequences (CDS) in these transcripts were predicted using *Transdecoder* v

5.5.0 (https://github.com/TransDecoder/TransDecoder). Predicted CDS and peptide sequences

were read from the assemblies using the genome-based coding region annotation produced with

Transdecoder and *gffread* v 0.12.7[102]. We investigated whether any inferred peptides or

transcripts were unique to *A* or *A* and *I* by comparing these sequences to the DToL reference

transcriptome and proteome (based on the *O* haplotype). We then searched for homologous and

annotated proteins in other taxa in the Swissprot database using *Blast* v 2.9.0[86]. We found three

gene models which are protein-coding and present in both *A* and *O* females (see Results, Fig. 6).

We scanned these protein sequences for functional domains using *InterProScan*[103] and searched

for orthologous groups and functional annotations in *EggNOG* v 5.0[104].

Data availability

Sequencing data from this study have been submitted to the NCBI Sequence Read Archive

(SRA) (https://www.ncbi.nlm.nih.gov/sra/) under BioProject PRJNA940276. For details, please

see Supplementary Tables 1 and 2. Morph-specific genome assemblies and intermediate output

files required to reproduce the figures in the main text and Supporting Material are available on

Zenodo[105].

Code availability

All code necessary to reproduce the results of this study can be found on Zenodo

https://doi.org/10.5281/zenodo.8304055 and Github https://github.com/bwillink/Morph-locus.

Acknowledgements

We thank H. Dort, R.A. Steward, J. Haushofer, P. de Sessions, and the Monteiro Lab for

suggestions and helpful discussions. We are also grateful to M.P. Celorio-Mancera and H.M.

Low for invaluable technical support. Prof. A. Monteiro hosted B.W. at the National University

of Singapore during part of the duration of this study. Computation and data handling were

enabled by resources provided by the Swedish National Infrastructure for Computing (SNIC)

through the Uppsala Multidisciplinary Center for Advanced Computational Science (UPPMAX),

under projects 2022/6-230 and 2022/5-419 awarded to EIS. Specimens of *I. senegalensis* were

collected in Singapore, under a Research Permit (NP/RP22-015b) from the National Parks

Board, Singapore. This work was supported by an International Postdoc Grant from the Swedish

Research Council (VR; grant no. 2019-06444 to BW). Funding was also provided by the

Swedish Research Council (VR; grant no. 2017-04386 to CWW, and grant no. 2016-03356 to

EIS) and by the Stina Werners Foundation (grant no. 2018-017 to EIS) and Erik Philip Sörensens

Stiftelse (grant no. 2019-033 to EIS). SN was funded by a scholarship grant for Master's students

from Sven and Lily Lawski's Foundation.

Author contributions

BW conceived the study with input from **CWW**. **EIS** organized field work in the long-term

population study of *I. elegans* during 2019 and 2020, planned and prepared the outdoor rearing

experiments. **EIS** and **SN** collected DNA and RNA samples of *I. elegans*. **MT** and **YT** collected

samples for pool sequencing of *I. senegalensis*, and **BW** collected samples of *I. senegalensis* in

Singapore. **SN**, **BW**, and **KT** conducted laboratory work on *I. elegans*. **MT**, **YT**, and **BW**

conducted laboratory work on *I. senegalensis*. **BW** analyzed the data with input from **CWW**,

KT, **RC**, and **TL**. **BW** wrote the manuscript with contributions from all authors.

Competing interests

The authors declare no competing interests.

Figure 1. The evolution of female-limited colour polymorphisms in *Ischnura* damselflies. **a** A

previous phylogenetic study and ancestral state reconstruction[28] proposed that the genus

Ischnura had a sexually dimorphic ancestor, with *O*-like females (red circle). The *O* morph is

markedly different from males, having a bronze-brown thorax and faint stripes, instead of the

black thoracic stripes on a bright blue background of males. **b** Male mimicry (*A* females, blue

circle) has evolved more than once, for instance, in an ancestor of the (expanded) clade that

includes the Common Bluetail (*I. elegans*, encircled with solid line) and the Tropical Bluetail (*I.*

senegalensis, encircled with dashed line). **c** *I. elegans* is a trimorphic species, due to the recent

evolution of a third female morph, *I* (yellow circle). In *I. elegans*, morph inheritance follows a

dominance hierarchy, where the most dominant allele produces the *A* morph and two copies of

most recessive allele are required for the development of *O* females. In contrast, the *O* allele is

dominant in *I. senegalensis*[106]. Terminal nodes in the phylogeny represent different species. Gray

triangles represent other clades of *Ischnura* that are collapsed for clarity.

Figure 2. Morph determination in the Common Bluetail Damselfly (*Ischnura elegans*) is

controlled in a ~ 1.5 mb region of chromosome 13. **a** SNP-based genome-wide associations in all

pairwise analyses between morphs. Genomic DNA from 19 wild-caught females of each colour

morph and of unknown genotype was extracted and sequenced for these analyses. Illumina short

reads were aligned against an *A* morph genome assembly, generated from Nanopore long-read

data (Extended Data Fig. 1). **b** A closer look of the SNP associations on the unlocalized scaffold

2 of chromosome 13, which contained all highly significant SNPs. Transcripts expressed in at

least one adult of both *I. elegans* and *I. senegalensis* are shown at the bottom (see also Fig. 6).

Grey transcripts are shared by all morphs, whereas blue transcripts are uniquely present in *A* or *A*

and *I* samples (see *Shared and morph-specific genes reside in the morph locus*). The *y* axis in **a**

and **b** indicates unadjusted -$\log_{10}$ P-values calculated from chi-squared tests. **c** Fst values

averaged across 30 kb windows for the same pairwise comparisons as in the SNP based GWAS.

The dashed line marks the 95 percentile of all non-zero Fst values across the entire genome. The

red double arrow shows the region of elevated divergence between *O* and both *A* and *I* samples

(~50 kb - 0.2 mb). The blue double arrow shows the region of elevated divergence between *A*

and both *O* and *I* samples (~0.2 mb - 1.5 mb). Population-level estimates of **d** Tajima's D, and **e**

nucleotide diversity (π) averaged across 30 kb windows. The shaded area contains the 5-95

percentile of all genome-wide estimates.

Figure 3. Female morphs of *Ischnura elegans* differ in genomic content. Number of significant

k-mers (below the 5% false-positive threshold, see Methods) associated with pairwise genome-

wide analyses and mapped to the unlocalized scaffold 2 of chromosome 13, in **a** the *A* morph

assembly, and in **b** the *I* morph assembly. Standardized read depths along the unlocalized

scaffold 2 of chromosome 13, relative to background coverage of **c** the *A* morph assembly, and **d**

the *I* morph assembly. Solid lines (yellow, blue and red) show short-read data (19 samples per

morph) and black dashed lines represent long-read data (1 sample per morph).Grey areas show

regions of genomic content present in *A* and *I* individuals, but absent in all but one *O* sample.

Note that different regions of the scaffold are plotted for the two assemblies (see main text).

Figure 4. Structural variants differentiate morph haplotypes in the Common Bluetail Damselfly

(*Ischnura elegans*). **a** Alignment between morph-specific genomes assembled from long-read

Nanopore samples with genotypes *Ao*, *Io*, and *oo*. Grey lines represent alignments of at least 5 kb

and > 70% identity. The black line connects regions of genomic content shared by the three

morphs within the morph locus. The red to blue gradient represents a ~20 kb region that carries

an inversion signature in *A* and *I* females relative to the *O* haplotype (see Extended Data Fig. 2).

The blue to yellow gradient represents a ~ 150 kb alignment between the start of the unlocalized

scaffold 2 of chromosome 13 in *A* and a region ~ 3.5 mb apart in the *I* haplotype. Coordinates at

the bottom are based on the Darwin Tree of Life (DToL) reference assembly. **b** Schematic

illustration of the hypothetical sequence of events responsible for the evolution of novel female

morphs. First, a sequence originally present in *O* was duplicated and inverted in tandem,

potentially causing the initial divergence of the *A* allele. Second, part of this inversion was

subsequently duplicated in *A*, in association with a putative TE, leading to multiple inversion

signatures in the *A* haplotype relative to an *O* reference (see Extended Data Fig. 3). Finally, part

of the *A* duplications were translocated into a position ~ 3.5 mb downstream into an *O*

background, giving rise to the *I* morph. Currently, *A* females are also characterized by another

region of unique content and unknown origin (question mark). *A* female show elevated sequence

divergence in the internal region of the morph locus that is shared by all haplotypes (dark grey

bars, see also black line in **a**). Coordinates on the *O* haplotype are based on the (DToL) reference

assembly. Grey numbers in IV give the approximate size of genomic sequences in *A* and *I* that

are absent in *O*.

Figure 5. A shared genomic basis of *A* females in *Ischnura elegans* and *Ischnura senegalensis* **a**

I. senegalensis is a female-dimorphic species, where one female morph (*O*-like) is distinctly

different from males and resembles *O* females in *I. elegans*, and the other female morph (*A*) is a

male-mimic. Photo credit: Mike Hooper. **b** Standardized read depth of pool-seq samples (n = 30

females of each morph per pool) of *I. senegalensis*, against the *A* morph assembly of *I. elegans*,

calculated in 500 bp windows. The x-axis shows the first 1.5 mb of the unlocalized scaffold 2 of

chromosome 13. **c** Alignments between morph-specific genomes from a homozygous *O*-like

female of *I. senegalensis* (top), an *Ao* female of *I. elegans* (middle) and a homozygous *A* female

of *I. senegalensis* (bottom). Lines connecting the assemblies represent alignments of at least 500

bp and > 70% identity. The black line connects genomic content in the morph locus, which is

shared by the three morphs of *I. elegans*. In *I. elegans*, this region is rich in SNPs differentiating

A females from the other two morphs (see Fig. 2b). The blue-turquoise gradient connects

sequences uniquely present in the *A* morphs of *I. elegans* and *I. senegalensis*.

Figure 6. The morph locus of *Ischnura elegans* is situated in the unlocalized scaffold 2 of

chromosome 13 **a** Diagram of the ~ 1.5 mb morph locus on the *A*-morph assembly, showing

from top to bottom: morph-specific read depth coverage, the location of LINE retrotransposons

in the the Jockey family, the mapping locations of *A*-derived reads with a previously detected

inversion signature against *O* females, and transcripts expressed in at least one adult individual

of both *I. elegans* and *I. senegalensis*. Transcripts plotted in black are present in both the *A* and *O*

assemblies, while transcripts in blue are located in genomic regions that are unique to the *A*

haplotype or are shared between *A* and *I* but not the *O* allele. **b** Functional annotations and sex-

and morph-specific expression of transcripts. Square fill indicates whether transcript expression

was detected in each group. RNAseq data for *I. elegans* comes from whole-thorax samples from

sexually immature and sexually mature wild-caught adults (n = 3 females of each morph and 3

males). RNAseq data for *I. senegalensis* comes from a recent study in which the abdomen, head,

thorax, and wings were sampled in two females of each morph and two males (one individual of

each group sampled upon emergence and one sampled after two days).

Extended Data Table 1. Significant *k*-mers associated with morph comparisons in *I. elegans*.
For each comparison (*A* vs *O*, *A* vs *I* and *A* and *I* vs *O*), we show the total number of significant *k*-mers, and the total number of significant *k*-mers that map without any mismatching position to morph-specific reference assemblies. Of the mapping *k*-mers, we then show the number located in the unlocalized scaffold 2 of chromosome 13, which includes the putative morph locus. For the DToL assembly, we show the number of significant *k*-mers mapping to both the primary assembly (capturing the *O* allele) and the haplotigs, where the haplotig RAPID_106 comprises the *A* allele (see Supporting Text 2).

k-mers	*A* vs *O*	*A* vs *I*	*I* vs *O*
Total number	568,039	508,031	85,134
A assembly	435,509	383,679	45,580
A Chr 13_2	427,606	376,075	44,990
I assembly	46,733	1,111	49,093
I Chr 13_2	45,819	375	48,484
O assembly	1,478	762	1,452
O Chr 13_2	276	92	72
DToL primary assembly	915	756	676
DToL Chr 13_2	134	115	14
DToL haplotigs	542,571	489,912	66,861
DToL RAPID_106	539,588	486,943	65,999

 Extended data figure legends

Extended Data Figure 1. Outline of data and analyses used in this study. For our main study

species *Ischnura elegans*, we obtained short-read genomic data from 19 field-caught females per

morph, and long-read genomic data from three females with genotypes *Ao*, *Io*, and *oo*. The long-

read samples were used to assemble morph-specific genomes, scaffolded against the Darwin

Tree of Life reference assembly. We obtained whole-thorax RNAseq data from three females of

each morph in both sexually immature and sexually mature colour phases (n = 2). Immature and

mature males (n = 3 of each) were also sampled for whole-thorax RNAseq data. We used short-

read pool-seq data (n = 30 individuals of each morph per pool) of the close relative *Ischnura*

senegalensis to investigate whether the female polymorphisms in both species share a genomic

basis. We also analysed expression levels of candidate genes in this species, using samples from

a previously published study[38], which produced transcriptomic data from four body parts (head,

thorax, wing and abdomen) of each *A* females, *O* females and males (n = 2), sampled at adult

emergence and two days thereafter. The *k*-mer based GWAS is reference-free, but significant *k*-

mers were mapped to the morph-specific assemblies to determine their chromosomal context.

Extended Data Figure 2. An inversion signature differentiates *A* and *I* individuals from the *O*

morph. Read mapping and sample coverage at the start of the scaffold 2 of chromosome 13 in **a**

our *O* assembly and **b** the DToL reference assembly, showing a signature of a ~ 20 kb inversion

in *A* and *I* samples. A single *O* sample also exhibited this signature but was excluded here for

clarity (see Supporting Text 3). Note that the first 415 kb of the reference DToL assembly are

absent in our scaffolded *O* assembly, and therefore the x-axis is shifted by 415 kb in **b**.

Extended Data Figure 3. The *A* and *I* reads mapped to inversion break points on the *O* assembly

(see Extended Data Fig. 2) map to multiple locations on the *A* assembly. **a** Reads from the first

inversion breakpoint. **b** Reads from the second inversion breakpoint. Each row represents a

sample and each circle an individual read. The x-axis corresponds to coordinates on the *A*

assembly.

Extended Data Figure 4. Proportion of TE content in non-overlapping 1.5 mb regions. The gray

dots correspond to genomic windows outside chromosome 13. The main assembly and the

unlocalized scaffolds of chromosome 13 are depicted with different colours. The dashed line

marks the 95 percentile of TE coverage across all windows.

Extended Data Figure 5. Linkage disequilibrium (LD) in the genome of *Ischnura elegans*. LD

estimates are shown for the first 15 mb of each chromosome and all unlocalized scaffolds of

chromosome 13. The morph locus is found in the first ~ 1.5 mb of the unlocalized scaffold 2 of

chromosome 13, which has a total size of ~ 15 mb. Each dot represent the square correlation

coefficient (R^2) between two variant sites on the x axis, separated by the number of sites

indicated in the y axis.

Extended Data Figure 6. Evidence of a translocation between *A* and *I* haplotypes. Mapping and

coverage of long reads from an *Io* sample across the first 5.6 mb of the unlocalized scaffold 2 of

chromosome 13 in the *A* assembly, showing a signature consistent with either a 5.54 mb

inversion or a translocation of inverted *A* content. Absence of morph divergence beyond ~1.5 mb

on the *A* assembly supports the translocation scenario.

Extended Data Figure 7. Structural variants between *A* and *O*-like females of *I. senegalensis*

along the morph locus identified in *I. elegans.* **a** Read mapping and sample coverage of *I.*

senengalensis pool-seq data at the start of the unlocalized scaffold 2 of chromosome 13 in the *O*

assembly of *I. elegans.* The same ~ 20 kb inversion signature is found in *A* and *I* samples of *I.*

elegans (see Extended Data Fig. 2). **b-c** The *A*-pool reads mapped to break points on the *O*

assembly map to multiple locations on the *A* assembly. **b** Reads from the first breakpoint. **c**

Reads from the second breakpoint. Each row represents a pool of *I. senegalensis* and each circle

an individual read. The x-axis corresponds to the *A* assembly of *I. elegans.*

Extended Data Figure 8. Morph divergence using the DToL assembly (*O* haplotype) as

mapping reference. **a** SNP-based genome-wide associations in all pairwise analyses between

morphs. Genomic DNA from 19 wild-caught females of each colour morph and of unknown

genotype was extracted and sequenced for these analyses. Illumina short reads were aligned

against the DToL reference assembly. **b** A closer look of the SNP associations on the

unlocalized scaffold 2 of chromosome 13, which contained all highly significant SNPs. The *y*

axis in **a** and **b** indicates unadjusted -Log_{10} P-values calculated from chi-squared tests. **c** Fst

values averaged across 30 kb windows for the same pairwise comparisons as in the SNP based

GWAS. The dashed line marks the 95 percentile of all non-zero Fst values across the entire

genome. The red double arrow shows the region of elevated divergence between *A* and both *O*

and *I* samples. Population-level estimates of **d** Tajima's D, and **e** nucleotide diversity (π)

averaged across 30 kb windows. The shaded area contains the 5-95 percentile of all genome-

wide estimates.

REFERENCES

1. Mank, J. E. Sex chromosomes and the evolution of sexual dimorphism: Lessons from the
genome. *The American Naturalist* **173**, 141–150 (2009).

2. De Lisle, S. P. Understanding the evolution of ecological sex differences: Integrating
character displacement and the Darwin-Bateman paradigm. *Evolution Letters* **3**, 434–447
(2019).

3. Hopkins, B. R. & Kopp, A. Evolution of sexual development and sexual dimorphism in
insects. *Current opinion in Genetics & Development* **69**, 129–139 (2021).

4. Jukema, J. & Piersma, T. Permanent female mimics in a lekking shorebird. *Biology*
Letters **2**, 161–164 (2006).

5. Hurtado-Gonzales, J. L. & Uy, J. A. C. Alternative mating strategies may favour the
persistence of a genetically based colour polymorphism in a pentamorphic fish. *Animal*
Behaviour **77**, 1187–1194 (2009).

6. Gosden, T. P. & Svensson, E. I. Density-dependent male mating harassment, female
resistance, and male mimicry. *The American Naturalist* **173**, 709–721 (2009).

7. Falk, J. J., Rubenstein, D. R., Rico-Guevara, A. & Webster, M. S. Intersexual social
dominance mimicry drives female hummingbird polymorphism. *Proceedings of the*
Royal Society B **289**, 20220332 (2022).

8. Falk, J. J., Webster, M. S. & Rubenstein, D. R. Male-like ornamentation in female
hummingbirds results from social harassment rather than sexual selection. *Current*
Biology **31**, 4381–4387 (2021).

9. Mank, J. E. Sex-specific morphs: The genetics and evolution of intra-sexual variation.
Nature Reviews Genetics 1–9 (2022).

10. Lamichhaney, S. *et al.* Structural genomic changes underlie alternative reproductive
strategies in the ruff (*Philomachus pugnax*). *Nature Genetics* **48**, 84–88 (2016).

11. Andrade, P. *et al.* Regulatory changes in pterin and carotenoid genes underlie balanced
color polymorphisms in the wall lizard. *Proceedings of the National Academy of Sciences*
116, 5633–5642 (2019).

12. Kim, K.-W. *et al.* Genetics and evidence for balancing selection of a sex-linked colour
polymorphism in a songbird. *Nature Communications* **10**, 1–11 (2019).

13. Woronik, A. *et al.* A transposable element insertion is associated with an alternative life
history strategy. *Nature Communications* **10**, 1–11 (2019).

14. Hendrickx, F. *et al.* A masculinizing supergene underlies an exaggerated male
reproductive morph in a spider. *Nature Ecology & Evolution* **6**, 195–206 (2022).

15. Tuttle, E. M. *et al.* Divergence and functional degradation of a sex chromosome-like
supergene. *Current Biology* **26**, 344–350 (2016).

16. Sanchez-Donoso, I. *et al.* Massive genome inversion drives coexistence of divergent
morphs in common quails. *Current Biology* **32**, 462–469 (2022).

17. Villoutreix, R. *et al.* Inversion breakpoints and the evolution of supergenes. *Molecular
Ecology* **30**, 2738–2755 (2021).

18. Tunström, K. *et al.* Evidence for a single, ancient origin of a genus-wide alternative life
history strategy. *Science Advances* **9**, eabq3713 (2023).

19. Gutiérrez-Valencia, J., Hughes, P. W., Berdan, E. L. & Slotte, T. The genomic
architecture and evolutionary fates of supergenes. *Genome Biology and Evolution* **13**,
evab057 (2021).

20. Sandkam, B. A. *et al.* Extreme Y chromosome polymorphism corresponds to five male
reproductive morphs of a freshwater fish. *Nature Ecology & Evolution* **5**, 939–948
(2021).

21. Kunte, K. *et al.* Doublesex is a mimicry supergene. *Nature* **507**, 229–232 (2014).

22. Feiner, N. *et al.* A single locus regulates a female-limited color pattern polymorphism in
a reptile. *Science Advances* **8**, eabm2387 (2022).

23. Neff, B. D. & Svensson, E. I. Polyandry and alternative mating tactics. *Philosophical
Transactions of the Royal Society B: Biological Sciences* **368**, 20120045 (2013).

24. Takahashi, M., Takahashi, Y. & Kawata, M. Candidate genes associated with color
morphs of female-limited polymorphisms of the damselfly *Ischnura senegalensis*.
Heredity **122**, 81–92 (2019).

25. Willink, B., Duryea, M. C., Wheat, C. & Svensson, E. I. Changes in gene expression
during female reproductive development in a color polymorphic insect. *Evolution* **74**,
1063–1081 (2020).

26. Takahashi, M., Okude, G., Futahashi, R., Takahashi, Y. & Kawata, M. The effect of the
doublesex gene in body colour masculinization of the damselfly *Ischnura senegalensis*.
Biology Letters **17**, 20200761 (2021).

27. Cordero, A., Carbone, S. S. & Utzeri, C. Mating opportunities and mating costs are
reduced in androchrome female damselflies, *Ischnura elegans* (odonata). *Animal
Behaviour* **55**, 185–197 (1998).

28. Blow, R., Willink, B. & Svensson, E. I. A molecular phylogeny of forktail damselflies
(genus *ischnura*) reveals a dynamic macroevolutionary history of female colour
polymorphisms. *Molecular Phylogenetics and Evolution* **160**, 107134 (2021).

29. Henze, M. J., Lind, O., Wilts, B. D. & Kelber, A. Pterin-pigmented nanospheres create
the colours of the polymorphic damselfly *Ischnura elegans*. *Journal of the Royal Society
Interface* **16**, 20180785 (2019).

30. Svensson, E. I., Abbott, J. & Härdling, R. Female polymorphism, frequency dependence,
and rapid evolutionary dynamics in natural populations. *The American Naturalist* **165**,
567–576 (2005).

31. Le Rouzic, A., Hansen, T. F., Gosden, T. P. & Svensson, E. I. Evolutionary time-series
analysis reveals the signature of frequency-dependent selection on a female mating
polymorphism. *The American Naturalist* **185**, E182–E196 (2015).

32. Svensson, E. I., Willink, B., Duryea, M. C. & Lancaster, L. T. Temperature drives pre-
reproductive selection and shapes the biogeography of a female polymorphism. *Ecology*
Letters **23**, 149–159 (2020).

33. Willink, B., Duryea, M. C. & Svensson, E. I. Macroevolutionary origin and adaptive
function of a polymorphic female signal involved in sexual conflict. *The American*
Naturalist **194**, 707–724 (2019).

34. Price, B. W., Winter, M., Brooks, S. J. & Darwin Tree of Life Consortium. The genome
sequence of the blue-tailed damselfly, elegans (vander linden, 1820) [version 1; peer
review: Awaiting peer review]. *Wellcome Open Research* **7**, 66 (2022).

35. Voichek, Y. & Weigel, D. Identifying genetic variants underlying phenotypic variation in
plants without complete genomes. *Nature Genetics* **52**, 534–540 (2020).

36. Parr, M. J. The terminology of female polymorphs of *ischnura* (Zygoptera:
Coenagrionidae). *International Journal of Odonatology* **2**, 95–99 (1999).

37. Takahashi, Y. & Watanabe, M. Male mate choice based on ontogenetic colour changes of
females in the damselfly *Ischnura senegalensis*. *Journal of Ethology* **29**, 293–299 (2011).

38. Okude, G. *et al.* Molecular mechanisms underlying metamorphosis in the most-ancestral
winged insect. *Proceedings of the National Academy of Sciences* **119**, e2114773119
(2022).

39. Laity, J. H., Lee, B. M. & Wright, P. E. Zinc finger proteins: New insights into structural
and functional diversity. *Current Opinion in Structural Biology* **11**, 39–46 (2001).

40. Gudmunds, E., Wheat, C. W., Khila, A. & Husby, A. Functional genomic tools for
emerging model species. *Trends in Ecology & Evolution* (2022).

41. Chakraborty, M., Emerson, J. J., Macdonald, S. J. & Long, A. D. Structural variants
exhibit widespread allelic heterogeneity and shape variation in complex traits. *Nature*
Communications **10**, 1–11 (2019).

42. Quan, C., Lu, H., Lu, Y. & Zhou, G. Population-scale genotyping of structural variation
in the era of long-read sequencing. *Computational and Structural Biotechnology Journal*
(2022).

43. Schrader, L. & Schmitz, J. The impact of transposable elements in adaptive evolution.
Molecular Ecology **28**, 1537–1549 (2019).

44. Catlin, N. S. & Josephs, E. B. The important contribution of transposable elements to
phenotypic variation and evolution. *Current Opinion in Plant Biology* **65**, 102140 (2022).

45. Zhang, J. *et al.* Autotetraploid rice methylome analysis reveals methylation variation of
transposable elements and their effects on gene expression. *Proceedings of the National*
Academy of Sciences **112**, E7022–E7029 (2015).

46. Diehl, A. G., Ouyang, N. & Boyle, A. P. Transposable elements contribute to cell and
species-specific chromatin looping and gene regulation in mammalian genomes. *Nature*
Communications **11**, 1–18 (2020).

47. Sundaram, V. & Wysocka, J. Transposable elements as a potent source of diverse cis-
regulatory sequences in mammalian genomes. *Philosophical Transactions of the Royal*
Society B **375**, 20190347 (2020).

48. Jangam, D., Feschotte, C. & Betrán, E. Transposable element domestication as an
adaptation to evolutionary conflicts. *Trends in Genetics* **33**, 817–831 (2017).

49. McCue, A. D. & Slotkin, R. K. Transposable element small RNAs as regulators of gene
expression. *Trends in Genetics* **28**, 616–623 (2012).

50. Kiuchi, T. *et al.* A single female-specific piRNA is the primary determiner of sex in the
silkworm. *Nature* **509**, 633–636 (2014).

51. Van Gossum, H., Stoks, R. & De Bruyn, L. Frequency-dependent male mate harassment
and intra-specific variation in its avoidance by females of the damselfly *Ischnura*
elegans. Behavioral Ecology and Sociobiology **51**, 69–75 (2001).

52. Takahashi, Y., Kagawa, K., Svensson, E. I. & Kawata, M. Evolution of increased
phenotypic diversity enhances population performance by reducing sexual harassment in
damselflies. *Nature Communications* **5**, 1–7 (2014).

53. Rowe, L., Chenoweth, S. F. & Agrawal, A. F. The genomics of sexual conflict. *The*
American Naturalist **192**, 274–286 (2018).

54. Sayadi, A. *et al.* The genomic footprint of sexual conflict. *Nature Ecology & Evolution* **3**,
1725–1730 (2019).

55. Wellenreuther, M. *et al.* Molecular and ecological signatures of an expanding hybrid
zone. *Ecology and Evolution* **8**, 4793–4806 (2018).

56. Okude, G., Fukatsu, T. & Futahashi, R. Interspecific crossing between blue-tailed
damselflies *Ischnura elegans* and *I. Senegalensis* in the laboratory. *Entomological*
Science **23**, 165–172 (2020).

57. Montgomery, E. A., Huang, S. M., Langley, C. H. & Judd, B. H. Chromosome
rearrangement by ectopic recombination in *Drosophila melanogaster*: Genome structure
and evolution. *Genetics* **129**, 1085–1098 (1991).

58.   Delprat, A., Negre, B., Puig, M. & Ruiz, A. The transposon Galileo generates natural
985   chromosomal inversions in *Drosophila* by ectopic recombination. *PloS one* **4**, e7883
986   (2009).

987   59.   Andrés, J. A. & Cordero, A. The inheritance of female colour morphs in the damselfly
988   *Ceriagrion tenellum* (Odonata, Coenagrionidae). *Heredity* **82**, 328–335 (1999).

989   60.   Kent, T. V., Uzunović, J. & Wright, S. I. Coevolution between transposable elements and
990   recombination. *Philosophical Transactions of the Royal Society B: Biological Sciences*
991   **372**, 20160458 (2017).

992   61.   Thompson, M. J. & Jiggins, C. Supergenes and their role in evolution. *Heredity* **113**, 1–8
993   (2014).

994   62.   Willink, B. & Svensson, E. I. Intra-and intersexual differences in parasite resistance and
995   female fitness tolerance in a polymorphic insect. *Proceedings of the Royal Society B:*
996   *Biological Sciences* **284**, 20162407 (2017).

997   63.   Abbott, J. K. & Gosden, T. P. Correlated morphological and colour differences among
998   females of the damselfly ischnura elegans. *Ecological Entomology* **34**, 378–386 (2009).

64.   West-Eberhard, M. J. *Developmental plasticity and evolution*. (Oxford University Press,
2003).

65.   Svensson, E. I., Abbott, J. K., Gosden, T. P. & Coreau, A. Female polymorphisms, sexual
conflict and limits to speciation processes in animals. *Evolutionary Ecology* **23**, 93
(2009).

66.   Colgan, D. J. *et al.* Histone H3 and U2 snRNA DNA sequences and arthropod molecular
evolution. *Australian Journal of Zoology* **46**, 419–437 (1998).

67.   Shafin, K. *et al.* Nanopore sequencing and the Shasta toolkit enable efficient de novo
assembly of eleven human genomes. *Nature Biotechnology* **38**, 1044–1053 (2020).

68.   Phillippy, A. M., Schatz, M. C. & Pop, M. Genome assembly forensics: Finding the
elusive mis-assembly. *Genome Biology* **9**, 1–13 (2008).

69.   Li, H. Minimap2: Pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–
3100 (2018).

70.   Shafin, K. *et al.* Haplotype-aware variant calling with PEPPER-Margin-DeepVariant
enables high accuracy in nanopore long-reads. *Nature Methods* **18**, 1322–1332 (2021).

71.   Guan, D. *et al.* Identifying and removing haplotypic duplication in primary genome
assemblies. *Bioinformatics* **36**, 2896–2898 (2020).

72.   Zimin, A. V. *et al.* Hybrid assembly of the large and highly repetitive genome of -
textitAegilops tauschii, a progenitor of bread wheat, with the MaSuRCA mega-reads
algorithm. *Genome Research* **27**, 787–792 (2017).

73. Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E. M.
BUSCO: Assessing genome assembly and annotation completeness with single-copy
orthologs. *Bioinformatics* **31**, 3210–3212 (2015).

74. Alonge, M. *et al.* Automated assembly scaffolding elevates a new tomato system for
high-throughput genome editing. *bioRxiv* (2021) doi:10.1101/2021.11.18.469135.

75. Delcher, A. L., Phillippy, A., Carlton, J. & Salzberg, S. L. Fast algorithms for large-scale
genome alignment and comparison. *Nucleic Acids Research* **30**, 2478–2483 (2002).

76. Hao, Z. *et al. Rideogram: Drawing SVG graphics to visualize and map genome-wide*
data on idiograms. (2020).

77. R Core Team. *R: A language and environment for statistical computing.* (R Foundation
for Statistical Computing, 2021).

78. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv preprint arXiv:1303.3997 (2013).

79. Van der Auwera, G. A. *et al.* From FastQ data to high-confidence variant calls: The
genome analysis toolkit best practices pipeline. *Current Protocols in Bioinformatics* **43**,
11.10.1–11.10.33 (2013).

80. Li, H. A statistical framework for SNP calling, mutation discovery, association mapping
and population genetical parameter estimation from sequencing data. *Bioinformatics* **27**,
2987–2993 (2011).

81. Smith, A. F. A., Hubley, R. & Green, P. RepeatMasker open-4.0. (2013-2015)
doi:http://www.repeatmasker.org.

82. Girgis, H. Z. Red: An intelligent, rapid, accurate tool for detecting repeats de-novo on the
genomic scale. *BMC Bioinformatics* **16**, 1–19 (2015).

83. Purcell, S. *et al.* PLINK: A tool set for whole-genome association and population-based
linkage analyses. *The American Journal of Human Genetics* **81**, 559–575 (2007).

84. Kokot, M., Długosz, M. & Deorowicz, S. KMC 3: Counting and manipulating *k*-mer
statistics. *Bioinformatics* **33**, 2759–2761 (2017).

85. Zhou, X. & Stephens, M. Genome-wide efficient mixed-model analysis for association
studies. *Nature Genetics* **44**, 821–824 (2012).

86. Camacho, C. *et al.* BLAST+: Architecture and applications. *BMC Bioinformatics* **10**, 1–9
(2009).

87. Li, H. *et al.* The sequence alignment/map format and SAMtools. *Bioinformatics* **25**,
2078–2079 (2009).

88. Pedersen, B. S. & Quinlan, A. R. Mosdepth: Quick coverage calculation for genomes and
exomes. *Bioinformatics* **34**, 867–868 (2018).

89. Korunes, K. L. & Samuk, K. pixy: Unbiased estimation of nucleotide diversity and
divergence in the presence of missing data. *Molecular Ecology Resources* **21**, 1359–1368
(2021).

90. Hudson, R. R., Slatkin, M. & Maddison, W. P. Estimation of levels of gene flow from
DNA sequence data. *Genetics* **132**, 583–589 (1992).

91. Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158
(2011).

92. Sedlazeck, F. J. *et al.* Accurate detection of complex structural variations using single-
molecule sequencing. *Nature Methods* **15**, 461–468 (2018).

93. Belyeu, J. R. *et al.* Samplot: A platform for structural variant visual validation and
automated filtering. *Genome Biology* **22**, 1–13 (2021).

94. Bailly-Bechet, M., Haudry, A. & Lerat, E. 'One code to find them all': A perl tool to
conveniently parse RepeatMasker output files. *Mobile DNA* **5**, 1–15 (2014).

95. Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology* **37**,
907–915 (2019).

96. Pertea, M. *et al.* StringTie enables improved reconstruction of a transcriptome from
RNA-seq reads. *Nature Biotechnology* **33**, 290–295 (2015).

97. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. EdgeR: A Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–
140 (2010).

98. Robinson, M. D. & Oshlack, A. A scaling normalization method for differential
expression analysis of RNA-seq data. *Genome Biology* **11**, 1–9 (2010).

99. Robinson, M. D. & Smyth, G. K. Moderated statistical tests for assessing differences in
tag abundance. *Bioinformatics* **23**, 2881–2887 (2007).

100. Robinson, M. D. & Smyth, G. K. Small-sample estimation of negative binomial
dispersion, with applications to SAGE data. *Biostatistics* **9**, 321–332 (2008).

101. Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: A practical and
powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B
(Methodological)* **57**, 289–300 (1995).

102. Pertea, G. & Pertea, M. GFF utilities: GffRead and GffCompare. *F1000Research* **9**,
(2020).

103. Jones, P. *et al.* InterProScan 5: genome-scale protein function classification.
Bioinformatics **30**, 1236–1240 (2014).

104.   Huerta-Cepas, J. *et al.* eggNOG5.0: A hierarchical, functionally and phylogenetically
1089   annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids*
1090   *Research* **47**, D309–D314 (2019).

1091   105.   Willink, B. *et al.* Data from: The genomics and evolution of inter-sexual mimicry and
1092   female-limited polymorphisms in damselflies. *Zenodo* (2023).

1093   106.   Takahashi, Y. Testing negative frequency-dependent selection: Linking behavioral
1094   plasticity and evolutionary dynamics. *Bull Kanto Branch Ecol Soc Jpn* **59**, 8–14 (2011).

1095

1096   Figure 1

1097

1098   Figure 2

1099

1100   Figure 3

1101

1102   Figure 4

1103

1104   Figure 5

1105

1106   Figure 6

1107

1108　　　　　　　　　　Extended Data Figure 1

1109

1110   Extended Data Figure 2

1111

1112   Extended Data Figure 3

1113

1114   Extended Data Figure 4

1115

1116   Extended Data Figure 5

1117

1118   Extended Data Figure 6

1119

1120   Extended Data Figure 7

1121

1122   Extended Data Figure 8