A single-cell atlas of the miracidium larva of the human blood fluke *Schistosoma*

***mansoni*: cell types, developmental pathways and tissue architecture**

Teresa Attenborough[1,2]*, Kate A. Rawlinson[1,3✉]*, Carmen L. Diaz Soria[1], Kirsty Ambridge[1],

Geetha Sankaranarayanan[1], Jennie Graham[1], James A. Cotton[1,4], Stephen R. Doyle[1], Gabriel

Rinaldi[1,5], Matthew Berriman[1,2✉] (*These authors contributed equally).

1. Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, CB10 1SA, UK

2. Wellcome Centre for Integrative Parasitology, School of Infection and Immunity,

College of Medical, Veterinary & Life Sciences, University of Glasgow, 120

University Place, Glasgow G12 8TA, UK

3. Josephine Bay Paul Center, Marine Biological Laboratory, Woods Hole,

Massachusetts, USA.

4. Wellcome Centre for Integrative Parasitology, School of Biodiversity, One Health

and Veterinary Medicine, College of Medical, Veterinary & Life Sciences, University

of Glasgow, Bearsden Road, Glasgow G61 1QH, UK

5. Department of Life Sciences, Aberystwyth University, Edward Llwyd Building,

Penglais Campus, Aberystwyth SY23 3DA, UK

Abstract

Schistosomes are parasitic flatworms that cause the water-borne disease schistosomiasis,

affecting millions of people worldwide. The miracidium larva of schistosomes represents the

first post-embryonic stage of development and is critical to transmission. After hatching, a

miracidium infects a freshwater snail and transforms into a mother sporocyst, where its stem

cells generate daughter sporocysts that give rise to many human-infective cercariae larvae. To

understand this important life cycle stage at the cellular and molecular levels, we have used

single-cell RNA sequencing, *in situ* hybridisation and image analysis to create a whole-body

cell atlas of the miracidium larva of *Schistosoma mansoni*. Our atlas shows that each

miracidium is composed of ~365 cells and 19 transcriptionally distinct cell types. We show

that 93% of miracidium cells are somatic (57% neural, 19% muscle, 13% epidermal

(tegument), 2% parenchyma, 2% protonephridia), and the remaining 7% are stem cells.

Cellular diversity within tissue types is revealed, and is highest in neurons. There are two

stem cell populations, and they show different activation and potency states. Trajectory

analysis indicates that one stem cell population is the origin of the tegument lineage and the

other likely contains pluripotent cells. Furthermore, each of these stem populations is

transcriptionally distinct based on sex-linked gene expression in male and female larvae.

Through single cell transcriptomics and *in-situ* hybridisation we identified every cell in the

whole organism revealing the organisation of the miracidium. This single cell atlas provides

the foundation to understand the development and interaction of cell types and tissues as they

change over a life cycle that is characterised by complex morphological changes.

Introduction

The parasitic blood fluke *Schistosoma mansoni* (Phylum Platyhelminthes) is a causative

agent of the neglected tropical disease schistosomiasis, which is responsible for a chronic

disease burden equivalent to 1.0–2.6 million Disability Adjusted Life Years and over 11,000

human deaths annually (WHO, 2023; IHME, 2019). The life cycle of *S. mansoni* is complex,

involving parasitic stages in two hosts (a snail and a mammal), with short-lived, freshwater

infectious larval stages in between. Mature male and female worms reside in the portal veins

of the mammalian host, and fertilisation of the zygote occurs inside the female worm (Jurberg

et al., 2009). Embryogenesis proceeds as the egg is laid in the host and continues over the

ensuing six days (Ashton *et al*., 2001). During this time, the egg passes through the host

vascular endothelium into the intestine and is passed in faeces into the outside environment

(Costain *et al*., 2018). If the egg lands in freshwater, a miracidium larva hatches out, swims

and infects a snail host. Once inside the snail, the miracidium transforms into the mother

sporocyst, and its stem cells (historically known as 'germinal cells') divide and differentiate

to produce many daughter sporocysts that migrate within the snail. In the daughter

sporocysts, a second round of embryogenesis generates the cercariae larvae by asexual

reproduction from a stem cell. The cercariae emerge from the snail, swim towards and infect

water-exposed mammals including humans, penetrating the skin. At the point of infection,

the cercariae shed their tails, and once inside the mammal the tegument (skin) is remodelled

during metamorphosis to the schistosomulum stage. The schistosomulum then moves into the

blood vessels and, over the following five weeks, grows into a juvenile stage that migrates

from the lungs to the liver and to the hepatic portal vein where sexually dimorphic male and

female worms pair, mature and mate. Thus, during the development from egg to adult, the

body axes are established three times, once during the development of the miracidia, once

during the development of the daughter sporocyst and finally during the development of the

cercariae. Additionally, there are two rounds of embryogenesis, first from fertilised zygote to

the mature miracidium, and second, the neo-embryogenesis from a single stem cell in the

daughter sporocyst to a mature cercaria.

Enabled by the advent of single-cell RNA sequencing (scRNA-seq), molecular classification

of cell types has been initiated for *Schistosoma mansoni* (Wang *et al*., 2018; Tarashansky *et*

al., 2019; Li *et al*., 2020; Diaz *et al*., 2020; Wendt *et al*., 2020, Diaz & Attenborough *et al*.,

2024). These studies have opened the door to investigate schistosome developmental cell

biology and to identify new targets for controlling infection and transmission of the disease.

To date, scRNA-seq studies have identified heterogeneity amongst stem cells that drive

schistosome development and reproduction, classified a variety of differentiated cell types

across the life cycle and revealed several key genes and their native cell types that are

essential for parasite survival and propagation (Nanes Sarfati *et al*., 2021). Single-cell RNA-

seq studies for the whole animal have only been carried out on intra-molluscan and intra-

mammalian stages to date: the mother sporocyst (Diaz & Attenborough *et al*., 2024), the

schistosomulum (Diaz *et al*., 2020), juvenile (3.5 weeks post-infection of mouse, Li *et al*.,

2021) and adult (Wendt *et al*., 2020). Single cell studies on stem cells specifically, however,

have only been undertaken in the mother sporocyst (Wang *et al*., 2018) and juvenile stages

(2.5 and 3.5 weeks post-infection)(Tarashansky *et al*., 2019).

An obvious gap in the coverage of the life cycle is the lack of single cell data for the earliest

developmental time point currently accessible – the miracidium larva; earlier embryonic

stages are inaccessible due to the impenetrable egg capsule. The miracidium is a short-lived,

non-feeding stage that relies on internal stores of energy to sustain swimming, and host

location and infection (Maldonado and Matienzo, 1948; Chernin, 1968, Pan, 1980). The role

of the miracidium and its differentiated somatic cells is to carry stem cells to a favourable

environment inside the snail where they can proliferate to generate daughter sporocysts, and

subsequently cercariae. If the miracidium fails to infect a snail within 12 hours it dies.

At the ultrastructural level, a rich set of cell types have been previously identified in the

miracidium and classified into eight major systems: musculature, nervous, penetration gland

cells, excretory, interstitial (parenchyma), stem/germinal cells, epithelial (ciliary plates and

epidermal ridges of the tegument cells) and terebratorium (an apical modified epidermal plate

with sensory structures and gland openings) (Pan, 1980).

To characterise this critical transmission stage, we have generated a single cell transcriptome

atlas for the miracidium. Our atlas provides molecular definitions for all described cell types

(Pan, 1980) and indicates the functions of different cell types within tissues. Multiplexed *in*

situ validation of cell type marker genes enabled the cell types to be spatially resolved within

the larva. Furthermore, the nuclear segmentation of the cells expressing tissue-specific genes

has revealed the relative contribution of each tissue type to the animal and each cell in the

larva has been assigned to a tissue. A key finding is the identification of two stem cell

populations in a miracidium, and advances in our understanding of the genome (Buddenborg

et al., 2021) has enabled us to determine that these two stem populations are transcriptionally

distinct between male and female larvae due to sex-linked gene expression. To predict the

fate of these two stem populations we carried out a trajectory analysis, combining scRNA-seq

data from miracidia and mother sporocysts, and this indicated that one stem cell population

likely contains the pluripotent cells that will develop into the cercariae, while the other

population gives rise to the somatic tissues, including the tegument. Similarities in gene

expression with the adult suggest that there is a tegument developmental program that is

redeployed at multiple stages of the complex life cycle.

Results

Single-cell RNA-seq of 20,478 miracidia cells reveals 19 cell types.

There are ~365 nuclei in a miracidium (median = 365, larva #1: 342, larva #2: 365, larva #3:

377). Some cell types, such as the apical gland and epithelial/tegument cells, are

multinucleate; and others, such as the ciliary plates may be anucleate (Pan, 1980, but cf

Meulemen *et al*., 1978). Therefore, the number of nuclei provides an approximation for the

actual number of cells (Figure 1A).

We performed scRNA-seq on a pool of ~20,000 mixed-sex miracidia, collected within four

hours of hatching from eggs (Figure 1B). Following dissociation, four samples of the cell

suspensions were collected; two were enriched for live cells using fluorescence-activated cell

sorting (FACS), and two were left unsorted to capture the maximum cellular diversity for the

atlas (Figure 1B). Using the droplet-based 10X Genomics Chromium platform,

transcriptome-sequencing data were generated from a total of 33,391 cells, of which 20,478

passed strict quality-control filters, resulting in a median of 858 genes and 1,761 median UMI

counts per cell (Supplementary Table 1). Given that an individual miracidium comprises

~365 cells (Figure 1A), the number of quality-controlled cells theoretically represents >56-

fold coverage of each cell in the larva.

Using Seurat (version 4.3.0) (Hao *et al*., 2021), 19 distinct clusters of cells were identified,

along with putative marker genes best able to discriminate between the populations (Figure

1C & D and Supplementary Table 2 and 3). We used Seurat's JackStraw and ElbowPlot,

along with molecular cross-validation to select the number of principal components, and

Seurat's clustree to select a resolution where clusters were stable (Hao *et al*., 2021). Manually

curated lists of previously defined cell-type-specific genes from later developmental stages

(Diaz *et al*., 2020; Wendt *et al*., 2020; Nanes Sarfati *et al*., 2021) were compared against the

list of putative markers generated in this analysis. The expression of genes from both these

lists in the miracidia single-cell data was then examined to identify the cell types that each

Seurat cluster represented. Based on these markers, the 19 clusters were resolved into the

following cell populations: two muscle-like (3,934 cells), five resembling neurons (3,255

cells), one ciliary plate (40 cells), one tegumental (366 cells), one protonephridial (539 cells),

two parenchymal (1698 cells) and seven clusters resembling stem cells (10,686 cells) (Figure

1C).

There were contributions from sorted and unsorted samples in almost all clusters (except

ciliary plates). We found that some cell/tissue types had similar recovery from both methods

(e.g. Stem A, Muscle 2, and Tegument), others were preferentially recovered by sorting (e.g

Neuron 1, Neuron 4, and Stem E), and some were depleted by sorting (e.g. Parenchyma 1,

Protonephridia, and Ciliary plates) (Supplementary Figure 1), Supplementary Table 4). This

variation in recovery, therefore, enabled us to maximise the discovery and inclusion of

different cell types in the atlas.

Orthogonal body wall muscle fibres are transcriptionally distinct

The two muscle clusters were identified based on the expression of previously described

muscle-specific genes; *paramyosin* (Smp_085540), *troponin t* (Smp_179810) and *titin*

(Smp_126240) (Wendt *et al*., 2020; Diaz *et al*., 2020; Diaz & Attenborough *et al*., 2024), as

well as other differentially expressed markers; e.g. a putative collagen alpha chain

(Smp_159600) (Figure 2A). A crp1/csrp1/crip1 homolog (Smp_087250), encoding a

conserved transcriptional regulator of muscle (Tarashansky *et al*., 2021) was also expressed

in both muscle clusters. High expression of *paramyosin* (*PRM*; Smp_085540) was seen in

muscle cell clusters 1 and 2 (Figure 2A). *In situ* hybridisation (ISH) (Figure 2B & C;

Supplementary Video 1) and image analysis (Supplementary Video 2) identified 74 nuclei

surrounded by *PRM* transcripts (i.e. ~19% of cells in a miracidium are muscle cells).

Counterstaining with phalloidin (to label F-actin filaments) revealed an orthogonal grid of

circular and longitudinal body wall muscles and from this we identified which *PRM*⁺ nuclei

belonged to each muscle subtype. The nuclei of the circular muscles formed two distinct

bilaterally symmetrical lines that ran peripherally from pole to pole of the larva (~28 nuclei in

total). Thirty-three more *PRM*⁺ nuclei sat regularly spaced between the circular muscle nuclei

and corresponded to longitudinal muscles. Another 13 *PRM*⁺ nuclei formed a unilateral

cluster adjacent to the apical gland cell (identifiable by its four nuclei; Pan, 1980) (Figure

2Cii; Supplementary Video 1 and 2).

Muscle cluster 1 (2,407 cells) was distinguished from muscle cluster 2 by markedly higher

expression of a Kunitz-type protease inhibitor (Smp_052230; Figure 2A), and ISH showed

Smp_052230 transcripts along the two peripheral lines of circular muscle nuclei (Figure 2D).

Subpopulations of muscle cluster 1 also expressed orthologs of *Wnt-11-1* (Smp_156540) and

Notum (Smp_015820), known anterior-posterior body axis patterning genes, where Notum

acts as an inhibitor of Wnt signalling (Peterson and Reddien, 2009; Kakugawa *et al.,*

2015)(Figure 2A). *Notum* (Smp_015820) was highly expressed in the cluster of 13 muscle

nuclei adjacent to the apical gland cell (Figure 2E). *Wnt-11-1* (Smp_156540) was expressed

at the posterior pole, in seven of the Kunitz-type protease inhibitor-expressing circular muscle

cells (Figure 2D; Supplementary Video 3). *Notum* was also expressed, albeit weakly, in some

nuclei of circular muscles, including those expressing *Wnt-11-1* (Supplementary Video 4).

Muscle cluster 2 (1,355 cells) was distinguished by higher expression of genes encoding a

PDZ domain protein (Smp_311170) and *Calcium-transporting ATPase* (Smp_136710), as

well as the unique expression of a putative calcium-activated potassium channel

(Smp_156150), and the transcription factor *MyoD* (Smp_167400)(Figure 2A); ISH of the

latter two genes showed expression only in the longitudinal body wall muscles (Figure 2F;

Supplementary Video 5). Orthologs of other axial patterning genes, *Netrin receptor UNC5*

(Smp_172470), *Wnt5* (Smp_145140)*,* and a putative wnt inhibitor frizzled, frzb2

(Smp_062560)(Witchley *et al*., 2013), showed distinct expression in Muscle 2 cells.

The higher expression of Muscle 1 markers in circular body wall muscles (BWM) and Muscle 2 genes in the longitudinal BWM revealed distinct transcriptomic signatures for these two types of myocyte that make up the orthogonal body wall musculature. In fact, differential expression analysis showed that *Wnt-11-1* (circular) and *MyoD* (longitudinal) were among the most differentially expressed genes between the two muscle clusters (Supplementary Table 5).

Neural abundance and diversity

We uncovered five clusters expressing the neural markers *Complexin* (*cpx*; Smp_050220)*, prohormone convertase 2* (*PC2*; Smp_077980) and Smp_068500 (a neural marker in adult worms, Wendt *et al*., 2020) (Figure 3A). *Complexin* was expressed in and around at least 209 nuclei, indicating that 57% of cells in the miracidium were neurons; out of which 129 cpx^+ nuclei formed the nuclear rind of the brain (or neural mass/ring), and the remaining 80 were situated peripherally, either anterior or posterior to the brain (Figure 3B). Sensory organelles were identified for peripheral neural cell types. We were able to validate neuron clusters 2 - 5 using ISH of specific marker genes, and performed an *in silico* analysis of Neuron 1.

Neuron cluster 1 was large (1,470 cells), and expressed neuron-marker *complexin*, the neuroendocrine precursor *7B2* (Smp_073270) and an uncharacterised gene (Smp_068500). At this resolution, there were no clear marker genes restricted to Neuron 1. We explored whether Neuron 1 could be further subdivided into transcriptionally distinct cells by subclustering (Supplementary Figure 2; Supplementary Table 6) using the self-assembling manifold (SAM) algorithm (Tarashansky *et al*., 2019) with ScanPy (Wolf *et al*., 2018), given its reported strength in discerning subtle variation in gene expression (Tarashansky *et al*., 2019), although a similar topology was subsequently found using Seurat. Three subclusters of Neuron 1 expressed neuropeptide precursor genes amongst their top five markers; *PWamide* (Smp_340860)(Wang *et al*., 2016) in subcluster 14, *Neuropeptide F* (Smp_088360) in subcluster 8*,* and *Neuropeptide Y* (Smp_159950) in subcluster 13 (Supplementary Figure 2); and each was expressed in the nuclear rind of the brain (Supplementary Figure 3). Another subcluster was composed of putative photoreceptor cells (subcluster 7 - Supplementary Figure 2, Supplementary Table 6) because their top-ten markers include two opsins (Smp_180350 and Smp_332250) – identified here as a rhabdomeric opsin and a peropsin, respectively (Supplementary Figure 4). Subcluster 7 also expressed other conserved phototransduction cascade genes, such as a 1-phosphatidylinositol 4,5-bisphosphate

phosphodiesterase (Smp_245900) and a beta-arrestin (Smp_152280) (Supplementary Figure

2, Supplementary Table 6).

Neuron 2 expressed two further rhabdomeric opsins (Smp_180030 and Smp_104210;

Supplementary Figure 4), but across the cluster of 282 cells, the most expressed marker was

the uncharacterised gene Smp_201600. ISH revealed three strongly Smp_201600$^+$ cells, with

nuclei in a median posterior location of the brain, and three cells with weaker expression in a

more anterior position (Figure 3C, Supplementary Video 6). Other top markers include *7B2,*

a guanylate kinase (Smp_137940) and a putative homeobox protein (Smp_126560).

Neuron 3 was a small cluster (47 cells) expressing *complexin* and *7B2*. The uncharacterised

gene, Smp_071050, was unique to and highly expressed throughout this population (Figure

3A). ISH showed expression in ~6 cells with projections that extended into the brain and to

the periphery, three of which sat adjacent to three Neuron 2 cells. There was also weak

expression in the four nuclei of the apical gland cell (Figure 3Ci, Supplementary Video 6).

STRING analysis of the top 100 markers of Neuron 3 predicted two protein interaction

networks with functional enrichment; these were associated with 'synapses' and 'calcium'

(Supplementary Figure 6). In Neuron clusters 1, 2 and 3, there was an enrichment for genes

annotated with Gene Ontology (GO) terms related to 'neuropeptide signalling pathway' and

'neurotransmitter transport' (Supplementary Table 7). There were, therefore, at least 5, and

likely several more, transcriptionally distinct cell types in the brain mass (we validated three

subclusters of Neuron 1, as well as Neurons 2 & 3).

Cells from neuron clusters 4 and 5 had nuclei peripheral to the brain. Neuron 4 formed a

large (1,140 cells), discrete cluster. The top marker was an uncharacterised gene,

Smp_319480 (Figure 3A), that, based on protein-structure prediction using I-TASSER (Yang

& Zhang, 2015), was similar to human mTORC1 (PDB entry 5H64; TM-score 0.85). *In situ*

expression revealed ~26 Smp_319480$^+$ cells; four at the anterior end, with projections

leading to the body wall in the first tier of ciliary plates, and ~22 sat lateral and directly

posterior to the brain. These latter cells formed four clusters of ~3–7 cells, with Smp_319480

transcripts distributed around the nuclei and along projections leading into the brain and

extending out to the body wall terminating between the second and third tier of ciliary plates

(Figure 3D, Supplementary Video 9 and 10). Twenty-three multi-ciliated nerve endings

formed a girdle around the larva between the second and third tier of ciliary plates

(Supplementary Figure 3C), and Smp_319480 transcripts were expressed at the base of these

cilia (Supplementary Figure 3D); some transcripts also reached the nuclei of circular body

wall muscles (Supplementary Figure 3E), suggesting these cells may function as sensory-

motor neurons. Two genes, identified as top markers and co-expressed in many Neuron 4

cells, encode likely orthologues of *Polycystin 1* (Smp_165230) and *Polycystin 2*

(Smp_165660) (Supplementary Figure 5A), with the latter prediction strongly supported by a

Foldseek search (van Kempen *et al*., 2023) against the protein database (PDB). Polycystins 1

and 2 form a heterodimeric ion channel required for oscillating calcium concentrations within

vertebrate cilia (Kim *et al*., 2016). A Sox transcription factor, Smp_301790, was also

expressed in Neuron 4 and co-expressed with *Polycystin 1* and *2* (Supplementary Figure 5B).

Other top markers included a putative potassium channel (Smp_141570), a cAMP-dependent

protein kinase (Smp_079010), a GTP-binding protein (Smp_045990) and an ortholog of

human *fibrocystin-L* (Smp_303980). The GO terms 'cilium', 'regulation of apoptotic

process' and 'potassium ion transmembrane transport' were significantly enriched, and

STRING analysis showed networks enriched in 'plasma membrane-bound cell projection'

(Supplementary Table 7 Supplementary Figure 6).

Neuron 5 cluster comprised 276 cells. An EF-hand domain-containing protein, Smp_343710,

was highly expressed in 55% of these cells. ISH showed expression in 10-20 cells whose

nuclei sat outside of, and anterior to, the brain, in and around the paired lateral glands and

their secretory ducts (Supplementary Video 11). There was also expression in a pair of

bilaterally symmetrical bulbous protrusions (called lateral sensory papilla and thought to be

depth sensors, Pan, 1980)(Figure 3F). STRING analysis revealed networks associated with

'synapses', 'ion channel binding' and 'cation transmembrane transporter activity'

(Supplementary Figure 6).

We expected to find a discrete cluster(s) for the penetration glands, and that it would show

similarities to the neural clusters (as glandular cells arise from neuroglandular precursor cells

in other animals, such as the sea anemone, *Nematostella vectensis*, Steger *et al*., 2022). There

are three gland cells per miracidium - one apical and two lateral (Pan, 1980). While the top

marker gene for Neuron 3 was expressed in the four nuclei of the apical gland (Figure 3Ci),

and there was expression of a Neuron 5 marker in the lateral glands (Figure 3Fi), these genes

were expressed in many other cells as well. This suggests that gland cells are either sub-

clusters of Neurons 3 and 5 or they were not captured in the 10X GEMs. To characterise the

transcriptomes of the gland cells, we carried out plate-based single cell RNA-seq (New England BioLabs) on 37 manually selected gland cells (Supplementary Table 8). The biological process GO terms enriched in the most abundant genes (top 200 genes by median transcripts per million, Supplementary Table 8) included translation, protein folding and ATP synthesis coupled electron transport (Supplementary Table 9). The gland cells showed co-expression of neural markers (*Complexin* and *7B2*) and muscle markers *Myosin light chain 1* (Smp_045200), *Troponin t* (Smp_179810) and *Actin-2* (Smp_307020) (Supplementary Figure 7). To identify genes that were specific to gland cells, we searched for differentially expressed genes in the gland cells compared to the other cells in the plate-based scRNA-seq data (i.e. ciliary plates and unknown cells) (Supplementary Table 10 and 11), and then looked at their expression in the 10X-based scRNA-seq data. A lack of, or minimal, expression of these markers in the 10X-based scRNA-seq data, would support the hypothesis that they were specific to the gland cells. Five venom antigen-like (VAL) genes (belonging to the SCP/TAPS or CAP) superfamily) were expressed in the gland cells; Smp_120670 (VAL 5) and Smp_070250 (VAL 15) in over 60% of the gland cells, and Smp_176180 (VAL 9), Smp_002630 (VAL 2) and Smp_331830 (SCP protein) in less than 50%. Many gland cells expressed multiple VAL genes (Supplementary Figure 7). In the 10X-based scRNA-seq data, very few cells expressed these VAL genes (1-5 cells for each gene, distributed between Neurons 1 and 5, Muscle 1, Parenchyma 2, and Stem F and G clusters). The VAL genes expressed in the gland cells are uniquely expressed during the egg, miracidia and sporocyst stages (Yoshino *et al*., 2014; Lu *et al*., 2018; Buddenborg *et al*., 2023), and it has been reported that VAL 9 (Smp_176180) induces the differential expression of an extracellular matrix remodelling gene in the snail host (Yoshino *et al*., 2014). Other differentially expressed genes in the gland cells included Smp_303400 (an HSP70 homolog), Smp_179250 (*α-galactosidase/α-N-acetylgalactominidase*), Smp_317530 (uncharacterised protein), and two putative major egg antigens Smp_185680 and Smp_303690 (Supplementary Figure 7). Eight putative secreted proteins previously identified in miracidia and *in vitro* transformed miracidia (Wang *et al*., 2016; Wu *et al*., 2006) were expressed in the glands cells (and not in other cell types based on the 10X data), and five of these showed relatively high expression; Smp_302170 (*HSP70* homolog), Smp_303400 (*HSP70* homolog), Smp_049230 (*HSP20*), Smp_304250 (*purine nucleoside phosphorylase*), and Smp_302180 (putative heat shock protein 70)(Supplementary Figure 7). Typical of venom glands in other animals, the transcriptomes of the miracidia gland cells are enriched in genes involved in protein

translation, stabilisation and stress response to cope with mass protein production (Zancolli *et*
al., 2021).

Ciliary plate gene marker identifies six submuscular cell bodies

A small cluster of 40 cells (almost exclusively from the unsorted samples) was identified near
the tegument cluster (Figure 1C). The top markers were Smp_096390, a 16 kDa calcium-
binding protein (CaBP) and *p25 alpha* (Smp_097490, a marker for ciliated neurons in adult
worms, Wendt *et al*., 2020) (Figure 3A). *In situ* hybridisation revealed CaBP and *p25 alpha*
expression in the ciliary plates; epithelial cells with thousands of motile cilia used for
swimming (Figure 3G and Supplementary Figure 3). Both genes were also expressed in six
previously undescribed cells that sent a projection out to the exterior (between the first and
second tier of ciliary plates as seen with phalloidin staining, Figure 3G) and whose nuclei sat
below the body wall musculature (Supplementary Videos 7 & 8; Supplementary Figure 3F).
Given the association of *p25 alpha* with ciliated neurons in adults (Wendt *et al.*, 2020),
expression in the sub-muscular cells suggests they project cilia into the external environment
and may be sensory cells. The expression of CaBP and *p25* in both the ciliary plates and
submuscular cells suggests that these anatomical structures are functionally linked. GO terms
for this cluster were related to 'mitochondrial transmembrane transport' and 'microtubule-
based process' (Supplementary Table 7), and STRING analysis showed two interaction
networks: one associated with 'carbon metabolism and mitochondria envelope', the second
associated with 'cytoplasmic dynein complex' and 'dynein light chain superfamily'
(Supplementary Figure 6). Further plate-based scRNA-seq of 21 individual ciliary plates
confirmed the high expression of CaBP (Smp_096390) and *p25 alpha* (Smp_097490), with
more than two-fold higher expression of CaBP than any other gene by median TPM
(transcripts per million) (Supplementary Table 12). Although the ciliary plates expressed *p25*
(a ciliated neuron marker in the adult, Wendt *et al.,* 2020), they did not express many of the
typical neural markers, e.g. *complexin* and *7B2* (Supplementary Figure 7).
362
Tegumental cells expressed *MEG6* and are located in the posterior two-thirds of the
larva.
We identified a cluster of 366 cells as tegumental based on the expression of markers for
tegument (skin) in the adult worm (Wendt *et al.*, 2020), e.g. genes encoding *dynein light*
chain (Smp_302860, Smp_200180), a DUF3421 domain-containing protein (Smp_331910)

and micro exon gene *MEG6* (Smp_163710). Two other dynein light chain genes were top

markers (Smp_201060, Smp_312630), and both a tetraspanin (Smp_303290) and *Gelsolin*

(Smp_008660) were specific to this cluster. *Meg6* (Smp_163710) was expressed in 66% of

the cells in this cluster; ISH showed 46 *Meg6*[+] cells, and the nuclei were in the posterior two-

thirds of the larva. The nuclei sat below the body wall muscle, and cytoplasmic protrusions

reached between muscle filaments and formed the epidermal ridges between the ciliary plates

(Figures 4C & D). These cells correspond to the epidermal ridge cells described by Pan

(1980). A second tegument marker, a dynein light chain (Smp_200180), showed overlapping

expression with *Meg6* (Supplementary Video 15). Enriched GO terms for tegument genes

included 'microtubule-based process' and 'protein folding' (Supplementary Table 7).

STRING analysis revealed a network of 25 genes associated with protein localisation to the

ER, folding, export and post-translational protein targeting to the membrane (Supplementary

Figure 8).

Spatial localisation of top marker transcripts identifies a simple protonephridial system

There were 536 cells in the protonephridial cluster, expressing genes common to excretory

systems in other animals (Gąsiorowski *et al.,* 2021), e.g., the structural protein *nephrin*

(Smp_130070) and the transcription factor *odd-skipped-related* (Smp_152210). Eighty

percent of the cells expressed the protonephridia marker Smp_335600 (ShKT-domain

containing protein) (Nanes Sarfati *et al.*, 2021) (Figure 4A). Smp_335600 was expressed

around six nuclei in a bilaterally symmetrical pattern (Figure 4Ei) (Supplementary Video 12).

On each side, Smp_335600 transcripts were distributed along the s-shaped path of an

excretory tubule, passing its nucleus, and connecting the anterior and posterior flame cells

(Figure 4Ei). Within the flame cells, expression was detected around the nuclei rather than

the barrel (Figure 4Eii). The four ciliated flame cells and two excretory ducts comprise the

excretory system of the miracidium. Accordingly, STRING (Supplementary Figure 8) and

GO analyses (Supplementary Table 7) of this cluster confirmed enrichment for terms

including 'cilium organisation', 'axoneme assembly', 'cytoskeleton organisation*'* and

'microtubule-based process*'*.

Two distinct parenchyma clusters predicted in the miracidium

We identified two clusters of cells (1,093 cells from Parenchyma 1 and 605 cells from

Parenchyma 2) that were transcriptionally similar to the parenchymal cells found in later

developmental stages; e.g. they express hypothetical protein (Smp_063330), *cathepsin B-like*

cysteine proteinase (Smp_141610) and *serpin* (Smp_090080) (Diaz *et al.,* 2020; Wendt *et al.*, 2020; Diaz & Attenborough *et al*., 2024). All top marker genes for Parenchyma 1 were also expressed in Parenchyma 2 but at lower levels, whereas the DUF2955 domain-containing protein (Smp_191970) was uniquely expressed in Parenchyma 2 (Figure 4A). A gene encoding a Zinc transporter Slc39a7 (Smp_318890) was a top marker for both clusters, with expression confirmed *in situ* within 7–9 cells; two of which were located anteriorly on either side of the brain and 5-7 located posterior to the brain (Figures 4B & F). These parenchyma cells have long cytoplasmic protrusions that reach between all the other cells, filling the intercellular space (Figure 4Fii-iii; Supplementary Videos 13 & 14). Pan (1980) called these the interstitial cells and, observing abundant glycogen particles and lipid droplets, proposed they function as sources of energy for the non-feeding larva and as nurse cells for the stem cells during the first few days post-infection of the snail. We performed ISH for two Parenchyma 2 markers, Smp_191970 and Smp_320330, but the first showed no detectable signal, and the second had broad expression across all cells with no specific overlap with the pan-parenchymal marker Smp_318890. Therefore, by ISH, we could only identify parenchymal cells in general and could not identify the two clusters *in situ*. STRING analysis for the Parenchyma 1 showed enrichment in networks for 'carbohydrate metabolic process' and 'hydrolase', and for Parenchyma 2: 'phagosome', 'mTOR signalling pathway', 'mitochondrial inner membrane' and 'proton-transporting two sector atpase complex' (Supplementary Figure 9). Analysis of GO annotations was able to distinguish between the two parenchyma sub-populations, with Parenchyma 1 genes enriched for 'extracellular region' and 'endopeptidase inhibitor activity', and Parenchyma 2 genes enriched for processes that are 'dynein complex' and 'microtubule-based process' (Supplementary Table 7; Supplementary Figure 10).

Two major stem cell populations in the miracidium that clustered by sex

We identified stem cells based on the expression of known stem cell markers for the mother sporocyst stage, e.g. *ago2-1* (Smp_179320), *nanos-2* (Smp_051920), *fgfrA* (Smp_175590), *fgfrB* (Smp_157300), *klf* (Smp_172480) and *hesl* (Smp_024860)(Wang *et al*., 2013; 2018)(Figure 5A). The mother sporocyst has been reported to have three stem cell populations: Kappa (*klf*[+], *nanos-2*[+]), Delta (*fgfrA*[+], *fgfrB*[+] and *nanos-2*[+]) and Phi (*fgfrA*[+], *fgfrB*[+] and *hesl*[+]) based on scRNAseq of 35 stem cells (Wang *et al*., 2018). Wang *et al*. (2018) suggest that the Kappa population serve as 'embryonic' stem cells that give rise to both the Delta and Phi populations; Delta cells generate somatic tissues and Phi produce temporary

larval tissues, including tegument. In the miracidium, we identified seven stem cell clusters

by expression of *ago2-1* (10,686 cells): Stem A and B are *klf*[+] and *nanos-2*[+] (i.e. Kappa-like);

Stem C and D are *nanos-2*[+], *fgfrA*[+], *fgfrB*[+], and *hesl*[+] (i.e. they resembled both Delta and

Phi)(Figure 5A, Supplementary Figure 11); as well as Stem E, F and G (Figure 5B).

Working with the latest version of *S. mansoni* genome (v10), where the sex chromosomes Z

and W have been resolved (Buddenborg *et al*., 2021), we were able to determine that the

Delta/Phi-like and Kappa-like populations are each composed of two well-separated clusters

that can be distinguished based on the expression of sex-linked genes (Figure 5A;

Supplementary Figure 12). *Schistosoma mansoni* has 7 pairs of autosomes and one pair of sex

chromosomes; males are homogametic (ZZ) and females heterogametic (ZW) (Buddenborg

et al., 2021). The sex chromosomes are composed of sex-specific regions that are unique to

each chromosome flanked by pseudoautosomal regions that are common to both Z and W. To

unveil the genes contributing to the sex-driven clustering we removed in turn different sex

chromosome regions from the analysis. When all Z-specific region (ZSR) and W-specific

region (WSR) genes were removed from the dataset and the cells reclustered, the two

Delta/Phi-like and two Kappa-like clusters each collapsed into single stem cell clusters,

giving five stem cell clusters in total (i.e. Delta/Phi, Kappa, E, F and G; Figure 5B,

Supplementary Figure 13). Removing from the dataset genes located solely on the WSR ($n \approx$

35), the ZSR ($n \approx 900$), or just the ~35 pairs of genes with homologous copies on the WSR

and ZSR did not impact the overall UMAP topology, i.e. in all three scenarios the stem cell

clusters remained separated by sex (Supplementary Figure 12). ZSR or WSR expression is

therefore sufficient on their own to split each of these stem clusters in two (Supplementary

Figure 12).

The unexpected subclustering of Delta/Phi and Kappa stem cells by sex could be either due to

the contribution to each cell's transcriptome by WSR genes (in females) that have diverged

from their ZSR homologues, or by differences in expression of ZSR genes between male or

female cells (see GO analysis of ZSR and WSR genes, Supplementary Table 13). Given the

small number of WSR genes, differences in ZSR genes — caused by a difference in copy-

number — seems likely to play a larger part. In many animals, dosage compensation

equalises gene expression across the sex chromosomes that are unevenly distributed between

the sexes. To investigate whether there is an incomplete or a lack of dosage compensation in

the ZSR genes in the *S. mansoni* miracidium, we classified all cells based on the expression

of WSR genes (i.e. only found in female individuals), so each cell was designated as W^+

(likely female) or W^- (likely male) (Supplementary Figure 12 A&B, Supplementary Figure

14A&B). The difference in ZSR and WSR gene expression was most pronounced in the

Delta/Phi-like and Kappa-like cells, shown by the separation of W^+ and W^- cells

(Supplementary Figure 14B). However, this analysis revealed other tissue-specific sex

differences in ZSR gene expression e.g. in the protonephridia, which was evident through

segregation of W^+ and W^- cells within that cluster. We found that W^+ and W^- cells were well

mixed in the majority of the remaining clusters (Supplementary Figure 14B). Male cells have

two copies of the ZSR (ZZ) and produce higher coverage across genes in this region

compared to female cells, which have one ZSR and one WSR (ZW). These findings suggest

that sex-linked gene dosage is not fully compensated in the miracidium, and there may be

tissue-specific variation.

To identify the stem cells *in situ*, we investigated the expression of the pan-stem marker

ago2-1. *Ago2-1*$^+$ cells were distributed lateral and posterior to the brain (Figure 5Ci) and

there was variation in the number between larvae, ranging from 20-29 (segmentation of

Ago2-1$^+$ nuclei in 5 larvae, median = 23, mean = 23.4)(i.e. ~7% of cells that make up the

larva are stem cells). Our count is slightly higher than previous estimates of 10-20

stem/germinal cells (Pan, 1980; Wang *et al*., 2013). Cells in the Delta/Phi-like cluster

expressed a canonical *p53 tumour suppressor family gene* (*p53-1*, Smp_139530) as a top

marker; *in situ* expression showed 12–14 *p53-1*$^+$ cells (segmentation of 3 larvae)(Figure 5D).

Cells in the Kappa-like cluster expressed *Uridine phosphorylase A* (*UPPA*, Smp_308145) as

a top marker; *in situ* expression showed 8–9 *UPPA*$^+$ cells, and they were located amongst the

p53-1$^+$ cells (segmentation of 3 larvae)(Figure 5D). We were not able to detect cells from the

Stem E, F and G clusters in the miracidium using ISH as there were no predicted marker

genes that were unique to these clusters. Although *Phosphomevalonate kinase* (*pmk*)

(Smp_072460) was a predicted marker for the Stem E cluster (Figure 5A), when multiplexed

as *p53/UPPA/pmk*, no *pmk*$^+$-only cells could be found (Supplementary Video 16). Likewise,

multiplexing *ago2-1/ p53/ UPPA* revealed no *ago2-1*$^+$-only cells (Supplementary Video 17).

Overall, we could spatially validate two stem cell populations within the miracidia: Delta/Phi

and Kappa. Cells in Stem E, F and G *in silico* clusters might be stressed/damaged/dying cells

or cells in transcriptionally transitional states (e.g. some *complexin* expression was also found

in a sub-population of Stem F).

A lack of EdU incorporation into miracidia cells (Supplementary Figure 15) indicated that DNA replication does not take place in the stem or somatic cells during this free-swimming stage. As the first stem cells start proliferating in the mother sporocyst ~20 hours post-infection (Wang *et al.,* 2013), this suggests that both stem cell populations pause cell division for at least 26 hours.

Many top markers for the Delta/Phi and Kappa stem clusters were ribosomal proteins (Figure 5A). STRING analysis showed that most of the top 100 markers of each cluster both formed large single networks enriched for genes involved in translation, ribosome biogenesis and gene expression, and this was particularly striking in the Delta/Phi cluster (Supplementary Figure 16). Amongst the top genes differentially expressed between Kappa and Delta/Phi (Supplementary Figure 17A; Supplementary Table 14) were six transcription factors up-regulated in Delta/Phi: an ortholog of *lin-39* (Smp_247590), *tiptop* (Smp_180690), *hes-related* (Smp_024860), *p53-1* (Smp_139530), *Zfp-1* (Smp_145470) and a putative *homothorax homeobox protein/six-3-1* (Smp_147790). The simultaneous co-expression of multiple transcription factors, indicates a multipotent progenitor stage (Kumar *et al*., 2019). Enriched in, or specific to, Kappa cells were genes involved in glycolysis and lipid metabolism, i.e. malate dehydrogenase (Smp_129610), 14 kDa fatty acid-binding protein (Smp_095360) and a very long chain fatty acid elongase (Smp_051810). GO analysis of differentially expressed genes (Supplementary Table 15) revealed that the most significantly upregulated genes in Delta/Phi were related to the structural component of the ribosome and translation; these are the first markers of a pre-activation stage in stem cells (Shin *et al*., 2015). Whereas GO analysis showed that up-regulated genes in Kappa are involved in transcription and metabolism (Supplementary Figure 17B), which indicated quiescent rather than primed and activated stem cells (Llorens-Babadilla *et al*., 2015).

One miracidia stem population contributes to somatic tissues and the other to the germline.

The Delta/Phi stem population simultaneously co-expressed three transcription factors that are known to be specific to the flatworm tegument/epidermal lineage, i.e *p53-1, zfp-1, six3-1* (Cheng *et al*., 2018; Collins *et al*., 2016; Wendt *et al*., 2018; 2022). This suggests that the Delta/Phi stem population may have been the origin of the miracidia tegument cells and may be the source of new tegument cells in the mother sporocyst stage. To explore the fates of the

two miracidia stem populations, we investigated the dynamics of their gene expression in

stem and tegument cells across the transition from miracidia to sporocyst. To do this, we

combined single-cell data from these developmental stages and cell types (this study and

Diaz & Attenborough *et al*., 2024) and used RNA velocity analysis (Bergen *et al*., 2020) to

estimate relative induction/repression and timing of gene expression in this dataset.

A UMAP plot of the combined clustering shows the gene expression dynamics of the

miracidia and sporocyst tegument and stem cells (Figure 6) (batch-corrected samples showed

similar results, Supplementary Figure 18). The velocity field and latent time representations

indicate the progression of some miracidia Delta/Phi stem cells towards the miracidia

tegument cells and a high speed of differentiation was inferred from the miracidia tegument

cells to the sporocyst tegument (Figure 6B & C). This suggests that the miracidia tegument

cluster may contain cells that range from post-mitotic stem cell progeny to tegument

progenitors to terminally differentiated tegumental cells.

Investigations into tegument development in adult schistosomes has revealed key regulators

and genes associated with different development states (Collins *et al*., 2016; Wendt *et al*.,

2018; Wendt *et al*., 2022). We examined the dynamics of these genes in the combined

miracidia/mother sporocyst dataset using RNA velocity and found that the velocities of these

genes show different temporal dynamics. *P53-1* (Smp_139530) was induced in the Delta/Phi

stem cells but repressed in some miracidia tegument cells and expression decreased in a

continuum across the miracidia tegument cells (Figure 6D). The flatworm-specific

transcription factor *Zfp-1* (Smp_145470) showed a similar expression pattern to *p53-1*,

except that the predominance of spliced copies (Figure 6E) suggested that transcription

happened earlier, during embryogenesis. A second zinc finger protein, *Zfp-1-1*

(Smp_049580) was expressed in tegumental cells more than stem cells, which suggested a

role in stem cell progeny and tegument progenitors (Supplementary Figure 19). In adults,

tegument cells are renewed continuously by a population of *tsp2+* (Smp_335630) progenitor

cells that fuse with the tegument (Wendt *et al*., 2018). In the miracidia, *tsp2* was expressed in

the Delta/Phi stem cells as well as tegument cells (Supplementary Figure 19), suggesting

earlier expression than in the adult, or that Delta/Phi are progenitor cells. Several markers of

mature tegument cells in the adult were expressed in the miracidia and sporocyst tegument

cells but not in the stem cells, e.g. *annexin* (Smp_077720), *calpain* (Smp_241190) and

endophilin B (Smp_003230) (Supplementary Figure 19). *Annexin* is specific to the tegument

cells, and the phase plot showed the accumulation of mature transcripts in the sporocyst
tegument, while there was evidence of recent transcription in the miracidia tegument
(Supplementary Figure 19). A putative ortholog of a planaria transcription factor *six-3-1*
(Smp_147790), a marker for early epidermal progenitors (Cheng *et al.*, 2018) was expressed
in the Delta/Phi stem cells and miracidia tegument (Supplementary Figure 19). All of these
genes showed little expression in the Kappa stem population (Supplementary Figure 19). The
expression of these known tegument/epidermal development and lineage-specific genes in
the Delta/Phi stem cluster, and the indication that *zfp-1* transcription occurred pre-hatching,
suggested that this stem population was the origin of the miracidia tegument cells and is the
source of the tegumental lineage. This, together with the observation that more than 80% of
protein-coding genes in the genome are detected in these miracidia data, suggests that similar
molecular programs for the development of tissues in the miracidia and mother sporocyst
may be redeployed multiple times across the life cycle.

As well as allowing us to interrogate the dynamics of known marker genes (Supplementary
Figure 19), the RNA velocity analysis also revealed genes previously unknown to have
dynamic behaviours across these stages; these included genes up-regulated from miracidia
tegument to sporocyst tegument, e.g. *epidermal growth factor receptor* (Smp_035260), genes
down-regulated from miracidia tegument to sporocyst tegument, e.g. *galectin domain*
(Smp_140590), and dynamic behaviour across stem cells and tegument, e.g. *unc44/ankyrin*
2,3 (Smp_145700)(Supplementary Figure 19, Supplementary Table 16).

In the Kappa stem population, there was transcription and expression of genes detected in
germline stem cells (GSCs) and germ cells in the intra-mammalian stages of the life cycle
(Wang *et al.*, 2018; Li *et al.*, 2021). A second P53 homolog, *p53-2* (Smp_136160), was
highly expressed and actively transcribed in Kappa cells (Figure 6F). In adults, this gene is
enriched in reproductive organs and has a genotoxic stress response that is thought to defend
the integrity of the genome by eliminating germ cells that have acquired mutations (Wendt *et*
al., 2022). Also expressed solely in Kappa cells (though some at low levels) are the germline-
specific regulatory genes *eled* (Smp_041540)(Figure 6G), *onecut1* (Smp_196950), *irx*
(Smp_063520), *akkn* (Smp_131630), and *boule* (Smp_144860)(expressed in the female
Kappa cells and Stem E, Supplementary Figure 22)(Wang *et al.*, 2018; Li *et al.*, 2021), as

well as *klf* (Smp_172480) (Supplementary Figure 20), a homolog of KLF4, which is an
important transcriptional regulator governing pluripotency (Takahashi & Yamanaka, 2006).
604
605 Together, these data suggest that the Delta/Phi population is the origin of the tegumental
606 lineage and likely contributes to extra-embryonic somatic tissues during intra-molluscan
607 development, while the Kappa population likely contains pluripotent cells that will contribute
608 to the development of cercariae and the germline in intra-mammalian stages (Figure 7).
609
610 **Discussion**
611
Schistosomes are complex metazoan parasites, with adult worms of *Schistosoma mansoni*
characterised into 68 transcriptionally distinct cell clusters (Wendt *et al*., 2020).
Understanding the genes involved in the development and maintenance of these cell types is
a promising approach for identifying novel therapeutic targets (Wendt *et al*., 2020). Here we
have shown that the miracidium larva is composed of ~365 cells. Compared to later life cycle
stages, where the cercariae is made up of ~1,600 cells, the schistosomulum ~900 (Diaz *et al*.,
2020) and the adults tens of thousands of cells (unpublished data), the miracidium is a simple
stage of the life cycle. We identified and spatially resolved 19 transcriptionally distinct cell
types, revealing the relative contribution of each tissue type to the larva (Figure 8). The
miracidium, therefore, with its diversity of differentiated cells present in relatively low
numbers, makes a simple and tractable system with which to gain a fundamental
understanding of the biology and spatial architecture of schistosome cells, and their
transcriptomes.
625
Nineteen cell types may be a conservative characterisation, but at this resolution most cell
types formed clearly defined clusters. There was sufficient sensitivity to detect cell types with
as few as six cells per larva, for example, Neurons 2 and 3, and the protonephridial system.
Some rarer cell types are embedded in existing clusters, particularly in Neuron 1, and a
complete neural cell atlas and a study of the putative photoreceptor cells is in progress
(Rawlinson *et al*., in prep). Some cell types were preferentially recovered, e.g. the stem cells;
ISH revealed that they comprise only 7% of the total cell number in the larva, yet they
accounted for ~52% of the scRNA-seq data (Supplementary Table 17). Preferential recovery
may be due to differences in the robustness of different cell types to dissociation, cell sorting,
and/or the microfluidics of the 10X GEM capture process. The ciliary plates, for instance,

were only found in sufficient numbers to be defined as a cluster in unsorted samples.
Therefore, there are clear benefits to including unsorted cell samples to discover the diversity
of cell types in a tissue, organ, or organism, especially when cell types may be fragile, sticky,
or unusually shaped. Even with the inclusion of unsorted cells, some known cell types were
not captured, i.e. the penetration glands. With three gland cells per larva and a theoretical 56x
coverage of each cell, we would have expected ~168 gland cells in the dataset. These cells
were easily identified after dissociation by their distinct flask-shape but were not identifiable
in these 10X data. Perhaps, because of their large size (20x50μm) and irregular shape, they
were not incorporated into the 10X GEMs. Therefore, we used plate-based scRNA-seq for
the penetration glands to complete the cell type transcriptome atlas for all known cell types.
Together, we have transcriptomically characterised the anatomical systems described by Pan
(1980), and we discuss key findings below.
648
The two muscle clusters in our dataset correspond to the circular and longitudinal muscles
that comprise the orthogonal body wall muscles of the miracidium. We identified differences
between circular and longitudinal muscle transcriptomes that have also been observed in
planaria; e.g. *MyoD* and *Wnt-11-1* are expressed in longitudinal and circular muscles,
respectively (Scimone *et al*., 2017). Posterior Wnt signalling and anterior Wnt inhibition are
commonly observed features of anterior-posterior body patterning in animals (Peterson &
Reddien, 2009). The expression of *Wnt-11-1* and a Wnt inhibitor, *Notum,* in the circular
muscles at opposite poles of the miracidium suggested these muscles are involved in defining
the anterior-posterior axis. Furthermore, the unilateral, asymmetrical expression of *Notum* at
the anterior end is suggestive of a dorso-ventral axis and/or a left-right axis, neither of which
have been identified in the miracidium before. Many axial patterning genes expressed in
planaria muscles are essential for patterning during regeneration (Witchley *et al*., 2013; Vogg
et al., 2014). The expression of orthologs in *S. mansoni* muscles, a nonregenerative animal,
suggests that these genes may regulate stem cells during homeostasis in the adult (Wendt *et*
al., 2020). As the body axes of the miracidium were defined during embryogenesis, the
expression of these genes in the muscles of the larva suggest they will provide positional
information to the stem cells of the mother sporocyst in the axial patterning of the daughter
sporocysts.
667
The nervous system accounts for over 50% of cells that make up the larva and is the most
transcriptionally heterogeneous tissue, as observed in other animals (Siebert *et al*., 2019;

Vergara *et al*., 2021). At the end of embryogenesis, at hatching, most of the miracidia

neurons are fully differentiated and form the functional neural circuits of the larva. This is

evident in our scRNA-seq data by the presence of discrete, well-defined neural cell clusters.

In situ validation of marker genes revealed many of the nerves identified ultrastructurally; for

example, the location and shape of Neuron 4 cells resemble that of the bipolar ganglion cells

with multi-ciliated nerve endings described by Pan, 1980. Examination of their

transcriptional profiles indicates their function; top markers include a putative *mTORC1*, two

Polycystin genes*, a *GTP-binding ras-like* and *fibrocystin.* Mammalian orthologs of these

function in cells with primary cilia in the detection of flow sheer and regulate mTORC1

signalling to maintain the cell in a quiescent and functional state (Lai & Jiang, 2020).

Polycystins 1 and 2 expressed in ciliated neurons of the larva of the annelid *Platynereis*

dumerilli are involved in mechanosensation, detection of water vibrations and mediate the

startle response (Bezares-Calderón *et al.,* 2018). Furthermore, in planaria, SoxB1-2 regulates

polycystin genes and together they function in ciliated cells that are involved in water flow

sensation (Ross *et al*., 2018). The gene expression profile of Neuron 4 cells in the miracidium

suggests they could function as flow and/or vibration sensors and may be specified by a Sox

transcription factor. These multiciliated receptors degenerate 12 hours after the miracidia

transform into a mother sporocyst (Poteaux *et al*., 2023).

In situ validation of marker genes also revealed an undescribed cell type, namely the six

submuscular cells that express the top marker genes for the ciliary plates (i.e. a 16KDa

calcium binding protein and *p25 alpha*). The expression of these genes in both the ciliary

plates and the submuscular cells suggests that these latter cells are transcriptionally similar

and might play a role in the development and functioning of the ciliary plates. Their

projection into the external environment suggests a sensory role, which, together with their

location between the first and second tier of ciliary plates, could suggest a role in

coordinating the movement of the motile cilia across the plates and tiers in response to

sensory cues (i.e. similar in function to the ciliomotor pacemaker neurons in *Platynereis*

larvae, Verasztó *et al*., 2017). The ciliary plates are a miracidium-specific cell type (it is the

only life cycle stage that uses motile cilia for locomotion), and once inside the snail, these

plates are shed within 2 – 48 hours (Meulement *et al*., 1978; Poteaux *et al*., 2023). As

expected, they are transcriptomically typified by enrichment with dynein and high metabolic

activity. The 16KDa calcium binding protein (Smp_096390) is clearly important to the

function of ciliary plates as it is highly expressed in data collected from both droplet- and

plate-based scRNA-seq platforms. Accordingly, bulk RNAseq data shows this gene was

previously detected in the late embryonic stage, miracidia and early mother sporocyst stage

(Lu *et al.*, 2021; Buddenborg *et al.*, 2023).

Between all the ciliary plates sit the ridges of the tegumental cells. The ridges connect to the

cell bodies beneath the body wall musculature, and their transcriptomes show a predicted

network of genes involved in protein folding, export and translocation. This is consistent with

ultrastructural studies that have observed abundant RER, ribosomes, Golgi apparatus, and

complex carbohydrates in these cell bodies, suggesting they are active in producing

exportable proteins (Meulement *et al.*, 1978; Pan, 1980). Large numbers of membrane-bound

vesicles were also observed and thought to become part of the new tegument membrane

(syncytium) that rapidly forms around the mother sporocyst during the early stages of

transformation, protecting the mother sporocyst from attack by the snails' amoebocytes (Pan,

1980). The RNA velocity analysis has revealed dynamic genes that may be crucial in the

maturation of the tegument cells across this life cycle transition, and in turn, for evasion of

host defences and parasite survival. This analysis also suggests a developmental trajectory for

the tegument cells from the miracidia Delta/Phi stem cells and follows their progeny to

terminally differentiated tegument cells in the miracidium and five-day-old sporocyst. Many

of the molecular regulators and markers of adult tegument biogenesis (Collins *et al.*, 2016;

Wendt *et al.*, 2018; Wendt *et al.*, 2022) are expressed in the miracidia and sporocyst stem and

tegument cells, indicating that there is a conserved molecular program for the development of

the syncytial tegument in the mother sporocyst and its maintenance in the adult worm.

The transcriptomes of the two stem populations in the miracidium suggest they are in

different activation states. This finding aligns with the discovery of two stem cell populations

in the mother sporocyst that have different proliferation kinetics (Wang *et al.*, 2013). The pre-

activated state of Delta/Phi stem cells in the miracidia may signify that they are the first

population to proliferate once the miracidium transforms into the mother sporocyst inside the

snail. These two stem populations may also have different potency states; the co-expression

of multiple transcription factors in Delta/Phi, including those specific to the tegument lineage

(Wendt *et al.*, 2018; 2022), indicates a multipotent progenitor stage. Whereas in the Kappa

stem cells, the expression of genes that define the germline in the intra-mammalian stages

(Wang *et al.*, 2018) suggest pluripotency. Supporting this, the RNA velocity analysis shows

that *p53-2*, the genome protection gene (Wendt *et al.*, 2022), is most highly expressed, and

being actively transcribed, in the Kappa population. The incorporation of miracidia single-cell data into existing stem cell scenarios on *Schistosoma mansoni* development (Wang *et al.*, 2013; Wang *et al.*, 2018; Li *et al.*, 2021) shows the continuum of the Kappa population from hatching through the intra-molluscan stages to the adult stage inside the mammalian host (Figure 7). The finding that *eled*, and many of the germline-specific regulatory program components, are expressed in some miracidia Kappa cells suggests that when a miracidium hatches, the cells that will contribute to the germline may already be segregated and the molecular regulatory program that differentiates somatic (Delta/Phi) and germ cell (Kappa) lineages is present. Intriguingly, the presence of spliced-only copies of the germline defining genes *eled* and *boule* could suggest that they are maternal transcripts that have been restricted to the primordial germ cells during embryogenesis, as is the case in Zebrafish embryos (Fishman et al., 2023). An alternative explanation is that unspliced transcripts exist for these lowly expressed genes but their abundance was below our threshold for detection.

An unexpected finding regarding the stem cells was the further clustering of the two populations by sex. This is the first identification of any sex differences in the miracidium. This degree of transcriptomic sexual difference was most evident in the Delta/Phi and Kappa stem cell clusters, with no distinct clustering by sex observed in other cell clusters. This could be explained by incomplete gene dosage compensation in the miracidium. Following investigations of differences in gene dosage compensation across developmental stages using bulk RNA-seq (Picard *et al.*, 2019), our study suggests tissue-specific variation could add a further dimension to the role of dosage compensation in schistosomes. A similar phenomenon has been described in mouse embryonic stem cells and in *Drosophila,* where a precise regulation of a gene-by-gene dosage compensation mechanism is critical to the proper development of specific tissues and organs, such as the wing and eye (Valsecchi *et al.*, 2018). In contrast to the miracidium, sex differences in undifferentiated stem cells have not been reported in later stages of the schistosome life cycle (Wang *et al.,* 2018; Diaz *et al.*, 2020; Wendt *et al.*, 2020). In fact, it may not have been possible to detect these differences in earlier studies because scRNA-seq data were mapped onto older versions of the genome, where the sex chromosomes were not as well defined (Buddenborg *et al.*, 2021). In addition to using an improved reference annotation, the present study has also benefited from a high level of coverage, enabling differences in stem cells to be more easily discerned. Whether these sex-linked gene expression and dosage differences have functional implications in developing male and female sporocysts remains to be investigated.

The current understanding of schistosome development is that differentiated somatic cells of

the miracidium do not continue into the daughter sporocyst stage, the cercariae and beyond;

it is the stem cells that persist and differentiate into the cell types of the subsequent life stages

(Wang *et al*., 2018). In this respect, all miracidium somatic cells are miracidium-only cells.

Our atlas shows that their transcriptional signatures are similar to comparable tissue types at

later stages (Diaz *et al*., 2020; Wendt *et al*., 2020, Diaz & Attenborough *et al*., 2024) and that

there may be conserved tissue-specific molecular developmental programs that are deployed

multiple times during the complex life cycle. Through this atlas, we have shown that even in

this simple and sexually monomorphic short-lived stage, there are different populations of

muscle, a diverse and distinct range of neural cell types, and that stem cell transcriptomes are

already different based on the sex of the miracidium. The molecular definitions and spatial

analyses of the miracidia cell types we have presented here allow us to explore the molecular

processes and pathways to reveal cell functions and interactions. In addition, high coverage

data from individual cells has allowed us to examine how gene expression changes across

and within tissue types. Already, a picture is emerging of how the peripheral neurons may

detect environmental cues, connect to the brain, muscles and ciliary plates to orientate in the

water column; how the parenchymal cells supply nutrients to the surrounding cells; what the

contents of the penetration glands may be; and how the tegumental and stem cells are primed

for the transition to the intra-molluscan stage.

Methods

Parasite material

Schistosoma mansoni (NMRI strain) eggs were isolated from livers collected from

experimentally infected mice perfused 40 days post-infection. All animal-regulated

procedures, including the experimental mouse infections at the Wellcome Sanger Institute,

were conducted under Home Office Project Licence No. P77E8A062 held by GR. All the

protocols were presented and approved by the Animal Welfare and Ethical Review Body

(AWERB) of the Wellcome Sanger Institute. The AWERB is constituted as required by the

UK Animals (Scientific Procedures) Act 1986 Amendment Regulations 2012.

Nuclei count

Schistosoma mansoni eggs isolated from infected mouse livers were hatched under light in

freshwater (Mann *et al.,* 2010), and the miracidia collected within four hours of hatching. The

sample was put on ice for an hour to sink and concentrate the larvae, and all but 1 mL of the

water was removed. One mL of 8% paraformaldehyde (4% final PFA concentration) was

added to fix the miracidia for 30 minutes at room temperature. The larvae were rinsed in

PBS/Tween (0.1%) and mounted in DAPI Fluoromount for confocal imaging. The whole

mounts were imaged on a Leica Sp8 using a 40X objective with the pinhole set to 1AU, and a

30-50% overlap between Z slices. The Z-stacks were imported into ImageJ, and each nucleus

was manually segmented and counted using the plugin TrakEM2 (Cardona *et al.*, 2012).

Single-cell tissue dissociation

Schistosoma mansoni miracidia for single-cell dissociation were collected as described

above. Eggs isolated from mouse livers were hatched in nuclease-free water (NFW) and

collected within four hours of hatching. Two samples of miracidia were collected; sample 1

contained approximately ~8250 individuals, and sample 2 ~11,700. The samples (live

miracidia in NFW in Low Protein Binding 15 mL Falcon Tubes [cat# 30122216]) were

placed on ice for an hour to sink and concentrate the larvae, and all but 20 µl of water was

then removed.

Three hundred µl of digestion buffer (2.5 µl of 0.5 M EDTA in 300 µl of Hank's Buffered

Salt Solution without Ca2+ and Mg2+ [HBSS-/-]) was added to the larvae. We reconstituted

250 mg of the enzyme Native *Bacillus licheniformis* Protease (NATE-0633, Creative

Enzymes)(Adam *et al.*, 2017) in 2.5 mL of HBSS-/- (to give a stock concentration of 100

mg/ml) and added 200 µl of this stock to the larvae in buffer (final enzyme concentration =

40 mg/ml). The samples were kept on ice for 45 minutes and pipetted gently ~20 times every

five minutes. Two hundred µl of 25% BSA in 1X HBSS-/- was added to the sample to

quench the enzyme, and 10 ml of ice-cold PBS +/+ was added to bring the BSA

to ~0.5%. The samples were centrifuged at 350 g for 5 minutes at 4℃, and all but 20 µl of

the supernatant was removed. We reconstituted 5 mg of the enzyme Liberase™ (Thermolysin

Medium) Research Grade (LIBTM-RO Roche) in 2 mL of PBS (to give a stock concentration

of 2.5 mg/ml) and added 100 µl of this stock to the sample (final enzyme concentration =

2.08 mg/ml). The samples were incubated at room temperature for 15 minutes,

pipetting gently ~10 times every five minutes. The suspension was passed through 60 µm and

30 µm cell strainers (Falcon) and resuspended in 5 mL of cold 1× PBS supplemented with

0.5% BSA. The samples were centrifuged at 350 g for 10 minutes at 4°C, and all but 700 µl

of the supernatant was removed.

Sample 1 cell suspension was sorted; 200 µl was used for a no-stain control for

autofluorescence during cell counting and sorting. The remaining 500 µl of cell suspension

was stained with 1 µg/ml of Propidium Iodide (PI; Sigma P4864) to label dead/dying cells.

Sample 1 was sorted into an eppendorf tube using the BD Influx™ cell sorter (Becton

Dickinson, NJ) by enriching for PI negative cells (i.e., live cells). It took ~3 hours from the

beginning of the enzymatic digestion to generating single-cell suspensions ready for library

preparation on the 10X Genomics Chromium platform. Sample 2 cell suspension was not

sorted.

10X Genomics library preparation and sequencing

The 10X Genomics protocol ("Chromium Next GEM Single Cell 3' Reagent Kits v3.1 User

Guide" available from

https://assets.ctfassets.net/an68im79xiti/1eX2FPdpeCgnCJtw4fj9Hx/7cb84edaa9eca04b607f

91193162994de/CG000204_ChromiumNextGEMSingleCell3_v3.1_Rev_D.pdf) was followed

to create gel in emulsion beads (GEMs) containing single cells, hydrogel beads and

reagents for reverse transcription, perform barcoded cDNA synthesis, and produce

sequencing libraries from pooled cDNAs. The concentration of single-cell suspensions

was approximately 119 cells/µl (sample 1) and 122 cells/µl (sample 2), as estimated by flow

cytometry-based counting (Beckman Coulter Cytoflex S). We concentrated both samples to

732 cells/µl. We loaded two aliquots of sample 1 and two of sample 2 into four 10X reactions

according to the 10X protocol (Chromium Single Cell 3' Reagent Kits v3.1), intending to

capture approximately 10,000 cells per reaction.

Library construction (following GEM breakage) using 10X reagents following the "Single

Cell 3'Reagent Kits v3.1 User Guide". The four libraries were sequenced on one lane of

Illumina Novaseq XP (28 bp read 1, 90 bp read 2, 10 bp index 1, 10 bp index 2). All raw

sequence data will be deposited in the ENA.

Mapping and quantification of single-cell RNA-seq

We mapped the single-cell RNA-seq data to version 10 of the *S. mansoni* reference genome

(Buddenborg *et al*., (2021) and WormBase ParaSite ((Howe *et al*., 2016; 2017)) using 10X

Genomics Cell Ranger (version 6.0.1) (Zheng *et al*., 2017). Before mapping, exon eight was

removed from Smp_323380 and the 3' end of was trimmed from Smp_337410 to reduce

erroneous mapping to the W chromosome from Z gametologs. The Cell Ranger

implementation of EmptyDrops (Lun *et al*., 2019) was used to detect cells and empty

droplets. Across the four samples, on average 63.88% of the reads mapped confidently to the

transcriptome (63.4% for sorted, 64.35% for unsorted), and an average 83.65% of reads were

associated with a cell. There were an average of 118,395 reads per cell. Across the four

samples, we captured an estimated 33,319 cells: 8,815 and 24,504 cells from sorted and

unsorted samples, respectively. The median gene count was 1,660, and the median UMI

count was 5,343. Following QC, 20,478 cells remained for onward analysis.

Quality control (QC) and clustering using Seurat

The four samples were imported into Seurat (version 4.3.0) (Hao *et al*., 2021) using R

(version 4.1.3) in RStudio (version 1.4.1106) for quality control, clustering, and marker gene

identification. The four samples were initially processed separately. Cells with less than 200

genes expressed and genes in less than three cells were removed. QC metrics, including gene

distribution and UMI counts per cell, were inspected visually and used to select QC cut-offs.

Following a preliminary analysis without further cell filtering, cells with fewer than 500

UMIs were removed, and a maximum mitochondrial threshold of 5% was applied. This was

based on data distribution, best practices from the authors of the software, and other similar

studies; here, we were aiming to remove low-quality cells while being relatively permissive

as there may be cell types in these data outside of standard filtering parameters. Based on

existing knowledge of this larva, we were expecting both multinucleate and anucleate cell

type(s) as well as single-nuclei cells (Pan, 1980). The data were normalised, scaled

(regressing for mitochondrial gene percentage), and variable genes were identified for

clustering following Seurat guidelines. Throughout the analysis of these data, several tools

were used to select the optimal number of principal components (PCs) to use: Seurat's

ElbowPlot and JackStraw, and molecular cross-validation

(https://github.com/constantAmateur/MCVR/blob/master/code.R), and Seurat's clustree tool

was used to assess the stability of clusters with increasing resolution. We employed

DoubletFinder (McGinnis *et al*., 2019) to identify multiplets, which can lead to spurious

signals. The estimated doublet rate was calculated based on 10X loading guidelines and

adjusted for our sample concentrations; we estimated multiplet rates at 8% for sorted samples

and 52.8% for unsorted samples. We used paramSweep, as recommended, to identify the

optimum parameters for each sample in our data and removed cells classified as multiplets.

DoubletFinder removed 299 and 303 cells from the two sorted samples, and 6,069 and 5,496

cells from the two unsorted samples.

We then combined the two sorted samples and the two unsorted samples (producing one

sorted object and one unsorted object). We used SCTransform to normalise and scale these

two objects (with regression for mitochondrial percentage and sample ID). Seurat's

IntegrateData function was then used to integrate the sorted and unsorted samples (with

normalization.method = "SCT"). This resulted in 20,478 cells passing all the QC filters and

going onto the primary analysis, 3,849 and 3,962 from the sorted samples and 6,702 and

5,965 from the unsorted samples.

Following integration, principal component analysis (PCA) was run for up to 100 PCs. The

data were explored using a range of PCs and resolutions and using the tools listed above, 55

PCs and a resolution of 1 were selected for the analysis. We manually segmented out the cells

in the Neuron 3 cluster based on clear differences in gene expression. Top markers were

identified using Seurat's FindAllMarkers (min.pct = 0.0, logfc.threshold = 0.0, test.use =

"roc") as recommended by Seurat best practices (https://satijalab.org/seurat/) to capture subtle

differences in gene expression, and were sorted by AUC. We used the identity and expression

of top marker genes (as identified by AUC), along with previously published cell and tissue

type marker gene expression, to determine the identity of the cell clusters. To link gene

description and chromosome locations to gene IDs, the GFF for the latest publicly available

genome version was downloaded from WBPS (WBPS18 release, Howe *et al*., 2016; 2017)).

Sex-effect analysis

To investigate sex effects on the single cell transcriptome, the cells were categorised as W+

or W- (an approximation of female and male cells respectively, as males lack WSR genes).

As ground-truth data was absent, the % of reads mapping to ZSR and WSR was calculated

using Seurat's PercentageFeatureSet. Cells with 0.2% or more reads mapping to the WSR

were classified as W+, and the remaining cells were classified as W-. The 0.2% threshold

was selected (Supplementary Figure 14) to avoid classifying ZZ cells with a few transcripts

erroneously mapping to a gene on the WSR as W+. This produced 8,702 cells labelled as

W+, and 11,776 cells labelled as W-.

Self-Assembling Manifold (SAM) algorithm

The count data from the QC-filtered Seurat analysis was exported for SAM (v1.0.1) analysis. For each SAM analysis, the raw count data was imported as a SAM object, standard preprocessing done (using default SAM parameters), and the SAM algorithm (Tarashansky *et al*., 2019) run. For further analysis, the object was converted into a Scanpy (v 1.8.2) (Wolf *et al*., (2018)) object, top markers extracted, and metadata from the Seurat analysis was attached. This method was repeated on the Neuron 1 cluster alone, to better understand the transcriptome heterogeneity.

RNA velocity

All four miracidia samples and the raw data from the mother sporocyst (Diaz & Attenborough *et al*., 2024) were pseudo-mapped using kallisto bustools (v 0.27.3) (Melsted *et al*., 2021) using the lamanno workflow to the version 10 *S. mansoni* genome (WBPS18 release (Howe *et al*., 2016; 2017, with the same amendments as listed above). The filtered spliced and unspliced matrices for each sample individually were imported to R. DropletUtils (v 1.20.0) (Lun *et al*., 2019) and BUSpaRse (v 1.14.1) (Moses and Pachter, 2023) were used to process and QC the matrices, and filter out cells and genes with few reads. The matrices were then converted to Seurat objects and further filtered so only cells with the cell barcodes that passed QC filtering in the primary Seurat analysis were retained. The metadata and raw counts of the unspliced and spliced reads following filtering were exported as CSV files. The same method was performed on the mother sporocyst samples, which were combined before exporting. For the joint analyses of miracidia and sporocyst cells, the combined sporocyst Seurat object was merged with each of the four miracidia samples separately (due to the mismatch in sample size and batch effect between sorted and unsorted samples) and all were exported.

The CSV files containing spliced and unspliced reads, together with sample metadata were combined to make an AnnData object using Scanpy (v 1.8.2, Python v 3.8.5) (Wolf *et al*., 2018). Using scvelo (v 0.2.4) (Bergen *et al*., 2020), the standard parameters were used to filter and normalise the data, compute moments, and recover dynamics. Velocity was calculated using the dynamical mode. Following PCA, a UMAP was generated (n_neighbors=10, n_pcs=40) and Leiden clustering was added (resolution=0.8). The top-ranked dynamical genes for each cluster and known genes of interest were plotted. This was

performed for each of the four miracidia samples combined with the sporocyst samples, with

and without the use of combat for batch correction. As the sporocyst samples were also

sorted, we used the sorted miracidia samples to reduce technical effects.

To focus on the stem/germinal and tegument clusters, the same method as above was used

after subsetting the data in Seurat to contain only cells from the clusters Stem/germinal,

Tegument 1, and Tegument 2 in the sporocyst samples, and only Stem A-G and Tegument

clusters from the miracidia clusters.

We present the stem and tegument cells from the sporocyst stage merged with sample 1

(sorted). The equivalent results from sporocysts merged with sample 4 of the miracidia and

results with all cells from those two conditions are available in the supplementary data

(Supplementary Figures 21-23).

EdU-staining experiment

Miracidia were collected as described above for 2 hours. As miracidia continually hatched

over this time period, they ranged from 0 to 2 hours post-hatching. The miracidia were

collected and split into two samples: one for EdU labelling and the second for a negative

control (miracidia in water without EdU). Sample one was pulsed with 500 µM EdU in water

for 4 hours. Sample two was kept in water for 4 hours. Both samples were fixed in 4% PFA

for 30 minutes at room temperature. EdU incorporation was detected by click reaction with

25 µM Alexa Fluor488 Azide conjugates (Invitrogen) for 30 minutes. As a positive control,

we pulsed freshly-transformed mother sporocysts in EdU for 3 days (Wang *et al*., 2013).

Three biological replicates were carried out, with 20 miracidia and 20 mother sporocysts per

treatment examined for each replicate.

Plate-based scRNAseq analysis of individually isolated cells

A sample of thousands of miracidia was dissociated into a single cell suspension using the

dissociation protocol described above. Individual gland cells, ciliary plates and unknown

cells were picked up one at time using an EZ Grip transfer pipette set to aspirate a 3µl

volume (Research Instruments Ltd 7-72-2800). Each cell was placed in a well of a 96-well

plate containing a lysis buffer (NEBNext Cell Lysis Buffer, E6428). cDNA was generated

using the NEB low input RNA kit (E6420). DNA library construction was carried out using

the NEBNext® Ultra™ II FS DNA Library Prep Kit for Illumina (E7805). The libraries were

sequenced on one lane of NovaSeq SP. The reads from the 96 samples were pseudoaligned to

the version 10 *S. mansoni* genome (WBPS18 release (Howe *et al*., 2016; 2017, with the same

amendments as listed above) using kallisto (v 0.44.0) (Bray *et al*., 2016)). These were

imported into R (v4.3.1) and the metadata were inspected, after which only samples coming

from a single cell with a minimum of 20% reads pseudoaligned were retained (n=76). The

samples were then processed using sleuth (v0.30.1) (Pimentel *et al*., 2017) following

guidance from sleuth tutorials, including the steps for normalising, fitting error models, and

differential analysis. Differential analysis was used to identify significant genes between the

cell types, which were visualised, along with known genes of interest, using sleuth's heatmap

function. Furthermore, the kallisto outputs were used to calculate the top 200 genes by

median transcripts per million (TPM) of the two key cell types of interest, ciliary plates and

gland cells, and these were used for GO term analysis following the method described below.

1018

***In situ* hybridisation**

Miracidia collected as described above within 4 hours of hatching were fixed in 4% PFA in 2

μm filtered PBSTw (1x PBS + 0.1% Tween) at room temperature for 30 minutes on a rocker.

The larvae were transferred to incubation baskets (Intavis, 35 μm mesh) in a 24-well plate,

rinsed 3 x 5 minutes in PBSTw, and incubated in 5 μg/ml Proteinase K (Invitrogen) in 1x

PBSTw for 5 minutes at room temperature. They were post-fixed in 4% Formaldehyde in

PBSTx for 10 minutes at room temperature and then rinsed in PBSTw for 10 minutes. From

this point on, the *in situ* hybridisation experiments followed the protocol described by Choi *et*

al., (2016; 2018) and developed for wholemount nematode larvae. We carried out fluorescent

in situ hybridisation on 27 cell-type marker genes. Probes, buffers, and hairpins for third-

generation *in situ* hybridization chain reaction (HCR) experiments were purchased from

Molecular Instruments (Los Angeles, California, USA). Molecular Instruments designed

probes against the sequences on WormBase ParaSite (WBPS version 18, WS271, Howe *et*

al., 2017)(Supplementary Table 18). We subsequently labelled some larvae with phalloidin

by adding 1 μl phalloidin to 1 ml 5X SSCT for an hour at room temperature in the dark, then

rinsing 6 x 30 minutes in SSCT, incubating overnight at 4°C in DAPI fluoromount-G

(Southern Biotech) and then mounting and imaging on a confocal laser microscope (Leica

Sp8). We carried out at more than three *in situ* hybridisation experiments for each marker

gene we validated (each experiment was a biological replicate). From each experiment we

imaged (by confocal microscopy) 10 miracidia (technical replicates) per marker gene

experiment. To count the number of cells belonging to each tissue type we imported confocal

Z-stacks of tissue marker gene expression (Supplementary Videos 18-20) into ImageJ, and

using the plugin TrakEM2 (Cardona *et al*., 2012), manually segmented each nuclei that was

expressing, and was surrounded, by the transcripts.

1043

1044   **Opsin phylogeny**

1045

Amino acid sequences for selected metazoan opsin proteins were obtained from the GenBank

database and aligned using the —auto mode of MAFFT v7.450 (Katoh & Standley, 2013),

then trimmed using trimAl v1.4.rev22 (Capella-Gutierrez *et al*., 2009) to remove alignment

columns with more than 50% gap characters ensuring at least 50% of alignment columns

were maintained in the final alignment. Phylogenetic analysis of this alignment was

performed using RaxML v8.2.12 (Stamatakis, 2014). The best-fitting empirical amino acid

substitution model was identified as maximising the log-likelihood of the data under a model

incorporating gamma-distributed variation in substitution rates across sites, and this optimal

model (LG) was used for subsequent inference. The maximum-likelihood tree and branch

lengths were identified as the highest likelihood from 25 independent searches starting from

different random starting tree topologies, and support for partitions on this tree was estimated

using 1000 bootstrap replicates of resampling alignment positions with replacement, using

the fast bootstrap (-x) algorithm implemented in RaxML.

1059

Protein-Protein interaction analysis

Protein interactions were predicted using the online search tool STRING (www.string-db.org;

V 11) (Szklarczyk *et al.*, 2019) for the following well-defined cell clusters: Neuron 3-5,

ciliary plates, protonephridia, Parenchyma 1 & 2, tegument and the two main stem cell

populations, Delta/Phi and Kappa. Protein sequences for the top 100 marker genes for each

cell cluster were collected from WormBase ParaSite. The protein sequences were entered as a

multiple protein search. Default settings were used to predict interactions with either a

minimum interaction (confidence) score of 0.4 or 0.7, corresponding to a medium or high

level of confidence.

1069

Gene Ontology enrichment analysis

Each cluster's list of marker genes was filtered to focus on specific cluster markers. Genes

were retained for analysis if they had a minimum AUC score of 0.7. To compare the

functional enrichment in Stem 1 v Stem 2, Stem A was combined with Stem B and Stem C

was combined with Stem D, and the DE markers were identified using Seurat's FindMarkers.

These markers were filtered (with adjusted p-values <0.001) and used as the basis for GO

enrichment analysis in the Delta/Phi and Kappa stem clusters. TopGO (version 2.46.0)

(Alexa & Rahnenfuhrer, 2023) was used in R with the weight01 method to identify enriched

GO terms. Node size was set to a minimum of 5 terms, and analysis was run for each cluster

on the BP and MF categories. Fisher's exact test was used to assess the statistical significance

of overrepresented terms, with a FDR threshold set < 0.05. The same method was used to

identify overrepresented terms in the WSR genes, ZSR genes, and WSR together with ZSR

genes for the BP, MF and CC categories.

1083

Data (and Software) Availability

1084

All raw sequence data is available at the ENA under study accession PRJEB45615. The

scripts used to perform the analyses presented here are available at

https://github.com/tessatten/singlecell-miracidia. For the purpose of Open Access, the authors

have applied a CC BY public copyright licence to any Author Accepted Manuscript version

arising from this submission. All supplementary material (figures, tables and videos) is

available on BioRxiv.

1091

Funding

1092

1093 The project was funded by the Wellcome Trust (grant 098051 and 206194). KR also received

1094 transition funds from the Marine Biological Laboratory in Woods Hole. SRD and GR are

1095 supported by UKRI Future Leaders Fellowships [MR/T020733/1] and [MR/W013568/1],

1096 respectively.

1097

Author Contribution

1098

KR, TA, KA, GR and MB designed the research

CD developed and optimised the dissociation protocol

KR and GR carried out the dissociation and cell collection

JG carried out cytometry

GS optimised the EdU protocol

TA analysed data

KR performed ISH experiments

SD annotated marker genes in v10 of the genome

TA and KR drafted the complete manuscript, with sections of text contributed by GR and

MB

JC carried out the opsin phylogeny

All authors read and approved the final manuscript.

1111

Acknowledgements

We thank Simon Clare and his team for technical assistance with the life cycle maintenance

and collection of parasite material at the Wellcome Sanger Institute, and Karl Hoffmann,

Josephine Fforde-Thomas and Benjamin Hulme for additional ongoing parasite material. We

thank Sarah Buddenborg for work on the annotation of the *Schistosoma mansoni* genome;

David Goulding for assistance with Microscopy; teams within DNA Pipelines for data

production; and core Informatics teams and Pathogen Informatics for the infrastructure used

for data analysis. We thank members of the Parasite Genomics team at the Wellcome Sanger

Institute for their comments and input on this study. Finally, we thank the reviewers and the

editor for their valuable feedback on this manuscript.

1122

1123

1124 **Figure legends**

1125

Figure 1. **Identification of 19 transcriptionally distinct cell types in the miracidium.**

A) The miracidium is composed of ~365 cells; **i)** DIC image of miracidium, **ii)** a Maximum

Intensity Projection (MIP) of a confocal z-stack of DAPI-stained miracidium with nuclei

segmented to enable counting (larval anterior pole at the top in all images). **B)** Experimental

scheme describing the parasite's life cycle (images of developmental stages not to scale),

parasite dissociation, single-cell analysis of miracidium and validation pipeline. An average

of 9975 miracidia per sample were dissociated; two samples were enriched for live cells

(propidium iodide negative) using fluorescence-activated cell sorting (FACS), another two

samples were unsorted. Cells were loaded according to the 10X Chromium single-cell 3'

protocol. Clustering was carried out to identify distinct populations and population-specific

markers. Validation of population-specific markers was performed by *in situ* hybridisation

(ISH). **C)** Uniform Manifold Approximation and Projection (UMAP) representation of

20,478 miracidium single cells. Cell clusters are coloured and distinctively labelled by cluster

identity. **D)** Gene-expression profiles of the top population markers identified for each cell

cluster (gene identifiers shown in parenthesis but with "Smp_" prefix removed for brevity).

The colours represent the expression level from yellow (high expression) to dark blue (low

expression). Gene expression has been log-normalised and scaled using Seurat(v. 4.3.0). The

sizes of the circles represent the percentages of cells in those clusters that expressed a specific

gene.

1145

Figure 2. Orthogonal body wall muscles are transcriptionally distinct. A) Dot plot

depicting the expression profiles of specific- or enriched- marker genes in the two muscle

clusters. Genes validated by *in situ* hybridisation (ISH) are marked in red. Gene identifiers

shown in parenthesis but with "Smp_" prefix removed for brevity. Gene expression has been

log-normalised and scaled using Seurat(v. 4.3.0). **B) i-ii)** Wholemount ISH (WISH) of

paramyosin *PRM*$^+$ and counterstaining with phalloidin reveals the nuclei of the circular body

wall muscles (BWM), which form two distinct bilaterally symmetrical lines that run

peripherally from the pole of the larva to the other (arrowheads in ii). The longitudinal BWM

nuclei (ii - arrows) are spaced regularly between the lines of the circular BWM nuclei and

their actin fibres run orthogonally to the circular muscles. These patterns were seen in 100%

of individuals examined, n = 50. **C) i)** Segmentation of the *PRM*$^+$ cells in one miracidium

shows 74 muscle cells in total: ~28 circular (segmented in orange), 33 longitudinal (yellow)

and **ii)** 13 in an anterior unilateral cluster (inside dashed cyan line) adjacent to the apical

gland cell (identifiable by its four nuclei, inside dashed red circle). **D)** Expression of markers

for muscle 1 cluster; **i)** *Kunitz-type protease inhibitor* is expressed in circular BWM and *wnt-*

11-1 is expressed in 7 *Kunitz-type protease inhibitor*$^+$ cells at the posterior pole; **ii)** close-up

of posterior end of another miracidium showing the co-expression of *Kunitz-type protease*

inhibitor and *wnt-11-1*. In 100% of individuals examined, n = 30. **E) i-iii)** *Notum,* an inhibitor

of *wnt* signalling, is expressed highly at the opposite pole to *wnt-11-1* in the cluster of 13

muscle nuclei adjacent to the apical gland cell (nuclei in dashed red circle). In 100% of

individuals examined, n = 30. **F)** Expression of markers for muscle 2 cluster; a *calcium*

activated potassium channel (Smp_156150) and the transcription factor *myoD* (Smp_167400)

show co-expression in longitudinal BWMs (arrowheads). In 100% of individuals examined, n

= 30. Scale shown in B also applies to C, Di, E and Fi.

1170

1171

Figure 3. Neural complexity and ciliary plates in a simple larva. A) Expression profiles of

cell marker genes that are specific or enriched in the five neuronal clusters and ciliary plates.

Genes validated by *in situ* hybridisation (ISH) are marked in red. Gene identifiers shown in

parenthesis but with "Smp_" prefix removed for brevity. Gene expression has been log-

normalised and scaled using Seurat(v. 4.3.0). *Complexin* (*cpx*) is expressed in all five nerve

clusters and is a neural marker in the adult (Wendt *et al*., 2020). **B) i)** A dorso-ventral view

of WISH of *cpx* shows expression in the brain, to the anterior - with projections around the

three gland cells, and in two posterior clusters (100% of individuals examined, n = 50). **ii)**

Segmentation of *cpx*⁺ cells (in magenta) in one miracidium (lateral view) reveals that at least

209 of the 365 cells that make up the larva are neural and that the clusters of *cpx* cells that are

located posterior to the brain, and send projections to the posterior pole, sit perpendicular to

the circular muscle cell nuclei (blue dashed lines). **C)** Multiplexed ISH for top markers for

neuron 2 (Smp_201600) and neuron 3 (Smp_071050) shows expression in multiple cells of

the brain, including cells adjacent to each other. Smp_071050 is also expressed in the apical

gland cell (red dashed circle). 100% of individuals examined, n = 30. **D)** The top marker for

neuron 4 Smp_319480 is expressed in four cells anteriorly (arrows) and ~22 that sit lateral

and just posterior to the brain; these later cells send projections into the brain and out to the

body wall at the latitude between the 2nd and 3rd tier of ciliary plates **(iii)** seen here expressing

the ciliary plate marker gene Smp_096390. 100% of individuals examined, n > 30. **E)** Neuron

5 marker Smp_343710*,* an EF-hand domain-containing protein, is expressed in **(i)** 10-20 cells

whose nuclei sit outside of, and anterior to, the brain, in and around the paired lateral glands

and their secretory ducts, and **(Eii & iii, F)** in a pair of bilaterally symmetrical bulbous

protrusions. 100% of individuals examined, n = 30. **F)** Summary schematic of the *in situ*

expression of marker genes for the neural cell clusters. **G) i)** A top marker for ciliary plates, a

calcium binding protein (Smp_096390), shows transcripts expressed in all the ciliary plates

of all four tiers (brackets) and in 6 cells towards the anterior pole. **(ii & iii)** Counterstaining

with phalloidin shows that the nuclei of these anterior cells sit beneath the body wall muscle

and send a projection externally between the 1st and 2nd tier of ciliary plates (arrowheads).

100% of individuals examined, n > 30. Scale shown in B also applies to C, D, Ei and Gi.

1201

1202

Figure 4. Identification of the tegument, protonephridia, parenchymal cells in the

miracidium. **A)** Expression profiles of cell marker genes specific or enriched in these cell

type clusters. Genes validated by *in situ* hybridisation are in red. Gene identifiers shown in

parenthesis but with "Smp_" prefix removed for brevity. Gene expression has been log-

normalised and scaled using Seurat(v. 4.3.0). **B** Segmentation of nuclei of the cells

expressing the marker genes for; **i)** protonephridia and tegument (multiplexed) show that 6

cells express Smp_335600 (*ShKT-domain protein*, a marker for protonephridia), 46 cells

express *Meg-6* (a marker for tegument) (segmentation of 1 larva), and **ii)** seven cells that

express an uncharacterised gene, Smp_318890 (a marker for parenchymal

cells)(segmentation of 1 larva). **C & D)** The tegument marker *Meg-6* shows expression

around nuclei in the posterior two-thirds of the larvae. The nuclei are below the body wall

muscle, and cytoplasmic protrusions reach between muscle filaments (arrows) and form the

epidermal ridges (arrowheads) between the ciliary plates (which are visible in **Dii & iii**

expressing the ciliary plate marker Smp_096390) (asterisks). 100% of individuals examined,

n > 30. **E)** The protonephridial marker Smp_335600 shows 'S'-shaped expression with

transcripts extending from the nucleus of the anterior flame cell (AFC) along the excretory

tubule and its nucleus (ETN) to the posterior flame cell (PFC), **Eii)** it is expressed around the

nuclei (arrow) rather than the barrels (arrowhead) of the flame cells. 100% of individuals

examined, n > 30. **Fi)** The pan-parenchymal marker, Smp_318890, was expressed in 2

anterior cells (one on either side of the brain) and 5-7 cells posterior to the brain. **Fii)** These

cells have long cytoplasmic protrusions that reach between all the other cells, **Fiii)** including

the *ago2-1*[+] stem cells. 100% of individuals examined, n > 30. Scale shown in B also applies

to C, D, Ei and F.

1226

Figure 5. Two defined populations of stem cells cluster by sex. A) Expression profiles of

cell marker genes that are specific or enriched in the stem cell clusters. Genes specific to the

W (female-specific) sex chromosome are highlighted in blue. Genes validated by ISH are

marked in red. Gene identifiers shown in parenthesis but with "Smp_" prefix removed for

brevity. Gene expression has been log-normalised and scaled using Seurat(v. 4.3.0). **B i)**

UMAP including all genes shows that there are two Delta/Phi and two Kappa stem clusters,

ii) UMAP showing that removal of all genes specific to the W and Z sex chromosomes

results in one Delta/Phi and one Kappa cluster, indicating that the stem cells are

transcriptionally different in male and female miracidia due to the expression of sex-linked

genes. **C&D)** Multiplexed ISH showing three stem cell markers simultaneously in the same

individual: *ago2-1* (Smp_179320) (pan-stem), *p53* (Smp_139530) (Delta/Phi) and *Uridine*

phosphorylase A (*UPPA*, Smp_308140) (Kappa). **Ci)** *Ago2-1* expression reveals stem cells

lateral and posterior to the brain (Optical section). 100% of individuals examined, n > 30. **ii)**

MIP with segmentation of the 23 *ago2-1*[+] cells. **Di)** *P53* and *UPPA* expression shows the

two stem cell populations intermingled in 100% of individuals examined, n > 30 **ii)** MIP and

segmentation reveal there are more *p53*[+] cells than *UPPA*[+] cells (in this larva,15 are *p53*[+] and

9 are *UPPA*[+]). Scale shown in C also applies to D.

1244

Figure 6. RNA velocity analysis of stem and tegument cells from the miracidium and

sporocyst show lineage-specific gene dynamics. A) UMAP shows the life cycle stage origin

of cells and cell cluster identity from Seurat analysis. **B)** RNA velocity analysis flow field

show the generalised direction of RNA velocity. **C)** Latent time analysis shows an estimated

temporal-relationship between cells. **B)** and **C)** are based on the expression of both spliced

and unspliced transcripts, and their expression dynamics across cells and genes. The phase

plot, velocity, and expression were calculated for **D)** *p53-1*, **E)** *Zfp-1*, **F)** *p-53-2* and **G)** *eled*

(no batch correction). Gene expression has been normalised (CPM) and log-transformed

using scvelo(v. 0.2.4) For all of **D) - G)**, the phase plot (left plot) shows the proportion of

spliced and unspliced transcripts in each cell, where each point is a cell and is coloured by the

clusters in **B)**. The purple almond-shape overlaid represents the processes of transcription,

splicing, and degradation, where this can be modelled. The dashed line shows the estimated

steady state where RNA transcription is constant. The middle panel shows the RNA velocity,

which for each gene is based on how the observations deviated from the estimated steady

state towards induction or repression. The right panel shows gene expression. *p53-1* and *zfp-1*

are predominantly expressed in the Delta/Phi-like miracidia stem cells and some miracidia

tegument, and velocity indicates active expression of *p53-1* but downregulation of *zfp-1* in

the stem cells. *p53-2* is most highly expressed in the Kappa-like miracidia stem cells, and

velocity indicates this gene is being actively transcribed. *eled* expression is very low and only

spliced transcripts have been detected, but is generally restricted to the Kappa-like miracidia

stem cells.

Figure 7. **A model for the fate of the two stem populations in the miracidium**. Adding

single-cell data for miracidia stem cells (this study) to existing stem cell scenarios on

Schistosoma mansoni development (Wang *et al*., 2018; Li *et al*., 2021) shows the continuum

of the Kappa (κ) population from the miracidium, through the intra-molluscan stages to the

juvenile and adult stages inside the mammalian host. Wang *et al*. (2018) proposed that the κ

cells give rise to epsilon (*ε*), *eled*[+], cells in the juvenile primordial testes, ovaries, and

vitellaria (germline), as well as in a gradient increasing toward the posterior growth zone

(soma). They suggested that germ cells may be derived from ε-cells early in juvenile

development, and *eled* is the earliest germline marker yet identified in schistosomes. This led

to the idea that *S. mansoni* does not specify its germline until the juvenile stage (Wang *et al*.,

2018) and a germline-specific regulatory program (including *eled, oc-1, akkn, nanos-1,*
boule) was identified in intra-mammalian stages (Wang *et al.*, 2018; Li *et al.*, 2021). We
show expression of these genes in κ stem cells in the miracidium. This suggests that after ~6
days of embryogenesis, at hatching of the miracidium, the cells that may contribute to the
germline might already be segregated into the κ population and the molecular regulatory
program that differentiates somatic (delta/phi, $\delta/_\varphi$) and germ cell (κ) lineages is present.
Furthermore, as *p53-2* plays a genotoxic stress response role in adult reproductive cells
(Wendt *et al.*, 2022), its expression in κ cells in the miracidia is another line of evidence that
indicates that this population may contain the pluripotent stem cells that likely give rise to the
germline.
1286
1287 **Figure 8**. **Tissue level segmentation of a miracidium reveals the contribution of each**
1288 **tissue to this simple larva**. *In situ* hybridisation using HCR (hybridisation chain reaction)
1289 enabled multiple tissue-level marker genes to be visualised simultaneously in the same larva.
1290 The nuclei of the cells of each tissue type were manually segmented using TrakEM2 in
1291 ImageJ (Cardona *et al.*, 2012).
1292
1293 **<u>Supplementary figure legends</u>**
1294
Supplementary Figure 1. Contributions of sorted and unsorted cells to Seurat clusters.
This figure uses the same UMAP topology as presented in Figure 1C. **A)** Shows contribution
of sorted cells to labelled clusters, **B)** shows contribution of unsorted cells to labelled
clusters. **C)** UMAP where cells are coloured by sorted and unsorted treatment. **D)** UMAP
where cells are coloured by sample of origin (samples 1 and 4 are sorted, samples 2 and 3 are
unsorted).
1301
Supplementary Figure 2. Subclustering the Neuron 1 cluster using SAM revealed
multiple distinct subpopulations. **A)** The Neuron 1 cluster was analysed using Self-
Assembling Manifold (SAM) algorithm, and Leiden clustering indicated 16 subpopulations.
Library size was normalised and gene expression values were log-normalised using SAM
(v1.0.1) and Scanpy (v1.8.2). **B)** The top five marker genes for each of the subpopulations.
C) Expression of NPP PWamide, the top marker for subcluster 14. **D)** Expression of
neuropeptide Y, the top marker for subcluster 13. **E)** Expression of neuropeptide F, the

second top marker for subcluster 8. Although this cluster was difficult to define, this analysis

suggests that there may be several small neural subpopulations within this cluster.

1311

Supplementary Figure 3. Neurons. A & B) Top markers for cell cluster Neuron 1 include

the neuropeptides precursor genes for Neuropeptide F, PWamide, and Neuropeptide Y. ISH

of these genes shows expression in cells that make up the brain. **C)** G-alpha-q antibody

staining reveals a girdle of 23 multi-ciliated nerve endings that sit between the 2nd and 3rd tier

of ciliary plates. **D)** Neuron 4 marker Smp_319480 (putative mTORC1) transcripts are

expressed at the base of these cilia; **E)** and some transcripts also reach the nuclei of circular

body wall muscles. **F)** The expression of *p25 alpha* (which is a marker for ciliated cells in

adult worms, Wendt *et al*., 2020) in the submuscular cells suggests that they may be ciliated.

1320

Supplementary Figure 4. **Phylogeny of metazoan opsins.** There are four *Schistosoma*

mansoni opsins that are expressed in two sub-clusters of Neuron 1; our analysis shows that

three belong to the rhabdomeric opsin clade and the fourth is a peropsin. Scale bar shows

branch lengths in expected amino acid substitutions per site. Shading of circles at internal

nodes shows percentage support for the partition implied by that node from 1,000 non-

parametric bootstrap samples.

1327

Supplementary Figure 5. **The expression of two Polycystin genes and a Sox**

transcription factor in Neuron 4 cells. A) Polycystin 1 and 2 are co-expressed in Neuron 4

cells, orthologs in other animals form heterodimers in ciliated cells that function to detect

flow sheer and vibrations (Kim *et al*., 2016; Calderon *et al*., 2018). Gene expression has been

log-normalised and scaled using Seurat(v. 4.3.0). **B)** In planaria, a transcription factor Sox

B1-2 regulates polycystin genes and together function in sensory neurons that detect flow.

The co-expression of polycystin genes with a Sox transcription factor in Neuron 4 cells

indicates that this transcription factor may regulate these polycystin genes in the miracidium.

1336

1337 **Supplementary Figure 6. STRING analysis of neural clusters and ciliary plates.**

1338 STRING analysis of the top 100 marker genes for each of the neural populations whose

1339 predicted networks have functional enrichment. **A)** Neuron 4, **B)** Neuron 5, **C)** Ciliary plates,

D) Neuron 3. Neuron 1 is not a defined cluster and Neuron 2 showed no functional enrichment. Lines (edges) connecting nodes are based on evidence of the function of homologs. Functional enrichment (*FDR*) as provided by STRING. (PPI = predicted protein interaction). Minimum interaction (confidence) score of 0.7, corresponding to a high level of confidence, 0.4 medium level of confidence.

Supplementary Figure 7. Gene expression by transcripts per million in the ciliary plates and gland cells. Heatmap showing the gene expression, by transcripts per million, detected in the ciliary plates and gland cells hand-picked for plate-based scRNA-seq. Genes included; the top 20 significant genes between gland cell and ciliary plates, two neural markers, five VAL genes, five significant genes between gland cells and other cells, eight genes previously reported to produce secreted proteins in the miracidia (Wang *et al.*, 2016; Wu *et al.*, 2006), five top marker genes from the ciliary plate cluster.

Supplementary Figure 8. STRING analysis of tegument and protonephridia clusters. STRING analysis of the top 100 marker genes for **A)** tegument, **B)** protonephridia. Lines (edges) connecting nodes are based on evidence of the function of homologs. Functional enrichment (*FDR*) as provided by STRING. (PPI = predicted protein interaction). Minimum interaction (confidence) score of 0.7, corresponding to a high level of confidence, 0.4 medium level of confidence.

Supplementary Figure 9. STRING analysis of parenchyma clusters. STRING analysis of the top 100 marker genes for **A)** Parenchyma 1, **B)** Parenchyma 2. Lines (edges) connecting nodes are based on evidence of the function of homologs. Functional enrichment (*FDR*) as provided by STRING (PPI = predicted protein interaction). Minimum interaction (confidence) score of 0.7, corresponding to a high level of confidence, 0.4 medium level of confidence.

Supplementary Figure 10. Gene ontology (GO) analysis of marker genes identified in the miracidia cell clusters. GO analysis was performed on each cluster using genes with a minimum AUC score of 0.7. Enriched biological process (BP) GO terms are shown for each cluster, and have been filtered to only show terms supported by a minimum of two genes.

1372 Top panels (left to right) show parenchyma, muscle, and tegument clusters, middle panels

1373 (left to right) show ciliary plate, neuron, and protonephridia clusters, and the bottom panel

1374 shows stem clusters. Full list of GO terms available in Supplementary Table 7.

1375

Supplementary Figure 11. Expression of known stem cell markers in miracidia cells.

A-F) Three stem cell classes were identified in the mother sporocyst stage and were named

based upon their respective markers: Kappa (*klf*[+], *nanos-2*[+]), Delta (*nanos-2*[+] and *fgfrA*[+],

fgfrB[+] and *nanos-2*[+]) and Phi (*fgfrA*[+] and, *fgfrB*[+] and *hesl*[+]), as well as pan-stem marker *ago2-*

1 (Wang *et al.*, 2018; Safarti *et al.*, 2021). In the miracidium, we identified seven stem cell

clusters by expression of *ago2-1*: stem A and B are *klf*[+] and *nanos-2*[+] (i.e. Kappa-like), stem

C and D are *nanos-2*[+], *fgfrA*[+], *fgfrB*[+], and *hesl*[+] (i.e. they resembled both Delta and Phi).

Gene expression has been log-normalised and scaled using Seurat(v. 4.3.0).

1384

Supplementary Figure 12 - The effect of sex-linked genes on cell clustering.

Schistosoma mansoni has 7 pairs of autosomes and one pair of sex chromosomes; males have

ZZ and females have ZW (Buddenborg *et al.*, 2021). The sex chromosomes are composed of

sex-specific regions that are unique to each chromosome flanked by pseudoautosomal regions

that are common to both Z and W. The W-specific region (WSR) is a large, highly repetitive

region that contains 38 genes, most of which are gametologs (genes with homologous copies

on the WSR and ZSR). The Z-specific region (ZSR) contains 941 genes, of which only 35 are

gametologs. **A & B)** The proportion of reads mapping to the WSR and ZSR across all cell

type clusters shows **A)** putative male cells (Stem A and C i.e. those that lack WSR gene

expression) and **B)** evidence of incomplete ZSR gene dosage compensation in the male stem

cells. **C)** To determine whether any of the WSR genes were responsible for the clustering

pattern, we excluded them from the mapping reference. The impact on the overall UMAP

clustering was minimal. We repeated this excluding the **D)** ZSR genes, **E)** gametologues and

F) ZSR genes, except the gametologues, in turn. In all four scenarios the stem cell clusters

remained separated by sex. ZSR or WSR expression is therefore sufficient on their own to

split these stem clusters in two.

1401

1402 **Supplementary Figure 13. Identification of new clusters after removing ZSR and WSR**

1403 **genes.** To identify whether stem cell clusters A and B, and C and D, still split down A/B and

1404 C/D lines despite the apparent collapsing of clusters after removing ZSR and WSR genes, we

1405 ran de novo clustering at several resolutions. Even once clusters were oversplitting, there was

not a separation between stem cluster A/B and C/D. This was run using Seurat (v5.0.2), so the overall topology is not identical to that shown in Figure 5, though the same number of PCs are used. **A)** Shows the UMAP topology labelled by original cluster ID, **B)** shows the de novo clusters at resolution = 0.6, and **C)** shows the de novo clusters at resolution = 1.

Supplementary Figure 14. Classification of cells based on percentage of reads mapping to W-specific region indicates incomplete dosage compensation of Z-specific region genes. **A)** The percentage of reads mapping to W-specific genes by cluster. The blue line at 0.2% indicates where the data were split into W+ and W- categories, i.e. Stem B and D contain cells with reads mapping to W-specific genes (and are likely cells from female miracidia), whereas most cells in Stem A and C do not have reads mapping to W-specific genes (cells from male miracidia). **B)** UMAP showing the clusters by W+ and W- categories. Almost complete separation can be seen in the main stem cell clusters, and the categories are mixed in most other clusters.

Supplementary Figure 15. EdU labelling shows that there are no dividing cells during the free-swimming miracidia stage. **A)** 4 hour EdU pulse on miracidia 0-6 hours post-hatching shows no EdU incorporation in any cells. **B)** Negative control. **C)** Positive control, three day EdU pulse on freshly-transformed mother sporocysts shows many dividing cells incorporating EdU, all treatments n = 30.

Supplementary Figure 16. STRING analysis of the top 100 marker genes for the two stem cell populations in the miracidium (combining the female and male cells). A) 61 of the top markers of Stem C and D combined (i.e. Delta/Phi-like) and **B)** 37 for Stem A and B combined (i.e. Kappa-like) form large predicted interactions enriched for translation, ribosome biogenesis and nucleic acid binding. Lines (edges) connecting nodes are based on evidence of the function of homologues. Functional enrichment (*FDR*) as provided by STRING. PPI = predicted protein interaction. Minimum interaction (confidence) score of 0.7, corresponding to a high level of confidence.

Supplementary Figure 17. Differential gene expression between the two stem cell populations, Delta/Phi and Kappa. **A)** Heatmap of top differentially expressed genes (by adjusted p-value) from each of Delta/Phi and Kappa populations show a high number of transcription factors (bold) in Delta/Phi and enrichment of lipid and glycolytic metabolism

1440   genes in Kappa. Gene expression has been log-normalised and scaled using Seurat(v. 4.3.0).

1441   **B)** Gene ontology analysis of DEG (adj p<0.001) revealed that upregulated genes in

1442   Delta/Phi were related to the structural constituent of the ribosome and translation, while up-

1443   regulated genes in Kappa were related to transcription and metabolism**.**

1444

Supplementary Figure 18. RNA velocity analysis of stem and tegument cells from the

miracidium and sporocyst (with batch correction) show lineage-specific gene dynamics.

A) UMAP shows the life cycle stage origin of cells and cell cluster identity from Seurat

analysis. **B)** RNA velocity analysis flow field shows generalised direction of RNA velocity.

C) Latent time analysis shows an estimated temporal-relationship between cells. The phase

plot, velocity, and expression were calculated for **D)** *p53-1*, **E)** *Zfp-1*, **F)** *p-53-2* and **G)** *eled*.

Gene expression has been normalised (CPM) and log-transformed using scvelo(v. 0.2.4).

p53-1 and *zfp-1* are predominantly expressed in the Delta/Phi-like miracidia stem cells and

miracidia tegument, and velocity indicates active expression of *p53-1* but downregulation of

zfp-1 in the stem cells. *p53-2* is most highly expressed in the Kappa-like miracidia stem cells,

and velocity indicates this gene is being actively transcribed. *eled* expression is very low and

only spliced transcripts have been detected, but is generally restricted to the Kappa-like

miracidia stem cells.

1458

Supplementary Figure 19. RNA velocity analysis of stem and tegument cells from the

miracidium and sporocyst (without batch correction). A) UMAP shows the life cycle

stage origin of cells and cell cluster identity from Seurat analysis. **B)** RNA velocity analysis

flow field shows generalised direction of RNA velocity. Phase plot, velocity and expression

as calculated for genes mentioned in text: blue outline = known molecular regulators of

tegument development and tegument markers in adult *S. mansoni* and planaria. Green outline

= miracidia and sporocyst tegument marker genes. Red outline = genes revealed by RNA

velocity analysis to be highly dynamic in the Delta/Phi, miracidia and sporocyst tegument

cells. Gene expression has been normalised (CPM) and log-transformed using scvelo(v.

0.2.4).

1469

Supplementary Figure 20. RNA velocity analysis of stem and tegument cells from the

miracidium and sporocyst (without batch correction). A) UMAP shows the life cycle

stage origin of cells and cell cluster identity from Seurat analysis. **B)** RNA velocity analysis

flow field shows generalised direction of RNA velocity. Phase plot, velocity and expression

as calculated for genes that are mentioned in the text: red outline = known molecular

regulators of germline stem cells in adult *S. mansoni*. Blue outline = pluripotency gene. *Oc-1*

expression is low with little velocity signal, but the expression is predominantly seen in the

Kappa-like miracidia stem cells (and Stem G). *Irx* is also primarily expressed in the Kappa-

like miracidia stem cells with velocity results indicating an accumulation of spliced

transcripts. *Irx* transcripts are also detected in Stem G and the sporocyst stem/germinal cells.

Akkn expression is low and largely seen in the Kappa-like and Stem G miracidia stem

clusters. Lsm14-domain containing protein transcripts are most consistently expressed in the

Kappa-like miracidia stem cells. *Klf* is most highly expressed in sporocyst stem/germinal

cells, and the Kappa-like miracidia stem cluster. Gene expression has been normalised

(CPM) and log-transformed using scvelo(v. 0.2.4).

1485

Supplementary Figure 21. RNA velocity analysis of all tissue types from the miracidium

and sporocyst show lineage-specific gene dynamics. A) UMAP shows the life cycle stage

origin of cells and **B)** shows cell cluster identity from Seurat analysis. **C)** RNA velocity

analysis flow field shows the generalised direction of RNA velocity. **D)** Latent time analysis

shows an estimated temporal-relationship between cells. Phase plot, velocity and expression

as calculated for **E)** *p53-1*, **F)** *Zfp-1*, **G)** *p-53-2* and **H)** *eled* (no batch correction, sporocyst

cells combined with miracidia sample 1 cells). *P53-1* is most highly expressed in the

Delta/phi-like miracidia stem clusters and some of the miracidia tegument cells, and the

velocity indicates active transcription. *Zfp-1* is expressed most highly in the same clusters,

but velocity suggests that the majority of transcripts are spliced. *P53-2* expression is highest

in the Kappa-like miracidia stem clusters, but the expression is detected in several other

clusters, including muscle, neuron, and other stem clusters. Velocity indicates that this gene

is being actively transcribed in the majority of these clusters. *Eled* expression is very low, but

restricted to the Kappa-like miracidia stem clusters. Gene expression has been normalised

(CPM) and log-transformed using scvelo(v. 0.2.4)

1501

Supplementary Figure 22. RNA velocity analysis of all tissue types from the miracidium

and sporocyst shows the dynamics of described germline-related genes. A) UMAP shows

the life cycle stage origin of cells and **B)** shows cell cluster identity from Seurat analysis. **C)**

RNA velocity analysis flow field shows the generalized direction of RNA velocity. Phase

plot, velocity and expression as calculated for **D)** *oc-1*, **E)** *boule*, **F)** *irx,* **G)** *akkn,* and **H)** an

FDF domain-containing protein (no batch correction, sporocyst cells combined with

miracidia sample 1 cells.). *Oc-1* expression is low but generally specific to the two Kappa-like miracidia stem cell clusters, and Stem G. Only spliced *Boule* transcripts were detected, and with low expression, predominantly in the Stem B and Stem E clusters. *Irx* was most highly expressed in Kappa-like miracidia stem clusters, as well as Stem E, Stem G, and sporocyst stem/germinal cells. *Akkn* is also largely expressed in the Kappa-like miracidia stem clusters. The FDF domain-containing protein expression is low, but seen most in Kappa-like miracidia stem clusters, Stem G, and several neural clusters. Gene expression has been normalised (CPM) and log-transformed using scvelo(v. 0.2.4).

Supplementary Figure 23. RNAvelocity analysis of all tissue types from the miracidium and sporocyst show lineage-specific gene dynamics. A) UMAP shows the life cycle stage origin of cells and **B)** shows cell cluster identity from Seurat analysis. **C)** RNA velocity analysis flow field shows the generalised direction of RNA velocity. **D)** Latent time analysis shows an estimated temporal-relationship between cells. Phase plot, velocity and expression as calculated for **E)** *p53-1*, **F)** *Zfp-1*, **G)** *p-53-2* and **H)** *eled* (no batch correction, sporocyst cells combined with miracidia sample 4 cells). *p53-1* and *zfp-1* are predominantly in the Delta/Phi-like miracidia stem cell clusters (Stem C and D) and also in the miracidia tegument. Velocity indicates that *p53-1* is being induced in the stem cells and repressed in the tegument, while *zfp-1* shows a gradient of velocity in the stem cells and repression in the tegument. *P53-2* is most highly expressed in the Kappa-like miracidia stem cells (Stem A and B) and Stem F, with velocity results indicating active transcription. *eled* expression was very low, but primarily in the Kappa-like miracidia stem cells (particularly Stem A). Gene expression has been normalised (CPM) and log-transformed using scvelo(v. 0.2.4).

References

1. Adam M, Potter AS, Potter SS. Psychrophilic proteases dramatically reduce single-cell RNA-seq artifacts: a molecular atlas of kidney development. Development. 2017;144: 3625–3632. doi:10.1242/dev.151142

2. Alexa A, Rahnenfuhrer J. topGO: Enrichment Analysis for Gene Ontology. In: R package version 2.54.0 [Internet]. 2023. doi:10.18129/B9.bioc.topGO

3. Ashton PD, Harrop R, Shah B, Wilson RA. The schistosome egg: development and

secretions. Parasitology. 2001;122: 329–338. doi:10.1017/s0031182001007351

4. Bergen V, Lange M, Peidli S, Wolf FA, Theis FJ. Generalizing RNA velocity to

transient cell states through dynamical modeling. Nat Biotechnol. 2020;38:

1408–1414. doi:10.1038/s41587-020-0591-3

5. Bezares-Calderón LA, Berger J, Jasek S, Verasztó C, Mendes S, Gühmann M, et al.

Neural circuitry of a polycystin-mediated hydrodynamic startle response for

predator avoidance. Elife. 2018;7. doi:10.7554/eLife.36262

6. Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq

quantification. Nat Biotechnol. 2016;34: 525–527. doi:10.1038/nbt.3519

7. Buddenborg SK, Lu Z, Sankaranarayan G, Doyle SR, Berriman M. The stage- and

sex-specific transcriptome of the human parasite Schistosoma mansoni. Sci

Data. 2023;10: 775. doi:10.1038/s41597-023-02674-2

8. Buddenborg SK, Tracey A, Berger DJ, Lu Z, Doyle SR, Fu B, et al. Assembled

chromosomes of the blood fluke *Schistosoma mansoni* provide insight into the

evolution of its ZW sex-determination system. bioRxiv. 2021;

2021.08.13.456314-2021.08.13.456314. doi:10.1101/2021.08.13.456314

9. Cardona A, Saalfeld S, Schindelin J, Arganda-Carreras I, Preibisch S, Longair M, et

al. TrakEM2 software for neural circuit reconstruction. PLoS One. 2012;7:

e38011–e38011. doi:10.1371/journal.pone.0038011

10. Chernin E. Interference with the capacity of Schistosoma mansoni miracidia to

infect the molluscan host. J Parasitol. 1968;54: 509–516.

11. Choi HMT, Calvert CR, Husain N, Huss D, Barsi JC, Deverman BE, et al.

Mapping a multiplexed zoo of mRNA expression. Development. 2016;143:

3632–3637. doi:10.1242/dev.140137

12. Choi HMT, Schwarzkopf M, Fornace ME, Acharya A, Artavanis G, Stegmaier J,

et al. Third-generation in situ hybridization chain reaction: multiplexed,

quantitative, sensitive, versatile, robust. Development. 2018;145.
doi:10.1242/dev.165753

13. Collins JJ 3rd, King RS, Cogswell A, Williams DL, Newmark PA. An atlas for
Schistosoma mansoni organs and life-cycle stages using cell type-specific
markers and confocal microscopy. PLoS Negl Trop Dis. 2011;5: e1009–e1009.
doi:10.1371/journal.pntd.0001009

14. Costain AH, MacDonald AS, Smits HH. Schistosome Egg Migration:
Mechanisms, Pathogenesis and Host Immune Responses. Front Immunol.
2018;9: 3042. doi:10.3389/fimmu.2018.03042

15. Diaz Soria CL, Attenborough T, Lu Z, Fontenla S, Graham J, Hall C, Thompson S,
Andrews TGR, Rawlinson KA, Berriman M, Rinaldi G. Single-cell
transcriptomics of the human parasite Schistosoma mansoni first intra-
molluscan stage reveals tentative tegumental and stem-cell regulators. Sci Rep.
2024 Mar 12;14(1):5974. doi: 10.1038/s41598-024-55790-3. PMID: 38472267;
PMCID: PMC10933418.

16. Diaz Soria CL, Lee J, Chong T, Coghlan A, Tracey A, Young MD, et al. Single-
cell atlas of the first intra-mammalian developmental stage of the human
parasite Schistosoma mansoni. Nat Commun. 2020;11: 6411.
doi:10.1038/s41467-020-20092-5

17. Fishman L, Nechooshtan G, Erhard F, Regev A, Farrell JA, Rabani M. Single-cell
temporal dynamics reveals the relative contributions of transcription and
degradation to cell-type specific gene expression in zebrafish embryos. bioRxiv
[Preprint]. 2023 Apr 21:2023.04.20.537620. doi: 10.1101/2023.04.20.537620.
PMID: 37131717; PMCID: PMC10153228.

18. Gąsiorowski L, Andrikou C, Janssen R, Bump P, Budd GE, Lowe CJ, et al.
Molecular evidence for a single origin of ultrafiltration-based excretory organs.
Curr Biol. 2021;31: 3629-3638.e2. doi:10.1016/j.cub.2021.05.057

19. Hao Y, Hao S, Andersen-Nissen E, Mauck WM 3rd, Zheng S, Butler A, et al.
1595   Integrated analysis of multimodal single-cell data. Cell. 2021;184: 3573-
1596   3587.e29. doi:10.1016/j.cell.2021.04.048

1597   20. Howe KL, Bolt BJ, Cain S, Chan J, Chen WJ, Davis P, et al. WormBase 2016:
1598   expanding to enable helminth genomic research. Nucleic Acids Res. 2016;44:
1599   D774-80. doi:10.1093/nar/gkv1217

1600   21. Howe KL, Bolt BJ, Shafie M, Kersey P, Berriman M. WormBase ParaSite - a
1601   comprehensive resource for helminth genomics. Mol Biochem Parasitol.
1602   2017;215: 2–10. doi:10.1016/j.molbiopara.2016.11.005

1603   22. IHME. 2019 Global Burden of Disease (GBD) study [Internet]. 2019. Available:
1604   https://vizhub.healthdata.org/gbd-results?params=gbd-api-2019-
1605   permalink/d30124bc2eaad3d977583362d4d37591 [vizhub.healthdata.org]

1606   23. Jurberg AD, Gonçalves T, Costa TA, de Mattos ACA, Pascarelli BM, de Manso
1607   PPA, et al. The embryonic development of Schistosoma mansoni eggs: proposal
1608   for a new staging system. Dev Genes Evol. 2009;219: 219–234.
1609   doi:10.1007/s00427-009-0285-9

1610   24. Kakugawa S, Langton PF, Zebisch M, Howell S, Chang T-H, Liu Y, et al. Notum
1611   deacylates Wnt proteins to suppress signalling activity. Nature. 2015;519: 187–
1612   192. doi:10.1038/nature14259

1613   25. Kharchenko P V, Silberstein L, Scadden DT. Bayesian approach to single-cell
1614   differential expression analysis. Nat Methods. 2014;11: 740–742.
1615   doi:10.1038/nmeth.2967

1616   26. Kim S, Nie H, Nesin V, Tran U, Outeda P, Bai C-X, et al. The polycystin complex
1617   mediates Wnt/Ca2+ signalling. Nat Cell Biol. 2016;18: 752–764.
1618   doi:10.1038/ncb3363

1619   27. Kumar P, Tan Y, Cahan P. Understanding development and stem cells using single
1620   cell-based analyses of gene expression. Development. 2017;144: 17–32.
1621   doi:10.1242/dev.133058

28. La Manno G, Soldatov R, Zeisel A, Braun E, Hochgerner H, Petukhov V, et al.
RNA velocity of single cells. Nature. 2018;560: 494–498. doi:10.1038/s41586-
018-0414-6

29. Li P, Nanes Sarfati D, Xue Y, Yu X, Tarashansky AJ, Quake SR, et al. Single-cell
analysis of Schistosoma mansoni identifies a conserved genetic program
controlling germline stem cell fate. Nat Commun. 2021;12: 485.
doi:10.1038/s41467-020-20794-w

30. Llorens-Bobadilla E, Zhao S, Baser A, Saiz-Castro G, Zwadlo K, Martin-Villalba
A. Single-Cell Transcriptomics Reveals a Population of Dormant Neural Stem
Cells that Become Activated upon Brain Injury. Cell Stem Cell. 2015;17: 329–
340. doi:10.1016/j.stem.2015.07.002

31. Lun ATL, Riesenfeld S, Andrews T, Dao TP, Gomes T, Marioni JC, et al.
EmptyDrops: distinguishing cells from empty droplets in droplet-based single-
cell RNA sequencing data. Genome Biol. 2019;20: 63. doi:10.1186/s13059-
019-1662-y

32. Mair GR, Maule AG, Day TA, Halton DW. A confocal microscopical study of the
musculature of adult Schistosoma mansoni. Parasitology. 2000;121 (Pt 2: 163–
170. doi:10.1017/s0031182099006174

33. Maldonado JF, Acosta-Matienzo J. Biological Studies on the Miracidium of
Schistosoma Mansoni. Am J Trop Med. Arlington VA, USA: The American
Society of Tropical Medicine and Hygiene; 1948;s1-28: 645–657.
doi:10.4269/ajtmh.1948.s1-28.645

34. Mann VH, Morales ME, Rinaldi G, Brindley PJ. Culture for genetic manipulation
of developmental stages of Schistosoma mansoni. Parasitology. 2010;137: 451–
462. doi:10.1017/S0031182009991211

35. McGinnis CS, Murrow LM, Gartner ZJ. DoubletFinder: Doublet Detection in
Single-Cell RNA Sequencing Data Using Artificial Nearest Neighbors. Cell
Syst. 2019;8: 329-337.e4. doi:https://doi.org/10.1016/j.cels.2019.03.003
[doi.org]

36. McManus DP, Dunne DW, Sacko M, Utzinger J, Vennervald BJ, Zhou X-N.
Schistosomiasis. Nat Rev Dis Prim. 2018;4: 13. doi:10.1038/s41572-018-0013-
8

37. Melsted P, Booeshaghi AS, Liu L, Gao F, Lu L, Min KH (Joseph), et al. Modular,
efficient and constant-memory single-cell RNA-seq preprocessing. Nat
Biotechnol. 2021;39: 813–818. doi:10.1038/s41587-021-00870-2

38. Moses L LP. BUSpaRse: kallisto | bustools R utilities [Internet]. 2023.
doi:10.18129/B9.bioc.BUSpaRse

39. Nanes Sarfati D, Li P, Tarashansky AJ, Wang B. Single-cell deconstruction of
stem-cell-driven schistosome development. Trends Parasitol. 2021;37: 790–802.
doi:https://doi.org/10.1016/j.pt.2021.03.005 [doi.org]

40. Orosz F, Ovádi J. TPPP orthologs are ciliary proteins. FEBS Lett. 2008;582: 3757–
3764. doi:10.1016/j.febslet.2008.10.011

41. Pan SC. The fine structure of the miracidium of Schistosoma mansoni. J Invertebr
Pathol. 1980;36: 307–372. doi:https://doi.org/10.1016/0022-2011(80)90040-3
[doi.org]

42. Petersen CP, Reddien PW. Wnt signaling and the polarity of the primary body
axis. Cell. 2009;139: 1056–1068. doi:10.1016/j.cell.2009.11.035

43. Picard MAL, Vicoso B, Roquis D, Bulla I, Augusto RC, Arancibia N, et al. Dosage
Compensation throughout the Schistosoma mansoni Lifecycle: Specific
Chromatin Landscape of the Z Chromosome. Genome Biol Evol. 2019;11:
1909–1922. doi:10.1093/gbe/evz133

44. Pimentel H, Bray NL, Puente S, Melsted P, Pachter L. Differential analysis of
RNA-seq incorporating quantification uncertainty. Nat Methods. 2017;14: 687–
690. doi:10.1038/nmeth.4324

45. Poteaux P, Gourbal B, Duval D. Time series analysis of tegument ultrastructure of
in vitro transformed miracidium to mother sporocyst of the human parasite

Schistosoma mansoni. Acta Trop. 2023 Apr;240:106840. doi:
10.1016/j.actatropica.2023.106840. Epub 2023 Jan 18. PMID: 36681315.

46. Scimone ML, Cote LE, Reddien PW. Orthogonal muscle fibres have different
instructive roles in planarian regeneration. Nature. 2017;551: 623–628.
doi:10.1038/nature24660

47. Shin J, Berg DA, Zhu Y, Shin JY, Song J, Bonaguidi MA, et al. Single-Cell RNA-
Seq with Waterfall Reveals Molecular Cascades underlying Adult
Neurogenesis. Cell Stem Cell. 2015;17: 360–372.
doi:https://doi.org/10.1016/j.stem.2015.07.013 [doi.org]

48. Siebert S, Farrell JA, Cazet JF, Abeykoon Y, Primack AS, Schnitzler CE, et al.
Stem cell differentiation trajectories in Hydra resolved at single-cell resolution.
Science (80-). American Association for the Advancement of Science;
2019;365: eaav9314–eaav9314. doi:10.1126/science.aav9314

49. Steger J, Cole AG, Denner A, Lebedeva T, Genikhovich G, Ries A, Reischl R,
Taudes E, Lassnig M, Technau U. Single-cell transcriptomics identifies
conserved regulators of neuroglandular lineages,Cell Reports, 40, 12,2022,
111370, https://doi.org/10.1016/j.celrep.2022.111370.

50. Szklarczyk D, Gable AL, Lyon D, Junge A, Wyder S, Huerta-Cepas J, et al.
STRING v11: protein–protein association networks with increased coverage,
supporting functional discovery in genome-wide experimental datasets. Nucleic
Acids Res. 2019;47: D607–D613. doi:10.1093/nar/gky1131

51. Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, Tomoda K, et al.
Induction of Pluripotent Stem Cells from Adult Human Fibroblasts by Defined
Factors. Cell. 2007;131: 861–872. doi:https://doi.org/10.1016/j.cell.2007.11.019
[doi.org]

52. Tarashansky AJ, Musser JM, Khariton M, Li P, Arendt D, Quake SR, et al.
Mapping single-cell atlases throughout Metazoa unravels cell type evolution.
Shalek AK, Barkai N, editors. Elife. eLife Sciences Publications, Ltd; 2021;10:
e66747–e66747. doi:10.7554/eLife.66747

53. Tarashansky AJ, Xue Y, Li P, Quake SR, Wang B. Self-assembling manifolds in
single-cell RNA sequencing data. Shalek AK, Barkai N, Yanai I, editors. Elife.
eLife Sciences Publications, Ltd; 2019;8: e48994–e48994.
doi:10.7554/eLife.48994

54. Valsecchi CIK, Basilicata MF, Semplicio G, Georgiev P, Gutierrez NM, Akhtar A.
Facultative dosage compensation of developmental genes on autosomes in
Drosophila and mouse embryonic stem cells. Nat Commun. 2018;9: 3626.
doi:10.1038/s41467-018-05642-2

55. van Kempen M, Kim SS, Tumescheit C, Mirdita M, Lee J, Gilchrist CLM, et al.
Fast and accurate protein structure search with Foldseek. Nat Biotechnol. 2023;
doi:10.1038/s41587-023-01773-0

56. Verasztó C, Jasek S, Gühmann M, Shahidi R, Ueda N, Beard JD, et al. Whole-
animal connectome and cell-type complement of the three-segmented
Platynereis dumerilii larva. bioRxiv. 2020; 2020.08.21.260984-
2020.08.21.260984. doi:10.1101/2020.08.21.260984

57. Vergara HM, Pape C, Meechan KI, Zinchenko V, Genoud C, Wanner AA, et al.
Whole-body integration of gene expression and single-cell morphology. Cell.
2021;184: 4819-4837.e22. doi:https://doi.org/10.1016/j.cell.2021.07.017
[doi.org]

58. Vogg MC, Owlarn S, Pérez Rico YA, Xie J, Suzuki Y, Gentile L, et al. Stem cell-
dependent formation of a functional anterior regeneration pole in planarians
requires Zic and Forkhead transcription factors. Dev Biol. 2014;390: 136–148.
doi:https://doi.org/10.1016/j.ydbio.2014.03.016 [doi.org]

59. Wang B, Collins III JJ, Newmark PA. Functional genomic characterization of
neoblast-like stem cells in larval Schistosoma mansoni. Sánchez Alvarado A,
editor. Elife. eLife Sciences Publications, Ltd; 2013;2: e00768–e00768.
doi:10.7554/eLife.00768

60. Wang B, Lee J, Li P, Saberi A, Yang H, Liu C, et al. Stem cell heterogeneity
drives the parasitic life cycle of Schistosoma mansoni. Sánchez Alvarado A,

1736 editor. Elife. eLife Sciences Publications, Ltd; 2018;7: e35449–e35449.

1737 doi:10.7554/eLife.35449

1738 61. Wang T, Zhao M, Rotgans BA, Strong A, Liang D, Ni G, et al. Proteomic Analysis

1739 of the Schistosoma mansoni Miracidium. PLoS One. Public Library of Science;

1740 2016;11: e0147247–e0147247.

1741 62. Wendt GR, Collins JNR, Pei J, Pearson MS, Bennett HM, Loukas A, et al.

1742 Flatworm-specific transcriptional regulators promote the specification of

1743 tegumental progenitors in Schistosoma mansoni. Patel NH, editor. Elife. eLife

1744 Sciences Publications, Ltd; 2018;7: e33221–e33221. doi:10.7554/eLife.33221

1745 63. Wendt GR, Shiroor DA, Adler CE, Collins JJ. Convergent evolution of a genotoxic

1746 stress response in a parasite-specific p53 homolog. Proc Natl Acad Sci.

1747 Proceedings of the National Academy of Sciences; 2022;119: e2205201119–

1748 e2205201119. doi:10.1073/pnas.2205201119

1749 64. Wendt G, Zhao L, Chen R, Liu C, O'Donoghue AJ, Caffrey CR, et al. A single-

1750 cell RNA-seq atlas of Schistosoma mansoni identifies a key regulator of blood

1751 feeding. Science (80-). 2020;369. doi:10.1126/science.abb7709

1752 65. WHO. Schistosomiasis [Internet]. 2023. Available: https://www.who.int/news-

1753 room/fact-sheets/detail/schistosomiasis [who.int]

1754 66. Witchley JN, Mayer M, Wagner DE, Owen JH, Reddien PW. Muscle Cells

1755 Provide Instructions for Planarian Regeneration. Cell Rep. 2013;4: 633–641.

1756 doi:https://doi.org/10.1016/j.celrep.2013.07.022 [doi.org]

1757 67. Wolf FA, Angerer P, Theis FJ. SCANPY: large-scale single-cell gene expression

1758 data analysis. Genome Biol. 2018;19: 15. doi:10.1186/s13059-017-1382-0

1759 68. Zheng GXY, Terry JM, Belgrader P, Ryvkin P, Bent ZW, Wilson R, et al.

1760 Massively parallel digital transcriptional profiling of single cells. Nat Commun.

1761 2017;8: 14049. doi:10.1038/ncomms14049

1762

1763

Supplementary Video legends

Supplementary Video 1 – Confocal z-stack of wholemount *in situ* hybridisation (WISH) of pan-muscle marker *paramyosin* (*PRM* Smp_085540) (cyan), counterstained with phalloidin (green) and DAPI (white), reveals which *PRM*[+] nuclei belong to the circular and longitudinal body wall muscles.

Supplementary Video 2 – Segmentation of *paramyosin* (Smp_085540) (cyan) positive cells (using the z-stack from Supplementary Figure 2); the nuclei of the circular body wall muscles are highlighted in orange, the nuclei of the longitudinal body wall muscles are highlighted in yellow.

Supplementary Video 3 - Confocal z-stack of WISH of muscle cluster 1 markers *Kunitz-type protease inhibitor* (Smp_052230) (cyan) and *Wnt-11-1* (Smp_156540) (magenta), counterstained with phalloidin (green) and DAPI (white), show expression in circular body wall muscles.

Supplementary Video 4 – Maximum intensity projection from confocal z-stack of WISH of muscle cluster 1 markers *notum* (Smp_015820) (cyan) and *Wnt-11-1* (Smp_156540) (magenta) counterstained with phalloidin (green) showing expression at opposite poles.

Supplementary Video 5 - Confocal z-stack of WISH of muscle cluster 2 markers *calcium activated potassium channel* (Smp_156150) (cyan) and *myoD* (Smp_167400) (magenta), counterstained with phalloidin (green) and DAPI (white), shows expression in longitudinal body wall muscles.

Supplementary Video 6 - Confocal z-stack of WISH of markers for neuron cluster 2 (Smp_201600) (magenta) and neuron 3 (Smp_071050)(cyan), counterstained with DAPI (white), shows expression of both genes in cells whose nuclei form part of the nuclear rind of the brain.

Supplementary Video 7 - Confocal z-stack of WISH of ciliary plate marker *16 kDa calcium*

binding protein (Smp_096390), counterstained with phalloidin (green) and DAPI (white),

shows expression in the ciliary plates and six previously undescribed cells that we term sub-

muscular cells. The nuclei of the sub-muscular cells sit beneath the body wall muscles, and

each cell has a protrusion (possible cilia) that extends beyond the body wall muscles, between

the first and second tiers of ciliary plates, and out to the exterior of the larva.

1803

1804 **Supplementary Video 8 –** Maximum intensity projection of confocal z-stack of WISH of

1805 ciliary plate marker *16 kDa calcium binding protein* (Smp_096390), counterstained with

1806 phalloidin (green) and DAPI (white), shows expression in ciliary plates and sub-muscular

1807 cells.

1808

1809 **Supplementary Video 9 –** Maximum intensity projection from confocal z-stack of WISH of

1810 neuron 2 marker Smp_319480 (magenta), counterstained with DAPI (white), showing

1811 expression in clusters of cells posterior to the brain that extend into the brain and out to the

1812 periphery.

1813

1814 **Supplementary Video 10 –** Maximum intensity projection from confocal z-stack of WISH

1815 of neuron 4 marker Smp_319480 (magenta) and ciliary plate marker *16 kDa calcium binding*

1816 *protein* (Smp_096390) (green), counterstained with DAPI (white), showing that the clusters

1817 of neuron 4 cells reach the periphery at the latitude between the second and third tiers of

1818 ciliary plates.

1819

1820 **Supplementary Video 11 -** Maximum intensity projection from confocal z-stack of WISH of

1821 neuron 5 marker Smp_319480 (cyan), counterstained with DAPI (white), showing expression

1822 in cells anterior to the brain and in and around the paired lateral glands and their secretory

1823 ducts.

1824

1825 **Supplementary Video 12 -** Maximum intensity projection from confocal z-stack of WISH of

1826 protonephridia marker *ShKT-domain containing protein* (Smp_335600) (magenta),

1827 counterstained with DAPI (white). This shows the transcripts in a bilaterally symmetrical

1828 pattern distributed along the s-shaped path of an excretory tubule that connects the anterior

1829 and posterior flame cells.

1830

1831   **Supplementary Video 13 -** Confocal z-stack of WISH of markers for parenchyma 1 and 2

1832   (Smp_318890) (green) and stem cells *ago2-1* (Smp_179320) (magenta), counterstained with

1833   DAPI (white), shows the long cytoplasmic protrusions of the parenchymal cells reaching

1834   between the stem and other cells.

1835

1836   **Supplementary Video 14 -** Maximum intensity projection from confocal z-stack of WISH of

1837   marker for parenchyma 1 and 2 (Smp_318890) (green) counterstained with DAPI (white),

1838   shows the distribution of the parenchymal cells and their reach in the intercellular space of

1839   the larva.

1840

1841   **Supplementary Video 15 -** Confocal z-stack of WISH of markers for tegument *Meg6*

1842   (Smp_163710) (magenta) and *dynein light chain* (Smp_200180) (cyan), counterstained with

1843   phalloidin (green) and DAPI (white), shows the nuclei sit below the body wall muscle, and

1844   cytoplasmic protrusions reach between muscle filaments and form the epidermal ridges

1845   between the ciliary plates.

1846

1847   **Supplementary Video 16 -** Confocal z-stack of WISH of markers for stem clusters Delta/Phi

1848   *p53* (Smp_139530) (cyan), Kappa *UPPA* (Smp_308145) (green) and stem E

1849   *Phosphomevalonate kinase* (Smp_072460) (magenta), counterstained with DAPI (white),

1850   shows there are no cells that are *pmk*-positive but negative for *UPPA* and *p53*. Stem E cells

1851   cannot be validated *in situ*.

1852

1853   **Supplementary Video 17 -** Confocal z-stack of WISH of markers for stem clusters Delta/Phi

1854   *p53* (Smp_139530) (cyan), Kappa *UPPA* (Smp_308145) (magenta) and the pan-stem marker

1855   *ago2-1* (Smp_179320) (green), counterstained with DAPI (white), shows there are no cells

1856   that are *ago2-1* positive but negative for *UPPA* and *p53* that could be Stem F and G cells.

1857

1858   **Supplementary Video 18 -** Confocal z-stack of multiplexed WISH of tissue markers for

1859   stem *ago2-1* (Smp_179320) (green), muscle *paramyosin* (Smp_085540) (cyan) and nerve

1860   *complexin* (Smp_050220)(magenta).

1861

1862

1863 **Supplementary Video 19 -** Confocal z-stack of multiplexed WISH of tissue markers for

1864 parenchyma (Smp_318890) (green), stem *ago2-1* (Smp_179320) (cyan) and tegument *Meg6*

1865 (Smp_163710) (magenta).

1866

1867 **Supplementary Video 20 -** Confocal z-stack of multiplexed WISH of tissue markers for

1868 tegument *Meg6* (Smp_163710) (green) and protonephridia *ShKT-domain containing protein*

1869 (Smp_335600) (magenta).

1870

1871

1872

1873

1874

A)

i

ii

20μm

B)

Schistosoma mansoni life cycle

adult

schistosomulum

egg

Vertebrate host

miracidium

cercaria

Freshwater

sporocysts

Mollusc host

miracidia

enzymatic digestion

FACS

4 rxns
10X Chromium

ISH validation

Identification of cluster discriminating marker genes

C)

Protonephridia

Stem D

Stem B

Neuron 5

Stem G

Neuron 3

Stem A

Muscle 2

Neuron 1

Stem F

Stem E

Neuron 4

Ciliary plate

Tegument

Stem C

Muscle 1

Parenchyma 1

Neuron 2

Parenchyma 2

UMAP_2

UMAP_1

D)

Percent Expressed
• 0
• 25
• 50
• 75
• 100

Average Expression
2
1
0
-1

A) Dot plot showing Average Expression and Percent Expressed across cell types.

Rows: Muscle 1, Muscle 2, Neuron 1, Neuron 2, Neuron 3, Neuron 4, Neuron 5, Ciliary plate, Tegument, Protonephridia, Parenchyma 1, Parenchyma 2, Stem A, Stem B, Stem C, Stem D, Stem E, Stem F, Stem G

Columns: Myosin light chain 1 (045220), Troponin t (179810), *Paramyosin (PRM) (085540)*, Tropomyosin-1 (340760), NA (347710), Collagen alpha chain (159600), Titin (126240), *Kunitz-type protease inhibitor (052230)*, *Protein Wnt (156540)*, Notum (015820), PDZ domain protein (311170), *Ca-activated K channel (156150)*, *Myogenic factor MyoD (167400)*, Ca-transporting ATPase (136710)

Average Expression: 2.5, 2.0, 1.5, 1.0, 0.5, 0.0
Percent Expressed: 0, 25, 50, 75

Bi) / **ii)** Pan muscle *paramyosin (Smp_085540)* DAPI Phalloidin — 20µm

Ci) / **ii)** Pan muscle *paramyosin (Smp_085540)* DAPI Phalloidin

Di) / **ii)** Muscle 1 *protease inhibitor (Smp_052230)* *Wnt (Smp_156540)* — 10µm

Ei) / **ii)** / **iii)** Muscle 1 *notum (Smp_015820)* *Wnt (Smp_156540)*

Fi) / **ii)** Muscle 2 *Ca-activated K channel (Smp_156150)* *MyoD (Smp_167400)* — 5µm

A) clustering of cells

Kappa-like
Stem A
Stem B
Stem G
Tegument
Stem
Stem F
Stem E
Tegument
Stem C
Stem D
Delta/Phi-like

○ *sporocyst*
● miracidia

B)

C) latent time

1
0

D) Smp_139530 *p53-1*

unspliced
0.8
0.4
0.0
0   3   6
spliced

velocity

1
0
-1

expression

4
2
0

E) Smp-145470 *zfp-1*

unspliced
0.16
0.08
0.00
0   8
spliced

velocity

0.8
0
-0.8

expression

4
2
0

F) Smp-136160 *p53-2*

unspliced
1
0
0   4
spliced

velocity

0.3
0
-0.3

expression

2
1
0

G) Smp-041540 *eled*

unspliced
0.04
0.02
0.00
-0.02
-0.04
0.00   0.02   0.04
spliced

velocity

0.08
0
-0.08

expression

0.012
0.006
0.000

Muscle (19%)
Nerve (57%)
Tegument (13%)
Protonephridia (2%)
Parenchyma (2%)
Stem (7%)

20μm

Ci *ago2-1 (Smp_179320)*

20μm

Di *p53 (Smp_139530)* stem 1 Delta/Phi
UPPA (Smp_308140) stem 2 Kappa