1

Unpacking the functional heterogeneity of the Emotional Face Matching Task:

a normative modelling approach

Hannah S. Savage[1,2], Peter C. R. Mulders [1,3], Philip F. P. van Eijndhoven[1,3], Jasper van Oort[1,3], Indira Tendolkar[1,3], Janna N. Vrijsen[1,3,4], Christian F. Beckmann[1,2,5], Andre F. Marquand[1,2]

[1]Donders Institute of Brain, Cognition and Behaviour, Radboud University, Nijmegen, The Netherlands

[2]Department of Cognitive Neuroscience, Radboud University Medical Centre, Nijmegen, The Netherlands

[3]Department of Psychiatry, Radboud University Medical Centre, Nijmegen, The Netherlands

[4]Depression Expertise Centre, Pro Persona Mental Health Care, Nijmegen, The Netherlands

[5]Centre for Functional MRI of the Brain (FMRIB), Nuffield Department of Clinical Neurosciences, Wellcome Centre for Integrative Neuroimaging, University of Oxford, Oxford, UK

Word count main text: 4745

Figures: 6

Tables: 1

Corresponding author:

Hannah S. Savage | Andre F. Marquand

Kappitelsweg 29,

Nijmegen, 6525EN

The Netherlands

E: hannah.savage@donders.ru.nl | andre.marquand@donders.ru.nl

Author contribution statement:

H.S.S – conceptualization, formal analysis, data curation, writing – original draft, writing – review & editing, visualisation. P.C.R.M – formal analysis, data curation, resources, writing – review & editing. P.F.P.E – resources, writing – review & editing, funding acquisition. J.O. – investigation, data curation, writing – review & editing. I.T. – resources, writing – review & editing. J.N.V. – resources, writing – review & editing. C.F.B. – resources, funding acquisition, writing – review & editing. A.F.M – conceptualization, supervision, funding acquisition, formal analysis, writing – original draft, writing – review & editing.

Abstract:

Functional neuroimaging has contributed substantially to understanding brain function but is dominated by group analyses that index only a fraction of the variation in these data. It is increasingly clear that parsing the underlying heterogeneity is crucial to understand individual differences and the impact of different task manipulations. We estimate large-scale (N=7641) normative models of task-evoked activation during the Emotional Face Matching Task, which enables us to bind heterogeneous datasets to a common reference and dissect heterogeneity underlying group-level analyses. We apply this model to a heterogenous patient cohort, to map individual differences between patients with one or more mental health diagnoses relative to the reference cohort and determine multivariate associations with transdiagnostic symptom domains. For the face>shapes contrast, patients have a higher frequency of extreme deviations with unique spatial distributions depending on diagnosis. In contrast, normative models for faces>baseline have greater predictive value for individuals' transdiagnostic functioning.

Introduction:

Task-based functional neuroimaging (functional magnetic resonance imaging; fMRI) has been widely applied in foundational and clinical neuropsychology to characterise neural processes that underpin a behaviour or process of interest. The typical approach in such studies is based on comparing mean differences in the magnitude and location of activation (measured by changes in BOLD signal), which has helped us to understand how these processes may differ between groups defined by biological and sociocultural factors, psychopathologies, or therapeutic interventions. The majority of prior research has reported group-level summary statistics, which inform us of those regions most consistently activated across participants/groups during task conditions. This method assumes that the neural mechanisms facilitating the process of interest are consistent across individuals within and between groups. This assumption enables our understanding to reach only so far as 'the average brain' of an 'average control', or 'average patient'.

In order to better understand how the brain relates to behaviour it is essential to move our focus from the group-level to studying individual differences and consider the neural activation of these processes within the context of multiple sources of heterogeneity. For example: (i) natural variation within the general population, including potentially heterogenous yet functionally convergent processes, and (ii) heterogeneity within groups of interest, such as within mental health diagnoses. Furthermore, when comparing between independent studies, the influence of task design (i.e. small modifications to an original task) and acquisition parameters should also be considered but are seldom investigated.

One approach that can provide insight into individual differences is normative modelling[1,2]. The normative modelling framework provides statistical inference at the level of each subject with respect to an expected pattern across the population, highlighting variation within populations in terms of the mapping between biological variables and other measures of interest. This framework has previously been employed by our group and others to dissect structural variation within large healthy populations[3] and clinical psychiatric populations (e.g. in autism [4-6], schizophrenia and bipolar disorder[7]), and in relation to dimensions of psychopathology[8]. Applying this method to task-based fMRI data we will be able to characterize how functional activity within each voxel or ROI in the brain differs between individuals, and hence show with greater nuance the range of task-evoked activation within the general population[2]. Further, applying this model to patients with a current diagnosis (mood and anxiety disorders, autism spectrum disorders (ASD) and/or attention deficit hyperactivity disorder (ADHD)) we will be able to map differences in these individual participants with respect to the reference cohort. This may reveal unique clusters of deviation patterns, within and/or across diagnostic categories.

In this study, we use the Emotional Face Matching Task (EFMT) to demonstrate the potential of the normative modelling method to identify individual differences in task-based fMRI. The EFMT, also commonly referred to as the 'Hariri task', has been used in over 250 fMRI studies since it was most notably introduced in 2002[9,10]. This task asks participants to match one of two images that are simultaneously

presented at the bottom of the screen, to a third target image displayed at the top of the screen; participants
match images of facial configurations consistent with the common view of prototypic facial expressions,
most frequently of fear or anger, or similarly positioned geometric shapes. Matching faces, as compared to
matching shapes, evokes explicit and/or implicit emotional face processing, which has previously been
shown to engage the amygdala, fusiform face area, anterior insula cortex, the pregenual and dorsal anterior
cingulate cortex, the dorsomedial and dorsolateral prefrontal cortex, and visual input areas. Previous work
has related activity to biological and demographic variables, and compared between many different clinical
groups and developmental spectrums.

Due to its experimental simplicity and focus on subcortical circuitry relevant to brain disorders, the
EFMT has been implemented in a number of large-scale neuroimaging initiatives including the UK
Biobank[11], the Human Connectome Project (HCP)[12,13], HCP Development[14], the Amsterdam Open MRI
Collection Population Imaging of Psychology (AOMIC PIOP2)[15], and the Duke Neurogenetics Study (DNS).
We take advantage of these large open-access/shared datasets to collate a large representative sample of
over 7500 participants from six sites to first (1) build reference normative models that highlight the natural
variation of functional activity evoked by the EFTM [as measured by the task contrasts faces > shapes and
faces>baseline], and (2) determine how the model's prediction relates to age, sex, and variations in task
design. We then apply these models to over 200 participants with a current mental health condition or who
are neurodivergent from the MIND-Set cohort (Measuring Integrated Novel Dimensions in
neurodevelopmental and stress-related psychiatric disorder)[16], to (3) map deviations in patients with a
current diagnosis (mood and anxiety disorders, ASD and/or ADHD) relative from the reference cohort, both
at the group level and at the level of the individual. We show that despite the ostensible simplicity of this
task and robust group effects, there is considerable inter-individual heterogeneity in the nature of the elicited
activation patterns and that such differences are both highly interpretable and predict cross-domain
symptomatology in a naturalistic clinical cohort.

Results:
Group level comparisons show consistent effects across cohorts
First, we performed a classical group comparison to provide a reference against which to understand the
inter-individual differences in subsequent analyses. To achieve this, we randomly selected 100 random
individuals' FSL pre-processed data into fixed-effects general linear models to create group level maps for
the faces>shapes (Fig. 1a) and faces>baseline (Fig. 1b) contrasts (see methods). This also served as a
sanity check to ensure the data was comparable to past literature. Overall, positive task effects (activations)
for faces>shapes were found in the bilateral inferior and middle occipital lobe and the calcarine cortex (V1)
extending anterior-ventrally to the bilateral lingual and fusiform gyrus, and anterior-dorsally to the middle
and inferior temporal gyrus; the bilateral amygdala extending into the hippocampus; the bilateral temporal
pole; a dorsal region of the vmPFC; and the bilateral middle and inferior frontal gyrus. Task-related

deactivations were found across regions comprising the default mode network, including the anterior and
posterior cingulate cortex and precuneus, the precentral gyrus and supplementary motor area and the
inferior temporal lobe.

Figure 1: Task evoked activation. Two representative groups maps (from HCP Young Adult and UK Biobank), illustrating regions where participants show greater BOLD signal (z-statistic maps, thresholded at>±2.6) to (a) faces, as compare to shapes (faces>shapes), and (b) faces, as compared to baseline (faces>baseline). x,y,z = -4,-6,-16.

Fitting reference normative models for emotional face processing

Next, we estimated normative models of EFMT-related BOLD activation for the face>shapes and
faces>baseline contrast using data from 7641 individuals across the lifespan. To achieve this, we split the
data into training (n = 3877) and test splits, stratified by site (n = 3764), then fit a Bayesian Linear
Regression model that predicted the single subject level activation for each voxel of the brain, as a function
of sex, age, and acquisition and task parameters (see methods). Explained variance in the test set was
good, especially in regions that showed activation at the group level (Fig. 1) including the occipital
lobe/visual cortex and the bilateral amygdala (faces>shapes: Fig. 2a; faces>baseline: Fig. 2d). As shown
in Supplementary Fig. 2a and 2c in most voxels the skew and kurtosis was acceptable (i.e. -1 < skew < 1
and kurtosis around zero). For a very small proportion of voxels this was not the case; the most ventral
region of the vmPFC (i.e. the bottom border of the brain) was the most negatively skewed which we interpret
to reflect the varying degrees of signal dropout, more so than biological variation. The few voxels with
positive kurtosis were spatially overlapping with regions that were negatively skewed, which likely reflects
the extended negative tails of the distributions in these voxels.

Figure 2: Evaluation and deviation scores from the faces>shapes (left) and faces>baseline (right) normative models.
Explained variance is high in the normative models, irrespective of whether they are built using the face>shapes contrast (a), or the faces>baseline contrast (x,y,z = -4,-6,-16). (d). Histograms show the relative frequency of the total number of deviations that a participant has for each model (b,e). Normative Probability Maps illustrate the percentage of participants of the total sample who had positive (hot colours) or negative deviations (cool colours) > ±2.6 within each voxel, for the faces > shapes (c) and faces>baseline (f) models. x,y,z = -5,6,-15.

Voxel-wise deviations show considerable inter-individual variability

We then used these normative models to quantify the degree of inter-individual variability. To achieve this,

for each participant we created a thresholded normative probability map (NPM; deviation scores >±2.6)

which indicates the difference between the predicted activation and true activation scaled by the prediction

variance, and therefore shows the voxels where that participant had greater or less activation than would

be expected by the normative models. Figures 2b and 2e show the frequency of the total number of

deviations that individuals had from the faces>shapes, and faces>baseline models, respectively. Within

each voxel, we then counted how many participants had positive or negative deviations (>±2.6). The

resulting brain maps illustrate the variability in the magnitude of functional activation per voxel, across the

population for the two task contrasts (Fig. 2c + f). This shows that: (i) there is considerable inter-individual

variability underlying the mean effects and (ii) that the spatial distribution of individual deviations mostly

occurs within the task network. Every voxel of the brain had at least one subject with a deviation >±2.6 (not

shown), although, as illustrated, there were regions including the medial occipital lobe extending to the

bilateral fusiform gyrus and inferior temporal lobe, the bilateral inferior frontal gyrus extending to the

precentral gyrus, and the posterior region of the vmPFC, wherein deviations were more frequently

observed. As there were minimal differences in the evaluation metrics between models built using either

contrast, and as the contrast faces>shapes is most commonly reported in prior literature, we use this as

our primary contrast for our further analysis of the reference model.

Separable effects of input variables on model predictions

Next, we examine structure coefficients from our models to disentangle the effects of different input

variables. As shown in Figure 3, the direction of the relationship between input variables and the predicted

BOLD activation, and the fraction of the explained variability can be meaningfully separated for

interpretation. Some input variables, namely acquisition parameters, showed overlapping effects (with

sensical direction flips) likely due to their relatively high correlation and limited variability across sites. For

example, the number of target blocks, volumes acquired (not shown), use of multiband sequence (not

shown), and the length of the TR (not shown) all showed a similar relation to predicted activity.

Increased age was related to decreased predicted activity across the peripheral/surface of the

brain, as well as regions surrounding the ventricles, and increased activity in midline regions of the default

mode network, the bilateral insula, the fusiform face area extending to the para-hippocampal gyrus and the

superior temporal gyrus. Predictions were only minimally influenced by sex, and the spatial mapping of this

relationship was broadly overlapping with that of intra-cranial volume (not shown).

We further illustrate the ability of this method to disentangle the influence of task design choices,

on predicted activation. For example, the influence of the matching rule and the stimuli presented. Being

told to match the emotional expression, as compared to matching the faces, related to increased predicted

BOLD activity within subcortical areas including the bilateral putamen, caudate body and medio-dorsal

thalamus. Attending to the emotional expression also predicted increased activity within the superior frontal
gyrus extending to the supplementary motor area, the posterior medial temporal gyrus the inferior temporal
gyrus, and the medial temporal pole. Conversely, when participants were asked to match faces, the model
predicted greater activation within the bilateral fusiform gyrus, the middle temporal gyrus, the superior
temporal pole, the dorsolateral prefrontal cortex, and a large area of the inferior parietal gyrus extending to
the supramarginal and angular gyrus. Additionally, when stimuli from the Ekman series were used the
model predicted greater activation within the bilateral inferior occipital gyrus and the calcarine cortex (V1),
the bilateral lingual and fusiform gyrus extending to the inferior temporal gyrus, as well as in the medial
cingulate cortex, an anterior region of the vmPFC, the superior medial prefrontal cortex, and subcortical
regions including the ventral posterior thalamus, the posterior putamen, para-hippocampus, hippocampus
and amygdala. Conversely, the use of the Nim-Stim Set stimuli related to greater activity within default
mode regions, including a large area of the ventromedial/medial prefrontal cortex, precuneus, cuneus, as
well as the supramarginal gyrus which extended medially to the anterior and posterior insula, which in turn
extended laterally to the superior and medial temporal gyri.

Figure 3: The relationship between input variables and the predicted BOLD activation for faces>shapes. Maps show the
correlation coefficients (rho) thresholded by their respective coefficients of determination (rho²>0.3) of selected model input variables.
This can be interpreted as showing how predicted BOLD activation for the faces>shapes contrast relates to the input variables of the
normative models. Positive correlations (warm colours) indicate greater activation for higher values of the input variable and negative
correlations (cool colours) greater activation for lower values of the input variable (note that some variables are dummy coded, e.g.
target stimuli, instructions) x,y,z = -4,-6,-16.
208
209

 A traditional case-control comparison identifies few differences between patients and
controls

We then performed a voxel-wise case-control comparison on the raw data in order to test for group level
differences between a heterogeneous patient cohort and matched controls from the naturalistic MIND-Set
sample. As evidenced in Table 1 (see Diagnoses), the naturalistic MIND-Set sample has many patients
with co-occurring and heterogenous mental health diagnosis, with or without neurodivergence, and is
therefore representative of diverse clinical populations. This analysis revealed very few differences between
the patient cohort, and unaffected controls for faces>shapes and faces>baseline (Fig. 4a and b – bottom
rows). More specifically, comparing patients' task activation (Fig. 4a – top row) to controls (Fig. 4a – middle
row) for the faces>shapes contrast showed patients had greater activation in the left temporal medial gyrus
and bilateral posterior cingulate cortex, as well as in small regions of the supplementary motor area, and
the genus of the anterior cingulate cortex (Fig. 4a – bottom row). There were negligible differences between
patients and controls for the faces>baseline contrast (Fig. 4b – bottom row).

224
225 ***Figure 4: General linear model results comparing patients to controls for the faces>shapes and faces>baseline contrasts.***
226 Maps show regions activated (warm colours) and deactivated (cool colours) for faces>shapes (a) and faces>baseline (b), for
227 patients (top row) and unaffected controls (middle row) from the MIND Set cohort. (c) Regions where patients have more activation
228 than controls (bottom row) (z-statistic maps, thresholded at > ±2.6). x,y,z = -14,-13,-9.
229

Application of normative model to a naturalistic clinical sample
Next, we aimed to relate the deviations from these normative models to psychopathology. To achieve this,
we evaluated the patient cohort with respect to the normative models estimated from the large reference
cohort. For the faces>shapes and faces>baseline models, the explained variance of the clinical test data
was quite low. This was expected given that this cohort is quite homogenous with respect to the covariates

included in the model (i.e., all subjects were scanned on the same scanner, using the same experimental
paradigm and had an age range considerably narrower than the reference cohort). This suggests that the
variance in BOLD signal was driven more by individual differences, as opposed to the variables included in
the model. The skew and kurtosis of the models were centred around zero. See Supplementary Figure 4a-
f for histograms of these evaluation metrics, and their respective illustration on the brain.

Frequency of deviations differentiates patients from unaffected controls
Next, we compared the frequency of extreme deviations (NPMs thesholded at > ±2.6), at the level of each
individual, between patients from the MIND-Set cohort and unaffected controls. For each model type
(faces>shapes: Fig. 5b,c; faces>baseline: Fig. 5e,f). MIND-Set patients had a greater frequency of
deviations relative to the reference cohort for the faces>shapes contrast (Mann-Whitney U test = 358986.5,
p = 1.55^{-8} ; Fig. 5b). These deviations were most frequently identified in the lateral ventral prefrontal cortex,
and the bilateral medial and inferior temporal lobe (Fig. 5a). In contrast, for the faces>baseline contrast
individuals from the reference cohort had a greater frequency of deviations relative to MIND-Set patients
(Mann-Whitney U test = 509017.0, p = 0.0007; Fig. 5e). For this contrast, these deviations are, however,
strongly localised within the most ventral region of the vmPFC (Fig. 5d) which is well-known to be
problematic area for signal distortion artefacts in fMRI, therefore we do not interpret this difference as being
biologically meaningful.

Associations of patterns of deviation with cross-diagnostic symptom domains
We then aimed to determine whether multivariate patterns of deviation from the reference models were
associated with cross-diagnostic symptomatology. To achieve this, we input whole-brain (unthresholded)
deviation maps and factor loadings for negative valence, cognitive function, social processes and
arousal/inhibition domains from prior work[17] to an established penalised canonical correlation analysis
(CCA) framework that enforces sparsity (sparce CCA, SCCA; functional domain loading scores were
available for 217 patients)[18,19]. Significant out of sample associations (10 fold 70% - 30% training- test split)
were detected both for faces>shapes and faces>baseline contrasts (mean r of test splits 0.133 and 0.211
respectively, both p < 0.001 by permutation test; Fig. 6b, e) but with distinct patterns of effects both in terms
of symptom domains and associated brain regions. More specifically, for the faces>shapes contrast,
decreased functioning predominantly in the negative valence and arousal/inhibition domains (Fig. 6a) was
associated with a pattern of deviations including the right insula, the bilateral medial prefrontal cortex and
pre- and post- central gyri, the bilateral inferior temporal gyrus, lingual gyrus, bilateral hippocampus and
the right thalamus, as well as the regions in the medial and left lateral cerebellum (Fig. 6c). By comparison,
for the faces>baseline contrast factor loadings for cognitive functioning and arousal/inhibition (Fig. 6d) were
most strongly related to a pattern comprising bilateral insula, the anterior-to-medial cingulate cortex
extending to the dorsal medial prefrontal cortex, the pre- and post- central gyri, the right middle frontal and
bilateral inferior frontal gyrus, and the bilateral hippocampus, caudate, putamen and amygdala, and the

medial and left lateral cerebellum (Fig. 6f). The SCCA was repeated to relate participant's diagnoses with
their whole-brain (unthresholded) deviation maps. In contrast to the cross-diagnostic symptom domains,
there was no association between diagnostic labels and deviation scores. Mean canonical correlations were
small (mean r of test splits <0.09 for both faces>shapes and faces>baseline models), and this was not
statistically significant as determined by 1000-fold permutation testing.
278
279

281 ***Figure 5: Testing the faces>shapes (left) and faces>baseline normative models with the MIND-Set cohort.*** Normative
282 Probability Maps illustrate the percentage of participants of the clinical sample who had positive (hot colours) or negative deviations
283 (cool colours) > ±2.6 within each voxel, for the faces>shapes (a) and faces>baseline (d) models. Histograms and box plots show the
284 relative frequency and mean number of the total deviations that a participant has for faces>shapes (b,c) and faces>baseline (e,f)
285 models. Positive: x,y,z = 9,-16,-16, Negative: x.y,z = 5,-29,-24.

286
Figure 6: Sparse canonical correlation analyses (SCCA) between functional domains, and deviation scores from
faces>shapes or faces>baseline normative models. Weights per factor to latent variable of psycho-social functioning (a,d).
Canonical correlation between 4 functional domains and whole-brain deviation scores from (b) faces>shapes and (e) faces>baseline
normative models (regularisation 10%). Mean voxel-wise weights to latent variable of deviation scores from (c) faces>shapes
normative models and from (f) faces>baseline. All results are statistically significant with 1000-fold permutation tests (*** = p<0.001).
x,y,z, = [-42,-17,8,33], [29,4,-21,-46,-71], [47,22,-3,-28].
293
294
295
Spatial extent of deviations highlights similarities across, and differences between diagnoses
Finally, we were interested in mapping the spatial distribution of the deviations within the clinical sample,
and whether this varied according to the participant's mental health diagnosis or neurodivergence (note
that subjects can be in multiple categories; Sup Fig. 5). For each diagnosis, the pattern of deviations was
highly heterogeneous, providing further support for high degree of inter-individual heterogeneity we have
reported previously for mental disorders[4,5,7], and underlining the need to move beyond case-control
comparisons at the level of diagnostic groups.
303
304

Discussion:

In this study we made use of six large publicly available datasets of participants completing the fMRI EFMT to build a reference normative model of functional activation underlying emotional face processing. We collated data from over 7500 participants and show that our voxel-wise models can explain up to 50% of variance in observed BOLD signal, with the remaining unexplained variance representative of individual differences in functional activation (deviation scores). We unpacked the variance explained by the models, to show how the predicted activation related to the models' input variables, namely demographics, variations in task design, and acquisition parameters. Lastly, we tested our reference model with data from a sample of patients with heterogenous and frequently co-occurring psychiatric conditions (mood and anxiety disorders, and neurodevelopmental conditions). Our analyses show that: (i) there is considerable inter-individual variation superimposed on the group effects customarily reported in fMRI studies, (ii) that such variation is predictive of psychiatric symptom domains in a cross-diagnostic fashion and (iii) while an overall effect of diagnosis was evident, this was highly individualised in that the overlap of deviations amongst individuals with the same diagnosis was low. This implies that there are brain regions wherein patients more frequently have deviations irrespective of the type of diagnoses, and other regions wherein the frequency of deviations appears specific to the mental-health condition or neurodivergent diagnosis.

A key feature of the normative modelling framework in the context of multi-site fMRI data is that it allows us to aggregate data across multiple samples by binding them to a common reference model. This provides multiple benefits: it removes site effects from the data without requiring the data to be harmonized[20], which avoids the introduction of certain biases due to harmonisation[21] and allows meaningful comparisons to be drawn across studies. For example, this allows aggregation of different studies to better understand variation across cohorts or across the lifespan and to understand the effect of different task parameters on functional activity across cohorts. Moreover, by placing each individual within the same reference model this provides the ability to quantify, compare and ultimately parse heterogeneity across studies.

Traditional group-level task contrasts, as shown in Fig. 1, inform us of the region's most consistently activated across participants/groups during task conditions. Their interpretation has relied heavily on the assumption of spatial homogeneity of activation between subjects; an assumption that the deviation scores from our reference model show to be largely untrue (Fig. 2). We show that such group effects reflect a small proportion of the variation amongst individuals and using the normative modelling framework we map the underlying heterogeneity, separating variation in the intensity and spatial extent of task-evoked functional activation between-subjects attributable to known factors such as site effects, demographics, acquisition parameters, and differences in EFMT paradigm design. More importantly, we show that residual differences in the neuronal effects elicited by the task are highly meaningful in that they were predictive of psychiatric

symptomatology and can be used to understand inter-individual differences in functional anatomy and its
relation to clinical variables. In our test reference population, while every voxel of the brain had at least one
participant with a large deviation, some regions considered active during the faces condition (as compared
to shapes), such as the medial occipital lobe, fusiform gyrus and inferior temporal lobe, were also regions
in which positive deviations were frequently observed.

When building our reference models, we chose to include and control for multiple variables that we
reasoned may influence the BOLD signal observed. These included demographic factors such as age and
sex, and task design choices that could influence the BOLD signal generated, as well as acquisition
parameters that could influence the BOLD signal recorded. Some effects, such as that of age and task
instructions were relatively strong and interpretable, for example, increased age predicted decreased
activity in surface areas of the brain and regions surrounding the ventricles likely reflecting decreased signal
due to age-related atrophy, and instructing participants to match emotional expressions, as opposed to
matching faces, increased the predicted activity in the thalamus which may reflect increased engagement
of regions associated with affective processing. On the other hand, other variables explained relatively little
variance in the predictions (e.g. sex). In our sample many predictor variables were collinear across sites
which limited our ability to detect systematic differences resulting for example from differences in the task
paradigm. In this work we decided to keep all variables in the model and used structure coefficients to
identify the importance of different variables, which are relatively insensitive to collinearity. This follows prior
work to identify specific effects of input variables on model predictions, for example the influence of specific
adversity types on predicted morphometric changes (Holz et al., *in prep*). However future researchers may
consider reducing the dimensionality of their inputs prior to model construction. Future studies with larger
numbers of more diverse samples (e.g. more variations on the basic task design), as is possible in
consortium such as ENIGMA, will allow for more fine grained analyses of the effect of task parameters on
inter-individual variation within the population.

We demonstrated that distinct patterns of deviations, derived from each model type (faces>shapes or
faces>baseline), were associated with unique profiles of functioning across four transdiagnostic domains.
The distinct patterns of effects, in terms of the implicated symptom domains and associated brain regions,
make sense in the context of relevant existing literature. For example, negative affect, impulsivity and
emotional liability have previously been related to functional activity within the bilateral insula, motor cortex
and hippocampus[22], and cognitive functioning has been linked to activity within the medial prefrontal cortex,
anterior-to-medial cingulate cortex, superior frontal gyrus. This not only validates the interpretability of
findings from these normative modelling analyse, but also illustrates the potential for future researchers to
use individualised deviation maps to better understand the neural processes that underly cognitive and
affective functioning, within and across diagnostic boundaries. Furthermore, approaching dysfunction
through the normative modelling framework and transdiagnostic functional domains appears to more

closely relate to underlying biology. This reflects practitioners implementation of clinical care and the use
of overlapping treatments for differing disorders, which often does not fit a binary classification paradigm.
Using this modelling approach may also better allow for the quantification of neurodivergence, not as being
'disordered' but rather as varying phenotypic expressions along a characterised spectrum.

It should be noted, however, that within any one voxel of the brain, only ~20% of the clinical sample
(be that in the total sample, or within disorders) had large deviations. This suggests that the exact location
of deviations is very variable between individuals, and could explain why many prior studies have not found
significant differences when performing traditional case-control analyses. In this study, we aimed to
estimate the degree to which the deviations from the normative models were associated with cross-
diagnostic symptomatology, but other approaches may also be useful, as outlined in Rutherford, et al. [23].
For example, clustering algorithms could be applied to derive a stratification of individuals[4,5] or to identify
heterogenous yet convergent functional processes (many-to-one functional mappings)[24], and supervised
learning methods may be useful to assess the degree to which specific clinical variables can be predicted
from the patterns of deviation we report.

Interestingly, the normative models of functional activation built using the faces>shapes have a different
pattern of association with symptomatology relative to the faces>baseline contrast. This suggests that the
two contrasts carry complementary information about psychopathology. The frequency of deviation scores
was significantly greater in the clinical cohort, compared to the reference cohort, when using the
faces>shapes contrast, and the weights attributed to each of these deviations (at a voxel-level) in the SCCA
were associated with different symptom domains. Neither contrast was significantly predictive of diagnosis.
By comparison, the relationship between the frequency of deviation scores and domains of function was
stronger when using models built using the faces>baseline contrast, which was further supported by the
stronger canonical correlation between factor loadings for functional domains and deviations from the
faces>baseline models. Taken together, this could be interpreted to suggest that widespread deviations,
best captured by the faces>shapes contrast, are indicative of global alterations in functioning which are
broadly linked to different clinical diagnoses. By comparison, fewer but more focal deviations and/or the
ability to detect abnormal baselines of activation[25,26], best revealed using the faces>baseline, have greater
relation to specific functional domains. Future researchers should carefully consider the task contrast used
to construct their normative models.

Concerns for the within-subject reliability of task-based fMRI data[27] are not to be dismissed in the context
of our models which are currently built on cross-sectional data. While we acknowledge the limitations
imposed due to the limits of test-retest reliability of task fMRI, our results nevertheless provide encouraging
evidence for the use of task fMRI readouts as individualised biomarkers as we show by their ability to predict

clinical variables in the context of SCCA. While we were not able to assess test-retest reliability directly within the context of normative models due to a lack of test-retest data for the EFMT, one great strength of the normative modelling approach is the shift from looking only at mean activity, to estimating the underlying variance. This can implicitly down-weight regions or individuals that are less reliable. Indeed, it could be expected that repeated sampling would fall within the same variance distribution, and as such our broad understanding of brain function remains quite stable. The normative modelling framework is also ideally positioned to directly test the reproducibility of fMRI within subjects. In follow-on work to the present manuscript, we are currently developing an extension to explicitly include test-retest variability in the model by testing reference models with repeat scans from participants, and compare individuals' deviation scores between the two tests, whilst explicitly quantifying within subject variance, such that it provides a lower bound on the size of deviation that can be considered meaningful (Bučková et. al., *in prep*). Alternatively, where multiple repeats are available, hierarchical models can be used to accommodate dependencies between subjects[20] which would provide more precise estimates of individual deviations. The application of the normative modelling method to fMRI can easily be generalised to other tasks (e.g. the monetary incentive delay incentive processing task or n-back work memory task) and need not stop at predicting functional activation. With the right data sets, this method could use fMRI data to predict many other variables including psychophysiological responses or subjective ratings of affect.

With this work, we show the potential for the normative modelling framework to be applied to large task-based fMRI data sets to bind heterogeneous datasets to a common reference model and enable meaningful comparisons between them. Using this approach, we illustrate the heterogeneous intensity and spatial location of task-evoked activation within the general population[2] using the EFMT in a sample of over 7500 participants. Further, we applied this model to patients with a current diagnosis (mood and anxiety disorders, ASD and/or ADHD) and demonstrate the transdiagnostic clinical relevance and further potential for deviation scores derived from this method. The potential of this method is clear; normative modelling of task-based functional activation can facilitate a better understanding of individual differences in complex brain-behaviour relationships, and further our understanding of how these differences relate to mental health and neurodivergence.

Methods:

Data sets: We collated a large reference sample from 6 independent sites for whom high quality fMRI data for the EFMT are available: AOMIC PIOP2, Duke Neurogenetics Study, HCP Development, HCP Young Adult (1200 release), UK Biobank, and the MIND-Set cohort which also includes a clinical population. For sample details per site see Table 1.

fMRI task paradigms: All sites collected a variant of the EFMT[9]. Although specific parameters varied, the overall design was consistent: in each face trial participants were presented with three images of human faces in a triangular formation. Participants were instructed to identify which of two faces/expressions presented at the bottom of the screen matched the one presented at the top of the screen by pushing a button with the index finger of their left or right hand. Multiple face trials were presented in one face block, and the task included multiple face blocks (see Table 1 for the number of trials per block, and number of blocks per site). As a somatomotor control, participants also completed shape trials, wherein they were presented with three geometric shapes (circles and ovals) and asked to indicate which of the two shapes presented at the bottom of the screen matched the one at the top. Multiple shape trials were concatenated to form one shape block, which were interspersed between face blocks.

Two paradigms (HCP Young Adult and HCP Development) included an inter-trial interval (white fixation cross on black screen), and three sites (HCP Young Adult, HCP Development and AOMIC PIOP2) had an instruction trial that preceded the start of each block. Tasks varied in their duration from 150 to 290 seconds, which indirectly corresponded to the acquisition of between 135 and 336 functional volumes.

fMRI data acquisition:

Site specific acquisition parameters per site are detailed in Table 1, and in the following site specific protocols: AOMIC PIOP2[15], HCP Young Adult[13], HCP Development[28], UKBiobank[29], Duke Neurogenetics Study (https://www.haririlab.com/methods/amygdala.html) and MIND-Set[30].

fMRI pre-processing: Data pre-processing was harmonised across all sites; a FSL-based pipeline[31] was consistently applied to decrease the likelihood of introducing variance due to pre-processing differences. Since the HCP young adult, HCP development and UKB Biobank data were already processed relatively consistently, we reused the processing pipelines provided by the respective consortia (for HCP sites we used the minimal processing pipeline)[29,32], with additional steps taken as necessary (e.g. matching smoothing kernels across studies). At a within-subject level, all functional data underwent gradient unwarping, motion correction, fieldmap-based EPI distortion correction (where fieldmaps were available), boundary-based registration of EPI to structural T1-weighted scan, denoising for secondary head motion-related artifacts using automatic noise selection, as implemented in ICA-AROMA[33], non-linear registration into MNI152 space, and grand-mean intensity normalization. All data were spatially smoothed using a 5 mm FWHM Gaussian kernel.

Quality control: Participants were excluded if their mean relative RMS was greater than 0.5mm. Additional quality control was performed for signal coverage in the prefrontal cortex for the UK Biobank sample (see supplementary methods).

fMRI general linear modelling (GLM) – single subject: We matched the methodological approach used to estimate the parameters within a GLM-based analysis, given evidence to suggest this analytic step can significantly contribute to the variability of reported results between sites[34]. Therefore, for each site, the linear model parameter were estimated using the FSL software package version 6.03 (HCP Young Adult, HCP Development, MIND-Set, Duke Neurogenetics Study; http://fsl.fmrib.ox.ac.uk/) or as downloadable form UK Biobank[29]. Two regressors were constructed from the faces and shapes blocks which were then convolved with a canonical double-gamma haemodynamic response function and combined with the temporal derivatives of each main regressor. These were treated as nuisance regressors and served to accommodate slight variations in slice timing or in the haemodynamic response. Data were pre-whitened using a version of FSL-FILM customized to accommodate surface data, the model and data were high-pass filtered (200s cut-off). Fixed-effects GLMs were estimated using FSL-FLAME 1: first for independent runs, then when necessary combining two runs into a single model for each participant (HCP Young Adult). and the AOMIC, DNS and MIND-Set maps were transformed into standard space using FNIRT[35]. We created summary group level maps per site (for a random sample of 100 participants), as a sanity check to ensure the data was otherwise comparable to past literature, and performed a case-control comparison between patients with a current diagnosis (mood and anxiety disorders, ASD and/or ADHD) and unaffected controls in the MIND-Set cohort.

Normative models: The z-statistic maps from the contrast face>shapes (5mm smoothed in standard space), for each subject, were used as response variables for the normative models. That is, we specified a functional relationship between a vector of covariates and responses. We created normative models of EFMT-related BOLD activation maps, as a function of sex, age, acquisition and task parameters (task duration (s), number of target blocks, instructions given to participants, the task stimuli), by training a Bayesian Linear Regression (BLR) model to predict BOLD signal for the faces>shapes contrast. Generalisability was assessed by using a half-split train-test sample (train: n = 3877, test: n = 3764). In preliminary analyses, we compared a warped model which can model non-Gaussianity with a vanilla Gaussian BLR model. Since the fit was comparable across most metrics and regions, we focus on the simpler Gaussian model below. We included dummy coded site-related variables as additional covariates of no-interest. We also created models to predict BOLD signal for the faces condition alone (i.e. face>baseline contrast). This was performed in the Predictive Clinical Neuroscience toolkit (PCNtoolkit) software v0.26 (https://pcntoolkit.readthedocs.io/en/latest) implemented in python 3.8.

Quantifying voxel-wise deviations from the reference normative model: To estimate a pattern of
regional deviations from typical brain function for each participant, we derived a normative probability map
(NPM) that quantifies the voxel-wise deviation from the normative model. The subject-specific *Z*-score
indicates the difference between the predicted activation and true activation scaled by the prediction
variance. We thresholded participant's NPM at Z = ±2.6 (i.e. p < .005)[7] using fslmaths and summed the
number of significantly deviating voxels for each participant, and then across the total sample.

Effects of input variables on model predictions: In order to probe the magnitude of the influence of task
design parameters on the predicted BOLD signal, we illustrated the structure coefficients (correlation
coefficients) of each task parameter-related variable (task duration (s), number of target blocks, instructions
given to participants, the task stimuli), as well as for age, sex and ICV. This approach is preferable to
regression coefficients when variables are collinear[36].

Clinical application: We tested the normative models we created using the reference data, with a
heterogeneous patient sample from the MIND-Set cohort (n = 236, mean age = 37. 1±13.27; 41.94%
female). This is a naturalistic and highly co-morbid sample derived from out-patients of the psychiatry
department of Radboud University Medical Centre. This included 150 patients diagnosed with a current
mood disorder (unipolar or bipolar depressive disorder), 12 with generalised anxiety disorder, 22 with social
phobia, 14 with panic disorder, 71 with attention-deficit-hyperactive-disorder, and 55 autistic individuals
(see Table 1 for full details and note that subjects can be in multiple diagnostic categories). The clinical
relevance of our models can also be tested in the context of transdiagnostic symptom domains; a
conceptualisation of mental functioning that transcends diagnostic boundaries and allows for nuanced
brain-behaviour interpretations. As such, for 217 (of our 236) patients for whom all required data was
available, we repeated a previously validated factor analysis method (performed in SPSS v24.0, oblique
rotation)[17] to obtain individual factor loadings on 4 functional domains: (1) negative valence, (2) cognitive
function, (3) social processes and (4) arousal/inhibition.

Quantifying patients' voxel-wise deviations from the reference normative model: As for the reference
cohort, we generated NPMs to estimate the pattern of regional deviations from typical brain function for
each participant, and summed across the sample. We then used a Mann-Whitney U test to compare the
frequency of deviations (>±2.6) between the reference controls and patients from the MIND-Set cohort.

Relating deviations to transdiagnostic functional domains: In order to map the association of the
deviation scores with cross-diagnostic symptomatology, we performed sparse canonical correlation
analyses (SCCA) to relate participant's scores in the four aforementioned functional domains or their
diagnoses, to their whole-brain (unthresholded) deviation maps using an established penalised CCA
framework that enforces sparsity[18,19]. Specifically, we applied variable shrinkage by adding an l_1-norm

penalty term to stabilise the CCA estimation and ensure the weights for the deviation scores were more

interpretable. We follow the formulation outlined in Witten, et al. [18], where we refer to for details. In brief,

given two data matrices X and Y with dimensions $n \times p$ and $n \times q$ respectively (here, these are the cross-

diagnostic factor loadings and whole-brain deviations), and two weight vectors u and v this involves

maximising the quantity $\rho = u^T X^T Y v$ subject to the constraints $\|u\|_2^2 \leq 1$, $\|v\|_2^2 \leq 1$, $\|u\|_1 \leq c_1$ and $\|v\|_1 \leq$

c_2, where the penalties $p(u)$ and $p(v)$ are the standard L1-norm. We set the regularisation parameters for

each view heuristically ($c_1 = 0.9p$ corresponding to light regularisation for the factor scores, $c_1 = 0.1q$,

corresponding to heavy regularisation for the deviation maps such that no more than 10% of voxels were

selected). While it is possible that better performance would be obtained by optimising the regularisation

parameters across a grid, we did not pursue that here due to the moderate sample size for the clinical

dataset. We assessed generalisability of SCCA by splitting the data in to 70% training data and 30% test

10 times. Finally, we wrapped the entire procedure in a permutation test where we randomly permuted the

rows of one of the matrices 1000 times to compute an empirical null distribution for significance testing.

Spatial patterns of deviations by primary and co-occurring diagnoses: We illustrated the spatial

heterogeneity in deviations between different diagnoses (note that subjects can be in multiple categories),

and further, compared patients with a single diagnoses to those with two, three, or more than three

diagnoses, to determine whether and if so, how the location of deviations related to the number of co-

occurring diagnoses a patient has.

571   Data availability:

572   Scripts for running the analysis and visualizations are available on GitHub (https://github.com/predictive-

573   clinical-neuroscience/EFMT_Norm_Models).

575   Acknowledgements:

The Duke Neurogenetics Study was supported by Duke University and National Institutes of Health (NIH)

grants R01DA031579 and R01DA033369. Ahmad R. Hariri (Duke University) is further supported by NIH

grant R01AG049789. The Duke Brain Imaging and Analysis Center's computing cluster, upon which all

DNS analyses heavily rely, was supported by the Office of the Director, NIH, under Award Number

S10OD021480. CFB gratefully acknowledges funding from the Wellcome Trust Collaborative Award in

Science 215573/Z/19/Z and the Netherlands Organization for Scientific Research Vici Grant No. 17854 and

NWO-CAS Grant No. 012-200-013.

Table 1: Sample details, functional scan acquisition parameters and Emotional Face Matching Task parameters for data included in the normative models.

Sample details				Functional scan acquisition parameters				Emotional Face Matching Task parameters									
Site	Sample size	Sex	Age (mean+ S.D) [range]	Scanner	TE/TR (ms)	Multi-band Factor	Flip angle	Matching Rule /Instructions	Target Stimulus	Trials per block/ Blocks/ Total Trials	Trial duration (s)	Instruction duration (s)	Inter-trial interval (s)	Block duration (s)	Task duration (s)	Inter-block-interval (s)	Volumes acquired
Human Connectome Project Young Adult	1044	561 F (53.7%)	28.76±3.70 [22-37]	3T Siemens Skyra	33.1/720	8	52	*Match faces*: Decide which of two faces presented on the bottom of the screen match the face at the top of the screen.	Angry and fearful faces: Nim-Stim Face Stimulus Set	6/3/18	2	3	1	21	156	NA	176
Human Connectome Project Development	201	110 F (54.7%)	13.86±3.83 [8-21]	3T Siemens Prisma	37/800												178
UK Biobank	5000	2487 F (49.7%)	63.99±7.45 [46- 82]	3T Siemens Skyra	39/735			*Match faces*: Indicate which face [or shape] on the bottom row matches the face on the top row.		NA/5/NA	NA	NA	NA		253	8	366
Amsterdam Open MRI Collection Population Imaging of Psychology	200	114 F (57.0%)	22.16±1.79 [18.25– 26.25]	3T Phillips Achieva dStream	28/2000	NA	76.1	*Match expression*: Match the emotional expression of the target face as quickly as possible.			when selected or up to 4.8s	10	5s – Reaction Time	~25	290	5	135
Duke Neurogenetics Study	1246	707 F (56.7%)	20.22±1.21 [18.09-23.07]	3T GE MR750	30/2000		NA	*Match faces*: Decide which of two faces presented on the bottom of the screen match the face at the top of the screen.	Angry, fearful, surprised, neutral faces: Ekman and Friesen, 1976	6/4/24	4	2	Faces: 2–6 (mean= 4) Shapes: 2	Faces: 48 Shapes: 36	390	NA	195
MIND-Set	Reference: 37/309 Clinical Test: 36/309	21 F (56.7%) 21 F (58.3%)	38.0±16.11 [20-74] 37.1±16.50 [20-70]	3T Siemens Magentom Prisma	34/1000	6	60	*Match expression*: Indicate which one of the bottom two faces matched the top face in terms of emotional expression.	Angry and fearful faces: Nim-Stim Face Stimulus Set	6/2/12	5	NA	NA	30	150	NA	150

Additional details of clinical samples:	Sample size	Sex	Age (mean+ S.D) [range]	Current diagnoses		Number of Diagnoses (% of total sample)
MIND-Set	236/309	99 F (41.9%)	37.1±13.27 [20-74]	150 Mood Disorder 71 Attention deficit hyperactivity disorder (ADHD) 55 Autism spectrum disorder (ASD) 22 Social Phobia 14 Panic Disorder	12 Generalised Anxiety Disorder 7 Anxiety disorder NOS* 6 Obsessive Compulsive Disorder 5 Post Traumatic Stress Disorder 4 Specific Phobia 2 Agoraphobia	**1:** 66 (27.9%) **2:** 65 (27.5%) **3:** 39 (16.5%) **>3:** 8 (3.38%)

Note: Underline indicates that this parameter was input as a variable in the normative models. *NOS (Not otherwise specified).

SUPPLEMENTARY METHODS – *Sample Details*:

Supplementary Figure 1: **Age and sex distributions** of (a) the total reference sample, (b) the total clinical test sample (MIND-Set) (c) the faces>shapes train (left) and test (right) split, and the (d) the faces>baseline train (left) and test (right) split.

SUPPLEMENTARY METHODS – *Signal coverage of the prefrontal cortex (PFC)*:

Due to air-tissue inhomogeneities which can diminish the acquired BOLD signal to such a degree that no activations are visible, a notorious effect within the ventral PFC, we performed targeted quality control for this extended region. Binary ROI masks were created for the dorsal ventro-medial PFC (d-vmPFC), ventral vmPFC (v-vmPFC), lateral vmPFC (l-vmPFC) and the dorso-medial PFC (dmPFC), as defined by the AAL2 atlas regions 25, 27, 33 and 1 respectively (see Supplementary Figure 1C). The percentage of voxels with an absolute value greater than 0 for the contrast faces > shapes within each ROI was determined (i.e. where any signal was present regardless of its relative direction; see Supplementary Figure 1A,B). While most sites had good coverage, the coverage within the ventral and lateral vmPFC regions were particularly variable for the UK Biobank data. We therefore performed this step *only on data from the UK Biobank site*; this selectivity was made possible by the large number of participants we had access to, and our need to include but a fraction of the total available sample. We ranked participants in descending order of the percent of their v-vmPFC, l-vmPFC, d-vmPFC, and dmPFC covered, respectively, and selected the first 5000 participants. We also collected the percentage covered value for a bilateral amygdala ROI mask, but made no exclusion/inclusions on this basis as coverage was very high across all participants and all sites.

Supplementary Table 1: Motion QC

Site	Full sample size	Included	Excluded
AOMIC	217	217	0
DNS	1263	1246	17
HCP Young Adult	1044	1044	0
HCP Development	256	256	0
UK Biobank	26167	5000	N/A
MIND-Set	393	389	4

Supplementary Table 2: vmPFC QC

ROI	Site	Sample size (n)	Mean percentage of ROI covered	Standard deviation
Bilateral Amygdala	AOMIC	217	100	0
	DNS	1246	100	0
	HCP Development	256	99.94936	0.162845
	HCP Young Adult	1044	99.98188	0.098589
	MIND Set	389	99.99966	0.006707
	UK Biobank	26120	99.47363	2.399575
dmPFC	AOMIC	217	99.10345	0.830591
	DNS	1246	99.61285	0.411399
	HCP Development	256	94.6524	2.460686
	HCP Young Adult	1044	96.85686	3.318827
	MIND Set	389	99.069	1.785034
	UK Biobank	26120	99.0493	1.405906
Dorsal vmPFC	AOMIC	217	85.08059	12.23881
	DNS	1246	94.68589	4.406996
	HCP Development	256	72.90154	16.11571
	HCP Young Adult	1044	98.58104	2.192929
	MIND Set	389	99.75218	0.886996
	UK Biobank	26120	94.50698	8.107753
Lateral vmPFC	AOMIC	217	98.90096	1.063159
	DNS	1246	99.82392	0.33135
	HCP Development	256	99.14376	1.003451
	HCP Young Adult	1044	99.46709	0.978404
	MIND Set	389	98.23297	1.745961
	UK Biobank	26120	89.56122	4.919904
Ventral vmPFC	AOMIC	217	96.06479	3.960334
	DNS	1246	99.45099	0.964661
	HCP Development	256	95.999	4.42593
	HCP Young Adult	1044	99.63665	0.85699
	MIND Set	389	98.83508	2.868568
	UK Biobank	26120	61.17452	15.9091

Supplementary Figure 2: vmPFC QC metrics. (a) Mean percentage of each ROI with signal greater than 0, used for quality control.
Error bars show +/- standard deviation (b) Stacked histograms (raw participant count) of the percentage of each ROI covered, coloured
by site.

SUPPLEMENTARY RESULTS – *Evaluation of reference normative models*:

Supplementary Figure 3: Evaluation of the faces>shapes (left) and faces>baseline (right) reference normative models.
Histograms show the skew (a,c), and kurtosis (b,d) of the normative models, and their respective illustration on the brain (x,y,z= 4,-6,-
15).
633

 SUPPLEMENTARY RESULTS – *Evaluation of normative models when applied to MIND-Set cohort*:

636
637 ***Supplementary Figure 4: Evaluation of the faces>shapes (left) and faces>baseline normative models when applied to MIND-***
638 ***Set cohort.*** Histograms show the explained variance (a,d), skew (b,e), and kurtosis (c,f) of the clinical data, as tested on reference
639 normative models of EFMT related BOLD activation, and their respective illustration on the brain (x,y,z= 4,-6,-15).

SUPPLEMENTARY RESULTS – *Location of deviations for diagnoses:*

644
645
646 ***Supplementary Figure 5: Heterogeneous location of deviations in predicted BOLD signal for different types of***
647 ***neurodivergence, and mental health diagnoses.*** Maps illustrate the percentage of participants with a neurodivergence or mental
648 health condition who had positive (left; hot colours) or negative deviations (right; cool colours) > ±2.6 within each voxel [minimum =
649 %5 of sample, or 1 participant where 5% was a participant count less than 1, maximum = 20% of disorder sample size]. x,y,z, = 5, -
650 28, -6.

SUPPLEMENTARY RESULTS – *Location of deviations for increasing levels of co-occurring*

diagnoses:

Supplementary Figure 6: Heterogeneous location of deviations in predicted BOLD signal for increasing levels of co-occurring diagnoses. Maps illustrate the percentage of participants with a neurodivergence or mental health condition who had positive (left; hot colours) or negative deviations (right; cool colours) > ±2.6 within each voxel [minimum = %5 of sample, or 1 participant where 5% was a participant count less than 1, maximum = 20% of sample size].

References:

1 Marquand, A. F. *et al.* Conceptualizing mental disorders as deviations from normative
functioning. *Molecular Psychiatry* **24**, 1415-1424, doi:10.1038/s41380-019-0441-1 (2019).
2 Marquand, A. F., Rezek, I., Buitelaar, J. & Beckmann, C. F. Understanding Heterogeneity in
Clinical Cohorts Using Normative Models: Beyond Case-Control Studies. *Biological psychiatry* **80**,
552-561, doi:10.1016/j.biopsych.2015.12.023 (2016).
3 Rutherford, S. *et al.* Charting brain growth and aging at high spatial precision. *eLife* **11**, e72904,
doi:10.7554/eLife.72904 (2022).
4 Zabihi, M. *et al.* Fractionating autism based on neuroanatomical normative modeling.
Translational Psychiatry **10**, 384, doi:10.1038/s41398-020-01057-0 (2020).
5 Zabihi, M. *et al.* Dissecting the Heterogeneous Cortical Anatomy of Autism Spectrum Disorder
Using Normative Models. *Biological psychiatry. Cognitive neuroscience and neuroimaging* **4**,
567-578, doi:10.1016/j.bpsc.2018.11.013 (2019).
6 Bethlehem, R. A. I. *et al.* A normative modelling approach reveals age-atypical cortical thickness
in a subgroup of males with autism spectrum disorder. *Communications Biology* **3**, 486,
doi:10.1038/s42003-020-01212-9 (2020).
7 Wolfers, T. *et al.* Mapping the Heterogeneous Phenotype of Schizophrenia and Bipolar Disorder
Using Normative Models. *JAMA Psychiatry* **75**, 1146-1155,
doi:10.1001/jamapsychiatry.2018.2467 (2018).
8 Cropley, V. L. *et al.* Brain-Predicted Age Associates With Psychopathology Dimensions in Youths.
Biological Psychiatry: Cognitive Neuroscience and Neuroimaging **6**, 410-419,
doi:https://doi.org/10.1016/j.bpsc.2020.07.014 (2021).
9 Hariri, A. R., Tessitore, A., Mattay, V. S., Fera, F. & Weinberger, D. R. The amygdala response to
emotional stimuli: a comparison of faces and scenes. *Neuroimage* **17**, 317-323,
doi:10.1006/nimg.2002.1179 (2002).
10 Hariri, A. R. *et al.* Serotonin Transporter Genetic Variation and the Response of the Human
Amygdala. **297**, 400-403, doi:doi:10.1126/science.1071829 (2002).
11 Miller, K. L. *et al.* Multimodal population brain imaging in the UK Biobank prospective
epidemiological study. *Nat Neurosci* **19**, 1523-1536, doi:10.1038/nn.4393 (2016).
12 Van Essen, D. C. *et al.* The WU-Minn Human Connectome Project: an overview. *Neuroimage* **80**,
62-79, doi:10.1016/j.neuroimage.2013.05.041 (2013).
13 Van Essen, D. C. *et al.* The Human Connectome Project: a data acquisition perspective.
Neuroimage **62**, 2222-2231, doi:10.1016/j.neuroimage.2012.02.018 (2012).
14 Harms, M. P. *et al.* Extending the Human Connectome Project across ages: Imaging protocols for
the Lifespan Development and Aging projects. *Neuroimage* **183**, 972-984,
doi:10.1016/j.neuroimage.2018.09.060 (2018).
15 Snoek, L. *et al.* The Amsterdam Open MRI Collection, a set of multimodal MRI datasets for
individual difference analyses. *Scientific Data* **8**, 85, doi:10.1038/s41597-021-00870-6 (2021).
16 van Eijndhoven, P. *et al.* Measuring Integrated Novel Dimensions in Neurodevelopmental and
Stress-Related Mental Disorders (MIND-SET): Protocol for a Cross-sectional Comorbidity Study
From a Research Domain Criteria Perspective. *JMIRx Med* **3**, e31269, doi:10.2196/31269 (2022).
17 Mulders, P. C. R. *et al.* Striatal connectopic maps link to functional domains across psychiatric
disorders. *Translational Psychiatry* **12**, 513, doi:10.1038/s41398-022-02273-6 (2022).
18 Witten, D. M., Tibshirani, R. & Hastie, T. A penalized matrix decomposition, with applications to
sparse principal components and canonical correlation analysis. *Biostatistics* **10**, 515-534,
doi:10.1093/biostatistics/kxp008 (2009).

19   Witten, D. M. & Tibshirani, R. J. Extensions of Sparse Canonical Correlation Analysis with
Applications to Genomic Data. **8**, doi:doi:10.2202/1544-6115.1470 (2009).
20   Bayer, J. M. M. *et al.* Accommodating site variation in neuroimaging data using normative and
hierarchical Bayesian models. *NeuroImage* **264**, 119699,
doi:https://doi.org/10.1016/j.neuroimage.2022.119699 (2022).
21   Nygaard, V., Rødland, E. A. & Hovig, E. Methods that remove batch effects while retaining group
differences may lead to exaggerated confidence in downstream analyses. *Biostatistics* **17**, 29-39,
doi:10.1093/biostatistics/kxv027 (2016).
22   Kebets, V. *et al.* Fronto-limbic neural variability as a transdiagnostic correlate of emotion
dysregulation. *Translational Psychiatry* **11**, 545, doi:10.1038/s41398-021-01666-3 (2021).
23   Rutherford, S. *et al.* Evidence for embracing normative modeling. *eLife* **12**, e85082,
doi:10.7554/eLife.85082 (2023).
24   Westlin, C. *et al.* Improving the study of brain-behavior relationships by revisiting basic
assumptions. *Trends in cognitive sciences*, doi:10.1016/j.tics.2022.12.015 (2023).
25   Everaerd, D., Klumpers, F., Oude Voshaar, R., Fernández, G. & Tendolkar, I. Acute Stress
Enhances Emotional Face Processing in the Aging Brain. *Biological Psychiatry: Cognitive*
Neuroscience and Neuroimaging **2**, 591-598, doi:https://doi.org/10.1016/j.bpsc.2017.05.001
(2017).
26   Mizzi, S., Pedersen, M., Lorenzetti, V., Heinrichs, M. & Labuschagne, I. Resting-state
neuroimaging in social anxiety disorder: a systematic review. *Molecular Psychiatry*,
doi:10.1038/s41380-021-01154-6 (2021).
27   Elliott, M. L. *et al.* What Is the Test-Retest Reliability of Common Task-Functional MRI Measures?
New Empirical Evidence and a Meta-Analysis. *Psychological Science* **31**, 792-806,
doi:10.1177/0956797620916786 (2020).
28   Somerville, L. H. *et al.* The Lifespan Human Connectome Project in Development: A large-scale
study of brain connectivity development in 5-21 year olds. *Neuroimage* **183**, 456-468,
doi:10.1016/j.neuroimage.2018.08.050 (2018).
29   Alfaro-Almagro, F. *et al.* Image processing and Quality Control for the first 10,000 brain imaging
datasets from UK Biobank. *Neuroimage* **166**, 400-424 (2018).
30   van Eijndhoven, P. F. P. *et al.* Measuring Integrated Novel Dimensions in Neurodevelopmental
and Stress-related Mental Disorders (MIND-Set): a cross-sectional comorbidity study from an
RDoC perspective. *medRxiv*, 2021.2006.2005.21256695, doi:10.1101/2021.06.05.21256695
(2021).
31   Oldehinkel, M. *et al.* Attention-Deficit/Hyperactivity Disorder symptoms coincide with altered
striatal connectivity. *Biological psychiatry. Cognitive neuroscience and neuroimaging* **1**, 353-363,
doi:10.1016/j.bpsc.2016.03.008 (2016).
32   Glasser, M. F. *et al.* The minimal preprocessing pipelines for the Human Connectome Project.
Neuroimage **80**, 105-124, doi:10.1016/j.neuroimage.2013.04.127 (2013).
33   Pruim, R. H. R. *et al.* ICA-AROMA: A robust ICA-based strategy for removing motion artifacts
from fMRI data. *Neuroimage* **112**, 267-277, doi:10.1016/j.neuroimage.2015.02.064 (2015).
34   Botvinik-Nezer, R. *et al.* Variability in the analysis of a single neuroimaging dataset by many
teams. *Nature* **582**, 84-88, doi:10.1038/s41586-020-2314-9 (2020).
35   Andersson, J. L., Jenkinson, M., Smith, S. & Oxford, F. A. G. o. t. U. o. Non-linear registration, aka
Spatial normalisation FMRIB technical report TR07JA2. **2**, e21 (2007).
36   Kraha, A., Turner, H., Nimon, K., Zientek, L. R. & Henson, R. K. Tools to support interpreting
multiple regression in the face of multicollinearity. *Frontiers in psychology* **3**, 44,
doi:10.3389/fpsyg.2012.00044 (2012).