Therapeutic targeting of ACLY in T-ALL *in vivo*

Authors: Victoria da Silva-Diz[1,#], Amartya Singh[1,2,#], Olga Lancho[1,#], Maya

Aleksandrova[1], Komal Mandleywala[1], Patricia Renck Nunes[1], Jesminara Khatun[1],

Oekyung Kim[1], Hayley Affronti[3], Eric Chiles[1], Xiaoyang Su[1,4], Hossein Khiabanian[1,2,5],

Kathryn E. Wellen[3], and Daniel Herranz[1,6,7*]

Affiliations:

[1]Rutgers Cancer Institute of New Jersey, Rutgers University, New Brunswick, NJ, 08901,

USA.

[2]Center for Systems and Computational Biology, Rutgers Cancer Institute of New Jersey,

Rutgers University, New Brunswick, NJ, 08901, USA.

20892, USA.

[3]Department of Cancer Biology, University of Pennsylvania Perelman School of Medicine,

Philadelphia, PA, 19104, USA.

[4]Department of Medicine, Rutgers Robert Wood Johnson Medical School, Rutgers

University, New Brunswick, NJ, 08901, USA

[5]Department of Pathology and Laboratory Medicine, Rutgers Robert Wood Johnson

Medical School, Rutgers University, New Brunswick, NJ, 08901, USA.

[6]Department of Pharmacology, Rutgers Robert Wood Johnson Medical School, Rutgers

University, Piscataway, NJ, 08854, USA.

[7]Department of Pediatrics, Rutgers Robert Wood Johnson Medical School, Rutgers

University, New Brunswick, NJ, 08901, USA.

[#]Equal contribution

*Corresponding author

Running title: Identification of ACLY as a therapeutic target in T-cell leukemia

Keywords: ACLY, NOTCH1, T-cell acute lymphoblastic leukemia, T-ALL

Financial support: Work in the laboratory of DH is supported by the US National

Institutes of Health (R01CA236936), the American Cancer Society (RSG-19-161-01-

TBE), The Leukemia & Lymphoma Society (Scholar Award 1386-23), the Alex's

Lemonade Stand Foundation, the Ludwig Cancer Research, the New Jersey Commission

on Cancer Research (COCR23PRG006) and the Rutgers Cancer Institute of New Jersey.

Work in the laboratory of H.K. is supported by the National Institutes of Health

(R01CA233662), the V Foundation (T2019-012), as well as by Rutgers Office of

Advanced Research Computing (NIH 1S10OD012346-01A1). In addition, Rutgers

Cancer Institute of New Jersey shared resources supported in part by the National Cancer

Institute Cancer Center Support Grant P30CA072720 were instrumental for this project,

including Biomedical Informatics Shared Resource (P30CA072720-5917), Metabolomics

Shared Resource (P30CA072720-5923) and the Pilot Award/New Investigator Award

(P30CA072720-5931). Moreover, purchase of the Eclipse and QE instruments was

supported by NIH grants S10OD025140 and S10OD01640. Fellowships from the New

Jersey Commission on Cancer Research supported the work of V.dD. (DCHS19PPC008)

and P.R.N. (COCR22PDF002). V.dD. was also supported by the Pediatric Cancer and

Blood Disorders Research Center at the Rutgers Cancer Institute of New Jersey.

Corresponding author:

Daniel Herranz

Assistant Professor of Pharmacology and Pediatrics

Robert Wood Johnson Medical School

Rutgers, Cancer Institute of New Jersey

Rutgers, The State University of New Jersey

195 Little Albany Street

Office Room 3037, Lab Room 3026

New Brunswick, NJ, 08901

Phone: +1-732-235-4064

e-mail: dh710@cinj.rutgers.edu

Conflict of interest disclosure:

The authors have declared that no conflict of interest exists.

Word count: 1153

Number of figures and tables: 2 main figures and 1 supplemental table.

Abstract

T-cell Acute Lymphoblastic Leukemia (T-ALL) is a hematological malignancy in need of novel therapeutic approaches. Here, we identify the ATP-citrate lyase ACLY as a novel therapeutic target in T-ALL. Our results show that ACLY is overexpressed in T-ALL, and its expression correlates with NOTCH1 activity. To test the effects of ACLY in leukemia progression and the response to NOTCH1 inhibition, we developed an isogenic model of NOTCH1-induced *Acly* conditional knockout leukemia. Importantly, we observed intrinsic antileukemic effects upon loss of ACLY, which further synergized with NOTCH1 inhibition *in vivo.* Gene expression profiling analyses showed that the transcriptional signature of ACLY loss very significantly correlates with the signature of NOTCH1 inhibition *in vivo*, with significantly downregulated pathways related to oxidative phosphorylation, electron transport chain, ribosomal biogenesis and nucleosome biology. Consistently, metabolomic profiling upon ACLY loss revealed a metabolic crisis with accumulation of nucleotide intermediates and reduced levels of several amino acids. Overall, our results identify a link between NOTCH1 and ACLY and unveil ACLY as a novel promising target for T-ALL treatment.

T-cell acute lymphoblastic leukemia (T-ALL) is an aggressive hematological disease of

immature T-cell progenitor cells. Although most patients nowadays get cured thanks to

intensive multiagent chemotherapy regimens, 20-50% patients still relapse, and the

prognosis at this stage is dismal[1]. Thus, identifying novel targets that could translate into

improved therapeutic effects in T-ALL is an urgent medical need. Related to this, targeting

different metabolic routes in patients has resulted in notable antileukemic effects, as

exemplified by the clinical use of methotrexate, 6-mercaptopurine or L-asparaginase[2]. In

this context, we recently identified the antileukemic effects of either serine

hydroxymethyltransferase inhibition[3] or mitochondrial uncoupling[4], highlighting the value

of further exploring metabolic vulnerabilities in leukemia. Thus, since NOTCH1 activating

mutations occur in over 60% of patients[5], and we previously demonstrated that the

metabolic effects of NOTCH1 inhibition are also critical for its therapeutic effects[6], we

hypothesized that thorough analyses of the metabolic enzymes being downregulated

upon NOTCH1 inhibition might yield novel therapeutic targets for T-ALL treatment.

Indeed, we found that NOTCH1 inhibition in T-ALL *in vivo* led to downregulation of the

ATP-citrate lyase *ACLY* (**Fig. 1A**). ACLY controls both metabolic and epigenetic

processes, given its critical role in fatty acid synthesis as well as its generation of acetyl-

CoA (critical for histone acetylation) as a product of its catalytic reaction[7]. Since both

metabolic and epigenetic mechanisms of resistance to NOTCH1 inhibition in T-ALL have

been previously described[6,8], our findings suggested a potential role for ACLY in the

response to NOTCH1 inhibition and, more broadly, as a relevant player in T-ALL. Next,

we analyzed available gene expression data from T-ALL pediatric patients[9] and observed

a broad upregulation of *ACLY* as compared to normal T-cell subsets (**Fig. 1B**). In line with

this, *ACLY* levels were similar across different T-ALL clinical subgroups[10] (**Fig. 1C**).

However, samples with NOTCH1 or FBXW7 mutations showed a trend towards increased

ACLY levels, which reached statistical significance in the cases with simultaneous

NOTCH1/FBXW7 mutations (**Fig. 1D**). Finally, ACLY overexpression in T-ALL was

confirmed at the protein level, with human T-ALL cell lines showing higher expression

that normal human thymus or peripheral blood cells (**Fig. 1E**). Taken together, our results

suggest a previously unknown but important role for ACLY in T-ALL.

Next, we hypothesized that targeting ACLY in T-ALL might confer therapeutic effects. To

test this hypothesis, we generated a model of NOTCH1-induced *Acly* conditional

knockout leukemia using a well-established protocol[4,6,11,12] of retroviral transduction of an

oncogenic form of NOTCH1 in bone marrow progenitor cells from *Acly* conditional

knockout mice, followed by transplantation into lethally irradiated recipients (**Fig. 2A**). In

time, these mice developed NOTCH1-induced leukemias, which were subsequently

transplanted into a secondary cohort of mice and treated with vehicle control or tamoxifen,

to induce isogenic loss of *Acly*. Mice were further subdivided into control-treated or treated

with the gamma-secretase inhibitor DBZ, to test the response to NOTCH1 inhibition *in*

vivo with or without ACLY expression (**Fig. 2A**). Notably, our results demonstrated

intrinsic antileukemic effects for ACLY loss, which further synergized with NOTCH1

inhibition resulting in a 20% cure rate of leukemic mice (**Fig. 2B**). Interestingly, leukemias

eventually developing in tamoxifen-treated mice were not genetic escapers and did not

show a concomitant feedback upregulation of ACSS2 (observed in some other contexts

upon ACLY loss[7]) either (**Fig. 2C**), suggesting the T-ALL cells can still survive ACLY loss

via alternative mechanisms. To explore the mechanistic effects of ACLY loss, we next

performed an acute deletion experiment in which mice harboring *Acly* conditional

knockout leukemias were first allowed to become fully leukemic before treating them with

vehicle-control or tamoxifen, followed by euthanasia 48h later. In this setting, acute ACLY

loss translated into reduced *Acly* levels (**Fig. 2D**) and led to reduced tumor burden (**Fig.**

2E), which was mainly driven by cytotoxic effects as revealed by increased AnnexinV

staining (**Fig. 2F**). Next, gene expression profiling analyses revealed a strong

transcriptional signature upon acute ACLY loss (**Fig. 2G**). Pathway analyses using the

STRING database[13] revealed strong downregulation of pathways related to TCA cycle,

electron transport chain and ribosomal biogenesis, together with a concomitant decrease

in nucleosome-biology related genes (**Fig. 2H** and **Supplemental Table 1**). Given the

correlation and effects on *ACLY* observed previously upon NOTCH1 inhibition (**Fig. 1**),

we next performed gene set enrichment analyses (GSEA)[14] against the signature

obtained upon NOTCH1 inhibition in mouse leukemias *in vivo*[6]. Of note, these analyses

revealed that the genes upregulated upon ACLY loss strongly correlate with the genes

upregulated by NOTCH1 inhibition and, conversely, genes downregulated upon ACLY

loss strongly correlate with the genes downregulated by NOTCH1 inhibition (**Fig. 2I**),

overall supporting a link between NOTCH1 and ACLY in T-ALL. Finally, given the strong

downregulation of transcriptional signatures linked to central metabolism, as well as the

well-known metabolic role of ACLY, we performed global metabolomic analyses upon

acute ACLY loss *in vivo*. Interestingly, while we did not observe significant changes in

pyruvate or acetyl CoA levels at this early time point, our results showed a strong

accumulation of AICAR (a purine intermediate and a well-described activator of AMPK[15])

together with concomitant accumulation of N-carbamoyl-L-aspartate (a pyrimidine

intermediate) and reduced levels of several amino acids, including aspartate, asparagine,

threonine and methionine (**Fig. 2J**), overall suggesting ACLY loss leads to a metabolic

crisis with compromised nucleotide biosynthesis in T-ALL cells.

ACLY has previously been suggested as an important therapeutic target in solid tumors

such as pancreatic cancer[16], and recent studies have described certain roles for ACLY in

the normal physiological function of CD4 and CD8 T-cells[17,18]. However, its putative role

in the normal development of T-cells and other hematological lineages, as well as in

leukemic transformation and progression, remains largely unknown. Our results

demonstrate that ACLY is upregulated in human T-ALL and, moreover, plays an important

role in leukemia progression *in vivo*. Our findings suggest a NOTCH1-ACLY link as

NOTCH1 inhibition leads to reduced *ACLY* levels, while NOTCH1 mutations in patients

correlate with higher levels of *ACLY*. In line with this, ACLY loss strongly synergized with

NOTCH1 inhibition *in vivo*, and the gene expression signature driven by ACLY loss very

significantly correlates with the signature obtained upon NOTCH1 inhibition. Intriguingly,

AKT is known to mediate NOTCH1-induced transformation[19], and AKT has also been

previously shown to directly phosphorylate and regulate ACLY[20]. Similarly, MYC is

another strong direct target of NOTCH1 in T-ALL[21], and recent studies support that MYC

might transcriptionally regulate ACLY[22], thus providing putative links between NOTCH1

and ACLY.

Taken together, our results demonstrate that targeting ACLY constitutes a novel and

attractive therapeutic option for leukemia treatment. While several ACLY inhibitors have

been described to date, these are either relatively weak and unspecific inhibitors, such as

bempedoic acid or BMS-303141[23,24], or potent inhibitors but not cell-permeable, such as

NDI-091143[25]. In this context, our unequivocal genetic results demonstrating a

therapeutic role for targeting ACLY in leukemia *in vivo* will hopefully spur additional efforts

to develop clinical-grade ACLY inhibitors that can be used in leukemia patients in the near

future.

METHODS

Cell lines and culture conditions

Cells were cultured in standard conditions in a humidified atmosphere in 5 % CO_2 at 37C

in HyClone RPMI 1640 Media (Fisher Scientific, SH3002701) with 10 % FBS (Gemini Bio-

Products, 900-108) and 100 U/mL penicillin and 100 μg/mL streptomycin (VWR, 45000-

652). DND41 (ACC 525), HPB-ALL (ACC 483), and JURKAT (ACC 282) cells were

obtained from Deutsche Sammlung von Mikroorganismen und Zellkulturen (DSMZ).

CUTTL1 and KOPTK1 were gifts from Adolfo Ferrando.

In vivo models of NOTCH1-driven mouse T-ALLs

Animals were maintained in ventilated caging in specific pathogen-free facilities at New

Brunswick RBHS Rutgers Campus. All animal housing, handling, and procedures

involving mice were approved by Rutgers Institutional Animal Care and Use Committee

(IACUC), in accordance with all relevant ethical regulations.

In order to generate *Acly* conditional inducible knockout leukemias, we used bone marrow

progenitor cells (fresh cells provided by K.E.W.) from *Acly*^{flox/flox} mice (JAX #043555)

harboring a tamoxifen-inducible Cre recombinase from the ubiquitous UBC locus (JAX

#007001). Then, we performed retroviral transduction of lineage-negative enriched cells

from with retrovirus encoding a ΔE-NOTCH1-GFP oncogenic activated form of NOTCH1

with concomitant expression of GFP, as previously described[6]. Cells were then

transplanted via retro-orbital injection into lethally irradiated (7.5 Gy) recipient mice.

For leukemia-progression studies, already generated ΔE-NOTCH1-GFP-induced

Acly^{flox/flox}-Cre-ERT2/+ leukemias (1 x 10⁶ leukemia cells) were transplanted from primary

recipients into sub-lethally irradiated (4.5 Gy) 6-8-week-old secondary recipient C57BL/6

mice (Taconic Farms) by retro-orbital injection. 48 hours after leukemic cell

transplantation, recipient mice were treated with vehicle only (corn oil; Sigma, C8267) or

tamoxifen (Sigma, T5648; 3 mg per mouse in corn oil), to induce isogenic loss of *Acly* in

established leukemias. Subsequently, 5 days post-transplantation, mice in each arm were

divided randomly into two different groups: control groups were subsequently treated with

vehicle only (2.3 % DMSO in 0.005% methylcellulose, 0.1 % Tween-80) or with DBZ (5

mg per Kg in vehicle solution; Syncom, 29762) on a 4-day-ON and 3-day-OFF schedule,

as previously described[6]. Investigators were not blinded to group allocation. Animals were

monitored for signs of distress or motor function at least twice daily until they were

terminally ill, whereupon they were euthanized.

For *Acly* acute deletion analyses in mouse primary leukemias, we transplanted

lymphoblasts from spleens of *Acly* conditional knockout NOTCH1-induced T-ALL-bearing

mice into a secondary cohort of recipient mice, as before. We monitored mice until they

presented clear leukemic signs with >60% GFP-positive leukemic cells in peripheral

blood; then, mice were treated with vehicle or tamoxifen and, 48 hours after treatment,

mice were euthanized and spleen samples were collected for further analyses.

Western blotting

Human peripheral blood mononuclear cells (PBMCs) and peripheral blood CD4-positive

cells were purchased from Lonza (product numbers CC-2702 and 2W-200, respectively).

Samples from normal human thymus were kindly provided by Dr. Adolfo Ferrando

(Columbia University). Whole-cell extracts were prepared using standard procedures.

After protein transfer, membranes were incubated with the antibodies anti-ACLY (1:1000,

15421-1-AP, Proteintech), anti-ACSS2 (1:1000, 3658, Cell Signaling) and anti-β-actin-

HRP (1:50000; A3854, Sigma). Antibody binding was detected with a secondary antibody

coupled to horseradish peroxidase (Sigma, NA934 and NA931) using enhanced

chemiluminescence (Thermo Scientific, 34578).

Flow cytometry analysis

To analyze leukemic spleen samples, single cell suspensions were prepared by

disrupting spleens through a 70 µm filter. Red cells were removed by incubation with

ammonium-chloride-potassium lysing buffer (155 mM NH4Cl, 12 mM KHCO3 and 0.1 mM

EDTA) for 5 minutes in ice. Apoptotic cells in leukemic spleens were quantified with PE-

AnnexinV Apoptosis Detection Kit I (BD Pharmingen, 559763). All flow cytometry data

was collected on an Attune NxT Flow Cytometer (ThermoFisher Scientific) and analyzed

with FlowJo v10.6.2 software (BD).

Metabolite extraction and LC-MS-based metabolomics

Leukemic spleen samples were processed and analyzed as previously described[4,11].

Quantitative RT-PCR

Total RNA was extracted from cells using RNeasy Plus Mini Kit (Qiagen), and cDNA was

generated with High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems).

Quantitative PCR was performed on a QuantStudio 3 Real-Time PCR System (Applied

Biosystems) using SYBR Green PCR Master Mix (Takara), and transcript levels were

normalized to actin as an internal control. Primers used: *Acly* (Forward: 5′-

TTCGTCAAACAGCACTTCC-3′; Reverse: 5′- ATTTGGCTTCTTGGAGGTG-3′) and *Actin*

(5′-GGCTGTATTCCCCTCCATCG-3′; 5′-CCAGTTGGTAACAATGCCATGT-3′).

RNA-seq gene expression profiling

Acly conditional knockout ∆E-NOTCH1-induced T-ALL-bearing mice were treated with

vehicle only (corn oil; Sigma, C8267) or tamoxifen (Sigma, T5648; 3 mg per mouse in

corn oil) to induce isogenic loss of *Acly* via intraperitoneal injection. 48 hours later, single-

cell suspensions of total leukemic splenocytes were prepared by pressing leukemic

spleens through a 70 μm filter. We removed red cells in spleen samples by incubation

with red blood cell lysis buffer (155 mmol/L NH4Cl, 12 mmol/L KHCO3, and 0.1 mmol/L

EDTA) for 5 minutes on ice. RNA was extracted using QIAshredder (QIAGEN, 79656)

and RNeasy Mini (QIAGEN, 74106) kits. RNA library preparations and next-generation

sequencing were performed using Illumina Next-Seq platform (Illumina). We estimated

gene-level raw counts and performed differential expression and/or GSEA analyses as

previously described[11].

RNA-seq analysis of Normal vs T-ALL

ACLY expression was analyzed among T-ALL samples (n=57) and physiological

thymocyte subsets (n=21) from published literature[9]. Quantile normalization was

performed across samples. Differential expression analysis was performed using the

Mann-Whitney U-Test and the resultant p-values were corrected for multiple hypothesis

testing using the Benjamini-Hochberg correction. Genes with FDR < 0.005 were

shortlisted as differentially expressed.

Statistical analysis

Statistical analyses were performed with Prism 8.0 (GraphPad). Unless otherwise

indicated in figure legends, statistical significance between groups was calculated using

an unpaired two-tailed Student's *t*-test. Survival in mouse experiments was represented

with Kaplan–Meier curves, and significance was estimated with the log-rank test.

Data availability

RNA-seq data from *Acly* conditional knockout leukemias was deposited in the NCBI Gene

Expression Omnibus repository under the following accession number: GSE227492.

Acknowledgements

We are grateful to Adolfo A. Ferrando (Columbia University Medical Center) for sharing

with us normal human thymocyte samples, and we thank both Adolfo A. Ferrando,

Antonio Maraver (IRCM, Montpellier) and Laura Belver (IJC, Barcelona) for their constant

constructive criticism and support. We also thank everyone involved with JuanLord for

their support. Figure 2A was created using BioRender.com.

Author contributions

V.dS. and O.L. performed most molecular biology experiments. A.S. performed all

computational analyses. M.A., P.R.N., J.K., K.M., O.K. and H.A. assisted with several

experiments. E.C. assisted with metabolomic experiments. X.S. supervised metabolomic

analyses. H.K. supervised all computational analyses. K.E.W. contributed critical

reagents and gave critical input. D.H. designed the study, supervised the research, and

wrote the manuscript with input from all authors.

References

1.   Hunger SP, Mullighan CG. Acute Lymphoblastic Leukemia in Children. *N Engl J*

Med. 2015;373(16):1541-1552.

2.   Rashkovan M, Ferrando A. Metabolic dependencies and vulnerabilities in

leukemia. *Genes Dev*. 2019;33(21-22):1460-1474.

3.   Garcia-Canaveras JC, Lancho O, Ducker GS, et al. SHMT inhibition is effective

and synergizes with methotrexate in T-cell acute lymphoblastic leukemia. *Leukemia*.

2021;35(2):377-388.

4.   da Silva-Diz V, Cao B, Lancho O, et al. A novel and highly effective mitochondrial

uncoupling drug in T-cell leukemia. *Blood*. 2021;138(15):1317-1330.

5.   Weng PA, Ferrando A, Lee W, et al. Activating Mutations of NOTCH1 in Human T

Cell Acute Lymphoblastic Leukemia. *Science*. 2004;306:269-271.

6.   Herranz D, Ambesi-Impiombato A, Sudderth J, et al. Metabolic reprogramming

induces resistance to anti-NOTCH1 therapies in T cell acute lymphoblastic leukemia. *Nat*

Med. 2015;21(10):1182-1189.

7.   Guertin DA, Wellen KE. Acetyl-CoA metabolism in cancer. *Nat Rev Cancer*.

2023;23(3):156-172.

8.   Knoechel B, Roderick JE, Williamson KE, et al. An epigenetic mechanism of

resistance to targeted therapy in T cell acute lymphoblastic leukemia. *Nat Genet*.

2014;46(4):364-370.

9.   Van Vlierberghe P, Ambesi-Impiombato A, Perez-Garcia A, et al. ETV6 mutations

in early immature human T cell leukemias. *J Exp Med*. 2011;208(13):2571-2579.

10.   Liu Y, Easton J, Shao Y, et al. The genomic landscape of pediatric and young adult

T-lineage acute lymphoblastic leukemia. *Nat Genet*. 2017;49(8):1211-1218.

11.   Lancho O, Singh A, da Silva-Diz V, et al. A Therapeutically Targetable NOTCH1-

SIRT1-KAT7 Axis in T-cell Leukemia. *Blood Cancer Discov*. 2023;4(1):12-33.

12.   Tottone L, Lancho O, Loh JW, et al. A Tumor Suppressor Enhancer of PTEN in T-

cell development and leukemia. *Blood Cancer Discov*. 2021;2(1):92-109.

13. Szklarczyk D, Morris JH, Cook H, et al. The STRING database in 2017: quality-

controlled protein-protein association networks, made broadly accessible. *Nucleic Acids*

Res. 2017;45(D1):D362-D368.

14. Subramanian A, Tamayo P, Mootha VK, et al. Gene set enrichment analysis: a

knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl*

Acad Sci U S A. 2005;102(43):15545-15550.

15. Vakana E, Altman JK, Glaser H, Donato NJ, Platanias LC. Antileukemic effects of

AMPK activators on BCR-ABL-expressing cells. *Blood*. 2011;118(24):6399-6402.

16. Carrer A, Trefely S, Zhao S, et al. Acetyl-CoA Metabolism Supports Multistep

Pancreatic Tumorigenesis. *Cancer Discov*. 2019;9(3):416-435.

17. Chowdhury S, Kar A, Bhowmik D, et al. Intracellular Acetyl CoA Potentiates the

Therapeutic Efficacy of Antitumor CD8+ T Cells. *Cancer Res*. 2022;82(14):2640-2655.

18. Osinalde N, Mitxelena J, Sanchez-Quiles V, et al. Nuclear Phosphoproteomic

Screen Uncovers ACLY as Mediator of IL-2-induced Proliferation of CD4+ T lymphocytes.

Mol Cell Proteomics. 2016;15(6):2076-2092.

19. Herranz D, Ferrando AA. An oncogenic enhancer enemy (N-Me) in T-ALL. *Cell*

Cycle. 2015;14(2):167-168.

20. Lee JV, Carrer A, Shah S, et al. Akt-dependent metabolic reprogramming

regulates tumor cell histone acetylation. *Cell Metab*. 2014;20(2):306-319.

21. Herranz D, Ambesi-Impiombato A, Palomero T, et al. A NOTCH1-driven MYC

enhancer promotes T cell development, transformation and acute lymphoblastic

leukemia. *Nat Med*. 2014;20(10):1130-1137.

22.   Singh KB, Hahm ER, Kim SH, Wendell SG, Singh SV. A novel metabolic function

of Myc in regulation of fatty acid synthesis in prostate cancer. *Oncogene*. 2021;40(3):592-

602.

23.   Granchi C. ATP citrate lyase (ACLY) inhibitors: An anti-cancer strategy at the

crossroads of glucose and lipid metabolism. *Eur J Med Chem*. 2018;157:1276-1291.

24.   Shah S, Carriveau WJ, Li J, et al. Targeting ACLY sensitizes castration-resistant

prostate cancer cells to AR antagonism by impinging on an ACLY-AMPK-AR feedback

mechanism. *Oncotarget*. 2016;7(28):43713-43730.

25.   Wei J, Leit S, Kuai J, et al. An allosteric mechanism for potent inhibition of human

ATP-citrate lyase. *Nature*. 2019;568(7753):566-570.

   Figures and Figure Legends

Figure 1. ACLY is overexpressed in human T-ALL. **A,** mRNA levels of *Acly* upon

NOTCH1 inhibition with the DBZ gamma-secretase inhibitor in mouse T-ALL *in vivo*[6]

(**P<0.01 calculated with 2-tailed Student's *t*-test). **B,** Box-plot showing *ACLY* expression

among T-ALL samples (n=57) and physiological thymocyte subsets (n=21)[9]. Quantile

normalization was performed across samples. Boxes represent first and third quartiles

and lines represent the median. Whiskers represent the upper and lower limits (P<0.001

using Mann-Whitney U-Test; FDR<0.05 using Benjamini-Hochberg correction). **C,** *ACLY*

expression levels in human primary T-ALL cases from different subclinical groups[10]. **D,**

ACLY expression levels in human primary T-ALL cases with/without *NOTCH1* and

FBXW7 mutations[10] (Wilcoxon rank-sum test *P < 0.05). **E,** Western blot analysis of

ACLY, ACSS2 and ACTIN expression in human peripheral blood mononuclear cells

(PBMNC), CD4+ T-cells or healthy human thymocytes, as compared to human T-ALL.

Figure 2. Targeting ACLY shows antileukemic and synergistic effects with NOTCH1 inhibition *in vivo*. **A,** Schematic of retroviral-transduction protocol for the generation of NOTCH1-induced T-ALLs from inducible *Acly*-conditional knockout mice, followed by transplant into secondary recipients treated with vehicle (*Acly*$^{+/+}$) or tamoxifen (*Acly*$^{-/-}$), with or without DBZ. **B,** Kaplan-Meier survival curves of mice harboring *Acly*-positive and *Acly*-deleted isogenic leukemias treated with 4 cycles of vehicle or DBZ (5 mg/kg) on a 4-days-ON (blue blocks at the bottom) and 3-days-OFF schedule (log-rank test; **$P < 0.01$; n = 10 per group). **C,** Western blot analysis of ACLY, ACSS2 and ACTIN expression in leukemic spleens from terminally ill mice from survival curve in B. **D**, Quantitative RT-PCR analysis of *Acly* mRNA expression in tumor cells isolated from ΔE-NOTCH1-induced *Acly* conditional knockout leukemia–bearing mice 48 h after being treated with vehicle only (*Acly*$^{+/+}$) or tamoxifen (*Acly*$^{-/-}$) *in vivo*. **E,** Tumor burden in ΔE-NOTCH1-induced *Acly* conditional knockout leukemia–bearing mice 48 h after being treated with vehicle only (*Acly*$^{+/+}$) or tamoxifen (*Acly*$^{-/-}$) *in vivo* as revealed by total spleen weight. **F**, Representative flow cytometry plots from of annexin V (apoptotic cells) and 7-AAD (dead cells) staining (left) and quantification of apoptosis (right) in leukemic spleens from ΔE-NOTCH1-induced *Acly* conditional knockout leukemia–bearing mice 48 h after being treated with vehicle only (*Acly*$^{+/+}$) or tamoxifen (*Acly*$^{-/-}$) *in vivo*. (n = 5 per treatment; *$P < 0.05$ and ***$P < 0.005$ using two-tailed Student's *t*-test. **G,** Heat map representation of the top differentially expressed genes between control (*Acly*$^{+/+}$) and tamoxifen-treated (*Acly*$^{-/-}$) *Acly*-conditional knockout ΔE-NOTCH1–induced leukemias. Cutoffs used: top 40 up and downregulated genes; *P*-adjusted value < 0.0001; log fold change >1. Scale bar shows color-coded differential expression, with red indicating higher levels of expression

and blue indicating lower levels of expression **H,** Pathway analyses of significantly

downregulated genes upon ACLY loss (flod change > 1.7; *P* < 0.001) using the STRING

database[13]. **I,** GSEA of genes regulated by ACLY in control (vehicle only–treated;

NOTCH1 ON) compared with DBZ-treated (NOTCH1 OFF) ΔE-NOTCH1–induced mouse

leukemias *in vivo*[6]. **J,** Relative abundance of indicated metabolites upon tamoxifen-

induced isogenic loss of *Acly* in leukemic spleens from mice treated as in D (n = 5 per

treatment; *P < 0.05; **P < 0.01; *** P < 0.005 using two-tailed Student's *t*-test.).