1
2 Transcriptional regulation of the synthesis and secretion of farnesol and aromatic fusel alcohols in the
3 fungus *Candida albicans:* Examination of the Homann transcription regulator knockout collection

11 Daniel J. Gutzmann*, Jaxon J. Kramer, Brigid M. Toomey, Cory H. T. Boone, Audrey L. Atkin* & Kenneth
12 W. Nickerson*
18 School of Biological Sciences
19 University of Nebraska
20 Lincoln, NE USA
27 *For correspondence. E-mail : knickerson1@unl.edu, aatkin@unl.edu
28 daniel.gutzmann@huskers.unl.edu
29 Tel (402) 472-2253; Fax (402) 472-8722.

Keywords

- *Candida albicans*

- Farnesol

- Aromatic fusel alcohol

- Transcription factor mutant

Abstract

Candida albicans is an efficient colonizer of human gastrointestinal tracts and skin and is an

opportunistic pathogen. *C. albicans* exhibits morphological plasticity and the ability to switch between

yeast and filamentous morphologies is associated with virulence. One regulator of this switch is the

quorum sensing molecule farnesol which is produced in *C. albicans* cells throughout growth. In addition,

aromatic fusel alcohols accumulate in *C. albicans* supernatants and are implicated in regulating

morphogenesis. The synthesis, secretion, and turnover of these molecules is not fully understood. To

address this, we used our improved assay for farnesol and aromatic fusel alcohol quantification of whole

cultures, pellets, and supernatants, to screen a transcription regulator knockout library of 164 mutants

for differences in farnesol and aromatic fusel alcohol accumulation and localization. We found that all

mutants produced farnesol with an average of 9.2X more farnesol localizing to the pellet in most

mutants. Nineteen mutants had significant differences in accumulation and nine differed in localization.

Ten mutants produced more farnesol than their SN152^{+} parent, while nine produced less. Seven

mutants exhibited greater secretion of farnesol while two exhibited less. Several of these mutants have

known roles in white/opaque switching and cell cycle progression such as the farnesol overproducing

mutants *ahr1* and *swi4*. The aromatic fusel alcohols were similarly characterized by four over- and

eleven under-producing mutants. Phenethyl alcohol and tryptophol accumulation were closely

proportional to one another, but despite their structural similarity, phenethyl alcohol and tyrosol

accumulation were often regulated differently.

Introduction

Candida albicans is an opportunistic pathogen that is present in most human gastrointestinal

tracts and is an efficient colonizer of mucosal surfaces (Neville et al. 2015). A weakened immune system

(HIV, chemotherapy, organ transplantation) or loss of competing flora (antibiotic treatment) allows for

C. albicans to colonize and invade host tissues leading to candidiasis (Pfaller and Diekema 2007).

Virulence of *C. albicans* is strongly linked to its ability to switch between yeast and filamentous

morphologies (Romano 1966; Kadosh 2019) as mutants that have lost the ability to switch remain

avirulent (Lo et al. 1997; Saville et al. 2003). Dimorphism is also important for adaptation to different

host environments (Noble et al. 2017; Alves et al. 2020) and evasion of host immune responses (Gow et

al. 2012). One regulator of dimorphic switching is the quorum sensing molecule (QSM) farnesol (Hornby

et al. 2001). Farnesol is produced by *C. albicans* cells continuously throughout growth (Boone et al.

2022). However, despite numerous publications on farnesol's role as a QSM and a virulence factor, the

actual mechanisms regulating its synthesis and secretion remain unclear.

C. albicans also secretes the aromatic fusel alcohols phenethyl alcohol, tyrosol, and tryptophol

(Ghosh et al. 2008), with most accumulation occurring in stationary phase (Boone et al. 2022). The

aromatic fusel alcohols are a product of the Ehrlich pathway (Hazelwood et al. 2008) in *S. cerevisiae* and

C. albicans. These molecules may also participate in yeast-mycelia morphogenesis (Chen et al. 2004;

Chen and Fink 2006) and biofilm formation (Alem et al. 2006), but regulation of their accumulation

remains unclear.

As an attractive approach to identifying these genes and mechanisms, we tested the hypothesis

that farnesol and fusel alcohol synthesis and secretion are regulated by the transcriptional networks

involved in morphogenesis and virulence. The transcription regulator knockout library generated by

Homann and colleagues consists of deletion strains for 165 different transcriptional regulators, in most

cases two independently derived knockouts (Homann et al. 2009). Thus, in the present paper the strains

in the transcription regulator knockout library were grown in liquid culture and then assayed by the

improved methods of Boone et al. (2022) for total, extracellular, and intracellular farnesol and the

aromatic fusel alcohols. Key features of this assay include prevention of analyte loss by avoiding

filtration and evaporation, while incorporating simultaneous cell lysis and analyte extraction by ethyl

acetate with the ability to compare whole cultures with the sum of their cell pellets and supernatants. It

is appropriate to screen for farnesol and the aromatic fusel alcohols together because there have been

numerous reports that they are all QSM for the yeast-mycelial transition in *Candida albicans*.

Materials and Methods

Strains and media

The homozygous transcription regulator deletion mutants were obtained from the transcription

regulator knockout library provided by Dr. Alexander Johnson's lab (Homann et al. 2009) available

through the Fungal Genomics Stock Center (Manhattan, Kansas, USA; McCluskey et al. 2010). All assays

with these mutants include the paired wild-type strain located in the H12 well of the X2 and Y2

independently derived deletion collections. The strains were maintained at 30°C on plates with YPD

(1% yeast extract, 2% peptone, 2% glucose). The transcription regulator knockout mutants were

constructed from *C. albicans* SN152, an auxotroph for arginine, histidine, and leucine. Each of the

transcription regulator (TR) mutants is auxotrophic for arginine. A wild-type control strain was created

by reintroduction of a single allele of HIS1 and LEU2 into the parent strain (Homann et al. 2009).

Throughout this paper, we will refer to this control strain simply as SN152^{+}.

Screening the transcription regulator deletion mutants for differences in farnesol and aromatic fusel

alcohol accumulation and secretion

The transcription regulator mutant collection was transferred by a 48-pin replicator to YPD

plates and grown at 30°C for 48 hours. Mutants were then transferred to 3 mL YPD liquid medium and

grown for 16 hours at 30°C with rotary agitation at 250 RPM. This culture was used as the inoculum

(1:100) for 75 mL YPD in 250 mL flasks which were incubated 24 hours at 30°C, 250 RPM. 50 mL of

culture was used for dry weight determination and 20 mL (2 X 10 mL) was used for determination of

farnesol and aromatic fusel alcohol production as described by Boone et al. (2022). In this method, 10

mL was used to measure whole culture values while the other was centrifuged to obtain the pellet and

supernatant values. Due to the large number of mutants in the collection, the mutants were assayed in

batches of 10-20 mutants with the SN152[+] control assayed in triplicate as apart of each batch. Mutants

with farnesol or aromatic fusel alcohol accumulation of greater than 1 $\log_2$ fold above or below the

mean value for SN152[+] were chosen for follow-up analysis. In follow-up analyses, E,E-farnesol and

aromatic fusel alcohol accumulation were measured in at least three independent experiments using

both the X and Y independently-derived TR deletion mutants.

Statistical analysis

Statistical analyses were performed using Microsoft Excel (Version 16.61, Microsoft Office, Las

Vegas, NV, USA) and GraphPad Prism Software (Version 9.5.0, San Diego, CA, USA). All biological data

are represented as mean $\pm$ SD of at least 3 biological replicates in both the X and Y independent mutants

unless otherwise stated. Mutants with significant differences between the X and Y mutant were

excluded and considered a false positive. Normality and homogeneity of variance were assessed by the

D' Agostino-Pearson omnibus (K2) test and Brown-Forsythe test, respectively. Differences between

groups that were normally distributed and homoscedastic (equal variances) were assessed by One-way

ANOVA with Dunnett's multiple comparisons test. Differences between groups that were not normally

distributed or heteroscedastic were assessed by Kruskal-Wallis test with Dunn's multiple comparisons

tests. Both analyses were performed in GraphPad Prism. Differences were considered significant at

$p<0.05$ (*$p<0.05$, **$p<0.01$, ***$p<0.001$).

Results

Identification of transcription regulator mutants defective in accumulation of intracellular and

extracellular farnesol and aromatic fusel alcohols

Homann et al. (2009) screened their collection of 165 independent transcription regulator mutants

for possible differences in 55 growth conditions. It was a monumental piece of work. However, for

practical reasons they limited themselves to phenotypes discernible via colonies on agar plates. We

have now extended the list of phenotypes to include accumulation of both intracellular and extracellular

farnesol as well as the aromatic fusel alcohols, as determined by our improved GC assay (Boone et al,

2022). For the initial screen, 198 cultures were screened, consisting of 164 mutants from the X plates

and 34 replicate controls for the SN152$^+$ parent, assayed in 12 batches including duplicate or triplicate of

the parent in each run, for a total of 594 GC-FID runs.

147   For the rescreen, 54 strains were selected because their farnesol and/or fusel alcohol values

differed from the average values for SN152$^+$ by a $\log_2$ fold change of > 1.0. Each of these values was

normalized on a per cell basis using the 50 mL dry weight value of the culture assayed. The initial screen

data set contains the same information for all 164 TR mutants with n=1. The rescreen data set examines

54 TR mutants, both raw data and their means ± SD with n=6, reporting normalized accumulation values

for phenethyl alcohol, farnesol, tyrosol, and tryptophol as detected in whole cultures, pellets, and

supernatants. For each mutant we also report the mean $\log_2$ fold change vs the SN152$^+$ control for each

of the four analytes as detected in whole cultures, pellets, and supernatants, as well as the

supernatant/pellet ratio for each analyte. Composite data sets for the initial screen (File S1) and

rescreen (File S2) are available at Figshare. We now describe several highlights from these data sets.

General Design and Reproducibility

Our initial screen of 164 strains, all from the X plate, was done with an n = 1 comparing whole

culture values versus the sum of the cell pellet and supernatant values. Strain *hfl1Δ/Δ* [orf19.3063] was

excluded as it exhibited a severe growth defect on YPD. The 164 TR mutants were assayed in 12 batches

which always included a biological duplicate or triplicate of the SN152[+] parent. In this initial screen, the

paired WT controls were run a total of 34 times and their whole culture mean values were 5.29 ± 1.12,

115.32 ± 31.32, 30.24 ± 5.60, and 9.52 ± 3.00 ng/μL/ 50 mL dry weight for E,E-farnesol, phenethyl

alcohol, tryptophol, and tyrosol, respectively. Thus, the ratio of mean values for the three fusel alcohols

were 12:3:1 for phenethyl alcohol, tryptophol, and tyrosol. The distribution of these 34 values for

SN152[+] is shown in Fig. 1 where they can be seen in comparison with the means and distribution of the

164 TR mutants. For reference, these values are denoted by the red X in Figs. 2-5. Fig. 1 confirms that

the vast majority of the 164 TR mutants were unchanged from the SN152[+] replicates. As expected

(Boone et al. 2022), the whole culture (W) and pellet (P) + supernatant (S) values were always very close

to one another with the average WPS relative error values (W-(P+S)/W) being: phenethyl alcohol 0.12 ±

0.09, E,E-farnesol 0.07 ± 0.06, tyrosol 0.21 ± 0.15, and tryptophol 0.19 ± 0.28.

We then identified all mutants which were > 1 $\log_2$ above or below the mean values, as

indicated by the 4 dotted lines in Fig. 2, and designated them as mutants of interest to be rescreened

with greater accuracy. There were 31 mutants with higher or lower farnesol accumulation and 40

mutants with higher or lower aromatic fusel alcohols; they made a cohort of 49 strains because 22

strains had overlapping differences. To these we added 5 strains (*upc2Δ/Δ, tac1Δ/Δ, dal81Δ/Δ,*

aro80Δ/Δ, and csr1Δ/Δ) because prior reports from the literature indicated they were likely to have

altered farnesol or aromatic fusel alcohol accumulations. These 54 mutant strains were then rescreened

five more times, two from the X plate and three times from the Y plate (total n =6) in 10 batches

including the SN152[+] parent in biological duplicate or triplicate, requiring 292 flasks and 876 GC

measurements. Ten mutants (*cta7Δ/Δ, cwt1Δ/Δ, rim101Δ/Δ, crz1Δ/Δ, zcf1Δ/Δ, ofi1Δ/Δ, leu3Δ/Δ,*

tea1Δ/Δ, ume7Δ/Δ, rme1Δ/Δ) were designated as false positives because their rescreened values from

the X and Y plates differed significantly or their rescreen analyte values differed significantly from the

initial screen. Thus, the TR mutants which were identified as being statistically different included ten

with elevated farnesol and nine with reduced farnesol (Fig. 6A-B, Table 1), four with elevated aromatic

fusel alcohols and 11 with reduced aromatic fusel alcohols (Table 2, Fig. 7A-B), and seven with elevated

farnesol supernatant/pellet ratios and two with lowered farnesol supernatant/pellet ratios (Fig. 6C).

These constituted only 35 mutants of interest because three mutants (*swi4Δ/Δ, rap1Δ/Δ, and cas5Δ/Δ*)

were on all three lists, and another ten mutants were on two.

Mutants altered in farnesol accumulation

Of the 19 mutants with statistically significant differences in farnesol accumulation, only

nrg1Δ/Δ had been identified in prior literature (Kebaara et al. 2008). Seven mutants (*swi4Δ/Δ, ahr1Δ/Δ,*

nrg1Δ/Δ, rap1Δ/Δ, pho4Δ/Δ, tye7Δ/Δ, aaf1Δ/Δ) had increased farnesol accumulation either in the whole

culture or supernatant fraction with a $\log_2$ fold change of 1 or greater compared to the SN152[+] parent

strain (Table 1). The two highest accumulators (*swi4Δ/Δ* and *ahr1Δ/Δ)* have known roles in cell cycle

progression (Hussein et al. 2011; Sellam et al. 2019). Two farnesol over accumulating mutants (*tye7Δ/Δ*

and rtg1*Δ/Δ*) have roles in carbohydrate metabolism (Askew et al. 2011; Moreno-Velásquez et al. 2020).

Four farnesol over accumulating mutants (*ahr1Δ/Δ, nrg1Δ/Δ, ash1Δ/Δ, ssn6Δ/Δ*) have a role in

dimorphism (Fu et al. 1998; Murad et al. 2001; Inglis and Johnson 2002; García-Sánchez et al. 2005).

Three farnesol over accumulating mutants (*ahr1Δ/Δ, ssn6Δ/Δ, wor2Δ/Δ)* have roles in white/opaque

switching (Zordan et al. 2007; Wang et al. 2011; Hernday et al. 2016). This is of particular interest as

another regulator of white/opaque switch, Czf1, has a known role in coordinating the response to

farnesol (Langford et al. 2013), and farnesol production is turned off in both anaerobically grown cells

(Dumitru et al. 2004) and opaque cells (Dumitru et al. 2007).

Eight mutants (*orf19.6874Δ/Δ, msn4Δ/Δ, cph2Δ/Δ, fcr1Δ/Δ, sko1Δ/Δ, gzf3Δ/Δ, cas5Δ/Δ,*

cup9Δ/Δ) had decreased farnesol accumulation either in the whole culture or supernatant fraction with

a $\log_2$ fold change of <1 or greater compared to the SN152[+] parent strain (Table 1). Three of these low

farnesol accumulating mutants (*msn4Δ/Δ, sko1Δ/Δ, cas5Δ/Δ*) have known roles in stress response

(Nicholls et al. 2004; Bruno et al. 2006; Rauceo et al. 2008; Xie et al. 2017) and two mutants (*sko1Δ/Δ,*

cup9Δ/Δ) have known roles in farnesol response (Lu et al. 2014).

Three of the four mutants chosen because of literature precedent had elevated farnesol, but did

not reach statistical significance (*csr1Δ/Δ, upc2Δ/Δ, tac1Δ/Δ*) (Table 1). One mutant (*tup1Δ/Δ*) did not

appear to over accumulate farnesol in these assay conditions (Table 1). At present we have no

explanation why the *tup1Δ/Δ* mutant did not over produce farnesol at the levels we had reported

previously (Kebaara et al. 2008).

Mutants altered in the farnesol supernatant/pellet ratio

The supernatant/pellet (S/P) ratios for each analyte were determined for each mutant in

comparison to those for their SN152[+] control (Fig. 3). The value for SN152[+] (n= 22 batches) was 0.12 for

farnesol. As we had observed previously (Boone et al. 2022), farnesol was predominantly retained in the

cell pellet. The S/P ratios were all tightly clustered from 0.11-0.16 except for four (*msn4Δ/Δ, fcr1Δ/Δ,*

gzf3Δ/Δ, and orf19.6874Δ/Δ) which exhibited less secretion (0.05-0.08) and seven (*swi4Δ/Δ, pho4Δ/Δ,*

cas5Δ/Δ, hap5Δ/Δ, ssn6Δ/Δ, tye7Δ/Δ, and rap1Δ/Δ) which exhibited greater secretion (0.19-0.30) (Fig.

6C).

Among these mutants, *hap5Δ/Δ* and *cas5Δ/Δ* are of special interest in that they exhibit greater

secretion without being accompanied by greater total farnesol production (Fig. 3). Thus, they target

secretion specifically. Mutants such as *swi4Δ/Δ, ahr1Δ/Δ, and rap1Δ/Δ* with much greater farnesol

production (Fig. 6A, Table 1) could exhibit a higher supernatant to pellet ratio because internal and

external farnesol are in equilibrium or there is a need to keep internal farnesol levels below a critical

threshold value. In both cases, the greater secretion would be an indirect consequence of greater

production.

Mutants altered in aromatic fusel alcohol accumulation

Nineteen mutants were identified with statistically significant differences in aromatic fusel

alcohol accumulation (Table 2, Fig. 7); 15 exhibited decreased production while only four exhibited

increased production. The four with increased production were *CR_02120CΔ/Δ, fgr15Δ/Δ, gzf3Δ/Δ,* and

hap5Δ/Δ. *CR_02120CΔ/Δ* and *fgr15Δ/Δ* were increased for all three aromatic fusel alcohols, *gzf3Δ/Δ* for

only phenethyl alcohol, and *hap5Δ/Δ* for only tyrosol (Table 2, Fig. 7A). Gzf3 has a known role in

negatively regulating nitrogen catabolite repression (Soussi-Boudekou et al. 1997).

Fifteen mutants had statistically significant decreased accumulation (*tup1Δ/Δ, ssn6Δ/Δ, stp2Δ/Δ,*

rap1Δ/Δ, dal81Δ/Δ, nrg1Δ/Δ, stp4Δ/Δ, swi4Δ/Δ, ahr1Δ/Δ, czf1Δ/Δ, ndt80Δ/Δ, cph2Δ/Δ, bcr1Δ/Δ,

cas5Δ/Δ, aro80Δ/Δ). Of these, 5 mutants (*tup1Δ/Δ, ssn6Δ/Δ, rap1Δ/Δ, stp4Δ/Δ, ndt80Δ/Δ*) were

decreased for all three aromatic fusel alcohols and five mutants (*stp2Δ/Δ, dal81Δ/Δ, nrg1Δ/Δ, swi4Δ/Δ,*

cph2Δ/Δ) were decreased for phenethyl alcohol and tryptophol (Table 2, Fig. 7B). A plot of phenethyl

alcohol accumulation vs tryptophol accumulation for SN152[+] and all 164 TR mutants (Fig. 4) showed that

these two fusel alcohols were closely proportional to one another, at least when grown on YPD. Tup1

and Ssn6 form a corepressor complex in yeast and have a known role in regulating amino acid

permeases (Tanaka and Mukai 2015), while Stp2 is an amino acid regulated TR that regulates amino acid

permeases (Martínez and Ljungdahl 2005). Dal81 has a known role in positively regulating nitrogen

degradation in *S. cerevisiae* (Bricmont et al. 1991), but has a different role in regulating arginine

synthesis in *C. parapsilosis* (Turner et al. 2018).

Finally, two mutants (*ahr1Δ/Δ, czf1Δ/Δ*) were decreased in phenethyl alcohol only, one mutant

aro80Δ/Δ in tyrosol only, one mutant *cas5Δ/Δ* in tryptophol only, and one mutant *bcr1Δ/Δ* in both

tryptophol and tyrosol (Table 2). Aro80 has a known role in regulating the Ehrlich pathway (Ghosh et al.

2008). An intriguing suggestion from these data is that despite their close structural similarity, phenethyl

alcohol and tyrosol accumulation are regulated differently. This idea is further supported by comparison

of their respective syntheses in all the TR mutants (Fig. 5).

Mutants altered in the aromatic fusel alcohol supernatant/pellet ratio

The three fusel alcohols were mostly secreted. The S/P ratios for SN152[+] (n=22) were 13.9, 8.9,

and 6.6 for phenethyl alcohol, tyrosol, and tryptophol, respectively. For phenethyl alcohol, the S/P ratios

were all tightly clustered from 11-20 except for *tup1Δ/Δ* at 2.34, *nrg1Δ/Δ* at 5.45, and *ssn6Δ/Δ* at 9.53. In

contrast, the S/P values for tryptophol were more diverse; 36 mutants were in the 5-8 range, close to

SN152[+], while 10 exhibited much less secretion (1.1-3.8) and another 10 exhibited much greater

secretion (11.6-34) (File S2).

Discussion

We examined each of the 164 independent transcriptional regulator knockout mutants for their

accumulation and secretion of farnesol and the three aromatic fusel alcohols. This project was

undertaken to define the genes needed for their synthesis, secretion, and regulation. For logistical

reasons, all cultures were grown at 30°C in liquid YPD medium and sampled after 24 hours when the

cells were in early stationary phase. The 24 hour time point was chosen based on our prior

measurements with wild type *C. albicans* SC5314 over 72 hours of culture which showed that the time

courses for farnesol and fusel alcohol production differed dramatically (Boone et al. 2022). Farnesol

generally followed increased cell mass, peaked in early stationary phase, and then declined sharply after

~ 30 hours. In contrast, accumulation of the aromatic fusel alcohols only started about 12 hours after

inoculation, just as the cells were entering stationary phase, and continued throughout stationary phase

(Boone et al. 2022). Thus, the 24 hour time point was intended to cover the four analytes of interest.

Similarly, our choice of YPD as a growth medium was to insure that all of the TR mutants grew well by

24hrs and indeed 164 out of 165 had done so.

The choice of a single 24 hour time point also impacts the observed fusel alcohol production

ratios. The SN152[+] ratios at 24 hours were 12:3:1 for phenethyl alcohol, tryptophol, and tyrosol,

respectively (Figs. 4 and 5), but they would likely be different if we had chosen a 36 or 48 hour time

point instead. In an earlier study with *C. albicans* SC5314, we found that the appearance of phenethyl

alcohol and tryptophol closely paralleled one another, plateauing in accumulation by 36 hours, but the

appearance of tyrosol was delayed, with much of it accumulating after 36 hours (Boone et al. 2022).

With the advantage of now having reliable measures for analyte presence in whole cultures, cell

pellets, and supernatants, we can now approach the genetic basis for their synthesis, secretion,

regulation, and turnover. As expected, most of the TR mutants were unchanged for these parameters

(Figs. 1-5) while a few potentially useful mutants gave significantly higher and lower levels of the

analytes of interest. Significantly, no mutants were entirely devoid of farnesol; its levels often varied but

they never dropped to zero. A tentative conclusion from this observation is that farnesol has an

essential physiological function over and above acting as a quorum sensing molecule and a virulence

factor or is the product of a redundant biosynthetic pathway.

There are many questions which can now be approached via the genetic information gleaned

from this screen. Among them are: 1/ How is carbon flow through the farnesyl pyrophosphate (FPP)

branch point regulated? 2/ Is farnesol made directly from FPP by an appropriate pyrophosphatase (e.g.

Dpp1p, Dpp2p, or Dpp3p) or are there other more easily regulated and/or safer biosynthetic pathways?

A method suggested from *Arabidopsis* invokes the necessary turnover of farnesylated proteins, where

proteolytic degradation of those proteins releases farnesylated cysteine which can be cleaved by a ligase

to release farnesaldehyde which is then converted to farnesol by a farnesol dehydrogenase (Bhandari et

al. 2010). 3/ Is the disappearance of farnesol later in stationary phase due strictly to evaporation of a

volatile molecule or to enzymatic conversion of farnesol to, for instance, 2,3-dihydrofarnesol (Brasch et

al. 2013; Costa et al. 2020). Significantly, regarding its ability to block germ tube formation and hyphal

development, 2,3-dihydrofarnesol is inactive; with only 0.34% of the activity of farnesol (Shchepin et al.

2003). 4/ Large amounts of the aromatic fusel alcohols are produced, usually in the ratios of phenethyl

alcohol >> tyrosol and tryptophol. Are these fusel alcohols synthesized via the well-known Ehrlich

pathway as in *S. cerevisiae* (Hazelwood et al. 2008), through the biosynthetic shikimic acid pathway

using prephenate dehydratase or dehydrogenase (Dai et al. 2021), or as in Arabidopsis from tyrosine to

tyrosol via a tyramine intermediate (Chung et al. 2017; Rodrigues and Černáková 2020)? These are just

four examples. Likely many more will surface following a careful examination of the collected data now

available in this mutant screen.

Author contributions:

DJG: conceptualization, methodology, validation, investigation, writing — original draft, writing —

review and editing, visualization, formal analysis, data curation, project administration; JJK: validation,

investigation, writing — review and editing; BMT: validation, investigation, writing — review and editing;

CHTB: methodology, validation, writing – review and editing; ALA: conceptualization, methodology,

resources, supervision, writing — review and editing, project administration; KWN: conceptualization,

methodology, resources, supervision, writing — original draft, writing — review and editing, project

administration, funding acquisition.

Data availability:

The datasets generated during and/ or analyzed during the current study are available from the

corresponding author on reasonable request. The initial screen (File S1) and rescreen dataset (File S2)

are available at Figshare.

Acknowledgments:

We are grateful to the Fungal Genetics Stock Center (FGSC) for providing strains.

Compliance with ethical standards:

Funding: This work was supported by Ann L. Kelsall and the Farnesol and *Candida albicans* Research

Fund, University of Nebraska Foundation. This research did not receive any specific grant from funding

agencies in the public, commercial, or not-for-profit sectors.

Conflict of interest: Daniel J. Gutzmann declares that he has no conflict of interest. Jaxon J. Kramer

declares that he has no conflict of interest. Brigid Toomey declares that she has no conflict of interest.

Cory H.T. Boone declares that he has no conflict of interest. Audrey L. Atkin declares that she has no

conflict of interest. Kenneth W. Nickerson declares that he has no conflict of interest.

Ethical approval: This article does not contain any studies with human participants or animals

performed by any of the authors.

References

Alem MAS, Oteef MDY, Flowers TH, Douglas LJ. 2006. Production of tyrosol by *Candida albicans* biofilms
and its role in quorum sensing and biofilm development. Eukaryot Cell. 5(10):1770–1779.
doi:10.1128/EC.00219-06.

Alves R, Barata-Antunes C, Casal M, Brown AJP, Van Dijck P, Paiva S. 2020. Adapting to survive: How
Candida overcomes host-imposed constraints during human colonization. Filler SG, editor. PLoS Pathog.
16(5):e1008478. doi:10.1371/journal.ppat.1008478.

Askew C, Sellam A, Epp E, Mallick J, Hogues H, Mullick A, Nantel A, Whiteway M. 2011. The zinc cluster
transcription factor Ahr1p directs Mcm1p regulation of *Candida albicans* adhesion. Mol Microbiol.
79(4):940–953. doi:10.1111/j.1365-2958.2010.07504.x.

Bhandari J, Fitzpatrick AH, Crowell DN. 2010. Identification of a Novel Abscisic Acid-Regulated Farnesol
Dehydrogenase from Arabidopsis. Plant Physiology. 154(3):1116–1127. doi:10.1104/pp.110.157784.

Boone CHT, Gutzmann DJ, Kramer JJ, Atkin AL, Nickerson KW. 2022. Quantitative assay for farnesol and
the aromatic fusel alcohols from the fungus *Candida albicans*. Appl Microbiol Biotechnol.
doi:10.1007/s00253-022-12165-w.

Brasch J, Horter F, Fritsch D, Beck-Jendroschek V, Tröger A, Francke W. 2013 Aug. Acyclic sesquiterpenes
released by *Candida albicans* inhibit growth of dermatophytes. Med Mycol.:1–10.
doi:10.3109/13693786.2013.814174.

Bricmont PA, Daugherty JR, Cooper TG. 1991. The DAL81 gene product is required for induced
expression of two differently regulated nitrogen catabolic genes in Saccharomyces cerevisiae. Mol Cell
Biol. 11(2):1161–1166. doi:10.1128/mcb.11.2.1161-1166.1991.

Bruno VM, Kalachikov S, Subaran R, Nobile CJ, Kyratsous C, Mitchell AP. 2006. Control of the C. albicans
cell wall damage response by transcriptional regulator Cas5. PLoS Pathog. 2(3):e21.
doi:10.1371/journal.ppat.0020021.

Chen H, Fink GR. 2006. Feedback control of morphogenesis in fungi by aromatic alcohols. Genes Dev.
20(9):1150–1161. doi:10.1101/gad.1411806.

Chen H, Fujita M, Feng Q, Clardy J, Fink GR. 2004. Tyrosol is a quorum-sensing molecule in *Candida*
albicans. Proc Natl Acad Sci USA. 101(14):5048–5052. doi:10.1073/pnas.0401416101.

Chung D, Kim SY, Ahn J-H. 2017. Production of three phenylethanoids, tyrosol, hydroxytyrosol, and
salidroside, using plant genes expressing in Escherichia coli. Sci Rep. 7(1):2578. doi:10.1038/s41598-017-
02042-2.

Costa CP, Bezerra AR, Almeida A, Rocha SM. 2020. Candida Species (Volatile) Metabotyping through
Advanced Comprehensive Two-Dimensional Gas Chromatography. Microorganisms. 8(12):1911.
doi:10.3390/microorganisms8121911.

Dai J, Xia H, Yang C, Chen X. 2021. Sensing, Uptake and Catabolism of L-Phenylalanine During 2-
Phenylethanol Biosynthesis via the Ehrlich Pathway in Saccharomyces cerevisiae. Front Microbiol.
12:601963. doi:10.3389/fmicb.2021.601963.

Dumitru R, Hornby JM, Nickerson KW. 2004. Defined anaerobic growth medium for studying *Candida*
albicans basic biology and resistance to eight antifungal drugs. Antimicrob Agents Chemother.
48(7):2350–2354. doi:10.1128/AAC.48.7.2350-2354.2004.

Dumitru R, Navarathna DHMLP, Semighini CP, Elowsky CG, Dumitru RV, Dignard D, Whiteway M, Atkin
AL, Nickerson KW. 2007. In vivo and in vitro anaerobic mating in *Candida albicans*. Eukaryot Cell.
6(3):465–472. doi:10.1128/EC.00316-06.

Fu Y, Filler SG, Spellberg BJ, Fonzi W, Ibrahim AS, Kanbe T, Ghannoum MA, Edwards JE. 1998. Cloning
and characterization of CAD1/AAF1, a gene from *Candida albicans* that induces adherence to endothelial
cells after expression in Saccharomyces cerevisiae. Infect Immun. 66(5):2078–2084.
doi:10.1128/IAI.66.5.2078-2084.1998.

García-Sánchez S, Mavor AL, Russell CL, Argimon S, Dennison P, Enjalbert B, Brown AJP. 2005. Global
roles of Ssn6 in Tup1- and Nrg1-dependent gene regulation in the fungal pathogen, Candida albicans.
Mol Biol Cell. 16(6):2913–2925. doi:10.1091/mbc.e05-01-0071.

Ghosh S, Kebaara BW, Atkin AL, Nickerson KW. 2008. Regulation of aromatic alcohol production in
Candida albicans. Appl Environ Microbiol. 74(23):7211–7218. doi:10.1128/AEM.01614-08.

Gow NAR, van de Veerdonk FL, Brown AJP, Netea MG. 2012. *Candida albicans* morphogenesis and host
defence: discriminating invasion from colonization. Nat Rev Microbiol. 10(2):112–122.
doi:10.1038/nrmicro2711.

Hazelwood LA, Daran J-M, van Maris AJA, Pronk JT, Dickinson JR. 2008. The Ehrlich pathway for fusel
alcohol production: a century of research on Saccharomyces cerevisiae metabolism. Appl Environ
Microbiol. 74(8):2259–2266. doi:10.1128/AEM.02625-07.

Hernday AD, Lohse MB, Nobile CJ, Noiman L, Laksana CN, Johnson AD. 2016. Ssn6 Defines a New Level of
Regulation of White-Opaque Switching in *Candida albicans* and Is Required For the Stochasticity of the
Switch. mBio. 7(1):e01565-01515. doi:10.1128/mBio.01565-15.

Homann OR, Dea J, Noble SM, Johnson AD. 2009. A phenotypic profile of the *Candida albicans*
regulatory network. PLoS Genet. 5(12):e1000783. doi:10.1371/journal.pgen.1000783.

Hornby JM, Jensen EC, Lisec AD, Tasto JJ, Jahnke B, Shoemaker R, Dussault P, Nickerson KW. 2001.
Quorum sensing in the dimorphic fungus *Candida albicans* is mediated by farnesol. Appl Environ
Microbiol. 67(7):2982–2992. doi:10.1128/AEM.67.7.2982-2992.2001.

Hussein B, Huang H, Glory A, Osmani A, Kaminskyj S, Nantel A, Bachewich C. 2011. G1/S transcription
factor orthologues Swi4p and Swi6p are important but not essential for cell proliferation and influence
hyphal development in the fungal pathogen *Candida albicans*. Eukaryot Cell. 10(3):384–397.
doi:10.1128/EC.00278-10.

Inglis DO, Johnson AD. 2002. Ash1 protein, an asymmetrically localized transcriptional regulator,
controls filamentous growth and virulence of *Candida albicans.* Mol Cell Biol. 22(24):8669–8680.
doi:10.1128/MCB.22.24.8669-8680.2002.

Kadosh D. 2019. Regulatory mechanisms controlling morphology and pathogenesis in *Candida albicans.*
Curr Opin Microbiol. 52:27–34. doi:10.1016/j.mib.2019.04.005.

Kebaara BW, Langford ML, Navarathna DHMLP, Dumitru R, Nickerson KW, Atkin AL. 2008. *Candida*
albicans Tup1 is involved in farnesol-mediated inhibition of filamentous-growth induction. Eukaryot Cell.
7(6):980–987. doi:10.1128/EC.00357-07.

Langford ML, Hargarten JC, Patefield KD, Marta E, Blankenship JR, Fanning S, Nickerson KW, Atkin AL.
2013. *Candida albicans* Czf1 and Efg1 coordinate the response to farnesol during quorum sensing, white-
opaque thermal dimorphism, and cell death. Eukaryot Cell. 12(9):1281–1292. doi:10.1128/EC.00311-12.

Lo H-J, Köhler JR, DiDomenico B, Loebenberg D, Cacciapuoti A, Fink GR. 1997. Nonfilamentous *C.*
albicans Mutants Are Avirulent. Cell. 90(5):939–949. doi:10.1016/S0092-8674(00)80358-X.

Lu Y, Su C, Unoje O, Liu H. 2014. Quorum sensing controls hyphal initiation in *Candida albicans* through
Ubr1-mediated protein degradation. Proc Natl Acad Sci U S A. 111(5):1975–1980.
doi:10.1073/pnas.1318690111.

Martínez P, Ljungdahl PO. 2005. Divergence of Stp1 and Stp2 transcription factors in *Candida albicans*
places virulence factors required for proper nutrient acquisition under amino acid control. Mol Cell Biol.
25(21):9435–9446. doi:10.1128/MCB.25.21.9435-9446.2005.

McCluskey K, Wiest A, Plamann M. 2010. The Fungal Genetics Stock Center: a repository for 50 years of
fungal genetics research. J Biosci. 35(1):119–126. doi:10.1007/s12038-010-0014-6.

Moreno-Velásquez SD, Tint SH, Del Olmo Toledo V, Torsin S, De S, Pérez JC. 2020. The Regulatory
Proteins Rtg1/3 Govern Sphingolipid Homeostasis in the Human-Associated Yeast *Candida albicans*. Cell
Rep. 30(3):620-629.e6. doi:10.1016/j.celrep.2019.12.022.

Murad AM, Leng P, Straffon M, Wishart J, Macaskill S, MacCallum D, Schnell N, Talibi D, Marechal D,
Tekaia F, et al. 2001. NRG1 represses yeast-hypha morphogenesis and hypha-specific gene expression in
Candida albicans. EMBO J. 20(17):4742–4752. doi:10.1093/emboj/20.17.4742.

Neville BA, d'Enfert C, Bougnoux M-E. 2015. *Candida albicans* commensalism in the gastrointestinal
tract. Munro C, editor. FEMS Yeast Research. 15(7):fov081. doi:10.1093/femsyr/fov081.

Nicholls S, Straffon M, Enjalbert B, Nantel A, Macaskill S, Whiteway M, Brown AJP. 2004. Msn2- and
Msn4-like transcription factors play no obvious roles in the stress responses of the fungal pathogen
Candida albicans. Eukaryot Cell. 3(5):1111–1123. doi:10.1128/EC.3.5.1111-1123.2004.

Noble SM, Gianetti BA, Witchley JN. 2017. *Candida albicans* cell-type switching and functional plasticity
in the mammalian host. Nat Rev Microbiol. 15(2):96–108. doi:10.1038/nrmicro.2016.157.

Pfaller MA, Diekema DJ. 2007. Epidemiology of Invasive Candidiasis: a Persistent Public Health Problem.
Clin Microbiol Rev. 20(1):133–163. doi:10.1128/CMR.00029-06.

Rauceo JM, Blankenship JR, Fanning S, Hamaker JJ, Deneault J-S, Smith FJ, Nantel A, Mitchell AP. 2008.
Regulation of the *Candida albicans* cell wall damage response by transcription factor Sko1 and PAS
kinase Psk1. Mol Biol Cell. 19(7):2741–2751. doi:10.1091/mbc.e08-02-0191.

Rodrigues CF, Černáková L. 2020. Farnesol and Tyrosol: Secondary Metabolites with a Crucial quorum-
sensing Role in Candida Biofilm Development. Genes. 11(4):444. doi:10.3390/genes11040444.

Romano A. 1966. Dimorphism. The fungi. 2:181–209.

Saville SP, Lazzell AL, Monteagudo C, Lopez-Ribot JL. 2003. EngineeredControl of Cell Morphology In Vivo
Reveals Distinct Roles for Yeast andFilamentous Forms of *Candida albicans* duringInfection. Eukaryot
Cell. 2(5):1053–1060. doi:10.1128/EC.2.5.1053-1060.2003.

Sellam A, Chaillot J, Mallick J, Tebbji F, Richard Albert J, Cook MA, Tyers M. 2019. The p38/HOG stress-
activated protein kinase network couples growth to division in *Candida albicans.* PLoS Genet.
15(3):e1008052. doi:10.1371/journal.pgen.1008052.

Shchepin R, Hornby JM, Burger E, Niessen T, Dussault P, Nickerson KW. 2003. Quorum sensing in
Candida albicans: probing farnesol's mode of action with 40 natural and synthetic farnesol analogs.
Chem Biol. 10(8):743–750. doi:10.1016/s1074-5521(03)00158-3.

Skrzypek MS, Binkley J, Binkley G, Miyasato SR, Simison M, Sherlock G. 2017. The Candida Genome
Database (CGD): incorporation of Assembly 22, systematic identifiers and visualization of high
throughput sequencing data. Nucleic Acids Res. 45(D1):D592–D596. doi:10.1093/nar/gkw924.

Soussi-Boudekou S, Vissers S, Urrestarazu A, Jauniaux JC, André B. 1997. Gzf3p, a fourth GATA factor
involved in nitrogen-regulated transcription in Saccharomyces cerevisiae. Mol Microbiol. 23(6):1157–
1168. doi:10.1046/j.1365-2958.1997.3021665.x.

Tanaka N, Mukai Y. 2015. Yeast Cyc8p and Tup1p proteins function as coactivators for transcription of
Stp1/2p-dependent amino acid transporter genes. Biochemical and Biophysical Research
Communications. 468(1–2):32–38. doi:10.1016/j.bbrc.2015.11.001.

Turner SA, Ma Q, Ola M, Martinez de San Vicente K, Butler G. 2018. Dal81 Regulates Expression of
Arginine Metabolism Genes in *Candida parapsilosis*. Mitchell AP, editor. mSphere. 3(2):e00028-18.
doi:10.1128/mSphere.00028-18.

Wang H, Song W, Huang G, Zhou Z, Ding Y, Chen J. 2011. *Candida albicans* Zcf37, a zinc finger protein, is
required for stabilization of the white state. FEBS Lett. 585(5):797–802.
doi:10.1016/j.febslet.2011.02.005.

Xie JL, Qin L, Miao Z, Grys BT, Diaz JDLC, Ting K, Krieger JR, Tong J, Tan K, Leach MD, et al. 2017. The
Candida albicans transcription factor Cas5 couples stress responses, drug resistance and cell cycle
regulation. Nat Commun. 8(1):499. doi:10.1038/s41467-017-00547-y.

Zordan RE, Miller MG, Galgoczy DJ, Tuch BB, Johnson AD. 2007. Interlocking transcriptional feedback
loops control white-opaque switching in *Candida albicans.* PLoS Biol. 5(10):e256.
doi:10.1371/journal.pbio.0050256.

Tables
Table 1. Summary of transcription regulator mutants with statistically significant differences in farnesol
accumulation.[a/]

Mutant	*S. cerevisiae* ortholog or best hit from CGD (Skrzypek et al. 2017)	Mean whole culture E,E-farnesol log$_2$ fold change vs WT control ± SD (n=6)	Mean supernatant fraction E,E-farnesol log$_2$ fold change vs WT control ± SD (n=6)	Predicted or known biological process(es) in *Candida albicans* adapted from CGD (Skrzpek et al. 2017)
E,E-farnesol over accumulating mutants				
swi4Δ/Δ	SWI4	2.57*** ± 0.39	3.11*** ± 0.50	Component of SBF complex, G1/S progression
ahr1Δ/Δ	LYS14	2.27*** ± 0.42	2.56 ***± 0.32	Regulation of adhesion genes, white/opaque switch, repressor of START
nrg1Δ/Δ	NRG2	1.73*** ± 0.45	1.93*** ± 0.49	Regulator of hyphal gene induction, virulence, chlamydospore formation, stress response
rap1Δ/Δ	RAP1	1.15*** ± 0.62	2.42*** ± 0.48	Telomere maintenance
pho4Δ/Δ	PHO4	1.07*** ± 0.53	1.68*** ± 0.23	Phosphate acquisition
tye7Δ/Δ	TYE7	0.95** ± 0.21	1.83*** ± 0.07	Regulator of glycolysis and carbohydrate metabolism
ash1Δ/Δ	ASH1	0.82* ± 0.43	0.92** ± 0.28	GATA-like transcription factor required for WT virulence and filamentous growth
aaf1Δ/Δ	none	0.76* ± 0.31	1.21*** ± 0.19	Expression in *S. cerevisiae* increases adherence
rtg1Δ/Δ	RTG1	0.63 ± 0.49	0.96** ± 0.20	Regulation of sphingolipid homeostasis, galactose catabolism
ssn6Δ/Δ	CYC8	0.20 ± 0.90	0.99*** ± 0.73	Forms a corepressor complex with Tup1, regulates a distinct set of

				genes different than Tup1 in *C. albicans*
wor2Δ/Δ	UME6	0.43 ± 0.82 (p=0.999)	0.59 ± 0.83 (p=0.217)	Regulator of white/opaque switching, maintenance of opaque state
tac1Δ/Δ [#]	HAL9	0.23 ± 0.56 (p=0.999)	0.36 ± 0.26 (p=0.919)	Transcriptional activator of drug-responsive genes *CDR1* and *CDR2*
csr1Δ/Δ [#]	ZAP1	0.32 ± 0.31 (p=0.984)	0.60 ± 0.13 (p=0.195)	Transcription factor, role in zinc homeostasis
upc2Δ/Δ [#]	UPC2	0.11 ± 0.43 (p=0.999)	0.31 ± 0.46 (p=0.980)	Regulator of ergosterol biosynthetic genes and sterol uptake
tup1Δ/Δ [#]	TUP1	0.0 ± 0.15 (p=0.999)	-0.02 ± 0.47 (p=0.999)	Transcriptional corepressor with *Ssn6p*, regulates morphological switching, farnesol response, germ tube formation
E,E-farnesol under accumulating mutants				
orf19.6874 Δ/Δ (C2_05640 W_A)	BAS1	-2.10*** ± 0.73	-2.65*** ± 1.03	Transcription factor, null mutant exhibits increase in invasive growth
msn4Δ/Δ	MSN2	-1.89*** ± 0.37	-2.40*** ± 0.65	Zinc finger transcription factor, not a significant stress response regulator in *C. ablicans*
cph2Δ/Δ	HMS1	-1.93*** ± 0.63	-2.04*** ± 0.91	Transcriptional activator of hyphal growth
fcr1Δ/Δ	CAT8	-1.97*** ± 0.88	-3.68*** ± 3.83	Negative regulator of fluconazole/ketoconazole/brefeldin A resistance
sko1Δ/Δ	SKO1	-1.93*** ± 0.39	-1.41*** ± 0.46	Cell wall damage response, morphological switching, oxidative stress response
gzf3Δ/Δ	GZF3	-1.32*** ± 0.32	-1.63*** ± 0.54	GATA type transcription factor, ortholog regulates nitrogen catabolic gene expression

cas5Δ/Δ	*MOT3*	-1.16** ± 0.13	-0.51 ± 0.32	Cell wall homeostasis, adherence, stress response, cell cycle
cup9Δ/Δ	*TOS8*	-1.08** ± 0.63	-0.85 ± 0.86	Represses SOK1 expression in response to farnesol inhibition
tec1Δ/Δ	*TEC1*	-0.75* ± 0.66	-0.58 ± 0.76	Hyphal gene regulator, white cell pheromone response

a/ Log_2 fold change production values ± SD represent the mean farnesol accumulation value of 3 X and 3
Y (n=6) independently derived mutant compared to the mean farnesol accumulation value of the paired
SN152 parent (n=22. Both over and under accumulating mutants are ordered by how much they exceed
the SN152 parent. The four mutants added because of literature precedent are indicated by #.
Differences between mutants (n=6) and the SN152[+] parent (n=22 batches) were accessed by One-way
ANOVA with Dunnett's multiple comparisons test using GraphPad Prism. Differences were considered
significant at $p < 0.05$ (*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$).

520 Table 2. Summary of transcription regulator mutants with statistically significant differences in fusel
521 alcohol accumulation.[a/]

Mutant	*S. cerevisiae* ortholog or best hit from CGD (Skrzypek et al. 2017)	Mean supernatant fraction log$_2$ fold change vs WT control ± SD (n=6)			Predicted or known biological process(es) in *Candida albicans* adapted from CGD (Skrzpek et al. 2017)
		Phenethyl alcohol	Tyrosol	Tryptophol	
Aromatic fusel alcohol over accumulating mutants					
19.2612Δ/Δ (CR_02120C)	SWI5	1.38*** ± 0.32	1.14* ± 0.62	1.39*** ± 0.40	zinc finger transcription factor of unknown function
fgr15Δ/Δ	*none*	1.06*** ± 0.37	0.63 ± 1.27	1.01** ± 0.22	Transposon mutation affects filamentous growth
gzf3Δ/Δ	GZF3	0.76*** ± 0.26	-0.24 ± 0.34	-0.04 ± 1.62	Ortholog negatively regulates nitrogen catabolic gene expression
hap5Δ/Δ	HAP5	0.13 ± 0.17	1.49*** ± 0.25	-0.15 ± 0.26	Regulator of genes involved in filamentous and respiratory growth; iron homeostasis
Aromatic fusel alcohol under accumulating mutants					
tup1Δ/Δ	TUP1	-4.90*** ± 0.20	-3.45*** ± 2.70	-3.56*** ± 0.45	Transcriptional corepressor with Ssn6, regulates morphological switching, farnesol response, germ tube formation
ssn6Δ/Δ	CYC8	-3.82*** ± 0.18	-2.26*** ± 1.61	-3.25*** ± 0.57	Forms a corepressor complex with Tup1, regulates a distinct set of genes different than Tup1 in *C. albicans*
stp2Δ/Δ	STP2	-3.10*** ± 0.19	0.20 ± 0.49	-2.58*** ± 0.38	Amino-acid-regulated; activator of amino acid permease genes

rap1Δ/Δ	RAP1	-2.01*** ± 0.61	-1.31* ± 1.05	-1.71*** ± 0.45	Telomere maintenance
dal81Δ/Δ	DAL81	-1.75*** ± 0.70	-1.17 ± 1.00	-1.87*** ± 0.89	Ortholog involved in activation of nitrogen-degradation genes
nrg1Δ/Δ	NRG2	-1.67*** ± 0.44	-0.24 ± 0.16	-2.39*** ± 1.31	Regulator of hyphal gene induction, virulence, chlamydospore formation, stress response
stp4Δ/Δ	STP3	-1.44*** ± 0.47	-1.37* ± 0.86	-1.32*** ± 0.64	transcription factor, paralogs have a role in amino acid metabolism
swi4Δ/Δ	SWI4	-1.33*** ± 0.30	0.00 ± 0.33	-0.94* ± 0.38	Component of SBF complex, G1/S progression
ahr1Δ/Δ	LYS14	-0.87*** ± 0.26	-0.70 ± 0.13	0.52 ± 0.17	Regulation of adhesion genes, white/opaque switch, repressor of START
czf1Δ/Δ	UME6	-0.81*** ± 0.28	-1.07 ± 0.73	-0.86 ± 0.31	Regulator of white/opaque switching and hyphal growth
ndt80Δ/Δ	NDT80	-0.81*** ± 0.22	-1.56** ± 0.81	-1.05** ± 0.21	meiosis-specific transcription factor; activator of *CDR1* induction by antifungal drugs
cph2Δ/Δ	HMS1	-0.78** ± 0.54	-1.12 ± 0.71	-1.16** ± 0.32	Transcriptional activator of hyphal growth
bcr1Δ/Δ	USV1	-0.58 ± 0.47	-1.51** ± 0.19	-1.01* ± 0.43	Regulator of a/α biofilm formation, adherence, cell surface genes, fluconazole resistance
cas5Δ/Δ	MOT3	-0.49 ± 0.28	0.12 ± 0.41	-1.11** ± 0.39	Cell wall homeostasis, adherence, stress response, cell cycle
aro80Δ/Δ [#]	ARO80	0.10 ± 0.19	-1.45** ± 0.82	0.16 ± 0.27	Activator of aromatic amino acid catabolism; regulator of aromatic alcohol biosynthesis

a/ Log_2 fold change production values ± SD represent the mean aromatic fusel alcohol accumulation
value of 3 X and 3 Y (n=6) independently derived mutant compared to the mean aromatic fusel alcohol
accumulation value of the paired SN152 parent (n=22). False positives are excluded. Both over and
under accumulating phenethyl alcohol mutants are ordered by how much they exceed the SN152
parent. Differences between mutants (n=6) and the SN152[+] parent (n=22 batches) were accessed by
One-way ANOVA with Dunnett's multiple comparisons test using GraphPad Prism. Differences were
considered significant at $p<0.05$ (*$p<0.05$, **$p<0.01$, ***$p<0.001$). Note that *fgr15Δ/Δ, stp2Δ/Δ,*
dal81Δ/Δ, and aro80Δ/Δ only had a 1 independent mutant, so these were not accessed in the Y plate,
but instead 6 replicates in the X plate.

Figure Captions

Fig. 1. Distribution of wild-type and transcription regulator mutant whole culture analytes from the

initial screen. Phenethyl alcohol, E,E-farnesol, tyrosol, and tryptophol were measured 24 hours post

inoculation following growth at 30°C in YPD. Data shown are the means of the initial screen

accumulation values for the SN152$^+$ parent (n=34) and transcription regulator mutants (n=164).

Fig. 2. Initial screen whole culture accumulation of phenethyl alcohol vs E,E-farnesol. Analytes were

measured 24 hours post inoculation at 30°C in YPD. Accumulation values are for each transcription

regulator mutant (black) (n=164) and each SN152$^+$ replicate (red) (n=34). The red X indicates the mean

value for the SN152$^+$ replicates (n=34).Dotted lines represent a log2 fold change increase or decrease, of

1 for each analyte from the means for SN152$^+$.

Fig. 3. Initial screen E,E-farnesol supernatant to pellet ratio vs E,E- farnesol whole culture accumulation

at 24 hours post inoculation. Data represents the accumulation values for each transcription regulator

mutant (black) (n=164) and each SN152$^+$ replicate (red) (n=34). The red X indicates the mean value for

the SN152$^+$ replicates (n=34). Dotted lines represent a log2 fold change increase or decrease of 1 for

each analyte from the mean for SN152$^+$.

Fig. 4. Initial screen whole culture accumulation of phenethyl alcohol vs tryptophol. Values were

measured 24 hours post inoculation at 30°C in YPD for each transcription regulator mutant (black)

(n=164) and each SN152$^+$ replicate (red) (n=34). The red X indicates the mean value for the SN152$^+$

replicates (n=34).

Fig. 5. Initial screen whole culture accumulation of phenethyl alcohol vs tyrosol. Values were measured

at 24 hours post inoculation at 30°C in YPD medium for each transcription regulator mutant (black)

(n=164) and each SN152$^+$ replicate (red) (n=34). The red X indicates the mean value for the SN152$^+$

replicates (n=34).

Fig. 6. Transcription regulator mutants differing in farnesol accumulation and localization. Accumulation

values were measured 24 hours post inoculation at 30°C in YPD for each transcription regulator mutant

in both the X (3) and Y (3) independent mutants (total n=6) and the SN152[+] parent (n=22 batches). Data

represents the mean accumulation value ± SD (A,B) or mean farnesol supernatant to pellet ratio ± SD

(C). The mutants added because of literature precedent are indicated by #. Differences between groups

that were normally distributed and that had equal variances (C) were accessed by One-way ANOVA with

Dunnett's multiple comparisons test. Differences between groups that were not normally distributed or

heteroscedastic (A,B) were accessed by Kruskal-Wallis test with Dunn's multiple comparisons tests.

Differences were considered significant at $p<0.05$ (*$p<0.05$, **$p<0.01$, ***$p<0.001$).

Fig. 7. Transcription regulator mutants differing in aromatic fusel alcohol accumulation. Accumulation

values were measured 24 hours post inoculation at 30°C in YPD for each transcription regulator mutant

in both the X (3) and Y (3) independent mutants (total n=6) and the SN152[+] parent (n=22 batches). Data

represents the mean accumulation value ± SD for mutants over-accumulating (A) or under-accumulating

(B) aromatic fusel alcohols. The mutants added because of literature precedent are indicated by #.

Differences between groups that were not normally distributed or heteroscedastic (A,B) were accessed

by Kruskal-Wallis test with Dunn's multiple comparisons tests. Differences were considered significant at

$p<0.05$ (*$p<0.05$, **$p<0.01$, ***$p<0.001$).

Initial Screen: Analyte Distribution

Normalized Analyte Amount (ng/uL) / 50 mL Dry weight (g)

Analyte

WT - Phenethyl alcohol
Mutant - Phenethyl alcohol
WT - E,E-Farnesol
Mutant - E,E-Farnesol
WT - Tyrosol
Mutant - Tyrosol
WT - Tryptophol
Mutant - Tryptophol

Initial Screen: Phenethyl alcohol vs E,E-farnesol Whole Culture Accumulation

Initial Screen: E,E-Farnesol Supernatant to Pellet Ratio VS Whole Culture Accumulation

Initial Screen: Phenethyl alcohol vs Tryptophol Whole Culture Accumulation

Initial Screen: Phenethyl alcohol vs Tyrosol Whole Culture Accumulation